Lysine methylation shields an intracellular pathogen from ubiquitylation

One Sentence Summary: Lysine methylation blocks the ubiquitin attack

Patrik Engström[1*], Thomas P. Burke[1], Anthony T. Iavarone[2], Matthew D. Welch[1*]

[1]Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA.

[2]QB3/Chemistry Mass Spectrometry Facility, University of California, Berkeley, CA 94720, USA.

*Correspondence: pengstrom@berkeley.edu (P.E.), welch@berkeley.edu (M.D.W)

ABSTRACT

Many intracellular pathogens stay invisible to the host detection machinery to promote their survival. However, it remains unknown how pathogen surfaces are disguised from host ubiquitin tagging, a first step in anti-microbial autophagy. We determined that outer membrane proteins (OMPs) of the intracellular bacterial pathogen *Rickettsia parkeri* are protected from ubiquitylation by protein-lysine methyltransferases (PKMTs) and the bacterial O-antigen. Analysis of the lysine-methylome revealed that PKMTs modify a subset of OMPs including surface protein OmpB. Mechanistically, methylation of lysines in OmpB camouflaged the same residues from ubiquitylation. Lysine methylation also prevented autophagy recognition and elimination by the autophagy factor ATG5 in macrophages and was critical for disease in mice. These findings suggest that lysine methylation shields proteins from ubiquitylation to evade autophagy targeting.

MAIN

Intracellular bacterial pathogens generally avoid the host immune surveillance machinery. This includes avoidance of surface targeting by the host ubiquitylation machinery and subsequent formation of a polyubiquitin coat, a first step in cell-autonomous immunity (*1-5*). The ubiquitin coat recruits autophagy receptors that engage with the autophagy machinery to target cytosol-exposed microbes for destruction (*1, 4-8*). Bacterial outer membrane proteins (OMPs) are targets for the host ubiquitylation machinery (*9, 10*), and the tick-borne obligate intracellular pathogen *Rickettsia parkeri* (*R. p.*) requires the abundant surface protein OmpB to protect OMPs from ubiquitylation (*11*). However, the detailed mechanisms that *R. p.* and other pathogens use to block lysine ubiquitylation of surface proteins, including OMPs, are unknown. We hypothesized that cell surface structures or modifications could provide protection at the molecular level. One such modification is lysine methylation, which is widespread in prokaryotes, archaea, and eukaryotes. This modification involves the transfer of one, two, or three methyl groups to the amino group of a lysine side chain, the same amino group that can also be modified by ubiquitin (*12*). Whether lysine methylation of bacterial surfaces prevents host detection and promotes intracellular survival has not been explored.

To identify bacteria-derived surface modifications that protect against ubiquitin coating, we screened pools of *R. p.* transposon mutants (*13*) (**Table S1**) for increased polyubiquitylation (pUb) relative to wild-type (WT) in Vero cells (an epithelial cell line commonly used to propagate and study intracellular pathogens) by immunofluorescence microscopy (**Fig. 1A**). We then analyzed individual mutants from pUb-positive pools and identified 4 mutants that were ubiquitylated, similar to *ompB* mutant bacteria (*11*) (**Fig. 1B, D,** and **E**). However, in contrast to the *ompB* mutant, these 4 strains expressed OmpB (**Fig. S1**). Two of the strains had insertions in the protein-lysine methyltransferase genes *pkmt1* and *pkmt2*, which are located at two distinct

chromosomal regions. The remaining two strains had insertions in the *wecA* and *rmlD* genes,

which are required for the biosynthesis of O-antigen (**Fig. S2**), a common surface structure in

Gram-negative bacteria. As a control, we analyzed a strain with a mutation in the *mrdA* gene,

which is required for peptidoglycan biosynthesis and cell shape in other bacteria (*14*). This mutant

strain had altered shape but was not polyubiquitylated (**Fig. 1B, D** and **E**), suggesting that not all

bacterial cell envelope structures are required to avoid ubiquitylation. We also further quantified

the pUb levels and observed that the *pkmt1*::tn bacteria had the highest levels (**Fig. 1E**). These

data indicate that OmpB, PKMTs and the O-antigen protect *R. p.* from ubiquitylation.

The O-antigen was previously shown to be required for rickettsial pathogenesis (*15*), and

OmpB was previously found to be required for *R. p.* to cause lethal disease in *Ifnar*[-/-] *Ifngr*[-/-] mice

lacking the type I interferon receptor (IFNAR) and IFN-γ receptor (IFNGR) (*16*). We therefore

examined whether PKMT1 or PKMT2 are important for causing disease *in vivo* by infecting *Ifnar*

[-/-]*Ifngr*[-/-] mice. We observed that mice succumbed to WT but not to *pkmt1*::tn or *pkmt2*::tn bacteria

(**Fig. 1F**). Mice infected with the *pkmt1*::tn mutant showed no signs of disease, whereas mice

infected with *pkmt2*::tn showed a transient loss in body weight (**Fig. S3**). This indicates that

PKMT1 and PKMT2 are virulence factors that promote *R. p.* pathogenesis.

Because PKMT1 or PKMT2 had previously been shown to methylate OmpB *in vitro* (*17-*

20), we next examined whether methylation and the O-antigen protect OmpB, or another

abundant outer membrane protein OmpA (*21*), from ubiquitylation. First, Vero cells

overexpressing 6xHis-tagged ubiquitin were infected with WT and the *ompB*, *pkmt1*::tn, *pkmt2*::tn,

wecA::tn, and *rmlD*::tn strains. 6xHis-tagged ubiquitin was recruited to the surface of all of the

mutants, but not WT bacteria, as observed by immunofluorescence microscopy (**Fig. S4**). Then,

6xHis-ubiquitylated proteins were affinity purified from infected cells, and OmpA and OmpB were

detected by Western blotting. OmpA was shifted towards higher molecular weights in cells

infected with mutant, but not WT bacteria, indicating OmpA is ubiquitylated (**Fig. 2A**). Similarly, in

comparison with WT, OmpB was also shifted towards higher as well as lower molecular weights

in cells infected with the *pkmt1*::tn and *pkmt2*::t mutants, and to a lesser extent with the *wecA*::tn

and *rmlD*::tn mutants, suggesting that OmpB is ubiquitylated (**Fig. 2A**). To confirm that

methylation protects OmpB and OmpA from ubiquitylation on the bacterial surface, we performed

pUb-enrichments of surface fractions from purified bacteria followed by Western blotting. This

revealed that both OmpB and OmpA shifted towards higher molecular weights in the

methyltransferase mutants but not in WT bacteria (**Fig. 2B**). These data demonstrate that

methylation is critical to protect OMPs from ubiquitylation on the bacterial surface.

Based on our observation that methylation protects both OmpA and OmpB from

ubiquitylation, we set out to determine how frequently, and to what extent, lysines of *R. p.* OMPs

are methylated. Peptides with methylated lysines from whole WT bacteria were quantified using

label-free liquid chromatography-mass spectrometry (LC-MS). We then analyzed lysine

methylation frequency in abundant OMPs as well as other abundant *R. p.* proteins (**Table S2**).

This analysis revealed that *R. p.* OmpB, OmpA, and surface cell antigen 2 (Sca2) proteins had

the highest abundance of methylated peptides. Lysine methylation was also detected in the outer-

membrane assembly protein BamA and in a predicted outer membrane protein porin

(WP_014410329.1; from here on referred to as OMP-porin) (**Fig. 3A** and **Table S3**). We next

mapped both methylated and unmethylated lysines on the above-mentioned OMPs and found

that more than 50% of lysines detected from OmpB and OmpA were methylated, and a significant

fraction of lysines were also methylated in Sca2 (31%), OMP-porin (30%), and BamA (27%) (**Fig.**

3B and **Fig. S5**). Thus, in *R. p.*, lysine methylation of OMPs is common.

To identify OMPs that are methylated by PKMT1 and PKMT2 during infection, we

compared lysine methylation frequencies in WT with those of *pkmt1*::tn or *pkmt2*::tn mutant

bacteria using LC-MS. We found that monomethylation of OmpB, OmpA, the predicted OMP-

porin, and another surface cell antigen protein Sca1 was reduced in *pkmt1*::tn compared to WT

bacteria (**Fig. 3C** and **Fig. S6**). Dimethylation of rickettsial surface proteins was not reduced in

the mutants (**Fig. S6**). Although trimethylation was rare and therefore difficult to analyze at the

individual protein level (**Fig. S7**), OmpB had reduced trimethylation levels in both

methyltransferase mutants (**Fig. 3C**). Notably, the frequency of unmethylated lysines in OmpB

was specifically increased in *pkmt1*::tn bacteria (**Fig. 3C** and **Fig. S6**). Lysine methylation of five

other surface proteins (Sca1, Sca2, BamA, LomR, and Pal-lipoprotein), and 21 of 23 abundant

proteins with different predicted subcellular distributions, was not affected by mutations in the

pkmt1 or *pkmt2* genes (**Fig. S6**). These data indicate that the PKMTs are required for methylation

of a subset of OMPs including OmpB.

To determine which OmpB residues are modified by PKMT1 and PKMT2 during *R. p.*

infection, we analyzed the methylation frequency of individual lysines in the mutants compared to

WT using LC-MS. We observed reduced monomethylation frequencies of OmpB K418, K623,

K634, K902, K1061, K1294, and K1323 in *pkmt1*::tn bacteria compared with WT, and reduced

trimethylation frequencies on K1061 and K388 in the *pkmt2*::tn strain (**Fig. 3D**). This indicates

that several lysines in *R. p.* OmpB are methylated by PKMT1 and PKMT2 during infection.

Although these data are consistent with previous biochemical results indicating that PKMT1

monomethylates and PKMT2 trimethylates OmpB's lysines (*17, 19*), we find that total OmpB-

methylation is unaffected in the *pkmt2*::tn mutant. This suggests that PKMT1 is the primary

methyltransferase for *R. p.* OmpB and that it can compensate, at least partly, for a deficiency in

PKMT2.

To test the hypothesis that methylation of specific lysines in OmpB shields the same

residues from ubiquitylation, we performed pUb-enrichments of bacterial surface fractions

followed by LC-MS to quantify lysines with diglycine (diGly) remnants, a signature for ubiquitin

after trypsin digestion. A prediction of this hypothesis is that individual lysines that are heavily

methylated in OmpB of *Rickettsia* species (*17*) including WT *R. p.* (**Fig. 3D**) are targets for

ubiquitylation in *pkmt1*::tn bacteria. We confirmed this hypothesis as OmpB K634 and K623 in

pkmt1::tn exhibited 7 to 10000-fold increased ubiquitylation compared with WT bacteria (**Fig. 3E,**

F and **Fig. S8**; **Table S4**). Furthermore, we observed a 13-fold increase in ubiquitylation of the

OMP-porin in *pkmt1*::tn bacteria (**Table S4**), indicating that methylation also protects additional

OMPs. However, in *pkmt2*::tn, differential OMP-ubiquitylation was below detection limits (**Fig. 3E,**

F and **Fig. S8**; **Table S4**). Together, these data indicate that methylation of lysines in OMPs by

PKMT1 camouflages the same residues from ubiquitylation.

Because polyubiquitylation promotes recruitment of the autophagy receptors

p62/SQSTM1 and NDP52 (*7, 8*), we hypothesized that lysine methylation and the O-antigen

shield OMPs from ubiquitylation to block recruitment of these proteins. Consistent with this

hypothesis, we observed that the majority of *pkmt1*::tn, *pkmt2*::tn, O-antigen (*wecA*::tn, *rmlD*::tn),

as well as *ompB* mutant bacteria, co-localized with p62 and NDP52 by immunofluorescence

microscopy (**Fig. S9**). These data demonstrate that PKMTs and the O-antigen protect *R. p.* from

autophagy recognition.

Many pathogenic bacteria including *R. p.* grow in immune cells such as macrophages (*11*),

despite the fact that microbial detection in such cells triggers anti-bacterial pathways. We

therefore investigated whether PKMT1 or PKMT2 were required for evading autophagy targeting

and bacterial growth in cultured bone-marrow-derived macrophages (BMDMs), as was observed

for OmpB (*11*). BMDMs were generated from control mice and mice lacking the gene encoding

for autophagy related 5 (ATG5), a protein required for optimal membrane envelopment around

pathogens targeted by autophagy, and for their subsequent destruction (*6*). We observed that

pkmt1::tn mutant bacteria were unable to grow in control BMDMs (*Atg5^flox/flox^*), and that growth

was rescued in *Atg5*-deficient BMDMs (*Atg5^-/-^*). Further, >91% of *pkmt1*::tn bacteria were labeled

with both pUb and p62 in *Atg5*-deficient BMDMs (**Fig. 4A, B, C** and **D**), suggesting that detected

bacteria are not restricted when the autophagy cascade is prevented. In contrast, the *pkmt2*::tn

mutant was not grossly defective in growth compared with WT bacteria and 50% of the bacteria

were labeled p62, irrespective of host genotype (**Fig. 4A, B, C** and **D**), consistent with less

pronounced ubiquitylation phenotypes compared to *pkmt1*::tn bacteria. Altogether, these data

indicate that methylation is required for *R. p.* growth in macrophages by avoiding autophagy

targeting.

Our work reveals a molecular mechanism involving lysine methylation that camouflages

bacterial surface proteins from host detection. In particular, we found that lysine methylation is

essential for blocking ubiquitylation, a first step in cell-autonomous immunity (*1-4*). This highlights

an intricate evolutionary arms race between pathogens and hosts and reveals a strategy that

pathogens can adapt to counteract host responses. The lysine methyltransferases PKMT1 and

PKMT2 are conserved between rickettsial species (**Fig. S10** and **Fig. S11**) and contain a core

Rossmann fold found in the broader superfamily of class I methyltransferases that exist in diverse

organisms (*12, 18*). Thus, we propose that lysine methylation, and potentially other lysine

modifications, could be used by pathogens, symbionts, and perhaps even in eukaryotic

organelles, to prevent unwanted surface ubiquitylation and downstream consequences including

elimination by autophagy. Further study of microbial surface modifications will continue to

enhance our understanding of the pathogen-host interface and could ultimately lead to new

therapeutic interventions to treat human diseases including those caused by infectious agents.

REFERENCES AND NOTES

1. F. Randow, R. J. Youle, Self and nonself: how autophagy targets mitochondria and bacteria. *Cell Host Microbe* **15**, 403-411 (2014).
2. A. J. Perrin, X. Jiang, C. L. Birmingham, N. S. So, J. H. Brumell, Recognition of bacteria in the cytosol of Mammalian cells by the ubiquitin system. *Curr Biol* **14**, 806-811 (2004).
3. S. J. L. van Wijk *et al.*, Linear ubiquitination of cytosolic *Salmonella* Typhimurium activates NF-kappaB and restricts bacterial proliferation. *Nat Microbiol* **2**, 17066 (2017).
4. J. Noad *et al.*, LUBAC-synthesized linear ubiquitin chains restrict cytosol-invading bacteria by activating autophagy and NF-kappaB. *Nat Microbiol* **2**, 17063 (2017).
5. E. D. Case *et al.*, The *Francisella* O-antigen mediates survival in the macrophage cytosol via autophagy avoidance. *Cell Microbiol* **16**, 862-877 (2014).
6. J. Huang, J. H. Brumell, Bacteria-autophagy interplay: a battle for survival. *Nat Rev Microbiol* **12**, 101-114 (2014).
7. Y. T. Zheng *et al.*, The adaptor protein p62/SQSTM1 targets invading bacteria to the autophagy pathway. *J Immunol* **183**, 5909-5916 (2009).
8. T. L. Thurston, G. Ryzhakov, S. Bloor, N. von Muhlinen, F. Randow, The TBK1 adaptor and autophagy receptor NDP52 restricts the proliferation of ubiquitin-coated bacteria. *Nat Immunol* **10**, 1215-1221 (2009).
9. E. Fiskin, T. Bionda, I. Dikic, C. Behrends, Global analysis of host and bacterial ubiquitinome in response to *Salmonella* Typhimurium infection. *Mol Cell* **62**, 967-981 (2016).
10. M. Polajnar, M. S. Dietz, M. Heilemann, C. Behrends, Expanding the host cell ubiquitylation machinery targeting cytosolic *Salmonella*. *EMBO Rep* **18**, 1572-1585 (2017).
11. P. Engström *et al.*, Evasion of autophagy mediated by *Rickettsia* surface protein OmpB is critical for virulence. *Nat Microbiol* **4**, 2538-2551 (2019).
12. S. Lanouette, V. Mongeon, D. Figeys, J. F. Couture, The functional diversity of protein lysine methylation. *Mol Syst Biol* **10**, 724 (2014).
13. R. L. Lamason, N. M. Kafai, M. D. Welch, A streamlined method for transposon mutagenesis of *Rickettsia parkeri* yields numerous mutations that impact infection. *PLoS One* **13**, e0197012 (2018).
14. B. G. Spratt, A. B. Pardee, Penicillin-binding proteins and cell shape in *E. coli*. *Nature* **254**, 516-517 (1975).
15. H. K. Kim, R. Premaratna, D. M. Missiakas, O. Schneewind, *Rickettsia conorii* O antigen is the target of bactericidal Weil-Felix antibodies. *Proc Natl Acad Sci U S A* **116**, 19659-19664 (2019).
16. T. P. Burke *et al.*, Inflammasome-mediated antagonism of type I interferon enhances Rickettsia pathogenesis. *Nat Microbiol* **5**, 688-696 (2020).
17. D. C. H. Yang, A. H. Abeykoon, B. E. Choi, W. M. Ching, P. B. Chock, Outer membrane protein OmpB methylation may mediate bacterial virulence. *Trends Biochem Sci* **42**, 936-945 (2017).
18. A. H. Abeykoon *et al.*, Structural insights into substrate recognition and catalysis in outer membrane protein B (OmpB) by protein-lysine methyltransferases from *Rickettsia*. *J Biol Chem* **291**, 19962-19974 (2016).
19. A. Abeykoon *et al.*, Multimethylation of *Rickettsia* OmpB catalyzed by lysine methyltransferases. *J Biol Chem* **289**, 7691-7701 (2014).
20. A. H. Abeykoon *et al.*, Two protein lysine methyltransferases methylate outer membrane protein B from *Rickettsia*. *J Bacteriol* **194**, 6410-6418 (2012).

21. N. F. Noriea, T. R. Clark, T. Hackstadt, Targeted knockout of the *Rickettsia rickettsii* OmpA surface antigen does not diminish virulence in a mammalian model system. *mBio* **6**, (2015).

22. R. L. Lamason *et al.*, *Rickettsia* Sca4 Reduces Vinculin-Mediated Intercellular Tension to Promote Spread. *Cell* **167**, 670-683 e610 (2016).

23. U. Distler *et al.*, Drift time-specific collision energies enable deep-coverage data-independent acquisition proteomics. *Nat Methods* **11**, 167-170 (2014).

24. R. S. Plumb *et al.*, UPLC/MS(E); a new approach for generating molecular fragment information for biomarker structure elucidation. *Rapid Commun Mass Spectrom* **20**, 1989-1994 (2006).

25. P. V. Shliaha, N. J. Bond, L. Gatto, K. S. Lilley, Effects of traveling wave ion mobility separation on data independent acquisition in proteomics studies. *J Proteome Res* **12**, 2323-2339 (2013).

26. S. Nahnsen, C. Bielow, K. Reinert, O. Kohlbacher, Tools for label-free peptide quantification. *Mol Cell Proteomics* **12**, 549-556 (2013).

27. K. A. Neilson *et al.*, Less label, more free: approaches in label-free quantitative mass spectrometry. *Proteomics* **11**, 535-553 (2011).

28. C. I. Carlström *et al.*, (Per)chlorate-reducing bacteria can utilize aerobic and anaerobic pathways of aromatic degradation with (per)chlorate as an electron acceptor. *mBio* **6**, (2015).

ACKNOWLEDGEMENTS

We thank Dr. Hwan Kim (Stony Brook University) for kindly providing the O-antigen antibody and for fruitful discussions. We are also grateful for the *Atg5$^{flox/flox}$* and *Atg5$^{-/-}$* BMDMs provided by Dr. G. Golovkine, a member of the laboratory of Prof. Jeffery S. Cox (UC Berkeley), and Prof. Michael Rape at UC Berkeley for fruitful discussions. **Funding:** P.E. was supported by a postdoctoral fellowship from the Sweden-America Foundation. M.D.W. was supported by NIH/NIAID grants R01 AI109044 and R21 AI109270. A mass spectrometer used in this study was purchased with support from the NIH (grant 1S10 OD020062-01). **Author contributions:** P.E. conceived the study with the assistance of M.D.W. P.E. performed laboratory work and analysis, except for mass spectrometry, which was conducted together with A.T.I, and animal experiments, which were conducted by T.P.B. P.E. drafted the initial manuscript, and A.T.I, T.P.B, and M.W.D provided editorial feedback. **Data and materials availability:** All data used in the analysis are provided in the main text or supplementary materials. Materials are available upon request.

FIGURES

237
Figure 1. The O-antigen and lysine methylation are virulence factors that protect _R. p._ from
ubiquitylation. (**A**) Pools of _R. p._ (green) mutants screened for increased pUb (red) via

immunofluorescence microscopy. DNA, blue. *ompB* mutant (*11*), positive control; WT, negative
control. White circle, pUb-positive bacteria; arrows pUb-negative bacteria. Scale bar, 20 µm. (**B**)
Vero cells infected with bacterial strains at 72 hours post-infection (h.p.i.) stained as in **A**. Scale
bar, 5 µm (representative of *n* = 4). (**C**) A schematic of biological function of the genes identified
in **A** and **B**. (**D**) Percentage of bacteria co-localized with pUb at 72 h.p.i. Data are the mean ±
s.e.m.; *n* = 4 (WT, *ompB^STOP^*::tn, *pkmt1*::tn, *pkmt2*::tn, *wecA*::tn, and *rmlD*::tn) and 2 for *mrdA*::tn;
≥ 108 bacteria were counted for each infection. Statistical comparisons between WT and mutants
were performed using a one-way ANOVA with Dunnett's post-hoc test; **** $P < 0.0001$. (**E**) pUb
signal per bacteria as determined from micrographs. Lines indicate the means; *n* = 3 fields of
vision; ≥ 35 bacteria per field of vision were analyzed, totaling ≥ 122 bacteria. Statistical
comparisons were performed using a Kruskal-Wallis test with Dunn's post-hoc test; **** $P <$
0.0001. (**F**) Survival of *Ifnar^-/-^Ifngr^-/-^* mice intravenously infected with 5×10^6 WT, *pkmt1*::tn, or
pkmt2::tn bacteria (*n* = 5 mice, WT; *n* = 6 mice, *pkmt1*::tn and *pkmt2*::tn, combined from two
independent experiments. Statistical comparison between WT and mutants was performed using
a two-way ANOVA and a Log-rank (Mantel-Cox) test; ** $P < 0.01$).

Figure 2. Lysine methylation and the O-antigen protect OMPs from ubiquitylation. (**A**) Western blot of His-Ub input and pull-down samples from infected control and 6xHis-ubiquitin expressing cells, probed for OmpB, OmpA, and pUb (representative of n = 3). (**B**) pUb-enriched (TUBE-1, pan specific) samples from purified bacteria probed for OmpB, OmpA, pUb, and OmpW (OmpB and OmpA of endogenous molecular weight represent non-specific binding to TUBE-1 beads) (representative of n = 3). Asterisks indicate OmpB and OmpA that exhibit increased molecular weight, indicating ubiquitylation. Arrows indicate OmpB and OmpA of endogenous molecular weight.

Figure 3. Methylation camouflages lysines from ubiquitylation. (**A**) Percentage lysine
methylation abundance of total among the abundantly detected proteins in WT *R. p.* determined
by LC-MS (data are combined from *n* = 2). (**B**) Methylated lysines (K) are indicated with residue
number in each OMP, and lysines in *R. prowazekii* OmpB known to be methylated in green (*17*).
K78, K131, K149, K312, and K547 in the surface domain of OmpB were unmethylated (data are
combined from *n* = 5, using data-independent (*n* = 3) and data-dependent acquisition modes (*n*
= 2). (**C**) Percentage of total abundances of the lysines methylated in **B,** that is unmethylated,
mono-, di-, or tri-methylated (data are the mean of *n* = 2). (**D**) Percentage of individual lysines in
OmpB that are unmethylated, mono-, di- or tri-methylated. Only residues repeatedly detected in
all strains were analyzed (data are the mean of *n* = 2). (**E**) Heat map representation of OmpB
residues from *pkmt1*::tn or *pkmt2*::tn that have similar levels of K-diGly peptides (white boxes)
compared to WT, or with a 5-fold, or more, increase of K-diGly peptides (cyan) (*n* = 2). (**F**)
Abundances of ubiquitylated, unmethylated, mono-, di-, or tri-methylated OmpB K623 (data are
the mean of *n* = 2). Each experiment (*n*) was performed in technical triplicate.

Figure 4. Methylation prevents ATG5-dependent *R. p.* killing in macrophages. (**A**) Micrographs of infected control (*Atg5^flox/flox^*) and *Atg5*-deficient (*Atg5^-/-^*) BMDMs at 48 h.p.i., stained for DNA (blue), pUb (red), and p62 (green). Arrow indicate a bacterium positive for both pUb and p62. Scale bars, 5 μm (*n* = 3). (**B**) Growth curves of WT, *pkmt1*::tn, and *pkmt2*::tn bacteria in control and *Atg5^-/-^* BMDMs, as measured by genomic equivalents using qPCR (*n* = 3). (**C**) Quantification of the mean number of bacteria per cell (*n* = 3). (**D**) Quantification of the percentage of bacteria with p62 and pUb (*n* = 3). nd, not determined. Data are the mean ± s.e.m. Statistical comparisons in **B** and **C** were performed using a Brown-Forsyth and Welch ANOVA with Dunnett's post-hoc test; *, $P < 0.05$.

SUPPLEMENTARY MATERIAL FIGURE

Figure S1. Polyubiquitylated mutant *R. p.* strains are positive for OmpA and OmpB. (A) Micrographs of Vero cells infected with the indicated strains stained for OmpA (green, 13-3 antibody), OmpB (red, OmpB-antibody), and DNA (blue, Hoechst) at 72 h.p.i. (representative of

n = 3). Scale bar 5 µm. (**B**) Quantification of OmpB signal per bacteria. Lines indicate the means;

n = 3 fields of vision; ≥ 50 bacteria per field of vision were analyzed. Statistical comparisons were

performed using a Kruskal-Wallis test with Dunn's post-hoc test; **** *P* < 0.0001 between

indicated strains. Dashed line indicates that the majority of mutant bacterial populations, except

pkmt1::tn bacteria, have OmpB levels comparable to WT bacteria. (**C**) Quantification of OmpA

signal per bacteria as in **B**. (**D**) Western blot of $5\text{x}10^6$ purified bacteria probed for OmpB, pUb,

OmpW and EF-Tu (bacterial loading controls) (representative of *n* = 2).

303
Figure S2. The *wecA*::tn and *rmlD*::tn bacteria lack the O-antigen. (**A**) Micrographs of Vero
cells infected with the indicated strains stained for OmpA (green, 13-3 antibody), the O-antigen
(red, O-antigen antibody(*15*)) and DNA (blue, Hoechst) at 72 h.p.i. (representative of $n = 3$). Scale
bars, 5 µm. (**B**) Percentage of bacteria positive for the O-antigen at 72 h.p.i. $n = 2$; ≥ 90 bacteria
were counted in each infection (data are the mean ± s.e.m.; $n = 3$). (**C**) Western blot of 5x10^6
purified WT and mutant bacteria probed for the O-antigen. OmpW and EF-Tu were used as
bacterial loading controls (representative of $n = 3$).

Figure S3. PKMT1 plays a more significant role in causing disease *in vivo* compared to PKMT2. (A) Weight changes of *Ifnar^-/-^Ifngr^-/-^* mice intravenous infected with 5x10⁶ WT, *pkmt1*::tn, or *pkmt2*::tn bacteria (data are the mean ± s.e.m. *n* = 5, WT; *n* = 6, *pkmt1*::tn and *pkmt2*::tn, combined from two independent experiments). A two-way ANOVA from 0 to 25 days post-infection (d.p.i.) was used to statistically compare the weight changes between the *pkmt1*::tn and *pkmt2*::tn mutants. **** $P < 0.0001$.

318
319 **Figure S4. 6xHis-ubiquitin is recruited to the surface of mutant *R. p*.** Micrographs of infected
320 Vero cells expressing 6xHis-ubiquitin stained with anti-His antibody (red) and Hoechst (blue,
321 bacterial and host DNA), at 28 h.p.i. Scale bar, 5 µm (representative of $n = 2$).

OMP-porin K-CH₃(mono or di)

```
        65                     201        349   394
        64
        48
1 │ K KK  KK  KK     K KK        KKKK K K  KKK K │ 438
```

BamA K-CH₃(mono or di)

```
  51                 287      420              597      722
                     265      410                       710
                              408
1 │ K   K    K KK K K K   KKKKKKKKKKKKKKK       K K K  KK KK K     K K  KK │ 768
```

Sca4 -K-CH₃(mono or di)

```
  25    128                      476 525  589 648  757        912
  11    126                      461 568      634  740        901
        124                      455 560 555  723   714       890
        118                      445 560                      863
                                 438 476
1 │ KKK   KKKKKKK KKK    KKK K KKK K K   KKKK K KKKKKKKKK K KKKKKKKKKKKKK K KKKKKKKKKKKKKKKKKKKKK KKK  KK │ 1023
```

323 **Figure S5. Lysine-methylome reveals that *R. p.* OMP-porin, BamA, and the released factor**
324 **Sca4 are methylated.** Methylated lysines are indicated with residue number, and unmethylated
325 residues without, determined by LC-MS as in **Fig. 3B** (data is combined from five independent
326 experiments). See also Table S2.

327
328 **Figure S6. PKMT1 primarily modifies bacterial OMPs.** Percentage of total abundances of the
329 lysines methylated in WT, that is unmethylated, mono-, di-, or tri-methylated in respective strain.
330 Proteins with ≥3 lysines methylated detected in independent experiments are shown (mean of n
331 = 2, performed in technical triplicates). OmpB, OmpA and Sca2 data are the same as in **Fig. 3C**.
332 See also Table S2.

333
334 **Figure S7. Dimethylation is a common methylation state among the abundant _R. p._**
335 **proteins.** Percentage of total lysines abundances detected to as mono-, di-, or tri-methylated in
336 WT bacteria of the proteins indicated in Table S2 (mean of _n_ = 2, performed in technical
337 triplicates).

Figure S8. K623 and K634 are frequently ubiquitylated when PKMT1-mediated methylation
is reduced. (**A**) OmpB peptide abundance values after pUb-enrichments from respective strain,
as determined by LC-MS (data are mean pf $n = 2$, performed in triplicates). (**B**) Abundance values
of lysines in OmpB that are ubiquitylated, unmethylated, mono-, di-, or tri-methylated, in
respective strain after pUb-enrichments as in **Fig 3G.** Only residues detected in independent
experiments are shown (data are the mean of $n = 2$, performed in technical triplicates). K623 data
in **B** are the same as in **Fig. 3F**.

346
Figure S9. OmpB, methylation, and the O-antigen block recruitment of autophagy
receptors to *R. p*. (**A**) Micrographs of Vero cells infected with the indicated strains: stained for
bacterial and host DNA (blue, Hoechst), p62 (green, p62-antibody), and NDP52 (red, NDP52-
antibody) at 72 h.p.i ($n = 4$ for p62 staining; $n = 2$ for NDP52 staining). (**B**) Percentage of bacteria
that show rim-like surface localization of p62 at 72 h.p.i. Data are the mean ± s.e.m.; $n = 3$; ≥ 142
bacteria were counted for each infection. Statistical comparisons between WT and *ompB*^{STOP}::tn,
pkmt1::tn, *pkmt2*::tn, *wecA*::tn, *rmlD*::tn were performed using a one-way ANOVA with Tukey's
post-hoc test; **** $P < 0.0001$. (**C**) Quantification of p62 signal per bacteria from a representative
experiment. Lines indicate the means. ($n = 3$ fields of vision; ≥ 50 bacteria per field of vision were
analyzed). Statistical comparisons were performed using a Kruskal-Wallis test with Dunn's post-
hoc test; **** $P < 0.0001$ between indicated strains. (**D**) Quantification of NDP52 signal per
bacteria as in **C**.

Figure S10. The PKMT1 enzyme is highly conserved between diverse rickettsial species.
Amino acid sequence alignment of PKMT1 from *R. p.* (WP_014411082.1), *R. rickettsii*
(WP_012262600.1), *R. conorii* (WP_016926592.1), *R. typhi* (WP_011191207.1), *R. prowazekii*
(WP_004596928.1), *R. bellii* (WP_045799810.1), and a *Rickettsia* endosymbiont
(WP_062811822.1), using COBALT. Amino acids indicated in red are identical; blue, variation
between species; grey, one or more of the analyzed proteins are lacking this residue(s).

Rickettsia parkeri	1	MTKQANKISYDEVPYSPFTFSYTSPPYLRTIGKLFGLNPPPLETAKILELGCGIGVNLLNFAETYPKSQSLGVDLSKTQI	80
Rickettsia rickettsii	1	MTKQANKISYDEVPYSPFTFSYTSPPYLRTIGKLFGLNPPPLETAKILELGCGIGVNLLNFAETYPKSQSLGVDLSKTQI	80
Rickettsia conorii	1	MTKQANKISYDEVPYSPFTFSYTSPPYLRTIGKLFGLNPPPLETAKILELGCGIGVNLLNFAETYPKSQSLGVDLSKTQI	80
Rickettsia typhi	1	MIKKANKISYDEVPYPPFTFSYTYPPYLRTIGKLFGLNPPLLETAKVLDIGCGIGVNLLNFAETYPKSQSLGVDLSKTQI	80
Rickettsia prowazeki	1	MIKKTNKISYDEVPYPPFTFSYTYPPYLRTIGKLFGLNPPPLETAKILDIGCGVGVNLLNFAETYPKSQSLGVDLSKTQI	80

Rickettsia parkeri	81	ELGKKFISDLKIKNAELKALSILDLDESYGKFDYIVCHGVYSWVPEEVQDKILKVCNKLLNPNGIAFVSYNTLPGWNMQR	160
Rickettsia rickettsii	81	ELGKKIISDLKIKNAELKALSILDLDESYGKFDYIICHGVYSWVPEEVQDKILKVCNKLLNPNGIAFVSYNTLPGWNMQS	160
Rickettsia conorii	81	ELGKKIISDLKIKNAELKALSILDLDESYGKFDYIVCHGVYSWVPEEVQDKILKICNKLLNPNGIAFVSYNTLPGWNMQR	160
Rickettsia typhi	81	ELGKKTISDAKINNVELKALSILDLDESYGKFDYIVCHGVYSWVSQEVQDKILEVLNKLLNPNGIAFVSYNTLPGWNMQN	160
Rickettsia prowazeki	81	EIGKKTISDSKIKNVGLKALSILDLDESYGKFDYIVCHGVYSWVSKEVQDKILEVLNKLLNPNGIAFISYNTLPGWNMQN	160

Rickettsia parkeri	161	TIREMIMFHSELFNTSHDKLQQAKLLLKFINDSLESSTTPYSNFLRDETKLLSAYTDSYVLHEYLGEINTGIYFHQFIEK	240
Rickettsia rickettsii	161	TIREMIMFHSELFNTSHDKLQQAKLLLKFINDSLESSTTPYSNFLRDETKLLSAYTDSYVLHEYLGEINTGIYFHQFIEK	240
Rickettsia conorii	161	TIREMIMFHSELFNTSHDKLQQAKLLLKFINDSLESSTTPYSNFLRDETKLLSAYTDSYVLHEYLGEINTGIYFHQFIEK	240
Rickettsia typhi	161	TIREMMMFHSESFNTSHDKLQQARLLLKFINDSLGNSTTPYANFLRDEAKLISTYDDSYVLHEYLGEINTGTYFHQFIEK	240
Rickettsia prowazeki	161	TIREMMMFHSESFNTSHDKLQQSKLLLKFINDSLENSTTPYANFLREEAKLISTYADSYVLHEYLGEINTGTYFHQFIEK	240

Rickettsia parkeri	241	AQKNHLNYLGDTSLTAMFIGNLPTQAAEKLQAVNDIVRTEQYMDFITNRKFRSTLLCHQNIPINRKIEFNNLKEFFTSLN	320
Rickettsia rickettsii	241	AQKNHLNYLGDTSLTAMFIGNLPTQAAEKLQAVNDIVRTEQYMDFITNRKFRSTLLCHQNIPINRKIEFNNLKEFFTSLN	320
Rickettsia conorii	241	AQKNHLNYLGDTSLTAMFIGNLPTQAAEKLQAVNDIVRTEQYMDFITNRKFRSTLLCHQNIPINRKIEFNNLKEFFTSLN	320
Rickettsia typhi	241	AQKNHLNYLGDTSIAAMFIGNLPTKAASKLQAINDIVCTEQYMDFITNRKFRSTLLCHQNIPINRKIEFDNLKDFYTTFN	320
Rickettsia prowazeki	241	AQKNHLNYLGDTSITAMFIGNLPTKAAEKLQAINDIVRTEQYMDFITNRKFRSTLLCHQNIPINRKIEFENLKDFYTTFN	320

Rickettsia parkeri	321	IRPVILEKAVDLTNEQENVSFYYENLPEPFISTTSPIMKAILYVYAENISNPISLEQVAKEAFKKLGKYQLQDFLAALEQ	400
Rickettsia rickettsii	321	IRPVILEKAVDLTNEQENVSFYYENLPEPFISTTSPIMKAILYVYAENISNPISLEQVAKEAFKKLGKYQLQDFLAALEQ	400
Rickettsia conorii	321	IRPVILEKAVDLTNEQENVSFYYENLPEPFISTTSPIMKAILYVYAENISNPISLEQVAKEAFKKLGKYQLQDFLAALEQ	400
Rickettsia typhi	321	IRPISPENKIDLNNEQENISFYYENLPEPFISTTSAIMKAILYVYAENISNPIRLEQVAKEAFKKLGKYRLQDFLATLEQ	400
Rickettsia prowazeki	321	IRPISSENKIDLNNEQENISFYYENLPEPFISTTSAIMKAILYVYAENISNPIRLEQVAKEAFKKLGKYQLQDFLAILEQ	400

Rickettsia parkeri	401	HFIIFIFQGYLKIFETKPHAIATITEKPKTSEFARYQAKQAYFNNVTSVFSVTNRLNDMVGIPIHEKYILEMLDGTHNID	480
Rickettsia rickettsii	401	HFIIFIFQGYLKIFETKPHAIATITEKPKTSEFARYQAKQAYFNNVTSVFSVTNRLNDMVGIPIHEKYILEMLDGTHNID	480
Rickettsia conorii	401	HFIIFIFQGYLKIFETKPHAIATITEKPKTSEFARYQAKQAYFNNVTSVFSVTNRLNDMVGIPIHEKYILEMLDGTHNID	480
Rickettsia typhi	401	HFITLIFQGYLKIFETKPHAIATITEKPKTSQFARYQAKHAHFNNVTNMFSITNRLNDMIGIPIHEKYILEMLDGTHNID	480
Rickettsia prowazeki	401	HFITFIFQGYLKIFETKPHAIATITEKPKTSQFVRYQAKHAHFNNVTNMLSVTNRLNDMIGIPIHEKYILEMLDGTHNID	480

Rickettsia parkeri	481	DIKKGVLEKINSKLLTARDDKGQEVTDPKLLKEFVDYVVNTSLEKFRINYLLVE	534
Rickettsia rickettsii	481	DIKKGVLEKINSKLLTARDDKGQEVTDPKLLKEFVDYAVNTSLEKFRMNYLLVE	534
Rickettsia conorii	481	DIKKGVLEKINSKLLTARDDKGQEVTDPKLLKEFVDYVVNTSLEKFRINYLLVE	534
Rickettsia typhi	481	DIKKSIIEKINSKLLTACDNKGQVVTDPKLLKEFVDYVVAVSLEKFRINYLLVG	534
Rickettsia prowazeki	481	DIKKGMIEKINSKLLIACDNKGQAVTDPKLLKEFVDYIVNISLEKFRINYLLIG	534

Figure S11. The PKMT2 enzyme is highly conserved between virulent rickettsial species but absent from *R. bellii* and the *Rickettsia endosymbiont Proechinophthirus fluctus*. Amino acid sequence alignment of PKMT2 from *R. p.* (WP_014410272.1), *R. rickettsii* (WP_012150259.1), *R. conorii* (WP_016925880.1), *R. typhi* (WP_011190574.1), and *R. prowazekii* (WP_004596662.1) using COBALT. PKMT1 of *R. bellii* and the *Rickettsia endosymbiont Proechinophthirus fluctus* showed 51% identity to PKMT2 of *R. p.* using single alignment BLAST at NCBI; however, no PKMT2 variants could be found in these organisms, and therefore they were excluded from this analysis. Amino acids indicated in red are identical; blue, variation between species.

376
377 **Fig. S12**. Strain validation of clonality and insertion site using primers for flanking chromosomal
378 regions ($n = 1$).

Materials and Methods

Cell lines and primary mouse macrophages

Vero cells were purchased from the UC Berkeley Cell Culture Facility and the identity was repeatedly confirmed by mass-spectrometry analysis. Cells were grown at 37 °C and 5% CO_2 in DMEM plus 2% heat-inactivated (30 min, 56 °C, in a water-bath) fetal bovine serum (Gemcell). Vero cells were confirmed to be mycoplasma negative by DAPI staining and fluorescence microscopy screening at the UC Berkeley Cell Culture Facility.

BMDMs generated from the femurs of mutant $Atg5^{flox/flox}$ and matched $Atg5^{-/-}$ C57BL/6 mice were a kind gift from the laboratory of Jeffery S. Cox (UC Berkeley), and they were prepared as previously described (*11*) although in the absence of antibiotics. Genotypes were confirmed by PCR and Sanger sequencing at the UC Berkeley DNA Sequencing Facility, as previously described (*11*).

***Rickettsia parkeri* strain generation and validation**

R. parkeri strain Portsmouth (NCBI accession no. NC_017044.1; originally a gift from C. Paddock, Center for Disease Control and Prevention) were propagated and purified as described below, and bacterial stocks of WT, *ompB^{STOP}*::tn (the genome sequences of these bacterial strains are available at the Sequence Read Archive as accession no. SRP154218 (WT, SRX4401164; *ompB^{STOP}*::tn, SRX4401167), *pkmt1*::tn, *pkmt2*::tn, *wecA*::tn and *rlmD*::tn bacteria were prepared every ~6-10 months, and side-by-side experimental comparisons were made between stocks prepared at similar times.

R. parkeri pkmt1::tn, *pkmt2*::tn, *wecA*::tn, and 114 other mutant strains screened for pUb were previously isolated in a screen for small plaque mutants (*13, 22*). The *ompB^{STOP}*::tn was previously isolated in a suppressor screen and lacked expression of OmpB (*11*). The *rlmD*::tn, and 132 other mutant strains screened for pUb were isolated in an independent screen in which

mutants were isolated without regard for plaque size (**Table S1**). The genomic locations of

transposon insertion sites for all mutants were determined by semi-random nested PCR. To verify

the insertions and clonality, we used PCR reactions that amplified the transposon insertion site

using primers for flanking chromosomal regions: 5'GCTCACTAGATAGCACTCG'3 and

5'GCTCGATTTATCTCACTTTATG'3 for *rlmD*::tn, 5'CGTTTAATAGTCCAGTTAATTTGT'3 and

5'CCGTCTATACCGTCCATAAAAT'3 for *wecA*::tn, 5'GCATCGAAATAACCCTGAG'3 and

5'GCAAACTTCTCAAAGAAATTAACG'3 for *pkmt1*::tn,

5'GCTAAGAAATCTTCTAATTTGATATTTTAC'3 and 5'CGAAAATTTACCTGAGCCTT'3 for

pkmt2::tn, 5'CGACACATAATAGCACAAACTAC'3 and 5'GCGGAGGCGGTAGTAAAG'3 for

mrdA::tn (**Fig. S12**).

Screening for pUb-positive strains

To prepare the mutant library for screening, passage 1 (P1) transposon insertion mutants

were amplified one time in Vero cells using 24-well cell culture plates. At 5-12 d.p.i, when 50-70%

of the infected cells appeared to be rounded up (as a sign of infection) by visual inspection using

a light microscope, cell culture media were completely removed, and cells were subsequently

lysed in 500 µL cold sterile water for 2-3 minutes (min). Next, 500 µL of 2x cold sterile brain-heart-

infusion (BHI) broth (BD Difco, cat. no. 237500) was added to the lysed cells, resuspended, and

P2 bacteria were transferred to cryogenic storage vials and frozen at -80 °C.

To screen for pUb-positive strains, 10-40 µL of each of five to seven P2 mutant bacterial

strains were diluted in 1 mL of room temperature (RT) cell culture media supplemented with 2%

FBS. Subsequently, the pooled bacterial suspension was centrifuged at 250*g* for 4 min at RT onto

confluent Vero cells grown on coverslips in 24-well plates. Cells were then incubated at 33 °C and

fixed at 50-55 h.p.i. with pre-warmed 4% PFA for 10 min at RT. If cells were over-infected (i.e.,

individual infection foci had grown together) as determined by immunofluorescence microscopy,

infections of that specific pool were repeated using reduced volumes of P2 bacteria. Next, fixed

cells were permeabilized with 0.2% Triton-X for 5 min and then stained with the anti-*Rickettsia*

I7205 antibody (1:500 dilution; gift from Ted Hackstadt) and anti-polyubiquitin FK1 antibody (Enzo

Life Sciences, BML-PW8805-0500; 1:250 dilution), followed by Alexa 488 anti-rabbit antibody

(Invitrogen, A11008; 1:500 dilution) or goat anti-mouse Alexa-568 (Invitrogen, A11004). Whole

coverslips were manually inspected on a Nikon Ti Eclipse microscope with x60 (1.4 numerical

aperture) Plan Apo objective. The initial screen revealed that five out of 39 mutant pools contained

pUb-positive areas.

In a secondary screen, individual strains from the pUb-positive pools were used to infect

Vero cells, as stated above. Infected cells were also fixed and stained as above except that a

post-fixation step using 100% methanol for 5 min was included and that an OmpB antibody (*11*)

and Hoechst (Sigma, B2261, 1:2500 dilution) was used instead of the anti-*Rickettsia* I7205

antibody. Samples were inspected as above and strains were scored as following: **1**) pUb-

negative (49 strains), **2**) a few infection foci were pUb-positive (2 strains: Sp mutant 24, insertion

at bp position 753916; Sp mutant 94, insertion at bp position 774831), **3**) bacteria in the center of

foci were pUb-positive but not on the edges (2 strains: Sp mutant 43, insertion at bp position

651602-651604; Sp mutant 45, insertion at bp position 751156), **4**) almost all bacteria in all foci

were pUb-positive (4 strains: *pkmt1*::tn, insertion at bp position 1161553 (gene *MC1_RS06185*);

pkmt2::tn, insertion at bp position 34100 (gene *MC1_RS00180*); *wecA*::tn, insertion at bp position

1223170 (*MC1_RS06510*); and *rmlD*::tn, insertion at bp position 455753 (*MC1_RS02345*).

Rickettsia **purification**

R. parkeri strains were propagated as described previously (*11*). "Purified bacteria" were

from five T175 flasks of Vero cells growing in DMEM supplemented with 2% FBS that after 5-8

days of infection (normally ~75% infected as observed by light microscopy) were harvested in the

media using a cell scraper. Then, bacteria were centrifuged 12000g for 15 min at 4 °C in pre-

chilled tubes. Pellets were resuspended in cold K-36 buffer (0.05 M KH_2PO_4, 0.05 M K_2HPO_4, pH

7, 100 mM KCl and 15 mM NaCl) and a pre-chilled dounce-homogenizer (tight fit) were used for

60 strokes to release bacteria from host cells. The homogenate was then centrifuged at 200g for

5 min at 4 °C to remove cellular debris. The supernatant was overlaid onto cold 30% v/v MD-76R

(Mallinckrodt Inc., 1317-07) diluted in K-36, and centrifuged at 58300g for 20 min at 4 °C in an

SW-28 swinging-bucket rotor. The pellet was resuspended in cold 1x BHI broth (0.5 mL BHI per

infected T175 flask), and after letting DNA precipitates sediment to the bottom of the tubes,

bacterial suspensions were collected, aliquoted and frozen at -80 °C.

"Gradient-purified bacteria" were from ten T175 flasks of Vero cells, purified as above with

the addition of a 40/44/54% v/v MD-76R (diluted in K-36 buffer) gradient step centrifuged at

58300g for 25 min at 4 °C using the SW-28 swinging bucket rotor. The bacteria were then

collected from the 44-54% interface, diluted in K-36 buffer, and pelleted by centrifugation at

12000g for 15 min at 4 °C. The pellet was resuspended in cold 1x BHI broth and subsequently

aliquoted and frozen at -80 °C.

OmpW and EF-Tu antibody production

The sequence encoding amino acids 22-224 of outer membrane protein W (OmpW;

WP_014411122.1) (a protein that lacks the signal peptide), or full-length Elongation factor Tu (EF-

Tu; WP_004997779.1), were amplified by PCR from *R. parkeri* genomic DNA, and subsequently

cloned into plasmid pETM1, which encodes N-terminal 6xHis and maltose-binding proteins (MBP)

tags. From the resulting plasmids, fusion proteins were expressed in *E. coli* strain BL21 codon

plus RIL-Cam[r] (DE3) (QB3 Macrolab, UC Berkeley) by induction with 1 mM isopropyl-β-D-thio-

galactoside (IPTG) for 2-2.5 hours at 37 °C. Bacterial pellets were resuspended in lysis buffer (50

mM NaH_2PO_4, pH 8.0, 300 mM NaCl, 1mM EDTA, and 1 mM dithiothreitol (DTT)) and stored at -

80 °C. For protein purification, bacteria were thawed, lysozyme was added to 1 mg/mL (Sigma,

L4919), and lysis was carried out by sonication. Lysates containing 6xHis-MBP-OmpW and 6xHis-

MBP-OmpW were incubated on amylose resin (New England Biolabs, E8021L) (Qiagen,

1018244) and bound proteins were eluted in lysis buffer lacking EDTA and DTT but containing 10

mM maltose. Fractions were analyzed by SDS-PAGE and those with the highest concentrations

of fusion proteins were pooled to generate rabbit antibodies against OmpW and EFTU. 1.2 mg of

purified 6xHis-MBP-OmpW and 6xHis-MBP-EFTU proteins were sent to Pocono Rabbit Farm and

Laboratory (Canadensis, PA), and immunization was carried out according to their 91-d protocol.

Western blotting

To determine the levels of bacterial and host proteins in purified bacterial samples, 30%-

purified bacterial samples were boiled in 1x SDS loading buffer (150 mM Tris pH 6.8, 6% SDS,

0.3% bromophenol blue, 30% glycerol, 15% β-mercaptoethanol) for 10 min, then 1×10^6 PFUs

were resolved on an 8-12% SDS-PAGE gel and transferred to an Immobilon-FL polyvinylidene

difluoride membrane (Millipore, IPEL00010). Membranes were probed for 30 min at room

temperature or 4°C overnight with antibodies as follows: affinity-purified rabbit anti-OmpB

antibody (*11*) diluted 1:200-30000 in TBS-T (20 mM Tris, 150 mM NaCl, pH 8.0, 0.05% Tween 20

(Sigma, P9416)) plus 5% dry milk (Apex, 20-241); mouse monoclonal anti-OmpA 13-3 antibody

diluted 1:10000-50000 in TBS-T plus 5% dry milk; rabbit anti-OmpW serum diluted 1:8000 in TBS-

T plus 5% dry milk; mouse monoclonal FK1 anti-polyubiquitin antibody diluted 1:2500 in TBS-T

plus 2% BSA; rabbit anti-OmpW serum diluted 1:15000 in TBS-T plus 5% dry milk; or rabbit anti-

O-antigen serum 1:5000 in TBS-T plus 5% dry milk. Secondary antibodies were: mouse anti-

rabbit horseradish peroxidase (HRP) (Santa Cruz Biotechnology, sc-2357), or goat anti-mouse

HRP (Santa Cruz Biotechnology, sc-2005), diluted 1:1000-2500 in TBS-T plus 5% dry milk.

Secondary antibodies were detected with ECL Western Blotting Detection Reagents (GE,

Healthcare, RPN2106) for 1 min at room temperature, and developed using Biomax Light Film

(Carestream, 178-8207).

Immunofluorescence microscopy

R. parkeri infections were carried out in 24-well plates with sterile circle 12-mm coverslips

(Thermo Fisher Scientific, 12-545-80). To initiate infection, 30%-purified bacteria were diluted in

cell culture media at room temperature to a MOI of 0.01 for Vero cells, and a MOI of 0.1 for

BMDMs. Bacteria were centrifuged onto cells at 300g for 5 min at room temperature and

subsequently incubated at 33 °C. Next, infected cells were fixed for 10 min at room temperature

in pre-warmed (37 °C) 4% paraformaldehyde (Ted Pella Inc., 18505) diluted in PBS, pH 7.4, then

washed 3 times with PBS. Primary antibodies were the following: for staining with the guinea pig

polyclonal anti-p62 antibody (Fitzgerald, 20R-PP001; 1:500 dilution), mouse polyclonal anti-

NDP52 antibody (Novus Biologicals, H00010241-B01P; 1:100 dilution), a rabbit anti-*Rickettsia*

I7205 antibody (1:500 dilution; gift from Ted Hackstadt), or anti-polyubiquitin FK1 antibody (1:250

dilution), cells were permeabilized with 0.5% Triton-X100 for 5 min prior to staining. For staining

with mouse monoclonal anti-OmpA 13-3 antibody (1:5000 dilution), anti-OmpB antibody (*11*)

(1:1,000 dilution), or rabbit anti-O-antigen serum (*15*) (1:500 dilution), infected cells were post-

fixed in methanol for 5 min at RT (no Triton-X). Cells were then washed 3 x with PBS and

incubated with the primary antibody diluted as already indicated in PBS with 2% BSA for 30 min

at RT. To detect the primary antibodies, secondary goat anti-rabbit Alexa-568 (Invitrogen,

A11036), goat anti-mouse Alexa-568 (Invitrogen, A11004), or goat anti-guinea pig Alexa-568

(Invitrogen, A11075), Alexa 488 anti-rabbit antibody (Invitrogen, A11008; 1:500 dilution), Alexa

488 anti-mouse antibody (Invitrogen, A11001) antibodies were incubated at room temperature for

30 min (all 1:500 in PBS with 2% BSA). Images were captured as 15-25 z-stacks (0.1-µm step

size) on a Nikon Ti Eclipse microscope with a Yokogawa CSU-XI spinning disc confocal with 100X

(1.4 NA) Plan Apo objectives, and a Clara Interline CCD Camera (Andor Technology) using

MetaMorph software (Molecular Devices). Images were processed using ImageJ using z-stack

average maximum intensity projections and assembled in Adobe Photoshop. For quantification

of the percentage of bacteria with pUb and p62, only bacteria that co-localized with rim-like

patterns of the respective marker were scored as positive for staining. To quantify pUb, p62,

NDP52, OmpB, and OmpA signal per bacteria, z-stacks were projected as stated above, and the

edges of individual bacteria were marked by the freehand region of interest (ROI) function in

ImageJ. Subsequently, the average pixel intensity within that ROI was measured.

pUb/p62/NDP52 signal intensities were calculated by subtracting the average pUb/p62/NDP52

signal of WT bacteria from the pUb/p62/NDP52-value of each bacterium. OmpB signal intensity

was calculated by subtracting the average OmpB-signal of $ompB^{STOP}$::tn bacteria from the OmpB-

value of each bacterium. OmpA signal intensity was calculated by subtracting the average

background-signal (areas with no bacteria) from the OmpA-value of each bacterium.

Sample preparation for mass spectrometry to determine the lysine methylome

$5x10^7$ gradient-purified WT (Passage 6), $pkmt1$::tn (P4) and $pkmt2$::tn (P4) bacteria were

centrifuged at 11,000g for 3 min. Each pellet was resuspended in 50 µL Tris (10 mM)-EDTA (10

mM), pH 7.6, and incubated for 45 min in a 45 °C water bath. Bacterial surface fractions were

recovered from the supernatant after centrifugation at 11,000g for 3 min. Pellet was resuspended

as above and incubated for additional 45 min at 45 °C before resuspension in 50 µL Tris (10 mM)-

EDTA (10 mM). Both pellet and surface fractions were boiled at 95 °C for 10 min. Samples were

cooled to RT prior to addition of 20 µL 50 mM NH_4HCO_3, pH 7.5, and 50 µL of a 0.2% solution of

RapiGest (diluted in NH_4HCO_3, Waters, 186001861). Next, samples were heated at 80 °C for 15

min and cooled to RT before addition of 1 µg of trypsin (Promega, V511A). Samples were digested

at 37 °C overnight. To hydrolyze the RapiGest, 20 µL of 5% trifluoroacetic acid (TFA) was added

to samples which were incubated at 37 °C for 90 min prior centrifugation at 15000g for 25 min at

4 °C. Supernatant samples were desalted using C18 OMIX tips (Agilent Technologies,

A57003100) according to the manufacturer's instructions and sample volume was decreased to

20 µL using a SpeedVac vacuum concentrator. Samples were stored at 4 °C prior to analysis.

TUBE assay and sample preparation for mass spectrometry

To enrich for polyubiquitylated proteins, 3x10^8 PFUs of "purified" WT (P6) and *pkmt1*::tn

(P4) and *pkmt2*::tn (P3) bacteria were centrifuged at 14,000g for 3 min at room temperature. Next,

to release the surface protein fraction, the bacterial pellets were resuspended in lysis buffer (50

mM Tris-HCl, pH 7.5, 150 mM NaCl, 1 mM EDTA and 10% glycerol), supplemented with 0.0031%

v/v lysonase (Millipore, 71230), the deubiquitylase inhibitor PR619 at a final concentration of 20

µM (Life Sensor, SI9619) and 0.8% w/v octyl β-D-glucopyranoside (Sigma, O8001), and

incubated on ice for 10 min with occasional pipetting of samples to break pellet into smaller pieces.

Subsequently, the lysate was cleared by centrifugation at 14,000g at 4 °C for 5 min and incubated

with equilibrated agarose TUBE-1 (Life Sensor, UM401) for 3 h, at 4 °C. After binding of

polyubiquitylated proteins to TUBE-1, agarose beads were washed 1 time with TBS

supplemented with 0.05% Tween and 5 mM EDTA, and subsequently 3 times with TBS only (no

Tween or EDTA) and centrifuged at 5,000g for 5 min. To prepare samples for MS analysis,

enriched proteins were digested at 37 °C overnight on agarose beads in RapiGest SF solution

(Waters, 186001861) supplemented with 0.75 µg trypsin (Promega, V511A). The reaction was

stopped using 1% TFA (Sigma, T6508). Octyl β-D-glucopyranoside was extracted using water-

saturated ethyl acetate. Prior to submission of samples for mass spectrometry analysis, samples

were desalted using C18 OMIX tips (Agilent Technologies, A57003100), according to the

manufacturer's instructions.

Liquid chromatography-mass spectrometry

Samples of proteolytically digested proteins were analyzed using a Synapt G2-S*i* ion

mobility mass spectrometer that was equipped with a nanoelectrospray ionization source

(Waters). The mass spectrometer was connected in line with an Acquity M-class ultra-

performance liquid chromatography system that was equipped with trapping (Symmetry C18,

inner diameter: 180 µm, length: 20 mm, particle size: 5 µm) and analytical (HSS T3, inner

diameter: 75 µm, length: 250 mm, particle size: 1.8 µm) columns (Waters). Data-independent, ion

mobility-enabled, high-definition mass spectra and tandem mass spectra were acquired in the

positive ion mode (*23-25*). Data acquisition was controlled using MassLynx software (version 4.1),

and tryptic peptide identification and relative quantification using a label-free approach (*26, 27*)

were performed using Progenesis QI for Proteomics software (version 4.0, Waters). Raw data

were searched against *Rickettsia parkeri* and *Chlorocebus sabaeus* protein databases (National

Center for Biotechnology Information, NCBI) to identify tryptic peptides, allowing for up to three

missed proteolytic cleavages, with diglycine-modified lysine (i.e., ubiquitylation remnant) and

methylated lysine as variable post-translational modifications. Data-dependent analysis was

performed using an UltiMate3000 RSLCnano liquid chromatography system that was connected

in line with an LTQ-Orbitrap-XL mass spectrometer equipped with a nanoelectrospray ionization

source, and Xcalibur (version 2.0.7) and Proteome Discoverer (version 1.3, Thermo Fisher

Scientific, Waltham, MA) software, as described elsewhere (*28*).

Localization of tagged ubiquitin and ubiquitin pull-downs

To assess localization of 6xHis-ubiquitin during infection, confluent Vero cells grown in 24-

well plates with coverslips were transfected with 2 µg of pCS2-6xHis-ubiquitin plasmid DNA using

Lipofectamine 2000 (Invitrogen, 11668-019) for 6 h in Opti-MEM (Gibco, 31985-070).

Subsequently, media was exchanged to media without transfection reagent and cells were

incubated overnight at 37 °C and 5% CO_2. The following day (~16 h after transfection), transfected

cells were infected with purified WT or mutant bacteria at a MOI of 1. At 28 h.p.i., infected cells

were fixed with 4% paraformaldehyde (Ted Pella Inc., 18505) diluted in PBS, pH 7.4 for 10 min,

then washed 3 times with PBS. Primary anti-6xHis monoclonal mouse antibody (Clontech,

631212, diluted 1:1,000) was used to detect 6xHis-ubiquitin in samples permeabilized with 0.5%

Triton-X100, and a goat anti-mouse Alexa-568 (Invitrogen, A11004) to detect the primary 6xHis

antibody. Hoechst (Thermo Scientific, 62249, diluted 1:2500) was used to detect bacterial DNA.

Samples were imaged as already described.

For ubiquitin pull-downs, confluent Vero cells grown in 6-well plates were transfected and

infected as described above. At 28 h.p.i., cells were washed once with 1x PBS, pH 7.4, and

subsequently lysed in urea lysis buffer (8 M urea, 50 mM Tris-HCl, pH 8.0, 300 mM NaCl, 50 mM

Na_2HPO_4, 0.5% Igepal CA-630 (Sigma, I8896)) for 20 min at RT. Subsequently, samples were

sonicated, and lysate was cleared by centrifugation at 15000g for 15 min at room temperature.

Prior to incubation with Ni-NTA resin, an aliquot was saved for the input sample. 6xHis-ubiquitin

conjugates were purified by incubation and rotation with Ni-NTA resin for 3 h, at RT, in the

presence of 10 mM imidazole. Beads were washed 3 times with urea lysis buffer and 1 time with

urea lysis buffer lacking Igepal CA-630. Ubiquitin conjugates were eluted at 65 °C for 15 min in

2x Laemmli buffer containing 200 mM imidazole and 5% 2-mercaptoetanol (Sigma, M6250),

vortexed for 90 seconds, and centrifuged at 5000g for 5 min at RT. Eluted and input proteins were

detected by SDS-PAGE followed by Western blotting, as described above.

Animal experiments

Animal research using mice was conducted under a protocol approved by the UC Berkeley

Institutional Animal Care and Use Committee (IACUC) in compliance with the Animal Welfare Act.

The UC Berkeley IACUC is fully accredited by the Association for the Assessment and

Accreditation of Laboratory Animal Care International and adheres to the principles of the Guide

for the Care and Use of Laboratory Animals. Infections were performed in a biosafety level 2

facility. Mice were age-matched between 8 and 18 weeks old. Mice were selected for experiments

based on their availability, regardless of sex. All mice were healthy at the time of infection and

were housed in microisolator cages and provided chow and water. Littermates of the same sex

were randomly assigned to experimental groups. For infections, *R. parkeri* was prepared by

diluting 30%-prep bacteria into cold sterile 1x PBS to $5x10^6$ PFU per 200 μL. Bacterial

suspensions were kept on ice during injections. Mice were exposed to a heat lamp while in their

cages for approximately 5 min and then each mouse was moved to a mouse restrainer (Braintree,

TB-150 STD). The tail was sterilized with 70% ethanol, and 200 µL of bacterial suspensions were

injected using 30.5-gauge needles into the lateral tail vein. Body temperatures were monitored

using a rodent rectal thermometer (BrainTree Scientific, RET-3). Mice were monitored daily for

clinical signs of disease, such as hunched posture, lethargy, or scruffed fur. If a mouse displayed

severe signs of infection, as defined by a reduction in body temperature below 90 ˚F or an inability

to move around the cage normally, the animal was immediately and humanely euthanized using

CO_2 followed by cervical dislocation, according to IACUC-approved procedures (*16*).

Statistical analysis, experimental variability and reproducibility

Statistical parameters and significance are reported in the legends. Data were considered

to be statistically significant when $p < 0.05$, as determined by a one-way ANOVA with Dunnett's

post-hoc test, a Kruskal-Wallis test with Dunn's post-hoc test, a Brown-Forsyth and Welch

ANOVA with Dunnett's post-hoc test, or a two-way ANOVA (all two-sided). Statistical analysis

was performed using PRISM 6 software (GraphPad Software). If not otherwise described, *n*

indicates the number of independent biological experiments executed at different times.