Lysine methylation shields an intracellular pathogen from ubiquitylation

Patrik Engström[1*], Thomas P. Burke[1], Anthony T. Iavarone[2], Matthew D. Welch[1*]

[1]Department of Molecular and Cell Biology, University of California, Berkeley, CA 94720, USA.

[2]QB3/Chemistry Mass Spectrometry Facility, University of California, Berkeley, CA 94720, USA.

*Correspondence to: pengstrom@berkeley.edu (P.E.), welch@berkeley.edu (M.D.W).

Abstract

Many intracellular pathogens avoid detection by their host cells. However, it remains

unknown how they avoid being tagged by ubiquitin, an initial step leading to anti-microbial

autophagy. Here, we show that the intracellular bacterial pathogen *Rickettsia parkeri* uses

two protein-lysine methyltransferases (PKMTs) to modify outer membrane proteins (OMPs) and

prevent their ubiquitylation. Mutants deficient in the PKMTs were avirulent in mice and failed to

grow in macrophages due to ubiquitylation and autophagy. Analysis of the lysine-methylome

revealed that PKMTs modify a subset of OMPs by methylation at the same sites that are recognized

by host ubiquitin. These findings show that lysine methylation is an essential determinant of

rickettsial pathogenesis that shields bacterial proteins from ubiquitylation to evade autophagic

targeting.

Main

13  Intracellular pathogens generally evade the host immune surveillance machinery. This

14 includes avoidance of surface targeting by the host ubiquitylation machinery and subsequent

15 formation of a polyubiquitin coat, a first step in cell-autonomous immunity (*1-5*). The ubiquitin

16 coat recruits autophagy receptors that engage with the autophagy machinery to target cytosol-

17 exposed microbes for destruction (*1, 4-8*). Bacterial outer membrane proteins (OMPs) are targets

18 for the host ubiquitylation machinery (*9, 10*), and the tick-borne obligate intracellular pathogen

19 *Rickettsia parkeri* requires the abundant surface protein OmpB to protect OMPs from

20 ubiquitylation (*11*). However, the detailed mechanisms that *R. parkeri* and other pathogens use to

21 block lysine ubiquitylation of surface proteins, including OMPs, are unknown. We hypothesized

22 that cell surface structures or modifications could provide protection at the molecular level. One

23 such modification is lysine methylation, which is widespread in all domains of life. This

24 modification involves the transfer of one, two, or three methyl groups to the amino group of lysine

25 side chains, the same amino group that can also be modified by ubiquitin (*12*). Whether lysine

26 methylation of bacterial surfaces prevents host detection and promotes intracellular survival has

27 not been explored.

28  To identify bacteria-derived surface modifications that protect against ubiquitin coating,

29 we screened a library of *R. parkeri* transposon mutants (*13*) (**Table S1**) for increased

30 polyubiquitylation (pUb) relative to wild-type (WT) in Vero cells (an epithelial cell line commonly

31 used to propagate and study intracellular pathogens) by immunofluorescence microscopy (**Fig.

32 1A**). We subsequently analyzed individual mutants and identified 4 that were ubiquitylated, similar

33 to *ompB* mutant bacteria (*11*) (**Fig. 1B** and **D**). Importantly, these 4 strains expressed OmpB (**Fig.

34 S1**). Two of the strains had insertions in the protein-lysine methyltransferase genes *pkmt1* and

pkmt2, which are located in two distinct chromosomal regions. The remaining two strains had

insertions in the *wecA* and *rmlD* genes, which are required for the biosynthesis of O-antigen (**Fig.**

S2), a common surface molecule in Gram-negative bacteria. The O-antigen also protects

Francisella tularensis from ubiquitylation (*5*), suggesting it performs a conserved function in this

process. As a control, we analyzed a strain with a mutation in the *mrdA* gene, which is required

for peptidoglycan biosynthesis and cell shape in other bacteria (*14*). This mutant strain had altered

shape but was not polyubiquitylated (**Fig. 1B** and **D**), suggesting that not all bacterial cell envelope

structures are required to avoid ubiquitylation. In addition, we further quantified the pUb levels

and observed that the *pkmt1*::tn mutant, followed by the *ompB* and *pkmt2*::tn mutants, had the

highest levels (**Fig. 1E**). These data indicated that OmpB, PKMTs, and the O-antigen protect *R.*

parkeri from ubiquitylation.

The O-antigen was previously shown to be required for rickettsial pathogenesis (*15*), and

OmpB was previously found to be required for *R. parkeri* to cause lethal disease in *Ifnar*$^{-/-}$*Ifngr*$^{-/-}$

mice lacking the type I interferon receptor (IFNAR) and IFN-g receptor (IFNGR) (*16*). We

therefore examined whether PKMT1 or PKMT2 are important for causing disease *in vivo*. We

observed that *Ifnar*$^{-/-}$*Ifngr*$^{-/-}$ mice succumbed to infection with WT but not to *pkmt1*::tn or *pkmt2*::tn

bacteria (**Fig. 1F**). Mice infected with the *pkmt1*::tn mutant showed no signs of disease, whereas

mice infected with *pkmt2*::tn showed a transient loss in body weight (**Fig. S3**). This indicates that

PKMT1 and PKMT2 are determinants of *R. parkeri* pathogenesis.

Because PKMT1 or PKMT2 had previously been shown to methylate OmpB *in vitro* (*17-*

20), we next examined whether methylation protects OmpB, or another abundant outer membrane

protein OmpA (*21*), from ubiquitylation. First, Vero cells overexpressing 6xHis-tagged ubiquitin

were infected with WT and the *ompB*, *pkmt1*::tn, and *pkmt2*::tn strains. 6xHis-tagged ubiquitin

was recruited to the surface of all of the mutants, but not WT bacteria, as observed by

immunofluorescence microscopy (**Fig. 2A**). Then, 6xHis-ubiquitylated proteins were affinity-

purified from infected cells, and OmpA and OmpB were detected by western blotting. OmpA was

shifted towards higher molecular weights in cells infected with mutant, but not WT bacteria,

indicating OmpA is ubiquitylated (**Fig. 2B**). Similarly, in comparison with WT bacteria, OmpB

was also shifted towards higher as well as lower molecular weights in cells infected with the

pkmt1::tn and *pkmt2*::tn mutants, suggesting that OmpB is ubiquitylated (**Fig. 2B**). To confirm that

methylation protects OmpB and OmpA from ubiquitylation on the bacterial surface, we performed

pUb-enrichments of surface fractions from purified bacteria followed by western blotting. This

revealed that both OmpB and OmpA shifted towards higher molecular weights in the

methyltransferase mutants but not in WT bacteria (**Fig. 2C**). Thus, methylation is critical to protect

OMPs from ubiquitylation on the bacterial surface.

Based on our observation that methylation protects both OmpA and OmpB from

ubiquitylation, we set out to determine how frequently, and to what extent, lysines of *R. parkeri*

OMPs are methylated. Peptides with methylated lysines from whole WT bacteria were quantified

using label-free liquid chromatography-mass spectrometry (LC-MS). We then analyzed lysine

methylation frequency in abundant OMPs as well as other abundant *R. parkeri* proteins (**Table

S2**). This analysis revealed that *R. parkeri* OmpB, OmpA, and surface cell antigen 2 (Sca2)

proteins had the highest abundance of methylated peptides. Lysine methylation was also detected

in the outer-membrane assembly protein BamA and in a predicted outer membrane protein porin

(WP_014410329.1; from here on referred to as OMP-porin) (**Fig. 3A** and **Table S3**). We next

mapped both methylated and unmethylated lysines on the above-mentioned OMPs and found that

more than 50% of lysines detected from OmpB and OmpA were methylated, and a significant

fraction of lysines were also methylated in Sca2 (31%), OMP-porin (30%), and BamA (27%) (**Fig.**

3B and **Fig. S4**). Thus, in *R. parkeri*, lysine methylation of OMPs is common.

To identify OMPs that are methylated by PKMT1 and PKMT2 during infection, we

compared lysine methylation frequencies in WT with those of *pkmt1*::tn or *pkmt2*::tn mutant

bacteria using LC-MS. We found that monomethylation of OmpB, OmpA, the predicted OMP-

porin, and another surface cell antigen protein Sca1 were reduced in *pkmt1*::tn compared to WT

bacteria (**Fig. 3C** and **Fig. S5**). Dimethylation of rickettsial surface proteins was not reduced in the

mutants (**Fig. S6**). Although trimethylation was rare and therefore difficult to analyze at the

individual protein level (**Fig. S6**), OmpB had reduced trimethylation levels in both

methyltransferase mutants (**Fig. 3C**). Notably, the frequency of unmethylated lysines in OmpB

was specifically increased in *pkmt1*::tn bacteria (**Fig. 3C** and **Fig. S5**). Lysine methylation of five

other surface proteins (Sca1, Sca2, BamA, LomR, and Pal-lipoprotein), and 21 of 23 abundant

proteins with different predicted subcellular distributions, was not affected by mutations in the

pkmt1 or *pkmt2* genes (**Fig. S5**). These data indicate that the PKMTs are required for methylation

of a specific subset of OMPs including OmpB.

To determine which OmpB residues are modified by PKMT1 and PKMT2 during *R.*

parkeri infection, we analyzed the methylation frequency of individual lysines in the mutants

compared to WT using LC-MS. We observed reduced monomethylation frequencies of OmpB

K418, K623, K634, K902, K1061, K1294, and K1323 in *pkmt1*::tn bacteria compared with WT,

and reduced trimethylation frequencies on K1061 and K388 in the *pkmt2*::tn strain (**Fig. 3D**).

These data indicate that several lysines in *R. parkeri* OmpB are methylated by PKMT1 and

PKMT2 during infection. Although these results were consistent with previous biochemical results

indicating that PKMT1 monomethylates and PKMT2 trimethylates OmpB's lysines (*17, 19*), we

found that total OmpB-methylation is unaffected in the *pkmt2*::tn mutant. This suggests that

PKMT1 is a primary methyltransferase for OmpB and that it can compensate, at least partly, for a

deficiency in PKMT2.

To test the hypothesis that methylation of specific lysines in OmpB shields the same

residues from ubiquitylation, we performed pUb-enrichments of bacterial surface fractions

followed by LC-MS to quantify lysines with diglycine (diGly) remnants, a signature for ubiquitin

after trypsin digestion. A prediction of this hypothesis is that individual lysines that are heavily

methylated in OmpB of *Rickettsia* species (*17*), including WT *R. parkeri* (**Fig. 3D**), are targets for

ubiquitylation in *pkmt1*::tn bacteria. In support of this hypothesis, OmpB K634 and K623 in

pkmt1::tn exhibited 7 to 10,000-fold increased ubiquitylation compared with WT bacteria (**Fig.**

3E, F and **Fig. S7**; **Table S4**). Furthermore, we observed a 13-fold increase in ubiquitylation of

the OMP-porin in *pkmt1*::tn bacteria (**Table S4**), indicating that methylation also protects

additional OMPs. However, in the *pkmt2*::tn mutant, differential OMP-ubiquitylation was below

detection limits (**Fig. 3E, F** and **Table S4**). Together, these data indicate that methylation by

PKMT1 camouflages lysines in OMPs from ubiquitylation.

Because polyubiquitylation promotes recruitment of the autophagy receptors

p62/SQSTM1 and NDP52 (*7, 8*), we hypothesized that lysine methylation shields OMPs from

ubiquitylation to block recruitment of these proteins. Consistent with this hypothesis, we observed

that the majority of *pkmt1*::tn, *pkmt2*::tn, as well as *ompB* mutant bacteria, co-localized with p62

and NDP52 by immunofluorescence microscopy (**Fig. S8**). These data demonstrate that PKMTs

protect *R. parkeri* from autophagy recognition.

Many pathogenic bacteria including *R. parkeri* grow in immune cells such as macrophages

(*11*), despite the fact that microbial detection in such cells triggers anti-bacterial pathways. We

therefore investigated whether PKMT1 or PKMT2 were required for evading autophagy targeting

and bacterial growth in cultured bone marrow-derived macrophages (BMDMs), as was observed

for OmpB (*11*). BMDMs were generated from control mice and mice lacking the gene encoding

for autophagy related 5 (ATG5), a protein required for optimal membrane envelopment around

pathogens targeted by autophagy, and for their subsequent destruction (*6*). We observed that

pkmt1::tn mutant bacteria were unable to grow in control BMDMs (*Atg5$^{flox/flox}$*), and that growth

was rescued in *Atg5*-deficient BMDMs (*Atg5$^{-/-}$*). Further, >91% of *pkmt1*::tn bacteria were labeled

with both pUb and p62 in *Atg5*-deficient BMDMs (**Fig. 4A, B, C** and **D**), suggesting that ubiquitin-

tagged bacteria were not restricted when the autophagy cascade was prevented. In contrast, the

pkmt2::tn mutant did not have a major growth defect compared with WT bacteria and 50% of the

bacteria were labeled with p62, irrespective of host genotype (**Fig. 4A, B, C** and **D**), consistent

with less pronounced ubiquitylation phenotypes compared to *pkmt1*::tn bacteria. Collectively,

these data indicate that methylation is required for *R. parkeri* growth in macrophages by avoiding

autophagic targeting.

Our work reveals a detailed molecular mechanism that camouflages bacterial surface

proteins from host detection. In particular, we found that lysine methylation was essential for

blocking ubiquitylation, a first step in cell-autonomous immunity (*1-4*). This highlights an intricate

evolutionary arms race between pathogens and hosts and reveals a strategy that pathogens can

adapt to counteract host responses. The lysine methyltransferases PKMT1 and PKMT2 are

conserved between rickettsial species (**Fig. S9** and **Fig. S10**) and contain a core Rossmann fold

found in the broader superfamily of class I methyltransferases that exist in diverse organisms (*12,*

18). Thus, we propose that lysine methylation, and potentially other lysine modifications, could be

used by pathogens, symbionts, and perhaps even in eukaryotic organelles, to prevent undesirable

surface ubiquitylation and downstream consequences including elimination by autophagy. Further

study of microbial surface modifications will continue to enhance our understanding of the

pathogen-host interface and could ultimately lead to new therapeutic targets for treating human

diseases including those caused by infectious agents.

References and Notes

1. F. Randow, R. J. Youle, Self and nonself: how autophagy targets mitochondria and bacteria. *Cell Host Microbe* **15**, 403-411 (2014).
2. A. J. Perrin, X. Jiang, C. L. Birmingham, N. S. So, J. H. Brumell, Recognition of bacteria in the cytosol of Mammalian cells by the ubiquitin system. *Curr Biol* **14**, 806-811 (2004).
3. S. J. L. van Wijk *et al.*, Linear ubiquitination of cytosolic *Salmonella* Typhimurium activates NF-kappaB and restricts bacterial proliferation. *Nat Microbiol* **2**, 17066 (2017).
4. J. Noad *et al.*, LUBAC-synthesized linear ubiquitin chains restrict cytosol-invading bacteria by activating autophagy and NF-kappaB. *Nat Microbiol* **2**, 17063 (2017).
5. E. D. Case *et al.*, The *Francisella* O-antigen mediates survival in the macrophage cytosol via autophagy avoidance. *Cell Microbiol* **16**, 862-877 (2014).
6. J. Huang, J. H. Brumell, Bacteria-autophagy interplay: a battle for survival. *Nat Rev Microbiol* **12**, 101-114 (2014).
7. Y. T. Zheng *et al.*, The adaptor protein p62/SQSTM1 targets invading bacteria to the autophagy pathway. *J Immunol* **183**, 5909-5916 (2009).
8. T. L. Thurston, G. Ryzhakov, S. Bloor, N. von Muhlinen, F. Randow, The TBK1 adaptor and autophagy receptor NDP52 restricts the proliferation of ubiquitin-coated bacteria. *Nat Immunol* **10**, 1215-1221 (2009).
9. E. Fiskin, T. Bionda, I. Dikic, C. Behrends, Global analysis of host and bacterial ubiquitinome in response to *Salmonella* Typhimurium infection. *Mol Cell* **62**, 967-981 (2016).
10. M. Polajnar, M. S. Dietz, M. Heilemann, C. Behrends, Expanding the host cell ubiquitylation machinery targeting cytosolic *Salmonella*. *EMBO Rep* **18**, 1572-1585 (2017).
11. P. Engström *et al.*, Evasion of autophagy mediated by *Rickettsia* surface protein OmpB is critical for virulence. *Nat Microbiol* **4**, 2538-2551 (2019).
12. S. Lanouette, V. Mongeon, D. Figeys, J. F. Couture, The functional diversity of protein lysine methylation. *Mol Syst Biol* **10**, 724 (2014).
13. R. L. Lamason, N. M. Kafai, M. D. Welch, A streamlined method for transposon mutagenesis of *Rickettsia parkeri* yields numerous mutations that impact infection. *PLoS One* **13**, e0197012 (2018).
14. B. G. Spratt, A. B. Pardee, Penicillin-binding proteins and cell shape in *E. coli*. *Nature* **254**, 516-517 (1975).
15. H. K. Kim, R. Premaratna, D. M. Missiakas, O. Schneewind, *Rickettsia conorii* O antigen is the target of bactericidal Weil-Felix antibodies. *Proc Natl Acad Sci U S A* **116**, 19659-19664 (2019).
16. T. P. Burke *et al.*, Inflammasome-mediated antagonism of type I interferon enhances Rickettsia pathogenesis. *Nat Microbiol* **5**, 688-696 (2020).
17. D. C. H. Yang, A. H. Abeykoon, B. E. Choi, W. M. Ching, P. B. Chock, Outer membrane protein OmpB methylation may mediate bacterial virulence. *Trends Biochem Sci* **42**, 936-945 (2017).
18. A. H. Abeykoon *et al.*, Structural insights into substrate recognition and catalysis in outer membrane protein B (OmpB) by protein-lysine methyltransferases from *Rickettsia*. *J Biol Chem* **291**, 19962-19974 (2016).
19. A. Abeykoon *et al.*, Multimethylation of *Rickettsia* OmpB catalyzed by lysine methyltransferases. *J Biol Chem* **289**, 7691-7701 (2014).

20. A. H. Abeykoon *et al.*, Two protein lysine methyltransferases methylate outer membrane
protein B from *Rickettsia*. *J Bacteriol* **194**, 6410-6418 (2012).
21. N. F. Noriea, T. R. Clark, T. Hackstadt, Targeted knockout of the *Rickettsia rickettsii*
OmpA surface antigen does not diminish virulence in a mammalian model system. *mBio*
6, (2015).
22. R. L. Lamason *et al.*, *Rickettsia* Sca4 Reduces Vinculin-Mediated Intercellular Tension to
Promote Spread. *Cell* **167**, 670-683 e610 (2016).
23. U. Distler *et al.*, Drift time-specific collision energies enable deep-coverage data-
independent acquisition proteomics. *Nat Methods* **11**, 167-170 (2014).
24. R. S. Plumb *et al.*, UPLC/MS(E); a new approach for generating molecular fragment
information for biomarker structure elucidation. *Rapid Commun Mass Spectrom* **20**, 1989-
1994 (2006).
25. P. V. Shliaha, N. J. Bond, L. Gatto, K. S. Lilley, Effects of traveling wave ion mobility
separation on data independent acquisition in proteomics studies. *J Proteome Res* **12**, 2323-
2339 (2013).
26. S. Nahnsen, C. Bielow, K. Reinert, O. Kohlbacher, Tools for label-free peptide
quantification. *Mol Cell Proteomics* **12**, 549-556 (2013).
27. K. A. Neilson *et al.*, Less label, more free: approaches in label-free quantitative mass
spectrometry. *Proteomics* **11**, 535-553 (2011).
28. C. I. Carlström *et al.*, (Per)chlorate-reducing bacteria can utilize aerobic and anaerobic
pathways of aromatic degradation with (per)chlorate as an electron acceptor. *mBio* **6**,
(2015).
22. R. L. Lamason *et al.*, *Rickettsia* Sca4 Reduces Vinculin-Mediated Intercellular Tension to
Promote Spread. *Cell* **167**, 670-683 e610 (2016).
23. U. Distler *et al.*, Drift time-specific collision energies enable deep-coverage data-
independent acquisition proteomics. *Nat Methods* **11**, 167-170 (2014).
24. R. S. Plumb *et al.*, UPLC/MS(E); a new approach for generating molecular fragment
information for biomarker structure elucidation. *Rapid Commun Mass Spectrom* **20**, 1989-
1994 (2006).
25. P. V. Shliaha, N. J. Bond, L. Gatto, K. S. Lilley, Effects of traveling wave ion mobility
separation on data independent acquisition in proteomics studies. *J Proteome Res* **12**, 2323-
2339 (2013).
26. S. Nahnsen, C. Bielow, K. Reinert, O. Kohlbacher, Tools for label-free peptide
quantification. *Mol Cell Proteomics* **12**, 549-556 (2013).
27. K. A. Neilson *et al.*, Less label, more free: approaches in label-free quantitative mass
spectrometry. *Proteomics* **11**, 535-553 (2011).
28. C. I. Carlström *et al.*, (Per)chlorate-reducing bacteria can utilize aerobic and anaerobic
pathways of aromatic degradation with (per)chlorate as an electron acceptor. *mBio* **6**,
e02287-14 (2015).

Acknowledgements

We thank Prof. Dan Portnoy (UC Berkeley) and Dr. Lilliana Radoshevich (University of Iowa) for providing comments on this manuscript, and Neil Fischer (UC Berkeley) for critically reading the manuscript. We also thank Dr. Hwan Kim (Stony Brook University) for kindly providing the O-antigen antibody and for fruitful discussions. We are also grateful for the *Atg5^{flox/flox}* and *Atg5^{-/-}* BMDMs provided by Dr. G. Golovkine (UC Berkeley), and Profs. Jeffery S. Cox and Michael Rape (both UC Berkeley) for fruitful discussions. **Funding:** P.E. was supported by a postdoctoral fellowship from the Sweden-America Foundation. M.D.W. was supported by NIH/NIAID grants R01 AI109044 and R21 AI138550. A mass spectrometer used in this study was purchased with support from NIH grant 1S10 OD020062-01. **Author contributions:** P.E. conceived the study with the assistance of M.D.W. P.E. performed laboratory work and analysis, except for mass spectrometry, which was conducted together with A.T.I, and animal experiments, which were conducted by T.P.B. P.E. drafted the initial manuscript, and A.T.I, T.P.B, and M.W.D provided editorial feedback. **Data and materials availability:** All data used in the analysis are provided in the main text or supplementary materials. Materials are available upon request.

256
Fig. 1. The O-antigen and lysine methylation are virulence factors that protect *R. parkeri*
from ubiquitylation. (**A**) Pools of *R. parkeri* (green) mutants screened for increased pUb (red)
via immunofluorescence microscopy. DNA, blue. *ompB* mutant (*11*), positive control; WT,

negative control. Circle, pUb-positive bacteria; arrows pUb-negative bacteria. Scale bar, 20 μm.
(**B**) Infected cells at 72 hours post-infection (h.p.i.) stained as in **A**. Scale bar, 5 μm ($n = 4$). (**C**)
Biological function of the genes identified. (**D**) Percentage of bacteria co-localized with pUb at 72
h.p.i. (Data are the mean ± s.e.m.; $n = 4$ for WT and mutant bacteria; *mrdA*::tn; $n = 2$). Statistical
comparisons between WT and mutants were performed using a one-way ANOVA with Dunnett's
post-hoc test; **** $P < 0.0001$. (**E**) pUb signal per bacteria. (Lines indicate the means; $n = 3$ fields
of vision). Statistical comparisons were performed using a Kruskal-Wallis test with Dunn's post-
hoc test; **** $P < 0.0001$. (**F**) Survival of *Ifnar*$^{-/-}$*Ifngr*$^{-/-}$ mice intravenously infected with 5×10^6
WT or mutant bacteria ($n = 5$ mice, WT; $n = 6$ mice, *pkmt1*::tn and *pkmt2*::tn). Statistical
comparison between WT and mutants was performed using a two-way ANOVA; ** $P < 0.01$.

Fig. 2. **Lysine methylation protects OMPs from ubiquitylation.** (**A**) Micrographs of infected Vero cells expressing 6xHis-ubiquitin stained with anti-His antibody (red) and Hoechst (blue, bacterial and host DNA), at 28 h.p.i. Scale bar, 5 μm (representative of $n = 2$). (**B**) Western blot of His-Ub input and pull-down samples from infected control and 6xHis-ubiquitin expressing cells, probed for OmpB, OmpA, and pUb (representative of $n = 3$). (**C**) pUb-enriched (TUBE-1, pan specific) samples from purified bacteria probed for OmpB, OmpA, and OmpW (OmpB and OmpA of endogenous molecular weight represent non-specific binding to TUBE-1 beads) (representative of $n = 3$). Asterisks indicate OmpB and OmpA that exhibit increased molecular weight, indicating ubiquitylation. Arrows indicate OmpB and OmpA of endogenous molecular weight.

Fig. 3. Methylation camouflages lysines from ubiquitylation. (A) Percentage lysine methylation
abundance of total among the abundantly detected proteins in WT *R. parkeri* determined by LC-
MS (data are combined from $n = 2$). **(B)** Methylated lysines (K) are indicated with residue number
in each OMP, and lysines in *R. prowazekii* OmpB known to be methylated in green (*17*). K78,
K131, K149, K312, and K547 in OmpB were unmethylated ($n = 5$, using data-independent and
data-dependent acquisition modes). **(C)** Percentage of total abundances of the lysines methylated
in **B**, that is unmethylated, mono-, di-, or tri-methylated (data are the mean of $n = 2$). **(D)**
Percentage of individual lysines in OmpB that are unmethylated, mono-, di- or tri-methylated.
Only residues repeatedly detected in all strains were analyzed (data are the mean of $n = 2$). **(E)**
Heat map representation of OmpB residues from *pkmt1*::tn or *pkmt2*::tn that have similar levels of
K-diGly peptides (white boxes) compared to WT, or with a 5-fold, or more, increase of K-diGly
peptides (cyan) after pUb-enrichments ($n = 2$). **(F)** Abundances of ubiquitylated (diGly),
unmethylated, mono-, di-, or tri-methylated OmpB K623 (data are the mean of $n = 2$). Each
experiment (n) was performed in technical triplicate.

Fig. 4. Methylation prevents ATG5-dependent *R. parkeri* killing in macrophages. (**A**)
Micrographs of infected control (*Atg5^{flox/flox}*) and *Atg5*-deficient (*Atg5^{-/-}*) BMDMs at 48 h.p.i.,
stained for DNA (blue), pUb (red), and p62 (green). Arrow indicate a bacterium positive for both
pUb and p62. Scale bars, 5 μm (*n* = 3). (**B**) Bacterial growth curves in control and *Atg5^{-/-}* BMDMs,
as measured by genomic equivalents using qPCR (*n* = 3). (**C**) Quantification of the mean number
of bacteria per cell (*n* = 3). (**D**) Quantification of the percentage of bacteria with p62 and pUb (*n*
= 3). nd, not determined. Statistical comparisons in **B** and **C** were performed using a Brown-
Forsyth and Welch ANOVA with Dunnett's post-hoc test; *, *P* < 0.05. Presented data are the mean
± s.e.m.

Supplementary Figures

Fig. S1. Polyubiquitylated mutant *R. parkeri* strains are positive for OmpA and OmpB. (A) Micrographs of Vero cells infected with the indicated strains stained for OmpA (green, 13-3 antibody), OmpB (red, OmpB-antibody), and DNA (blue, Hoechst) at 72 h.p.i. (representative of $n = 3$). Scale bar 5 μm. (**B**) Quantification of OmpB signal per bacteria. (Lines indicate the means; $n = 3$ fields of vision; ≥ 50 bacteria per field of vision were analyzed). Statistical comparisons were performed using a Kruskal-Wallis test with Dunn's post-hoc test; **** $P < 0.0001$ between indicated strains. Dashed line indicates that the majority of mutant bacterial populations, except *pkmt1*::tn bacteria, have OmpB levels comparable to WT bacteria. (**C**) Quantification of OmpA signal per bacteria as in **B**. (**D**) Western blot of 5×10^6 purified bacteria probed for OmpB, pUb, OmpW and EF-Tu (bacterial loading controls) (representative of $n = 2$).

Fig. S2. The *wecA*::tn and *rmlD*::tn bacteria lack the O-antigen. (**A**) Micrographs of Vero cells
infected with the indicated strains stained for OmpA (green, 13-3 antibody), the O-antigen (red,
O-antigen antibody(*15*)) and DNA (blue, Hoechst) at 72 h.p.i. (representative of n = 3). Scale bars,
5 μm. (**B**) Percentage of bacteria positive for the O-antigen at 72 h.p.i. n = 2; ≥ 90 bacteria were
counted in each infection (data are the mean ± s.e.m.; n = 3). (**C**) Western blot of 5×10^6 purified
WT and mutant bacteria probed for the O-antigen. OmpW and EF-Tu were used as bacterial
loading controls (representative of n = 3).

Fig. S3. PKMT1 plays a more significant role in causing disease *in vivo* compared to PKMT2.
(**A**) Weight changes of *Ifnar⁻/⁻Ifngr⁻/⁻* mice intravenous infected with 5×10^6 WT, *pkmt1*::tn, or
pkmt2::tn bacteria (data are the mean ± s.e.m. *n* = 5, WT; *n* = 6, *pkmt1*::tn and *pkmt2*::tn, combined
from two independent experiments). A two-way ANOVA from 0 to 25 days post-infection (d.p.i.)
was used to statistically compare the weight changes between the *pkmt1*::tn and *pkmt2*::tn mutants.
**** $P < 0.0001$.

Fig. S4. Lysine-methylome reveals that _R. parkeri_ OMP-porin, BamA, and the released factor

Sca4 are methylated. Methylated lysines are indicated with residue number, and unmethylated

residues without, determined by LC-MS as in **Fig. 3B** (data is combined from five independent

experiments). See also **Table S2**.

338
339 **Fig. S5. PKMT1 primarily modifies bacterial OMPs.** Percentage of total abundances of the
340 lysines methylated in WT, that is unmethylated, mono-, di-, or tri-methylated in respective strain.
341 Proteins with ≥3 lysines methylated detected in independent experiments are shown (mean of n =
342 2, performed in technical triplicates). OmpB, OmpA and Sca2 data are the same as in **Fig. 3C**. See
343 also **Table S2**.

Fig. S6. Dimethylation is a common methylation state among the abundant *R. parkeri* proteins. Percentage of total lysines abundances detected to as mono-, di-, or tri-methylated in WT bacteria of the proteins indicated in **Table S2** (mean of *n* = 2, performed in technical triplicates).

Fig. S7. OmpB K623 and K634 are frequently ubiquitylated when PKMT1-mediated methylation is reduced. (**A**) OmpB peptide abundance values after pUb-enrichments from respective strain, as determined by LC-MS (data are mean of n = 2, performed in triplicates). (**B**) Abundance values of lysines in OmpB that are ubiquitylated (diGly), unmethylated, mono-, di-, or tri-methylated, in respective strain after pUb-enrichments as in **Fig 2C.** Only residues detected in independent experiments are shown (data are the mean of n = 2, performed in technical triplicates). K623 data in **B** are the same as in **Fig. 3F.**

Fig. S8. OmpB and lysine methylation block recruitment of autophagy receptors to *R.
***parkeri*.** (**A**) Micrographs of Vero cells infected with the indicated strains: stained for bacterial and
host DNA (blue, Hoechst), p62 (green, p62-antibody), and NDP52 (red, NDP52-antibody) at 72
h.p.i ($n = 4$ for p62 staining; $n = 2$ for NDP52 staining). (**B**) Percentage of bacteria that show rim-
like surface localization of p62 at 72 h.p.i. (Data are the mean ± s.e.m.; $n = 3$; ≥ 142 bacteria were
counted for each infection). Statistical comparisons between WT and *ompB^{STOP}*::tn, *pkmt1*::tn,
pkmt2::tn were performed using a one-way ANOVA with Tukey's post-hoc test; **** $P < 0.0001$.
(**C**) Quantification of p62 signal per bacteria from a representative experiment. Lines indicate the
means. ($n = 3$ fields of vision; ≥ 50 bacteria per field of vision were analyzed). Statistical
comparisons were performed using a Kruskal-Wallis test with Dunn's post-hoc test; **** $P <$
0.0001 between indicated strains. (**D**) Quantification of NDP52 signal per bacteria as in **C**.

```
Rickettsia parkeri                                     1   MSPKVTNSSSTPNGHDKMAKKTHSAQSVVN--GAVSDHNTYDEIPYESYPYALTNPYHLSTLATLFGVNAPEVENAKILE  78
Rickettsia rickettsii                                  1   MSPKATNSSSTPNGHDKMAKKTHSAQSVVN--GAVSDHNTYDEIPYESYPYALTNPYHLSTLATLFGVNAPEVENAKILE  78
Rickettsia conorii                                     1   MSPKATNSSTPNGHDKMAKKTHSAQSVVN--GAVLDHNTYDEIPYESYPYALTHPYHLSTLATLFGVNAPEVENAKILE   78
Rickettsia typhi                                       1   MSLKSST----TNDHDK-TTKINSIQSLVNctDTVADHNTYDEIPYESYPYAITNPYHLSTLATLFGINAPEVENSKILE  75
Rickettsia prowazeki                                   1   MSLKSTTSSLTTNNHDK-T--INSVQSLVNgtGTVADHNPYDEVPYESYPYAITNPYHLSTLATLFGINAPEVENSKILE  77
Rickettsia bellii                                      1   MSAKASNSSNLPNGHDKTTKEKHNTQPVIN--GAIK-HNTYDEVPYESYPYSFTNPFHLSTLATLFGVDAPNVETAKILE  77
Rickettsia endosymbiont of Proechinophthirus fluctus   1   MSPKATNSSSTPNSHDKMAKKTHSVQSVVN--SAVSDHNTYDEIPYESYPYAFTHPYHLSTLATLFGVNAPEVENAKILE  78

Rickettsia parkeri                                     79  LGCAAGGNLIPHAVLYPKAYFVGVDLSKVQIDEANKNVKALGLKNIEFHHCSITDINDSFGKFDYIICHGVISWVPKTVR  158
Rickettsia rickettsii                                  79  LGCAAGGNLIPHAVLYPKAYFVGVDLSKVQIDEANKNVKALGLKNIEFHHCSITDINDSFGKFDYIICHGVISWVPKTVR  158
Rickettsia conorii                                     79  LGCAAGGNLIPHAVLYPKAYFVGVDLSKVQIDEANKNVKALGLKNIEFHHCSITDINDSFGKFDYIICHGVISWVPKTVR  158
Rickettsia typhi                                       76  LGCAAGGNLIPHAVLYPKAHFVGVDLSKVQIDEANKNVRALGLKNIEFHHCSITDINDSFGKFDYIICHGVISWVPKIVR  155
Rickettsia prowazeki                                   78  LGCAAGGNLIPHAVLYPNAHFVGVDLSKVQIDEANKNVRALGLKNIEFHHCSITDIDDSFGKFDYIICHGVISWVPKIVR  157
Rickettsia bellii                                      78  LGCAAGGNLIPHAVLYPKAHFVGIDLSKVQIDEANKTVKELGLKNIEFHHCSILDIDDSLNKFDYIICHGVMSWVSKNVR  157
Rickettsia endosymbiont of Proechinophthirus fluctus   79  LGCAAGGNLIPHAVLYPKAYFVGVDLSKVQIDEANKNVKALGLKNIEFHHCSITDINDSFGKFDYIICHGVISWVPKTVR  158

Rickettsia parkeri                                     159 DKIFEVCNKNLSPNGIAYISYNTLPGWNMVRTIRDMMMYHSSSFANVRDRIAQSRLLLEFVKDSLENSKTPYAEALKTEA  238
Rickettsia rickettsii                                  159 DKIFEVCNKNLSPNGIAYISYNTLPGWNMVRTIRDMMMYHSSSFANVRDKIAQSRLLLEFVKDSLENSKTPYAEALKTEA  238
Rickettsia conorii                                     159 DKIFEVCNKNLSPNGIAYISYNTLPGWNMVRTIRDMMMYHSSSFANVRDRIAQSRLLLEFVKDSLENSKTPYAEALKTEA  238
Rickettsia typhi                                       156 DKIFKVCNSNLSTNGIAYISYNTLPGWNMVRTIRDMMLYHSSSFTNVRDRIAQSRLLLEFVKDSLENSKTPYAEVLKTEA  235
Rickettsia prowazeki                                   158 DKIFKVCNRNLSTNGIAYISYNTLPGWNMVRTIRDMMLYHSSSFTNIRDRIAQSRLLLEFVKDSLEHSKTPYAEVLKTEA  237
Rickettsia bellii                                      158 DKIFDVCNKNLSKNGIAYISYNTLPGWNMVRTVRDMMLYHSSSFVNVRDKIAQSRLLLDFVKDSLENSKTPYAEVLKTEA  237
Rickettsia endosymbiont of Proechinophthirus fluctus   159 DKIFEVCNKNLSPNGIAYISYNTLPGWNMVRTIRDMMMYHSSSFANVRDRIAQSRLLLEFVKDSLENSKTPYAEALKTEA  238

Rickettsia parkeri                                     239 GLLAKQTDHYLRHDHLEEENAQFYFHEFMNEARKHNLQYLADCNLSTMYLGNMPPKVVEQLKAVNDIVRTEQYMDFITNR  318
Rickettsia rickettsii                                  239 GLLAKQTDHYLRHDHLEEENAQFYFHEFMNEARKHNLQYLADCNLSTMYLGNMPPKVVEQLKAVNDIVRTEQYMDFITNR  318
Rickettsia conorii                                     239 GLLAKQTDHYLRHDHLEEENAQFYFHEFMNEARKHNLQYLADCNLSTMYLGNMPPKVVEQLKAVNDIVRTEQYMDFITNR  318
Rickettsia typhi                                       236 GLLAKQTDHYLRHDHLEEENAQFYFHEFMNEARKYNLQYLADCNISTMYLGNMPPKVVEQLKAVNDIVRTEQYMDFITNR  315
Rickettsia prowazeki                                   238 GLLAKQTDHYLRHDHLEEENAQFYFHEFMNEARKHNLQYLADCNLSTMYLGNMPPKVVEQLKAVNDIVRTEQYMDFITNR  317
Rickettsia bellii                                      238 GLLSKQTDHYLRHDHLEEENAQFYFHEFMDEARKHNLQYLADCNLSTMYLGNMPQKVVEQLKVVNDIVRTEQYMDFITNR  317
Rickettsia endosymbiont of Proechinophthirus fluctus   239 GLLAKQTDHYLRHDHLEEENAQFYFHEFMNEARKHNLQYLADCNLSTMYLGNMPPKVVEQLKAVNDIVRTEQYMDFITNR  318

Rickettsia parkeri                                     319 RFRTTLLCHSDVKINRNINNDDITKFNIIFNIVPEKPLKEVDLNNASENLKFFLNGNQDSNLTTSSPYMKAILYTFSENL  398
Rickettsia rickettsii                                  319 RFRTTLLCHSDVKINRNINNDDITKFNIIFNIVPEKPLKEVDLNNASENLKFFLNGNQDSNLTTSSPYMKAILYTFSENL  398
Rickettsia conorii                                     319 RFRTTLLCHSDVKINRNINNDDITKFNIIFNIVPEKPLKEVDLNNASENLKFFLNGNQDSNLTTSSPYMKAILYTFSENL  398
Rickettsia typhi                                       316 RFRTTLLCHNDLKINRNINNEDIKKFNIIFNVVPEKPLQEVDLDNAAENLQFFLNGNKECNLSTTSPYMKAILYTFSENL  395
Rickettsia prowazeki                                   318 RFRTTLLCHNDLKINRNINNDDIKKFNIIFNVIPEKPLKEVDLNNATENLQFFLNGNKESNLSTTSPYMKAILYTFSENL  397
Rickettsia bellii                                      318 RFRSTLLCHNDVKINRNINNEDIMKFNMIFNVVPEKSLKEVDLNNSSESLAFFLNGNKDSNLTTASPYMKAILYTFSEHL  397
Rickettsia endosymbiont of Proechinophthirus fluctus   319 RFRTTLLCHSDVKINRNINNDDITKFNIIFNIVPEKPLKEVDLNNASENLKFFLNGNQDSNLTTSSPYMKAILYTFSENL  398

Rickettsia parkeri                                     399 NNPLSFEKITTEANKKLHNTKLNEIKAEFLNNAMKLVLQGYISITNQKHRNNPELDKPKTTKMVIHQATHTPSMWVTNLK  478
Rickettsia rickettsii                                  399 NNPLSFEKITTEANKKLHNTKLNEIKAELLNNAMKLVLQGYISITNQKHRNNPELDKPKTTKMVIHQATHTPSMWVTNLK  478
Rickettsia conorii                                     399 NNPLSFEKITTEANKKLHNTKLNEIKAELLNNAMKLVLQGYISITNQKHRNNPELDKPKTTKMVIHQATHTPSMWVTNLK  478
Rickettsia typhi                                       396 NNPLSFKQVTSEANKKLNNNKLNEIKNELLNNAMKLVLQGYISITNQKHRNKPVLDKPKTTQMVLYQAKHTPSMWVTNLK  475
Rickettsia prowazeki                                   398 NNPLSFKQVTSEANKKLNNTKLNEIKNELLNNAMKLVLQGYISITNQKHRSKPVLDKPKTTQMVIYQAKYTPSMWVTNLK  477
Rickettsia bellii                                      398 NNPLSFEKITEANKKLHGTKLNEIKAELLENAMKLVLQGYINITTQKHRETPELNKPKTTKLVIHQAIHTPYMWVTNLK   477
Rickettsia endosymbiont of Proechinophthirus fluctus   399 NNPLSFEKITTEANKKLHNTKLNEIKAELLNNAMKLVLQGYISITNQKHRNNPELDKPKTTKMVIHQATHTPSMWVTNLK  478

Rickettsia parkeri                                     479 HEPIGVNFFEKFALRYMDGKHDKKAIIEAVLGHVEKGELTLSKEGQKVENKEEIRKELESLFIPMIKKFSSNALLV  554
Rickettsia rickettsii                                  479 HEPIGVNFFEKFALRYMDGKHDKKAIIEAVLGHVEKGELTLSKEGQKVENKEEIRKELESLFIPMIKKFSSNALLV  554
Rickettsia conorii                                     479 HEPIGVNFFEKFALRYMDGKHDKKAIIEAVLGHVEKGELTLSKEGQKVENKEEIRKELESLFIPMIKKFSSNALLV  554
Rickettsia typhi                                       476 HEPIGVNFFEKFALRYMDGKNDKKAIIEAILGHVEKGELTLSKEGQKIENKEEIRKELESLFTPMIEKFASNALLV  551
Rickettsia prowazeki                                   478 HEPIGVNFFEKFALRYMDGRNDKKAIIEAILGHVEKGELTLSREGQKIENKEEIRKELESLFTPMIEKFCSNALLV  553
Rickettsia bellii                                      478 HEPVGVNFFEKLAIRYMDGKHDKKAIIDAVMGHVVKGEINLSKDGQKIEDQEVIRKQLEVLFMPMIDKFAANALLV  553
Rickettsia endosymbiont of Proechinophthirus fluctus   479 HEPIGVNFFEKFALRYMDGKHDKKAIIEAVLGHVEKGELTLSKDGQKVENKEEIRKELESLFIPMIKKFSSNALLV  554
```

Fig. S9. The PKMT1 enzyme is highly conserved between diverse rickettsial species. Amino acid sequence alignment of PKMT1 from *R. parkeri* (WP_014411082.1), *R. rickettsii* (WP_012262600.1), *R. conorii* (WP_016926592.1), *R. typhi* (WP_011191207.1), *R. prowazekii* (WP_004596928.1), *R. bellii* (WP_045799810.1), and a *Rickettsia* endosymbiont (WP_062811822.1), using COBALT. Amino acids indicated in red are identical; blue, variation between species; grey, one or more of the analyzed proteins are lacking this residue(s).

Fig. S10. The PKMT2 enzyme is highly conserved between virulent rickettsial species but absent from *R. bellii* and the *Rickettsia endosymbiont Proechinophthirus fluctus*. Amino acid sequence alignment of PKMT2 from *R. parkeri* (WP_014410272.1), *R. rickettsii* (WP_012150259.1), *R. conorii* (WP_016925880.1), *R. typhi* (WP_011190574.1), and *R. prowazekii* (WP_004596662.1) using COBALT. PKMT1 of *R. bellii* and the *Rickettsia endosymbiont Proechinophthirus fluctus* showed 51% identity to PKMT2 of *R. parkeri* using single alignment BLAST at NCBI; however, no PKMT2 variants could be found in these organisms, and therefore they were excluded from this analysis. Amino acids indicated in red are identical; blue, variation between species.

386
Fig. S11. Strain validation of clonality and insertion site using primers for flanking chromosomal
regions ($n = 1$).

Materials and Methods

Cell lines and primary mouse macrophages

Vero cells were purchased from the UC Berkeley Cell Culture Facility and the identity was

repeatedly confirmed by mass-spectrometry analysis. Cells were grown at 37 °C and 5% CO_2 in

DMEM (Gibco, cat. no. 11965) with high glucose (4.5 g l^{-1}) and 2% heat-inactivated (30 min, 56

°C, in a water-bath) fetal bovine serum (Gemcell). Vero cells were confirmed to be mycoplasma

negative by DAPI staining and fluorescence microscopy screening at the UC Berkeley Cell Culture

Facility.

BMDMs generated from the femurs of mutant $Atg5^{flox/flox}$ and matched $Atg5^{-/-}$ C57BL/6

mice were a kind gift from the laboratory of Jeffery S. Cox (UC Berkeley), and they were prepared

as previously described (*11*) although in the absence of antibiotics. Genotypes were confirmed by

PCR and Sanger sequencing at the UC Berkeley DNA Sequencing Facility, as previously

described (*11*).

***Rickettsia parkeri* strain generation and validation**

WT *R. parkeri* strain Portsmouth (NCBI accession no. NC_017044.1; originally a gift from

C. Paddock, Center for Disease Control and Prevention) and bacterial stocks of $ompB^{STOP}$::tn (the

genome sequences of these bacterial strains are available at the Sequence Read Archive as

accession no. SRP154218 (WT, SRX4401164; $ompB^{STOP}$::tn, SRX4401167)), *pkmt1*::tn,

pkmt2::tn, *wecA*::tn and *rlmD*::tn mutants were propagated and purified every ~6-10 months as

described below. Side-by-side experimental comparisons were made between stocks prepared at

similar times.

R. parkeri pkmt1::tn, *pkmt2*::tn, *wecA*::tn, and 114 other transposon insertion mutant

strains, screened for pUb, were previously isolated in a screen for small plaque mutants (*13, 22*).

The *ompB*^{STOP}::tn was previously isolated in a suppressor screen and lacked expression of OmpB

(*11*). The *rmlD*::tn, and 132 other transposon insertion mutant strains, screened for pUb, were

isolated in an independent screen in which mutants were isolated without regard for plaque size

(**Table S1**). The genomic locations of transposon insertion sites for all mutants were determined

by semi-random nested PCR. To verify the insertions and clonality, we used PCR reactions that

amplified the transposon insertion site using primers for flanking chromosomal regions:

5'GCTCACTAGATAGCACTCG'3 and 5'GCTCGATTTATCTCACTTTATG'3 for *rlmD*::tn,

5'CGTTTAATAGTCCAGTTAATTTGT'3 and 5'CCGTCTATACCGTCCATAAAAT'3 for

wecA::tn, 5'GCATCGAAATAACCCTGAG'3 and 5'GCAAACTTCTCAAAGAAATTAACG'3

for *pkmt1*::tn, 5'GCTAAGAAATCTTCTAATTTGATATTTTAC'3 and

5'CGAAAATTTACCTGAGCCTT'3 for *pkmt2*::tn, 5'CGACACATAATAGCACAAACTAC'3

and 5'GCGGAGGCGGTAGTAAAG'3 for *mrdA*::tn (**Fig. S11**).

Screening for pUb-positive strains

To prepare the mutant library for screening, passage 1 (P1) transposon insertion mutants

were amplified one time in Vero cells using 24-well cell culture plates. At 5-12 d.p.i, when 50-

70% of the infected cells appeared to be rounded up (as a sign of infection) by visual inspection

using a light microscope, cell culture media were completely removed, and cells were subsequently

lysed in 500 µL cold sterile water for 2-3 minutes (min). Next, 500 µL of 2x cold sterile brain-

heart-infusion (BHI) broth (BD Difco, 237500) was added to the lysed cells, resuspended, and P2

bacteria were transferred to cryogenic storage vials and frozen at -80 °C.

To screen for pUb-positive strains, 10-40 µL of each of five to seven P2 mutant bacterial

strains were diluted in 1 mL of room temperature (RT) cell culture media supplemented with 2%

FBS. Subsequently, the pooled bacterial suspension was centrifuged at 250g for 4 min at RT onto

confluent Vero cells grown on coverslips in 24-well plates. Cells were then incubated at 33 °C and

fixed at 50-55 h.p.i. with pre-warmed 4% paraformaldehyde (PFA) (Ted Pella Inc., 18505) for 10

min at RT. If cells were over-infected (i.e., individual infection foci had grown together) as

determined by immunofluorescence microscopy, infections of that specific pool were repeated

using reduced volumes of P2 bacteria. Next, fixed cells were permeabilized with 0.2% Triton-X

for 5 min and then stained with the anti-*Rickettsia* I7205 antibody (1:500 dilution; gift from Ted

Hackstadt) and anti-polyubiquitin FK1 antibody (Enzo Life Sciences, BML-PW8805-0500; 1:250

dilution), followed by Alexa 488 anti-rabbit antibody (Invitrogen, A11008; 1:500 dilution) or goat

anti-mouse Alexa-568 (Invitrogen, A11004). Whole coverslips were manually inspected on a

Nikon Ti Eclipse microscope with 60x (1.4 numerical aperture) Plan Apo objective. The initial

screen revealed that five out of 39 mutant pools contained pUb-positive areas.

In a secondary screen, individual strains from the pUb-positive pools were used to infect

Vero cells, as stated above. Infected cells were also fixed and stained as above except that a post-

fixation step using 100% methanol for 5 min was included and an OmpB antibody (*11*) and

Hoechst (Sigma, B2261, 1:2500 dilution) was used instead of the anti-*Rickettsia* I7205 antibody.

Samples were inspected as above and strains were scored as follows: **1**) pUb-negative (49 strains),

2) a few infection foci were pUb-positive (2 strains: Sp mutant 24, insertion at bp position 753916;

Sp mutant 94, insertion at bp position 774831), **3**) bacteria in the center of foci were pUb-positive

but not on the edges (2 strains: Sp mutant 43, insertion at bp position 651602-651604; Sp mutant

45, insertion at bp position 751156), **4**) almost all bacteria in all foci were pUb-positive (4 strains:

pkmt1::tn, insertion at bp position 1161553 (gene *MC1_RS06185*); *pkmt2*::tn, insertion at bp

position 34100 (*MC1_RS00180*); *wecA*::tn, insertion at bp position 1223170 (*MC1_RS06510*); and

rmlD::tn, insertion at bp position 455753 (*MC1_RS02345*).

***Rickettsia* purification**

R. parkeri strains were propagated as described previously (*11*). "Purified bacteria" were

from five T175 flasks of Vero cells that after 5-8 days of infection (normally ~75% infected as

observed by light microscopy) were harvested in the media using a cell scraper. Bacteria were tehn

centrifuged 12000g for 15 min at 4 °C in pre-chilled tubes. Pellets were resuspended in cold K-36

buffer (0.05 M KH_2PO_4, 0.05 M K_2HPO_4, pH 7, 100 mM KCl and 15 mM NaCl) and a pre-chilled

dounce-homogenizer (tight fit) were used for 60 strokes to release bacteria from host cells. The

homogenate was then centrifuged at 200g for 5 min at 4 °C to remove cellular debris. The

supernatant was overlaid onto cold 30% v/v MD-76R (Mallinckrodt Inc., 1317-07) diluted in K-

36, and centrifuged at 58300g for 20 min at 4 °C in an SW-28 swinging-bucket rotor. The bacterial

pellet was resuspended in cold 1x BHI broth (0.5 mL BHI per infected T175 flask), and after letting

DNA precipitates sediment to the bottom of the tubes, bacterial suspensions were collected,

aliquoted and frozen at -80 °C.

"Gradient-purified bacteria" were from ten T175 flasks of Vero cells, purified as above

with the addition of a 40/44/54% v/v MD-76R (diluted in K-36 buffer) gradient step centrifuged

at 58300g for 25 min at 4 °C using the SW-28 swinging bucket rotor. The bacteria were then

collected from the 44-54% interface, diluted in K-36 buffer, and pelleted by centrifugation at

12000g for 15 min at 4 °C. The pellet was resuspended in cold 1x BHI broth and subsequently

aliquoted and frozen at -80 °C.

OmpW and EF-Tu antibody production

The sequence encoding amino acids 22-224 of outer membrane protein W (OmpW; WP_014411122.1) (a protein that lacks the signal peptide), or full-length Elongation factor Tu (EF-Tu; WP_004997779.1), were amplified by PCR from *R. parkeri* genomic DNA, and subsequently cloned into plasmid pETM1, which encodes N-terminal 6xHis and maltose-binding proteins (MBP) tags. From the resulting plasmids, fusion proteins were expressed in *E. coli* strain BL21 codon plus RIL-Cam^r (DE3) (QB3 Macrolab, UC Berkeley) by induction with 1 mM isopropyl-β-D-thio-galactoside (IPTG) for 2-2.5 hours at 37 °C. Bacterial pellets were resuspended in lysis buffer (50 mM NaH$_2$PO$_4$, pH 8.0, 300 mM NaCl, 1mM EDTA, and 1 mM dithiothreitol (DTT)) and stored at -80 °C. For protein purification, bacteria were thawed, lysozyme was added to 1 mg/mL (Sigma, L4919), and lysis was carried out by sonication. Lysates containing 6xHis-MBP-OmpW and 6xHis-MBP-EF-Tu were incubated on amylose resin (New England Biolabs, E8021L) (Qiagen, 1018244) and bound proteins were eluted in lysis buffer lacking EDTA and DTT but containing 10 mM maltose. Fractions were analyzed by SDS-PAGE and those with the highest concentrations of fusion proteins were pooled to generate rabbit antibodies against OmpW and EF-Tu. 1.2 mg of purified 6xHis-MBP-OmpW and 6xHis-MBP-EF-Tu proteins were sent to Pocono Rabbit Farm and Laboratory (Canadensis, PA), and immunization was carried out according to their 91-d protocol.

Western blotting

To determine the levels of bacterial and host proteins in purified bacterial samples, 30%-purified bacterial samples were boiled in 1x SDS loading buffer (150 mM Tris pH 6.8, 6% SDS,

0.3% bromophenol blue, 30% glycerol, 15% β-mercaptoethanol) for 10 min, then $5x10^6$ PFUs

were resolved on an 8-12% SDS-PAGE gel and transferred to an Immobilon-FL polyvinylidene

difluoride membrane (Millipore, IPEL00010). Membranes were probed for 30 min at room

temperature or 4°C overnight with antibodies as follows: affinity-purified rabbit anti-OmpB

antibody (*11*) diluted 1:200-30000 in TBS-T (20 mM Tris, 150 mM NaCl, pH 8.0, 0.05% Tween

20 (Sigma, P9416)) plus 5% dry milk (Apex, 20-241); mouse monoclonal anti-OmpA 13-3

antibody diluted 1:10000-50000 in TBS-T plus 5% dry milk; rabbit anti-OmpW serum diluted

1:8000 in TBS-T plus 5% dry milk; mouse monoclonal FK1 anti-polyubiquitin antibody diluted

1:2500 in TBS-T plus 2% BSA; rabbit anti-EF-Tu serum diluted 1:15000 in TBS-T plus 5% dry

milk; or rabbit anti-O-antigen serum 1:5000 in TBS-T plus 5% dry milk. Secondary antibodies

were: mouse anti-rabbit horseradish peroxidase (HRP) (Santa Cruz Biotechnology, sc-2357), or

goat anti-mouse HRP (Santa Cruz Biotechnology, sc-2005), all diluted 1:1000-2500 in TBS-T plus

5% dry milk. Secondary antibodies were detected with ECL Western Blotting Detection Reagents

(GE, Healthcare, RPN2106) for 1 min at room temperature, and developed using Biomax Light

Film (Carestream, 178-8207).

Immunofluorescence microscopy

R. parkeri infections were carried out in 24-well plates with sterile circle 12-mm coverslips

(Thermo Fisher Scientific, 12-545-80). To initiate infection, 30%-purified bacteria were diluted in

cell culture media at room temperature to a MOI of 0.01 for Vero cells, and a MOI of 0.1 for

BMDMs. Bacteria were centrifuged onto cells at 300*g* for 5 min at room temperature and

subsequently incubated at 33 °C. Next, infected cells were fixed for 10 min at room temperature

in pre-warmed (37 °C) 4% PFA diluted in PBS, pH 7.4, then washed 3 times with PBS. Primary

antibodies were the following: for staining with the guinea pig polyclonal anti-p62 antibody (Fitzgerald, 20R-PP001; 1:500 dilution), mouse polyclonal anti-NDP52 antibody (Novus Biologicals, H00010241-B01P; 1:100 dilution), a rabbit anti-*Rickettsia* I7205 antibody (1:500 dilution; gift from Ted Hackstadt), or anti-polyubiquitin FK1 antibody (1:250 dilution), cells were permeabilized with 0.5% Triton-X100 for 5 min prior to staining. For staining with mouse monoclonal anti-OmpA 13-3 antibody (1:5000 dilution), anti-OmpB antibody (*11*) (1:1000 dilution), or rabbit anti-O-antigen serum (*15*) (1:500 dilution), infected cells were post-fixed in methanol for 5 min at RT (no Triton-X). Cells were then washed 3 times with PBS and incubated with the primary antibodies for 30 min at RT. To detect the primary antibodies, secondary goat anti-rabbit Alexa-568 (Invitrogen, A11036), goat anti-mouse Alexa-568 (Invitrogen, A11004), or goat anti-guinea pig Alexa-568 (Invitrogen, A11075), Alexa 488 anti-rabbit antibody (Invitrogen, A11008; 1:500 dilution), Alexa 488 anti-mouse antibody (Invitrogen, A11001) antibodies were incubated at room temperature for 30 min (all 1:500 in PBS with 2% BSA). Images were captured as 15-25 z-stacks (0.1-μm step size) on a Nikon Ti Eclipse microscope with a Yokogawa CSU-XI spinning disc confocal with 100x (1.4 NA) Plan Apo objectives, and a Clara Interline CCD Camera (Andor Technology) using MetaMorph software (Molecular Devices). Images were processed using ImageJ using z-stack average maximum intensity projections and assembled in Adobe Photoshop. For quantification of the percentage of bacteria with pUb and p62, only bacteria that co-localized with rim-like patterns of the respective marker were scored as positive for staining. To quantify pUb, p62, NDP52, OmpB, and OmpA signal per bacteria, z-stacks were projected as stated above, and the edges of individual bacteria were marked by the freehand region of interest (ROI) function in ImageJ. Subsequently, the average pixel intensity within that ROI was measured. pUb/p62/NDP52 signal intensities were calculated by subtracting the average pUb/p62/NDP52

signal of WT bacteria from the pUb/p62/NDP52-value of each bacterium. OmpB signal intensity

was calculated by subtracting the average OmpB-signal of *ompB^STOP*::tn bacteria from the OmpB-

value of each bacterium. OmpA signal intensity was calculated by subtracting the average

background-signal (areas with no bacteria) from the OmpA-value of each bacterium.

Sample preparation for mass spectrometry to determine the lysine methylome

$5x10^7$ gradient-purified WT (Passage 6), *pkmt1*::tn (P4) and *pkmt2*::tn (P4) bacteria were

centrifuged at 11,000*g* for 3 min. Each pellet was resuspended in 50 µL Tris (10 mM)-EDTA (10

mM), pH 7.6, and incubated for 45 min in a 45 °C water bath. Bacterial surface fractions were

recovered from the supernatant after centrifugation at 11,000*g* for 3 min. Pellet was resuspended

as above and incubated for additional 45 min at 45 °C before resuspension in 50 µL Tris (10 mM)-

EDTA (10 mM). Both pellet and surface fractions were boiled at 95 °C for 10 min. Samples were

cooled to RT prior to addition of 20 µL 50 mM NH_4HCO_3, pH 7.5, and 50 µL of a 0.2% solution

of RapiGest (diluted in NH_4HCO_3, Waters, 186001861). Next, samples were heated at 80 °C for

15 min and cooled to RT before addition of 1 µg of trypsin (Promega, V511A). Samples were

digested at 37 °C overnight. To hydrolyze the RapiGest, 20 µL of 5% trifluoroacetic acid (TFA)

was added to samples which were incubated at 37 °C for 90 min prior centrifugation at 15000*g* for

25 min at 4 °C. Samples were desalted using C18 OMIX tips (Agilent Technologies, A57003100)

according to the manufacturer's instructions and sample volume was decreased to 20 µL using a

SpeedVac vacuum concentrator. Samples were stored at 4 °C prior to analysis.

TUBE assay and sample preparation for mass spectrometry

To enrich for polyubiquitylated proteins, 3x10^8 PFUs of "purified" WT (P6), $ompB^{STOP}$::tn

(P6) and $pkmt1$::tn (P4) and $pkmt2$::tn (P3) bacteria were centrifuged at 14,000g for 3 min at room

temperature. Next, to release the surface protein fraction, the bacterial pellets were resuspended in

lysis buffer (50 mM Tris-HCI, pH 7.5, 150 mM NaCl, 1 mM EDTA and 10% glycerol),

supplemented with 0.0031% v/v lysonase (Millipore, 71230), the deubiquitylase inhibitor PR619

(Life Sensor, SI9619) at a final concentration of 20 μM, and 0.8% w/v octyl β-D-glucopyranoside

(Sigma, O8001), and incubated on ice for 10 min with occasional pipetting of samples to break

pellet into smaller pieces. Subsequently, the lysate was cleared by centrifugation at 14,000g at 4

˚C for 5 min and incubated with equilibrated agarose TUBE-1 (Life Sensor, UM401) for 3 h, at 4

˚C. After binding of polyubiquitylated proteins to TUBE-1, agarose beads were washed 1 time

with TBS supplemented with 0.05% Tween and 5 mM EDTA, and subsequently 3 times with TBS

only (no Tween or EDTA) and centrifuged at 5,000g for 5 min. To prepare samples for MS

analysis, enriched proteins were digested at 37 ˚C overnight on agarose beads in RapiGest SF

solution (Waters, 186001861) supplemented with 0.75 μg trypsin (Promega, V511A). The reaction

was stopped using 1% TFA (Sigma, T6508). Octyl β-D-glucopyranoside was extracted using

water-saturated ethyl acetate (Sigma, 34858). Prior to submission of samples for mass

spectrometry analysis, samples were desalted using C18 OMIX tips (Agilent Technologies,

A57003100), according to the manufacturer's instructions.

Liquid chromatography-mass spectrometry

Samples of proteolytically digested proteins were analyzed using a Synapt G2-Si ion

mobility mass spectrometer that was equipped with a nanoelectrospray ionization source (Waters).

The mass spectrometer was connected in line with an Acquity M-class ultra-performance liquid

chromatography system that was equipped with trapping (Symmetry C18, inner diameter: 180 μm,

length: 20 mm, particle size: 5 μm) and analytical (HSS T3, inner diameter: 75 μm, length: 250

mm, particle size: 1.8 μm) columns (Waters). Data-independent, ion mobility-enabled, high-

definition mass spectra and tandem mass spectra were acquired in the positive ion mode (*23-25*).

Raw data acquisition was controlled using MassLynx software (version 4.1), and tryptic peptide

identification and relative quantification using a label-free approach (*26, 27*) were performed using

Progenesis QI for Proteomics software (version 4.0, Waters). Raw data were searched against

Rickettsia parkeri and *Chlorocebus sabaeus* protein databases (National Center for Biotechnology

Information, NCBI) to identify tryptic peptides, allowing for up to three missed proteolytic

cleavages, with diglycine-modified lysine (i.e., ubiquitylation remnant) and methylated lysine as

variable post-translational modifications. Calculation of the percentage of lysine methylation

(mono, di, tri or unmethylated), for each bacterial strain, was performed by dividing the abundance

of a residue/protein bearing a modification by the total abundance and multiplying by 100. Data-

dependent analysis was performed using an UltiMate3000 RSLCnano liquid chromatography

system that was connected in line with an LTQ-Orbitrap-XL mass spectrometer equipped with a

nanoelectrospray ionization source, and Xcalibur (version 2.0.7) and Proteome Discoverer

(version 1.3, Thermo Fisher Scientific, Waltham, MA) software, as described elsewhere (*28*).

Localization of tagged ubiquitin and ubiquitin pull-downs

To assess localization of 6xHis-ubiquitin during infection, confluent Vero cells grown in

24-well plates with coverslips were transfected with 2 μg of pCS2-6xHis-ubiquitin plasmid DNA

using Lipofectamine 2000 (Invitrogen, 11668-019) for 6 h in Opti-MEM (Gibco, 31985-070).

Subsequently, media were exchanged to media without transfection reagent, and cells were

incubated overnight at 37 ˚C and 5% CO_2. The following day (~16 h after transfection), transfected

cells were infected with purified WT or mutant bacteria at a MOI of 1. At 28 h.p.i., infected cells

were fixed with 4% PFA diluted in PBS, pH 7.4 for 10 min, then washed 3 times with PBS. Primary

anti-6xHis monoclonal mouse antibody (Clontech, 631212, diluted 1:1,000) was used to detect

6xHis-ubiquitin in samples permeabilized with 0.5% Triton-X100, and a goat anti-mouse Alexa-

568 (Invitrogen, A11004) to detect the primary 6xHis antibody. Hoechst (Thermo Scientific,

62249, diluted 1:2500) was used to detect host and bacterial DNA. Samples were imaged as

already described.

For ubiquitin pull-downs, confluent Vero cells grown in 6-well plates were transfected and

infected as described above. At 28 h.p.i., cells were washed once with 1x PBS, pH 7.4, and

subsequently lysed in urea lysis buffer (8 M urea, 50 mM Tris-HCI, pH 8.0, 300 mM NaCI, 50

mM Na_2HPO_4, 0.5% Igepal CA-630 (Sigma, I8896)) for 20 min at RT. Subsequently, samples

were sonicated, and lysate was cleared by centrifugation at 15000*g* for 15 min at room temperature.

Prior to incubation with Ni-NTA resin, an aliquot was saved for the input sample. 6xHis-ubiquitin

conjugates were purified by incubation and rotation with Ni-NTA resin for 3 h, at RT, in the

presence of 10 mM imidazole. Beads were washed 3 times with urea lysis buffer and 1 time with

urea lysis buffer lacking Igepal CA-630. Ubiquitin conjugates were eluted at 65 ˚C for 15 min in

2x Laemmli buffer containing 200 mM imidazole and 5% 2-mercaptoetanol (Sigma, M6250),

vortexed for 90 seconds, and centrifuged at 5000*g* for 5 min at RT. Eluted and input proteins were

detected by SDS-PAGE followed by western blotting, as described above.

Animal experiments

Animal research using mice was conducted under a protocol approved by the UC Berkeley

Institutional Animal Care and Use Committee (IACUC) in compliance with the Animal Welfare

Act. The UC Berkeley IACUC is fully accredited by the Association for the Assessment and

Accreditation of Laboratory Animal Care International and adheres to the principles of the Guide

for the Care and Use of Laboratory Animals. Infections were performed in a biosafety level 2

facility. Mice were age-matched between 8 and 18 weeks old. Mice were selected for experiments

based on their availability, regardless of sex. All mice were healthy at the time of infection and

were housed in microisolator cages and provided chow and water. Littermates of the same sex

were randomly assigned to experimental groups. For infections, *R. parkeri* was prepared by

diluting 30%-prep bacteria into cold sterile 1x PBS to 5×10^6 PFU per 200 mL. Bacterial

suspensions were kept on ice during injections. Mice were exposed to a heat lamp while in their

cages for approximately 5 min, and then each mouse was moved to a mouse restrainer (Braintree,

TB-150 STD). The tail was sterilized with 70% ethanol, and 200 µL of bacterial suspensions were

injected using 30.5-gauge needles into the lateral tail vein. Body temperatures were monitored

using a rodent rectal thermometer (BrainTree Scientific, RET-3). Mice were monitored daily for

clinical signs of disease, such as hunched posture, lethargy, or scruffed fur. If a mouse displayed

severe signs of infection, as defined by a reduction in body temperature below 90 ˚F or an inability

to move around the cage normally, the animal was immediately and humanely euthanized using

CO_2 followed by cervical dislocation, according to IACUC-approved procedures (*16*).

Statistical analysis, experimental variability and reproducibility

Statistical parameters and significance are reported in the legends. Data were considered

to be statistically significant when $p < 0.05$, as determined by a one-way ANOVA with Dunnett's

post-hoc test, a Kruskal-Wallis test with Dunn's post-hoc test, a Brown-Forsyth and Welch

ANOVA with Dunnett's post-hoc test, or a two-way ANOVA (all two-sided). Statistical analysis

was performed using PRISM 6 software (GraphPad Software). If not otherwise described, *n*

indicates the number of independent biological experiments executed at different times.