A unifying framework for quantifying and comparing n-dimensional hypervolumes

Muyang Lu, Kevin Winner, Walter Jetz

Abstract

The quantification of Hutchison's n-dimensional hypervolume has enabled substantial progress in community ecology, species niche analysis and beyond. While non-parametric methods for quantifying and comparing hypervolumes are popular and require no assumptions about the underlying distributions of data, they have limited mechanistic interpretability. Here, we propose the use of multivariate normal distributions for the assessment and comparison of niche hypervolumes and introduce this as the multivariate-normal hypervolume (MVNH) framework. The framework provides parametric measures of the size and dissimilarity of niche hypervolumes, each of which can be partitioned into mechanistically interpretable components. Specifically, we propose the use of 1) the determinant of the covariance matrix (e.g. the generalized variance) of a MVNH as a measure of total niche size, which can be partitioned into the components of univariate niche variances and a correlation component; and 2) the Bhattacharyya distance between two MVNHs as a measure of niche dissimilarity, which can be partitioned into the components of Mahalanobis distance between hypervolume centroids and the determinant ratio which measures size difference. We use empirical data at the community (functional diversity) and species (environmental niche breadth) levels to demonstrate the new insights provided by these metrics and their components. Our approach overcomes several operational and computational limitations of non-parametric methods and provides a framework that offers both unification and granularity in the assessment of niche volumes and differences.

Introduction

The n-dimensional hypervolume is one of the most fundamental (Hutchinson, 1957; Whittaker, Levin, & Root, 1973; Pulliam, 2000) and commonly used concepts (Díaz et al., 2015; Blonder, 2017; Pironon et al., 2018) in ecology and evolutionary biology. Hutchinson first described the niche as the n-dimensional hypervolume in which a species can survive and reproduce (Hutchinson, 1957). The use of hypervolume was later extended to describe functional space and trait space (Lamanna et al., 2014; Laughlin, 2014; Pigot et al., 2020). The geometric features of n-dimensional hypervolumes (especially their size and dissimilarity) are associated with a wide range of hypotheses and applications in ecology and evolution. For example, the size of a climatic niche hypervolume is hypothesized to drive species diversification rates: in mammals, species with small niches (specialists) have been shown to have higher speciation rates and lower extinction rates than those with large niches (generalists) (Gómez-Rodríguez, Baselga, & Wiens, 2015; Rolland & Salamin, 2016). The size of climatic niche hypervolume is also hypothesized to drive the variations of geographic range size (Slatyer, Hirst, & Sexton, 2013; Saupe et al., 2015; Cardillo, Dinnage, & McAlister, 2018; Ficetola, Lunghi, & Manenti, 2020). Similarity between species' environmental niche hypervolumes or functional trait hypervolumes is used to measure niche divergence or niche packing among co-occurring species, which is hypothesized to determine species' coexistence and species richness patterns (Serra-Varela et al., 2015; Pigot, Trisos, & Tobias, 2016; Castro-Insua, Gómez-Rodríguez, Wiens, & Baselga, 2018; Read et al., 2018). Niche similarity is also used in within-species comparisons such as assessing the impact of climate change (Tayleur et al., 2015; Gómez, Tenorio, Montoya, & Cadena, 2016; Zurell, Gallien, Graham, & Zimmermann, 2018) and whether there are niche shifts during biotic

invasion (Lauzeral et al., 2011; Early & Sax, 2014; Guisan, Petitpierre, Broennimann, Daehler, & Kueffer, 2014; Davies, Hill, McGeoch, & Clusella-Trullas, 2019).

The quantification of niche volume and similarity has thus become an urgent pursuit for both theoretical investigation and real-world applications. Recently, non-parametric methods, such as kernel-density estimates (Blonder, Lamanna, Violle, & Enquist, 2014; Blonder et al., 2017; Eckrich, Albeke, Flaherty, Bowyer, & Ben-David, 2020; Mammola & Cardoso, 2020), dynamic range boxes (Junker, Kuppler, Bathke, Schreyer, & Trutschnig, 2016), minimum convex hulls, and support vector machines (Blonder et al., 2017), have been widely favored because of their minimal assumptions around the distribution of data and the existence of well-documented statistical packages supporting their exploration and use. In these methods, an n-dimensional distribution over environmental or trait space is typically converted to a hypervolume with a hard boundary defined as the region enclosed by a particular isopleth (often 95%). Niche size is calculated as the volume of the hard-boundaried space. Niche similarity of two hypervolumes can be quantified in multiple ways, either with the union and intersection of hypervolumes (Jaccard and Sorensen indices) or with the distances between the two hypervolumes (Loiseau et al., 2017; Mammola, 2019).

These non-parametric methods of quantifying hypervolumes have several methodological limitations: 1) niche volumes and similarity measures are sensitive to the parameterization of the underlying distribution and depend additionally on the choice of isopleth (e.g. 95% or 50%), 2) computational costs are high when the dimension of variables and sample size are large (Blonder et al., 2014), and most importantly 3) the non-parametric methods generally cannot say anything about the relative importance of the constituent components of the various measures (but see Junker et al., 2016; Loiseau et al., 2017). For example, most non-

parametric methods do not provide information on the impact of each individual dimension on the size of a hypervolume. For those that do (e.g. dynamic range box; Junker et al., 2016), the shape information (correlation among variables) on hypervolume size is ignored. Another example is the use of Jaccard/Sorensen indices and distance metrics for hypervolume similarity comparison. Though the two classes of metrics (Jaccard and Sorensen indices, Euclidean distances between niche centroids) are complementary in capturing hypervolume dissimilarity (Mammola, 2019), no analytical link exists between them for comparing their relative importance on overall hypervolume similarity. The Jaccard and Sorensen indices are used to calculate how much hypervolume is shared between the two hypervolumes, which are more informative when two hypervolumes intersect with each other. On the contrary, the Euclidean distance between hypervolume (centroid distance or minimum distance) is more informative when two hypervolumes are disjunct (Mammola, 2019).

The prevalence of non-parametric methods in quantifying hypervolumes also hampers the integration of empirical niche studies with niche theories because empirical metrics are disconnected from theoretical derivations. For example, in MacArthur and Levin's work on limiting similarity (MacArthur & Levins, 1967) and in Rappoldt and Hogeweg's work on niche packing (Rappoldt & Hogeweg, 1980), the outcomes of competition exclusion are calculated based on competition coefficients derived from one-dimensional Gaussian utilization curves. However, the testable theoretical predictions derived by these studies rely on quantitative analyses in higher dimensions, meaning that questions such as "Does higher niche similarity among species lead to more competitive exclusion" are to date only able to be studied in a limited or qualitative way with non-parametric methods (D'Andrea & Ostling, 2016). Although various parametric dissimilarity metrics have been proposed (Morisita, 1959; Pianka, 1974; Lu, Smith, & Good, 1989), they were largely ignored because of their

conceptual disconnection with the prevalent size metrics of hypervolume. Parametric

measures that have clear analytical links to theory and which unify both size and dissimilarity

measurements are urgently needed.

To fill in this gap, we propose a framework to quantify niche volume and dissimilarity

based on the assumption of multivariate normal (MVN) distribution of the niche variables

under study and call this the MVN hypervolume or MVNH framework. We chose the

multivariate normal distribution because it is the most widely used assumption for niche

assessment and modeling both in theoretical (MacArthur & Levins, 1967; Rappoldt &

Hogeweg, 1980; Soberon & Nakamura, 2009; Jiménez, Soberón, Christen, & Soto, 2019) and

empirical work (Swanson et al., 2015; La Sorte, Fink, & Johnston, 2018), which provides

room for integration and synthesis of size and dissimilarity measures (Lu et al., 1989). We

show that in the MVN hypervolume framework niche size is derived from the covariance

matrix (Soberon & Nakamura, 2009), and that niche dissimilarity can be quantified from the

Bhattacharyya distance (Bhattacharyya, 1946; Lu et al., 1989; Winner et al., 2018). We

demonstrate through analytical derivation and empirical examples how the MVNH

framework's metrics support a natural partitioning that yields more mechanistic insights for

hypervolume comparisons than existing methods, and allows a more rigorous assessment of

hypervolume variation.

Methods and Materials

In our MVNH framework, we propose that a multi-dimensional niche space be described by a

multivariate normal distribution with probability density function:

$$f_X(\mathbf{X}) = \frac{\exp\left(-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})^\mathrm{T}\Sigma^{-1}(\mathbf{x}-\boldsymbol{\mu})\right)}{\sqrt{(2\pi)^n|\Sigma|}},$$

… (1)

where $\mathbf{x} = \{x_1, ..., x_n\}$ is a vector in the n-dimensional environmental niche space, $\mathbf{\mu} = \{\mu_1, ..., \mu_n\}$ is the environmental niche centroid, Σ is the covariance matrix for the n-dimensional environmental niche, and $|\Sigma|$ denotes the determinant of the covariance matrix.

The size of a hypervolume

We define the size of an MVN hypervolume as the determinant of the covariance matrix $|\Sigma|$, also called the "generalized variance" (La Sorte et al., 2018). It has a geometric interpretation as the volume of the n-dimensional parallelepiped spanned by the column or row vectors of the covariance matrix. For example, the covariance matrix Σ and the generalized variance $|\Sigma|$ of a two-dimensional MVNH are:

$$\Sigma = \begin{bmatrix} \sigma_x{}^2 & \rho_{xy}\sigma_x\sigma_y \\ \rho_{xy}\sigma_x\sigma_y & \sigma_y{}^2 \end{bmatrix},$$

... (2)

$$|\Sigma| = \sigma_x{}^2\sigma_y{}^2\left(1 - \rho_{xy}{}^2\right),$$

... (3)

where $\sigma_x{}^2$ is the variance of variable x, $\sigma_y{}^2$ is the variance of variable y, and ρ_{xy} is the correlation coefficient between the two environmental variables x and y. When log-transformed, Equation 3 enables us to additively partition the size of a 2-dimensional hypervolume (in terms of the determinant) into three separate components (Fig. 1a): the variance of the first niche axis $\sigma_x{}^2$, the variance of the second niche axis $\sigma_y{}^2$, and a correlation component $(1-\rho_{xy}{}^2)$. The correlation component could be seen as a shrinkage factor which shrinks the niche volume by a factor of $(1-\rho_{xy}{}^2)$. Thus, when two niche axes are independent, the niche size is just the product of the variances of individual axes $\sigma_x{}^2\sigma_y{}^2$. Similarly, for a three-dimensional hypervolume, the determinant of the covariance matrix is:

$$|\Sigma| = \sigma_x{}^2\sigma_y{}^2\sigma_z{}^2\left(1 - \rho_{xy}{}^2 - \rho_{yz}{}^2 - \rho_{xz}{}^2 + 2\rho_{xy}\rho_{yz}\rho_{xz}\right),$$

… (4)

where $\left(1 - \rho_{xy}{}^2 - \rho_{yz}{}^2 - \rho_{xz}{}^2 + 2\rho_{xy}\rho_{yz}\rho_{xz}\right)$ is the correlation component. The factoring of the determinant generalizes easily to higher dimensions: the determinant is just the product of all individual univariate variances and a correlation component.

The determinant also has a close relationship with the standardized ellipse area (SEA, the ellipse with its major and minor axes defined by the principal components; Jackson, Inger, Parnell, & Bearhop, 2011): it scales with the SEA by a constant (π for two-dimensional niche and $4\pi/3$ for three-dimensional niche). Factoring the determinant (as in Equations 3 and 4) can be used to assess the relative importance of the separate components in driving changes in the size of the hypervolume, such as when assessing whether individual trait axis or the constraint on the combination of traits is the major driver of community-wide trait diversity.

The dissimilarity of hypervolumes

We suggest measuring the dissimilarity of two MVN hypervolumes with the Bhattacharyya distance (Bhattacharyya, 1946; Lu et al., 1989; Minami & Shimizu, 1999; Fieberg & Kochanny, 2005; Winner et al., 2018). The Bhattacharyya distance (BD) of two continuous probability distributions, A and B defined on the same domain $\mathcal{X}$ is:

$$BD(A, B) = -\log BC(A, B),$$

… (5)

where BC is the Bhattacharyya coefficient:

$$BC(A, B) = \int_{x \in \mathcal{X}} \sqrt{A(x)B(x)}\, dx$$

… (6)

Therefore, for two probability distributions, the BC ranges from 0 to 1, and the BD ranges from 0 to ∞.

For two multivariate normal hypervolumes A and B, the BD has a closed form:

$$BD(A,B) = \underbrace{\frac{1}{8}(\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)^T \overline{\Sigma}^{-1}(\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)}_{MD} + \underbrace{\frac{1}{2}\ln\left(\frac{|\overline{\Sigma}|}{\sqrt{|\Sigma_A||\Sigma_B|}}\right)}_{DR},$$

… (7)

where μ_A and μ_B are the centroids of A and B respectively and $\overline{\Sigma}$ is the mean of the covariance matrices of A and B:

$$\overline{\Sigma} = \frac{\Sigma_A + \Sigma_B}{2}.$$

… (8)

The first term in Equation 7 is the Mahalanobis distance (MD) between two hypervolume centroids, and the second term in Equation 7 is the determinant ratio (DR) which measures size difference between A and B (see Fig. 1d). If two hypervolumes have the same size, then the second term in Equation 7 will equal 0. As with Equations 3 and 4, factoring the Bhattacharyya distance according to Equation 7 can be used to assess the relative contributions of the Mahalanobis distance (centroid difference) and the determinant ratio (size difference) to the overall niche dissimilarity. We suggest that the ratio of Mahalanobis distance to determinant ratio (MD/DR), can be used as measure of the relative importance of centroid distance and size difference in driving overall dissimilarity.

We recognize other dissimilarity metrics based on multivariate normal distributions which share a similar form with the Bhattacharyya distance and which could also be partitioned into a Mahalanobis distance component and a determinant ratio component (see Lu et al., 1989), such as the MacArthur-Levins measure (MacArthur & Levins, 1967) :

$$MacArthur - Levins(A, B)_A = \underbrace{(\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)^T \overline{\Sigma}^{-1} (\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)}_{MD} + \underbrace{\ln\left(\frac{|\Sigma_A|}{|\Sigma_A||\Sigma_B|}\right)}_{DR},$$

$$MacArthur - Levins(A, B)_B = \underbrace{(\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)^T \overline{\Sigma}^{-1} (\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)}_{MD} + \underbrace{\ln\left(\frac{|\Sigma_B|}{|\Sigma_A||\Sigma_B|}\right)}_{DR},$$

… (9)

the Pianka's measure (Pianka, 1974):

$$Pianka(A, B)_A = \underbrace{(\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)^T \overline{\Sigma}^{-1} (\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)}_{MD} + \underbrace{\frac{1}{2}\ln\left(\frac{|\overline{\Sigma}|}{\sqrt{|\Sigma_A||\Sigma_B|}}\right)}_{DR},$$

… (10)

and the Morisita's measure (Morisita, 1959):

$$Morisita(A, B) = \underbrace{(\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)^T \overline{\Sigma}^{-1} (\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)}_{MD} + \underbrace{\ln\left(\frac{\sqrt{|\overline{\Sigma}|}(\sqrt{|\Sigma_A|}+\sqrt{|\Sigma_B|})^2}{2\sqrt{|\Sigma_A||\Sigma_B|}}\right)}_{DR}.$$

… (11)

These metrics differ in their probabilistic interpretations, weighting of each component, and how they calculate the determinant ratio component. We will focus on the use of the Bhattacharyya distance in this paper because it was intensively studied with known statistical properties (Lu et al., 1989; Minami & Shimizu, 1999; Winner et al., 2018), and also has a straightforward interpretation as the integration of geometric means of two probability distributions (Equation 6).

It should also be noticed that the Mahalanobis distance component in Equations 7-11 could be linked to an existing partitioning framework which additively partitions the Mahalanobis distance into standardized Euclidean distances along PCA axes which measure the relative importance of each PCA axis (Calenge, Darmon, Basille, Loison, & Jullien, 2008; Mahony, Cannon, Wang, & Aitken, 2017).

Estimators

We used sample means and sample covariances as estimates of μ_A, μ_B, Σ_A, and Σ_B for the calculation of the determinant of the covariance matrices and the Bhattacharyya distance with its partitioned components. These plug-in estimates of the Bhattacharyya distance are known to be biased, but the bias is small with low dimensionality of data (Lu et al., 1989). For the purpose of demonstrating the utility of the partitioning framework, our empirical analysis proceeds without applying existing bias-correction methods which have been presented in the literature such as estimating the covariance matrix using Bayesian methods (Swanson et al., 2015), and bias-correction techniques for the Bhattacharyya distance (Minami & Shimizu, 1999; Winner et al., 2018).

Empirical examples

Community example: Functional diversity and turnover

We used a trait dataset of 36 annual plant communities in Western Australia (Dwyer & Laughlin, 2017) to demonstrate how functional diversity (alpha diversity) and turnover (beta diversity) can be measured in the MVN hypervolume framework using the determinant and the Bhattacharyya distance, respectively. The three measured traits for each species were specific leaf area (SLA), maximum height (MH), and seed mass (SM). We then assessed each component of the trait volume (SLA variance, MH variance, SM variance and the correlation component) as predicted by growing season available moisture, mean growing season minimum temperature, and soil surface dispersion (dispersion is a measure of heterogeneity) in multiple linear regression. We used the coefficient of variation of each component to assess the relative contribution of each trait volume component to the variation of total trait

volume, because the individual niche variances are scalable and not directly comparable with each other. We assessed the environmental drivers of the Bhattacharyya distance (functional turnover) components using multiple regression on distance matrices (MRM) based on a partial Mantel test (Lichstein, 2007).The regression coefficients of MRM are calculated with ordinary least square regression, but the p-values of the coefficients are calculated with matrix permutation tests. The predictor variables were calculated as the Euclidean distances for each environmental variable between sites.

Species example: Environmental niche breadth and dissimilarity

We studied 260 bird species in the Western Hemisphere with occurrence data extracted from eBird (Sullivan et al., 2009), environmental layers obtained from CHELSA (monthly mean temperature and precipitations for breeding seasons, https://chelsa-climate.org/), and MODIS EVI (https://modis.gsfc.nasa.gov/data/dataprod/mod13.php) at 1km resolution to assess the drivers of realized environmental niche breadth (volume) and niche dissimilarity between species. 1000 random points were sampled and spatially thinned to 10km distance using the 'spThin' package (Aiello-Lammens, Boria, Radosavljevic, Vilela, & Anderson, 2015) in R for extraction of environmental data. We then investigated the drivers of environmental niche breadth components using ordinary multiple linear regression, and assessed the drivers of the Bhattacharyya distance components using MRM. We obtained body mass and dietary niche breadth data from the 'EltonTrait' dataset (Wilman et al., 2014). Area and centroid absolute latitude of geographic range were calculated from IUCN range maps (https://www.iucnredlist.org/resources/spatial-data-download).

Comparisons to non-parametric metrics

For both the community and niche examples we compared the newly proposed size and dissimilarity metrics from the MVNH framework with measures based on the 95% isopleth of a kernel density estimator using the 'hypervolume' package in R (Blonder et al., 2017). For the species environmental niche example, we also compared the determinant of the MVNH with the volume of minimum convex hull; for the community functional diversity example, we additionally compared the determinant with three commonly used functional diversity measures: functional divergence, functional dispersion, and functional evenness (Laliberté, Legendre, & Shipley, 2015). For both examples, we compared the Bhattacharyya distance with Jaccard (Fig. 1e), Sorensen similarities, centroid distances (Fig. 1f) and minimum distances between hypervolumes.

Results

We use the MVNH framework to firstly analyze drivers of community-level functional diversity and functional turnover, and to secondly assess drivers of species-level environmental niche breadth and dissimilarity. Finally, we show the correlations between the proposed metrics with commonly used non-parametric measures.

Community level: Functional diversity

We compared functional diversity (measured as the determinant of the trait covariance matrix) and functional turnover (measured as Bhattacharyya distance) among 36 plant communities. For a two-community example (site 1 and site 2), we partitioned the functional diversity of each site into four components: the univariate variance of each of the three traits (specific leaf area, SLA; maximum height, MH; and seed mass, SM) and the correlation component following Equation 4. Component-wise differences between the two sites add up to the

difference in total trait volume (the determinant of the environmental niche covariance matrix) on the log-scale (Fig. 2a).

For the 36-community assessment, the coefficient of variation for each trait volume component across the 36 sites was 0.31 for the correlation component, and varied between 0.28 (MH) and 0.16 (SM) for the individual trait variances. This suggests that the correlation component is the largest driver of the variation in functional diversity among sites. By relating individual trait variances and the correlation component to environmental predictors, we identify mean moisture availability and, less so, minimum temperature as key correlates of trait diversity variation (Fig. 2c, d). The variances of all three traits differ substantially in the strength and direction of their respective associations with environmental predictors. Linear regression shows that none of the examined environmental variables predict the variation of SLA variance (Table 1). Both minimum temperature and moisture availability are positively correlated with maximum height variance, whereas only soil surface dispersion is positively correlated with seed mass variance. Moisture availability is also positively correlated with the correlation component. The overall trait volume (the determinant) is only correlated with moisture availability (Table 1).

We measured the pairwise functional turnover between any two sites as the Bhattacharyya distance between their MVN hypervolumes, and according to Equation 7, partitioned it into the Mahalanobis distance (which measures the standardized distance between community means) and the determinant ratio (which measures the differences in functional alpha diversity) (see a two-community example in Fig. 2b). The ratio of Mahalanobis distance to determinant ratio (MD/DR) reflects the relative importance of difference in community means and trait diversity differences in driving the overall community-level functional turnover. The median ratio MD/DR across all pairs of sites is 0.63, suggesting functional

turnover among communities is mainly driven by the variation in functional alpha-diversity. The functional turnover is more determined by moisture availability than by minimum temperature (Fig. 2e, f). Multiple regression on distance matrices suggests that the Mahalanobis distance is correlated with both minimum temperature and moisture availability, but the determinant ratio is only correlated with moisture availability (Table 1).

Species level: environmental niches

We used the presented methodology to analyze and compare the three-dimensional environmental niches (temperature, precipitation, and EVI) of 260 bird species in the New World. We partitioned the total environmental niche volume for each single species into the three environmental variable variance components and the correlation component. For a two-species example, we attributed the difference in niche volume between the white-throated swift and the tricolored blackbird to differences in temperature variance, precipitation variance, EVI variance and the correlation component (Fig. 3a). For the 260-species, assessment, we found that the coefficient of variation for each of the niche volume components among 260 bird species was 0.79 for precipitation variance, 0.67 for temperature variance, 0.44 for EVI variance, and 0.22 for the correlation component, suggesting that precipitation niche breadth variation contributes most to the species-level niche volume variation. Range size and centroid latitude are the two main drivers of realized environmental niche volume (Fig. 3c, d), but the predictors for each niche volume component are different: the precipitation variance and the correlation component are only correlated with the centroid latitude while the temperature variance and the EVI variance are correlated with both centroid latitude and range size (Table 2). The median ratio between Mahalanobis distance and determinant ratio is 2.43, suggesting that the distance between niche centroids is more important than niche volume differences in driving overall environmental niche dissimilarity

among the 260 bird species. When the multiple regression on distance matrices (MRM) is applied, different components of the Bhattacharyya distance have different determining factors: the Mahalanobis distance is only correlated with centroid latitude; while the determinant ratio is correlated with both range size and centroid latitude (Table 2).

Comparison with other metrics

Community-level functional traits: The determinant of trait covariance matrix as the measure of functional diversity had the highest correlation with the area of the 95% isopleth of a kernel density derived hypervolume (0.90), followed by functional dispersion (0.79) and minimum convex hull (0.65) (additional methods in Fig. 4a). The Bhattacharyya distance had a similar correlation with Euclidean centroid distance (0.69), Jaccard similarity (-0.69) and Sorensen similarity (-0.71). The Mahalanobis distance component was more correlated with Euclidean centroid distance (0.71) than with Jaccard similarity (-0.56) or Sorensen similarity (-0.59), while the determinant ratio component had a relatively low correlation with the Euclidean centroid distance (0.4; Fig. 4b). All components were uncorrelated with minimum distance between hypervolumes (Fig. 4b).

Species-level environmental niche: the determinant of the environmental niche covariance matrix correlates most strongly with the area of the 95% isopleth of the kernel density derived hypervolume (0.85), followed by the minimum convex hull (0.77, Fig. 4c). The Bhattacharyya distance and its Mahalanobis distance component are most correlated with the Euclidean centroid distance (0.87 and 0.85 respectively); the determinant ratio is similarly correlated with most alternative metrics except for minimum distance between hypervolumes (Fig. 4d).

Discussion

We proposed a new parametric hypervolume framework based on multivariate normal distributions and corresponding size and dissimilarity metrics. The greatest advantage of the MVN hypervolume approach is the analytical clarity and insights provided by the partitioned components of the niche size and dissimilarity metrics. The MVNH framework is also significantly faster than the widely used KDE approach. For example, using the 'Iris' dataset in R (a dataset with 150 rows and 3 columns), it takes 6 seconds to compute the hypervolume using KDE but only 0.001 seconds to compute the hypervolume using MVNH. Although the determinant of the covariance matrix has been used as a total niche volume measure in some studies (Soberon & Nakamura, 2009; La Sorte et al., 2018), it has never been illustrated that the determinant could be interpreted as the product of individual niche variances and a correlation component. The Bhattacharyya distance has been used as a measure of home range overlap in animal movements (Fieberg & Kochanny, 2005; Winner et al., 2018). We suggest that its use in trait or environmental space (Lu et al., 1989; Minami & Shimizu, 1999), and in particular the interpretation of the partitioned components, offers valuable biological insights. Specifically, the Mahalanobis distance between hypervolume centroids can be interpreted as the standardized distance between optimal traits of two communities or between the optimal environmental conditions of two species under the assumptions that mean community traits represents the optimal traits under equilibrium (Blonder et al., 2015) and that niche centroid has the highest fitness for a species (Sagarin, Gaines, & Gaylord, 2006). The determinant ratio can be interpreted as the functional diversity difference between communities or as niche volume differences between species. The Bhattacharyya distance (BD) not only avoids choosing ad hoc algorithm parameters, but more importantly reconciles the limitations of the two commonly used classes of dissimilarity metrics. When Jaccard or Sorensen indices become uninformative for disjunct hypervolumes, the BD incorporates the

distance information through the Mahalanobis distance component. When the centroid distance metric becomes uninformative for two close and overlapping hypervolumes, the BD incorporates the shape information through the determinant ratio component.

The comparison with existing metrics shows that the determinant of the covariance matrix is most correlated with kernel density derived volume estimates (0.90 for the trait example and 0.85 for the environmental niche example). The good concordance between our method and the kernel density method across each of our studies suggests that the determinant as a measure of niche volume is robust to different distributions of data. The BD metric is most correlated with the centroid Euclidean distance in the environmental niche example (0.87) but is similarly correlated with Jaccard, Sorensen and centroid Euclidean distances in the trait example (-0.69, -0.71, and 0.69, respectively). Both the partitioned components of BD show a moderate amount of correlation with the Jaccard, Sorensen and the centroid Euclidean distances (0.42 – 0.75), showing that the partitioned components of BD both capture a useful amount of information about the shape of the hypervolume. Therefore, our BD metrics combine the important interpretations of both centroid distance metrics and the Jaccard and Sorensen metrics of niche dissimilarity. More importantly, the relative importance of centroid difference and volume difference are directly comparable using the BD metrics.

Other metrics such as the MacArthur-Levins measure (Equation 9), the Pianka's measure (Equation 10) and the Morisita's measure (Equation 11) could be similarly partitioned into a Mahalanobis distance component and a determinant ratio component (Lu et al., 1989), though their properties are less well known than the Bhattacharyya distance (Minami & Shimizu, 1999; Winner et al., 2018). The shared Mahalanobis distance component in all these metrics could be further partitioned into standardized Euclidean distance along different PCA axes according to a previously proposed partitioning framework (Calenge et al., 2008; Mahony et

al., 2017), which measures the relative importance of each PCA axis in determining the overall Mahalanobis distance.

Case studies

Our methods provide mechanistic understanding of the variation of each component in driving total functional diversity or environmental niche volume: in the community functional diversity example, we demonstrated that the constraints on combination of traits is the major driver of functional alpha diversity variation rather than the change of individual trait variances, which in turn is shaped by moisture availability; the major driver of functional turnover among the 36 annual plant communities is the variation of functional alpha-diversity rather than the mean trait differences between communities. In the environmental niche example, our method shows that the major driver of niche volume variation is the precipitation niche breadth variation (Fig. 3d) which is determined by centroid latitude of species' range. We also show that the range size-niche breadth relationship of total environmental niche breadth is mainly driven by the range size-temperature niche breadth and range size-EVI niche breadth relationship (Table 2).

Caveats

A common limitation of all hypervolume metrics is the curse of dimensionality and collinearity among axes (Blonder et al., 2014; D'Andrea & Ostling, 2016). The use of the determinant is especially sensitive to collinearity because any pair of fully dependent columns or rows in a covariance matrix will necessarily result in a determinant of zero. Increasing dimensions of trait or environmental data will increase the chance of collinearity and is also more likely to violate the multivariate normal assumption. Such an issue could be resolved by choosing the biologically relevant variables a priori or using common dimension

reduction techniques applied in non-parametric methods such as PCA (Junker et al., 2016). Because PCA creates orthogonal axes, the correlation component will always equal 1 if PCA is applied, and thus it is only recommended when the correlation among axes is of no interest to the study.

Potential

The presented size and dissimilarity measures are to our knowledge the first parametric metrics that allow a partitioning of n-dimensional hypervolumes. The partitioning of total niche volume and niche dissimilarity enables more rigorous testing of ecological and evolutionary hypotheses beyond those we explored in our two case studies. For example, when investigating the spatial scaling of niche volume (Pearman, Guisan, Broennimann, & Randin, 2008; Connor et al., 2017), the scaling relationship could be broken down to the scaling of univariate niche variances and correlation component, each of which could be linked to different sets of predictors. When studying niche shifts during biotic invasion (Lauzeral et al., 2011; Hill, Hoffmann, Macfadyen, Umina, & Elith, 2012; Wiens, Litvinenko, Harris, & Jezkova, 2019), the BD could be used to determine whether the niche shift is mainly caused by shifts in niche centroids or caused by volume expansion or reduction; the determinant could be used to assess which niche axis is most responsible for the total volume expansion or reduction. In niche evolution studies, the evolution of niche dissimilarity (Nunes & Pearson, 2017; Castro-Insua et al., 2018) between two species could be traced to contributions by niche breadth or center differences to the total niche divergence. In other words, the newly proposed metrics have the potential to provide more mechanistic understanding of biodiversity patterns and niche evolution, and to pave the way for more accurate forecasts for biodiversity change.

Acknowledgement

We thank Jetz Lab members for discussion of the work, especially Richard Li, Aurore Maureaud and Ben Carlson for comments on the first draft.

Reference

Aiello-Lammens, M. E., Boria, R. A., Radosavljevic, A., Vilela, B., & Anderson, R. P. (2015). spThin: An R package for spatial thinning of species occurrence records for use in ecological niche models. *Ecography*, *38*(5), 541–545. doi:10.1111/ecog.01132

Bhattacharyya, A. (1946). On a Measure of Divergence between Two Multinomial Populations. *Sankhyā: The Indian Journal of Statistics*. doi:10.1038/157869b0

Blonder, B. (2017). Hypervolume concepts in niche- and trait-based ecology. *Ecography*, (August), 1–13. doi:10.1111/ecog.03187

Blonder, B., Lamanna, C., Violle, C., & Enquist, B. J. (2014). The n-dimensional hypervolume. *Global Ecology and Biogeography*, *23*(5), 595–609. doi:10.1111/geb.12146

Blonder, B., Morrow, C. B., Maitner, B., Harris, D. J., Lamanna, C., Violle, C., … Kerkhoff, A. J. (2017). New approaches for delineating n-dimensional hypervolumes. *Methods in Ecology and Evolution*, (June 2017), 305–319. doi:10.1111/2041-210X.12865

Blonder, B., Nogués-Bravo, D., Borregaard, M. K., Donoghue, J. C., Jørgensen, P. M., Kraft, N. J. B., … Enquist, B. J. (2015). Linking environmental filtering and disequilibrium to biogeography with a community climate framework. *Ecology*, *96*(4), 972–985. doi:10.1890/14-0589.1

Calenge, C., Darmon, G., Basille, M., Loison, A., & Jullien, J. M. (2008). The factorial decomposition of the Mahalanobis distances in habitat selection studies. *Ecology*, *89*(2), 555–566. doi:10.1890/06-1750.1

Cardillo, M., Dinnage, R., & McAlister, W. (2018). The relationship between environmental niche breadth and geographic range size across plant species. *Journal of Biogeography*, (November 2016), 1–13. doi:10.1111/jbi.13477

Castro-Insua, A., Gómez-Rodríguez, C., Wiens, J. J., & Baselga, A. (2018). Climatic niche divergence drives patterns of diversification and richness among mammal families. *Scientific Reports*, *8*(1), 8781. doi:10.1038/s41598-018-27068-y

Connor, T., Hull, V., Viña, A., Shortridge, A., Tang, Y., Zhang, J., … Liu, J. (2017). Effects of grain size and niche breadth on species distribution modeling. *Ecography*, (October), 1–12. doi:10.1111/ecog.03416

D'Andrea, R., & Ostling, A. (2016). Challenges in linking trait patterns to niche differentiation. *Oikos*, *125*(10), 1369–1385. doi:10.1111/oik.02979

Davies, S. J., Hill, M. P., McGeoch, M. A., & Clusella-Trullas, S. (2019). Niche shift and resource supplementation facilitate an amphibian range expansion. *Diversity and Distributions*, *25*(1), 154–165. doi:10.1111/ddi.12841

Díaz, S., Kattge, J., Cornelissen, J. H. C., Wright, I. J., Lavorel, S., Dray, S., … Gorné, L. D. (2015). The global spectrum of plant form and function. *Nature*, 1–17. doi:10.1038/nature16489

Dwyer, J. M., & Laughlin, D. C. (2017). Constraints on trait combinations explain climatic drivers of biodiversity: the importance of trait covariance in community assembly. *Ecology Letters*. doi:10.1111/ele.12781

Early, R., & Sax, D. F. (2014). Climatic niche shifts between species' native and naturalized ranges raise concern for ecological forecasts during invasions and climate change. *Global Ecology and Biogeography*, *23*(12), 1356–1365. doi:10.1111/geb.12208

Eckrich, C. A., Albeke, S. E., Flaherty, E. A., Bowyer, R. T., & Ben-David, M. (2020). rKIN: Kernel-based method for estimating isotopic niche size and overlap. *Journal of Animal*

Ecology, *89*(3), 757–771. doi:10.1111/1365-2656.13159

Ficetola, G. F., Lunghi, E., & Manenti, R. (2020). Microhabitat analyses support

relationships between niche breadth and range size when spatial autocorrelation is strong.

Ecography, 724–734. doi:10.1111/ecog.04798

Fieberg, J., & Kochanny, C. O. (2005). Quantifying home-range overlap: The importance of

the utilization distribution. *Journal of Wildlife Management*. doi:10.2193/0022-

541X(2005)69[1346:QHOTIO]2.0.CO;2

Gómez-Rodríguez, C., Baselga, A., & Wiens, J. J. (2015). Is diversification rate related to

climatic niche width? *Global Ecology and Biogeography*, *24*(4), 383–395.

doi:10.1111/geb.12229

Gómez, C., Tenorio, E. A., Montoya, P., & Cadena, C. D. (2016). Niche-tracking migrants

and niche-switching residents: evolution of climatic niches in New World warblers

(Parulidae). *Proceedings of the Royal Society B: Biological Sciences*, *283*(1824),

20152458. doi:10.1098/rspb.2015.2458

Guisan, A., Petitpierre, B., Broennimann, O., Daehler, C., & Kueffer, C. (2014). Unifying

niche shift studies : insights from biological invasions. *Trends in Ecology & Evolution*,

29(5), 260–269. doi:10.1016/j.tree.2014.02.009

Hill, M. P., Hoffmann, A. A., Macfadyen, S., Umina, P. A., & Elith, J. (2012). Understanding

niche shifts: using current and historical data to model the invasive redlegged earth mite,

Halotydeus destructor. *Diversity and Distributions*, *18*(2), 191–203. doi:10.1111/j.1472-

4642.2011.00844.x

Hutchinson, G. E. (1957). Concluding Remarks. *Cold Spring Harbor Symposia on

Quantitative Biology*, *22*, 415–427. doi:10.1101/SQB.1957.022.01.039

Jackson, A. L., Inger, R., Parnell, A. C., & Bearhop, S. (2011). Comparing isotopic niche

widths among and within communities: SIBER - Stable Isotope Bayesian Ellipses in R.

Journal of Animal Ecology, *80*(3), 595–602. doi:10.1111/j.1365-2656.2011.01806.x

Jiménez, L., Soberón, J., Christen, J. A., & Soto, D. (2019). On the problem of modeling a fundamental niche from occurrence data. *Ecological Modelling*, *397*(5), 74–83. doi:10.1016/j.ecolmodel.2019.01.020

Junker, R. R., Kuppler, J., Bathke, A. C., Schreyer, M. L., & Trutschnig, W. (2016). Dynamic range boxes - a robust nonparametric approach to quantify size and overlap of n-dimensional hypervolumes. *Methods in Ecology and Evolution*, (i), 1503–1513. doi:10.1111/2041-210X.12611

La Sorte, F. A., Fink, D., & Johnston, A. (2018). Seasonal associations with novel climates for North American migratory bird populations. *Ecology Letters*, 845–856. doi:10.1111/ele.12951

Laliberté, E., Legendre, P., & Shipley, B. (2015). FD: measuring functional diversity from multiple traits, and other tools for functional ecology. *R Package*.

Lamanna, C., Blonder, B., Violle, C., Kraft, N. J. B., Sandel, B., Šímová, I., … Enquist, B. J. (2014). Functional trait space and the latitudinal diversity gradient. *Proceedings of the National Academy of Sciences of the United States of America*, *111*(38), 13745–50. doi:10.1073/pnas.1317722111

Laughlin, D. C. (2014). The intrinsic dimensionality of plant traits and its relevance to community assembly. *Journal of Ecology*, *102*(1), 186–193. doi:10.1111/1365-2745.12187

Lauzeral, C., Leprieur, F., Beauchard, O., Duron, Q., Oberdorff, T., & Brosse, S. (2011). Identifying climatic niche shifts using coarse-grained occurrence data: a test with non-native freshwater fish. *Global Ecology and Biogeography*, *20*(3), 407–414. doi:10.1111/j.1466-8238.2010.00611.x

Lichstein, J. W. (2007). Multiple regression on distance matrices: A multivariate spatial

analysis tool. *Plant Ecology*, *188*(2), 117–131. doi:10.1007/s11258-006-9126-3

Loiseau, N., Legras, G., Gaertner, J.-C., Verley, P., Chabanet, P., & Mérigot, B. (2017). Performance of partitioning functional beta-diversity indices: Influence of functional representation and partitioning methods. *Global Ecology and Biogeography*, (January), 1–10. doi:10.1111/geb.12581

Lu, R. P., Smith, E. P., & Good, I. J. (1989). Multivariate measures of similarity and niche overlap. *Theoretical Population Biology*, *35*(1), 1–21. doi:10.1016/0040-5809(89)90007-5

MacArthur, R., & Levins, R. (1967). The limiting similarity, convergence, and divergence of coexisting species. *The American Naturalist*, *101*(921), 377–385. doi:10.1086/282505

Mahony, C. R., Cannon, A. J., Wang, T., & Aitken, S. N. (2017). A closer look at novel climates: new methods and insights at continental to landscape scales. *Global Change Biology*, *23*(9), 3934–3955. doi:10.1111/gcb.13645

Mammola, S. (2019). Assessing similarity of n‐ dimensional hypervolumes: Which metric to use? *Journal of Biogeography*, (January), jbi.13618. doi:10.1111/jbi.13618

Mammola, S., & Cardoso, P. (2020). Functional diversity metrics using kernel density n -dimensional hypervolumes . *Methods in Ecology and Evolution*, *2020*(May), 1–10. doi:10.1111/2041-210x.13424

Minami, M., & Shimizu, K. (1999). Estimation of similarity measure for multivariate normal distributions. *Environmental and Ecological Statistics*, *6*(3), 229–248. doi:10.1023/A:1009678412953

Morisita, M. (1959). Measuring of interspecific association and similarity between communities. *Memorandum, Faculty of Science, Kyushu Univ. Ser. E.*, 65–80. doi:10.18960/seitai.11.6_252_4

Nunes, L. A., & Pearson, R. G. (2017). A null biogeographical test for assessing ecological

niche evolution. *Journal of Biogeography*, *44*(6), 1331–1343. doi:10.1111/jbi.12910

Pearman, P. B., Guisan, A., Broennimann, O., & Randin, C. F. (2008). Niche dynamics in space and time. *Trends in Ecology and Evolution*, *23*(3), 149–158. doi:10.1016/j.tree.2007.11.005

Pianka, E. R. (1974). Niche overlap and diffuse competition. *Proceedings of the National Academy of Sciences of the United States of America*. doi:10.1073/pnas.71.5.2141

Pigot, A. L., Sheard, C., Miller, E. T., Bregman, T. P., Freeman, B. G., Roll, U., … Tobias, J. A. (2020). Macroevolutionary convergence connects morphological form to ecological function in birds. *Nature Ecology and Evolution*, *4*(2), 230–239. doi:10.1038/s41559-019-1070-4

Pigot, A. L., Trisos, C. H., & Tobias, J. A. (2016). Functional traits reveal the expansion and packing of ecological niche space underlying an elevational diversity gradient in passerine birds. *Proceedings of the Royal Society B: Biological Sciences*, *283*(1822), 1–9. doi:10.1098/rspb.2015.2013

Pironon, S., Villellas, J., Thuiller, W., Eckhart, V. M., Geber, M. A., Moeller, D. A., & García, M. B. (2018). The 'Hutchinsonian niche' as an assemblage of demographic niches: implications for species geographic ranges. *Ecography*, *41*(7), 1103–1113. doi:10.1111/ecog.03414

Pulliam, H. R. (2000). On the relationship between niche and distribution. *Ecology Letters*, *3*(4), 349–361. doi:10.1046/j.1461-0248.2000.00143.x

Rappoldt, C., & Hogeweg, P. (1980). Niche Packing and Number of Species. *The American Naturalist*, *116*(4), 480–492. doi:10.1086/283643

Read, Q. D., Grady, J. M., Zarnetske, P. L., Record, S., Baiser, B., Belmaker, J., … Thibault, K. M. (2018). Among-species overlap in rodent body size distributions predicts species richness along a temperature gradient. *Ecography*, 1–10. doi:10.1111/ecog.03641

Rolland, J., & Salamin, N. (2016). Niche width impacts vertebrate diversification. *Global Ecology and Biogeography*, 1252–1263. doi:10.1111/geb.12482

Sagarin, R. D., Gaines, S. D., & Gaylord, B. (2006). Moving beyond assumptions to understand abundance distributions across the ranges of species. *Trends in Ecology and Evolution*, *21*(9), 524–530. doi:10.1016/j.tree.2006.06.008

Saupe, E. E., Qiao, H., Hendricks, J. R., Portell, R. W., Hunter, S. J., Soberón, J., & Lieberman, B. S. (2015). Niche breadth and geographic range size as determinants of species survival on geological time scales. *Global Ecology and Biogeography*, *24*(10), 1159–1169. doi:10.1111/geb.12333

Serra-Varela, M. J., Grivet, D., Vincenot, L., Broennimann, O., Gonzalo-Jim??nez, J., & Zimmermann, N. E. (2015). Does phylogeographical structure relate to climatic niche divergence? A test using maritime pine (Pinus pinasterAit.). *Global Ecology and Biogeography*, *24*(11), 1302–1313. doi:10.1111/geb.12369

Slatyer, R. A., Hirst, M., & Sexton, J. P. (2013). Niche breadth predicts geographical range size: A general ecological pattern. *Ecology Letters*, *16*(8), 1104–1114. doi:10.1111/ele.12140

Soberon, J., & Nakamura, M. (2009). Niches and distributional areas: Concepts, methods, and assumptions. *Proceedings of the National Academy of Sciences*, *106*(Supplement_2), 19644–19650. doi:10.1073/pnas.0901637106

Sullivan, B. L., Wood, C. L., Iliff, M. J., Bonney, R. E., Fink, D., & Kelling, S. (2009). eBird: A citizen-based bird observation network in the biological sciences. *Biological Conservation*. doi:10.1016/j.biocon.2009.05.006

Swanson, H. K., Lysy, M., POwer, M., Stasko, A. D., Johnson, J. D., & Reist, J. D. (2015). A new probabilistic method for quantifying n-dimensional ecological niches and niche overlap. *Ecology*, *96*(2), 318–324. doi:10.1890/14-0235.1

Tayleur, C., Caplat, P., Massimino, D., Johnston, A., Jonzén, N., Smith, H. G., & Lindström, Å. (2015). Swedish birds are tracking temperature but not rainfall: Evidence from a decade of abundance changes. *Global Ecology and Biogeography*, *24*(7), 859–872. doi:10.1111/geb.12308

Whittaker, R. H., Levin, S. A., & Root, R. B. (1973). Niche, Habitat, and Ecotope. *The American Naturalist*. doi:10.1086/282837

Wiens, J. J., Litvinenko, Y., Harris, L., & Jezkova, T. (2019). Rapid niche shifts in introduced species can be a million times faster than changes among native species and ten times faster than climate change. *Journal of Biogeography*, (May), jbi.13649. doi:10.1111/jbi.13649

Wilman, H., J., B., J., S., C., de L. R., M., R., & W, J. (2014). EltonTraits 1 . 0 : Species-level foraging attributes of the world ' s birds and mammals. *Ecology*, *95*(October 2013), 2027.

Winner, K., Noonan, M. J., Fleming, C. H., Olson, K. A., Mueller, T., Sheldon, D., & Calabrese, J. M. (2018). Statistical inference for home range overlap. *Methods in Ecology and Evolution*, *9*(7), 1679–1691. doi:10.1111/2041-210X.13027

Zurell, D., Gallien, L., Graham, C. H., & Zimmermann, N. E. (2018). Do long-distance migratory birds track their niche through seasons? *Journal of Biogeography*, *45*(7), 1459–1468. doi:10.1111/jbi.13351

Figure 1. (a) The volume of the 2-dimensional niche measured by the determinant of the covariance matrix of environmental variables, which is the product of three components: two univariate variances (σ_x^2 and σ_y^2) and a correlation component which has the effect of shrinking the volume by a factor of $(1-\rho^2)$. Light blue indicates change of hypervolume. (b) Two commonly used nonparametric niche volume measures: the minimum convex hull and the kernel density estimated hypervolume. The filled area indicates 95% contour of the hypervolume while the dashed blue lines indicate 90% contour of the hypervolumes. (c) Niche dissimilarity of two hypervolumes (orange, blue) measured by the Bhattacharyya distance, which is the sum of the Mahalanobis distance between the two niche centroids and a determinant ratio component measuring the differences of the niche volumes. The ratio MD/DR indicates the relative importance of centroid distance and size difference. (d) Two commonly used dissimilarity measures based on non-parametric volume estimates: Jaccard similarity and Euclidean distance between centroids. White hatches indicate intersecting area between two hypervolumes.

Figure 2. Community-level functional diversity assessment. (a) Component-wise differences between site 1 and site 2 in a) functional trait volume (e.g. functional diversity) and b) Bhattacharyya distance of functional traits. (c)-(f) Change of trait volume and pairwise community-level trait dissimilarity among all 36 sites of annual plant communities with environmental variables in southwest Western Australia. Solid lines are the fitted linear regression. Shaded area represents 95% confidence interval of linear regression. See table 1 for summary statistics.

Figure 3. Species-level environmental niche assessment. (a) Component-wise environmental niche volume differences and (b) Bhattacharyya distance of environmental niche dissimilarity between two example species (White-throated Swift and Tricolored Blackbird) (c)-(f) Change of niche volume and pairwise niche dissimilarity among 260 bird species with range size and centroid latitude. Solid lines are the fitted linear regression. Shaded area represents 95% confidence interval of linear regression. See table 2 for summary statistics.

Figure 4. (a) Correlations among functional alpha-diversity measures for the 36 sites. (b) Correlations among pairwise functional turnover measures. (c) Correlations among environmental niche volume measures for the 260 bird species. (d) Correlations among pairwise environmental niche dissimilarity measures. 'det' stands for the determinant of the environmental niche or functional trait covariance matrix, 'hv' for kernel density derived hypervolume, 'convex' for minimum convex hull, 'FDiv' for functional divergence, 'FDis' for functional dispersion, 'FEve' for functional evenness. 'bd' stands for Bhattacharyya distance, 'bd_mah' for the Mahalanobis distance component of BD, 'bd_det' for the determinant ratio component of BD, 'dst_cent' for Euclidean distance between centroids, 'dst_min' for minimum distance between kernel density derived hypervolumes, 'jac' for Jaccard similarity, 'sor' for Sorensen similarity.

Table 1. Summary statistics for multiple linear regression of community-level trait volumes,

and multiple regression on distance matrices on community-level trait dissimilarity. Specific

leaf area, maximum height, seed mass variances, the correlation component and the

determinant are log-transformed. Adjusted R^2 is used. Only significant predictors are reported.

Response	Predictor	Estimate	p-value	R^2
Specific leaf area variance	--	--	--	--
Maximum height variance	Minimum temperature	0.125	<0.01	0.19
	Moisture availability	0.102	<0.05	
Seed mass variance	Soil surface dispersion	0.066	<0.05	0.25
Correlation component	Moisture availability	0.228	<0.001	0.37
Determinant	Moisture availability	0.369	<0.001	0.27
Mahalanobis distance	Distance (moisture availability)	0.18	<0.001	0.20
	Distance (minimum temperature)	-0.07	<0.05	
Determinant ratio	Distance (moisture availability)	0.08	<0.001	0.13
Bhattacharyya distance	Distance (moisture availability)	0.26	<0.001	0.18

Table 2. Summary statistics for multiple linear regression of environmental niche volumes, and Multiple regression on distance matrices on environmental niche dissimilarity. Area, temperature, precipitation, EVI variances, the correlation component and the determinant are log-transformed. Adjusted R^2 is used. Only significant predictors are reported.

response	predictor	estimate	p-value	R^2
Temperature variance	Area	1.13	<0.001	0.14
	Centroid latitude	0.84	<0.001	
Precipitation variance	Centroid latitude	-5.72	<0.001	0.69
EVI variance	Area	0.70	<0.001	0.13
	Centroid latitude	-0.46	<0.001	
Correlation component	Centroid latitude	-0.67	<0.001	0.22
Determinant	Area	2.09	<0.001	0.55
	Centroid latitude	-5.97	<0.001	
Mahalanobis distance	Distance (Centroid latitude)	32.62	<0.001	0.13
Determinant ratio	Distance (Area)	-2.66	<0.001	0.23
	Distance (Centroid latitude)	9.78	<0.001	
Bhattacharyya distance	Distance (Centroid latitude)	40.36	<0.001	0.15