A unifying framework for quantifying and comparing n-dimensional

hypervolumes

Muyang Lu[1,2], Kevin Winner[1,2], Walter Jetz[1,2*]

[1]Ecology and Evolutionary Biology, Yale University, New Haven, CT, USA

[2]Center for Biodiversity and Global Change, Yale University, New Haven, CT, USA

*Correspondence: walter.jetz@yale.edu (W.J.)

Abstract

1. The quantification of Hutchison's n-dimensional hypervolume has enabled substantial

progress in community ecology, species niche analysis and beyond. While non-

parametric methods for quantifying and comparing hypervolumes are popular, they do

not support a partitioning of the different components and drivers of hypervolume

variation. Here, we propose as alternative the use of multivariate normal distributions

for the assessment and comparison of niche hypervolumes and introduce this as the

multivariate-normal hypervolume (MVNH) framework.

2. The framework provides parametric measures of the size and dissimilarity of niche

hypervolumes, each of which can be partitioned into biologically interpretable

components. Specifically, We use 1) the determinant of the covariance matrix (i.e. the

generalized variance) of a MVNH as a measure of total niche size, which can be

partitioned into the components of univariate niche variances and a correlation

component; and 2) the Bhattacharyya distance between two MVNHs as a measure of

niche dissimilarity, which can be partitioned into the components of Mahalanobis

distance between hypervolume centroids and the determinant ratio which measures

hypervolume size difference.

3. We use empirical examples of community- and species-level analysis to demonstrate

the new insights provided by these metrics. We show that the newly proposed

framework enables us to quantify the relative contributions of different hypervolume

components and to identify the drivers of functional diversity and environmental

niche variation.

4. Our approach overcomes several operational and computational limitations of non-

parametric methods and provides a framework that offers both unification and

granularity in the assessment of niche volumes and differences, which has wide

implications for understanding niche evolution, niche shifts and expansion during

biotic invasions etc.

Keywords: hypervolume; environmental niche; functional diversity; generalized variance;

Bhattacharyya distance; beta diversity; standardized ellipse area

Introduction

The n-dimensional hypervolume is one of the most fundamental (Hutchinson, 1957;

Whittaker, Levin, & Root, 1973; Pulliam, 2000) and commonly used concepts (Díaz et al.,

2015; Blonder, 2018; Pironon et al., 2018) in ecology and evolutionary biology. Hutchinson

first proposed to describe species' niche as an n-dimensional hypervolume in which a species

can survive and reproduce (Hutchinson, 1957). The use of hypervolume was later extended to

describe functional space and trait space (Lamanna et al., 2014; Laughlin, 2014; Pigot et al.,

2020). The geometric features of n-dimensional hypervolumes (especially their size and

dissimilarity) are associated with a wide range of hypotheses and applications in ecology and

evolution. For example, the size of a climatic niche hypervolume is hypothesized to drive

species diversification rates: in mammals, species with small niches (specialists) have been

shown to have higher speciation rates and lower extinction rates than those with large niches

(generalists) (Gómez-Rodríguez, Baselga, & Wiens, 2015; Rolland & Salamin, 2016). The

size of climatic niche hypervolume is also hypothesized to drive the variation of geographic

range size (Slatyer, Hirst, & Sexton, 2013; Saupe et al., 2015; Cardillo, Dinnage, &

McAlister, 2019; Ficetola, Lunghi, & Manenti, 2020). Similarity between species'

environmental niche hypervolumes or functional trait hypervolumes is used to measure niche

divergence or niche packing, which is hypothesized to determine species' coexistence and

species richness patterns (Serra-Varela et al., 2015; Pigot, Trisos, & Tobias, 2016; Castro-

Insua, Gómez-Rodríguez, Wiens, & Baselga, 2018; Read et al., 2018). Niche similarity is

also used for within-species comparisons such as assessing the impact of climate change

(Tayleur et al., 2015; Gómez, Tenorio, Montoya, & Cadena, 2016; Zurell, Gallien, Graham,

& Zimmermann, 2018) and whether there are niche shifts during biotic invasion (Lauzeral et

al., 2011; Early & Sax, 2014; Guisan, Petitpierre, Broennimann, Daehler, & Kueffer, 2014;

Davies, Hill, McGeoch, & Clusella-Trullas, 2019).

The quantification of niche volume and similarity has thus become an urgent pursuit for both

theoretical investigation and real-world applications. Recently, non-parametric methods, such

as kernel-density estimates (Blonder, Lamanna, Violle, & Enquist, 2014; Blonder et al., 2018;

Eckrich, Albeke, Flaherty, Bowyer, & Ben-David, 2020; Mammola & Cardoso, 2020),

dynamic range boxes (Junker, Kuppler, Bathke, Schreyer, & Trutschnig, 2016), minimum

convex hulls, and support vector machines (Blonder et al., 2018), have been widely favored

because of their minimal assumptions around the distribution of data and the existence of

well-documented statistical packages supporting their exploration and use. In these methods,

an n-dimensional distribution over environmental or trait space is typically converted to a

hypervolume with a boundary defined by a particular isopleth (often 95%) of the distribution.

Niche size is calculated as the volume of the enclosed space. Niche similarity of two

hypervolumes can be quantified in multiple ways, either with the union and intersection of

hypervolumes (Jaccard and Sorensen indices) or with the distances between the two

hypervolumes (Loiseau et al., 2017; Mammola, 2019).

These non-parametric methods of quantifying hypervolumes have several methodological

limitations: 1) niche volumes and similarity measures are sensitive to the parameterization of

the underlying distribution and depend additionally on the choice of isopleth (e.g. 95% or

50%), 2) computational costs are high when the dimension of variables and sample size are

large (Blonder et al., 2014), and most importantly 3) the non-parametric methods generally

cannot say anything about the relative importance of the constituent components of the

measures (but see Junker et al., 2016; Loiseau et al., 2017). For example, most non-

parametric methods do not provide information on the impact of each individual dimension

on the size of a hypervolume. For those that do (e.g. dynamic range box; Junker et al., 2016),

the shape information (correlation among variables) on hypervolume size is ignored. Another

example is the use of Jaccard/Sorensen indices and distance metrics for hypervolume

similarity comparison. Though the two classes of metrics (Jaccard and Sorensen indices,

Euclidean distances between niche centroids) are complementary in capturing hypervolume

dissimilarity (Mammola, 2019), no analytical link exists between them for comparing their

relative importance on overall hypervolume similarity. The Jaccard and Sorensen indices are

used to calculate how much hypervolume is shared between the two hypervolumes, which are

more informative when two hypervolumes intersect with each other. On the contrary, the

Euclidean distance between hypervolume (centroid distance or minimum distance) is more

informative when two hypervolumes are disjunct (Mammola, 2019).

The prevalence of non-parametric methods in quantifying hypervolumes also hampers the

integration of empirical niche studies with niche theories because empirical metrics are

disconnected from theoretical derivations. For example, in the theory of limiting similarity,

the outcomes of competition exclusion and niche evolution are typically calculated based on

competition coefficients derived from one-dimensional utilization curves (MacArthur &

Levins, 1967; Leimar, Sasaki, Doebeli, & Dieckmann, 2013). However, the testable

theoretical predictions derived by these studies rely on quantitative analyses in higher

dimensions (Rappoldt & Hogeweg, 1980), meaning that with the prevalent non-parametric

methods, questions such as "Does higher niche similarity among species lead to more

competitive exclusion" could only be studied in a limited or qualitative way (D'Andrea &

Ostling, 2016). Although various parametric dissimilarity metrics have been proposed

(Morisita, 1959; Pianka, 1974; Lu, Smith, & Good, 1989), they were largely ignored because

of their conceptual disconnection with the prevalent size metrics of hypervolume. Parametric

measures that have clear analytical links to theory and which unify both size and dissimilarity

measurements are urgently needed.

To fill in this gap, we propose a framework to quantify and partition niche volume and

dissimilarity based on the assumption of multivariate normal (MVN) distribution of the niche

variables under study and call this the MVN hypervolume or MVNH framework. Different to

currently used non-parametric methods, this partitioning framework provides a powerful

quantitative assessment of relative contributions of the constituent components in driving

total niche variation. We chose the multivariate normal distribution because it is the most

widely used assumption for niche assessment and modeling both in theoretical (MacArthur &

Levins, 1967; Rappoldt & Hogeweg, 1980; Soberon & Nakamura, 2009; Jiménez, Soberón,

Christen, & Soto, 2019) and empirical work (Swanson et al., 2015; La Sorte, Fink, &

Johnston, 2018), which provides room for integration and synthesis of size and dissimilarity

measures (Lu et al., 1989). We show that in the MVN hypervolume framework niche size is

derived from the covariance matrix (Soberon & Nakamura, 2009), and that niche

dissimilarity can be quantified from the Bhattacharyya distance (Bhattacharyya, 1946; Lu et

al., 1989; Winner et al., 2018). We demonstrate through analytical derivation and empirical

examples how the partitioning of the metrics reveals the key drivers of variations in

functional diversity and environmental niche.

Materials and Methods

In our MVNH framework, we propose that a multi-dimensional niche space be described by a

multivariate normal distribution with probability density function:

$f_X(\mathbf{x}) = \dfrac{\exp\left(-\frac{1}{2}(\mathbf{x}-\boldsymbol{\mu})^{\mathrm{T}}\Sigma^{-1}(\mathbf{x}-\boldsymbol{\mu})\right)}{\sqrt{(2\pi)^n|\Sigma|}},$

… (1)

where $\mathbf{x} = \{x_1, \ldots, x_n\}$ is a vector in the n-dimensional environmental niche space, $\mu =$

$\{\mu_1, \ldots, \mu_n\}$ is the environmental niche centroid, Σ is the covariance matrix for the n-

dimensional environmental niche, and $|\Sigma|$ denotes the determinant of the covariance matrix.

The size of a hypervolume

We define the size of a MVN hypervolume as the determinant of the covariance matrix $|\Sigma|$,

also called the "generalized variance" (La Sorte et al., 2018). It has a geometric interpretation

as the volume of the n-dimensional parallelepiped spanned by the column or row vectors of

the covariance matrix. For example, the covariance matrix Σ and the generalized variance $|\Sigma|$

of a two-dimensional MVNH are:

$\Sigma = \begin{bmatrix} \sigma_x{}^2 & \rho_{xy}\sigma_x\sigma_y \\ \rho_{xy}\sigma_x\sigma_y & \sigma_y{}^2 \end{bmatrix},$

… (2)

$|\Sigma| = \sigma_x{}^2\sigma_y{}^2\left(1 - \rho_{xy}{}^2\right),$

… (3)

where $\sigma_x{}^2$ is the variance of variable x, $\sigma_y{}^2$ is the variance of variable y, and ρ_{xy} is the

correlation coefficient between the two environmental variables x and y. When log-

transformed, Equation 3 enables us to additively partition the size of a 2-dimensional

hypervolume (in terms of the determinant) into three separate components (Fig. 1c): the

variance of the first niche axis $\sigma_x{}^2$, the variance of the second niche axis $\sigma_y{}^2$, and a correlation

component $(1-\rho_{xy}{}^2)$. The correlation component could be seen as a shrinkage factor which

shrinks the niche volume by a factor of $(1-\rho_{xy}{}^2)$. Thus, when two niche axes are independent,

the niche size is just the product of the variances of individual axes $\sigma_x{}^2\sigma_y{}^2$. Similarly, for a

three-dimensional hypervolume, the determinant of the covariance matrix is:

$|\Sigma| = \sigma_x{}^2\sigma_y{}^2\sigma_z{}^2\left(1 - \rho_{xy}{}^2 - \rho_{yz}{}^2 - \rho_{xz}{}^2 + 2\rho_{xy}\rho_{yz}\rho_{xz}\right),$

… (4)

where   is the correlation component. The factoring

of the determinant generalizes easily to higher dimensions: the determinant is just the product

of all individual univariate variances and a correlation component.

Figure 1. (a) Two commonly used non-parametric niche volume measures: the minimum

convex hull and the kernel density estimated hypervolume. The filled area indicates 95%

contour of the hypervolume while the dashed blue lines indicate 90% contour of the

hypervolumes. (b) Two commonly used dissimilarity measures based on non-parametric

volume estimates: Jaccard similarity and Euclidean distance between centroids. Orange and

blue represent two hypervolumes. White hatches indicate intersecting area between two

hypervolumes. (c) The volume of the 2-dimensional niche measured by the determinant of

the covariance matrix of environmental variables, which is the product of three components:

137 two univariate variances (σ_x^2 and σ_y^2) and a correlation component which has the effect of

138 shrinking the volume by a factor of $(1-\rho^2)$. Light blue indicates change of hypervolume. (d)

139 Niche dissimilarity of two hypervolumes measured by the Bhattacharyya distance, which is

140 the sum of the Mahalanobis distance between the two niche centroids and a determinant ratio

141 component measuring the differences of the niche volumes. The ratio MD/DR indicates the

142 relative importance of centroid distance and size difference.

144 The determinant also has a close relationship with the standardized ellipse area (SEA, the

145 ellipse with its major and minor axes defined by the principal components; Jackson, Inger,

146 Parnell, & Bearhop, 2011): it scales with the SEA by a constant (π for two-dimensional niche

147 and $4\pi/3$ for three-dimensional niche). Factoring the determinant (as in Equations 3 and 4)

148 can be used to assess the relative importance of the separate components in driving changes

149 in the size of the hypervolume, such as when assessing whether individual trait axis or the

150 constraint on the combination of traits is the major driver of community-wide trait diversity.

152 The dissimilarity of hypervolumes

153 We suggest measuring the dissimilarity of two MVN hypervolumes with the Bhattacharyya

154 distance (Bhattacharyya, 1946; Lu et al., 1989; Minami & Shimizu, 1999; Fieberg &

155 Kochanny, 2005; Winner et al., 2018). The Bhattacharyya distance (BD) of two continuous

156 probability distributions, A and B defined on the same domain $\mathcal{X}$ is:

157 $BD(A,B) = -\log BC(A,B),$

158 … (5)

159 where BC is the Bhattacharyya coefficient:

$$BC(A,B) = \int_{x\in\mathcal{X}} \sqrt{A(x)B(x)}\,dx$$

160 … (6)

Therefore, for two probability distributions, the BC ranges from 0 to 1, and the BD ranges

from 0 to ∞.

For two multivariate normal hypervolumes A and B, the BD has a closed form:

$$BD(A,B) = \underbrace{\frac{1}{8}(\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)^T \overline{\Sigma}^{-1}(\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)}_{MD} + \underbrace{\frac{1}{2}\ln\left(\frac{|\overline{\Sigma}|}{\sqrt{|\Sigma_A||\Sigma_B|}}\right)}_{DR},$$

… (7)

where μ_A and μ_B are the centroids of A and B respectively and $\overline{\Sigma}$ is the mean of the

covariance matrices of A and B:

$$\overline{\Sigma} = \frac{\Sigma_A + \Sigma_B}{2}.$$

… (8)

The first term in Equation 7 is the Mahalanobis distance (MD) between two hypervolume

centroids, and the second term in Equation 7 is the determinant ratio (DR) which measures

size difference between A and B (see Fig. 1d). If two hypervolumes have the same size, then

the second term in Equation 7 will equal 0. As with Equations 3 and 4, factoring the

Bhattacharyya distance according to Equation 7 can be used to assess the relative

contributions of the Mahalanobis distance (centroid difference) and the determinant ratio

(size difference) to the overall niche dissimilarity. We suggest that the ratio of Mahalanobis

distance to determinant ratio (MD/DR), can be used as measure of the relative importance of

centroid distance and size difference in driving overall dissimilarity.

It should also be noticed that the Mahalanobis distance component in Equations 7 could be

linked to an existing partitioning framework which additively partitions the Mahalanobis

distance into standardized Euclidean distances along PCA axes which measure the relative

importance of each PCA axis (Calenge, Darmon, Basille, Loison, & Jullien, 2008; Mahony,

Cannon, Wang, & Aitken, 2017).

Estimators

We used sample means and sample covariances as estimates of μ_A, μ_B, Σ_A, and Σ_B for the

calculation of the determinant of the covariance matrices and the Bhattacharyya distance with

its partitioned components. These plug-in estimates of the Bhattacharyya distance are known

to be biased, but the bias is small with low dimensionality (less than 4) of data (Lu et al.,

1989; Minami & Shimizu, 1999). For the purpose of demonstrating the utility of the

partitioning framework, our empirical analysis proceeds without applying existing bias-

correction methods which have been presented in the literature such as estimating the

covariance matrix using Bayesian methods (Swanson et al., 2015), and bias-correction

techniques for the Bhattacharyya distance (Minami & Shimizu, 1999; Winner et al., 2018).

Calculation of both metrics only requires a few lines of codes in basic R, which is attached in

the supplement (Appendix 1).

We also recognize other dissimilarity metrics based on multivariate normal distributions

which share a similar form with the Bhattacharyya distance and could be partitioned into a

Mahalanobis distance component and a determinant ratio component (see Lu et al., 1989),

such as the MacArthur-Levins measure (MacArthur & Levins, 1967) :

$$MacArthur - Levins(A,B)_A = \underbrace{(\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)^T \overline{\Sigma}^{-1}(\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)}_{MD} + \underbrace{\ln\left(\frac{|\Sigma_A|}{|\Sigma_A||\Sigma_B|}\right)}_{DR},$$

$$MacArthur - Levins(A,B)_B = \underbrace{(\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)^T \overline{\Sigma}^{-1}(\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)}_{MD} + \underbrace{\ln\left(\frac{|\Sigma_B|}{|\Sigma_A||\Sigma_B|}\right)}_{DR},$$

… (9)

the Pianka's measure (Pianka, 1974):

$$Pianka(A,B)_A = \underbrace{(\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)^T \overline{\Sigma}^{-1}(\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)}_{MD} + \underbrace{\frac{1}{2}\ln\left(\frac{|\overline{\Sigma}|}{\sqrt{|\Sigma_A||\Sigma_B|}}\right)}_{DR},$$

… (10)

and the Morisita's measure (Morisita, 1959):

$$Morisita(A, B) = \underbrace{(\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)^T \overline{\boldsymbol{\Sigma}}^{-1} (\boldsymbol{\mu}_A - \boldsymbol{\mu}_B)}_{MD} + \ln\left(\underbrace{\frac{\sqrt{|\overline{\boldsymbol{\Sigma}}|}(\sqrt{|\boldsymbol{\Sigma}_A|}+\sqrt{|\boldsymbol{\Sigma}_B|})^2}{2\sqrt{|\boldsymbol{\Sigma}_A||\boldsymbol{\Sigma}_B|}}}_{DR}\right).$$

… (11)

These metrics differ in their probabilistic interpretations, weighting of each component, and

how they calculate the determinant ratio component. We will focus on the use of the

Bhattacharyya distance in this paper because it was most studied with known statistical

properties (Lu et al., 1989; Minami & Shimizu, 1999; Winner et al., 2018), and also has a

straightforward interpretation as the integration of geometric means of two probability

distributions (Equation 6).

Empirical examples

We will use empirical examples of functional diversity and environmental niche to show the

novel insights provided by the partitioning framework especially in quantifying the relative

importance of univariate variance and correlation in driving size variation; the relative

importance of centroid distance and size difference in driving hypervolume dissimilarity.

Community example: Functional diversity and turnover

We used a trait dataset of 36 annual plant communities in Western Australia (Data availabile

from the Dryad Digital Repository https://doi.org/10.5061/dryad.76kt8 ;Dwyer & Laughlin,

2018) to demonstrate how functional diversity (alpha diversity) and turnover (beta diversity)

can be measured in the MVN hypervolume framework using the determinant and the

Bhattacharyya distance, respectively. The three measured traits for each species were specific

leaf area (SLA), maximum height (MH), and seed mass (SM). We then assessed each

component of the trait volume (SLA variance, MH variance, SM variance and the correlation

component) as predicted by growing season available moisture, mean growing season

minimum temperature, and soil surface dispersion (dispersion is a measure of heterogeneity)

in multiple linear regression. We used the coefficient of variation of each component to

assess the relative contribution of each trait volume component to the variation of total trait

volume, because the individual niche variances are scalable and not directly comparable with

each other. We assessed the environmental drivers of the Bhattacharyya distance (functional

turnover) components using multiple regression on distance matrices (MRM) based on a

partial Mantel test (Lichstein, 2007).The regression coefficients of MRM are calculated with

ordinary least square regression, but the p-values of the coefficients are calculated with

matrix permutation tests. The predictor variables were calculated as the Euclidean distances

for each environmental variable between sites.

Species example: Environmental niche breadth and dissimilarity

We studied 260 bird species in the Western Hemisphere with occurrence data extracted from

eBird (Sullivan et al., 2009), environmental layers obtained from CHELSA (monthly mean

temperature and precipitations for breeding seasons, https://chelsa-climate.org/), and MODIS

EVI (https://modis.gsfc.nasa.gov/data/dataprod/mod13.php) at 1km resolution to assess the

drivers of realized environmental niche breadth (volume) and niche dissimilarity between

species. 1000 random points were sampled and spatially thinned to 10km distance using the

'spThin' package (Aiello-Lammens, Boria, Radosavljevic, Vilela, & Anderson, 2015) in R

for extraction of environmental data. We then investigated the drivers of environmental niche

breadth components using ordinary multiple linear regression, and assessed the drivers of the

Bhattacharyya distance components using MRM. We obtained body mass and dietary niche

breadth data from the 'EltonTrait' dataset (Wilman et al., 2014). Area and centroid absolute

latitude of geographic range were calculated from IUCN range maps

(https://www.iucnredlist.org/resources/spatial-data-download).

Comparisons to non-parametric metrics

For both the community and niche examples we compared the newly proposed size and

dissimilarity metrics from the MVNH framework with measures based on the 95% isopleth

of a box kernel density estimator using the 'hypervolume' package in R (Blonder et al., 2018).

For the species environmental niche example, we also compared the determinant of the

MVNH with the volume of minimum convex hull (Fig. 1a); for the community functional

diversity example, we additionally compared the determinant with three commonly used

functional diversity measures: functional divergence, functional dispersion, and functional

evenness (Laliberté, Legendre, & Shipley, 2015). For both examples, we compared the

Bhattacharyya distance with Jaccard and Sorensen similarities, centroid distances (Fig. 1b)

and minimum distances between hypervolumes.

Results

We use the MVNH framework to firstly analyze drivers of community-level functional

diversity and functional turnover, and to secondly assess drivers of species-level

environmental niche breadth and dissimilarity. Finally, we show the correlations between the

proposed metrics with commonly used non-parametric measures.

Community level: Functional diversity

We compared functional diversity (measured as the determinant of the trait covariance matrix)

and functional turnover (measured as Bhattacharyya distance) among 36 plant communities.

For a two-community example (site 1 and site 2), we partitioned the functional diversity of

each site into four components: the univariate variance of each of the three traits (specific leaf

area, SLA; maximum height, MH; and seed mass, SM) and the correlation component

following Equation 4. Component-wise differences between the two sites add up to the

difference in total trait volume (the determinant of the environmental niche covariance matrix)

on the log-scale (Fig. 2a).

For the 36-community assessment, the coefficient of variation for each trait volume

component across the 36 sites was 0.31 for the correlation component, and varied between

0.28 (MH) and 0.16 (SM) for the individual trait variances. This suggests that the correlation

component is the largest driver of the variation in functional diversity among sites. By

relating individual trait variances and the correlation component to environmental predictors,

we identify mean moisture availability and, less so, minimum temperature as key correlates

of trait diversity variation (Fig. 2c, d). The variances of all three traits differ substantially in

the strength and direction of their respective associations with environmental predictors.

Linear regression shows that none of the examined environmental variables predict the

variation of SLA variance (Table 1). Both minimum temperature and moisture availability

are positively correlated with maximum height variance, whereas only soil surface dispersion

is positively correlated with seed mass variance. Moisture availability is also positively

correlated with the correlation component. The overall trait volume (the determinant) is only

correlated with moisture availability (Table 1).

Table 1. Summary statistics for multiple linear regression of community-level trait volumes,

and multiple regression on distance matrices on community-level trait dissimilarity. Specific

leaf area, maximum height, seed mass variances, the correlation component and the

determinant are log-transformed. Adjusted R^2 is used. Only significant predictors are reported.

Response	Predictor	Estimate	p-value	R^2
Specific leaf area variance	--	--	--	--
Maximum height variance	Minimum temperature	0.125	<0.01	0.19
	Moisture availability	0.102	<0.05	
Seed mass variance	Soil surface dispersion	0.066	<0.05	0.25
Correlation component	Moisture availability	0.228	<0.001	0.37
Determinant	Moisture availability	0.369	<0.001	0.27
Mahalanobis distance	Distance (moisture availability)	0.18	<0.001	0.20
	Distance (minimum temperature)	-0.07	<0.05	
Determinant ratio	Distance (moisture availability)	0.08	<0.001	0.13
Bhattacharyya distance	Distance (moisture availability)	0.26	<0.001	0.18

We measured the pairwise functional turnover between any two sites as the Bhattacharyya

distance between their MVN hypervolumes, and according to Equation 7, partitioned it into

the Mahalanobis distance (which measures the standardized distance between community

means) and the determinant ratio (which measures the differences in functional alpha

diversity) (see a two-community example in Fig. 2b). The ratio of Mahalanobis distance to

determinant ratio (MD/DR) reflects the relative importance of difference in community

means and trait diversity differences in driving the overall community-level functional

turnover. The median ratio MD/DR across all pairs of sites is 0.63, suggesting functional

turnover among communities is mainly driven by the variation in functional alpha-diversity.

The functional turnover is more determined by moisture availability than by minimum

temperature (Fig. 2e, f). Multiple regression on distance matrices suggests that the

Mahalanobis distance is correlated with both minimum temperature and moisture availability,

but the determinant ratio is only correlated with moisture availability (Table 1).

Figure 2. Community-level functional diversity assessment. (a) Component-wise differences

between site 1 and site 2 in a) functional trait volume (e.g. functional diversity) and b)

Bhattacharyya distance of functional traits. (c)-(f) Change of trait volume and pairwise

community-level trait dissimilarity among all 36 sites of annual plant communities with

environmental variables in southwest Western Australia. Solid lines are the fitted linear

regression. Shaded area represents 95% confidence interval of linear regression. See table 1

for summary statistics.

Species level: environmental niches

We used the presented methodology to analyze and compare the three-dimensional

environmental niches (temperature, precipitation, and EVI) of 260 bird species in the New

World. We partitioned the total environmental niche volume for each single species into the

three environmental variable variance components and the correlation component. For a two-

species example, we attributed the difference in niche volume between the white-throated

swift and the tricolored blackbird to differences in temperature variance, precipitation

variance, EVI variance and the correlation component (Fig. 3a). For the 260-species,

assessment, we found that the coefficient of variation for each of the niche volume

components among 260 bird species was 0.79 for precipitation variance, 0.67 for temperature

variance, 0.44 for EVI variance, and 0.22 for the correlation component, suggesting that

precipitation niche breadth variation contributes most to the species-level niche volume

variation. Range size and centroid latitude are the two main drivers of realized environmental

niche volume (Fig. 3c, d), but the predictors for each niche volume component are different:

the precipitation variance and the correlation component are only correlated with the centroid

latitude while the temperature variance and the EVI variance are correlated with both

centroid latitude and range size (Table 2). The median ratio between Mahalanobis distance

and determinant ratio is 2.43, suggesting that the distance between niche centroids is more

important than niche volume differences in driving overall environmental niche dissimilarity

among the 260 bird species. When the multiple regression on distance matrices (MRM) is

applied, different components of the Bhattacharyya distance have different determining

factors: the Mahalanobis distance is only correlated with centroid latitude; while the

determinant ratio is correlated with both range size and centroid latitude (Table 2).

Table 2. Summary statistics for multiple linear regression of environmental niche volumes,

and Multiple regression on distance matrices on environmental niche dissimilarity. Area,

temperature, precipitation, EVI variances, the correlation component and the determinant are

log-transformed. Adjusted R^2 is used. Only significant predictors are reported.

response	predictor	estimate	p-value	R^2
Temperature variance	Area	1.13	<0.001	0.14
	Centroid latitude	0.84	<0.001	
Precipitation variance	Centroid latitude	-5.72	<0.001	0.69
EVI variance	Area	0.70	<0.001	0.13
	Centroid latitude	-0.46	<0.001	
Correlation component	Centroid latitude	-0.67	<0.001	0.22
Determinant	Area	2.09	<0.001	0.55
	Centroid latitude	-5.97	<0.001	
Mahalanobis distance	Distance (Centroid latitude)	32.62	<0.001	0.13
Determinant ratio	Distance (Area)	-2.66	<0.001	0.23
	Distance (Centroid latitude)	9.78	<0.001	
Bhattacharyya distance	Distance (Centroid latitude)	40.36	<0.001	0.15

Figure 3. Species-level environmental niche assessment. (a) Component-wise environmental

niche volume differences and (b) Bhattacharyya distance of environmental niche dissimilarity

between two example species (White-throated Swift and Tricolored Blackbird) (c)-(f)

Change of niche volume and pairwise niche dissimilarity among 260 bird species with range

size and centroid latitude. Solid lines are the fitted linear regression. Shaded area represents

95% confidence interval of linear regression. See table 2 for summary statistics.

Comparison with other metrics

Community-level functional traits: The determinant of trait covariance matrix as the measure

of functional diversity had the highest correlation with the area of the 95% isopleth of a

kernel density derived hypervolume (0.89), followed by functional dispersion (0.79) and

minimum convex hull (0.65) (additional methods in Fig. 4a). The Bhattacharyya distance had

a similar correlation with Euclidean centroid distance (0.70), Jaccard similarity (-0.69) and

Sorensen similarity (-0.71). The Mahalanobis distance component was more correlated with

Euclidean centroid distance (0.73) than with Jaccard similarity (-0.57) or Sorensen similarity

(-0.59), while the determinant ratio component had a relatively low correlation with the

Euclidean centroid distance (0.42; Fig. 4b). All components were uncorrelated with minimum

distance between hypervolumes (Fig. 4b).

Species-level environmental niche: the determinant of the environmental niche covariance

matrix correlates most strongly with the area of the 95% isopleth of the kernel density

derived hypervolume (0.85), followed by the minimum convex hull (0.77, Fig. 4c). The

Bhattacharyya distance and its Mahalanobis distance component are most correlated with the

Euclidean centroid distance (0.85 and 0.82 respectively); the determinant ratio is similarly

correlated with most alternative metrics except for minimum distance between hypervolumes

(Fig. 4d).

Figure 4. (a) Correlations among functional alpha-diversity measures for the 36 sites. (b)

Correlations among pairwise functional turnover measures. (c) Correlations among

environmental niche volume measures for the 260 bird species. (d) Correlations among

pairwise environmental niche dissimilarity measures. 'DET' stands for the determinant of the

environmental niche or functional trait covariance matrix, 'HV' for kernel density derived

hypervolume, 'CONVEX' for minimum convex hull, 'FDiv' for functional divergence, 'FDis'

for functional dispersion, 'FEve' for functional evenness. 'BD' stands for Bhattacharyya

distance, 'BD_MD' for the Mahalanobis distance component of BD, 'BD_DR" for the

determinant ratio component of BD, 'DST_CENT' for Euclidean distance between centroids,

'DST_MIN' for minimum distance between kernel density derived hypervolumes, 'JAC' for

Jaccard similarity, 'SOR' for Sorensen similarity.

Discussion

We proposed a new parametric hypervolume framework based on multivariate normal

distributions and corresponding size and dissimilarity metrics. The greatest advantage of the

MVN hypervolume approach is the analytical clarity and insights provided by the partitioned

components of the niche size and dissimilarity metrics. The MVNH framework is also

significantly faster than the widely used KDE approach. For example, using the 'Iris' dataset

in R (a dataset with 150 rows and 3 columns), it takes 6 seconds to compute the hypervolume

using KDE but only 0.001 seconds to compute the hypervolume using MVNH. Although the

determinant of the covariance matrix has been used as a total niche volume measure in some

studies (Soberon & Nakamura, 2009; La Sorte et al., 2018), it has never been illustrated that

the determinant could be interpreted as the product of individual niche variances and a

correlation component. The Bhattacharyya distance has been used as a measure of home

range overlap in animal movements (Fieberg & Kochanny, 2005; Winner et al., 2018). We

suggest that its use in trait or environmental space (Lu et al., 1989; Minami & Shimizu, 1999),

and in particular the interpretation of the partitioned components, offers valuable biological

insights. Specifically, the Mahalanobis distance between hypervolume centroids can be

interpreted as the standardized distance between optimal traits of two communities or

between the optimal environmental conditions of two species under the assumptions that

mean community traits represents the optimal traits under equilibrium (Blonder et al., 2015)

and that niche centroid has the highest fitness for a species (Sagarin, Gaines, & Gaylord,

2006). The determinant ratio can be interpreted as the functional diversity difference between

communities or as niche volume differences between species. The Bhattacharyya distance

(BD) not only avoids choosing ad hoc algorithm parameters, but more importantly reconciles

the limitations of the two commonly used classes of dissimilarity metrics. When Jaccard or

Sorensen indices become uninformative for disjunct hypervolumes, the BD incorporates the

distance information through the Mahalanobis distance component. When the centroid

distance metric becomes uninformative for two close and overlapping hypervolumes, the BD

incorporates the shape information through the determinant ratio component.

The good concordance between our method and the kernel density method across each of our

studies suggests that the determinant as a measure of niche volume is robust to different

distributions of data. For niche dissimilarity measures, the moderate amount of correlation

between the partitioned components of BD with the Jaccard, Sorensen and the centroid

Euclidean distances shows that the partitioned components of BD both capture a part of the

shape information of the hypervolume. Therefore, our BD metrics combine the important

interpretations of both centroid Euclidean distance metrics and the Jaccard and Sorensen

metrics of niche dissimilarity. More importantly, the relative importance of centroid

difference and volume difference are directly comparable using the BD metrics.

Other metrics such as the MacArthur-Levins measure (Equation 9), the Pianka's measure

(Equation 10) and the Morisita's measure (Equation 11) could be similarly partitioned into a

Mahalanobis distance component and a determinant ratio component (Lu et al., 1989), though

their properties are less well known than the Bhattacharyya distance (Minami & Shimizu,

1999; Winner et al., 2018). The shared Mahalanobis distance component in all these metrics

could be further partitioned into standardized Euclidean distance along different PCA axes

according to a previously proposed partitioning framework (Calenge et al., 2008; Mahony et

al., 2017), which measures the relative importance of each PCA axis in determining the

overall Mahalanobis distance.

Case studies

Through the partitioning framework, our methods provide a novel understanding of each

component driver of community functional diversity or species environmental niche volume.

In the functional diversity example we demonstrated that for the 36 annual plant communities

the constraints on combination of traits is the major driver of functional alpha diversity

variation rather than the change of individual trait variances, which in turn is shaped by

moisture availability; the major driver of functional community turnover among is the

variation of functional alpha-diversity rather than the mean trait differences between

communities. In the species environmental niche example, our method shows that the major

driver of niche volume variation is the precipitation niche breadth variation (Fig. 3d) which is

determined by centroid latitude of species' range. We also show that the range size-niche

breadth relationship of total environmental niche breadth is mainly driven by the range size-

temperature niche breadth and range size-EVI niche breadth relationship (Table 2).

Caveats

A common limitation of all hypervolume metrics is the curse of dimensionality and

collinearity among axes (Blonder et al., 2014; D'Andrea & Ostling, 2016). The use of the

determinant is especially sensitive to collinearity because any pair of fully dependent

columns or rows in a covariance matrix will necessarily result in a determinant of zero.

Increasing dimensions of trait or environmental data will increase the chance of collinearity

and is also more likely to violate the multivariate normal assumption. Such an issue could be

resolved by choosing the biologically relevant variables a priori or using common dimension

reduction techniques applied in non-parametric methods such as PCA (Junker et al., 2016).

Because PCA creates orthogonal axes, the correlation component will always equal 1 if PCA

is applied, and thus it is only recommended when the correlation among axes is of no interest

to the study.

Potential

The presented size and dissimilarity measures are to our knowledge the first parametric

metrics that allow a partitioning of n-dimensional hypervolumes. The partitioning of total

niche volume and niche dissimilarity enables more rigorous testing of ecological and

evolutionary hypotheses beyond those we explored in our two case studies. For example,

when investigating the spatial scaling of niche volume (Pearman, Guisan, Broennimann, &

Randin, 2008; Connor et al., 2018), the scaling relationship could be broken down to the

scaling of univariate niche variances and correlation component, each of which could be

linked to different sets of predictors. When studying niche shifts during biotic invasion

(Lauzeral et al., 2011; Hill, Hoffmann, Macfadyen, Umina, & Elith, 2012; Wiens, Litvinenko,

Harris, & Jezkova, 2019), the BD could be used to determine whether the niche shift is

mainly caused by shifts in niche centroids or caused by volume expansion or reduction; the

determinant could be used to assess which niche axis is most responsible for the total volume

expansion or reduction. In niche evolution studies, the evolution of niche dissimilarity

(Nunes & Pearson, 2017; Castro-Insua et al., 2018) between two species could be traced to

contributions by niche breadth or center differences to the total niche divergence. The

parametric measures also have closer links to theory as theoretical predictions of niche

evolution are usually derived from covariance matrixes with multivariate normal

approximations (Lande & Arnold, 1983; Caetano & Harmon, 2019). In other words, the

newly proposed metrics have the potential to provide more mechanistic understanding of

biodiversity patterns and niche evolution, and to pave the way for more accurate forecasts for

biodiversity change.

Acknowledgement

We thank Jetz Lab members for discussion of the work, especially Richard Li, Aurore

Maureaud and Ben Carlson for comments on the first draft.

Authors' contributions

ML and WJ conceived the ideas. ML analyzed the data with substantial inputs from KW and

WJ. ML led the writing of the manuscript. All authors contributed critically to the drafts and

gave final approval for publication.

Reference

Aiello-Lammens, M. E., Boria, R. A., Radosavljevic, A., Vilela, B., & Anderson, R. P.

(2015). spThin: An R package for spatial thinning of species occurrence records for use

in ecological niche models. *Ecography*, *38*(5), 541–545. doi:10.1111/ecog.01132

Bhattacharyya, A. (1946). On a Measure of Divergence between Two Multinomial

Populations. *Sankhyā: The Indian Journal of Statistics*. doi:10.1038/157869b0

Blonder, B. (2018). Hypervolume concepts in niche- and trait-based ecology. *Ecography*,

41(9), 1441–1455. doi:10.1111/ecog.03187

Blonder, B., Lamanna, C., Violle, C., & Enquist, B. J. (2014). The n-dimensional

hypervolume. *Global Ecology and Biogeography*, *23*(5), 595–609.

doi:10.1111/geb.12146

Blonder, B., Morrow, C. B., Maitner, B., Harris, D. J., Lamanna, C., Violle, C., … Kerkhoff,

A. J. (2018). New approaches for delineating n dimensional hypervolumes. *Methods in*

Ecology and Evolution, *9*(2), 305–319. doi:10.1111/2041-210X.12865

Blonder, B., Nogués-Bravo, D., Borregaard, M. K., Donoghue, J. C., Jørgensen, P. M., Kraft, N. J. B., … Enquist, B. J. (2015). Linking environmental filtering and disequilibrium to biogeography with a community climate framework. *Ecology*, *96*(4), 972–985. doi:10.1890/14-0589.1

Caetano, D. S., & Harmon, L. J. (2019). Estimating correlated rates of trait evolution with uncertainty. *Systematic Biology*, *68*(3), 412–429. doi:10.1093/sysbio/syy067

Calenge, C., Darmon, G., Basille, M., Loison, A., & Jullien, J. M. (2008). The factorial decomposition of the Mahalanobis distances in habitat selection studies. *Ecology*, *89*(2), 555–566. doi:10.1890/06-1750.1

Cardillo, M., Dinnage, R., & McAlister, W. (2019). The relationship between environmental niche breadth and geographic range size across plant species. *Journal of Biogeography*, *46*(1), 97–109. doi:10.1111/jbi.13477

Castro-Insua, A., Gómez-Rodríguez, C., Wiens, J. J., & Baselga, A. (2018). Climatic niche divergence drives patterns of diversification and richness among mammal families. *Scientific Reports*, *8*(1), 8781. doi:10.1038/s41598-018-27068-y

Connor, T., Hull, V., Viña, A., Shortridge, A., Tang, Y., Zhang, J., … Liu, J. (2018). Effects of grain size and niche breadth on species distribution modeling. *Ecography*, *41*(8), 1270–1282. doi:10.1111/ecog.03416

D'Andrea, R., & Ostling, A. (2016). Challenges in linking trait patterns to niche differentiation. *Oikos*, *125*(10), 1369–1385. doi:10.1111/oik.02979

Davies, S. J., Hill, M. P., McGeoch, M. A., & Clusella-Trullas, S. (2019). Niche shift and resource supplementation facilitate an amphibian range expansion. *Diversity and Distributions*, *25*(1), 154–165. doi:10.1111/ddi.12841

Díaz, S., Kattge, J., Cornelissen, J. H. C., Wright, I. J., Lavorel, S., Dray, S., … Gorné, L. D.

(2015). The global spectrum of plant form and function. *Nature*, 1–17.

doi:10.1038/nature16489

Dwyer, J. M., & Laughlin, D. C. (2018). Data from: Constraints on trait combinations explain

climatic drivers of biodiversity: the importance of trait covariance in community

assembly. *Dryad, Dataset*. doi:10.5061/dryad.76kt8

Early, R., & Sax, D. F. (2014). Climatic niche shifts between species' native and naturalized

ranges raise concern for ecological forecasts during invasions and climate change.

Global Ecology and Biogeography, *23*(12), 1356–1365. doi:10.1111/geb.12208

Eckrich, C. A., Albeke, S. E., Flaherty, E. A., Bowyer, R. T., & Ben-David, M. (2020). rKIN:

Kernel-based method for estimating isotopic niche size and overlap. *Journal of Animal*

Ecology, *89*(3), 757–771. doi:10.1111/1365-2656.13159

Ficetola, G. F., Lunghi, E., & Manenti, R. (2020). Microhabitat analyses support

relationships between niche breadth and range size when spatial autocorrelation is strong.

Ecography, 724–734. doi:10.1111/ecog.04798

Fieberg, J., & Kochanny, C. O. (2005). Quantifying home-range overlap: The importance of

the utilization distribution. *Journal of Wildlife Management*. doi:10.2193/0022-

541X(2005)69[1346:QHOTIO]2.0.CO;2

Gómez-Rodríguez, C., Baselga, A., & Wiens, J. J. (2015). Is diversification rate related to

climatic niche width? *Global Ecology and Biogeography*, *24*(4), 383–395.

doi:10.1111/geb.12229

Gómez, C., Tenorio, E. A., Montoya, P., & Cadena, C. D. (2016). Niche-tracking migrants

and niche-switching residents: evolution of climatic niches in New World warblers

(Parulidae). *Proceedings of the Royal Society B: Biological Sciences*, *283*(1824),

20152458. doi:10.1098/rspb.2015.2458

Guisan, A., Petitpierre, B., Broennimann, O., Daehler, C., & Kueffer, C. (2014). Unifying

niche shift studies: insights from biological invasions. *Trends in Ecology & Evolution*,

29(5), 260–269. doi:10.1016/j.tree.2014.02.009

Hill, M. P., Hoffmann, A. A., Macfadyen, S., Umina, P. A., & Elith, J. (2012). Understanding

niche shifts: using current and historical data to model the invasive redlegged earth mite,

Halotydeus destructor. *Diversity and Distributions*, *18*(2), 191–203. doi:10.1111/j.1472-

4642.2011.00844.x

Hutchinson, G. E. (1957). Concluding Remarks. *Cold Spring Harbor Symposia on*

Quantitative Biology, *22*, 415–427. doi:10.1101/SQB.1957.022.01.039

Jackson, A. L., Inger, R., Parnell, A. C., & Bearhop, S. (2011). Comparing isotopic niche

widths among and within communities: SIBER - Stable Isotope Bayesian Ellipses in R.

Journal of Animal Ecology, *80*(3), 595–602. doi:10.1111/j.1365-2656.2011.01806.x

Jiménez, L., Soberón, J., Christen, J. A., & Soto, D. (2019). On the problem of modeling a

fundamental niche from occurrence data. *Ecological Modelling*, *397*(5), 74–83.

doi:10.1016/j.ecolmodel.2019.01.020

Junker, R. R., Kuppler, J., Bathke, A. C., Schreyer, M. L., & Trutschnig, W. (2016). Dynamic

range boxes - a robust nonparametric approach to quantify size and overlap of n-

dimensional hypervolumes. *Methods in Ecology and Evolution*, (i), 1503–1513.

doi:10.1111/2041-210X.12611

La Sorte, F. A., Fink, D., & Johnston, A. (2018). Seasonal associations with novel climates

for North American migratory bird populations. *Ecology Letters*, 845–856.

doi:10.1111/ele.12951

Laliberté, E., Legendre, P., & Shipley, B. (2015). FD: measuring functional diversity from

multiple traits, and other tools for functional ecology. *R Package*.

Lamanna, C., Blonder, B., Violle, C., Kraft, N. J. B., Sandel, B., Šímová, I., … Enquist, B. J.

(2014). Functional trait space and the latitudinal diversity gradient. *Proceedings of the*

National Academy of Sciences of the United States of America, *111*(38), 13745–50.

doi:10.1073/pnas.1317722111

Lande, R., & Arnold, S. J. (1983). The measurement of selection on correlated characters.

Evolution, *37*(6), 1210. doi:10.2307/2408842

Laughlin, D. C. (2014). The intrinsic dimensionality of plant traits and its relevance to

community assembly. *Journal of Ecology*, *102*(1), 186–193. doi:10.1111/1365-

2745.12187

Lauzeral, C., Leprieur, F., Beauchard, O., Duron, Q., Oberdorff, T., & Brosse, S. (2011).

Identifying climatic niche shifts using coarse-grained occurrence data: a test with non-

native freshwater fish. *Global Ecology and Biogeography*, *20*(3), 407–414.

doi:10.1111/j.1466-8238.2010.00611.x

Leimar, O., Sasaki, A., Doebeli, M., & Dieckmann, U. (2013). Limiting similarity, species

packing, and the shape of competition kernels. *Journal of Theoretical Biology*, *339*, 3–

13. doi:10.1016/j.jtbi.2013.08.005

Lichstein, J. W. (2007). Multiple regression on distance matrices: A multivariate spatial

analysis tool. *Plant Ecology*, *188*(2), 117–131. doi:10.1007/s11258-006-9126-3

Loiseau, N., Legras, G., Gaertner, J.-C., Verley, P., Chabanet, P., & Mérigot, B. (2017).

Performance of partitioning functional beta-diversity indices: Influence of functional

representation and partitioning methods. *Global Ecology and Biogeography*, *26*(6), 753–

762. doi:10.1111/geb.12581

Lu, R. P., Smith, E. P., & Good, I. J. (1989). Multivariate measures of similarity and niche

overlap. *Theoretical Population Biology*, *35*(1), 1–21. doi:10.1016/0040-

5809(89)90007-5

MacArthur, R., & Levins, R. (1967). The limiting similarity, convergence, and divergence of

coexisting species. *The American Naturalist*, *101*(921), 377–385. doi:10.1086/282505

Mahony, C. R., Cannon, A. J., Wang, T., & Aitken, S. N. (2017). A closer look at novel

climates: new methods and insights at continental to landscape scales. *Global Change*

Biology, *23*(9), 3934–3955. doi:10.1111/gcb.13645

Mammola, S. (2019). Assessing similarity of n☐ dimensional hypervolumes: Which metric

to use? *Journal of Biogeography*, *46*(9), 2012–2023. doi:10.1111/jbi.13618

Mammola, S., & Cardoso, P. (2020). Functional diversity metrics using kernel density n

☐dimensional hypervolumes. *Methods in Ecology and Evolution*, *11*(8), 986–995.

doi:10.1111/2041-210X.13424

Minami, M., & Shimizu, K. (1999). Estimation of similarity measure for multivariate normal

distributions. *Environmental and Ecological Statistics*, *6*(3), 229–248.

doi:10.1023/A:1009678412953

Morisita, M. (1959). Measuring of interspecific association and similarity between

communities. *Memorandum, Faculty of Science, Kyushu Univ. Ser. E.*, 65–80.

doi:10.18960/seitai.11.6_252_4

Nunes, L. A., & Pearson, R. G. (2017). A null biogeographical test for assessing ecological

niche evolution. *Journal of Biogeography*, *44*(6), 1331–1343. doi:10.1111/jbi.12910

Pearman, P. B., Guisan, A., Broennimann, O., & Randin, C. F. (2008). Niche dynamics in

space and time. *Trends in Ecology and Evolution*, *23*(3), 149–158.

doi:10.1016/j.tree.2007.11.005

Pianka, E. R. (1974). Niche overlap and diffuse competition. *Proceedings of the National*

Academy of Sciences of the United States of America. doi:10.1073/pnas.71.5.2141

Pigot, A. L., Sheard, C., Miller, E. T., Bregman, T. P., Freeman, B. G., Roll, U., … Tobias, J.

A. (2020). Macroevolutionary convergence connects morphological form to ecological

function in birds. *Nature Ecology and Evolution*, *4*(2), 230–239. doi:10.1038/s41559-

019-1070-4

Pigot, A. L., Trisos, C. H., & Tobias, J. A. (2016). Functional traits reveal the expansion and

packing of ecological niche space underlying an elevational diversity gradient in

passerine birds. *Proceedings of the Royal Society B: Biological Sciences*, *283*(1822), 1–

9. doi:10.1098/rspb.2015.2013

Pironon, S., Villellas, J., Thuiller, W., Eckhart, V. M., Geber, M. A., Moeller, D. A., &

García, M. B. (2018). The 'Hutchinsonian niche' as an assemblage of demographic

niches: implications for species geographic ranges. *Ecography*, *41*(7), 1103–1113.

doi:10.1111/ecog.03414

Pulliam, H. R. (2000). On the relationship between niche and distribution. *Ecology Letters*,

3(4), 349–361. doi:10.1046/j.1461-0248.2000.00143.x

Rappoldt, C., & Hogeweg, P. (1980). Niche Packing and Number of Species. *The American*

Naturalist, *116*(4), 480–492. doi:10.1086/283643

Read, Q. D., Grady, J. M., Zarnetske, P. L., Record, S., Baiser, B., Belmaker, J., … Thibault,

K. M. (2018). Among-species overlap in rodent body size distributions predicts species

richness along a temperature gradient. *Ecography*, 1–10. doi:10.1111/ecog.03641

Rolland, J., & Salamin, N. (2016). Niche width impacts vertebrate diversification. *Global*

Ecology and Biogeography, 1252–1263. doi:10.1111/geb.12482

Sagarin, R. D., Gaines, S. D., & Gaylord, B. (2006). Moving beyond assumptions to

understand abundance distributions across the ranges of species. *Trends in Ecology and*

Evolution, *21*(9), 524–530. doi:10.1016/j.tree.2006.06.008

Saupe, E. E., Qiao, H., Hendricks, J. R., Portell, R. W., Hunter, S. J., Soberón, J., &

Lieberman, B. S. (2015). Niche breadth and geographic range size as determinants of

species survival on geological time scales. *Global Ecology and Biogeography*, *24*(10),

1159–1169. doi:10.1111/geb.12333

Serra-Varela, M. J., Grivet, D., Vincenot, L., Broennimann, O., Gonzalo-Jim??nez, J., &

Zimmermann, N. E. (2015). Does phylogeographical structure relate to climatic niche

divergence? A test using maritime pine (Pinus pinasterAit.). *Global Ecology and*

Biogeography, *24*(11), 1302–1313. doi:10.1111/geb.12369

Slatyer, R. A., Hirst, M., & Sexton, J. P. (2013). Niche breadth predicts geographical range

size: A general ecological pattern. *Ecology Letters*, *16*(8), 1104–1114.

doi:10.1111/ele.12140

Soberon, J., & Nakamura, M. (2009). Niches and distributional areas: Concepts, methods,

and assumptions. *Proceedings of the National Academy of Sciences*, *106*(Supplement_2),

19644–19650. doi:10.1073/pnas.0901637106

Sullivan, B. L., Wood, C. L., Iliff, M. J., Bonney, R. E., Fink, D., & Kelling, S. (2009). eBird:

A citizen-based bird observation network in the biological sciences. *Biological*

Conservation. doi:10.1016/j.biocon.2009.05.006

Swanson, H. K., Lysy, M., POwer, M., Stasko, A. D., Johnson, J. D., & Reist, J. D. (2015). A

new probabilistic method for quantifying n-dimensional ecological niches and niche

overlap. *Ecology*, *96*(2), 318–324. doi:10.1890/14-0235.1

Tayleur, C., Caplat, P., Massimino, D., Johnston, A., Jonzén, N., Smith, H. G., & Lindström,

Å. (2015). Swedish birds are tracking temperature but not rainfall: Evidence from a

decade of abundance changes. *Global Ecology and Biogeography*, *24*(7), 859–872.

doi:10.1111/geb.12308

Whittaker, R. H., Levin, S. A., & Root, R. B. (1973). Niche, Habitat, and Ecotope. *The*

American Naturalist. doi:10.1086/282837

Wiens, J. J., Litvinenko, Y., Harris, L., & Jezkova, T. (2019). Rapid niche shifts in

introduced species can be a million times faster than changes among native species and

ten times faster than climate change. *Journal of Biogeography*, *46*(9), 2115–2125.

doi:10.1111/jbi.13649

Wilman, H., Belmaker, J., Simpson, J., de la Rosa, C., Rivadeneira, M. M., & Jetz, W. (2014).

EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals.

Ecology, *95*(7), 2027–2027. doi:10.1890/13-1917.1

Winner, K., Noonan, M. J., Fleming, C. H., Olson, K. A., Mueller, T., Sheldon, D., &

Calabrese, J. M. (2018). Statistical inference for home range overlap. *Methods in*

Ecology and Evolution, *9*(7), 1679–1691. doi:10.1111/2041-210X.13027

Zurell, D., Gallien, L., Graham, C. H., & Zimmermann, N. E. (2018). Do long-distance

migratory birds track their niche through seasons? *Journal of Biogeography*, *45*(7),

1459–1468. doi:10.1111/jbi.13351

706