ER bodies are induced by *Pseudomonas syringae* and negatively regulate

immunity

Running title: Role of ER bodies in the defense response

José S. Rufián[1,2,3], James M. Elmore[2,4], Eduardo R. Bejarano[1], Carmen R.

Beuzon[1], Gitta L. Coaker[2*]

[1] Instituto de Hortofruticultura Subtropical y Mediterránea, Universidad de

Málaga-Consejo Superior de Investigaciones Científicas (IHSM-UMA-CSIC),

Depto. Biología Celular, Genética y Fisiología, Campus de Teatinos, Málaga

E-29071, Spain

[2] Department of Plant Pathology, University of California Davis, Davis, CA

95616 USA

[3] Shanghai Center for Plant Stress Biology, CAS Center for Excellence in

Molecular Plant Sciences; Shanghai Institutes of Biological Sciences, Chinese

Academy of Sciences, Shanghai 201602, China

[4] Current address: Department of Plant Pathology and Microbiology, Iowa State

University, Ames, IA 5001 USA

*Address correspondence to: glcoaker@ucdavis.edu

ABSTRACT

ER bodies are endoplasmic reticulum-derived organelles present in plants

belonging to the *Brassicales* order. In *Arabidopsis thaliana*, ER bodies are

ubiquitous in cotyledons and roots, and present only in certain cell types in

rosette leaves. However, both wounding and jasmonic acid treatment induce

the formation of ER bodies in leaves. Formation of this structure is dependent

on the transcription factor *NAI1*. The main components of the ER bodies are β-

glucosidases (BGLUs), enzymes that hydrolyze specialized compounds. In

Arabidopsis, PYK10 (BGLU23) and BGLU18 are the most abundant ER body

proteins. In this work, we found that ER bodies are downregulated as a

consequence of the immune responses induced by bacterial flagellin

perception. *Arabidopsis* mutants defective in ER body formation show

enhanced responses upon flagellin perception and enhanced resistance to

bacterial infections. Furthermore, the bacterial toxin coronatine induces the

formation of *de novo* ER bodies in leaves and its virulence function is partially

dependent on this structure. Finally, we show that performance of the

polyphagous beet armyworm herbivore, *Spodoptera exigua*, increases in plants

lacking ER bodies. Altogether, we provide new evidence for the role of the ER

bodies in plant immune responses.

Keywords: ER bodies, plant immunity, *Pseudomonas syringae*, *Spodoptera*

exigua,

INTRODUCTION

ER bodies are rod-shaped endoplasmic reticulum (ER)-associated structures that are present in three families within the Brassicales order (Nakano et al., 2014a). ER bodies are constitutively present in roots and cotyledons of young plants (constitutive ER bodies), but their abundance decreases with senescence (Matsushima et al., 2002). In *Arabidopsis* rosette leaves, ER bodies are constitutively present in marginal cells, epidermal cells covering the midrib and extremely large pavement cells (Leaf ER bodies) (Nakazaki et al., 2019b). Furthermore, the formation of ER bodies in rosette leaves is induced by Methyl Jasmonate (MeJA) treatment or wounding (Inducible ER bodies) (Matsushima et al., 2002).

Formation of both constitutive and leaf ER bodies is dependent on the basic-helix-loop-helix transcription factor *NAI1* (Matsushima et al., 2004) and the essential protein *NAI2* (Yamada et al., 2008). Inducible ER bodies are formed in a *nai1* mutant, but differin shape and composition with the ER bodies induced in wild-type Col-0 (Ogasawara et al., 2009). The main components of the constitutive ER bodies are the β-glucosidases (BGLU) PYK10 (BGLU23), BGLU21 and BGLU22 (Matsushima et al., 2003), while inducible-ER bodies also contain BGLU18 (Ogasawara et al., 2009). These myrosinases belong to a subfamily of 8 β-glucosidases (*i. e.* BGLU18 to BGLU25) (Xu et al., 2004) with ER-retention signals (Yamada et al., 2011). Leaf ER bodies mainly contain PYK10 and BGLU18 (Nakazaki et al., 2019b). PYK10 has been described as an atypical myrosinase that hydrolyzes Indole Glucosinolates (IG) (Nakano et al., 2017). Upon tissue damage, PYK10 associates in the cytosol with PYK10-binding protein 1 (PBP1), activating PYK10 function (Nagano et al., 2005). An

Arabidopsis pyk10/blgu18 double mutant accumulates high amounts of the IG

4-methoxyindol-3-ylmethylglucosinolate (4MI3G) after tissue damage (Nakazaki

et al., 2019b). IGs and the products resulting from their breakdown have a role

in the defense responses against microbial pathogens and herbivory (Bednarek

et al., 2009; Clay et al., 2009; Millet et al., 2010; Müller et al., 2010; Johansson

et al., 2014; Nakazaki et al., 2019b). IGs are also required for the establishment

of beneficial interactions with the endophytic fungus *Piriformospora indica* by

appropriately regulating the level of root colonization through *PYK10* (Sherameti

et al., 2008; Lahrmann et al., 2015). Collectively, multiple independent studies

have supported a role in defense responses for ER bodies (Yamada et al.,

2011; Nakano et al., 2014b).

Plants are constantly exposed to a variety of biotic and abiotic stresses.

Surface localized plant pattern-recognition receptors (PRRs) act as one of the

first lines of defense against pathogens and are able to recognize conserved

pathogen features, such as flagellin, as non-self (Couto and Zipfel, 2016). One

of the best-characterized PRRs is the flagellin receptor FLS2 (Flagellin Sensing

2) which recognizes flg22, a 22 amino acid epitope of bacterial flagellin

(Gómez-Gómez and Boller, 2000). Perception of flg22 by FLS2 induces a

number of cellular events resulting in Pattern-Triggered Immunity (PTI), which

restricts pathogen proliferation. Some of the cellular events that take place

during PTI are the activation of MAP Kinases (MAPK) and calcium-dependent

protein kinases, the production of reactive oxygen species (ROS), stomatal

closure and the formation of callose (β-1,3-glucan) deposits in the cell wall

(Henry et al., 2013). However, successful pathogens have evolved different

mechanisms to suppress defense responses and successfully colonize their
host.

Pseudomonas syringae is a Gram-negative bacterium that has been
extensively used to study plant immune signaling. *P. syringae* virulence
depends on the Type III Secretion System (T3SS), a complex nanomachine that
translocates proteins, called effectors, into the cytosol of the plant cell. Most of
the effectors suppress PTI acting on different cellular pathways (Macho, 2015).
Another virulence mechanisms used by bacterial pathogens to subvert plant
immunity is the production of phytotoxins. *P. syringae* pv. tomato DC3000 (*Pst*
DC3000) produces coronatine, a polyketide toxin that acts as a molecular mimic
of the plant hormone jasmonic acid (JA) (Weiler et al., 1994). Coronatine
induces stomata re-opening upon PRR perception to allow bacterial
colonization (Melotto et al., 2006; Melotto et al., 2008; Gimenez-Ibanez et al.,
2017), and also suppresses callose deposition and induces bacterial growth
within the apoplast (Geng et al., 2012).

In this work, we investigated the biological function of ER bodies during
plant immunity. We found that ER bodies are downregulated after flg22-
triggered immune activation but induced after *P. syringae* infection in a
coronatine-dependent manner. Furthermore, loci underlying ER body formation
act as negative regulators of immunity against bacterial pathogens but are
required for a complete response against the chewing insect *Spodoptera*
exigua. With these results, we report a new role of ER bodies in virulence of
phytopathogenic bacteria and their suppression by plant immune perception..

RESULTS

ER bodies are rapidly inhibited upon flagellin perception

To investigate the role of ER body formation after plant pathogen perception,

we used an *Arabidopsis* transgenic line expressing GFP with the ER-retention

signal HDEL (GFP-HDEL), which allows the visualization of ER bodies

(Mitsuhashi et al., 2000). We vacuum-infiltrated 2 week-old *Arabidopsis* plants

with either 10 µM flg22 or water and visualized cotyledons three hours later by

confocal microscopy. We were able to detect and quantify the ER bodies in

cotyledons, where this structure is constitutively present, in mock-treated plants.

However, after flg22 treatment, only a small number of ER bodies were

detected (Fig. 1A). This decrease in ER body detection was statistically different

compared to water-treated controls (Fig. 1A). To further investigate the

disappearance of ER bodies within the same tissue, we monitored cotyledons

for 120 minutes after either water (mock) or flg22 treatment. ER bodies lose

their characteristic rod-shape within 60 minutes post-flg22 treatment, and the

GFP signal is diffused throughout the cell at later time points (Fig. 1B).

To analyze the impact of the flg22 treatment in ER body formation at the

transcriptional level, we analyzed the transcriptional expression of *NAI1* and

PYK10 under different biotic treatments using an available dataset

(NASCArrays-120). We found that flg22 treatment reduces the expression level

of *NAI1* and *PYK10* in an *FLS2*-dependent manner (Fig. 1C, Fig. S1), pointing

to regulation of ER bodies at the transcriptional level as a consequence of

activation of defense responses. However, inoculation with the virulent bacterial

strain *Pst* DC3000 induces the expression of both *NAI1* and *PYK10* at 24h, and

the same occurs with a strain expressing the avirulence effector AvrRpm1 (Fig.

1C). Interestingly, both the T3SS DC3000 mutant derivative Δ*hrcC* and the non-

host strain 1448A reduced the accumulation of *NAI1* transcripts (Fig. 1C).

Altogether, these results show a downregulation of ER bodies as a

consequence of the activation of flg22-triggered immunity and suggest a role of

this subcellular structure during the defense response against *P. syringae*.

***Arabidopsis* mutants impacting ER body formation and contents exhibit**

enhanced immune responses against bacterial pathogens

After flg22 perception, many cellular events take place to restrict pathogen

invasion, including activation of MAPKs and calcium dependent protein kinases

minutes after perception, followed by ROS production, activation of defense

gene expression and several hours later, callose deposition (Henry et al.,

2013). To further characterize the role of the ER bodies in flg22-induced

responses, we measured ROS production as an early event of PTI, in plants

affected in the formation of ER bodies. We treated 3-week-old Col-0, *nai1-1,*

pyk10-1 and *pyk10-1/blgu21* plants, with 100 nM flg22 and measured the ROS

burst. The total ROS produced in all the mutant backgrounds tested was higher

than the ROS produced in Col-0 (Fig. 2A). This increase in ROS production in

response to flg22 also occurs in both *bglu18-1* and *pbp1-1* mutants (Fig S2A).

To analyze a late event of PTI, we treated four-week-old plants with flg22 and

quantified callose deposition. Consistent with the ROS burst, the number of

callose deposits was two times higher in ER bodies mutants compared to Col-0

(Fig 2B). These data indicate that a loss of ER bodies or its main components

induce an enhanced response to bacterial flg22, pointing to a negative role of

ER bodies on PTI.

To determine the role of the ER bodies in bacterial growth, we quantified *Pst*

DC3000 titers in Col-0 and in the different mutant backgrounds (Fig. 2C). *Pst*

DC3000 exhibited decreased growth (>1 log) in the *nai1-1* mutant compared to

Col-0. Interestingly, both the *pyk10-1* and the *pyk10-1/bglu21* mutants exhibited

a stronger decrease in bacterial growth, with 1.5 Log reduction compared to

wild-type Col-0 (Fig 2C). No differences were detected between *pyk10* and

pyk10-1/bglu21, indicating that *PYK10* is a critical gene involved in negative

regulation of bacterial growth. Plants lacking *PBP1*, necessary for PYK10

function, or *BGLU18*, the main component of inducible ER bodies, also showed

enhanced resistance to bacterial infection (Fig, S2B).

A critical component controlling bacterial virulence is the ability to deliver

effectors through the type III secretion system (Macho, 2015). In order to

investigate the relationship between immune suppression and ER body

formation, we used DC3000 Δ*hrcC*, which is unable to translocate effectors to

the plant cell, and thus, does not suppress PTI. We measured the growth of

Δ*hrcC* in ER body mutants. In the *nai1-1* background, the growth of Δ*hrcC* was

not significantly different from that in Col-0. However, we observed a slight but

significant decrease in growth in the *pyk10-1/bglu21* background (Fig. 2D). The

increased *P. syringae* growth restriction in *pyk10-1* and *pyk10-1/bglu21*

backgrounds compared with *nai1* (Fig. 2C and 2D) could indicate a NAI1-

independent regulation for *PYK10* and/or *BGLU21*. Altogether, these results

show that the absence of ER bodies increases resistance against the bacterial

pathogen *P. syringae.*

NAI1 contributes to the defense response against *S. exigua*

ER bodies have been proposed to participate in the defense response against

herbivores (Yamada et al., 2011; Nakano et al., 2014b). A recent study

demonstrated increased susceptibility of the double mutant *bglu18 pyk10* to the

terrestrial isopod *Armadillidium vulgare* (Nakazaki et al., 2019b). The

pyk10/bglu21 mutant forms ER bodies, although they are larger and less

numerous than those in wild-type plants (Nagano et al. 2009). We measured

the performance of the generalist leaf-chewing herbivore *S. exigua* in different

ER body related mutants. Performance assays have been widely used to study

the defense mechanisms against herbivory (Cipollini et al., 2004; Chung et al.,

2008; Van Oosten et al., 2008; Müller et al., 2010; Santamaría et al., 2017).

We fed *S. exigua* larvae with Col-0 and the ER body mutants *nai1-1*, *pyk10-1*

and *pyk10-1/bglu21* plants, and measured the weight of the larvae after 7 days.

The fresh weight of *S. exigua* larvae was higher in *nai1-1* plants compared with

the Col-0 control, confirming a role of ER bodies in the defense response

against herbivore insects. Surprisingly, the performance of *S. exigua* in either

pyk10-1 or *pyk10-1/bglu21* plants was not different compared Col-0, suggesting

a redundant role of *PYK10* and *BGLU21* with other myrosinases in this context.

(Fig. 2E)

Coronatine induces the formation of ER bodies in rosette leaves

Transcriptomic analyses indicate that inoculation with virulent *Pst* DC3000

induces the expression of *NAI1* and *PYK10* (Fig. 1C). To check if this increased

expression results in *de novo* formation of ER bodies, we inoculated rosette

leaves of *Arabidopsis* GFP-HDEL plants with a relatively high inoculum (5×10^7

CFU/mL) of *Pst* DC3000, the derivative Δ*hrcC* and a coronatine mutant (*cor-*,

Pst DC3118). We applied the bacterial suspension using a brush to avoid

wounding tissue. Three days after inoculation, we observed the samples under

the confocal microscope (Fig. 3A). No ER bodies were found in the mock-

treated leaves, in agreement with the absence of ER bodies in certain rosette

leaves (Matsushima et al., 2002; Nakazaki et al., 2019b). Interestingly, a large

number of ER bodies was found in leaves inoculated with wild-type *Pst*

DC3000. The distribution of the ER bodies was patchy, suggesting that only

some cells were forming the structure. To investigate if the formation of ER

bodies localized surrounding bacterial colonies, we generated a *Pst* DC3000

derivative tagged with eYFP. Three days after inoculation, we observed a large

number of ER bodies surrounding the bacterial microcolony in Col-0 (Fig. 3B,C)

but only a few aberrant ER bodies in the *nai1-1* mutant (Fig. 3C), suggesting

that ER bodies induced by *Pst* DC3000 are dependent on *NAI1*. To determine

which bacterial factors are inducing the formation of ER bodies in rosette

leaves, we inoculated Col-0 GFP-HDEL leaves with the Δ*hrcC* derivative and

the *cor-* mutant, both tagged with eYFP. We were unable to detect Δ*hrcC* –

eYFP microcolonies due to their extremely small size, consistent with their

reduced growth *in planta*. Nevertheless, we did not detect ER bodies in leaves

inoculated with the Δ*hrcC* mutant (Fig. 3A). Interestingly, the *cor-* eYFP mutant

was unable to induce the formation of *de novo* ER bodies, consistent with the

role of coronatine mimicking JA (Fig. 3). This result indicates that induction of

ER bodies could be a virulence function of coronatine. Analyzing transcriptomic

data of plants inoculated either with *Pst* DC3000 or the *cor-* mutant, we

observed that many genes related to ER bodies, including *NAI1*, *PYK10* and

BGLU18, were upregulated by *Pst* DC3000 but not after inoculation with *Pst*

DC3118 (*cor-*) (Fig. S3). To investigate the ability of coronatine to promote

bacterial virulence in the absence of wild-type ER body formation, we measured

the growth of *Pst* DC3000 and the *Pst cor-* mutant in Col-0, *nai1-1*, *pyk10-1* and

pyk10-1/bglu21. In order to bypass the role of coronatine on re-opening stomata

(Melotto et al., 2006; Melotto et al., 2008), we performed syringe inoculation in

five-week old plant leaves. Four days after inoculation, a strong growth

restriction was observed in the coronatine mutant compared with wild-type *Pst*

DC3000 in Col-0 (Fig. 3D). In the ER body-defective backgrounds, a stronger

growth restriction was observed for *Pst* DC3000 (Fig. 2C, Fig. 3D). However,

the growth of the coronatine mutant was not significantly different between wild-

type Col-0 and the ER body mutants.

DISCUSSION

The main contents in ER bodies are β -glucosidases, enzymes which can

produce toxic compounds to protect the plant from herbivore attack

(Matsushima et al., 2003). Induction of ER bodies by wounding or by MeJA

treatment suggests an association with plant defense (Nakano et al., 2014b).

However, there are few experimental evidences of the association between ER

bodies and plant immunity. Plants are infected by diverse pathogens that often

utilize opposing virulence strategies and are differentially impacted by plant

immune responses (e.g. chewing insects versus biotrophic/hemibiotrophic

pathogens). In this study we demonstrate contrasting roles of ER bodies in the

plant response to pathogenic bacteria and herbivores.

Previous studies have implicated ER bodies in plant protection against fungi

and feeding insects. Sherameti et al. (2008) demonstrated that *PYK10* is

required to protect plants against excessive root colonization by the endophytic

fungus *Piriformospora indica*. Furthermore, Nakazaki et al. (2019b) showed that

bglu18 pyk10 double mutant plants are more susceptible than wild-type to the

attack of the terrestrial isopod *Armadillidium vulgare* (pill-bug). In this work, we

experimentally link the function of the ER bodies with the plant response

against the bacterial pathogen *P. syringae* and the herbivore *S. exigua*. We

found an increased weight acquisition of the generalist chewing herbivore *S.*

exigua in *nai1-1* plants compared with Col-0, but not in *pyk10-1* and *pyk10-*

1/bglu21 mutants (Fig. 2E). *S. exigua* feeds on leaves, where constitutive ER

bodies are absent (Matsushima et al., 2002). Constitutive leaf ER bodies, which

are dependent on *NAI1*, localize in the edges of the leaves and mainly contain

PYK10 and BGLU18 (Nakazaki et al., 2019b). The biogenesis and content of

leaf ER bodies is regulated by both Jasmonic Acid (JA) dependent and

independent pathways (Nakazaki et al., 2019a). While *NAI1* expression, and

therefore PYK10 accumulation is dependent on JA, BGLU18 accumulation in

leaf ER bodies is independent on JA (Nakazaki et al., 2019a). Since the

performance of *S. exigua* is not affected in *pyk10-1* mutant, the genetic analysis

points to *BGLU18* as the responsible of the defense response against *S.*

exigua. Furthermore, chewing induces JA accumulation (Rehrig et al., 2014)

and this signal would potentially lead to the formation of the inducible ER

bodies. The *nai1* mutant is compromised in formation of inducible local and

systemic ER bodies. In the *nai1* mutant, inducible ER bodies are long and

tubular and mainly observed in cells surrounding the wound site (Ogasawara et

al., 2009). We also observed that *Pst* DC3000 was unable to induce ER body

formation in the *nai1* mutant, indicating that the absence or altered

composition/structure of ER bodies in *NAI1* suppresses growth of the

hemibiotrophic pathogen *P. syringae*, while enhancing performance of a

chewing herbivore.

Pst DC3000 induces the formation of ER bodies in leaves. Since the flagellin

peptide flg22 downregulates the formation of constitutive ER bodies and we

have shown that ER bodies negatively regulate PTI, we hypothesize that *P.*

syringae-induction of ER bodies is a virulence strategy to suppress host

immune responses. The two main virulence factors of *P. syringae* pv. tomato

are the phytotoxin coronatine and the T3SS effectors. The bacterial phytotoxin

coronatine, which mimics the function of JA in the plant (Bender et al., 1999),

induces the formation of ER bodies in leaves (Fig 3A,B). JA is perceived in the

plant through degradation of JASMONATE ZIM-DOMAIN (JAZ) transcriptional

repressors and JAZ proteins suppress the expression of ER body components

as well as indole glucosinolates (Guo et al., 2018; Ghorbel et al., 2021). Thus,

JA or pathogen JA-mimics can regulate the abundance of this specialized

organelle to induce or suppress defense depending on pathogen type. The

most studied coronatine virulence function is to re-open stomata upon PAMP-

induced closure (Melotto et al., 2006; Melotto et al., 2008). However, coronatine

also has been shown to suppress immunity once the bacteria colonize the

apoplast (Fig. 3C; (Geng et al., 2012)). The T3SS defective mutant Δ*hrcC* is not

able to induce the formation of ER bodies, which would indicate an effect of

T3Es on the induction of ER bodies. However, we cannot rule out that the

absence of ER bodies is due to an insufficient amount of coronatine produced

by the Δ*hrcC* microcolonies.

Bacterial flg22 perception leads to a down regulation of the ER bodies at both

transcriptional and post-transcriptional levels (Fig. 1 and S1). We found that

Arabidopsis mutants defective in either ER body biogenesis or composition are

more resistant to pathogenic bacteria (Fig. 2 and S2). The main components of

ER bodies are myrosinases, enzymes that modify glucosinolates to initiate the

glucosinolate breakdown, which will result in different products depending on

the biochemical properties of the modifier protein and the chemical nature of the

glucosinolate side chain (Wittstock and Burow, 2010; Sugiyama and Hirai,

2019). Those products can function as defense compounds or as signaling

molecules.

PYK10/BGLU23 is an atypical myrosinase similar to PENETRATION 2 (PEN2,

BGLU26) that hydrolyzes indole glucosinolates (IGs) (Nakano et al., 2017;

Nakazaki et al., 2019b). PEN2-mediated conversion of 4-methoxyindol-3-

ylmethylglucosinolate (4MI3G) into glucosinolate-derived products plays an

important role in the defense response against fungi (Bednarek et al., 2009).

Interestingly, *PEN2* is required for callose deposition upon flg22 perception and

for resistance against *Pst* DC3000 infection in seedlings grown in liquid media

(Clay et al., 2009; Johansson et al., 2014). In contrast, in adult plants grown in

soil, callose deposition and *Pst* DC3000 growth is not affected in the *pen2-1*

mutant (Geng et al., 2012). The *pen2-1* mutant grown in soil exhibits enhanced

resistance to *P. syringae* pv. *maculicola* (Stahl et al., 2016). These apparently

contradictory results indicate that PEN2 activity is dependent on age and/or the

metabolic state of the plant. 4MI3G overaccumulates in homogenized tissue of

bglu18/pyk10 mutants indicating that one or both of these β-glucosidases

hydrolyze 4MI3G (Nakazaki et al., 2019). Our results indicate that the single

pyk10-1 mutant and the double *pyk10-1/bglu21* mutant accumulate an

increased number of callose deposits in response to flg22 treatment and are

more resistant to *Pst* DC3000. Although both PEN2 and PYK10 seem to have

the same substrate (4MI3G), their respective mutants differ on phenotypes such

as flg22-induced callose deposition and resistance to *P. syringae Pst* DC3000

(Clay et al., 2009; Johansson et al., 2014). Apart from the above-mentioned

effect related to growth conditions, one of the reasons explaining these

differences could be the different compartmentalization of both PEN2 and

PYK10. While PEN2 is localized in peroxisome and mitochondria (Lipka et al.,

2005; Fuchs et al., 2016), PYK10 localizes in ER bodies (Matsushima et al.,

2003). Furthermore, myrosinase activity is regulated by different Myrosinase-

Biding Proteins and Myrosinase-Associated Proteins, as well as different

specifier proteins, which lead to the formation of different breakdown products

(Wittstock et al., 2016). This variety of compounds can have a direct or indirect

effect on the plant defense response (Wittstock et al., 2016). It is possible that

mutants lacking ER bodies or core components of ER bodies also exhibit

heightened basal defense, resulting in enhanced ROS and callose production.

In conclusion, we have demonstrated the involvement of ER body-dependent

regulation of immunity to bacteria and induction of ER bodies as a pathogen

virulence strategy. Further work will shed light on how this novel function of ER

bodies influences bacterial-plant interactions and pave the way for the

discovery of the role of indole-glucosinolates breakdown on the response

against bacterial invasion in plants.

Experimental procedures

Plant material and growth conditions

A. thaliana Col-0, the transgenic line expressing GFP-HDEL (Hayashi et al.,

2001) and the mutants *nai1-1* (Matsushima et al., 2003), *pyk10-1* (Nagano et

al., 2008), *pyk10-1/bglu21* (Nagano et al., 2009), *bglu18-1* (Ogasawara et al.,

2009) and *pbp1-1* (Nagano et al., 2005) were grown in a controlled environment

chamber at 23°C, 70% relative humidity, and a 10h/14h light/dark photoperiod

with light intensity of $100\mu E.m^{-2}.S^{-1}$. All seeds were stratified for 3-4 days at 4ºC

before sowing into soil.

Bacterial strains, growth conditions and pathogen assays

Pseudomonas syringae pv. *tomato* DC3000 (Cuppels, 1986) and the derivative

strains Δ*hrcC* and *cor*⁻ (cma- cfa-; (Brooks et al., 2004)) were grown at 28°C in

Lysogeny Broth (LB) medium (Bertani, 1951). Antibiotics were used when

appropriate, at the following concentrations: Rifampicin (100 µg/ml), kanamycin

(25 µg/ml) and gentamycin (10 µg/ml).

The *Pst* DC3000, Δ*hrcC* and *cor*⁻ eYFP (enhanced yellow fluorescent protein)

derivatives were generated using a Tn7 delivery system as previously described

(Rufian et al., 2018)

For pathogen assays, five-week-old *Arabidopsis* plants were inoculated with a

P. syringae suspension of $5x10^4$ CFU/mL prepared in 10 mM $MgCl_2$. Three

leaves per plant were syringe infiltrated and five plants were used for each

experiment. Four days after inoculation, three leaf discs of 1 cm of diameter

(one per leaf were ground in 1 mL 10 mM $MgCl_2$, and serial dilutions were

plated.

Transcriptomic analyses

For the analysis of NAI1 and PYK10 expression under different biotic stresses,

we used transcriptomic data available in eFP Browser. Specifically, we used the

dataset AtGenExpress: Response to virulent, avirulent, type III□secretion

system deficient and nonhost bacteria (NASCArrays-120). 5-week-old Col-0

plants were infiltrated with the corresponding *P. syringae* strain. Samples were

taken 24h after treatment and RNA was isolated and hybridized to the ATH1

GeneChip. The data were normalized by GCOS normalization, TGT 100.

flg22-related assays

For ROS burst assays, leaf discs were collected using a cork borer of 5mm

diameter from three-week-old *Arabidopsis* plants and floated overnight in

deionized water. The next day, the water was replaced with an assay solution

containing 17 mg/ml luminol (Sigma), 10 mg/ml horse radish peroxidase

(Sigma) and 100 nM flg22 (Genscript). Luminescence was measured using

Tristar multimode reader (Berthold technology).

Callose deposits were quantified from four-week-old *Arabidopsis* leaves

infiltrated with 10 µM flg22 solution. Eighteen hours after treatment, leaves were

detached and incubated 15 minutes at 65ºC in 5 ml of alcoholic lactophenol (1

volume of phenol: glycerol: lactic acid: water (1:1:1:1) and 2 volumes of

ethanol). Leaves were transferred to fresh alcoholic lactophenol and incubated

overnight at room temperature. After two washes with 50% ethanol, leaves were

stained with aniline blue and callose deposits were visualized under a

fluorescence microscope. Quantification of callose deposits was performed

using the Fiji distribution of ImageJ.

Confocal microscopy

Two-week old *Arabidopsis* GFP-HDEL plants were vacuum infiltrated with either

water or 100 nM flg22. Three hours after the treatment, cotyledons were

detached and observed under a Zeiss LSM710 confocal microscope equipped

with a LDC-apochromat 40x/1.1W Korr M27 water-immersion objective (NA

1.1). GFP was excited at 488 nm and emission collected at 500-550 nm. For

monitoring of the dynamics of the ER bodies, cotyledons were treated as above

and visualized using a Leica SP8 confocal microscope (Leica Microsystems

GmbH, Wetzlar, Germany). GFP was excited at 488 nm and emission collected

at 500-550 nm.

For visualization of bacterial-induced ER bodies, five-week old *Arabidopsis*

GFP-HDEL plants were brush-inoculated with a 5×10^7 CFU/mL suspension of

the indicated strain (Fig. 3A). Three days after inoculation, leaves were

observed as described above.

For visualization of bacterial microcolonies, five-week old *Arabidopsis* GFP-

HDEL plants were syringe infiltrated with a 5×10^4 CFU/mL suspension and

leaves were visualized using the Leica SP5 II confocal microscope (Leica

Microsystems GmbH, Wetzlar, Germany). Images of eYFP and GFP were

sequentially obtained using the following conditions (excitation/ emission):

eYFP (514 nm/ 525 to 600 nm), GFP (488/ 500 to 525 nm). Z series imaging

was taken at 1 μm interval. Image processing was performed using Leica LAS

AF (Leica Microsystems). ER bodies were quantified using Fiji distribution of

ImageJ software.

***S. exigua* performance**

S. exigua eggs were kindly provided by Dr. Salvador Herrero (U. Valencia).

After eclosion, *S. exigua* larvae were fed with an artificial diet (Elvira et al.,

2010). Three days after eclosion, five larvae of approximately same size were

transferred to a four-to-five week old *Arabidopsis* plant grown in Jiffy-7 pods.

Larvae were transfer to new plants every 24 hours to avoid food privation.

Seven days after plant feeding, fresh weight of the larvae was measured. For

each experiment, five replicates were performed (25 larvae per genotype).

ACKNOWLEDGMENTS

J.S. Rufián has been supported by a FPI fellowship associated with a grant of

E.R. Bejarano (MICINN, Spain; AGL2010-22287-C02-2), funds from BIO2012-

35641 (MICINN, Spain) granted to C.R. Beuzón, and Plan Propio de la

Universidad de Málaga – Andalucía Tech. The work was co-funded by

European Regional Development Funds (FEDER). G. Coaker is supported by

grants from the National Institutes of Health (RO1GM092772, R35GM13640).

We thank Alberto Macho for hosting J.S. Rufián in his research laboratory and

providing feedback on the manuscript.

Figure legends

Figure 1. ER bodies are downregulated after activation of PTI.

(**A**) Constitutive ER bodies disappear after flg22 treatment. Two-week-old

Arabidopsis GFP-HDEL plants were vacuum infiltrated with water or 100 nM

flg22. Cotyledons were observed in a Zeiss LSM710 confocal microscope three

hours after treatment. Red autofluorescence correspond to chloroplasts. For ER

body quantification, three fields from three independent cotyledons were used.

ER bodies were quantified using Fiji. The experiment was repeated four times

with similar results and a mean value ± SE of all quantifications is shown.

Statistical differences were determined with a two-tailed t-test comparing water

treatment with flg22. Three asterisks indicate p<0.001. Scale bar: 20 μm. (**B**)

Two-week-old *Arabidopsis* GFP-HDEL plants were vacuum infiltrated with water

or 100 nM flg22. The infiltrated cotyledons were monitored for 120 minutes

taken photos every 5 minutes with a Leica SP8 confocal microscope. (**C**)

Expression patterns of *nai1* and *pyk10* genes under different biotic stress. Data

was obtained from available microarray in eFP Browser. Five week-old Col-0

plant leaves were syringe-infiltrated with either a 1 μM flg22 solution (using

water as mock treatment) or the indicated bacterial strain at 10^8 cfu/ml (using 10

mM $MgCl_2$ as mock treatment). Samples were taken 24 hours after treatment.

Fold change relative to mock treatment is shown.

Figure 2. ER body mutants exhibit enhanced resistance against *P.*

syringae.

(**A**) flg22-induced ROS burst in Col-0, *nai1-1*, *pyk10-1* and *pyk10-1/bglu21*.

Three-week-old *Arabidopsis* leaf discs were treated with 100 nM flg22 and ROS

was quantified using a luminescence based assay. The left graph represent the

dynamics of the ROS produced in the different genotypes and the right graph

represent the total relative lights units (RLU) detected over a 30 minutes period.

Error bars indicate SD, n = 16. Statistically differences were determined with

one-way ANOVA (α = 0.05) with Tukey's multiple comparisons test and different

letters indicate statistical significance. The experiment was repeated four times

with similar results and a representative experiment is shown. (**B**) Quantification

of flg22-induced callose deposition in Col-0, *nai1-1*, *pyk10-1* and *pyk10-*

1/bglu21. Four-week-old plants were infiltrated with 10 µM flg22 and callose

deposits were visualized under an epifluorescence microscope eighteen hours

after treatment. Four images were taken per leaf and nine leaves were used per

experiment. The mean number of deposits was quantified and is included ± SE

in the bottom-right corner of each representative image. The experiment was

repeated three times with similar results, and a representative experiment is

shown. White bar: 100 µm. (**C**) Growth of *P. syringae* pv. *tomato* DC3000 in

Col-0, *nai1-1*, *pyk10-1* and *pyk10-1/bglu21.* Five-week-old plant leaves were

syringe inoculated with a suspension of $5x10^4$ CFU/mL. Four days after

inoculation, bacteria were recovered and quantified. Values from three

independent replicates are shown (*n*=18). Different symbols represent individual

values from different replicates. Statistically differences were determined with

by ANOVA (α = 0.05) with Tukey's multiple comparisons test and different

letters indicate statistical significance. (**D**) Growth of the *P. syringae* pv. *tomato*

Δ*hrcC* derivative in Col-0, *nai1-1*, and *pyk10-1/bglu21.* Five-week-old plant

leaves were syringe inoculated with a suspension of $5x10^4$ CFU/mL. Four days

after inoculation, bacteria were recovered and quantified. Values from three

independent replicates are shown (*n*=18). Different symbols represent individual

values from different replicates. Statistically differences were determined by

ANOVA (α= 0.05) with Tukey's multiple comparisons test and different letters

indicate statistical significance. (E) Three-week-old *Arabidopsis* Col-0, *nai1-1*,

pyk10-1 and *pyk10-1/bglu21* plants were challenged with three-days-old *S.*

exigua larvae. Seven days after feeding, fresh weight of each larva was

measured. Bars represent the mean weight ± SE (*n*=20). Statistically

differences were determined with one-way ANOVA (α= 0.05) with Tukey's

multiple comparisons test and different letters indicate statistical significance.

The experiment was repeated three times with similar results, and a

representative experiment is shown.

Figure 3. Coronatine induces formation of ER bodies in leaves and its

virulence function is compromised in mutants impaired in ER body

formation.

(**A-C**) Confocal microscope images of bacterial micro-colonies (yellow) and ER

bodies (green). Five-week-old *Arabidopsis* GFP-HDEL plants were brush-

inoculated with a suspension containing 5×10^7 CFU/mL (A) or syringe

inoculated with a bacterial suspension containing 5×10^4 CFU/mL (B, C). Three

days after inoculation, leaf sections were visualized under the confocal

microscope. A representative z-stack projection image is shown. ER bodies

were quantified from the green channel using Fiji. Bars represent the mean

number of ER bodies in (A) or the number of ER bodies surrounding a single

bacterial micro colony in (B, C) ± SE. Red autofluorescence correspond to

chloroplasts. The experiments were repeated three times with similar results,

and the mean of all experiments is shown. ER bodies were quantified using Fiji

and statistically differences were determined with one-way ANOVA (α= 0.05)

with Tukey's multiple comparison test in (A) and different letters indicate

statistical significance. In (B, C) data were analyzed by a t-test (α= 0.001).

White bar: 20 µm (A), 100 µm (B), 50 µm (C). (**D**) Growth of *P. syringae* pv.

tomato DC3000 and its derivative coronatine mutant (*cor* -, DC3118) in Col-0,

nai1-1, *pyk10-1* and *pyk10-1/bglu21.* Five-week-old plant leaves were syringe

inoculated with a bacterial suspension of 5×10^4 CFU/mL. Four days after

inoculation, bacteria were recovered and quantified. Values from three

independent replicates are shown (*n*=12). Different symbols represent individual

values from different replicates. Statistically differences were determined by

ANOVA (α = 0.05) with Tukey's multiple comparisons test and different letters

indicate statistical significance.

Supplemental Figure Legends

Figure S1. Downregulation of *NAI1* and *PYK10* after flg22 treatment.

Expression pattern of *NAI1* and *PYK10* after flg22 treatment (5, 10, 30, 90 and

180 minutes) in Col-0 and *fls2* plants. Data were obtained from Bjornson et al.

(2021). Fourteen-day old Arabidopsis Col-0 or *fls2* seedlings were immersed in

1 µM flg22 and samples were taken at the indicated time points.

Figure S2. *bglu18-1* and *pbp1-1* show enhanced resistance to *P. syringae*

(**A**) Flg22-induced ROS burst in Col-0, *bglu18-1*, and *pbp1-1*. Three-week-old

Arabidopsis leaf discs were treated with 100 nM flg22 and ROS was quantified

using a luminescence based assay. Error bars indicate ±SE, n = 16. The

experiment was repeated twice with similar results and a representative

experiment is shown. (**B**) Growth of *P. syringae* pv. *tomato* DC3000 in Col-0,

bglu18-1, and *pbp1-1.* Five-week-old plant leaves were syringe inoculated with

a suspension of 5×10^4 CFU/mL. Four days after inoculation, bacteria were

recovered and quantified. Values for each individual plant are shown (*n*=6).

Bars represent the mean value ± SE. Statistically differences were determined

with one-way ANOVA (α= 0.05) with Tukey's multiple comparisons test and

different letters indicate statistical significance. The experiment was repeated

twice with similar results and a representative experiment is shown.

Figure S3. Coronatine induce the expression of ER body and indole

glucosinolate related genes.

(A) Transcriptomic analysis of several genes related with ER bodies formation

or indole-glucosinolate metabolism. Bars show Log_2 signal values from

microarrays datasets. Data was obtained from the study by Thilmony et al.

(2006), available in Genevestigator. *Arabidopsis* Col-5 leaves were inoculated

with water (Mock), or a 10^6 cfu/ml suspension of *Pst* DC3000 or *Pst* DC3118

(cor⁻). Samples were taken 24 hours after inoculation. (B) Coexpression

analyses of PYK10 (At3G09260). Analyses were performed using ATTED-II

software.

Literature Cited

Bednarek, P., Piślewska-Bednarek, M., Svatos, A., Schneider, B., Doubsky, J.,
Mansurova, M., Humphry, M., Consonni, C., Panstruga, R., Sanchez-
Vallet, A., Molina, A., and Schulze-Lefert, P. (2009). A glucosinolate
metabolism pathway in living plant cells mediates broad-spectrum
antifungal defense. In Science (American Association for the
Advancement of Science), pp. 101-106.
Bender, C.L., Alarcón-Chaidez, F., and Gross, D.C. 1999. Pseudomonas
syringae phytotoxins: mode of action, regulation, and biosynthesis by
peptide and polyketide synthetases. Microbiol Mol Biol Rev 63:266-292.
Bertani, G. 1951. Studies on lysogenesis. I. The mode of phage liberation by
lysogenic Escherichia coli. Journal of bacteriology 62:293-300.
Bjornson, M., Pimprikar, P., Nurnberger, T., and Zipfel, C. 2021. The
transcriptional landscape of Arabidopsis thaliana pattern-triggered
immunity. Nat Plants.
Brooks, D.M., Hernandez-Guzman, G., Kloek, A.P., Alarcon-Chaidez, F.,
Sreedharan, A., Rangaswamy, V., Penaloza-Vazquez, A., Bender, C.L.,
and Kunkel, B.N. 2004. Identification and characterization of a well-
defined series of coronatine biosynthetic mutants of Pseudomonas
syringae pv. tomato DC3000. Mol Plant Microbe Interact 17:162-174.
Chung, H.S., Koo, A.J.K., Gao, X., Jayanty, S., Thines, B., Jones, A.D., and
Howe, G.A. (2008). Regulation and function of Arabidopsis JASMONATE
ZIM-domain genes in response to wounding and herbivory. In Plant
physiology (American Society of Plant Biologists), pp. 952-964.
Cipollini, D., Enright, S., Traw, M.B., and Bergelson, J. (2004). Salicylic acid
inhibits jasmonic acid-induced resistance of Arabidopsis thaliana to
Spodoptera exigua. In Molecular Ecology, pp. 1643-1653.
Clay, N.K., Adio, A.M., Denoux, C., Jander, G., and Ausubel, F.M. 2009.
Glucosinolate metabolites required for an Arabidopsis innate immune
response. Science 323:95-101.
Couto, D., and Zipfel, C. 2016. Regulation of pattern recognition receptor
signalling in plants. Nature reviews. Immunology 16:537-552.
Cuppels, D.A. 1986. Generation and Characterization of Tn*5* Insertion
Mutations in *Pseudomonas syringae* pv. *tomato*. Appl. Environ. Microbiol.
51:323-327.
Elvira, S., Gorria, N., Munoz, D., Williams, T., and Caballero, P. 2010. A
simplified low-cost diet for rearing Spodoptera exigua (Lepidoptera:
Noctuidae) and its effect on S. exigua nucleopolyhedrovirus production. J
Econ Entomol 103:17-24.
Fuchs, R., Kopischke, M., Klapprodt, C., Hause, G., Meyer, A.J.,
Schwarzländer, M., Fricker, M.D., and Lipka, V. (2016). Immobilized
Subpopulations of Leaf Epidermal Mitochondria Mediate
PENETRATION2-Dependent Pathogen Entry Control in Arabidopsis. In
The Plant Cell … (American Society of Plant Biologists), pp. 130-145.
Geng, X., Cheng, J., Gangadharan, A., and Mackey, D. (2012). The coronatine
toxin of Pseudomonas syringae is a multifunctional suppressor of
Arabidopsis defense. In The Plant Cell …, pp. 4763-4774.
Ghorbel, M., Brini, F., Sharma, A., and Landi, M. 2021. Role of jasmonic acid in
plants: the molecular point of view. Plant Cell Rep.

Gimenez-Ibanez, S., Boter, M., Ortigosa, A., García-Casado, G., Chini, A., Lewsey, M.G., Ecker, J.R., Ntoukakis, V., and Solano, R. (2017). JAZ2 controls stomata dynamics during bacterial invasion. In New Phytol., pp. 1378-1392.

Gómez-Gómez, L., and Boller, T. 2000. FLS2: an LRR receptor-like kinase involved in the perception of the bacterial elicitor flagellin in Arabidopsis. Mol Cell 5:1003-1011.

Guo, Q., Yoshida, Y., Major, I.T., Wang, K., Sugimoto, K., Kapali, G., Havko, N.E., Benning, C., and Howe, G.A. 2018. JAZ repressors of metabolic defense promote growth and reproductive fitness in Arabidopsis. Proc Natl Acad Sci U S A 115:E10768-E10777.

Henry, E., Yadeta, K.A., and Coaker, G. 2013. Recognition of bacterial plant pathogens: local, systemic and transgenerational immunity. New Phytol 199:908-915.

Johansson, O.N., Fantozzi, E., Fahlberg, P., Nilsson, A.K., Buhot, N., Tör, M., and Andersson, M.X. (2014). Role of the penetration-resistance genes PEN1, PEN2and PEN3in the hypersensitive response and race-specific resistance in Arabidopsis thaliana. In Plant J., pp. 466-476.

Lahrmann, U., Strehmel, N., Langen, G., Frerigmann, H., Leson, L., Ding, Y., Scheel, D., Herklotz, S., Hilbert, M., and Zuccaro, A. 2015. Mutualistic root endophytism is not associated with the reduction of saprotrophic traits and requires a noncompromised plant innate immunity. New Phytol 207:841-857.

Lipka, V., Dittgen, J., Bednarek, P., Bhat, R., Wiermer, M., Stein, M., Landtag, J., Brandt, W., Rosahl, S., Scheel, D., Llorente, F., Molina, A., Parker, J., Somerville, S., and Schulze-Lefert, P. (2005). Pre- and postinvasion defenses both contribute to nonhost resistance in Arabidopsis. In Science (American Association for the Advancement of Science), pp. 1180-1183.

Macho, A.P. 2015. Subversion of plant cellular functions by bacterial type-III effectors: beyond suppression of immunity. New Phytol.

Matsushima, R., Fukao, Y., Nishimura, M., and Hara-Nishimura, I. (2004). NAI1 gene encodes a basic-helix-loop-helix-type putative transcription factor that regulates the formation of an endoplasmic reticulum-derived structure, the ER body. In THE PLANT CELL ONLINE (American Society of Plant Biologists), pp. 1536-1549.

Matsushima, R., Hayashi, Y., Kondo, M., Shimada, T., Nishimura, M., and Hara-Nishimura, I. (2002). An endoplasmic reticulum-derived structure that is induced under stress conditions in Arabidopsis. In Plant physiology (American Society of Plant Biologists), pp. 1807-1814.

Matsushima, R., Hayashi, Y., Yamada, K., Shimada, T., Nishimura, M., and Hara-Nishimura, I. (2003). The ER body, a novel endoplasmic reticulum-derived structure in Arabidopsis. In Plant Cell Physiol., pp. 661-666.

Melotto, M., Underwood, W., and He, S.Y. 2008. Role of stomata in plant innate immunity and foliar bacterial diseases. Annu Rev Phytopathol 46:101-122.

Melotto, M., Underwood, W., Koczan, J., Nomura, K., and He, S.Y. 2006. Plant stomata function in innate immunity against bacterial invasion. Cell 126:969-980.

Millet, Y.A., Danna, C.H., Clay, N.K., and Songnuan, W. (2010). Innate immune responses activated in Arabidopsis roots by microbe-associated molecular patterns. In The Plant Cell ….

Mitsuhashi, N., Shimada, T., Mano, S., Nishimura, M., and Hara-Nishimura, I. 2000. Characterization of organelles in the vacuolar-sorting pathway by visualization with GFP in tobacco BY-2 cells. Plant Cell Physiol 41:993-1001.

Müller, R., de Vos, M., Sun, J.Y., Sønderby, I.E., Halkier, B.A., Wittstock, U., and Jander, G. (2010). Differential Effects of Indole and Aliphatic Glucosinolates on Lepidopteran Herbivores. In J Chem Ecol, pp. 905-913.

Nagano, A.J., Matsushima, R., and Hara-Nishimura, I. (2005). Activation of an ER-body-localized beta-glucosidase via a cytosolic binding partner in damaged tissues of Arabidopsis thaliana. In Plant Cell Physiol., pp. 1140-1148.

Nagano, A.J., Fukao, Y., Fujiwara, M., Nishimura, M., and Hara-Nishimura, I. (2008). Antagonistic jacalin-related lectins regulate the size of ER body-type beta-glucosidase complexes in Arabidopsis thaliana. In Plant Cell Physiol., pp. 969-980.

Nagano, A.J., Maekawa, A., Nakano, R.T., Miyahara, M., Higaki, T., Kutsuna, N., Hasezawa, S., and Hara-Nishimura, I. (2009). Quantitative analysis of ER body morphology in an Arabidopsis mutant. In Plant Cell Physiol., pp. 2015-2022.

Nakano, R.T., Yamada, K., Bednarek, P., Nishimura, M., and Hara-Nishimura, I. 2014a. ER bodies in plants of the Brassicales order: biogenesis and association with innate immunity. Front Plant Sci 5:73.

Nakano, R.T., Yamada, K., Bednarek, P., Nishimura, M., and Hara-Nishimura, I. (2014b). ER bodies in plants of the Brassicales order: biogenesis and association with innate immunity. In Front. Plant Sci. (Frontiers), pp. 73.

Nakano, R.T., Piślewska-Bednarek, M., Yamada, K., Edger, P.P., Miyahara, M., Kondo, M., Böttcher, C., Mori, M., Nishimura, M., Schulze-Lefert, P., Hara-Nishimura, I., and Bednarek, P. (2017). PYK10 myrosinase reveals a functional coordination between endoplasmic reticulum bodies and glucosinolates in Arabidopsis thaliana. In Plant J., pp. 204-220.

Nakazaki, A., Yamada, K., Kunieda, T., Tamura, K., Hara-Nishimura, I., and Shimada, T. (2019a). Biogenesis of leaf endoplasmic reticulum body is regulated by both jasmonate-dependent and independent pathways. In Plant Signaling & Behavior (Taylor & Francis), pp. 1-3.

Nakazaki, A., Yamada, K., Kunieda, T., Sugiyama, R., Hirai, M.Y., Tamura, K., Hara-Nishimura, I., and Shimada, T. (2019b). Leaf Endoplasmic Reticulum Bodies Identified in Arabidopsis Rosette Leaves Are Involved in Defense against Herbivory. In Plant physiology, pp. 1515-1524.

Ogasawara, K., Yamada, K., Christeller, J.T., Kondo, M., Hatsugai, N., Hara-Nishimura, I., and Nishimura, M. (2009). Constitutive and inducible ER bodies of Arabidopsis thaliana accumulate distinct beta-glucosidases. In Plant Cell Physiol., pp. 480-488.

Rehrig, E.M., Appel, H.M., Jones, A.D., and Schultz, J.C. (2014). Roles for jasmonate- and ethylene-induced transcription factors in the ability of Arabidopsis to respond differentially to damage caused by two insect herbivores. In Front. Plant Sci. (Frontiers), pp. 407.

Rufian, J.S., Macho, A.P., Corry, D.S., Mansfield, J.W., Ruiz-Albert, J., Arnold,
D.L., and Beuzon, C.R. 2018. Confocal microscopy reveals in planta
dynamic interactions between pathogenic, avirulent and non-pathogenic
Pseudomonas syringae strains. Molecular plant pathology 19:537-551.
Santamaría, M.E., Martinez, M., Arnaiz, A., Ortego, F., Grbic, V., and Diaz, I.
(2017). MATI, a Novel Protein Involved in the Regulation of Herbivore-
Associated Signaling Pathways. In Front. Plant Sci., pp. 80-18.
Sherameti, I., Venus, Y., Drzewiecki, C., Tripathi, S., Dan, V.M., Nitz, I., Varma,
A., Grundler, F.M., and Oelmüller, R. (2008). PYK10, a beta-glucosidase
located in the endoplasmatic reticulum, is crucial for the beneficial
interaction between Arabidopsis thaliana and the endophytic fungus
Piriformospora indica. In Plant J. (Blackwell Publishing Ltd), pp. 428-439.
Sugiyama, R., and Hirai, M.Y. (2019). Atypical Myrosinase as a Mediator of
Glucosinolate Functions in Plants. In Front. Plant Sci. (Frontiers), pp.
1008.
Thilmony, R., Underwood, W., and He, S.Y. 2006. Genome-wide transcriptional
analysis of the Arabidopsis thaliana interaction with the plant pathogen
Pseudomonas syringae pv. tomato DC3000 and the human pathogen
Escherichia coli O157:H7. Plant J 46:34-53.
Van Oosten, V.R., Bodenhausen, N., Reymond, P., Van Pelt, J.A., Van Loon,
L.C., Dicke, M., and Pieterse, C.M.J. (2008). Differential effectiveness of
microbially induced resistance against herbivorous insects in
Arabidopsis. In Mol. Plant Microbe Interact. (The American
Phytopathological Society), pp. 919-930.
Weiler, E.W., Kutchan, T.M., Gorba, T., Brodschelm, W., Niesel, U., and Bublitz,
F. 1994. The Pseudomonas phytotoxin coronatine mimics octadecanoid
signalling molecules of higher plants. FEBS letters 345:9-13.
Wittstock, U., and Burow, M. (2010). Glucosinolate Breakdown in Arabidopsis:
Mechanism, Regulation and Biological Significance. In The Arabidopsis
Book, pp. e0134-0114.
Wittstock, U., Kurzbach, E., Herfurth, A.M., and Stauber, E.J. (2016).
Glucosinolate Breakdown. In Advances in Botanical Research (Elsevier
Ltd), pp. 125-169.
Xu, Z., Escamilla-Trevino, L., Zeng, L., Lalgondar, M., Bevan, D., Winkel, B.,
Mohamed, A., Cheng, C.L., Shih, M.C., Poulton, J., and Esen, A. 2004.
Functional genomic analysis of Arabidopsis thaliana glycoside hydrolase
family 1. Plant Mol Biol 55:343-367.
Yamada, K., Hara-Nishimura, I., and Nishimura, M. (2011). Unique defense
strategy by the endoplasmic reticulum body in plants. In Plant Cell
Physiol., pp. 2039-2049.
Yamada, K., Nagano, A.J., Nishina, M., Hara-Nishimura, I., and Nishimura, M.
(2008). NAI2 is an endoplasmic reticulum body component that enables
ER body formation in Arabidopsis thaliana. In THE PLANT CELL
ONLINE (American Society of Plant Biologists), pp. 2529-2540.
775