Antagonistic modules, SIB1 and LSD1, regulate photosynthesis-associated nuclear

genes via GOLDEN2-LIKE transcription factors in Arabidopsis

Mengping Li[1,2,a], Keun Pyo Lee[1,a], Tong Liu[1,2], Vivek Dogra[1,b], Jianli Duan[1],

Mengshuang Li[1,2], Weiman Xing[1], and Chanhong Kim[1,c]

[1] Shanghai Center for Plant Stress Biology, CAS Center for Excellence in Molecular

Plant Sciences, Chinese Academy of Sciences, Shanghai 200032, China.

[2] University of the Chinese Academy of Sciences, Beijing 100049, China.

[a] These authors contributed equally to this work.

[b] Present address: Biotechnology Division, CSIR-Institute of Himalayan Bioresource

Technology, Palampur 176061, India

[c] Correspondence: chanhongkim@cemps.ac.cn

Running title: SIB1 and LSD1 modulate GLK activity

ABSTRACT

GOLDEN2-LIKE (GLK) transcription factors drive the expression of photosynthesis-associated nuclear genes (PhANGs), indispensable for chloroplast biogenesis. We previously demonstrated that the salicylic acid (SA)-induced SIGMA FACTOR-BINDING PROTEIN 1 (SIB1), a transcription coregulator and positive regulator of cell death, interacts with GLK1 and GLK2 to reinforce their activities. The SIB1-GLK interaction raises the level of light-harvesting antenna proteins in photosystem II, aggravating photoinhibition and singlet oxygen ($^1O_2$) burst. $^1O_2$ then contributes to SA-induced cell death via EXECUTER 1 (EX1, $^1O_2$ sensor protein)-mediated retrograde signaling upon reaching a critical level. We now reveal that LESION-SIMULATING DISEASE 1 (LSD1), a transcription coregulator and negative regulator of SA-primed cell death, interacts with GLK1/2 to repress their activities. Consistently, the overexpression of LSD1 represses GLK target genes including PhANGs, whereas the loss of LSD1 increases their expression. Remarkably, LSD1 overexpression inhibits chloroplast biogenesis, resembling the characteristic *glk1glk2* double mutant phenotype. The subsequent chromatin immunoprecipitation analysis coupled with quantitative PCR further reveals that LSD1 inhibits the DNA-binding activity of GLK1 towards its target promoters. The SA-induced nuclear-targeted SIB1 appears to counteractively interact with GLK1/2, leading to the activation of EX1-mediated $^1O_2$ signaling. Taken together, we provide a working model that SIB1 and LSD1, mutually exclusive SA-signaling components, antagonistically regulate GLK1/2 to fine-tune the expression of PhANGs, thereby modulating $^1O_2$ homeostasis and related stress responses.

Key words: SIB1; LSD1; GLK; photosynthesis-associated nuclear genes; singlet oxygen; EXECUTER1; salicylic acid; retrograde signaling

INTRODUCTION

Chloroplasts communicate with the nucleus via retrograde signaling (RS) in response to the ever-changing environment. Upon exposure to unfavorable environmental conditions, chloroplasts downregulate photosynthesis-associated nuclear genes (PhANGs), referred to as biogenic RS, but stimulate the expression of stress-related genes via alternate RS pathways, collectively called operational RS. The nuclear-encoded chloroplast GENOMES UNCOUPLED 1 (GUN1) protein plays a pivotal role in the biogenic RS (Nott et al., 2006). GUN1 integrates various retrograde signals released by the disturbance in plastid gene expression, redox homeostasis, and tetrapyrrole biosynthesis in chloroplasts (Chan et al., 2016; Koussevitzky et al., 2007; Nott et al., 2006). The well-known downstream targets of GUN1-mediated RS are two nuclear genes encoding the GOLDEN2-LIKE (GLK) transcription factors (TFs) (Martin et al., 2016; Waters et al., 2009). In fact, GUN1-mediated RS represses *GLK* transcription. In *Arabidopsis thaliana* (Arabidopsis), GLK1 and GLK2 function redundantly to express PhANGs, promoting chloroplast biogenesis. Consistently, the loss of both GLKs significantly impairs chloroplast biogenesis (Fitter et al., 2002).

Recent studies discovered an unexpected function of GLKs towards plant immune responses. The steady-state levels of salicylic acid (SA)-responsive genes are significantly lower in GLK1-overexpressing (*oxGLK1*) Arabidopsis transgenic plants relative to wild-type (WT) plants (Savitch et al., 2007). Accordingly, the *oxGLK1* plants are susceptible to the biotrophic pathogen *Hyaloperonospora arabidopsidis* (*Hpa*) Noco2, while *glk1 glk2* double knockout mutant plants are more resistant compared to WT plants (Murmu et al., 2014). However, other studies reported that GLKs confer resistance towards the cereal fungal pathogen *Fusarium graminearum* (Savitch et al., 2007), necrotrophic fungal pathogen *Botrytis cinerea* (Murmu et al., 2014), and the Cucumber mosaic virus (Han et al., 2016). These findings indicate that multiple regulatory circuits (positive and negative) may differently modulate GLK activity towards various microbial pathogens.

We lately demonstrated that the nuclear-targeted SIGMA FACTOR-BINDING

PROTEIN 1 (SIB1), a defense-related transcription coregulator, interacts with GLK1/2

in response to an increase in foliar SA (Lai et al., 2011; Lv et al., 2019). In

Arabidopsis *lesion-simulating disease 1* (*lsd1*) mutant grown under continuous light

(CL) conditions, the transiently increased level of SA rapidly induces the otherwise

undetectable SIB1, leading to its accumulation in both the nucleus and the chloroplasts

(Lai et al., 2011; Lv et al., 2019). It is important to note that the extended daylength is

one of the lesion-triggering external factors evoking SA-dependent runaway

(uncontrolled) cell death (RCD) in the *lsd1* mutant (Dietrich et al., 1994; Lv et al., 2019).

The SA receptor Nonexpresser of PR genes 1 (NPR1) induces the expression of *SIB1*

and the dual targeting of SIB1 also occurs in WT plants after SA treatment (Lai et al.,

2011; Lv et al., 2019; Xie et al., 2010). Whereas the loss of NPR1 abolishes *lsd1* RCD,

the loss of SIB1 significantly delays RCD (Aviv et al., 2002; Lv et al., 2019), indicating

that SIB1 is one of the RCD-triggering components directed by NPR1. The SIB1-GLK

interaction in the nucleus enhances the expression of PhANGs, while chloroplast-

localized SIB1 (cpSIB1) represses the expression of photosynthesis-associated plastid

genes (PhAPGs) (Lv et al., 2019; Morikawa et al., 2002). This concurrent uncoupled

expression of PhANGs and PhAPGs increases singlet oxygen ($^1O_2$) levels in

chloroplasts through enhanced photoinhibition in PSII (Lv et al., 2019). EXECUTER

1 (EX1), a $^1O_2$ sensor protein (Dogra et al., 2019), then mediates $^1O_2$-triggered RS to

contribute to stress responses in *lsd1* mutant plants (Lv et al., 2019). It appears that

SIB1 undergoes co-translational N-terminal acetylation (NTA) and post-translational

ubiquitination (Li et al., 2020). While NTA renders the nuclear SIB1 (nuSIB1) more

stable, the latter modification promotes its turnover via the ubiquitin-proteasome

system (UPS). The interplay of NTA and UPS seems to regulate nuSIB1-mediated

stress responses finely. Nonetheless, earlier reports regarding the positive role of both

nuSIB1 and cpSIB1 to RCD suggest that LSD1 may be required to repress the

expression of PhANGs to sustain $^1O_2$ homeostasis.

Here, we demonstrate that LSD1, a transcription coregulator and negative regulator

of cell death, interacts with GLK1/2. LSD1 considerably diminishes the GLK binding

activity to promoters of the examined PhANGs in Arabidopsis. In agreement, LSD1-

overexpressing plants exhibit significantly reduced levels of PhANGs, whereas loss of

LSD1 causes a notable upregulation of PhANGs relative to WT plants. SA most likely

intervenes in the LSD1-GLK interaction through a rapid accumulation of nuSIB1,

leading to a nuSIB1-GLKs interaction, enhanced expression of PhANGs, and activation

of EX1-dependent $^1O_2$ signaling implicated in cell death. We thus concluded that the

stress-associated but mutually exclusive transcription coregulators nuSIB1 (positive

regulator) and LSD1 (negative regulator) antagonistically regulate the expression of

PhANGs through the physical interaction with GLKs. Such antagonistic regulation of

GLK activity by nuSIB1 and LSD1 might be instrumental in sustaining $^1O_2$ homeostasis

under SA-associated stress conditions.

RESULTS

LSD1 interacts with the GOLDEN2-LIKE transcription factors GLK1 and GLK2

The stress hormone SA primes cell death in the *lsd1* mutant in a light-dependent manner,

a typical characteristic of most lesion mimic mutants, as manifested by the abrogated

cell death by either loss of key SA signaling components (such as NPR1) or

overexpression of the bacterial salicylate hydroxylase NahG that metabolizes SA (Lv

et al., 2019; Muhlenbock et al., 2008). Upon exposure to various stimuli, including light,

cold, UV-C, red light, hypoxia, and pathogens (Chai et al., 2015; Dietrich et al., 1997;

Huang et al., 2010; Jabs et al., 1996; Karpinski et al., 2013; Muhlenbock et al., 2007;

Muhlenbock et al., 2008; Rusaczonek et al., 2015), *lsd1* mutant plants drastically

develop the foliar RCD phenotype. Among those differentially regulated genes prior to

the onset of RCD, the SA-induced transcription coregulator nuSIB1 potentiates the

expression of PhANGs and stress-related genes by modulating the TF activity of

GLK1/2 and WRKY33, respectively (Lai et al., 2011; Lv et al., 2019; Zarrinpar et al.,

2003). These data suggest a possible antagonism between LSD1 and nuSIB1 because

nuSIB1-driven stress responses occur in the absence of LSD1. In this regard, we sought

if LSD1 also interacts with GLK1/2 to modulate the expression of PhANGs.

We then generated Arabidopsis WT transgenic plants overexpressing GREEN

FLUORESCENT PROTEIN (GFP)-tagged LSD1 under the control of the CaMV 35S

promoter (35S) (hereafter *oxLSD1*) to unveil putative LSD1-associated proteins. The

immunoblot assay detected the LSD1-GFP fusion protein at the predicted molecular

mass of approximately 46 kD using an anti-GFP antibody (Supplemental Figure 1).

Next, using GFP antibody-conjugated magnetic beads, we co-immunoprecipitated

LSD1-GFP and its putative associated proteins from the transgenic plants. The trypsin-

digested protein samples were then subjected to tandem mass spectrometry (MS)

analyses. The co-immunoprecipitation (Co-IP) coupled to MS analysis using three

independent biological replicates identified 217 proteins, which were detected in at

least two independent biological replicates, but absent in protein samples of WT and

GFP-overexpressing transgenic plants (*35S:GFP*) (Supplemental Dataset 1).

Accordingly, among the 217 proteins, we identified both GLK1 and GLK2

(Supplemental Dataset 1). In Arabidopsis, GLK1 and its homolog GLK2 share around

50% amino acid sequence identity. Both contain two conserved domains, a DNA-

binding domain (DBD) and a GLK1/2-specific C-terminal GCT-box (Supplemental

Figure 2) (Fitter et al., 2002; Rossini et al., 2001). A domain comparison between GLK1

and GLK2 shows a 90% and 79% identity, respectively (Bravo-Garcia et al., 2009).

Therefore, it is not surprising that both GLK1 and GLK2 were detected as putative

LSD1-associated proteins. Next, we performed a bimolecular fluorescence

complementation (BiFC) assay in *Nicotiana benthamiana* (*N. benthamiana*) leaves.

Consistent with the previous report (Czarnocka et al., 2017), we confirmed the LSD1-

LSD1 interaction in the nucleus, as evident in the overlapped signals detected from

YELLOW FLUORESCENT PROTEIN (YFP) and blue-fluorescent DNA stain 4', 6-

diamidino-2-phenylindole (DAPI)-stained nucleus, as well as in the cytosol (Figure 1A).

Similarly, we observed a YFP signal in *N. benthamiana* leaf coexpressing LSD1-YFP[N]

and GLK1 (or GLK2)-YFP[C] (Figure 1A). All YFP signals were exclusively observed

in the nucleus (Figure 1A). The Co-IP and an ensuing immunoblot assay further

corroborated the LSD1-GLK interaction (Figure 1B). We also purified full-length

recombinant proteins of LSD1, GLK1, and GLK2 expressed in *Escherichia coli*.

Subsequent gel filtration assays demonstrated that GLK1 (Figure 1C and 1E) and GLK2

(Figure 1D and 1E) form a complex with LSD1, as shown by their co-migration.

Figure 1. LSD1 interacts with GLK1 and GLK2.

(A) Bimolecular fluorescence complementation (BiFC) assays. The GLK1 or GLK2 fused with the C-
terminal part of YFP (YFP^C) were coexpressed with the N-terminal part of the YFP (YFP^N) fused with
LSD1 in *N. benthamiana* leaves. The combinations of LSD1-YFP^N + LSD1-YFP^C and LSD1-YFP^N +
YFP^C were used as a positive and negative control, respectively. DAPI was used to stain the nucleus. All
images were taken at the same scale (scale bars: 25 μm).
(B) Co-immunoprecipitation (Co-IP) analyses using *N. benthamiana* leaves transiently coexpressing
LSD1-GFP and GLK1-Myc (or GLK2-Myc). Co-IP was performed using GFP-Trap beads, and the
interaction was evaluated by using Myc antibody.
(C–E) Gel filtration assays showing *in vitro* interaction between LSD1 and GLK proteins expressed in
E. coli. Gel filtration profiles of LSD1, GLK1, and LSD1-GLK1 complex **(C)** and of LSD1, GLK2, and
LSD1-GLK2 complex **(D)**. A$_{280}$(mAU), micro-ultraviolet absorbance at the wavelength of 280 nm.
Coomassie blue staining of the peak fractions following SDS-PAGE **(E)**. Numbers on top of SDS-PAGE
panels indicate elution volume (mL). M, molecular weight ladder (kD).

LSD1 interacts with GLK1 and GLK2 through the proline-rich domain

We then generated truncated GLK variants to determine which domain is required for the interaction with LSD1. Prior to the interaction analysis, GLK1/2 and their variants lacking either DBD, potential proline-rich domain (PRD, located between DBD and GCT-box; see discussion), or GCT-box were C-terminally fused with GFP to monitor their nuclear localization (Figure 2A; Supplemental Figure 2). Following transient expression, all intact and variants of Arabidopsis GLK1/2 localized in the nucleus in *N. benthamiana* leaves but with a weak cytosolic GFP signal of GCT-box-deleted GLK1 and GLK2 (Supplemental Figure 3). To examine their interaction with Arabidopsis LSD1 protein, various combinations of BiFC constructs, as shown in Figure 2A, were expressed in *N. benthamiana* leaves to observe their interactions under the confocal microscope. The result clearly showed that the PRD of GLK1/2 is indispensable for the interaction with LSD1 (Figure 2B), further verified by Co-IP analyses (Figure 2C). We then generated GLK1/2 variants by C-terminal serial deletions to ascertain the significance of PRD for the interaction (Supplemental Figure 4A). All GFP-tagged proteins transiently expressed in *N. benthamiana* leaves were localized to the nucleus (Supplemental Figure 4B). The resulting BiFC and Co-IP analyses confirmed the critical role of PRD for LSD1 interaction, as evidenced by the lack of YFP signal when coexpressing LSD1 and GLK1/2 variants lacking the PRD-including C-terminal part (Supplemental Figure 4C and 4D).

Figure 2. PRD is indispensable for the interaction with LSD1.

(A) Schematic diagrams show intact GLK1/2 as well as their domain-deleted variants.

(B) BiFC assays. Each of intact and domain-deleted GLK variants fused with YFP^C was coexpressed with LSD1 fused with YFP^N in *N. benthamiana* leaves. DAPI was used to stain the nucleus. All images were taken at the same scale (scale bars: 10 μm).

(C) Co-IP analyses using *N. benthamiana* leaves transiently coexpressing LSD1-GFP with the indicated domain-deleted variant of GLK1/2 fused with Myc-tag.

Loss of *LSD1* upregulates GLK target genes

Given that GLKs promote the expression of PhANGs (Waters et al., 2009), it is plausible that loss of LSD1 may primarily affect their abundance. To identify affected genes either by loss of LSD1 or of GLK1/2, we compared the RNA sequencing (RNA-seq) data of *lsd1* versus WT (Lv et al., 2019) and *glk1 glk2* versus WT (Ni et al., 2017). As shown in Supplemental Figure 5A, a total of 91 genes (Supplemental Dataset 2) are shared between the upregulated genes (395, at least twofold) in *lsd1* versus WT

(Supplemental Dataset 3) and the downregulated genes (936, at least twofold) in *glk1*

glk2 versus WT (Supplemental Dataset 4). The Gene Ontology enrichment analysis

with the 91 genes for the biological process revealed that photosynthesis and light-

harvesting in PSII are over-represented (*P*-value=1.42E-07) (Supplemental Figure 5B;

Supplemental Dataset 5). The potentiated expression of PhANGs in *lsd1* versus WT

plants is indicative of a negative role of LSD1 in GLK activity. To this end, we also

examined the transcript abundance of PhANGs in WT, *glk1 glk2*, and two independent

oxLSD1 transgenic lines (Supplemental Figure 6A) using reverse transcription-

quantitative PCR (RT-qPCR). The results indicated that the examined GLK target

genes, such as genes encoding light-harvesting chlorophyll *a/b* binding proteins

(LHCBs in PSII and LHCA1 in PSI) and chlorophyll synthesis enzymes were

substantially repressed in *oxLSD1* relative to WT plants (Figure 3A). *oxLSD1* plants

exhibited comparable levels of *GLK1* and *GLK2* transcripts relative to WT

(Supplemental Figure 6B), implying that the repression of PhANGs likely resulted from

the post-translational regulation of GLK1/2. Remarkably, the overexpression of LSD1-

GFP fusion proteins in WT prematurely terminated chloroplast development, which is

reminiscent of the phenotype observed in *glk1 glk2* double mutant (Figure 3B and 3C).

Some mesophyll cells with nearly undetectable LSD1-GFP signals showed WT-like

chloroplasts (Figure 3B). One explanation might be an ectopic cosuppression of the

transgene, which also dilutes the molecular phenotypes (e.g., the transcript levels of

PhANGs) in the examined leaf tissue. Two independent *oxLSD1* lines with higher *LSD1*

transgene expression than the endogenous *LSD1* in WT plants (Figure 3D) exhibited

similar phenotypes, such as partial cosuppression of the transgene, defect in chloroplast

biogenesis, and reduced levels of LHCB proteins (Figure 3B, 3C, and 3E). Regardless

of the promoters used (35S and native), all stable transgenic lines (over 20 lines)

showed nearly undetectable or detectable GFP signals but with partial cosuppression

(data not shown).

Figure 3. Overexpression of LSD1 negatively affects the expression of GLK target genes and chloroplast biogenesis.

(A) Relative transcript levels of GLK1 and GLK2 target genes, such as *LHCBs*, *LHCA1*, and chlorophyll synthesis genes including *glutamyl tRNA reductase* (*HEMA1/Glu-TR*), *genome uncoupled 4* (*GUN4*), *magnesium chelatase H subunit* (*CHLH/GUN5*), *protochlorophyllide oxidoreductase* (*POR*) *A*, *PORB*, *PORC*, and *chlorophyllide a oxygenase* (*CAO*) were examined in CL-grown 24-d-old WT, *glk1 glk2*, and *oxLSD1* (#1 and #5) plants using RT-qPCR.

(B) Plant phenotypes (left panels) and GFP fluorescence (green) of LSD1-GFP fusion proteins merged with chlorophyll autofluorescence signals (red; middle and right panels) in 24-d-old WT, *glk1 glk2*, and transgenic WT plants overexpressing GFP alone (*GFP*) or LSD1-GFP (*oxLSD1* #1 and #5) grown under CL conditions. The small white square boxes in the middle panels were enlarged (right panels).

(C) Means of chloroplast diameter. At least three confocal images taken from three independent leaves were used to measure the chloroplast diameter. For *oxLSD1*, only mesophyll cells with detectable GFP signals were chosen.

(D and E) Relative levels of *LSD1* transcript **(D)** and LHCB proteins **(E)** in 24-d-old CL-grown plants of WT, *glk1 glk2*, and *oxLSD1* #1 and #5. UGPase was used as a loading control for the immunoblot analysis in **(E)**. Numbers at the bottom of each immunoblot result indicate the relative quantities of LHCB1 or LHCB3 proteins against the control signal of UGPase. For the RT-qPCR analyses In **(A)** and **(D)**, *ACT2* was used as an internal standard. The value represents means ± standard deviation (SD) (n=3). Lowercase letters indicate statistically significant differences between mean values (P < 0.05, one-way ANOVA with posthoc Tukey's HSD test).

(E) ChIP–qPCR results showing the effect of LSD1 overexpression on GLK1 binding to the promoter regions of its target genes (*LHCB1.4*, *LHCB3,* and *LHCB6*). Myc-tagged GLK1 (GLK1-Myc) was transiently expressed with (+ LSD1) or without LSD1-RFP (– LSD1) in Arabidopsis leaf protoplasts isolated from *lsd1 glk1 glk2* triple mutant. The enrichment value was normalized to the input sample,

277 representing means ± SD from two independent ChIP assays. Asterisks denote statistically significant
278 differences by Student's *t*-test ($P < 0.01$) from the value of − LSD1.

LSD1 inhibits the DNA-binding activity of GLK1

Regarding that nuclear-localized LSD1 acts as a transcription coregulator and that

LSD1 interacts with GLKs, it is conceivable that LSD1 might directly regulate the

DNA-binding activity of GLK1/2. Therefore, we performed a chromatin

immunoprecipitation (ChIP) coupled with a qPCR analysis. The relative activity of

GLK1 towards its target promoters was examined in the presence or absence of LSD1

using protoplasts isolated from the rosette leaves of *lsd1 glk1 glk2* triple mutant plants.

Since GLK1 and GLK2 are highly unstable (Tokumaru et al., 2017; Waters et al., 2008),

we used a protoplast transient expression system to ensure sufficient protein expression

to elucidate the impact of LSD1 on GLK1 function. The *35S:GLK1-Myc* was transiently

coexpressed with either *35S:RED FLUORESCENT PROTEIN* (*RFP*) or *35S:LSD1-*

RFP in the protoplasts. ChIP assays were then performed with nuclear lysis from

transfected protoplasts. With anti-Myc antibody-conjugated agarose beads, the Myc-

tagged protein-DNA complex was pulled down. The immunoprecipitated DNA was

then analyzed using qPCR to compare the DNA-binding activity of GLK1 in the

presence or absence of LSD1. Afterward, we examined the relative expression levels

of well-established GLK target genes such as *LHCB1.4, LHCB3,* and *LHCB6* (Waters

et al., 2009). The results demonstrated that the presence of LSD1 markedly diminished

the DNA-binding activity of GLK1 to promoters of these *LHCB* genes (Figure 3F).

Loss of LSD1 potentiates $^1O_2$-triggered EX1-dependent RS in *flu* mutant

We next validated the above ChIP assay result *in planta*. Considering the positive

regulation of chlorophyll synthesis by GLK1 (Waters et al., 2009) and the repression

of GLK activity by LSD1 (Figure 3A and 3F), it is tempting to hypothesize that loss of

LSD1 might increase the rate of chlorophyll biosynthesis. By revisiting the previously

published RNA-seq data (Lv et al., 2019), we noticed that a set of chlorophyll synthesis

genes including *glutamyl tRNA reductase* (*HEMA1/Glu-TR*), *genome uncoupled 4*

(*GUN4*), *magnesium chelatase H subunit* (*GUN5/CHLH*), *magnesium protoporphyrin*

IX monomethyl ester cyclase (*CHL27/CRD1*), *protochlorophyllide oxidoreductase A* (*PORA)*, and *chlorophyllide a oxygenase* (*CAO*) were markedly upregulated in *lsd1* before the onset of RCD (Figure 4A and 4B). We then analyzed the 5-aminolevulinic acid (5-ALA, the common precursor of all tetrapyrroles) synthesis rate in light-grown *lsd1* mutant plants treated with levulinic acid (LA) (Nandi and Shemin, 1968), a competitive chemical inhibitor of 5-ALA dehydratase that catalyzes the synthesis of porphobilinogen through the asymmetric condensation of two 5-ALA molecules (Figure 4A). It is important to note that the 5-ALA synthesis is the rate-limiting step for chlorophyll synthesis (Beale and Castelfranco, 1974; Hou et al., 2019) (Figure 4A). In view of the fact that FLUORESCENT (FLU) protein directly represses Glu-TR activity to inhibit protochlorophyllide (Pchlide) accumulation in the dark (Goslings et al., 2004) (Figure 4A) and that *flu* mutant plants exhibit a higher 5-ALA synthesis rate in the presence of LA under continuous light (CL) conditions (Goslings et al., 2004), we used *flu* as a positive control.

The 5-ALA synthesis rate was almost comparable in *lsd1* and *flu* seedlings in the presence of LA (Figure 4C). Although it is yet unclear whether the transcriptional upregulation of *HEMA1* is responsible for the 5-ALA accumulation in *lsd1,* the concurrent loss of both *FLU* and *LSD1* further increased the 5-ALA synthesis rate under CL conditions (Figure 4C). Since Pchlide levels in the dark would indirectly reflect chlorophyll biosynthesis rate owing to the absence of the enzyme(s) involved in Pchlide turnover (Forreiter and Apel, 1993), we measured Pchlide levels in dark-incubated plants of WT, *flu*, *lsd1,* and *lsd1 flu* using high-performance liquid chromatography analysis. As anticipated, Pchlide was highly upregulated in the *flu* mutant background, as demonstrated earlier (Meskauskiene et al., 2001) (Figure 4D). The loss of LSD1 raises the Pchlide level (approximately a 1.5-fold increase) in both WT and *flu* mutant backgrounds. The presence of FLU protein in *lsd1* seems to prevent the drastic accumulation of Pchlide in the dark. Collectively, these results corroborate the negative role of LSD1 towards GLK activity, which seems to be, at least in part, required for tetrapyrrole homeostasis.

Figure 4. *LSD1* mutation leads to the transcriptional upregulation of chlorophyll synthesis genes.

(A) A schematic representation of the chlorophyll synthesis pathway.

(B) The expression levels of chlorophyll synthesis genes represented in **(A)** were obtained from our

previous study (Lv et al., 2019), and the relative transcript levels of chlorophyll synthesis genes in 17-d-

old CL-grown *lsd1* mutants compared to wild type (WT) are represented. Error bars indicate SD (n=3).

Asterisks indicate statistically significant differences (*$P < 0.05$; **$P < 0.01$) in the *lsd1* mutant

determined by Student's *t*-test relative to wild type. ND: non-detected. Red triangles in **(A)** and **(B)**

indicate the significantly upregulated genes in the *lsd1* mutant compared to WT.

(C) Levels of 5-ALA synthesis rate under CL conditions. The 5-ALA synthesis rate was measured in 16-

d-old plants of WT, *flu*, *lsd1*, and *lsd1 flu*.

(D) Relative levels of protochlorophyllide (Pchlide) in 10-d-old plants of WT, *flu*, *lsd1*, and *lsd1 flu*

grown under CL and then transferred to the dark for 8 hours. Values in **(C)** and **(D)** represent means ±

SD (n=3). Lowercase letters indicate significant differences between the indicated genotypes ($P < 0.05$,

one-way ANOVA with posthoc Tukey's HSD test).

We then hypothesized that the SA-induced nuSIB1-GLK-driven upregulation of

PhANGs and the higher chlorophyll synthesis rate by *FLU* mutation might further

enhance $^1O_2$ levels in chloroplasts in *lsd1 flu* plants grown under CL conditions before

the onset of RCD. Indeed, *lsd1 flu* double mutant plants exhibited accelerated RCD

than *lsd1* plants (Figure 5A), which was found to be radically reduced in *lsd1 flu ex1*,

indicating $^1O_2$ was the prime cause of the reinforced RCD in *lsd1 flu*. This result

coincided with the intensity of maximum fluorescence of PSII (Fm) (Figure 5A), PSII

maximum efficiency (Fv/Fm) (Figure 5B), and the abundance of $^1O_2$-responsive genes

(SORGs) (Dogra et al., 2017) (Figure 5C).

Figure 5. Loss of LSD1 potentiates $^1O_2$-triggered EX1-dependent cell death in *flu* mutant.

(A) WT, *flu*, *lsd1*, *lsd1 flu*, and *lsd1 flu ex1* plants were grown on Murashige and Skoog (MS) medium

under CL conditions (100 μmol·m^{-2}·s^{-1}). The RCD phenotype (Pheno, left panels) and the chlorophyll

maximum fluorescence (Fm) of PSII (middle panel) were monitored in the whole plants at the indicated

time points. The dead cells in the first or second leaves from the genotypes were visualized via trypan

blue staining (right panel). Images are representative phenotypes.

(B) The first or second leaves from each genotype were harvested at the indicated time points to measure

the maximum photochemical efficiency of PSII (Fv/Fm). Data represent means ± SD (n=10).

(C) Expression levels of selected $^1O_2$-responsive genes (SORGs) were examined by RT-qPCR in 18-d-

old plants. *ACT2* was used as an internal standard. Data represent means ± SD (n=3). Lowercase letters

in **(B)** and **(C)** indicate statistically significant differences between mean values (P < 0.05, one-way

ANOVA with post-hoc Tukey's HSD test).

SA-induced SIB1 interrupts LSD1-GLK1 interaction

Since SIB1-mediated genomes uncoupled expression of PhANGs and PhAPGs largely

contributes to *lsd1* RCD via $^1O_2$ signaling (Lv et al., 2019), we hypothesized that

nuSIB1 would counteractively modulate LSD1-GLK interaction to reinforce the

expression of PhANGs. Considering its rapid turnover via UPS (Li et al., 2020), nuSIB1

may promptly intervene in this LSD1-GLK interaction, resulting in nuSIB1-GLK

interaction and reinforced expression of PhANGs, thereby contributing to cell death

(Lv et al., 2019). Alternatively, SA per se may interfere with LSD1-GLK interaction,

for instance, through alteration of protein conformation of LSD1 or GLKs or both. In

fact, a previous report showed a redox-sensitive reconfiguration of LSD1 and

concurrent change of its interactome (Czarnocka et al., 2017). Thus, we examined how

SA impacts the LSD1-GLK1 interaction in Arabidopsis leaf protoplasts isolated from

WT and *sib1* mutant plants. The result that SA significantly hindered the LSD1-GLK1

interaction in WT but not in *sib1* (Figure 6A) suggested that nuSIB1 rather than SA per

se interrupts LSD1-GLK1 interaction. To further elucidate an antagonistic action of

nuSIB1 towards LSD1-GLK1 interaction, a dose-dependent impact of nuSIB1 was

examined. For this, LSD1-GFP and GLK1-Myc were transiently coexpressed in

Arabidopsis leaf protoplasts, along with different amounts of free RFP or SIB1-RFP. It

should be noted that increasing doses of RFP or SIB1-RFP reduce the expression of

LSD1-GFP and GLK1-Myc, probably as a consequence of diminished transfection

efficiency due to the presence of the additional constructs (Figure 6B). Nonetheless, the

relative amount of GLK1-Myc protein co-immunoprecipitated with LSD1-GFP was

quantified using ImageJ following immunoblot analysis (Figure 6C). The results

showed a SIB1 dose-dependent inhibition of the LSD1-GLK1 interaction.

Figure 6. SA-induced SIB1 intervenes in LSD1-GLK interaction.
(A) The effect of the SA-induced nuSIB1 on the LSD1-GLK1 interaction. For Co-IP analyses, *35S:LSD1-GFP* and *35S:GLK1-Myc* were transiently coexpressed in Arabidopsis leaf protoplasts. The protoplasts were treated with either mock or 0.2 mM SA for 5 hours.
(B) The dose-dependent impact of SIB1 on the LSD1-GLK1 interaction. As indicated, *35S:LSD1-GFP* and *35S:GLK1-Myc* were coexpressed in Arabidopsis leaf protoplasts isolated from WT plants together with different amounts (10 µg, 20 µg, or 40 µg, respectively) of a plasmid containing either free *35S:RFP* or *35S:SIB1-RFP*. The subsequent Co-IP and immunoblot results are shown. Three independent experiments were conducted with similar results, and representative results are shown in **(A)** and **(B)**.
(C) The signal intensity of eluted GLK1-Myc (from triplicate immunoblots in **B**) versus its input signal was quantified using the ImageJ software. Data are means ± SD (n=3). Asterisks denote statistically significant differences by Student's *t*-test (*$P < 0.05$, **$P < 0.01$, NS: not significant).

One possible scenario for the nuSIB1-dependent interruption of LSD1-GLK

interaction is that SA-induced nuSIB1 may directly interact with LSD1, releasing

GLK1 and GLK2 in the nucleus. The free GLK1/2 may interact with excess nuSIB1,

promoting the expression of PhANGs. However, while the LSD1-LSD1 interaction was

apparent, no interaction between LSD1 and nuSIB1 was observed (Supplemental

Figure 7). Then we assumed that SA-induced nuSIB1 might compete with LSD1 to

bind to the PRD of GLK1 and GLK2. We then carried out Co-IP analyses to investigate

if PRD is required for the interaction with SIB1. The result showed that the N-terminal

region excluding all three domains is sufficient to interact with nuSIB1 (Supplemental

Figure 8A and 8B). Since the N-terminal part contains a nuclear localization signal

(Zhang et al., 2021a), we ended further defining the minimum length of the N-terminal

necessitated for the interaction with nuSIB1. It is likely that SA-induced nuSIB1

competitively interacts with GLK1 and GLK2 through the N-terminal part, which

consequently enhances the expression of PhANGs and the $^1O_2$ level, thereby activating

an EX1-mediated cell death response (Figure 7). The rapid turnover of nuSIB1 via UPS

(Li et al., 2020) might result in LSD1-GLK interaction and restore the expression levels

of PhANGs.

Figure 7. Proposed model elucidating the counteractive regulation of GLKs by LSD1 and nuSIB1.
LSD1-GLK interaction is required for negative regulation of GLK activity to fine-tune the expression of
PhANGs, including LHCBs and chlorophyll synthesis genes. Under SA-increasing stress conditions, the
NPR1-induced and NTA-stabilized nuSIB1 intervenes in LSD1-GLK interaction to reinforce the
expression of PhANGs, while the cpSIB1 represses the expression of PhAPGs by interacting with SIG1
(Li et al., 2020; Lv et al., 2019; Morikawa et al., 2002). The resulting uncoupled expression of PhANGs
and PhAPGs aggravates PSII photoinhibition and increases $^1O_2$ level in chloroplasts, enabling EX1-
mediated retrograde signaling to activate the expression of SORGs and cell death response (Kim et al.,
2012; Lv et al., 2019). While NTA renders nuSIB1 more stable, UPS promotes the proteolysis of nuSIB1
(Li et al., 2020), restoring LSD1-GLK interaction to avoid an excess of PhANG expression and $^1O_2$
accumulation. The counteractive regulation of GLKs by nuSIB1 and LSD1, along with post-translational
regulation of nuSIB1 stability, seems vital to modulate $^1O_2$ levels in chloroplasts during and after SA-
increasing stress conditions.

DISCUSSION

Besides their essential role in chloroplast biogenesis and photosynthesis, multiple lines

of evidence demonstrate that GLKs function in plant stress responses, evoking an

intriguing proposal that GLK may serve as a master switch in synchronously regulating

photosynthesis and stress responses. We previously reported that the positive regulator

of SA signaling and transcription coregulator nuSIB1 interacts with GLKs and

WRKY33 to reinforce the expression of PhANGs and SA-responsive genes,

respectively, upon an increase in cellular SA level (Li et al., 2020; Lv et al., 2019). On

the contrary, cpSIB1 interacts with SIG1 polymerase to repress the expression of

PhAPGs (Lv et al., 2019; Morikawa et al., 2002; Xie et al., 2010). The genomes-

uncoupled expression of PhANGs and PhAPGs heightens the PSII photoinhibition,

thereby escalating the highly reactive oxygen species, specifically $^1O_2$. $^1O_2$ then

contributes to SA-driven plant stress responses via EX1-mediated RS, which is shown

to reinforce RCD phenotype in *lsd1* mutant (Dogra et al., 2019; Lv et al., 2019). Since

the SA receptor NPR1 is required to induce the expression of *SIB1* (Xie et al., 2010),

EX1-mediated $^1O_2$ signaling is likely to be one of the downstream events led by SA and

NPR1.

We now showed that LSD1 interacts with GLK1 and GLK2 TFs in the nucleus

(Figure 1A). Besides their transcriptional regulation (e.g., by GUN1-mediated RS),

multiple proteins post-translationally modulate GLK activity (Tang et al., 2016;

Tokumaru et al., 2017; Zhang et al., 2021a). The C-terminal GCT-box drives GLK

homo- or hetero-dimerization in maize (Rossini et al., 2001). The *turnip yellow mosaic*

virus (TYMV) protein P69 binds to the GLK1/2 GCT-box, repressing PhANGs and

chloroplast biogenesis in Arabidopsis (Ni et al., 2017). On the contrary,

BRASSINOSTEROID INSENSITIVE 2 (BIN2)-dependent GLK phosphorylation

promotes chloroplast biogenesis by stabilizing GLK proteins in Arabidopsis (Zhang et

al., 2021a). These reports indicate that both DBD and GCT-box in GLK1/2 are involved

in protein-protein interaction. Notably, the interdomain region of GLK1 and GLK2 are

proline-enriched (Figure 2A; Supplemental Figure 2). Since proline residues provide

protein-docking sites (Siligardi and Drake, 1995; Zarrinpar et al., 2003), we anticipated

the PRD as an additional candidate domain required for the interaction with LSD1. The

ensuing BiFC and Co-IP assays verified that the PRD is central for interacting with

LSD1. GLK1 and GLK2 lacking DBD and GCT-box but retaining PRD interacted with

LSD1, but complete loss of PRD abolished these interactions (Figure 2B and

2C). Besides, the association of GLK2 with the CUL4-DDB1-based E3 ligase complex

promotes UPS-mediated GLK2 turnover in tomato (Tang et al., 2016). The COP1 and

UPS-mediated GLK1 degradation was also reported in Arabidopsis plants with long-

term abscisic acid (ABA) treatment (Lee et al., 2021). Interestingly, one latest work

showed that WRKY75 directly represses *GLK* expression during leaf senescence

(Zhang et al., 2021b). The ABA-induced SIB1 and its close homolog SIB2 interact with

and inhibit WRKY75 activity, enabling the expression of GLKs in response to ABA.

The antagonistic regulation of GLKs expression by SIB1/2 and WRKY75 was

proposed to be essential in controlling ABA-mediated leaf senescence and seed

germination. These findings by other groups and our data suggest that the GLKs are

common targets of development or stress signaling to modulate chloroplast homeostasis.

As emerging notion strongly supports the role of chloroplasts as environmental sensors,

such modulation of GLK activity and stability would also significantly affect

chloroplast-mediated plant stress responses.

Loss of LSD1 potentiated the expression of GLK target genes such as PhANGs

(Supplemental Figure 5A and 5B) and increased the 5-ALA synthesis rate compared to

WT plants (Figure 4C). Conversely, LSD1 overexpression repressed the expression of

PhANGs (Figure 3A). These results were consistent with *oxLSD1* plant phenotypes

exhibiting prematurely terminated chloroplast development and reduced LHCB levels

(Figure 3B-3E). Consistently, LSD1 overexpression repressed GLK1 binding activity

to its target promoters (Figure 3F). The effects of loss- and gain-of-function of LSD1

towards the expression of GLK target genes also suggest a steady-state LSD1-GLK

interaction in WT plants grown under normal growth conditions. Given that the

elevated expression of PhANGs directed by nuSIB1-GLK interaction contributes to

lsd1 RCD (Lv et al., 2019), it was tempting to hypothesize that SA-induced nuSIB1

interferes with LSD1-GLK interaction. Indeed, the Co-IP assay confirmed the negative

impact of nuSIB1 accumulation on LSD1-GLK interaction (Figure 6). It has been

shown that the PRD domain provides the sequence-specific docking site for interacting

proteins without the requirement of a high-affinity interaction (Saraste and Musacchio,

1994; Siligardi and Drake, 1995; Zarrinpar et al., 2003). The sequence-specific but low-

affinity interaction at the proline-rich region might allow a highly reversible interaction

between LSD1-GLKs, enabling SA-induced nuSIB1 to rapidly intervene in this

interaction through the N-terminus of GLKs (Supplemental Figure 8). Such versatile

regulation of nuSIB1 stability and an antagonistic mode of action of nuSIB1 and LSD1

towards GLK1/2 might be vital to maintain $^1O_2$ homeostasis in chloroplasts and to

induce SA-driven stress responses under fluctuating environmental conditions (Figure

7).

Our findings also raise a plausible idea that GUN1-mediated RS would largely

contribute to plant stress responses because the signaling primarily represses the

expression of *GLK1* and *GLK2* once the foliar plastid function is interrupted (see

introduction). Consistently, *gun1* mutant plants exhibit an increased susceptibility

towards heat, water, drought, cold, and high-light stresses with enhanced cellular ROS

levels (Cheng et al., 2011; Miller et al., 2007; Tang et al., 2014; Zhang et al., 2013;

Zhang et al., 2011). The multifaceted interactions between GLK1/2 and the antagonistic

modules nuSIB1 and LSD1, as well as other stress-related proteins, may be accountable

for the altered *gun1* phenotype to various stress factors. The enhanced expression of

GLK1 and *GLK2* may increase the level of $^1O_2$ if nuSIB1 is accumulated and intervene

in LSD1-GLK interaction in *gun1*. The $^1O_2$-triggered EX1-mediated RS may then

modulate plant stress responses in *gun1* mutant plants under SA-increasing stress

conditions. In this regard, a new study of *gun1* may provide further insight into how

chloroplast RS pathways mediated by GUN1 and EX1 coordinate SA-mediated plant

stress responses through GLK1/2 and $^1O_2$, respectively.

Methods

Plant materials and growth conditions

The seeds used in this study were derived from *Arabidopsis thaliana* Columbia-0 (Col-

0) ecotype and were harvested from plants grown under continuous light (CL; 100

$\mu mol \cdot m^{-2} \cdot s^{-1}$) at 22 ± 2 °C. Arabidopsis mutant seeds used in this study, including *lsd1-*

2 (SALK_042687) (Lv et al., 2019), *glk1 glk2* (*Atglk1.1; Atglk2.1*) (Fitter et al., 2002),

and *ex1* (SALK_002088) (Lee et al., 2007) were obtained from the Nottingham

Arabidopsis Stock Centre (NASC). *flu5c* has been described previously (Meskauskiene

et al., 2001). The double and triple mutants in the *lsd1-2* background including *lsd1 flu*,

lsd1 flu ex1, and *lsd1 glk1 glk2* were generated by crossing the homozygous plants. The

genotypes of all mutants were confirmed by PCR-based analyses. Primer sequences

used for PCR are listed in Supplemental Table 1.

Seeds were surface sterilized with 70% (v/v) ethanol containing 0.05% (v/v) Triton

X-100 (Sigma-Aldrich) for 10 min and washed five times with sterile distilled water.

The sterile seeds were plated on Murashige and Skoog (MS) medium (Duchefa

Biochemie) with 0.7% (w/v) agar (Duchefa Biochemie) and stratified at 4 °C in

darkness for two days prior to placing in a growth chamber (CU-41L4; Percival

Scientific) with CL condition.

Generation of *LSD1* overexpression lines

The stop-codon-less *LSD1* coding sequence (CDS) was cloned into the modified

pCAMBIA3300 binary vector containing the 35S promoter, a NcoI restriction site, and

the *EGFP*. Arabidopsis stable transgenic lines were generated by a floral dip

transformation procedure (Clough and Bent, 1998) with *Agrobacterium tumefaciens*

strain GV3101. Homozygous transgenic lines were selected on MS medium containing

12.5 mg/L glufosinate-ammonium (Sigma-Aldrich).

RNA extraction and RT-quantitative PCR (RT-qPCR)

Total RNA was isolated from leaf tissues using the Spectrum Plant Total RNA Kit

(Sigma-Aldrich) according to the manufacturer's instructions. The concentration of

RNA was determined using the ultraviolet-visible spectrophotometer (NanoDrop™,

Thermo Fisher Scientific), and the quality of RNA was evaluated by measuring the

A260/A280 ratio. cDNA synthesis was performed with 1 μg of total RNA using the

PrimeScript™ RT Reagent Kit (Takara) following the manufacturer's instructions. The

RT-qPCR was performed on QuantStudio™ Flex Real-Time PCR System (Applied

Biosystems) using iTaq Universal SYBR Green PCR master mix (Bio-Rad). The

relative transcript level was calculated by the ddCt method (Livak and Schmittgen,

2001) and normalized to the *ACTIN2* (AT3G18780) gene transcript level. The

sequences of the primers used for RT-qPCR are listed in Supplemental Table 1.

Co-immunoprecipitation (Co-IP) assay

Co-IP assays were performed using *Nicotiana benthamiana* or Arabidopsis leaf protoplasts transiently coexpressed with the indicated combination of proteins. For the Co-IP assays in *N. benthamiana*, the *35S:LSD1-sGFP*, *35S:SIB1-sGFP*, *35S:GLK1-4×Myc,* and *35S:GLK2-4×Myc* constructs were created as described previously (Lv et al., 2019). Briefly, pDONR221/Zeo entry vector (Thermo Scientific) containing the stop codon-less full-length CDS of *LSD1, SIB1, GLK1*, or *GLK2* was recombined into the destination vector pGWB605 for C-terminal fusion with sGFP or into pGWB617 for C-terminal fusion with 4×Myc through the Gateway LR reaction (Thermo Scientific). For the *35S:GLK1-4×Myc* (or *sGFP*) and *35S:GLK2-4×Myc* (or *sGFP*) constructs, a linker DNA encoding Gly-Gly-Ser-Gly-Gly-Ser was added between *4xMyc* (or *sGFP*) tag and GLK1 or GLK2 to increase conformational flexibility of the fusion protein as described previously (Tokumaru et al., 2017). The same procedures were used to create the constructs containing CDSs encoding domain-deleted or C-terminally truncated variants of GLK1 and GLK2. The different combinations of selected vectors were coexpressed in 4-week-old *Nicotiana benthamiana* leaves by Agrobacterium-mediated leaf infiltration as previously described by Boruc et al. (2010). For the Co-IP assays in Arabidopsis leaf protoplasts, the *35S:LSD1-sGFP*, *35S:GLK1-4×Myc*, *35S:SIB1-RFP*, and *35S:LSD1-RFP* were cloned into the pSAT6 vector (Tzfira et al., 2005). The isolation and transfection of Arabidopsis leaf protoplasts were performed as described previously (Yoo et al., 2007). The indicated combination of vectors was cotransfected into protoplasts (3×10^6) isolated from 4-week-old plants of WT or *sib1*.

Total protein was extracted using an IP buffer containing 50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 0.5mM EDTA, 10% (v/v) glycerol, 1% (v/v) Nonidet P-40 (NP-40), 1% deoxycholate, 0.1% (w/v) SDS, $1 \times$ cOmplete protease inhibitor cocktail (Roche), 1 mM PMSF, and 50 μM MG132. The protein extracts were incubated with 20 μL of GFP-Trap magnetic agarose beads (GFP-TrapMA, Chromotek) for 2 h at 4 °C by vertical rotation (10 rpm). After incubation, the beads were washed five times with

the washing buffer containing 10 mM Tris-HCl (pH 7.5), 150 mM NaCl, 0.5 mM

EDTA, 1 mM PMSF, 50 μM MG132, and 1 × cOmplete protease inhibitor cocktail.

The immunoprecipitated proteins were then eluted with 2 × SDS protein sample buffer

[120 mM Tris-HCl (pH 6.8), 20% (v/v) glycerol, 4% (v/w) SDS, 0.04% (v/w)

bromophenol blue, and 10% (v/v) β-mercaptoethanol] for 10 min at 95 °C. The eluates

were subjected to 10% SDS-PAGE gels, and the interaction between coexpressed

proteins was examined by immunoblot analyses using a mouse anti-Myc monoclonal

antibody (1:10,000; Cell Signaling Technology), a rat anti-RFP monoclonal antibody

(1:10,000; Chromotek), and a mouse anti-GFP monoclonal antibody (1:5,000; Roche).

Protein Extraction and immunoblot analysis

Total proteins were extracted from 100 mg of foliar tissues with the IP buffer and

quantified with a Pierce BCA protein assay kit (Thermo Fisher Scientific). Afterward,

20 μg total protein was separated on 10% SDS-PAGE gels and blotted onto Immun-

Blot PVDF membrane (Bio-Rad). LSD1-GFP, LHCB1, and LHCB3 were

immunochemically detected with mouse anti-GFP (1:10,000; Roche), rabbit anti-

LHCB1 (1:5,000; Agrisera), and rabbit anti-LHCB3 (1:5,000; Agrisera) antibodies,

respectively. The UDP-glucose pyrophosphorylase (UGPase) detected with rabbit anti-

UGPase (1:3,000; Agrisera) was used as a loading control.

Confocal laser-scanning microscopy

611 The GFP, YFP, chlorophyll, and 4', 6'- diamidino-2-phenylindole (DAPI) fluorescence

612 signals were detected by confocal laser-scanning microscopy analysis using TCS SP8

613 (Leica Microsystems). All the images were obtained and processed with Leica LAS AF

614 Lite software, version 2.6.3 (Leica Microsystems).

Bimolecular fluorescence complementation (BiFC) assay

BiFC assays were conducted with a split-YFP system in *N. benthamiana* leaves, as

described previously (Lee et al., 2020; Lu et al., 2010). Briefly, the pDONR/Zeo entry

vectors (Thermo Fisher Scientific) containing CDSs lacking the termination codon of

intact forms, domain-deleted, or C-terminally truncated variants of GLK1 and GLK2

were recombined into the destination vector pGTQL1221 through Gateway LR reaction.

The same procedure was done to recombine the pDONR221/ZEO entry vector

containing the *LSD1* CDS lacking the terminal codon into the pGTQL1211. For the

BiFC assay, *A. tumefaciens* mixtures carrying the appropriate constructs were

infiltrated into 4-week-old *N. benthamiana* leaves. The presence of YFP fluorescence

signals was evaluated by confocal laser-scanning microscopy analysis.

ChIP-qPCR assays

ChIP assays were performed using Arabidopsis leaf protoplasts as described previously

(Lee et al., 2017; Lv et al., 2019; Yoo et al., 2007). Briefly, 1 mg of pSAT6 vectors

containing *35S:GLK1-4×Myc* DNA were transfected with or without pSAT6 vector

containing *35S:LSD1-RFP* DNA into Arabidopsis leaf protoplasts (2×10^7) isolated

from 4-week-old *lsd1 glk1 glk2* triple mutant plants grown under 10-h light/14-h dark

conditions at a light intensity of 100 μmol $m^{-2}s^{-1}$. Afterward, the protoplasts were

incubated at 24 °C for 16 h under dim light conditions. The protoplast chromatins were

crosslinked by 1% (v/v) formaldehyde in $1 \times$ PBS (pH 7.4) for 10 min and quenched

with 0.1 M glycine for 5 min. After isolating nuclei from the protoplasts, the chromatins

were sheared by sonication into an average size of around 500 bp. The lysates were

diluted with $10 \times$ ChIP dilution buffer [1% (v/v) Triton X-100, 2 mM EDTA, 20 mM

Tris-HCl (pH 8.0), 150 mM NaCl, 50 μM MG132, 1 mM PMSF, and $1 \times$ protease

inhibitor cocktail] and precleared by incubation with 50 μL Protein-A agarose

beads/Salmon sperm DNA (Millipore) at 4 °C for 1 h. The samples were then incubated

with anti-Myc monoclonal antibodies (1:10,000; Cell Signaling Technology) at 4 °C

overnight. To determine non-specific binding of DNA on beads, ChIP assays were also

performed without antibodies. After washing the beads, the immunocomplexes were

eluted with elution buffer containing 1% (w/v) SDS and 100 mM $NaHCO_3$. The eluates

were treated with proteinase K for 1 h at 37 °C after reverse cross-linking. The bound

DNA fragments were purified as previously described by Lee et al. (2017) and

precipitated with ethanol in the presence of glycogen. The purified DNA was dissolved

in water. qPCR analyses were performed on bound and input DNAs. The primers for

each tested gene are listed in Supplemental Table 1. The amount of DNA enriched by

the anti-Myc antibody was calculated in comparison with the respective input DNA

used for each ChIP analysis. Afterward, the enrichment was calculated by normalizing

against the corresponding control sample (without antibody).

Production of recombinant proteins

To produce recombinant proteins of LSD1, GLK1, and GLK2, the coding sequences

(CDS) of genes were cloned into the modified pET21b (Novagen) expression vector

after adding a cleavage site for the tobacco etch virus (TEV) protease to the 5' end of

the CDSs. The recombinant proteins with a cleavable N-terminal 10×His-MsyB tag

were expressed in *E. coli* BL21 (DE3). After culturing the cells at 37 °C until an OD_{600}

of 0.6, recombinant proteins were induced by adding 0.3 mM isopropyl-β-D-

thiogalactopyranoside (IPTG) for 12 h at 16 °C. Cells were pelleted by centrifugation

and resuspended with buffer A [(50 mM Tris (pH 8.0), 200 mM NaCl, and 1 mM

PMSF]. The cells were lysed by high-pressure homogenizer at 600-800 bar and then

centrifuged at 17,000 rpm for 50 min. Each soluble fraction was passed over a Ni-NTA

column (Novagen) and eluted with buffer containing 25 mM Tris (pH 8.0), 200 mM

NaCl, and 200 mM Imidazole. Subsequently, the eluates containing recombinant

proteins with 10×His-MsyB tag were further purified by an anion-exchange column

(Source-15Q; GE Healthcare). The 10×His-MsyB tag was cleaved by TEV protease at

4 °C overnight and removed by an anion-exchange column. Untagged recombinant

proteins were then concentrated and further purified by size-exclusion chromatography

(Superdex 200 Increase10/300 GL; GE Healthcare) in buffer containing 20 mM Tris

(pH 8.0), 200 mM NaCl, and 3 mM DTT. The peak fractions of each protein were

pooled together and used for gel filtration assay.

Gel filtration assay

The recombinant proteins purified as described above were subjected to gel filtration

assay (Superdex 200 Increase10/300 GL; GE Healthcare) in buffer containing 20 mM

Tris (pH 8.0), 200 mM NaCl, and 3 mM DTT. A mixture of the purified LSD1 and

GLK1 (or GLK2) proteins was incubated at 4 °C for 1 h before gel filtration. Samples

from relevant fractions were applied to SDS-PAGE and visualized by Coomassie blue

staining.

Measuring 5-ALA synthesis rate

The 5-ALA synthesis rate was quantified as previously described (Goslings et al., 2004).

CL-grown 16-day-old plants of WT, *flu*, *lsd1,* and *lsd1 flu* were vacuum-infiltrated for

5 min with an 80 mM levulinic acid (Sigma) solution containing 10mM KH_2PO_4 (pH

7.2) and 0.5% (v/v) Tween 20. After 1 hour incubation at room temperature under CL,

samples were immediately frozen in liquid nitrogen and then homogenized in 4% (v/v)

TCA. The homogenates were lysed at 95 °C for 15 min, cooled on ice for 2 min, and

filtrated with 0.45 μm cellulose acetate membrane filters (Sterlitech). The filtrated

lysates were neutralized with an equal volume of 0.5 M NaH_2PO_4 (pH 7.5). Afterward,

ethylacetoacetate (1/5) was added and then the samples were incubated at 95 °C for 10

min. After cooling on ice for 5 min, the extracts were mixed with the same volume of

fresh Ehrlich's reagent [0.2 g *p*-dimethylaminobenzaldehyde (Sigma-Aldrich), 8.4 mL

acetic acid, and 1.6 mL 70% (v/v) perchloric acid (Sigma-Aldrich)] and centrifuged at

14,000 g for 5 min at 4 °C. The OD of each supernatant was measured at 553 nm using

the NanoDrop 2000 (Thermo Fisher Scientific). The amount of 5-ALA was calculated

using a coefficient of 7.45×10^4 mol^{-1} cm^{-1}.

Determining photochemical efficiency

Measurements of photochemical efficiency of PSII (Fv/Fm) were conducted with a

FluorCam system (FC800-C/1010GFP; Photon Systems Instruments) containing a

CCD camera and an irradiation system according to the instrument manufacturer's

instructions.

Trypan blue staining

Cell death was determined by trypan blue (TB) staining as described previously (Lv et

al., 2019). The plant tissues were submerged in TB staining solution [25% (v/v) phenol,

25% (v/v) glycerol, 25% (v/v) lactic acid, 0.05% (w/v) trypan blue] diluted with ethanol

1:2 (v/v) and boiled for 2 min. After incubating for 16 h on a vertical shaker at room

temperature, the non-specific staining was removed using destaining solution (250 g

chloral hydrate dissolved in 100 ml H_2O, pH 1.2). Plant tissues were then kept in 50%

(v/v) glycerol before taking images.

Gene ontology (GO) enrichment analysis

The individual RNA-seq data using the *lsd1* and *glk1 glk2*, as analyzed in Supplemental

Figure 5A, were previously published by Li et al. (Lv et al., 2019) and Ni *et al.* (Ni et

al., 2017), respectively. The GO enrichment analysis of the selected genes shown in

Supplemental Dataset 5 was performed on gprofiler (https://biit.cs.ut.ee/gprofiler) and

represented the significantly enriched GO terms in the data set of biological processes

(BP) with a significance of *P*-value < 0.05.

Pigment analysis

The level of Pchlide was measured in 10-day-old plants of WT, *flu*, *lsd1*, and *lsd1 flu*

as described by Goslings *et al.* (Goslings et al., 2004).

FUNDING

This research was supported by the Strategic Priority Research Program from the

Chinese Academy of Sciences (Grant No. XDB27040102), the 100-Talent Program of

the Chinese Academy of Sciences, and the National Natural Science Foundation of

China (NSFC) (Grant No. 31871397) to C.K..

AUTHOR CONTRIBUTIONS

M.L., K.P.L., T.L., V.D., W.X., and C.K. designed the experiments. M.L., K.P.L., T.L.,

V.D., J.D., and M.S.L. performed the experiments. M.L., K.P.L., T.L., V.D., W.X., and

C.K. analyzed the data. C.K. wrote the manuscript with significant contributions from

M.L and K.P.L. All authors discussed the results and reviewed the manuscript.

ACKNOWLEDGMENTS

We thank the Core Facility of Proteomics in Shanghai Center for Plant Stress Biology

(PSC) for carrying out mass spectrometry. No conflict of interest declared.

REFERENCES

Aviv, D.H., Rusterucci, C., Holt, B.F., 3rd, Dietrich, R.A., Parker, J.E., and Dangl,
J.L. (2002). Runaway cell death, but not basal disease resistance, in lsd1 is SA- and
NIM1/NPR1-dependent. Plant J **29**:381-391.

Beale, S.I., and Castelfranco, P.A. (1974). The Biosynthesis of delta-Aminolevulinic
Acid in Higher Plants: I. Accumulation of delta-Aminolevulinic Acid in Greening Plant
Tissues. Plant Physiol **53**:291-296.

Boruc, J., Van den Daele, H., Hollunder, J., Rombauts, S., Mylle, E., Hilson, P.,
Inze, D., De Veylder, L., and Russinova, E. (2010). Functional modules in the
Arabidopsis core cell cycle binary protein-protein interaction network. Plant Cell
22:1264-1280.

Bravo-Garcia, A., Yasumura, Y., and Langdale, J.A. (2009). Specialization of the
Golden2-like regulatory pathway during land plant evolution. New Phytol **183**:133-141.

Chai, T., Zhou, J., Liu, J., and Xing, D. (2015). LSD1 and HY5 antagonistically
regulate red light induced-programmed cell death in Arabidopsis. Front Plant Sci **6**:292.

Chan, K.X., Phua, S.Y., Crisp, P., McQuinn, R., and Pogson, B.J. (2016). Learning
the Languages of the Chloroplast: Retrograde Signaling and Beyond. Annu Rev Plant
Biol **67**:25-53.

Cheng, J., He, C.X., Zhang, Z.W., Xu, F., Zhang, D.W., Wang, X., Yuan, S., and
Lin, H.H. (2011). Plastid signals confer Arabidopsis tolerance to water stress. Z
Naturforsch C **66**:47-54.

Clough, S.J., and Bent, A.F. (1998). Floral dip: a simplified method for
Agrobacterium-mediated transformation of Arabidopsis thaliana. Plant J **16**:735-743.

Czarnocka, W., Van Der Kelen, K., Willems, P., Szechynska-Hebda, M.,
Shahnejat-Bushehri, S., Balazadeh, S., Rusaczonek, A., Mueller-Roeber, B., Van
Breusegem, F., and Karpinski, S. (2017). The dual role of LESION SIMULATING
DISEASE 1 as a condition-dependent scaffold protein and transcription regulator. Plant
Cell Environ **40**:2644-2662.

Dietrich, R.A., Richberg, M.H., Schmidt, R., Dean, C., and Dangl, J.L. (1997). A
novel zinc finger protein is encoded by the Arabidopsis LSD1 gene and functions as a
negative regulator of plant cell death. Cell **88**:685-694.

Dietrich, R.A., Delaney, T.P., Uknes, S.J., Ward, E.R., Ryals, J.A., and Dangl, J.L.
(1994). Arabidopsis mutants simulating disease resistance response. Cell **77**:565-577.

Dogra, V., Li, M., Singh, S., Li, M., and Kim, C. (2019). Oxidative post-translational
modification of EXECUTER1 is required for singlet oxygen sensing in plastids. Nat
Commun **10**:2834.

Dogra, V., Duan, J., Lee, K.P., Lv, S., Liu, R., and Kim, C. (2017). FtsH2-Dependent
Proteolysis of EXECUTER1 Is Essential in Mediating Singlet Oxygen-Triggered
Retrograde Signaling in Arabidopsis thaliana. Front Plant Sci **8**:1145.

Fitter, D.W., Martin, D.J., Copley, M.J., Scotland, R.W., and Langdale, J.A.
(2002). GLK gene pairs regulate chloroplast development in diverse plant species. Plant
J **31**:713-727.
Forreiter, C., and Apel, K. (1993). Light-independent and light-dependent
protochlorophyllide-reducing activities and two distinct NADPH-protochlorophyllide
oxidoreductase polypeptides in mountain pine (Pinus mugo). Planta **190**:536-545.
Goslings, D., Meskauskiene, R., Kim, C., Lee, K.P., Nater, M., and Apel, K. (2004).
Concurrent interactions of heme and FLU with Glu tRNA reductase (HEMA1), the
target of metabolic feedback inhibition of tetrapyrrole biosynthesis, in dark- and light-
grown Arabidopsis plants. Plant J **40**:957-967.
Han, X.Y., Li, P.X., Zou, L.J., Tan, W.R., Zheng, T., Zhang, D.W., and Lin, H.H.
(2016). GOLDEN2-LIKE transcription factors coordinate the tolerance to Cucumber
mosaic virus in Arabidopsis. Biochem Biophys Res Commun **477**:626-632.
Hou, Z., Yang, Y., Hedtke, B., and Grimm, B. (2019). Fluorescence in blue light
(FLU) is involved in inactivation and localization of glutamyl-tRNA reductase during
light exposure. Plant J **97**:517-529.
Huang, X., Li, Y., Zhang, X., Zuo, J., and Yang, S. (2010). The Arabidopsis LSD1
gene plays an important role in the regulation of low temperature-dependent cell death.
New Phytol **187**:301-312.
Jabs, T., Dietrich, R.A., and Dangl, J.L. (1996). Initiation of runaway cell death in
an Arabidopsis mutant by extracellular superoxide. Science **273**:1853-1856.
Karpinski, S., Szechynska-Hebda, M., Wituszynska, W., and Burdiak, P. (2013).
Light acclimation, retrograde signalling, cell death and immune defences in plants.
Plant Cell Environ **36**:736-744.
Kim, C., Meskauskiene, R., Zhang, S., Lee, K.P., Lakshmanan Ashok, M.,
Blajecka, K., Herrfurth, C., Feussner, I., and Apel, K. (2012). Chloroplasts of
Arabidopsis are the source and a primary target of a plant-specific programmed cell
death signaling pathway. Plant Cell **24**:3026-3039.
Koussevitzky, S., Nott, A., Mockler, T.C., Hong, F., Sachetto-Martins, G., Surpin,
M., Lim, J., Mittler, R., and Chory, J. (2007). Signals from chloroplasts converge to
regulate nuclear gene expression. Science **316**:715-719.
Lai, Z., Li, Y., Wang, F., Cheng, Y., Fan, B., Yu, J.Q., and Chen, Z. (2011).
Arabidopsis sigma factor binding proteins are activators of the WRKY33 transcription
factor in plant defense. Plant Cell **23**:3824-3841.
Lee, J., Choi, B., Yun, A., Son, N., Ahn, G., Cha, J.Y., Kim, W.Y., and Hwang, I.
(2021). Long-term abscisic acid promotes golden2-like1 degradation through
constitutive photomorphogenic 1 in a light intensity-dependent manner to suppress
chloroplast development. Plant Cell Environ **44**:3034-3048.
Lee, J.H., Jin, S., Kim, S.Y., Kim, W., and Ahn, J.H. (2017). A fast, efficient
chromatin immunoprecipitation method for studying protein-DNA binding in
Arabidopsis mesophyll protoplasts. Plant Methods **13**:42.
Lee, K.P., Kim, C., Landgraf, F., and Apel, K. (2007). EXECUTER1- and
EXECUTER2-dependent transfer of stress-related signals from the plastid to the
nucleus of Arabidopsis thaliana. Proc Natl Acad Sci U S A **104**:10270-10275.

Lee, K.P., Liu, K., Kim, E.Y., Medina-Puche, L., Dong, H., Duan, J., Li, M., Dogra,
V., Li, Y., Lv, R., et al. (2020). PLANT NATRIURETIC PEPTIDE A and Its Putative
Receptor PNP-R2 Antagonize Salicylic Acid-Mediated Signaling and Cell Death. Plant
Cell **32**:2237-2250.
Li, Z., Dogra, V., Lee, K.P., Li, R., Li, M., Li, M., and Kim, C. (2020). N-Terminal
Acetylation Stabilizes SIGMA FACTOR BINDING PROTEIN1 Involved in Salicylic
Acid-Primed Cell Death. Plant Physiol **183**:358-370.
Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data
using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods
25:402-408.
Lu, Q., Tang, X., Tian, G., Wang, F., Liu, K., Nguyen, V., Kohalmi, S.E., Keller,
W.A., Tsang, E.W., Harada, J.J., et al. (2010). Arabidopsis homolog of the yeast
TREX-2 mRNA export complex: components and anchoring nucleoporin. Plant J
61:259-270.
Lv, R., Li, Z., Li, M., Dogra, V., Lv, S., Liu, R., Lee, K.P., and Kim, C. (2019).
Uncoupled Expression of Nuclear and Plastid Photosynthesis-Associated Genes
Contributes to Cell Death in a Lesion Mimic Mutant. Plant Cell **31**:210-230.
Martin, G., Leivar, P., Ludevid, D., Tepperman, J.M., Quail, P.H., and Monte, E.
(2016). Phytochrome and retrograde signalling pathways converge to antagonistically
regulate a light-induced transcriptional network. Nat Commun **7**:11431.
Meskauskiene, R., Nater, M., Goslings, D., Kessler, F., op den Camp, R., and Apel,
K. (2001). FLU: a negative regulator of chlorophyll biosynthesis in Arabidopsis
thaliana. Proc Natl Acad Sci U S A **98**:12826-12831.
Miller, G., Suzuki, N., Rizhsky, L., Hegie, A., Koussevitzky, S., and Mittler, R.
(2007). Double mutants deficient in cytosolic and thylakoid ascorbate peroxidase reveal
a complex mode of interaction between reactive oxygen species, plant development,
and response to abiotic stresses. Plant Physiol **144**:1777-1785.
Morikawa, K., Shiina, T., Murakami, S., and Toyoshima, Y. (2002). Novel nuclear-
encoded proteins interacting with a plastid sigma factor, Sig1, in Arabidopsis thaliana.
FEBS Lett **514**:300-304.
Muhlenbock, P., Plaszczyca, M., Plaszczyca, M., Mellerowicz, E., and Karpinski,
S. (2007). Lysigenous aerenchyma formation in Arabidopsis is controlled by LESION
SIMULATING DISEASE1. Plant Cell **19**:3819-3830.
Muhlenbock, P., Szechynska-Hebda, M., Plaszczyca, M., Baudo, M., Mateo, A.,
Mullineaux, P.M., Parker, J.E., Karpinska, B., and Karpinski, S. (2008).
Chloroplast signaling and LESION SIMULATING DISEASE1 regulate crosstalk
between light acclimation and immunity in Arabidopsis. Plant Cell **20**:2339-2356.
Murmu, J., Wilton, M., Allard, G., Pandeya, R., Desveaux, D., Singh, J., and
Subramaniam, R. (2014). Arabidopsis GOLDEN2-LIKE (GLK) transcription factors
activate jasmonic acid (JA)-dependent disease susceptibility to the biotrophic pathogen
Hyaloperonospora arabidopsidis, as well as JA-independent plant immunity against the
necrotrophic pathogen Botrytis cinerea. Mol Plant Pathol **15**:174-184.

Nandi, D.L., and Shemin, D. (1968). Delta-aminolevulinic acid dehydratase of
Rhodopseudomonas spheroides. 3. Mechanism of porphobilinogen synthesis. J Biol
Chem **243**:1236-1242.

Ni, F., Wu, L., Wang, Q., Hong, J., Qi, Y., and Zhou, X. (2017). Turnip Yellow
Mosaic Virus P69 Interacts with and Suppresses GLK Transcription Factors to Cause
Pale-Green Symptoms in Arabidopsis. Mol Plant **10**:764-766.

Nott, A., Jung, H.S., Koussevitzky, S., and Chory, J. (2006). Plastid-to-nucleus
retrograde signaling. Annu Rev Plant Biol **57**:739-759.

Rossini, L., Cribb, L., Martin, D.J., and Langdale, J.A. (2001). The maize golden2
gene defines a novel class of transcriptional regulators in plants. Plant Cell **13**:1231-
1244.

Rusaczonek, A., Czarnocka, W., Kacprzak, S., Witon, D., Slesak, I., Szechynska-
Hebda, M., Gawronski, P., and Karpinski, S. (2015). Role of phytochromes A and
B in the regulation of cell death and acclimatory responses to UV stress in Arabidopsis
thaliana. J Exp Bot **66**:6679-6695.

Saraste, M., and Musacchio, A. (1994). Backwards and forwards binding. Nat Struct
Biol **1**:835-837.

Savitch, L.V., Subramaniam, R., Allard, G.C., and Singh, J. (2007). The GLK1
'regulon' encodes disease defense related proteins and confers resistance to Fusarium
graminearum in Arabidopsis. Biochem Biophys Res Commun **359**:234-238.

Siligardi, G., and Drake, A.F. (1995). The importance of extended conformations and,
in particular, the PII conformation for the molecular recognition of peptides.
Biopolymers **37**:281-292.

Tang, H., Zhang, D., Yuan, S., Zhu, F., F., X., Fu, F.Q., Wang, S., and Lin, H.H.
(2014). Plastid signals induce ALTERNATIVE OXIDASE expression to enhance the
cold stress tolerance in Arabidopsis thaliana. Plant Growth Regul **74**:275-283.

Tang, X., Miao, M., Niu, X., Zhang, D., Cao, X., Jin, X., Zhu, Y., Fan, Y., Wang,
H., Liu, Y., et al. (2016). Ubiquitin-conjugated degradation of golden 2-like
transcription factor is mediated by CUL4-DDB1-based E3 ligase complex in tomato.
New Phytol **209**:1028-1039.

Tokumaru, M., Adachi, F., Toda, M., Ito-Inaba, Y., Yazu, F., Hirosawa, Y.,
Sakakibara, Y., Suiko, M., Kakizaki, T., and Inaba, T. (2017). Ubiquitin-
Proteasome Dependent Regulation of the GOLDEN2-LIKE 1 Transcription Factor in
Response to Plastid Signals. Plant Physiol **173**:524-535.

Tzfira, T., Tian, G.W., Lacroix, B., Vyas, S., Li, J., Leitner-Dagan, Y., Krichevsky,
A., Taylor, T., Vainstein, A., and Citovsky, V. (2005). pSAT vectors: a modular
series of plasmids for autofluorescent protein tagging and expression of multiple genes
in plants. Plant Mol Biol **57**:503-516.

Waters, M.T., Moylan, E.C., and Langdale, J.A. (2008). GLK transcription factors
regulate chloroplast development in a cell-autonomous manner. Plant J **56**:432-444.

Waters, M.T., Wang, P., Korkaric, M., Capper, R.G., Saunders, N.J., and
Langdale, J.A. (2009). GLK transcription factors coordinate expression of the
photosynthetic apparatus in Arabidopsis. Plant Cell **21**:1109-1128.

Xie, Y.D., Li, W., Guo, D., Dong, J., Zhang, Q., Fu, Y., Ren, D., Peng, M., and Xia,
Y. (2010). The Arabidopsis gene SIGMA FACTOR-BINDING PROTEIN 1 plays a
role in the salicylate- and jasmonate-mediated defence responses. Plant Cell Environ
33:828-839.
Yoo, S.D., Cho, Y.H., and Sheen, J. (2007). Arabidopsis mesophyll protoplasts: a
versatile cell system for transient gene expression analysis. Nat Protoc **2**:1565-1572.
Zarrinpar, A., Bhattacharyya, R.P., and Lim, W.A. (2003). The structure and
function of proline recognition domains. Sci STKE **2003**:RE8.
Zhang, D., Tan, W., Yang, F., Han, Q., Deng, X., Guo, H., Liu, B., Yin, Y., and Lin,
H. (2021a). A BIN2-GLK1 Signaling Module Integrates Brassinosteroid and Light
Signaling to Repress Chloroplast Development in the Dark. Dev Cell **56**:310-324 e317.
Zhang, H., Zhang, L., Ji, Y., Jing, Y., Li, L., Chen, Y., Wang, R., Zhang, H., Yu,
D., and Chen, L. (2021b). Arabidopsis SIGMA FACTOR BINDING PROTEINs
function antagonistically to WRKY75 in abscisic acid-mediated leaf senescence and
seed germination. J Exp Bot:erab391.
Zhang, Z.W., Yuan, S., Xu, F., Yang, H., Chen, Y.E., Yuan, M., Xu, M.Y., Xue,
L.W., Xu, X.C., and Lin, H.H. (2011). Mg-protoporphyrin, haem and sugar signals
double cellular total RNA against herbicide and high-light-derived oxidative stress.
Plant Cell Environ **34**:1031-1042.
Zhang, Z.W., Feng, L.Y., Cheng, J., Tang, H., Xu, F., Zhu, F., Zhao, Z.Y., Yuan,
M., Chen, Y.E., Wang, J.H., et al. (2013). The roles of two transcription factors, ABI4
and CBFA, in ABA and plastid signalling and stress responses. Plant Mol Biol **83**:445-
458.

 SUPPLEMENTAL INFORMATION

SUPPLEMENTAL FIGURES

Supplemental Figure 1. Detection of the LSD1-GFP fusion protein.
Total proteins were extracted from two-week-old plants of WT and *35S:LSD1-GFP* transgenic line
(*oxLSD1*) grown under CL. The proteins were subjected to an immunoblot assay to detect the LSD1-
GFP fusion protein using an anti-GFP antibody. Denaturing gel stained with Coomassie brilliant blue
(CBB) was used as a loading control.

Supplemental Figure 2. Sequence alignment of Arabidopsis GLK1 and GLK2 proteins.
The conserved amino acid sequences were aligned using the Clustal Omega software
(https://www.ebi.ac.uk/Tools/msa/clustalo/) before shading with the Jalview program. The boxes with
different colors indicate identical and similar amino acids (black: 100%; dark gray: ≥75%; light gray:
≥50%). Red, blue, and green lines indicate region of DNA-binding domain (DBD), proline-rich domain
(PRD), and GLK/C-terminal box (GCT-box), respectively.

Supplemental Figure 3. Domain-deleted GLK1 and GLK2 variants localize to the nucleus.
Subcellular localization of GFP-tagged intact GLK1/2 and their domain-deleted variants upon transient
expression in *N. benthamiana* leaves. DAPI was used to stain the nucleus. All images were taken at the
same scale (scale bars: 10 μm).

Supplemental Figure 4. The C-terminal PRD region of GLKs is critical for LSD1-GLKs interaction.
(A) Schematic diagram of GLK1/2 and their C-terminally truncated variant proteins.
(B) Subcellular localization of GFP-tagged protein variants **(A)** upon transient expression in *N.*
benthamiana leaves. **(C)** BiFC analysis. The intact or truncated variants of GLK1 or GLK2 fused with
YFP[C] were individually coexpressed with LSD1 fused with YFP[N] in *N. benthamiana* leaves. In **(B)** and
(C), DAPI was used to stain the nucleus. All images were taken at the same scale (scale bars: 10 μm).
(D) Co-IP analyses using *N. benthamiana* leaves coexpressing LSD1-GFP with indicated intact or
truncated variants of GLK1 (or GLK2) fused with Myc-tag. GFP-Trap beads were used, and the
interaction was evaluated by using Myc antibody.

Supplemental Figure 5. Loss of LSD1 leads to an upregulation of GLK target genes.
(A) Venn diagram showing the numbers of uncommon and overlapped genes between upregulated genes
(395) in 17-d-old *lsd1* (Lv et al., 2019) and downregulated genes (936) in *glk1 glk2* (Ni et al., 2017).
(B) Gene Ontology (GO) enrichment analysis towards the biological process of the overlapped genes in
(A). The GO enrichment analysis was done as described in Methods.

Supplemental Figure 6. Expression levels of *GLK1* and *GLK2* in *oxLSD1* lines.
(A and B) Total protein and RNA were extracted from 24-d-old CL-grown plants of WT, *glk1 glk2,* and
two independent transgenic lines overexpressing GFP-tagged LSD1 under the control of the CaMV 35S
promoter (*oxLSD1* #1 and #5). The proteins were subjected to an immunoblot assay to detect the LSD1-
GFP fusion proteins using an anti-GFP antibody **(A)**. UGPase was used as a loading control. The relative
expression levels of *GLK1* and *GLK2* were analyzed by RT-qPCR **(B)**. *ACT2* was used as an internal
standard. Data are means ± SD (n=3). Lowercase letters indicate statistically significant differences
between mean values (P < 0.01, one-way ANOVA with posthoc Tukey's HSD test).

Supplemental Figure 7. LSD1 does not interact with SIB1 *in vivo*.
Co-IP analyses using *Arabidopsis* leaf protoplasts transiently coexpressing LSD1−GFP and SIB1−RFP.
LSD1−GFP was also coexpressed with free RFP as a negative control or LSD1−RFP as a positive control.
The immunoblot result is one representative of three independent experiments with similar results.

Supplemental Figure 8. SIB1 interacts with GLK1 through the N-terminal region of GLK1.
(A and B) Co-IP analyses using *N. benthamiana* leaves transiently coexpressing SIB1-GFP with intact
form and C-terminally truncated **(A)**, or domain-deleted variants **(B)** of GLK1 fused with Myc-tag.
LSD1-GFP was also transiently coexpressed with GLK1$_{\Delta CT}$ -Myc as a negative control showing a lack
of interaction. Co-IP was performed with GFP-Trap beads, and the interaction was evaluated by using
the Myc antibody.

 SUPPLEMENTAL TABLE

 Supplemental Table 1. List of primer sets used in this study.

Gene ID	Gene name	Mutant alleles	Primer sequence (5' to 3')	Primer length	Size (bp)	Used for
At3g18780	*ACT2*	-	F: TATGTATGTCGCCATCCAA	19	76	
			R: ACCAGAATCCAGCACAATA	19		
At2g34430	*LHCB1.4*	-	F: AGCAGAGGACTTGCTTTACC	20	115	
			R: CATAGCCAACCTTCCGTTCT	20		
At2g05100	*LHCB2.1*	-	F: CAAAGCATCTGGTACGGACCAG	22	336	
			R: GATGCTTTGCGCGTGGATCAAG	22		
At2g05070	*LHCB2.2*	-	F: CGTCAAGTCTACTCCCCAAAG	21	71	
			R: TCTCCGAGAATGGTCCCAAG	20		
At3g27690	*LHCB2.3*	-	F: CGGAGAATACCCTGGAGACTA	21	104	
			R: CCCATCTACTGTGGATCACTTC	22		
At1g61520	*LHCB3*	-	F: CACGAGCTCAAGCAGTGTTC	20	92	
			R: CGAGAGAGACAACATCACGA	20		
At1g15820	*LHCB6*	-	F: CGGATTCTCAATCGGTTGAGT	21	112	
			R: CCCAACGGATCGAAGAATCTC	21		
At3g54890	*LHCA1*	-	F: CCCTTCGCTTCTCTCTTCTTC	21	98	
			R: AGTGAGCAGCCATTCTGATAC	21		
At2g20570	*GLK1*	-	F: GCTACGAGATTTAGAGCACCG	21	123	RT-qPCR
			R: TTGACGGATGTAAGTCTACC	20		
At5g44190	*GLK2*	-	F: GGCATCAGCAACCACTCTAT	20	113	
			R: ATGAATGTCGATGGGAGGATTAG	23		
At4g20380	*LSD1*	-	F: AAGGGTACCTCTCCCAACTAA	21	136	
			R: CACCAACTTTCCGCTTTCATC	21		
At1g44446	*CAO*	-	F: TAGGGGTGAAGACGGGAAACC	21	134	
			R: CTCCATCGGTTGAGTATTCCC	21		
At1g58290	*HEMA1*	-	F: TAATGGGGTTCGTGTTCTTCCG	22	111	
			R: ATGCTAGCTGCATTAGACGCAG	22		
At3g59400	*GUN4*	-	F: GCCATCCTGCGTTTGCGACAG	21	104	
			R: GTCTGCTCCTACTCCTGCCTG	21		
At5g13630	*GUN5*	-	F: CTACAGGGCGAACAGAGATAAG	22	102	
			R: GCTTGCATTAGACTCCCTAGTT	22		
At5g54190	*PORA*	-	F: CCCTCTTCCCTCCTTTCCAG	20	122	
			R: GCTCCAATACACTCCCGACTTC	22		
At4g27440	*PORB*	-	F: CAAACCGCTGCGACTTCAAGC	21	163	
			R: TGCACGCCATTATCACGTTCC	21		

Gene ID	Gene name	Mutant alleles	Primer sequence (5' to 3')	Primer length	Size (bp)	Used for
At1g03630	*PORC*	-	F: CAGACAGTTACAGCCACGCCG	21	133	RT-qPCR
			R: TGTCTGCTAAAGCTTTGGCCG	21		
At4g62830	*LSD1*	Salk_042687 (*lsd1-2*)	LP: CTGGGATTTGTAAAGCAGCTG	21	-	
			RP: TCAAGTTCCATGGAGCAAAAG	21		
At2g20570	*GLK1*	*Atglk1.1*	WT-F: CAATAGGCGGGCCTTATCTAG	23	-	
			WT-R: GATAAGATCTCAGGGTCGATCTCC	24		
			glk1.1-F: ACTGCAGGTTACTGATCCGATTGTTCTT	29		
			glk1.1-R: CGGGATCCGACACTCTTTAATTAACTGACACTC	33		Geno-typing
At5g44190	*GLK2*	*Atglk2.1*	WT-F: CGACGGAAGACTTGCCGGACTT	22	-	
			WT-R: GTGTAACTCCGGCGTCCAATCC	22		
			glk2.1-F-1: CCTATTTCAGTAAGAGTGTGGGGTTTTGG	29		
			glk2.1-R-1: GTGTAACTCCGGCGTCCAATCC	22		
At4g33630	*EX1*	Salk_002088 (*ex1*)	LP: TACCCCAATCACTCAAATTG	21	-	
			RP: CACTCCCTCCTCCAAAAGATC	21		
At2g34430	*LHCB1.4*	-	F: CTGTATCTGTTTAGTGATTGGC	22	256	
			R: TGAGAGCATGAAGTGGATTGG	21		
At1g61520	*LHCB3*	-	F: CCCACCTCTCTTCTCATCCA	20	141	ChIP-qPCR
			R: TGATGCCATTGTCTCTCTCG	20		
At1g15820	*LHCB6*	-	F: GCAATAAGCCACATAATGCAG	21	225	
			R: CTGACCAATTAGGAGTCAGAAACTAC	26		