Peptide Ancestry Informative Markers in Uterine Neoplasms from Women of European, African and Asian Ancestry

Nicholas W. Bateman, PhD*[1-3], Christopher Tarney, MD[1], Tamara Abulez, BS[1,3], Brian L. Hood, PhD[1,3], Kelly Conrads, PhD[1,3], Ming Zhou, PhD[4], Anthony R. Soltis, PhD[3,5], Pang-ing Teng, PhD[1,3], Amanda Jackson, MD[1], Chunqiao Tian, PhD[1,3], Clifton L. Dalgard, PhD[5,6], Matthew D. Wilkerson, PhD[2,3,5,6], Michael Kessler, PhD[7], Craig D, Shriver, MD[2], Hai Hu, PhD[8], Michele Cote, PhD[9], Glendon J. Parker, PhD[10], James Segars, MD[11], Ayman Al-Hendy, MD, PhD[12], John R. Risinger, PhD[13], Kathleen Darcy, PhD[1-3], Yovanni Casablanca, MD[1,2], G. Larry Maxwell, MD[1,2,4], Thomas P. Conrads, PhD[1,2,4], Timothy D. O'Connor, PhD*[6,14,15]

[1]Gynecologic Cancer Center of Excellence, Department of Obstetrics and Gynecology, Uniformed Services University and Walter Reed National Military Medical Center, 8901 Wisconsin Avenue, Bethesda, MD 20889, USA.

[2]The John P. Murtha Cancer Center, Uniformed Services University and Walter Reed National Military Medical Center, 8901 Wisconsin Avenue, Bethesda, MD 20889, USA.

[3]Henry M. Jackson Foundation for the Advancement of Military Medicine, Inc., 6720A Rockledge Dr., Suite 100, Bethesda, MD 20817, USA.

[4]Department of Obstetrics and Gynecology, Inova Fairfax Medical Campus, 3300 Gallows Rd. Falls Church, VA 22042, USA.

[5]The American Genome Center; Uniformed Services University of the Health Sciences, Bethesda, MD 20814, USA

[6]Department of Anatomy Physiology and Genetics, Uniformed Services University, 4301 Jones Bridge Road, Bethesda, MD, 20814 USA.

[7]Institute for Genome Sciences, University of Maryland School of Medicine, Baltimore, MD 21201 USA.

[8]Chan Soon-Shiong Institute of Molecular Medicine at Windber. Windber, PA 15963 USA.

[9]Wayne State University, Detroit, MI 48202 USA.

[10]University of California, Davis, One Shields Avenue, Davis, CA 95616 USA.

[11]Johns Hopkins University Medical Center, Baltimore, MD 21218 USA.

[12]The University of Illinois College of Medicine, Chicago, IL 60612 USA.

[13]Department of Obstetrics and Gynecology, Michigan State University, East, Lansing, MI 48824, USA.

[14]Program in Personalize and Genomic Medicine, University of Maryland School of Medicine, Baltimore, MD 21201 USA.

[15]Marlene and Stewart Greenebaum Comprehensive Cancer, University of Maryland School of Medicine, Baltimore, MD 21201 USA.

*Authors contributed equally to this work.

Keywords: Proteogenomics, Uterine Neoplasms, Ancestry Informative Markers.

Funding Statement: This study was supported in part by the U.S. Department of Defense -
Uniformed Services University of the Health Sciences (HU0001-16-2-0006 and HU0001-16-2-
0014)

Address correspondence to: Nicholas W. Bateman, PhD. 3289 Woodburn Rd, Suite 375,
Annandale, VA, 22003; Tel: 703-207-3362; Email: batemann@whirc.org

Abstract

Deep proteogenomic analyses of diverse populations will improve our understanding of molecular

drivers of disease, such as those underlying uterine neoplasms which exhibit marked racial

disparities. The characterization of ancestry-linked peptide variants in disease-relevant patient

tissues represents a foundational step towards investigating relationships linking patient ancestry

with the proteome, genome and disease pathogenesis. We selected 1,037 nonsynonymous single

nucleotide polymorphisms (nsSNPs) encoding missense amino acid substitutions occurring within

putative tryptic peptides exhibiting high allele frequencies ($\geq 50\%$) in European, African, and East

Asian populations, termed peptide ancestry informative markers (pAIMs) and investigated these

in cell line models and tissues of uterine neoplasms collected from women representing diverse

ancestries. We quantified 62 pAIMs in total across >100 patient-derived samples by multiplexed,

quantitative proteomic analyses. Our analyses of endometrial cancer cell lines established from

European (n=7), African (n=3), and East Asian (n=3) women resulted in the quantitation of 43

pAIMs that significantly correlated with population-level allele frequencies consistent with global

ancestry estimates defined by genotype analyses. We next investigated uterine leiomyoma (ULM)

tissues from self-described African-American (n=31) and European-American (n=35) women and

quantified 33 pAIMs that correlated with patient race. We validated 10 of these in an independent

cohort of ULM tissues collected from women determined to be of European (n=16) and African

(n=11) ancestry by genotype analyses. This validated subset was predominated by candidates of

undetermined clinical significance (ClinVar) and included a substitution in GC vitamin D-binding

protein, i.e. (D451E, rs7041), that exhibits a higher allele frequency in European (58%) versus

African (7.3%) populations and correlates with altered GC protein and vitamin D levels.

Comparison of cell line and tissue analyses revealed co-quantitation of pAIMs encoded within

Protein-L-isoaspartate(D-aspartate) O-methyltransferase (V178I, >80% AF in African and East

Asians), Serine/threonine-protein phosphatase CPPED1 (A19D, >75% AF in Africans) and AH

receptor-interacting protein (Q228K, >99% AF in European and East Asians). We further assessed

the performance of pAIMs to determine patient ancestry *in silico* relative to standard genotype

estimates by random sampling analyses and find that as few as 20 population-specific pAIMs can

determine ancestry proportions similarly to >260K SNPs (R^2=0.9905). These efforts have

quantified and validated pAIMs in diverse uterine neoplasms that represent a foundational set of

ancestry-linked variant peptides observable by multiplexed proteomics and provides proof of

concept that estimates of European, African and East Asian ancestry can be determined from

routine proteomic analysis of disease-relevant patient cell lines and tissues. This work will support

ongoing efforts focused on linking the proteome and genome with patient ancestry to better

understand relationships with disease mechanisms driving racial disparities in the pathogenesis of

uterine neoplasms.

Introduction

Proteogenomics aims to integrate protein-level measurements with companion transcriptome and genome sequencing data (1) to elucidate complex systems-level relationships between protein and transcript-level expression including identification of proteins encoding missense substitutions that may impact protein function (1,2). Understanding the impact of proteogenomic alterations on disease risk and patient outcome is a priority for population-level investigations (3), particularly those focusing on cancer, such as efforts by The Cancer Genome Atlas (TCGA) (4) and the recently initiated Applied Proteogenomics OrganizationaL Learning and Outcomes (APOLLO) research network (5). Many historic efforts have poor representation of U.S. minorities amongst patient cohorts analyzed (6) and there remains a paucity of systems-level molecular analyses within powered cohorts to elucidate race-related differences in disease etiology. In the context of gynecologic disease, patients with uterine leiomyomas (fibroids) and endometrial cancers experience significant racial disparities in disease incidence and outcome (7-13). Among possible molecular mechanisms that may give rise to racial disparities in gynecologic disease, one such manifestation could be from non-synonymous single nucleotide polymorphisms (nsSNPs) encoding missense substitutions that may alter protein function(s). Defining ancestry-linked peptide variants in disease-relevant cells and tissues will augment proteogenomic discovery efforts that aim to define molecular drivers underlying racial disparities in gynecologic diseases.

Ancestry informative markers (AIMs) comprise SNPs that are highly differentiated within discrete ancestral populations (14-16). Most of these markers occur within non-protein coding regions, but with large-scale sequencing now available for a diversity of ancestries (17,18), especially in protein coding regions (19,20), it is now possible to assess AIMs that are detectable through proteomics data alone. To this end, efforts to date have identified genetically variant peptides

differentiating individuals of European and African descent (21) as well as Asian populations (22)

and these have been explored for forensic human identification, such as through proteomic

analyses of human hair and bone samples. We leveraged population-level genotyping data from

the 1000 Genomes Project (23) to identify SNPs encoding missense substitutions that exhibit

altered allele frequencies with European, African, and East Asian ancestries and prioritized

candidates occurring within tryptic peptides observable by routine, bottom-up proteomic

workflows to explore these within disease-relevant models and tissues from uterine neoplasms.

Although these candidates are not tissue dependent and represent a generalized set of markers for

proteoancestry assessment in diverse tissues and cell lines, we show here the ability to ancestrally

characterize endometrial cancer cell line models as well as uterine fibroid tissues from women

representing diverse racial and ancestral backgrounds using multiplexed proteomic approaches.

Our study further defines a foundational set of ancestry-linked variant peptides within uterine

tissues that will support ongoing efforts to investigate germline as well as somatic proteogenomic

alterations underlying ancestry-linked disease biology and how this may further relate to racial

disparities in the pathogenesis of uterine neoplasms.

Results

Selection and *In Silico* Analyses of Peptide Ancestry Informative Markers (pAIMs).

We selected >640K SNPs from the 1000 Genomes Project (23) encoding nonsynonymous

substitutions and further filtered them to those exhibiting ≥50% allele frequency differences

between individuals of European (i.e. FIN, GBR, IBS, and TSI; N=404), African (i.e. ESN, GWD,

LWK, MSL, and YRI; N=504), and East Asian (i.e. CDX, CHB, CHS, and KHV; N=400) descent.

Candidates were also filtered to prioritize missense substitutions occurring in tryptic peptides (≥6

and ≤40 amino acids in length) that would be observable within "bottom-up", multiplexed analyses

of disease-relevant cell lines and tissues collected from individuals representing diverse ancestries

(Workflow Figure 1A, Supplemental Table 1A). This filtering approach resulted in 1,037 peptide

ancestry informative markers (pAIM) mapping to 831 unique proteins spanning all 22 autosomes

and exhibit the greatest frequencies on Chr1 (12.3%) and Chr19 (7.5%) (Supplemental Table 1A).

Most of these parent proteins encode a single pAIM (Supplemental Table 1A) with a subset of 136

candidates encoding ≥2 pAIMs as well as 3 candidates encoding >5 pAIMs; filaggrin (10 pAIMs),

cardiomyopathy-associated protein 5 (7 pAIMs) and HLA class II histocompatibility antigen, DP

alpha 1 chain precursor (6 pAIMs). Functional enrichment analyses of proteins encoding pAIMs

revealed significant enrichment of cellular pathways regulating the cellular matrisome as well as

keratinocyte differentiation and sensory perception (Supplemental Table 1B). Unique pAIMs

exhibiting ≥50% allele frequencies were most prevalent in African populations (344 total sites)

followed by East Asian (229) and European (79) ancestries and East Asian as well as European

populations exhibited nearly 2-fold greater conservation (208) of shared pAIMs relative to African

populations (92 and 85 respectively) (Figure 1B). The clinical significance and known disease

pathogenicity of SNPs encoding pAIMs within the ClinVar resource was assessed for 1,031 subset

pAIMs mapping to the Ensemble Variant Effect Predictor tool (24). Most of these pAIMs had no

previous associations with clinical significance. We did however identify 87 pAIMs of unknown

clinical significance that map to putative functional protein domains (Figure 2A, Supplemental

Table 1A). The remaining candidates were largely undocumented for disease association in

ClinVar with seven pAIMs candidates correlating with altered drug responses or as being risk

factors for disease. This latter subset included a substitution in kinesin family member 6 that is

associated with altered response to statin treatment (25) (W719R, rs20455) and has higher allele

frequencies within African (86%) and East Asian (53%) populations, or as being risk factors for

several diseases (Supplemental Table 1A). Risk factors included a substitution in aurora kinase A

(F31I, rs2273535) which has higher allele frequencies within East Asians (67%) and is associated

with an increased risk of developing colon cancer within these populations (26). We also assessed

the impact of pAIM substitutions on protein function via *in silico* prediction analyses using SIFT

(27), Poly-Phen-2 (28), and PROVEAN (29) and identified 23 pAIMs that may be deleterious

(Figure 2B). Although many of these candidates have unknown clinical significance, five

candidates have been shown to not be pathogenic for disease, i.e. categorized by ClinVar as benign

or likely benign (Figure 2C, Supplemental Table 1A). These candidates include a substitution in

ATP-binding cassette transporter sub-family C member 11 (G180R, rs17822931) that has high

allele frequencies within East Asian populations (75%) and is significantly correlated with

Axillary Osmidrosis in Chinese Han populations (30) as well as increased risk for breast cancer in

Japanese women (31). Further analyses of these 23 putatively deleterious pAIMs revealed they

exhibit higher allele frequencies in East Asian versus European and African populations (Figure

2D). We also utilized as a test set of populations CEU, a European population from Utah; ASW,

an admixed African-American population from the South West; ACB, an admixed African

Caribbean population; and JPT, a Japanese population from East Asia to assess the accuracy of

pAIMs to classify ancestral proportions relative to standard genotype estimates *in silico*. We apply

a random sampling analyses of nsSNPs encoding pAIMs and compared them to using 266,403

LD-pruned SNPs by standard estimates approaches. We find that as few as 20 variants can

recapitulate genome-wide ancestry proportions, using standard approaches (32-34) (error is

increased due to smaller feature set, but average of 500 random sampling assessments remains

robust is $R^2 > 0.99$) (Figure 3).

Quantitation of Peptide Ancestry Informative Markers (pAIMS) in Endometrial Cancer Cell Lines.

We determined the ancestry of thirteen endometrial cancer cell line models by standard estimates using comparison of whole genome sequencing (WGS) data to reference populations (Table 1). Most EC models were established from women of European descent (n=7), with a subset corresponding to women of African (n=3) and East Asian (n=3) descent. We performed quantitative bottom-up proteomic analyses employing a multiplexed tandem-mass tag (TMT) approach and quantified 133,473 total peptides that included 43 high-confidence pAIM variant peptides (Supplemental Table 2), where MS2 ions flanking the pAIM substitution of interest were confirmed using SpectrumAI (35). Among these pAIMS, 39 were quantified in >50% of all EC cell lines and this feature subset was assessed to investigate pAIM abundance relative to cell line ancestries and genotype for the nsSNP of interest. Of these pAIM candidates, 30 are of unknown clinical significance, 12 are undocumented for disease association in ClinVar and one candidate (rs8010699) is of uncertain significance for disease. This latter variant encodes a substitution in nesprin-2 (H3309R) that is associated with Emery-Dreifuss muscular dystrophy and has further been correlated with expression of the tumor suppressor cyclin-dependent kinase inhibitor 1 (CDK1/ p21) in HBV-related hepatocellular carcinomas (36). A subset of 17 pAIMs exhibited >50% allele frequency within European ancestry populations and we found that these candidates were significantly elevated in cell lines from individuals of European versus African ancestry (Mann Whitney U (MWU) Median = +1.003 median fold difference, p=0.0333) as well as East Asian ancestry (East Asian ancestry Median = +0.57 median fold difference, p=0.0167) (Figure 4A). We further find that these European ancestry-correlated pAIMs were most abundant in European ancestry cell lines that were homozygous and heterozygous for the pAIM variant

allele (+/+ and -/+ genotype) relative to those homozygous for the reference allele (-/-) (MWU

p=0.0006). Another subset of 21 pAIMs quantified exhibited >50% allele frequency within

African ancestry populations and were significantly elevated in cell lines from individuals of

African versus European (MWU p=0.0333) ancestry and trended as elevated in African versus

East Asian ancestry cell lines (+1.0676 median fold difference) (Figure 4B). These African

ancestry-associated pAIMs were most abundant in African cell lines homozygous and

heterozygous for the pAIM variant allele (+/+ and -/+ genotype) relative to those homozygous for

the reference allele (-/-), i.e. +/+ vs. -/- cell lines = +1.69 median fold difference. We also quantified

17 pAIMs exhibiting >50% allele frequency within East Asian populations and found that these

are elevated in cell lines from individuals of East Asian versus European ancestry (+0.4058 median

fold difference) and African (+0.84563 median fold difference) ancestry (Figure 4C). These East

Asian ancestry-related pAIMs further tracked with genotype, i.e. (+/+ pAIM allele versus -/- cell

lines = +1.99 median fold difference).

Quantitation of Peptide Ancestry Informative Markers (pAIMS) in Uterine Leiomyoma

Tissues.

We assessed pAIMs within multiplexed (TMT) quantitative proteomic analysis of formalin-fixed,

paraffin embedded uterine leiomyoma tissues from self-described European-American (n=35) or

African-American (n=31) women (Table 2, Discovery). We quantified 69,056 total peptides that

included 33 high-confidence pAIMs (Supplemental Table 3). Of these candidates, 24 are of

unknown clinical significance while 9 have been reported as not being pathogenic for disease

(Supplemental Table 1A, ClinVar). Several of these candidates map to putative functional protein

coding domains including intermediate filament domains within keratin 3 (R375G, >50% AF in

Europeans) and keratin 13 (A187V, >90% in Europeans and East Asians) as well as within the

dihydroxyacetone- (Dha-)binding domain of DHA kinase (A185T, >90% in Europeans and East

Asians) (Supplemental Table 1A). We further assessed the performance of a subset of 18 pAIMs

quantified within >50% of all uterine leiomyoma tissues to distinguish patient ancestry stratified

by self-described race. Unsupervised hierarchical cluster analyses revealed that the abundance of

pAIMs exhibiting high allele frequencies within European or African populations trended with

self-described European or African-American race (Figure 5A). We further assessed performance

of these candidates using a partial least squares (37) approach and identified, based on principle

component analysis, these candidates served to explain 21% and 7% of the variance between

African-American and European-American patient tissues (Figure 5B) and can further classify

African-American from European-American patients with high accuracy (AUC=0.9834, P=1.6E[-]

[11], Figure 5C). We also quantified pAIM abundance by multiplexed (TMT), quantitative proteomic

analyses within an independent cohort of fresh-frozen uterine leiomyoma tissues from women of

European (n=16) or African (n=10) ancestry confirmed using standard genotype estimates from

companion WGS analyses (Table 2, Validation, Supplemental Table 4). We quantified a total of

61,283 total peptides among which 15 pAIMs were quantified in this cohort and 10 of which were

co-quantified with discovery analyses (Supplemental Table 4). This validated subset was

predominated by candidates of unknown clinical significance (ClinVar) and included a substitution

in GC vitamin D-binding protein, i.e. (D451E, >50% Europeans). Assessment of pAIMs with

allele frequencies >50% in African populations were significantly more abundant in African

ancestry patients than pAIMS with allele frequencies of >50% in European ancestry populations

(+1.0381 median fold difference, MWU p=0.003) (Figure 6A). Conversely, we find that pAIMs

with allele frequencies >50% in European ancestry populations were significantly more abundant

in European ancestry patients than pAIMS with allele frequencies >50% in African populations

(+1.74 median fold difference, MWU p<0.0001 (Figure 6B). Further, as observed in the

endometrial cancer cell lines, we find that pAIMs abundances directly correlate within patients

homozygous and heterozygous for the pAIM variant allele (+/+ and -/+ genotype) relative to those

homozygous for the reference allele (-/-), i.e. African and European ancestry +/+ versus -/- patients

(p<0.0001).

Discussion

Comprehensive analysis of proteogenomic alterations correlating with patient ancestry are

necessary to better understand the relationship between racial disparities underlying disease

pathogenesis. In gynecologic malignancies, uterine neoplasms exhibit significant racial disparities,

specifically increased incidence of uterine leiomyomas (38) as well as more aggressive

endometrial cancers in African-American versus European-American patients (8,39,40). We and

others have described race-specific molecular alterations correlating with disease outcome in

endometrial cancers (41-43). As a foundational step to support our ongoing investigations of

molecular alterations underlying racial disparities in gynecologic malignancies, we have

investigated 1,037 nsSNPs exhibiting high allele frequencies within individuals of European,

African, and East Asian populations that encode missense substitutions occurring within putative

tryptic peptides, i.e. termed peptide ancestry informative markers (pAIMS) in cell line models and

tissues derived from uterine neoplasms collected from women representing diverse ancestral

backgrounds. Candidates prioritized were investigated *in silico* for their associations with disease

as well as prevalence within putative functional protein domains and for predicted impact on

protein function. We found that many pAIMs encode substitutions of unknown clinical

significance (ClinVar), occur within functional protein domains and include subsets that are

predicted to be deleterious to protein function. We further find that several pAIM variants are

linked to an increased risk to develop cancer, such as the F31I substitution in aurora kinase A

associated with colon cancer development in East Asian populations (16) and candidates that

further track with cancer cell signaling, such as a substitution in nesprin-2 (SYNE2, H3309,

rs8010699) that correlates with expression of cyclin-dependent kinase inhibitor 1 (CDK1/ p21) in

HBV-related hepatocellular carcinomas (36). We also find that several candidates of unknown

clinical significance occur within known functional protein domains including multiple

immunoglobulin-like domains, such as in the major histocompatibility complex, class II, DP alpha

1 (HLA-DPA1) protein. These candidates further represent ancestrally-linked, variant peptides

theoretically observable with routine LC-MS proteomic workflows encoding substitutions that

may alter functional protein domain interactions as well as protein complex architecture.

Our efforts have quantified and validated pAIMs in uterine neoplasms from >100 women

representing diverse ancestries and consist of a foundational set of ancestry-linked variant peptides

observable by multiplexed proteomic analyses of uterine cells and tissues. Many pAIMs quantified

are of unknown clinical significance in ClinVar and may warrant further investigation to better

understand possible functional roles in the pathogenesis of uterine disease. Validated candidates

were largely undocumented for disease pathogenesis in ClinVar but were associated with altered

biological functions. One such candidate encodes a substitution in GC vitamin D-binding protein,

i.e. (D451E, rs7041), a protein that is important in regulating transport of vitamin D, exhibiting a

higher allele frequency within individuals of European (58%) versus African (7.3%) descent.

rs7041 has been shown to correlate with altered levels of GC within European and African-

Americans (44) and further to impact vitamin D metabolism in pregnant women (45). Comparison

of cell line and tissue analyses revealed quantitation of pAIMs encoded within Protein-L-

isoaspartate(D-aspartate) O-methyltransferase (V178I, >80% AF in African and East Asians),

Serine/threonine-protein phosphatase CPPED1 (A19D, >75% AF in Africans) and AH receptor-

interacting protein (Q228K, >99% AF in European and East Asians) across all samples. Protein-

L-isoaspartate(D-aspartate) O-methyltransferase (PCMT1) regulates methyl esterification of L-

isoaspartyl and D-aspartyl residues and has been shown to repair damage to aspartate resides that

can occur in an age-dependent manner (46). Interestingly, the I178 variant of PCMT1 has been

shown to result in increased catalytic activity and a more thermostable version of PCMT1, while

the V178 variants exhibits greater substrate affinity. Serine/threonine-protein phosphatase

CPPED1 has been shown to function as a tumor suppressor in bladder cancer via inactivation of

Protein Kinase B (AKT) through dephosphorylation of S473 leading to inhibition of cell cycle

progression and promotion of apoptosis (47). Although the A19D ancestry-linked substitution we

have observed is not predicted to impact CPPED1 function, we have previously reported higher

mutational frequencies in the gene encoding tumor suppressor phosphatase and tensin homolog

(PTEN) in European-American versus African-American endometrial cancer patients, a regulator

of AKT signaling, that correlates with improved outcome in European-American women (42).

This intersection suggests that regulation of AKT signaling associated protein machinery may be

linked to patient ancestry and warrants further investigation of these relationships and their impact

on disease pathogenesis.

Our analyses provide proof of concept that estimates of European, African and East Asian ancestry

can be determined by routine proteomic analysis of disease-relevant patient tissues. We find that

pAIMs exhibiting higher allele frequencies within European, African or East Asian ancestry

individuals are quantified at significantly greater abundances in cell lines and tissue samples

derived from individuals of similar self-described ancestry, i.e. European-American or African-

American, and further exhibited abundance trends correlating with patient genotype. Our findings

also underscore that quantitation of pAIMs is feasible in formalin-fixed, paraffin embedded as well

as fresh frozen human tissue samples. These data extend previous efforts showing that genetically

variant peptides can provide estimates of global ancestry in human hair and bone and underscore

that these estimates can be determined using multiplexed proteomic data similar to that being

generated for population-scale proteogenomic studies focused on improving our understanding of

cancer. We further show that we can accurately estimate global ancestry by quantifying as few as

20 population-specific pAIMs in comparison with genotype analyses of ~266K genetic markers

using standard estimates (R^2>0.99). Although our focus has been to characterize pAIMs in uterine

neoplasms, these candidates are not tissue dependent and represent a generalized set of markers

for proteoancestry assessment in diverse tissues and cell lines. To that end, comparison of proteins

encoding pAIMs with large-scale proteomic analyses of diverse human cell and tissue types (48)

suggests that many of these protein targets are ubiquitously expressed across various tissues,

including diverse immune cell types (data not shown) and that quantitation of pAIMs in a wide

range of biological samples is likely possible. We further note that several pAIMs quantified also

map to largely predicted proteins (Uniprot.org and noted in Supplemental Tables 2 thru 4)

underscoring the need to further refine pAIMs panels with the goal of prioritizing candidates that

estimate patient ancestry with the greatest sensitivity and specificity. These findings will further

support future efforts focused on optimizing tissue-specific, stable isotope-labelled pAIMs panels

that can be co-quantified during primary sample analyses.

We have characterized ancestry-linked peptide variants in endometrial cancer cell lines and patient

tissues representing a foundational step towards investigating relationships linking patient ancestry

with the proteome, genome and disease pathogenesis. To this end, recent investigations have

revealed that clinical serum biomarkers can exhibit altered performance directly as a product of

differences in patient genetic ancestry, underscoring the need to integrate proteomics with genetic

ancestry assessments at the level of investigational analyses such as biomarker discovery (49).

Racial disparities in the incidence and outcome of uterine neoplasms persists following

adjustments for socioeconomic variables, such as patient access to care (50) and has fueled efforts

to define ancestry-linked molecular alterations that may drive these differences. Our ability to

discern these relationships will emerge through the assessment of more ancestrally-diverse

populations of women suffering from gynecologic disease as well as within disease-relevant *in*

vitro and *in vivo* models. We recently showed that historic cell line models of cancer have been

predominantly established from individuals of European ancestry (51), noting that cancers

impacted by racial disparities, such as endometrial cancer, often lacked cell line models from non-

European individuals. We confirm here three endometrial cancer cell line models, i.e. NCI-EC1,

ACI-80 and ACI-181 are derived from women of African ancestry representing *in vitro* models to

investigate ancestry-linked disease biology in endometrial cancers. This work will support ongoing

efforts focused on linking the proteome and genome with patient ancestry to better understand

relationships with disease mechanisms driving racial disparities in the pathogenesis of uterine

neoplasms.

Methods

Prioritization and *In Silico* Analyses of Peptide Ancestry Information Markers.

80,855,907 SNPs for European (i.e. FIN, GBR, IBS, and TSI; N=404), African (i.e. ESN, GWD,

LWK, MSL, and YRI; N=504), and East Asian (i.e. CDX, CHB, CHS, and KHV; N=400) were

downloaded from the 1000 Genomes Project (23). SNPs were further mapped to refSeq version

85 to identify candidate missense variants (646,768 nsSNPs) occurring within putative tryptic

peptides ≥6 or <40 amino acids in length that did not encode isoleucine to leucine substitutions

and were further filtered for those exhibiting ≥ 50% difference in allele frequency between

European, African, and East Asian populations. These substitutions were then mapped to several

RefSeq proteome databases to confirm amino acid substitution positions and tryptic specificity

resulting in a final subset of 1,037 candidates. nsSNPs encoding pAIMs were mapped to rsIDs

using Kaviar (http://db.systemsbiology.net/kaviar/cgi-pub/Kaviar.pl) against hg19 (GRCh37).

Candidates were further assessed using the Variant Effect Predictor tool

(https://useast.ensembl.org/Homo_sapiens/Tools/VEP) against hg19 (GRCh37) and extracted

metadata detailing putative clinical significance, i.e. ClinVar traits and clinical significance, as

well as predicted impact on protein function, i.e. SIFT (27) and Poly-Phen-2 (28) and PROVEAN

(29) outputs. Candidates were further mapped to the Uniprot resource and amino acid position of

pAIM substitutions were correlated with putative functional domains within parent proteins and

pAIM substitutions positions mapping with putative domain regions were prioritized for

downstream analyses. Functional enrichment analyses was performed using the "express analysis"

settings in MetaScape (52).

Random Sampling Analysis

After the 1037 pAIMs were selected from the 1000 Genomes Project training data, we then performed ADMIXTURE (33) randomly sampling 20 pAIMs without replacement, using the supervised learning version (32) with the training data as fixed for their respective ancestries (K=3 for African, European, and East Asian). Test data also came from the following 1000 Genomes Project populations (23), which were not included in the training: JPT to represent East Asian, CEU to represent European, and both ACB and ASW as admixed African ancestry. This procedure was repeated 500 times with the mean and standard deviation calculated and plotted against the estimates for these same samples from 266,403 SNPs. This larger set of sites was selected in the standard way (53), where they were LD pruned (using the PLINK(54) command --indep-pairwise 50 5 0.1), and filtered for allele frequency of 5%.

Sample Prep for Proteomics

Cell Lines – Commercial endometrial cancer cell lines, i.e. MFE-296 cells were purchased from DSMZ GmbH (Berlin, Germany) and cultured in MEM with 10% fetal bovine serum (FBS) and 1X penicillin/streptomycin (Pen/Strep). SNG-M cells were purchased from the JCRB cell bank (Osaka, Japan) and cultured in Ham's F12 median with 10% FBS and 1X Pen/Strep. HEC1A (McCoy's 5a with 10% FBS and 1X Pen/Strep), RL95-2 (DMEM:F12, 0.005 mg/mL insulin, 10% FBS and 1X Pen-Strep), Ishikawa (MEM with 2mM Glutamine, 5% Fetal Bovine Serum (FBS) and 1X Pen/Strep), AN3CA (EMEM with 10% FBS and 1X Pen/Strep) and KLE (DMEM:F12 with 10% FBS and 1X Pen/Strep) cells were from ATCC (Manassas, VA). NCI-EC1, ACI-80, ACI-181, ACI-52, ACI-61 and ACI-68 were obtained from John Risinger at Michigan State University (East Lansing, MI) and maintained in DMEM:F12 with 10% FBS and 1X Pen/Strep (ATCC). Cell lines were plated at equivalent densities and lysed at ~80% confluency in 1% SDS

and 10mM Tris-HCl, pH 7.4. Cell lysates (30 µg) were run approximately one inch into a 4-15%

bis-acrylamide gel (Bio-Rad) and processed for in-gel digestion as previously described (55).

Peptide extracts were labelled with TMT-11 reagents (Thermo Fisher Scientific) as described

below.

Uterine Leiomyoma Tissues (Discovery cohort) – Age-matched, formalin-fixed, paraffin

embedded uterine leiomyomas from self-described white and black women were obtained under

an institutional-review board approved protocol from Inova Fairfax Hospital (Falls Church, VA).

Tissue specimens were hematoxylin and eosin (H&E) stained and evaluated by a pathologist to

confirm diagnosis. Thin (10 µm) tissue sections were cut using a cryostat and placed on

polyethylene naphthalate membrane slides. After staining with aqueous H&E, laser

microdissection (LMD) was used to harvest tumor cells from thin sections.

Uterine Fibroid Tissues (Validation cohort) - Flash-frozen uterine leiomyomas were obtained

under an institutional-review board approved protocol from Johns Hopkins Memorial Institute

(Baltimore, MD). Tissues were cryopulverized, resuspended in 100 mM triethylammonium

bicarbonate (TEAB), and sonicated. Protein was quantified by BCA Protein Assay (Thermo

Scientific) and 50 µg of total protein in 100 mM TEAB and 10% acetonitrile was incubated at 99

ºC for 1 h.

Sample digestion, TMT labeling and offline fractionation

Sample digestion, TMT labeling and offline fractionation of tissue samples was performed as

recently described (56). Briefly, tissues were collected in 20 µL of 100 mM TEAB in MicroTubes

(Pressure Biosciences, Inc). Following the addition of 1 µg of SMART Digest Trypsin (Thermo

Scientific) to each sample, MicroTubes were capped with MicroPestles. Pressure-assisted lysis

and digestion was performed in a barocycler (2320 EXT, Pressure BioSciences, Inc) by

sequentially cycling between 45 kpsi and atmospheric pressure for 60 cycles at 50 ℃. The peptide

digests were transferred to 0.5 mL microcentrifuge tubes, vacuum-dried, resuspended in 100 mM

TEAB, pH 8.0 and the peptide concentration of each digest was determined using the bicinchoninic

acid assay (BCA assay). Forty-fifty micrograms of peptide from each sample, along with a pooled

reference sample assembled from equivalent amounts of peptide digests pooled from individual

patient samples for individual sample sets, were aliquoted into a final volume of 100 µL of 100

mM TEAB and labeled with tandem-mass tag (TMT) isobaric labels (TMT10 or 11plex™ Isobaric

Label Reagent Set, Thermo Fisher Scientific) according to the manufacturer's protocol. Each TMT

sample plex was fractionated by basic reversed-phase liquid chromatography (bRPLC) into 96

fractions through development of a linear gradient of acetonitrile (0.69%/min). Concatenated

fractions (36 total pooled samples for the endometrial cancer cell lines, FFPE ULMs, 24 total for

the frozen ULMs) were generated in a serpentine fashion for global LC-MS/MS analysis.

Quantitative Proteomics - LC-MS/MS Analyses

The TMT sample multiplex bRPLC fractions were resuspended in 100 mM NH_4HCO_3 and

analyzed by LC-MS/MS employing a nanoflow LC system (EASY-nLC 1200, Thermo Fisher

Scientific) coupled online with an Orbitrap Fusion Lumos MS or Q-Exactive HF-X (Thermo

Fisher Scientific). In brief, each fraction (~500 ng total peptide) was loaded on a nanoflow HPLC

system fitted with a reversed-phase trap column (Acclaim PepMap100 C18, 20 mm, nanoViper,

Thermo Scientific) and a heated (50 °C) reversed-phase analytical column (Acclaim PepMap

RSLC C18, 2 µm, 100 Å, 75 µm × 500 mm, nanoViper, Thermo Fisher Scientific) coupled online

with the MS. Peptides were eluted by developing a linear gradient of 2% mobile phase B (95%

acetonitrile, 0.1% formic acid) to 32% mobile phase B over 120 min at a constant flow rate of 250

nL/min. For both instrument platforms, the electrospray source capillary voltage and temperature

were set at 2.0 kV and 275 °C, respectively. High resolution (R=60,000 at m/z 200) broadband

(m/z 400-1600) mass spectra (MS) were acquired, followed by selection of the top 12 most intense

molecular ions in each MS scan for high-energy collisional dissociation (HCD). Instrument

specific parameters were set as follows for each instrument platform. Orbitrap Fusion Lumos -

Full MS: AGC, 5e5; RF Lens, 30%; Max IT, 50ms; Charge State, 2-4; Dynamic Exclusion,

10ppm/20 sec; MS2: AGC, 1e5; Max IT, 120ms; Resolution, 50k; Quadrupole Isolation, 0.8 m/z;

Isolation Offset, 0.2 m/z; HCD, 38; First Mass, 100. Q Exactive HF-X - Full MS: AGC, 3e6; RF

Lens, 40%; Max IT, 45ms; Charge State, 2-4; Dynamic Exclusion, 10ppm/20 sec; MS2: AGC,

1e5; Max IT, 95ms; Resolution, 45k; Quadrupole Isolation, 1.0 m/z; Isolation Offset, 0.2 m/z;

NCE, 34; First Mass, 100; Intensity Threshold, 2e5; TMT Optimization, On.

DNA extraction for endometrial cancer cell lines and uterine fibroid tissue samples

Cells were lysed in SNET Digestion buffer (20 mM Tris, 5 mM EDTA, 400 mM NaCl, 1% w/v

SDS) with Proteinase K (Thermo Fisher Scientific, Pittsburgh, PA) overnight at 55 ºC with

intermittent shaking. An equal volume of 25:24:1 phenol:chloroform:isoamyl alcohol (Thermo

Fisher Scientific, Pittsburgh, PA) was added, mixed by inversion and centrifuged at 21,000 x *g* for

5 min. The supernatant was removed to a new tube, half sample volume of 7.5 M ammonium

acetate and one and a half sample volumes of 100% ethanol were added, mixed by inversion and

centrifuged at 14,000 x *g* for 10 min. The DNA pellet was washed with 70% ethanol and

centrifuged at 14,000 x *g* for 5 min. The supernatant was removed and the pellet was air-dried and

resuspended in 50 μL of 10 mM Tris buffer. Tissue scrolls were generated from OCT-embedded

fresh-frozen tumors and collected in ATL buffer (Qiagen Sciences LLC, Germantown, MD)

followed by storage at -80 ºC until isolation. Samples were normalized to 360 μL ATL buffer, 40

μL of Proteinase K was added for lysis and incubated at 56 ºC for 4 h with intermittent shaking.

Isolation was performed according to the manufacturer's protocol (DNA Purification from

Tissues) using the QiAamp DNA Mini Kit (Qiagen Sciences LLC, Germantown, MD). DNA was

eluted after a 10 min incubation with 40 μL of Buffer AE, followed by another 10 min incubation

with 160 μL of nuclease-free water (Thermo Fisher Scientific) and reduced to 50 μL using a

CentriVap Concentrator (Labconco, Kansas City, MO). Quantity and 260/280 purity reading was

established using the Nanodrop 2000 Spectrophotometer (Thermo Fisher Scientific) and Quant-iT

PicoGreen dsDNA Assay Kit (Thermo Fisher Scientific) according to manufacturer's protocols

using eight standards from 0 to 50 ng/uL. Samples were run in triplicate and measurements were

taken on a SpectraMax M4 microplate reader (Molecular Devices, San Jose, CA).

DNA PCR-free library preparation and whole genome sequencing

TruSeq DNA PCR-free Library Preparation Kit (Illumina, San Diego, CA) was performed

following manufacturer's instructions. Briefly, genomic DNA (gDNA) was diluted to 20 ng/μL

using Resuspension Buffer (RSB, Illumina) and 55 μL were transferred to Covaris microTubes

(Covaris, Woburn, MA). The normalized gDNA was then sheared on an LE220 focused-

ultrasonication system (Covaris) to achieve target peak of 450 bp with an Average Power of 81.0

W (SonoLab settings: duty factor, 18.0%; peak incident power, 45.0 watts; 200 cycles per burst;

treatment duration, 60 s; water bath temperature, 5 °C – 8.5 °C). The quality of the final DNA

libraries was assessed with the High Sensitivity dsDNA (AATI). Per manufacturer's protocol,

library peak size was in the range of 550 to 620 bp. The DNA libraries were quantified by real-

time quantitative PCR, using the KAPA SYBR FAST Library Quantification Kit (KAPA

Biosystems, Boston, MA) optimized for the Roche LightCycler 480 instrument (Roche). DNA

libraries were then normalized to 2 nM and clustered on the Illumina cBot 2 at 200pM using a

HiSeq X Flowcell v2 and the HiSeq X HD Paired-End Cluster Generation Kit v2. Paired-end

sequencing was performed with the HiSeq X HD SBS Kit (300 cycles) on the Illumina HiSeq X.

Data Processing – Proteomics

Peptide ancestry informative marker identifications are generated by searching .RAW data files

with a publicly-available human proteome database (RefSeq human proteome, downloaded

9/22/17 and updated with entries from 8/19/2019 when pAIM amino acid positions did not map to

entries in 9/22/17 version) appended with 1,037 full-length proteins encoding predicted pAIM

substitutions of interested using Mascot (v2.6.0, Matrix Science) and Proteome Discoverer

(v2.2.0.388, Thermo Fisher Scientific) as previously described (56). Samples are searched using

the following parameters: precursor mass tolerance of 10 ppm, fragment ion tolerance of 0.05 Da,

a maximum of two tryptic miscleavages, dynamic modifications for oxidation (15.9949 Da) on

methionine residues and TMT reporter ion tags (229.1629 Da) on peptide N-termini and lysine

residues. Peptide spectral matches (PSMs) are filtered using a false-discovery rate (FDR) < 1.0%

(Percolator q-value < 0.01). TMT reporter ion intensities are extracted at a mass tolerance of 20

ppm and PSMs lacking TMT reporter ion signal in the pooled reference channel, in all patient

TMT channels or exhibiting an isolation interference of ≥50% were excluded from downstream

analyses. Normalized peptide abundance was determined by calculating TMT reporter ion ratios

relative to a pooled reference channel and mode centered, z-score transformation of each sample

channel per TMT multiplex, i.e. normalized (PSM (Log_2Ratio) = (PSM (Log_2Ratio) – ModeCenter

(PSM (Log_2Ratio) / σ (PSM (Log_2Ratio). Peptide fragment ion spectra for pAIM peptide variants

are confirmed to encode the substitution of interest by automated inspection of MS2 fragment ion

spectra in .mzML converted .RAW files (MSConvert, Proteowizard) using SpectrumAI (35) and

the following settings, fragment ion tolerance was set to 50 ppm, relative set to True and a

candidate was considered high-confidence if diagnostic ions flanking the substitution of interest

were identified in at least one sample multiplex in endometrial cancer cell line and uterine

leiomyoma cohorts. Differential analyses of pAIM abundance was performed using Mann-

Whitney U rank sum testing in MedCalc (version 19.0.3). pAIMs variants were visualized in

heatmaps and by principle component analysis (PCA) using default settings in the ClustVis web

tool (https://biit.cs.ut.ee/clustvis/). Sparse Partial Least Squares Discriminant analysis (sPLS-DA)

(37) was performed using mixOmics (ver 6.8.5) in RStudio (ver 3.6.0). The sPLS-DA model was

run in regression mode for an optimized 18 feature set for two principal components. Plot loadings

were calculated using the median to assess contribution onto the first principal component. The

AUC and ROC curve were generated from the sPLS-DA model.

DNA WGS processing and variant calling

WGS sample raw reads were aligned to the hg19 reference genome and further processed through

the Resequencing workflow within Illumina's HiSeq Analysis Software (HAS; Isis version

2.5.55.1311;   https://support.illumina.com/sequencing/sequencing_software/hiseq-analysis-

software-v2-1.html). This workflow utilizes the Isaac read aligner (iSAAC-SAAC00776.15.01.27)

and variant caller (starka-2.1.4.2) (57), the Manta structural variant caller (version 0.23.1) (58),

and the Canvas CNV caller (version 1.1.0.5) (59).).

Global ancestry proportion of EC cell lines

To estimate admixture proportions, we used reference samples with known ancestry from the 1000

Genome Project (23). These reference samples were comprised of 99 European samples from the

CEU population, 108 African samples from the YRI population, and 208 Asian samples from the

CHB and JPT populations (414 total references samples). Working with genotype data from these

reference samples and each of the endometrial cancer cell lines, we first used bcftools(60) to

remove Indels and non-biallelic variants. We then used Plink v1.90b3.32 (54)to remove singleton

sites (i.e. variants with only one alt allele copy in the dataset. After this processing, we merged

these datasets with Plink by taking the intersect of autosomal variants and removing any

ambiguous A/T and G/C variants from this combined data. We then pruned for linkage using the

plink linkage pruning algorithm command of --indep-pairwise 50 5 0.5, which uses a window of

50 with an r^2 greater than 0.5 and a SNP step of 5. Ancestry was estimated for each cell line

independently, and so this process was repeated independently for each cell line. At the end of this

processing, we were left with ~500,000 LD-pruned variants per reference cell line merged dataset

(range 459,592 – 701,192).

Global ancestry proportion of uterine fibroid patients

We predicted sample super population ancestries using the methods implemented in Peddy (61).

Briefly, principal component reduction was performed on genotype calls at specific loci from

2,504 samples in the 1000 Genomes project and a support vector machine (SVM) classifier was

trained on the resulting first four components, using known ancestries as the training labels.

Genotype calls at the same loci from each sample collected in this study were then mapped to

principal component space and the trained SVM was used to predict ancestries. All classifier

prediction probabilities were >0.89.

Data Availability: Supplemental data tables include chromosomal loci, missense substitutions

and amino acid position, reference SNP Ids (rsIDs) corresponding to peptide ancestry informative

markers (pAIMs), as well as pAIM abundances quantified in endometrial cancer cell lines and

uterine leiomyoma tissues. RAW LC-MS data files as well as peptide spectrum match exports

from Proteome Discoverer will be deposited in ProteomeXchange. Please contact

batemann@whirc.org with Supplemental Data inquiries.

Acknowledgments: The authors would like to thank Tracy Litzi, BS, Glenn Gist, BS, Julie Oliver,

MS, Dave Mitchell, MS, Domenic Tommarello, MS, Jeremy Loffredo, BS and Wei Ao, BS for

their technical contributions as well as Albert Dobi, PhD, Gyorgi Petrovics, PhD and Sara

Scannell, BS for critical review of this manuscript. A patent based on the concept presented in this

study has been awarded (US 8,877,455 B2, Australian Patent 2011229918, Canadian Patent CA

2794248, and European Patent EP11759843.3); GJP inventor. GLM is a consultant for Kiyatec,

GSK, and Merck. TPC is a ThermoFisher Scientific, Inc SAB member and receives research

funding from AbbVie. Disclaimer: The views expressed herein are those of the authors and do not

reflect the official policy of the Department of Army/Navy/Air Force, Department of Defense, or

U.S. Government.

Author Contributions: Contributed to conception: NWB, TDO, TPC, GLM. Contributed to

experimental design: NWB, TDO, TPC. Contributed to data acquisition, analysis and/or

interpretation of data: NWB, TDO, CMT, TA, BLH, KC, MZ, PT, ARS, AJ, CT, CD, MW. Drafted

and/or revised the article: NWB, TDO, ARS, MK, CDS, HH, MC, GJP, JS, AAH, JRR, KD, YC,

TPC, GLM. Acquired funding for the research: YC, GLM. All authors read and approved the final

manuscript.

References

1.   Gupta, N., Tanner, S., Jaitly, N., Adkins, J. N., Lipton, M., Edwards, R., Romine, M.,
Osterman, A., Bafna, V., Smith, R. D., and Pevzner, P. A. (2007) Whole proteome analysis
of post-translational modifications: applications of mass-spectrometry for proteogenomic
annotation. *Genome Res* **17**, 1362-1377
2.   Ng, P. C., and Henikoff, S. (2003) SIFT: Predicting amino acid changes that affect protein
function. *Nucleic Acids Res* **31**, 3812-3814
3.   Suhre, K., McCarthy, M. I., and Schwenk, J. M. (2020) Genetics meets proteomics:
perspectives for large population-based studies. *Nat Rev Genet*
4.   Mertins, P., Mani, D. R., Ruggles, K. V., Gillette, M. A., Clauser, K. R., Wang, P., Wang,
X., Qiao, J. W., Cao, S., Petralia, F., Kawaler, E., Mundt, F., Krug, K., Tu, Z., Lei, J. T.,

584   Gatza, M. L., Wilkerson, M., Perou, C. M., Yellapantula, V., Huang, K. L., Lin, C.,
585   McLellan, M. D., Yan, P., Davies, S. R., Townsend, R. R., Skates, S. J., Wang, J., Zhang,
586   B., Kinsinger, C. R., Mesri, M., Rodriguez, H., Ding, L., Paulovich, A. G., Fenyo, D., Ellis,
587   M. J., Carr, S. A., and Nci, C. (2016) Proteogenomics connects somatic mutations to
588   signalling in breast cancer. *Nature* **534**, 55-62

589 5.  Fiore, L. D., Rodriguez, H., and Shriver, C. D. (2017) Collaboration to Accelerate
590   Proteogenomics Cancer Care: The Department of Veterans Affairs, Department of
591   Defense, and the National Cancer Institute's Applied Proteogenomics OrganizationaL
592   Learning and Outcomes (APOLLO) Network. *Clin Pharmacol Ther* **101**, 619-621

593 6.  Spratt, D. E., Chan, T., Waldron, L., Speers, C., Feng, F. Y., Ogunwobi, O. O., and
594   Osborne, J. R. (2016) Racial/Ethnic Disparities in Genomic Sequencing. *JAMA Oncol* **2**,
595   1070-1074

596 7.  Deshmukh, S. K., Azim, S., Ahmad, A., Zubair, H., Tyagi, N., Srivastava, S. K., Bhardwaj,
597   A., Singh, S., Rocconi, R. P., and Singh, A. P. (2017) Biological basis of cancer health
598   disparities: resources and challenges for research. *Am J Cancer Res* **7**, 1-12

599 8.  Tarney, C. M., Tian, C., Wang, G., Dubil, E. A., Bateman, N. W., Chan, J. K., Elshaikh,
600   M. A., Cote, M. L., Schildkraut, J. M., Shriver, C. D., Conrads, T. P., Hamilton, C. A.,
601   Maxwell, G. L., and Darcy, K. M. (2018) Impact of age at diagnosis on racial disparities
602   in endometrial cancer patients. *Gynecol Oncol* **149**, 12-21

603 9.  Felix, A. S., Linkov, F., Maxwell, G. L., Ragin, C., and Taioli, E. (2011) Racial disparities
604   in risk of second primary cancers in endometrial cancer patients: Analysis of SEER Data.
605   *Int J Gynecol Cancer* **21**, 309-315

606 10.  Allard, J. E., and Maxwell, G. L. (2009) Race disparities between black and white women
607   in the incidence, treatment, and prognosis of endometrial cancer. *Cancer Control* **16**, 53-
608   56

609 11.  Farley, J., Risinger, J. I., Rose, G. S., and Maxwell, G. L. (2007) Racial disparities in blacks
610   with gynecologic cancers. *Cancer* **110**, 234-243

611 12.  Maxwell, G. L., Tian, C., Risinger, J., Brown, C. L., Rose, G. S., Thigpen, J. T., Fleming,
612   G. F., Gallion, H. H., Brewster, W. R., and Gynecologic Oncology Group, s. (2006) Racial
613   disparity in survival among patients with advanced/recurrent endometrial adenocarcinoma:
614   a Gynecologic Oncology Group study. *Cancer* **107**, 2197-2205

615 13.  Mahdi, H., Schlick, C. J., Kowk, L. L., Moslemi-Kebria, M., and Michener, C. (2014)
616   Endometrial cancer in Asian and American Indian/Alaskan Native women: tumor
617   characteristics, treatment and outcome compared to non-Hispanic white women. *Gynecol*
618   *Oncol* **132**, 443-449

619 14.  Enoch, M. A., Shen, P. H., Xu, K., Hodgkinson, C., and Goldman, D. (2006) Using
620   ancestry-informative markers to define populations and detect population stratification. *J*
621   *Psychopharmacol* **20**, 19-26

622 15.  Kosoy, R., Nassir, R., Tian, C., White, P. A., Butler, L. M., Silva, G., Kittles, R., Alarcon-
623   Riquelme, M. E., Gregersen, P. K., Belmont, J. W., De La Vega, F. M., and Seldin, M. F.
624   (2009) Ancestry informative marker sets for determining continental origin and admixture
625   proportions in common populations in America. *Hum Mutat* **30**, 69-78

626 16.  Galanter, J. M., Fernandez-Lopez, J. C., Gignoux, C. R., Barnholtz-Sloan, J., Fernandez-
627   Rozadilla, C., Via, M., Hidalgo-Miranda, A., Contreras, A. V., Figueroa, L. U., Raska, P.,
628   Jimenez-Sanchez, G., Zolezzi, I. S., Torres, M., Ponte, C. R., Ruiz, Y., Salas, A., Nguyen,
629   E., Eng, C., Borjas, L., Zabala, W., Barreto, G., Gonzalez, F. R., Ibarra, A., Taboada, P.,

Porras, L., Moreno, F., Bigham, A., Gutierrez, G., Brutsaert, T., Leon-Velarde, F., Moore, L. G., Vargas, E., Cruz, M., Escobedo, J., Rodriguez-Santana, J., Rodriguez-Cintron, W., Chapela, R., Ford, J. G., Bustamante, C., Seminara, D., Shriver, M., Ziv, E., Burchard, E. G., Haile, R., Parra, E., Carracedo, A., and Consortium, L. (2012) Development of a panel of genome-wide ancestry informative markers to study admixture throughout the Americas. *PLoS Genet* **8**, e1002554

17. Mathias, R. A., Taub, M. A., Gignoux, C. R., Fu, W., Musharoff, S., O'Connor, T. D., Vergara, C., Torgerson, D. G., Pino-Yanes, M., Shringarpure, S. S., Huang, L., Rafaels, N., Boorgula, M. P., Johnston, H. R., Ortega, V. E., Levin, A. M., Song, W., Torres, R., Padhukasahasram, B., Eng, C., Mejia-Mejia, D. A., Ferguson, T., Qin, Z. S., Scott, A. F., Yazdanbakhsh, M., Wilson, J. G., Marrugo, J., Lange, L. A., Kumar, R., Avila, P. C., Williams, L. K., Watson, H., Ware, L. B., Olopade, C., Olopade, O., Oliveira, R., Ober, C., Nicolae, D. L., Meyers, D., Mayorga, A., Knight-Madden, J., Hartert, T., Hansel, N. N., Foreman, M. G., Ford, J. G., Faruque, M. U., Dunston, G. M., Caraballo, L., Burchard, E. G., Bleecker, E., Araujo, M. I., Herrera-Paz, E. F., Gietzen, K., Grus, W. E., Bamshad, M., Bustamante, C. D., Kenny, E. E., Hernandez, R. D., Beaty, T. H., Ruczinski, I., Akey, J., Caapa, and Barnes, K. C. (2016) A continuum of admixture in the Western Hemisphere revealed by the African Diaspora genome. *Nat Commun* **7**, 12522

18. Kessler, M. D., Yerges-Armstrong, L., Taub, M. A., Shetty, A. C., Maloney, K., Jeng, L. J., Ruczinski, I., Levin, A. M., Williams, L. K., Beaty, T. H., Mathias, R. A., Barnes, K. C., Consortium on Asthma among African-ancestry Populations in the, A., and O'Connor, T. D. (2016) Challenges and disparities in the application of personalized genomic medicine to populations with African ancestry. *Nat Commun* **7**, 12521

19. Tennessen, J. A., Bigham, A. W., O'Connor, T. D., Fu, W., Kenny, E. E., Gravel, S., McGee, S., Do, R., Liu, X., Jun, G., Kang, H. M., Jordan, D., Leal, S. M., Gabriel, S., Rieder, M. J., Abecasis, G., Altshuler, D., Nickerson, D. A., Boerwinkle, E., Sunyaev, S., Bustamante, C. D., Bamshad, M. J., Akey, J. M., Broad, G. O., Seattle, G. O., and Project, N. E. S. (2012) Evolution and functional impact of rare coding variation from deep sequencing of human exomes. *Science* **337**, 64-69

20. Tennessen, J. A., O'Connor, T. D., Bamshad, M. J., and Akey, J. M. (2011) The promise and limitations of population exomics for human evolution studies. *Genome Biol* **12**, 127

21. Parker, G. J., Leppert, T., Anex, D. S., Hilmer, J. K., Matsunami, N., Baird, L., Stevens, J., Parsawar, K., Durbin-Johnson, B. P., Rocke, D. M., Nelson, C., Fairbanks, D. J., Wilson, A. S., Rice, R. H., Woodward, S. R., Bothner, B., Hart, B. R., and Leppert, M. (2016) Demonstration of Protein-Based Human Identification Using the Hair Shaft Proteome. *PLoS One* **11**, e0160653

22. Feng Lei, J. L., Li Shaun-Fei, Zhang Jian, Liu Hai-bo, Ji An-Quan, Ye Jian, Wang Gui-Qiang, Li Cai-Xia. (2019) Development and Validation of Protein-based Forensic Ancestry Inference Method Using Hair Shaft Proteome. *Progress in Biochemistry and Biophysics* **46（1）：81~88**

23. Genomes Project, C., Auton, A., Brooks, L. D., Durbin, R. M., Garrison, E. P., Kang, H. M., Korbel, J. O., Marchini, J. L., McCarthy, S., McVean, G. A., and Abecasis, G. R. (2015) A global reference for human genetic variation. *Nature* **526**, 68-74

24. McLaren, W., Gil, L., Hunt, S. E., Riat, H. S., Ritchie, G. R., Thormann, A., Flicek, P., and Cunningham, F. (2016) The Ensembl Variant Effect Predictor. *Genome Biol* **17**, 122

25. Ruiz-Iruela, C., Padro-Miquel, A., Pinto-Sala, X., Baena-Diez, N., Caixas-Pedragos, A.,
Guell-Miro, R., Navarro-Badal, R., Jusmet-Miguel, X., Calmarza, P., Puzo-Foncilla, J. L.,
Alia-Ramos, P., and Candas-Estebanez, B. (2018) KIF6 gene as a pharmacogenetic marker
for lipid-lowering effect in statin treatment. *PLoS One* **13**, e0205430
26. Xu, L., Zhou, X., Jiang, F., Xu, L., and Yin, R. (2014) STK15 rs2273535 polymorphism
and cancer risk: a meta-analysis of 74,896 subjects. *Cancer Epidemiol* **38**, 111-117
27. Sim, N. L., Kumar, P., Hu, J., Henikoff, S., Schneider, G., and Ng, P. C. (2012) SIFT web
server: predicting effects of amino acid substitutions on proteins. *Nucleic Acids Res* **40**,
W452-457
28. Adzhubei, I., Jordan, D. M., and Sunyaev, S. R. (2013) Predicting functional effect of
human missense mutations using PolyPhen-2. *Curr Protoc Hum Genet* **Chapter 7**, Unit7
20
29. Choi, Y., Sims, G. E., Murphy, S., Miller, J. R., and Chan, A. P. (2012) Predicting the
functional effect of amino acid substitutions and indels. *PLoS One* **7**, e46688
30. Ren, Y., Liu, W., Chen, J., Wang, J., Wang, K., Zhou, J., Cai, S., Zheng, M., Liu, J., Liu,
L., and Xue, D. (2017) A missense variant of the ABCC11 gene is associated with Axillary
Osmidrosis susceptibility and clinical phenotypes in the Chinese Han Population. *Sci Rep*
7, 46335
31. Ishiguro, J., Ito, H., Tsukamoto, M., Iwata, H., Nakagawa, H., and Matsuo, K. (2019) A
functional single nucleotide polymorphism in ABCC11, rs17822931, is associated with the
risk of breast cancer in Japanese. *Carcinogenesis* **40**, 537-543
32. Alexander, D. H., and Lange, K. (2011) Enhancements to the ADMIXTURE algorithm for
individual ancestry estimation. *BMC bioinformatics* **12**, 246
33. Alexander, D. H., Novembre, J., and Lange, K. (2009) Fast model-based estimation of
ancestry in unrelated individuals. *Genome Res* **19**, 1655-1664
34. Kessler, M. D., Yerges-Armstrong, L., Taub, M. A., Shetty, A. C., Maloney, K., Jeng, L.
J., Ruczinski, I., Levin, A. M., Williams, L. K., Beaty, T. H., Mathias, R. A., Barnes, K.
C., and O'Connor, T. D. (2016) Challenges and disparities in the application of
personalized genomic medicine to populations with African ancestry. *Nat Commun* **7**,
12521
35. Zhu, Y., Orre, L. M., Johansson, H. J., Huss, M., Boekel, J., Vesterlund, M., Fernandez-
Woodbridge, A., Branca, R. M. M., and Lehtio, J. (2018) Discovery of coding regions in
the human genome by integrated proteogenomics analysis workflow. *Nat Commun* **9**, 903
36. Han, C., Liao, X., Qin, W., Yu, L., Liu, X., Chen, G., Liu, Z., Lu, S., Chen, Z., Su, H., Zhu,
G., Lu, Z., Liu, Z., Qin, X., Gui, Y., Mo, Z., Li, L., and Peng, T. (2016) EGFR and SYNE2
are associated with p21 expression and SYNE2 variants predict post-operative clinical
outcomes in HBV-related hepatocellular carcinoma. *Sci Rep* **6**, 31237
37. Le Cao, K. A., Boitard, S., and Besse, P. (2011) Sparse PLS discriminant analysis:
biologically relevant feature selection and graphical displays for multiclass problems. *BMC*
Bioinformatics **12**, 253
38. Baird, D. D., Dunson, D. B., Hill, M. C., Cousins, D., and Schectman, J. M. (2003) High
cumulative incidence of uterine leiomyoma in black and white women: ultrasound
evidence. *Am J Obstet Gynecol* **188**, 100-107
39. American Cancer Society. (2016) Cancer Facts & Figures for African Americans 2016-
2018., American Cancer Society, Atlanta, GA

40.   Rocconi, R. P., Lankes, H. A., Brady, W. E., Goodfellow, P. J., Ramirez, N. C., Alvarez,
R. D., Creasman, W., and Fernandez, J. R. (2016) The role of racial genetic admixture with
endometrial cancer outcomes: An NRG Oncology/Gynecologic Oncology Group study.
Gynecol Oncol **140**, 264-269

41.   Bateman, N. W., Dubil, E. A., Wang, G., Hood, B. L., Oliver, J. M., Litzi, T. A., Gist, G.
D., Mitchell, D. A., Blanton, B., Phippen, N. T., Tian, C., Zahn, C. M., Cohn, D. E.,
Havrilesky, L. J., Berchuck, A., Shriver, C. D., Darcy, K. M., Hamilton, C. A., Conrads,
T. P., and Maxwell, G. L. (2017) Race-specific molecular alterations correlate with
differential outcomes for black and white endometrioid endometrial cancer patients.
Cancer **123**, 4004-4012

42.   Maxwell, G. L., Risinger, J. I., Hayes, K. A., Alvarez, A. A., Dodge, R. K., Barrett, J. C.,
and Berchuck, A. (2000) Racial disparity in the frequency of PTEN mutations, but not
microsatellite instability, in advanced endometrial cancers. *Clin Cancer Res* **6**, 2999-3005

43.   Dubil, E. A., Tian, C., Wang, G., Tarney, C. M., Bateman, N. W., Levine, D. A., Conrads,
T. P., Hamilton, C. A., Maxwell, G. L., and Darcy, K. M. (2018) Racial disparities in
molecular subtypes of endometrial cancer. *Gynecol Oncol* **149**, 106-116

44.   Powe, C. E., Evans, M. K., Wenger, J., Zonderman, A. B., Berg, A. H., Nalls, M., Tamez,
H., Zhang, D., Bhan, I., Karumanchi, S. A., Powe, N. R., and Thadhani, R. (2013) Vitamin
D-binding protein and vitamin D status of black Americans and white Americans. *N Engl*
J Med **369**, 1991-2000

45.   Ganz, A. B., Park, H., Malysheva, O. V., and Caudill, M. A. (2018) Vitamin D binding
protein rs7041 genotype alters vitamin D metabolism in pregnant women. *FASEB J* **32**,
2012-2020

46.   DeVry, C. G., and Clarke, S. (1999) Polymorphic forms of the protein L-isoaspartate (D-
aspartate) O-methyltransferase involved in the repair of age-damaged proteins. *J Hum*
Genet **44**, 275-288

47.   Zhuo, D. X., Zhang, X. W., Jin, B., Zhang, Z., Xie, B. S., Wu, C. L., Gong, K., and Mao,
Z. B. (2013) CSTP1, a novel protein phosphatase, blocks cell cycle, promotes cell
apoptosis, and suppresses tumor growth of bladder cancer by directly dephosphorylating
Akt at Ser473 site. *PLoS One* **8**, e65679

48.   Kim, M. S., Pinto, S. M., Getnet, D., Nirujogi, R. S., Manda, S. S., Chaerkady, R.,
Madugundu, A. K., Kelkar, D. S., Isserlin, R., Jain, S., Thomas, J. K., Muthusamy, B.,
Leal-Rojas, P., Kumar, P., Sahasrabuddhe, N. A., Balakrishnan, L., Advani, J., George, B.,
Renuse, S., Selvan, L. D., Patil, A. H., Nanjappa, V., Radhakrishnan, A., Prasad, S.,
Subbannayya, T., Raju, R., Kumar, M., Sreenivasamurthy, S. K., Marimuthu, A., Sathe, G.
J., Chavan, S., Datta, K. K., Subbannayya, Y., Sahu, A., Yelamanchi, S. D., Jayaram, S.,
Rajagopalan, P., Sharma, J., Murthy, K. R., Syed, N., Goel, R., Khan, A. A., Ahmad, S.,
Dey, G., Mudgal, K., Chatterjee, A., Huang, T. C., Zhong, J., Wu, X., Shaw, P. G., Freed,
D., Zahari, M. S., Mukherjee, K. K., Shankar, S., Mahadevan, A., Lam, H., Mitchell, C. J.,
Shankar, S. K., Satishchandra, P., Schroeder, J. T., Sirdeshmukh, R., Maitra, A., Leach, S.
D., Drake, C. G., Halushka, M. K., Prasad, T. S., Hruban, R. H., Kerr, C. L., Bader, G. D.,
Iacobuzio-Donahue, C. A., Gowda, H., and Pandey, A. (2014) A draft map of the human
proteome. *Nature* **509**, 575-581

49.   Sjaarda, J., Gerstein, H. C., Kutalik, Z., Mohammadi-Shemirani, P., Pigeyre, M., Hess, S.,
and Pare, G. (2020) Influence of Genetic Ancestry on Human Serum Proteome. *Am J Hum*
Genet **106**, 303-314

50. Oliver, K. E., Enewold, L. R., Zhu, K., Conrads, T. P., Rose, G. S., Maxwell, G. L., and
Farley, J. H. (2011) Racial disparities in histopathologic characteristics of endometrial
cancer are present in older, not younger blacks in an equal-access environment. *Gynecol*
Oncol **123**, 76-81
51. Kessler, M. D., Bateman, N. W., Conrads, T. P., Maxwell, G. L., Dunning Hotopp, J. C.,
and O'Connor, T. D. (2019) Ancestral characterization of 1018 cancer cell lines highlights
disparities and reveals gene expression and mutational differences. *Cancer* **125**, 2076-2088
52. Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O., Benner,
C., and Chanda, S. K. (2019) Metascape provides a biologist-oriented resource for the
analysis of systems-level datasets. *Nat Commun* **10**, 1523
53. Kessler, M. D., Yerges-Armstrong, L., Taub, M. A., Shetty, A. C., Maloney, K., Jeng, L.
J. B., Ruczinski, I., Levin, A. M., Williams, L. K., Beaty, T. H., Mathias, R. A., Barnes, K.
C., Consortium on Asthma among African-ancestry Populations in the, A., and O'Connor,
T. D. (2016) Challenges and disparities in the application of personalized genomic
medicine to populations with African ancestry. *Nat Commun* **7**, 12521
54. Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A., Bender, D., Maller,
J., Sklar, P., de Bakker, P. I., Daly, M. J., and Sham, P. C. (2007) PLINK: a tool set for
whole-genome association and population-based linkage analyses. *Am J Hum Genet* **81**,
559-575
55. Bateman, N. W., Jaworski, E., Ao, W., Wang, G., Litzi, T., Dubil, E., Marcus, C., Conrads,
K. A., Teng, P. N., Hood, B. L., Phippen, N. T., Vasicek, L. A., McGuire, W. P., Paz, K.,
Sidransky, D., Hamilton, C. A., Maxwell, G. L., Darcy, K. M., and Conrads, T. P. (2015)
Elevated AKAP12 in paclitaxel-resistant serous ovarian cancer cells is prognostic and
predictive of poor survival in patients. *J Proteome Res* **14**, 1900-1910
56. Lee, S., Zhao, L., Rojas, C., Bateman, N. W., Yao, H., Lara, O. D., Celestino, J., Morgan,
M. B., Nguyen, T. V., Conrads, K. A., Rangel, K. M., Dood, R. L., Hajek, R. A., Fawcett,
G. L., Chu, R. A., Wilson, K., Loffredo, J. L., Viollet, C., Jazaeri, A. A., Dalgard, C. L.,
Mao, X., Song, X., Zhou, M., Hood, B. L., Banskota, N., Wilkerson, M. D., Te, J., Soltis,
A. R., Roman, K., Dunn, A., Cordover, D., Eterovic, A. K., Liu, J., Burks, J. K., Baggerly,
K. A., Fleming, N. D., Lu, K. H., Westin, S. N., Coleman, R. L., Mills, G. B., Casablanca,
Y., Zhang, J., Conrads, T. P., Maxwell, G. L., Futreal, P. A., and Sood, A. K. (2020)
Molecular Analysis of Clinically Defined Subsets of High-Grade Serous Ovarian Cancer.
Cell Rep **31**, 107502
57. Raczy, C., Petrovski, R., Saunders, C. T., Chorny, I., Kruglyak, S., Margulies, E. H.,
Chuang, H. Y., Kallberg, M., Kumar, S. A., Liao, A., Little, K. M., Stromberg, M. P., and
Tanner, S. W. (2013) Isaac: ultra-fast whole-genome secondary analysis on Illumina
sequencing platforms. *Bioinformatics* **29**, 2041-2043
58. Chen, X., Schulz-Trieglaff, O., Shaw, R., Barnes, B., Schlesinger, F., Kallberg, M., Cox,
A. J., Kruglyak, S., and Saunders, C. T. (2016) Manta: rapid detection of structural variants
and indels for germline and cancer sequencing applications. *Bioinformatics* **32**, 1220-1222
59. Roller, E., Ivakhno, S., Lee, S., Royce, T., and Tanner, S. (2016) Canvas: versatile and
scalable detection of copy number variants. *Bioinformatics* **32**, 2375-2377
60. Narasimhan, V., Danecek, P., Scally, A., Xue, Y., Tyler-Smith, C., and Durbin, R. (2016)
BCFtools/RoH: a hidden Markov model approach for detecting autozygosity from next-
generation sequencing data. *Bioinformatics* **32**, 1749-1751

61.   Pedersen, B. S., and Quinlan, A. R. (2017) Who's Who? Detecting and Resolving Sample
Anomalies in Human DNA Sequencing Studies with Peddy. *Am J Hum Genet* **100**, 406-
413

Tables

Table 1: Endometrial Cancer Cell Lines Analyzed by Whole Genome Sequence and

Quantitative Proteomic Analyses. Cell lines include both primary and commercial cell lines and

global ancestry was determined by standard estimates using single nucleotide variant-derived

ancestry informative markers measured by whole genome sequence analyses.

Cell line	Histology	Ancestry (Standard Estimate)
ACI181	Endometrioid	African
ACI80	Endometrioid	African
NCIEC1	Endometrioid	African
HEC1A	Adenocarcinoma NOS	East Asian
ISHIKAWA	Adenocarcinoma NOS	East Asian
SNGM	Adenocarcinoma NOS	East Asian
ACI52	Endometrioid	European
ACI61	Endometrioid	European
ACI68	Endometrioid	European
AN3CA	Adenocarcinoma NOS	European
KLE	Adenocarcinoma NOS	European
MFE296	Adenocarcinoma NOS	European
RL952	Adenocarcinoma NOS	European

Table 2: Uterine Leiomyoma (ULM) Tissues Analyzed by Quantitative Proteomic (Discovery and Validation) as well as Whole Genome Sequencing (Validation) Analysis. Global ancestry was determined by standard estimates for the validation cohort using single nucleotide variant-derived ancestry informative markers measured by whole genome sequence analyses.

Discovery - ULM Tissues (n=66)	
Ancestry (Self-Described)	**#**
European-American	35
African-American	31

Validation - ULM Tissues (n=26)	
Ancestry (Standard Estimate)	**#**
European	16
African	10

Figure Legends

Figure 1: Selection of Peptide Ancestry Informative Markers (pAIMs) and Workflow for Analyses in Cell Lines Models and Tissues of Uterine Neoplasms collected from Women of Diverse Ancestries. A: >640K single nucleotide polymorphisms (SNPs) were filtered to prioritize 1,037 missense substitutions occurring within lysine or arginine-terminating tryptic peptides, ≥ 6 and ≤ 40 amino acids in length, "so-called" peptide ancestry informative markers (pAIMs). We then investigated pAIMs in endometrial cancer cell line models as well as uterine leiomyoma

tissues from women representing diverse ancestries. B: Comparison of pAIMs exhibiting ≥ 50%

allele frequencies within individuals of European (EUR), African (AFR) or East Asian (EAS).

Figure 2: *In silico* **analyses of Peptide Ancestry Informative Markers (pAIMs) to assess**

Clinical Significance, Localization to Protein Domains as well as to Predict Impact on Protein

Function. A: Clinical significance (ClinVar) and protein domain (Uniprot) localization of 1,037

pAIMs. B: Comparison of pAIMs predicted to be deleterious to protein function by PROVEAN,

PolyPhen and SIFT functional prediction tools. C: Clinical significance (ClinVar) and protein

domain (Uniprot) localization of pAIMs predicted to be deleterious to protein function. D: Allele

frequency of pAIMs predicted to be deleterious to protein function in European (EUR), African

(AFR) or East Asian (EAS) populations

Figure 3: Comparison of European (EUR), African (AFR) or East Asian (EAS) Global

Ancestry Classification using Peptide Ancestry Informative Markers (pAIMs) with

Standard Estimates. Random sampling of pAIMs in European, African and East Asian reference

populations from 1K genome project revealed that a minimum of 20 pAIM-encoding

nonsynonymous single nucleotide polymorphisms (SNPs) can classify global ancestry with

accuracy comparable to standard estimates using >266,000 SNPs, i.e. R^2=0.9943 ± 0.005.

Figure 4: Quantitation of Peptide Ancestry Informative Markers (pAIMs) within

Endometrial cancer cell Lines. A: Box plots detail the median logFC abundance of 17 EUR

pAIMs within cell lines stratified by European (EUR), African (AFR) or East Asian (EAS)

ancestry as well as by pAIM variant allele genotype observed by companion whole genome

sequence analyses of EUR cell lines. B: Box plots details the median logFC abundance of 21 AFR

pAIMs within cell lines stratified by European (EUR), African (AFR) or East Asian (EAS)

ancestry as well as by as well as by pAIM variant allele genotype observed by companion whole

genome sequence analyses of AFR cell lines. C: Box plots detail the median logFC abundance of

17 EAS pAIMs within cell lines stratified by European (EUR), African (AFR) or East Asian (EAS)

ancestry as well as by pAIM variant allele genotype observed by companion whole genome

sequence analyses of EAS cell lines.

Figure 5: Quantitation of Peptide Ancestry Informative Markers (pAIMs) within Uterine

Leiomyoma (ULM) Tissues Collected from a Cohort of Women of Self-Described Black or

White Race. A: Data reflects unsupervised cluster analyses of 18 pAIMs quantified across n66

patient tissue samples; heatmap reflects clustering by Euclidean distance and complete linkage. B:

Principle component analyses of 18 pAIMs, principal component 1 and principal component 2 that

explain 21% and 7% of the total variance, respectively. C: Receiver operator curve assessing the

performance of 18 pAIMs to classify African-American (n31) versus European-American (n35)

patients.

Figure 6: Validation of Peptide Ancestry Informative Markers (pAIMs) within Uterine

Leiomyoma (ULM) Tissues Collected from Women with Global Ancestry determined by

Standard Estimates. A: Box plots details the median logFC abundance for pAIMs exhibiting

>50% allele frequencies in African (8 of 15 pAIMs quantified) reference populations across n10

ULMs sample from women with AFR ancestry as well as by pAIM variant allele genotype

observed by companion whole genome sequence analyses. B: Box plots details the median logFC

abundance pAIMs exhibiting >50% allele frequencies for European (5 of 15 quantified) reference

populations quantified across n=16 ULMs sample from women with EUR ancestry as well as by

pAIM variant allele genotype observed by companion whole genome sequence analyses.

Supplemental Tables

Supplemental Table 1A – Peptide Ancestry Informative Markers. Column headers denote the following ProteinAccession_pAIM: Refseq protein accession and missense substitution and amino acid position encoded by the nonsynonymous single nucleotide variant encoding pAIM, SNP-ID: chromosome and genomic loci (hg19), GeneSymbol: Gene name, rsID: reference SNP, VAR_Position: amino acid and position of the reference and variant, AF-African: pAIM allele frequency in African reference populations (1000 genomes project), AF-European: pAIM allele frequency in European reference populations (1000 genomes project), AF-EastAsian: pAIM allele frequency in East Asian reference populations (1000 genomes project), SIFT-D: YES=SIFT score for rsID is predicted to be deleterious by the Variant Effect Predictor tool (Ensembl), PROVEAN-D: YES=PROVEAN score is predicted to be deleterious by the Variant Effect Predictor tool (Ensembl), PolyPhenHVAR=D: YES=PolyPhenHVAR score is predicted to be deleterious by the Variant Effect Predictor tool (Ensembl), Clinical_Significance(CLINVAR): clinical significance for rsID (CLINVAR/ Variant Effect Predictor tool), Trait(CLINVAR): clinical trait correlating with rsID (CLINVAR/ Variant Effect Predictor tool).

Supplemental Table 1B – Peptide Ancestry Informative Markers – Functional enrichment analyses (metascape.org).

Supplemental Table 2 – pAIMs Quantified and genotype for rsID encoding pAIM - Endometrial cancer cell Lines (n=13). UniprotAccMatch: Alternate protein accessions mapping to pAIMs (Uniprot.org). GT: genotype, GT=1_1 (homozygous for variant encoding pAIM), GT=0_1 (heterozygous for pAIM), GT=0_0 (homozygous for reference).

Supplemental Table 3 – pAIMs Quantified - Uterine Leiomyomas (Discovery, n=66).

UniprotAccMatch: Alternate protein accessions mapping to pAIMs (Uniprot.org).

Supplemental Table 4 – pAIMs Quantified and genotype for rsID encoding pAIM - Uterine

Leiomyomas (Discovery, n=26). UniprotAccMatch: Alternate protein accessions mapping to

pAIMs (Uniprot.org). GT: genotype, GT=1_1 (homozygous for variant encoding pAIM), GT=0_1

(heterozygous for pAIM), GT=0_0 (homozygous for reference).

A

B

A

A

B ○ African-American △ European-American

C

African-American vs.
European-American
(AUC=0.9834, P=1.6E^{-11})

A

B

European	MWU, p	Median Difference
EURvsAFR	<0.0001	1.74
+/+ vs -/+	0.053	0.7
+/+ vs -/-	<0.0001	2.37