A novel, ataxic mouse model of Ataxia Telangiectasia caused by a clinically relevant nonsense mutation

Perez, Harvey [¥1]; Abdallah, May, F.[¥1]; Chavira, Jose, I. [¥1]; Egeland, Martin, T. [1]; Vo, Karen, L. [1];

Buechsenschuetz, Callan, L. [1]; Sanghez, Valentina [1]; Kim, Jeannie, L. [1]; Pind, Molly [2], Nakamura,

Kotoka [3], Hicks, Geoffrey, G. [2]; Gatti, Richard, A. [3]; Madrenas, Joaquin [1,5]; Iacovino, Michelina [1,4];

McKinnon, Peter, J. [6]; Mathews, Paul, J. [*1,7]

1. *The Lundquist Institute for Biomedical Innovation, Harbor-UCLA Medical Center, Torrance, CA*
2. *Department of Biochemistry & Medical Genetics, Max Rady College of Medicine, University of Manitoba*
3. *Department of Pathology & Laboratory Medicine, David Geffen School of Medicine, University of California, Los Angeles, CA*
4. *Department of Pediatrics, Harbor-UCLA Medical Center, Torrance, CA*
5. *Department of Medicine, Harbor-UCLA Medical Center, Torrance, CA*
6. *Center for Pediatric Neurological Disease Research, St. Jude Translational Neuroscience, St. Jude Children's Research Hospital, Memphis, TN, USA*
7. *Department of Neurology, Harbor-UCLA Medical Center, Torrance, CA*

¥Contributed equally

*Corresponding Author

Abstract

Ataxia Telangiectasia (A-T) and ataxia with ocular apraxia type 1 (AOA1) are devastating neurological disorders caused by null mutations in the genome stability genes, A-T mutated (*ATM*) and aprataxin (*APTX*), respectively. Our mechanistic understanding and therapeutic repertoire for treating these disorders is severely lacking, in large part due to the failure of prior animal models with similar null mutations recapitulating the characteristic loss of motor coordination (i.e., ataxia) and associated cerebellar defects. By increasing genotoxic stress, through the insertion of null mutations in both the *Atm* (nonsense) and *Aptx* (knockout) genes in the same animal, we have generated a novel mouse model that for the first time progressively develops a severe ataxic phenotype associated with atrophy of the cerebellar molecular layer. We find biophysical properties of cerebellar Purkinje neurons are significantly perturbed (e.g., reduced membrane capacitance, lower action potential thresholds, etc.), while properties of synaptic inputs remain largely unchanged. These perturbations significantly alter Purkinje neuron

neural activity, including a progressive reduction in spontaneous action potential firing frequency that

correlates with both cerebellar atrophy and ataxia over the animal's first year of life. Double mutant mice

also exhibit a high predisposition to developing cancer (thymomas) and immune abnormalities (impaired

early thymocyte development and T-cell maturation), symptoms characteristic of the disorder A-T. Lastly,

by inserting a clinically relevant nonsense type null mutation in *Atm*, we demonstrate that small molecule

readthrough (SMRT) compounds can restore ATM production, indicating their potential as a future A-T

therapeutic.

1.0 Introduction

Ataxia Telangiectasia (A-T) is a rare (1 in ~100,000) (Swift et al. 1986), autosomal recessive genetic

disorder characterized by cancer predisposition, immune deficiency, and a highly penetrant progressive

and severe ataxia linked to cerebellar atrophy (Rothblum-Oviatt et al. 2016; Boder and Sedgwick 1958;

Levy and Lang 2018). A-T patients typically die in their second and third decade of life (Crawford et al.

2006) from lymphatic cancers, respiratory infections, or debilitating ataxia—unfortunately, survivability

has not dramatically changed since the 1950s (Micol et al. 2011; Rothblum-Oviatt et al. 2016). While

disease progression and cause of death vary widely across patients, the highly penetrant progressive

decline in motor coordination is reported as having the greatest negative impact on a patient's quality of

life (Jackson et al. 2016). Care is generally palliative, directed at reducing, limiting, or eliminating cancers

or infections. No long-term therapies are available for treating the ataxia and associated cerebellar

dysfunction and atrophy.

A-T is caused by deficiency or dysfunction of the ATM (A-T mutated) protein (Savitsky et al. 1995).

Premature termination codon (PTC) causing nonsense mutations account for up to a half of known cases

with missense and deletions also contributing (Concannon and Gatti 1997; Sandoval et al. 1999). ATM

is a serine/threonine PIKK family kinase that is an important regulator of gene stability, in particular, as a

regulator of the DNA damage response (DDR) to the tens of thousands of DNA lesions cells incur each

day (Lindahl and Barnes 2000; Kastan and Bartek 2004; Shiloh and Ziv 2013). In the active monomeric

form, ATM phosphorylates several key proteins halting the production of new DNA (cell cycle arrest)

(Ando et al. 2012), and then, depending on severity of the damage, initiating DNA repair or programmed

cell death (i.e., apoptosis) (Ando et al. 2012; Rashi-Elkeles et al. 2006). Several downstream DDR

pathway targets of ATM have been identified, including p53, CHK2, BRCA1, SMC1, and NBS1 (Matsuoka

et al. 2007). ATM's role in DNA repair is also implicated in normal immune system development, where

it is proposed to contribute to the recombination of natural DNA splicing that occurs during gene

rearrangement in T- and B-lymphocyte maturation (Chao, Yang, and Xu 2000; Matei, Guidos, and

Danska 2006; Vacchio et al. 2007; Schubert, Reichenbach, and Zielen 2002). Although its roles are still

emerging, ATM has also been implicated in oxidative stress homeostasis (Guo et al. 2010) and mitophagy

(Valentin-Vega and Kastan 2012; Pizzamiglio, Focchi, and Antonucci 2020).

A mechanistic understanding of why ATM deficiency causes ataxia is still under debate, but it is far from

the only DDR protein linked to ataxia, as aprataxin (APTX) (Aicardi et al. 1988), meiotic recombination

11 homolog 1 (MRE11) (Sedghi et al. 2018), nibrin (NBS1) (van der Burgt et al. 1996), senataxin (SETX)

(Moreira et al. 2004), and tyrosyl-DNA phosphodiesterase 1 (TDP1) (Takashima et al. 2002) when absent

or dysfunctional can cause cerebellar-related ataxia. This suggests that the neurological features of

genome instability syndromes have a common underlying cause, although this idea is still to be

mechanistically demonstrated (McKinnon 2009; Rass, Ahel, and West 2007).

A major factor limiting our ability to define why loss of DDR proteins, like ATM, selectively impacts the

cerebellum and causes progressive ataxia is the lack of an animal model that recapitulates these

neurological symptoms (Lavin 2013). Several A-T rodent models have been created over the past several

years by inserting gene mutations that cause protein dysfunction (lack kinase activity) or complete

deficiency (Herzog et al. 1998; Xu and Baltimore 1996; Elson et al. 1996; Spring et al. 2001; Campbell

et al. 2015; Quek et al. 2016; Tal et al. 2018; Lavin 2013); a minipig was also recently reported (Beraldi

et al. 2017). Unfortunately, none acquire an overt, progressive ataxia with cerebellar dysfunction and

atrophy that recapitulates the human disease, even though other aspects of the disorder like thyroid

cancers, infertility, and immune abnormalities do develop. It remains unclear why these prior animal

models fail to display the progressive ataxic phenotype (Lavin 2013). It is possible that species specific

molecular compensations in mice provide redundancies or alternative pathways minimizing the effects of

ATM deficiency in the brain (El-Brolosy and Stainier 2017). It is also possible that the shortened lifespan

of prior models (Barlow et al. 1996) is too brief for the stochastic mechanisms driving cerebellar

dysfunction and atrophy to accumulate and impact motor behavior. Other challenges include potentially

leaky genetic manipulations that result in low levels of ATM protein or active fragments with residual

kinase activity, thus limiting neuropathology (Li et al. 2011). The impact of missing such a crucial animal

model has been significant, severely limiting not only experimental studies from identifying the cellular

and molecular mechanisms but hampering pre-clinical development and testing of therapeutics.

We test here whether increasing the genotoxic stress on cells of the murine nervous system through

knocking out expression of ATM and an additional DDR protein leads to a more representative model of

A-T that displays cerebellar dysfunction, atrophy, and the development of progressive ataxia. We chose

to additionally knock-out APTX because its deficiency in humans causes an A-T like disorder called ataxia

with ocular apraxia type 1 (AOA1), but without A-T's other system defects that could increase the potential

for prenatal lethality or early death (e.g., immunodeficiency and cancer predisposition) (Coutinho P 2002).

Moreover, APTX is a phosphodiesterase involved in DNA reassembly after double and single stranded

repair, having a function downstream of, but not directly regulated or related to ATM (Gueven et al. 2004;

Schellenberg, Tumbale, and Williams 2015; Ahel et al. 2006). We therefore reasoned that deficiency of

both proteins would have an additive effect, increasing the amount of genotoxic stress that arises from

either protein alone to induce neurological dysfunction. Our results demonstrated that mice deficient in

ATM and APTX developed cerebellar dysfunction, atrophy, and a progressive and profound ataxia, while

deficiency in either protein alone did not. Additionally, double mutants acquired several additional

characteristic symptoms of A-T, including defects in immune maturation and a high incidence of cancer

(thymomas).

Finally, to improve the clinical and translational relevance of this new mouse model, we inserted a point

mutation (103C>T) in the *Atm* gene common to a large family North African A-T patients. This mutation

results in a premature termination codon (PTC) at what would normally be amino acid 35 resulting in loss

of ATM production. As a result of this genotypic mutation, we were then able to conduct proof-of-principle

experiments demonstrating the ability of so-called PTC readthrough therapeutics to restore production of

ATM protein. This includes the Small Molecule Readthrough (SMRT) compound we are developing (Du

et al. 2013).

113 2.0 Results

114 2.1 Creation of a new A-T mutant mouse model expressing a clinically relevant nonsense mutation

To create a more clinically relevant mouse model of A-T we used a gateway recombination cloning and

site-directed mutagenesis method to recapitulate a c.103C>T (p.R35X) mutation in the *ATM* gene found

in a large population of North African A-T patients (**Fig. 1A and Methods**) (Gilad et al. 1996). The

insertion of thymine in place of cytosine at this site in exon 3 results in a premature termination codon

(PTC) causing nonsense mutation in the ATM gene. Since the c.103C>T mutation results in different

PTCs in the human compared to the mouse *Atm* gene, TGA *vs.* TAG respectively, we created two

different mice by exchanging the mouse *Atm* exon 3 with either a human or mouse exon 3 variant with

the c.103C>T mutation (**Fig. 1B**). In the human variant, a 103C>T mutation of the mouse codon, where

the arginine (R) encoding codon (CGA) becomes a TGA stop codon, results in a mouse we denote as

Atm^{R35X}. In the mouse variant, the c.103C>T mutation transforms a glutamine (Q) encoding CAG codon

into a TAG stop codon and is denoted Atm^{Q35X}. The presence of the PTC results in a loss of ATM

expression, either partially in the heterozygote, or completely in the homozygote (**Fig. 1C**).

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice were created by first crossing single mutant $Atm^{R35X/R35X}$ (congenic on the

C57BL/6J background) and $Aptx^{-/-}$ (mixed C57BL/6J and 129 background) mice to generate double

mutant heterozygote $Atm^{R35X/Wt}$; $Aptx^{-/+}$ mice. F1-5 littermate $Atm^{R35X/Wt}$; $Aptx^{-/+}$ mice were then crossed

within litters to create sufficient numbers of the desired experimental and control genotypes to determine

how loss of different amounts of ATM and APTX affect the animal's phenotype (**Fig. 1D**). Like prior ATM

deficient A-T mouse models, ATM or APTX deficiency alone did not result in mice with ataxia (**Video S1**

and S2). However, deficiency in both proteins ($Atm^{R35X/R35X}$; $Aptx^{-/-}$) results in the development of a severe

and progressively ataxic phenotype (**Fig. 1E**, **Video S3 and S4**).

2.2 ATM deficient mice have lowered survivability and a high incidence of thymomas

We assessed the general health and development of control and experimental mice expressing different

levels of ATM and APTX (**Fig. 2**). We found that $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice grew ~55% slower and reached

estimated plateau weights that were ~35% less than control genotypes (log-rank, n = 21 to 40, p<0.0001;

Fig. 2A). These differences in weight were a postnatal phenomenon, as no significant weight differences

were detected just after birth (P8) across all genotypes (1-way ANOVA, n = 5 to 23, p>0.23). Adolescent

double mutant mice at postnatal day 45 (P45) weighed on average 30% less in males double mutant:

14.4 ± 1.0 g (n = 13) *vs.* wildtype: 20.2 ± 0.5 g (n = 16), *t*-test, p<0.0001] and 25% less in females [double

mutant: 12.7 ± 0.6 g (n = 17) *vs.* wildtype: 17.0 ± 0.2 g (n = 15), *t*-test, p<0.0001; **Fig. 1A**]. Differences

across the control genotypes were observed, but they were small and not consistent across time points

or sex and therefore judged to not be physiologically relevant (**Fig. 2A**). Survivability of the $Atm^{R35X/R35X}$;

$Aptx^{-/-}$ mice was significantly reduced compared to $Atm^{Wt/Wt}$; $Aptx^{+/+}$ mice, with 53% of mice still alive at

400 days of age, compared to 97% of $Atm^{Wt/Wt}$; $Aptx^{+/+}$ mice at the same time point (**Fig. 2B**). ATM

deficiency alone was sufficient to reduce survivability, as compared to $Atm^{Wt/Wt}$; $Aptx^{+/+}$ mice, both

$Atm^{R35X/R35X}$; $Aptx^{+/+}$ and $Atm^{R35X/R35X}$; $Aptx^{-/+}$ mice had significantly reduced survivability rates [42%, log-

rank, $\chi^2_{(1, 56)} = 13.49$, p=0.0002 and 52%, log-rank, $\chi^2_{(1, 53)} = 19.54$, p<0.0001, respectively]. No significant

difference between ATM deficient mice with partial or complete APTX deficiency was detected [log-rank,

$\chi^2_{(2, 85)} = 1.01$, p=0.6]. Conversely, mice harboring at least one functional copy of the *Atm* gene had normal

survivability rates, regardless of whether they expressed APTX or not [log-rank, $\chi^2_{(3, 131)} = 3.08$, p=0.4].

No significant difference between male and female mice was observed and thus data were pooled [log-

rank, p>0.4 for all pairwise comparisons; **Fig. 2-fig. S1B**]. Generally, a third of mice with ATM deficiency

died from complications related to large thymic cancers (thymoma) found in the thoracic cavity (**Fig. 2C**).

The presence or absence of APTX did not impact cancer prevalence, and mice with at least one *Atm*

transcript were cancer free up until at least P400. Overall, ATM, but not APTX deficiency had severe

effects on the health and survivability of mice.

2.3 Both ATM and APTX deficiency are necessary to produce progressive motor dysfunction

The progressive development of severe ataxia is a hallmark characteristic of A-T that is recapitulated in

the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice but none of the other control genotypes we tested. Overall, we find motor

coordination deficits emerge between 210 and 400 days after birth in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice and find

no evidence of ataxia in mice with at least one copy of the Atm or $Aptx$ gene (**Fig. 3A, B**). For the vertical

pole test, $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice took twice as long to descend at P400 compared to $Atm^{Wt/Wt}$; $Aptx^{+/+}$,

$Atm^{Wt/Wt}$; $Aptx^{-/-}$, $Atm^{R35X/R35X}$; $Aptx^{+/+}$, or $Atm^{Wt/R35X}$; $Aptx^{-/-}$ mice [Male: 29.1 ± 0.9 s (n = 3) $vs.$ 7.5 ± 0.4 s

(n = 12), 12.5 ± 2.5 s (n = 9), 9.2 ± 0.9 s (n = 10), 8.6 ± 0.9 s (n = 11), 1-way ANOVA, $F_{(4, 40)} = 19.9$,

p<0.0001; Female: 19.0 ± 4.0 s (n = 4) $vs.$ 7.5 ± 0.4 s (n = 12), 7.8 ± 0.4 s (n = 10), 10.5 ± 1.2 s (n = 6), 8.2

± 0.5 s (n = 8), 1-way ANOVA, $F_{(4, 35)} = 13.9$, p<0.0001]. An examination of gait indicated that $Atm^{R35X/R35X}$;

$Aptx^{-/-}$ mice at P400, but not P210 need additional stabilization during ambulation, as they spend twice

as much time with 3 paws, rather than the normal 2 in contact with the ground as they walk across the

gait analysis platform [Male: 56.2 $vs.$ 26.4 to 32.2 %, 1-way ANOVA, $F_{(4, 54)} = 14.3$, p<0.0001; Female:

58.4 $vs.$ 18.9 to 28.8 %, 1-way ANOVA, $F_{(3, 178)} = 95.5$, p<0.0001; **Fig. 3B**]. $Atm^{R35X/R35X}$; $Aptx^{-/-}$ also display

a slower cadence and average speed across the platform compared to all other genotypes at P400

[cadence, Male: 9.5 $vs.$ 13.3 to 15.9 steps/s, 1-way ANOVA, $F_{(3, 204)} = 36.8$, p<0.0001; Female: 9.1 $vs.$ 14.2

to 15.9 steps/s, 1-way ANOVA, $F_{(3, 204)} = 39.7$, p<0.0001; speed, Male: 8.8 $vs.$ 22 to 26 cm/s, 1-way

ANOVA, $F_{(4, 50)} = 28.3$, p<0.0001; Female: 58.4 $vs.$ 18.9 to 28.8 cm/s, 1-way ANOVA, $F_{(3, 178)} = 39.7$,

p<0.0001; **Fig. 3B; Fig. 3-fig. S1**]. This difference in speed and cadence is unlikely due to animal size,

as there are no significant differences in these parameters at earlier time points when the difference in

size is significant (**Fig. 2A**). These observations across the two behavioral tests were found in both male

and female mice at each of their respective time points, consistent with the lack of sex differences

observed in A-T patients.

We further examined behavioral differences between the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and $Atm^{Wt/Wt}$; $Aptx^{+/+}$ mice

using a standardized set of experimental procedures used to phenotype genetically modified mice (i.e.,

SHIRPA; **Fig. 3C**; **Fig. 3-fig. S1**) (Rogers et al. 1997). We first detected differences in motor function at

P8, where $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice took 3-4 times longer on average to right themselves compared to

$Atm^{Wt/Wt}$; $Aptx^{+/+}$ mice [Male: 6.4 ± 1.1 s (n = 24) *vs.* 1.5 ± 0.1 s (n = 23), *t*-test, p<0.0002; Female: $11.1 \pm$

1.9 s (n = 21) *vs.* 2.4 ± 0.3 s (n = 17), *t*-test, p<0.0002; **Fig. 3C bottom**]. At 30-days of age, we detected

significant differences between $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and $Atm^{Wt/Wt}$; $Aptx^{+/+}$ mice in behavioral tests that

qualitatively measure body position and spontaneous activity (**Fig. 3C**). Striking differences in

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ compared to $Atm^{Wt/Wt}$; $Aptx^{+/+}$ mice were observed at P400, especially for behaviors

related to movement, including locomotor activity, body position, and gait (**Fig. 3C**). The results from this

battery of tests demonstrates that $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice develop a severe change in behavior by P400,

consistent with purely visual observations of significant motor coordination deficits in the mice up to this

time point. Importantly, we do not find any significant differences between the other control genotypes,

including $Atm^{Wt/R35X}$; $Aptx^{-/-}$ mice that express at least some ATM but no APTX protein (**Fig. 3-fig. S1**).

2.4 The membrane and synaptic properties are perturbed in ATM and APTX deficient neurons of

the cerebellum

Purkinje neurons (PN) are a key neuronal subtype located in the cerebellar cortex. They display

considerable intrinsic excitability, firing action potentials spontaneously at rates >1,000 % higher than

other neurons in the brain. Their activity shapes cerebellar output via tonic inhibition of neurons of the

cerebellar nuclei, which project to motor coordination centers in the forebrain, brainstem, and spinal cord.

Dysfunction of cerebellar PNs is associated with several forms of ataxia and implicated in A-T (Hoxha et

al. 2018; Cook, Fields, and Watt 2020; Shiloh 2020). We therefore examined if the electrophysiological

properties of PNs in the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ cerebellum were abnormal.

Since PN baseline activity and responsivity to input is mediated by a baseline set of passive and active

membrane properties (**Fig. 4**), we directly recorded from and compared the membrane properties of PNs

in acute cerebellar slices harvested from $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and $Atm^{Wt/Wt}$; $Aptx^{+/+}$ mice (P350 to 400).

PNs recorded from $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice had significantly "tighter" membranes, displaying higher

membrane input resistances (R_m) than those from $Atm^{Wt/Wt}$; $Aptx^{+/+}$ mice [47.7 ± 5.6 (n = 15) *vs.* $30.2 \pm$

1.47 (n = 23) MΩ, *t*-test, p=0.008; **Fig. 4B**]. They also displayed a faster membrane time constant (τ) [3.6

± 0.4 (n = 15) *vs.* 5.1 ± 0.3 (n = 23) ms, t-test, p=0.009; **Fig. 4B**]. These results indicate that the total

membrane capacitance ($C_m = \tau/R_m$) of the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ PNs is significantly reduced [98.25 ± 19.23

(n = 15) *vs.* 175.6 ± 12.67 (n = 23) pF, *t*-test, p=0.0025; **Fig. 4B**]. At the cellular level, this suggests that

the ATM and APTX deficient PNs have less (i.e., smaller in size) or thinner membranes than that of

wildtype PNs; a result suggestive of a developmental deficit or neurodegenerative process (Dell'Orco et

al. 2015). We next assessed the intrinsic excitability of PNs in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and $Atm^{Wt/Wt}$; $Aptx^{+/+}$

mice by examining PN action potential (AP) generation and dynamics. Significant deficits in the ability of

PNs to fire continuously in response to current injection were observed in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice (**Fig.**

4C). These deficits were associated with significant perturbations to the amplitude, threshold, and area

of evoked action potentials [amplitude: 66.2 ± 0.7 (n = 14) *vs.* 72.1 ± 1.4 (n = 13) ΔmV, *t*-test, p = 0.003;

threshold: -55.2 ± 1.5 *vs.* -48.61 ± 1.9 mV, *t*-test, p=0.0196; area: 17.96 ± 0.6 *vs.* 20.63 ± 1.0 mV*ms, *t*-

test, p=0.048; **Fig. 4C**]. Together, these experiments demonstrate significant perturbations of PN

physiological properties that likely perturb their ability to function normally in the cerebellum of

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice.

We next tested whether extrinsic and/or synaptic PN properties were also impacted in $Atm^{R35X/R35X}$; $Aptx^{-}$

$^{/-}$ mice. We first examined spontaneous mini excitatory post synaptic currents (mEPSC) generated by

granule cell-to-PN synapses (i.e., parallel fiber inputs). No difference in mEPSC size was detected,

indicating the function of granule cell axon terminals (i.e. parallel fibers) was relatively normal in the

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ cerebellum [18.92 ± 1.3 (n = 11) *vs.* 23.4 ± 3.3 (n = 11) pA, *t*-test, p=0.477; **Fig. 4D**]

(Yamasaki, Hashimoto, and Kano 2006). mEPSC frequency however, was found to be significantly

increased, a phenomenon that could be attributed to either an increase in the total number of synapses,

an increase in the size of the readily releasable pool of synaptic vesicles, or an increase in the probability

of neurotransmitter release in PNs of $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice [18.75 ± 2.8 Hz (n = 11) *vs.* 11.4 ± 1.0 Hz

(n = 11), *t*-test, p=0.047; **Fig. 4D**]. We next explored evoked synaptic release and short-term plasticity by

simultaneously recording from PNs and electrically stimulating either granule cell (i.e., parallel fibers) or

inferior olivary (i.e., climbing fiber) axons with a paired-pulse burst (2-pulses, 50 ms apart). The synaptic

properties of parallel fibers were found to be normal, displaying no significant differences in the expected

239 short-term facilitation (Atluri and Regehr 1996) or halfwidth and decay time constant of the evoked EPSC

240 [PPR: 1.3 ± 0.03 (n = 10) *vs.* 1.4 ± 0.05 (n = 13), *t*-test, p=0.162; halfwidth: 3.9 ± 0.6 *vs.* 4.9 ± 0.4 ms, *t*-

241 test, p=0.175; time constant: 3.5 ± 0.5 *vs.* 4.7 ± 0.4 ms, *t*-test, p=0.054; **Fig. 4E**]. In comparison, we found

242 climbing fiber-to-PN synaptic responses, which normally displays pair-pulse depression (Hansel and

243 Linden 2000), to depress at significantly greater magnitudes in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice [PPR: 0.6 ± 0.03

244 (n = 6) *vs.* 0.7 ± 0.02 (n = 9), *t*-test, p=0.03*;* **Fig. 4F**]. The overall width and decay time constant of the

245 evoked currents were also smaller [halfwidth: 2.3 ± 0.6 (n = 6) *vs.* 3.0 ± 0.2 (n = 9) ms, *t*-test, p=0.004;

246 time constant (fast): 1.1 ± 0.14 *vs.* 2.9 ± 0.4 ms, t-test, p=0.001]. While these results could be caused by

247 a presynaptic deficit, such as reduced vesicle stores in the climbing fiber axon terminal, the fact that the

248 initial magnitude of the EPSC is unaffected [23.5 ± 3.6 (n = 6) *vs.* 19.1 ± 1.9 (n = 9), *t*-test, p=0.3], points

249 to a more intrinsic deficit, such as a reduced Ca^{2+} influx from the endoplasmic reticulum, which could

250 significantly impact long term synaptic plasticity critical to cerebellar function (Hoxha et al. 2018; Kano

251 and Watanabe 2017). Overall, the perturbations we observed here to the passive and active PN

252 properties likely give rise to significant cerebellar dysfunction in the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice.

253 **2.5 ATM and APTX deficiency causes a progressive perturbation of PN neural activity that is**

254 **associated with dendritic shrinking and overall cerebellar atrophy.**

255 Decreased rates of spontaneous PN action potential firing, which can be indicative of PN dysfunction,

256 have been observed in several mouse models of ataxia, including SCAs 2, 3, 5, 6, 13, 27, several models

257 of episodic ataxia (e.g., leaner, ducky, and tottering), and autosomal-recessive spastic ataxia of the

258 Charlevoix-Saguenay ((Hourez et al. 2011; Hansen et al. 2013; Dell'Orco, Pulst, and Shakkottai 2017;

259 Kasumu and Bezprozvanny 2012; Liu et al. 2009; Perkins et al. 2010; Shakkottai et al. 2011a; Jayabal

260 et al. 2016; Stoyas et al. 2020; Hurlock, McMahon, and Joho 2008; Shakkottai et al. 2009; Bosch et al.

261 2015; Walter et al. 2006; Alviña and Khodakhah 2010; Ady et al. 2018; Larivière et al. 2019; Cook, Fields,

262 and Watt 2020). We therefore used this biomarker to characterize the progression of PN perturbation in

263 $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice and assess whether deficits were restricted to ATM and APTX deficient mice,

264 consistent with the behavioral results (**Fig. 2,3**). We additionally examined whether decreased PN activity

differed across the cerebellum, as anecdotal clinical pathology reports suggest degeneration may occur

asymmetrically across the cerebellum, with the anterior and posterior vermis and middle cerebellar

hemispheres affected the most, although no systematic analysis has been performed and the consistency

of results across patients is highly variable (Verhagen et al. 2012; De Leon, Grover, and Huff 1976;

Amromin, Boder, and Teplitz 1979; Monaco et al. 1988; Terplan and Krauss 1969; Strich 1966; Solitare

1968; Solitare and Lopez 1967; Aguilar et al. 1968a; Paula-Barbosa et al. 1983).

Using extracellular recording methods in the acute slice, we recorded spontaneous action potentials from

3,300 PNs (**Fig. 5A**), across 188 animals, encompassing $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and 3 other genotypes at 4

different time points (P45, 120, 210, and 400). We visually selected "healthy" cells (see Methods), deeper

in the slice, that consistently fired during the extent of the 60 second recording period. Qualitatively, tissue

and cell quality did not visually differ across genotypes under DIC microscopy. Cells were sampled in a

distributed fashion across the lateral, intermediate and medial (vermis) cerebellum of each mouse to

assess whether changes in PN firing activity was ubiquitous or anatomically restricted. Regions were

segregated based on gross anatomical domains in the mouse defined by natural anatomical boundaries

(e.g., foliation) and their general connectivity with different regions of the nervous system (e.g., forebrain,

brainstem, etc.) (Voogd and Glickstein 1998).

We found that complete deficiency of both ATM and APTX, consistent with the behavioral results, was

necessary to produce a significantly reduced spontaneous PN firing frequency (**Fig. 5A, B**). Although the

trend of slower PN firing rates was observed across most regions of the cerebellum, some subregions

appeared to be less or minimally impacted, including several areas of the lateral cerebellum, including

the paraflocculus, paramedian, and crus I and II (**Fig. 5-fig. S1**). Significant age dependent changes in

firing frequency were also only observed in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice (**Fig. 5B**). The most significant

decline occurring between P120 and 210 [medial: 50.3 ± 2.4 Hz (n = 61) vs. 36.9 ± 2.2 Hz (n = 31), t-test,

p=0.0006]. No significant difference in PN firing frequency was detected between male and female mice

within each genotype, thus the data were pooled (2-way ANOVA, p>0.3 across all pairwise comparisons;

Fig. 5-fig. S2). Previous studies across several mouse models of heritable ataxia, including episodic

ataxia and several variants of spinocerebellar ataxia find that physiological disruption in PN firing not only

changes its frequency, but also its regularity (Kasumu and Bezprozvanny 2012; Jayabal et al. 2016;

Stoyas et al. 2020; Cook, Fields, and Watt 2020). We compared both the coefficient of variation (CV) and

variability in adjacent intervals (CV2) between $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and control mice (**Fig. 5-figs. S3, S4**).

No difference in these parameters across sex, age, or genotype was detected. Consistent with the

behavioral results, cerebellar dysfunction was found only in the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice that developed

ataxia and not in mice with at least some expression of ATM or APTX.

2.5 ATM and APTX deficiency induces cerebellar atrophy

Ataxia in A-T patients is usually detected between 1 to 2-years of age and is associated with mild to no

cerebellar atrophy (Tavani et al. 2003; Taylor et al. 2015). Significant structural changes are usually

apparent via neuroimaging between 5 to 10-years of age (Demaerel, Kendall, and Kingsley 1992; Tavani

et al. 2003). Comprehensive characterization of the progression of cerebellar atrophy is however severely

lacking in large part due challenges in imaging young patients (<2-years) and the rarity of the disorder.

We therefore assessed the gross developmental progression of cerebellar size and pathohistology in

mice deficient in both ATM and APTX that progressively develop a severe ataxia and perturbations in

cerebellar function. Structural changes in the overall size of the cerebellum were examined in

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ and several different control mice over 5 time points (P45, 120, 210, 400, 460; **Fig.**

5C). Cerebellar size was defined within each animal by the ratio of 2-dimensional surface area of the

dorsal cerebellum to the forebrain (i.e., cerebellum area divided by forebrain area). The cerebellar size

of control mice slightly increased during adolescence and early adulthood (P45-P210), was generally

stable through adulthood (P210-400), and then declined slightly in older age (P400-460). In stark

contrast, relative cerebellar size in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice progressively declined after P120. No

cerebellar atrophy was observed in mice with at least one copy of the Atm gene [i.e., $Atm^{Wt/R35X}$; $Aptx^{-/-}$;

1-way ANOVA, $F_{(3,44)}$= 1.2, p=0.32; **Fig. 5C**]. To rule out the possibility that reduced cerebellar size was

related to the smaller stature of $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice, we examined, but did not find a correlation

between animal weight and actual cerebellar size [Pearson's correlation, p>0.3 for all 4 genotypes at

P460, n = 10 to 20]. Furthermore, we found that cerebellar size did not differ between male and the on

average 22% smaller female mice across genotypes at this age [2-way ANOVA, $F_{(2, 153)}$= 1.9, p=0.2].

Therefore, cerebellar neurodegeneration in the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice, which begins after P120, is

correlated with ATM and APTX deficiency.

In humans, cerebellar atrophy is associated with a variety of changes in the macro-structure of the

cerebellar cortex, this includes various reports of abnormalities in the density and distribution of granule

cells (GC) and PNs (Verhagen et al. 2012; De Leon, Grover, and Huff 1976; Amromin, Boder, and Teplitz

1979; Monaco et al. 1988; Terplan and Krauss 1969; Strich 1966; Solitare 1968; Solitare and Lopez

1967; Aguilar et al. 1968a; Paula-Barbosa et al. 1983; Gatti and Vinters 1985). In the $Atm^{R35X/R35X}$; $Aptx^{-/-}$

mice we do not qualitatively observe severe pathological changes in the anatomy of their cerebellum

(**Fig. 5-fig. S5B**). To then understand the anatomical basis of the gross atrophy, we measured the width

of the cortical layer of the cerebellum that contains the massive dendritic arbors of PNs called the

molecular layer (ML). Consistent with the temporal changes in gross cerebellar size, PN firing frequency,

and behavior, ML width in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice was reduced and declined in size from P120 to P400

[P400: 120.2 ± 2.1 μm (n = 5) *vs.* 140.2 ± 4.8 μm (n = 5), Sidak test, p=0.0004; **Fig. 5D**]. In contrast, the

width of the GC layer did not change with age or differ significantly between $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and

$Atm^{Wt/Wt}$; $Aptx^{+/+}$ mice [P400: 135.5 ± 2.4 μm (n = 6) *vs.* 127.5 ± 4.3 μm (n = 5), Sidak test, p=0.5; **Fig. 5D**].

These results, along with the finding that PN density did not significantly differ between $Atm^{R35X/R35X}$; $Aptx^{-/-}$

and $Atm^{Wt/Wt}$; $Aptx^{+/+}$ mice [P400: 3.9 ± 0.3 (n = 4) *vs.* 4.4 ± 0.3 (n = 5) PNs/4000 μm^2, Welch's test, p=0.3;

Fig. 5-fig. S5B] indicate that cerebellar atrophy is in large part due to a reduction in the extent of PN

dendrites. At the anatomical level, we were able to qualitatively observe a few other PN abnormalities.

This includes abnormally large diameter primary dendrites and axonal swellings in the $Atm^{R35X/R35X}$; $Aptx^{-/-}$

mice (**Fig. 5-fig. S5C**). Overall, we find a good correlation between the abnormal structural and

electrophysiological properties and the progression of motor behavioral deficits.

2.6 Differential disruption of thymocyte development in ATM-deficient vs. APTX-deficient mice

Chronic sinopulmonary infections associated with immunodeficiency are one of the leading causes of

death in A-T patients (Morrell, Cromartie, and Swift 1986; Bhatt and Bush 2014). Immunodeficiency is

linked to deficits in the generation of B- and T-lymphocytes that have been linked to defects in the antigen

receptor gene rearrangement processes during the generation of these cells in bone marrow and thymus,

respectively (Staples et al. 2008). The resulting defects in mature lymphocyte numbers include decreases

in CD4$^+$ helper T-cells and killer CD8$^+$ T-cells (Schubert, Reichenbach, and Zielen 2002). We therefore

examined the percentages of T-cells in peripheral blood and of different subpopulations in the thymus of

Atm$^{R35X/R35X}$; Aptx$^{-/-}$ mice using T-cell antigen receptor (TCR) and CD4/CD8 co-receptor expression.

In the peripheral blood, we observed a significant reduction in the total fraction of CD3$^+$ T-cells in mice

with reduced or absent ATM expression compared to wildtype mice (**Fig. 6**). This reduction was further

compounded by concomitant deficiency of APTX. ATM and APTX deficiencies reduced T-cells in

peripheral blood by over 65% decrease compared to wild type controls. The effect of APTX deficiency

was additive to that of ATM deficiency suggesting a different mechanism of action for each of these two

proteins on T-cell generation. The reduction in the percentage of T-cells in peripheral blood was mostly

associated with reduction in the CD4+ helper T-cell population (**Fig. 6B**). Of interest, the proportion of

CD8+ T-cells was increased only in Atm$^{R35X/R35X}$; Aptx$^{-/-}$ mice (**Fig. 6B**). Again, we observed a differential

effect of ATM and APTX deficiencies as seen for the effects of these mutations on the total T-cell fraction.

Given the reduction in T-cell populations in the blood, we next assessed T-cell development in the

thymus. In this organ, bone marrow-derived T-cell progenitors undergo TCR gene rearrangement

followed by positive selection for MHC restriction and negative selection of autoreactive clones. The

phases of thymocyte development can be followed by monitoring expression of CD4 and CD8 expression

in thymocytes. The progression of this developmental program goes from double negative (CD4$^-$CD8$^-$)

thymocytes, to double positive (CD4$^+$CD8$^+$) thymocytes and then to single positive (CD4$^+$ or CD8$^+$)

thymocytes. In addition, within the double negative stage, four different subpopulations can be identified,

based on expression of CD25 and CD44, known as DN1 (CD44$^+$CD25$^-$), DN2 (CD44$^+$CD25$^+$), DN3

(CD25$^+$CD44$^-$) and DN4 (CD44$^-$CD25$^-$) (Germain 2002).

Gene rearrangement during thymocyte development occurs twice, once at the double negative thymocyte

stage in the CD25$^+$CD44$^-$ stage (Krangel 2009) and then again in double positive thymocyte stage before

progressing into separate CD4$^+$ and CD8$^+$ single positive populations (Livák et al. 1999). ATM deficiency

has been linked to defects in both bouts of rearrangement in mice (Vachio 2007, Hathcock 2013).

Therefore, we compared the proportion of cells in the thymus expressing these different developmental

cell surface markers in our ATM deficient and control mice (**Fig. 7**). *Atm$^{R35X/R35X}$; Aptx$^{-/-}$* and *Atm$^{Wt/R35X}$;*

Aptx$^{-/-}$, but not *Atm$^{R35X/R35X}$; Aptx$^{+/+}$* mice had significantly elevated proportions of CD44$^+$CD25$^-$,

CD44$^+$CD25$^+$, and CD44$^-$CD25$^+$ cells compared to wildtype (**Fig. 7A**). These increased proportions

appear to be due in part to an impediment of CD44$^-$CD25$^+$ cells maturing into CD44$^-$CD25$^-$ double

negative cells, as the fraction of cells from *Atm$^{R35X/R35X}$; Aptx$^{-/-}$* and *Atm$^{Wt/R35X}$; Aptx$^{-/-}$* mice is significantly

lower than wildtype (**Fig. 7A**). Of interest, APTX deficiency by itself had the greatest effect on the loss of

DN4 cells suggesting that APTX deficiency, rather than ATM deficiency, is responsible for this effect. To

our knowledge, this finding implicates for the first time APTX in gene rearrangement during the process

of TCRβ recombination.

Next, we looked at the proportions of CD4$^+$CD8$^+$ thymocytes compared to CD4$^+$CD8$^-$ and CD4$^-$CD8$^+$

single positive thymocytes in these four different strains. In agreement with our results in the blood and

prior studies, we found that ATM-deficient mice but not control mice displayed decreased expression of

CD4$^+$CD8$^-$ and CD4$^-$CD8$^+$ single positive thymocytes (**Fig. 7B**). These results support the role of ATM in

TCR α/δ gene rearrangement during thymocyte development (Bredemeyer et al. 2006), a role that is

independent of the role played by APTX in early thymocyte maturation.

2.7 Readthrough molecules overcome PTC to restore ATM expression

Our primary rationale for inserting a clinically relevant nonsense mutation in the *Atm* gene was to

generate a mouse amenable to critical pre-clinical testing of a novel set of small molecule readthrough

(SMRT) compounds. We previously demonstrated SMRT compounds recover production of ATM protein

in A-T patient derived lymphoblastoid cell lines by overcoming premature termination codons (PTC)

caused by nonsense mutations (Du et al. 2013). To demonstrate suitability of this new A-T animal model

394 for SMRT compound testing we chose to directly examine their ability to restore ATM expression using

395 an explant approach that circumvents challenges related to *in vivo* delivery (e.g., bioavailability, route of

396 delivery, etc.). ATM expression was measured in samples from the spleen, where ATM is normally

397 expressed at high levels, and the cerebellum, a key target tissue for the disorder. We exposed these

398 explant tissues, harvested from both ATM^{R35X} and ATM^{Q35X} mice with either a candidate SMRT compound

399 (GJ103), or an aminoglycoside previously known to have readthrough properties (G418) for 72 hrs. and

400 then measured ATM expression by immunoblot to assess restoration. In both types of ATM deficient

401 mice, ATM expression was consistently restored in the spleen and cerebellum by both G418, GJ103 (**Fig.**

402 **8**). These results demonstrate that our SMRT compounds can enable readthrough of at least 2 of the 3

403 possible nonsense mutation causing PTCs and provide the rationale for *in vivo* efficacy testing in follow-

404 on studies.

405 3.0 Discussion

406 By increasing genotoxic stress through the addition of a secondary hit to the DDR pathway, we generated

407 a novel mouse model that displays the most comprehensive set of A-T symptoms of any model to date.

408 This includes a severe and progressive ataxia associated with cerebellar atrophy and perturbations of

409 PN properties along with a high incidence of cancer and defects in immune cell development. Together,

410 these comorbidities encompass the three leading causes of premature death in A-T — each contributing

411 to roughly a third. Of these, the incapacitating effect of ataxia is the most penetrant and is reported by

412 patients and caregivers as having the greatest impact on their quality of life. For this reason, the presence

413 of ataxia and cerebellar atrophy in this new mouse model is of great significance as it provides for the

414 very first time a resource to not only elucidate the mechanisms of neurological dysfunction, but also a

415 critically needed *in vivo* model to test severely needed A-T therapeutics like the readthrough compounds

416 we describe here.

417 We found several similarities between the overall progression of ataxia in the *Atm$^{R35X/R35X}$*; *Aptx$^{-/-}$* mice

418 and A-T patients. In clinical A-T, motor deficits are observable by roughly 2-years of age, when parents

419 and doctors detect a lowered ability to transition from toddling to a smooth, reflexively coordinated gate—

unfortunately, little is known about motor defects at earlier stages due to the diseases low prevalence

and current lack of early diagnostic testing (Rothblum-Oviatt et al. 2016). Patients usually learn to walk

without assistance and neurological symptoms tend to remain stable through the first 4 to 5 years of life

(Rothblum-Oviatt et al. 2016). We found a similar early progression of motor deficits in $Atm^{R35X/R35X}$; $Aptx^{-/-}$

mice, detecting mild motor deficits early at P8 (righting reflex deficit), followed by a period of relative

stability, prior to onset of a progressive and severe ataxia developing after p210 that included changes

in gait, startle reflex, tremor, and locomotor activity. Several important questions arise out of these

findings, including whether ATM and/or APTX have a neurodevelopmental role in the cerebellum. Future

studies focused on the early phase of the disorder will be critical in understanding if the cerebellum

develops normally prior to dysfunction or whether developmental defects are an initial cause. We also

found, similar to A-T patients, that the severity of the late developing ataxia was variable with some mice

ambulating with a clumsy, high stepping rear gate (**Video S3**) and others moving almost entirely via

contortion of the rear trunk (**Fig. 1E** and **Video S4**) (Rothblum-Oviatt et al. 2016; Levy and Lang 2018;

Boder and Sedgwick 1958). Overall, we found that $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice developed a visually profound

and measurable progressive loss in motor coordination that was similar to that observed in A-T patients

and that was rescued by expression of at least one copy of the Atm or $Aptx$ gene.

The loss of motor coordination in A-T has been attributed to cerebellar degeneration due to its relatively

selective neuropathology across the brain and its causal role in several different forms of ataxia (Hoche

et al. 2012). Consistent with A-T patient neuroimaging studies (Wallis et al. 2007; Sahama et al. 2015;

Sahama et al. 2014; Dineen et al. 2020; Tavani et al. 2003; Quarantelli et al. 2013), we find that cerebellar

size in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice is initially normal, but progressively atrophies in line with changes in

neurological function. While loss of cerebellar tissue has been considered a main cause of ataxia in

humans, it is unclear from clinical data if ataxia severity is a good predictor of the extent of cerebellar

degeneration found postmortem (Aguilar et al. 1968b; Crawford et al. 2006; Dineen et al. 2020). In the

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice, we find clear atrophy associated with thinning of the Purkinje neuron dendrite

layer that precedes the late, severe behavioral deficits. However, since the neuropathology observed in

the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice is not severe, our findings suggest that like several SCAs (e.g.,1 AND 3),

changes in cerebellar function itself, rather than profound loss of cerebellar tissue, is likely sufficient to

cause the ataxic phenotype (Shakkottai et al. 2011b; Lorenzetti et al. 2000; Clark et al. 1997).

Why ATM and APTX deficiency is required to generate ataxia in mice, when loss of either is sufficient to

cause ataxia in humans, remains an open question. It is possible that the rodent brain may more flexibly

utilize compensatory pathways or redundant proteins to reduce the impact of impaired DNA repair as it

responds to the 10-20k DNA lesions that assault it per day (Lindahl and Barnes 2000). Several forms of

DNA repair exist to potentially meet this challenge, including base excision repair (BER), nucleotide

excision repair (NER), as well as homologous and non-homologous end joining (HEJ and NHEJ,

respectively), all of which ATM and APTX have been implicated in (Chou et al. 2015; Çaglayan et al.

2017; Wakasugi et al. 2014; Tumbale et al. 2018; Chatterjee and Walker 2017). Alternatively, it may be

the case that deficiency in ATM or APTX alone does not adequately impact cell health during the mouse's

comparatively short lifespan. The use of various DDR pathways depends on the extent and form of DNA

damage (e.g., radiation, genotoxic agents, metabolism, etc.), and in cases like double stranded breaks,

multiple pathways are necessary to fully address the damage. Thus, loss or reduced efficiency of repair

caused by missing DDR pathway proteins could have a graded impact on the severity and/or accrual rate

of deleterious damage sites. This possibility is strengthened by the fact that ATM and APTX have distinct

biochemical properties and functional roles in the DNA damage response, and thus deficiency in both

would be predicted to cause a broader hit to genome stability (i.e., increased genotoxic stress). This may

also account for why we find Purkinje neuron dysfunction and atrophy in 12 to 14-month-old mice, but

not more severe degeneration and cell death as is characteristic of human patients near their second

decade of life (Demaerel, Kendall, and Kingsley 1992; Tavani et al. 2003).

Our finding, that two genome stability pathway proteins are required to induce neurological defects in

mice strongly suggests that it is the loss of ATM's role in DNA repair, rather than potential functions in

oxidative stress signaling, mitophagy, or mitochondrial function that cause the cerebellar defects (Shiloh

2020). Alternatives however, cannot be completely ruled out, as APTX, like ATM has been observed

within the mitochondria of brain cells, where it has been proposed to be important in the processing of

mitochondrial DNA (Meagher and Lightowlers 2014; Sykora et al. 2011). Future experiments, enabled by

this new mouse model, will need to be conducted to mechanistically link ATM and ATPX deficiency

directly to cerebellar dysfunction.

We find several similarities between the biophysical and neuronal response properties of PNs in

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice and other mouse models of ataxia that support a role for PN dysfunction as a

significant contributing factor in the ataxic phenotype. The changes in PN membrane properties we

observed are similar to those found in several SCA mouse models, like 1, 3, and 7 (Stoyas et al. 2020;

Shakkottai et al. 2011b; Dell'Orco et al. 2015). Of note, is the significant reduction in membrane

capacitance that is consistent with the progressive shrinking of the PN dendritic layer we observed (i.e.,

ML). Moreover, changes in the active PN membrane properties were found to impact action potential

generation and properties (e.g., threshold, width, etc.). These intrinsic changes likely underlie the

progressive reduction in PN action potential firing frequency that we find well correlates with the

development of ataxia—a common finding across several other ataxic mouse model studies, including

those for spinocerebellar ataxias (SCA) 1, 2, 3, 5, 6, and 13 as well as a few episodic forms (see review

(Cook, Fields, and Watt 2020)). While it would be tempting to design experiments to simply increase the

spontaneous firing rate of PNs as a solution to all ataxias, it is likely that while this phenomenon serves

as a biomarker for PN defects, it only superficially highlights more distinct neural signaling defects across

the different cerebellar ataxias. Several key questions remain to be answered for not just A-T but

cerebellar ataxias as a whole, including whether disease-causing cerebellar defects primarily cause

ataxia through a loss of cerebellar function (e.g., loss of coordinating signals during movement), or a

dominant negative effect (e.g., disruptive signaling that prevents the normal function of downstream

neural circuits). While there is no doubt that a broad therapeutic strategy to address cerebellar related

ataxias, it is likely that a directed approach, that addressed the genetic and molecular causes of cellular

function will be required to have significant therapeutic success.

While the link between protein deficiency and neuronal dysfunction is straightforward for some types of

ataxia, like the loss of voltage gated ion channels that directly shape membrane excitability in SCA6 (i.e.,

Ca$_{v2.1}$) (Jayabal et al. 2016), the mechanistic link between deficiency in DNA stability proteins like ATM

and APTX and PN dysfunction is far less obvious. Our results suggest the effects of ATM and APTX loss

on PNs is intrinsic, as we don't find changes in the presynaptic properties of granule cells, nor evidence

of cellular loss (no change in GCL thickness). Moreover, while we observed differences in short term

plasticity of inferior olivary inputs in ATM and APTX deficient PNs and wildtype, these results likely point

to a disruption in Ca^{2+} homeostasis potentially via reductions in Inositol 1,4,5-triphosphate receptor 1

(*Itpr1*) expression, similar to those observed in SCAs 1, 2, and 3 mouse models as well as ATM-deficient

mice (Kim et al. 2020; Chen et al. 2008; Liu et al. 2009; Shakkottai et al. 2011b). While this provides a

promising avenue for future examination and comparison, it is as of yet unclear, even for the SCAs,

whether changes in Ca^{2+} homeostasis is the causal factor or just another symptom or even compensatory

response of diseased or disturbed PNs (Dell'Orco et al. 2015).

In the immune system, ATM is implicated in the repair of DNA breaks that naturally occur during gene

rearrangement of antigen receptor genes in B- and T-cell precursors, a phenomenon critical for antigen

receptor (Ig and TCR) diversity of these cells. Our finding that T-cell proportions in the blood are

significantly reduced is consistent with prior studies in humans and A-T knockout mice (Schubert,

Reichenbach, and Zielen 2002; Hathcock et al. 2013; Chao, Yang, and Xu 2000; Barlow et al. 1996).

This reduction of T-cells in the periphery likely correlates with a defect in both cellular and humoral

immunity. Importantly, we found that expression of at least one copy of the ATM gene is enough to restore

CD4+ deficits in the blood indicating that therapies able to restore at least some ATM expression would

have therapeutic efficacy. Although we have not assessed B-cell development in this paper, it is likely

that similar conclusions would apply to that process given their mechanistic similarities (Marshall et al.

2018).

As expected, the reduction of T-cells in peripheral blood correlated with defective thymocyte

development. In the thymus, we found two main defects. One, induced primarily by APTX deficiency,

manifests as a defect in the DN3 to DN4 transition coinciding with early rearrangement of TCR β locus.

The other defect, primarily caused by ATM deficiency, correlates with decreased progression of double

positive CD4$^+$CD8$^+$ to single positive cells, primarily CD4$^+$ thymocytes. While the APTX finding was

surprising, as its deficiency (AOA 1) is not associated with immune deficits, APTX is known to interact

with TCR β gene rearrangement proteins, including XRCC4 (Clements et al. 2004). Future studies aimed

at defining APTX's role in end-joining mechanisms during TCR gene rearrangement will be important,

and the possibility that alternative end-joining mechanisms, like the use of microhomologies account for

the lack of an immune deficit in its absence need further investigation (Bogue et al. 1997).

The survivability of $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice is considerably longer than prior A-T mouse models. In

comparison, the first A-T KO mouse model reported by Barlow et al. died from thymomas usually within

2-4 months after birth (Barlow et al. 1996). The increased severity of cancer survivability in this, and many

other knockout A-T mouse models is likely genetic, as the background strain harboring the mutation has

been shown to have significant effects on cancer prevalence and survivability, with A/J and C57BL/6

backgrounds having significantly increased survivability over the BALBC and 129S strains (Genik et al.

2014). The fact that our ATM deficient mice were created on a C57BL/6 background likely underlies their

comparatively long lifespan. Given that the $Atm^{R35X/R35X}$; $Aptx^{+/+}$ mice do not develop ataxia, it is unlikely

that the early death in A-T KO mice prevents observation of an ataxic phenotype that would otherwise

develop in these mice. On the other hand, it is unknown whether the C57BL/6 background confers a

resilience to developing ataxia, as it does for cancer. Defining the genetic or possibly epigenetic factors

that influence the severity of the disease could provide avenues for future therapeutic development.

Given the global nature of the ATM and ATPX knockout in our mouse model, we cannot entirely rule out

that extra-cerebellar defects may also contribute to the severe ataxic phenotype, and thus future

examination outside the cerebellum, in both the forebrain, brainstem, spinal cord, and even muscle will

need to be conducted. Within the cerebellum, while we found some anatomical differences in the PN

firing properties within different regions of the cerebellum, we didn't detect anatomical differences in ML

width or PN density. In general, there are significant challenges in using anatomy as a grouping factor in

the cerebellum, as the physical folds of the tissue do not necessarily correlate well with different

functional, molecular expression, or biophysical property domains that have been described (Apps and

Hawkes 2009; Tsutsumi et al. 2015; Gao, van Beugen, and De Zeeuw 2012; Zhou et al. 2014).

Experiments focused on examining the extent of cerebellar defects within these domains will be important

in future studies and compared to the anecdotal reports of anatomical differences in A-T patients

(Verhagen et al. 2012; De Leon, Grover, and Huff 1976; Amromin, Boder, and Teplitz 1979; Monaco et

al. 1988; Terplan and Krauss 1969; Strich 1966; Solitare 1968; Solitare and Lopez 1967; Aguilar et al.

1968a; Paula-Barbosa et al. 1983).

While we detect two potential stages in the progression of ataxia in the $Atm^{R35X/R35X}$; $Aptx^{+/+}$ mice, the

later stage of severe ataxia develops in adulthood in mice, compared to the childhood onset in humans.

This may limit its use in some neurodevelopmental based studies. Also, the interpretation of future

experiments must carefully factor in the fact that this new model expresses null mutations in two genome

stability genes at the same time, a situation that has not been detected in human patients with either A-

T or AOA1.

Finally, pinpointing where, when, and how ATM deficiency causes cerebellar pathology and ataxia has

been a challenge as prior ATM deficient mice generally lack the characteristic features needed to causally

link cellular and molecular deficits to the ataxic phenotype. Multiple promising avenues of investigation

have been defined, including those focused at the neuronal level where ATM is implicated in oxidative

stress signaling (Chen et al. 2003) and synaptic function (Li et al. 2009; Vail et al. 2016), as well as glial

function, where recent evidence suggests glial pathology may be a leading factor in cerebellar pathology

(Kaminsky et al. 2016; Campbell et al. 2016; Petersen, Rimkus, and Wassarman 2012; Weyemi et al.

2015). This novel animal model provides a new tool to test mechanistic hypotheses regarding how ATM

deficiency causes cerebellar pathology and ataxia. Additionally, this model may serve most importantly

as a critical preclinical tool for testing both previously proposed therapeutic candidates (Browne et al.

2004; Chen et al. 2003) and our own SMRT compounds (Du et al. 2013). The severe limitations of not

having a suitable preclinical model for therapeutic testing, especially for a rare disorder like A-T and

AOA1, cannot be overstated.

4.0 Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Strain, strain background (*Mus musculus*)	*Atm^{R35X}*	This paper	103C>T mutation, human exon replacement	Generated by Hicks laboratory. Has been backcrossed into C57b/6 9 times. Contact pmathews@lundquist.org
Strain, strain background (*Mus musculus*)	*Atm^{Q35X}*	This paper	103C>T mutation, targeted premature termination signal in the mouse codon	Generated by Hicks laboratory. Has been backcrossed into C57b/6 9 times. Contact pmathews@lundquist.org
Strain, strain background (*Mus musculus*)	*Aptx^{-/-}*	Ahel et al. 2006	Aptx^{tm1Pmc}; MGI:3686875	Contact peter.mckinnon@stjude.org
gene (*Mus musculus*)	*Atm*	MGI	MGI:107202; C030026E19Rik; ENSMUSG00000034218	
gene (*Homo Sapien*)	*ATM*	OMIM	OMIM: 607585 MGI: 107202 HomoloGene: 30952; ENSG00000149311	

sequence-based reagent	Atm gene	Transnetyx	PCR primers	F-5'-CCTTTGAGGCATAAGTTGCAACTTG-3'
sequence-based reagent	Atm gene	Transnetyx	PCR primers	R- 5'-GTACAGTGTATCAGGTTAGGCATGC-3'
Chemical compound/ drugs	GJ103 Formula: c16H14N403S	TargetMol	T3448	Treatment
antibody	anti-ATM (D2E2) (Rabbit-monoclonal)	Cell signalling	Cat#: mAb-2873	IF(1:1000), WB(1:500) WB(1:1000)
antibody	anti-GAPDH (14C10) (Rabbit-monoclonal)	Cell signalling	Cat#: mAb 2118	WB(1:4000)
antibody	Anti β-Actin (D6A8) (Rabbit-monoclonal)	Cell signalling	Cat#: mAb 8457	WB(1:5000)
antibody	Anti-rabbit IgG,HRP-linked Antibody (monoclonal-polyclonal)	Cell signalling	Cat#:7074	WB(1:5000)
antibody	Calbindin D-28k	Swant	Cat# CB-38a	IF (1:1000)
antibody	Goat Anti-Rabbit488	ThermoFisher Invitrogen	Cat# A11034	IF(1:1000)
antibody	Fluoromount-G with DAPI	Southern Biotech	Cat# 0100-20	

antibody	CD4	ThermoFisher Invitrogen	Cat# 50-0041-82	FACS (5 ul per test)
antibody	CD8	ThermoFisher Invitrogen	Cat# 53-0081-82)	FACS (5 ul per test)
antibody	CD3	ThermoFisher Invitrogen	Cat#12-0031-83	FACS (5 ul per test)
antibody	CD44	ThermoFisher Invitrogen	Cat# 25-0441-82	FACS (5 ul per test)
antibody	CD25	ThermoFisher Invitrogen	Cat# 47-0251- 82	FACS (5 ul per test)
commercial assay or kit	BCA Protein Assay Kit	ThermoFisher Pierce	Cat# 23225	Protein Assay
commercial assay or kit	SuperSignal West Pico Chemiluminescent Substrate	ThermoFisher Pierce	Cat# 34580	Chemiluminescent Substrate
commercial assay or kit	Radiance plus	Azure Biosystems	Cat# AC2103	Chemiluminescent Substrate
Software	FlowJo	https://www.flowjo.com/solutions/flowjo	RRID:SCR_008520	
Software	ImageJ software	ImageJ (http://imagej.nih.gov/ij/)	RRID:SCR_003070	
Software	IgorPro	http://www.wavemetrics.com/products/igorpro/igorpro.htm	RRID:SCR_000325	Version 7; Tarotools procedures
Software	Neuroexpress	https://www.researchgate.net/project/NeuroExpress-	https://www.researchgate.net/project/NeuroExpress-	Version 21.1.13; Used for

		Analysis-software-for-whole-cell-electrophysiological-data	Analysis-software-for-whole-cell-electrophysiological-data	mEPSC analyses
Software	GraphPad, Prism	GraphPad Prism (https://graphpad.com)	RRID:SCR_015807	Versions 8 and 9
Software	MBF, Stereo investigator	https://www.mbfbioscience.com/stereology	RRID:SCR_017667	Version 2021
Software	Microsoft Excel	https://www.microsoft.com/en-us/microsoft-365/excel	Mircosoft 365	
Software	Catwalk XT	https://www.noldus.com/applications/rodent-gait-analysis	N/A	Catwalk XT v10.6

4.1 Ethics Statement

This study was performed in strict accordance with the recommendations in the Guide for the Care and

Use of Laboratory Animals of the National Institutes of Health. All the animals were handled according to

approved institutional animal care and use committee (IACUC) protocols at The Lundquist Institute

(31374-03, 31773-02) and UCLA (ARC-2007-082, ARC-2013-068). The protocol was approved by the

Committee on the Ethics of Animal Experiments of the Lundquist Institute (Assurance Number: D16-

00213). Every effort was made to minimize pain and suffering by providing support when necessary and

choosing ethical endpoints.

4.2 Mice

All mice were group housed and kept under a 12-h day/night cycle with food and water available *ad libitum*.

Animals were housed within the general mouse house population, and not in specialized pathogen free

rooms. Older animals were made available wetted food or food gel packs on the ground of the cages as

ataxia developed. *Atm^R35X* and *Atm^Q35X* mice were created and provided by Dr. Hicks and colleagues at

the University of Manitoba.

These mice were created to contain the c.103C>T mutation found in a large population of North African

AT patients, using recombineering Gateway technology and site-directed mutagenesis. A C>T mutation

at this position in the mouse *Atm* gene creates a TAG G stop codon. The same mutation in the human

ATM gene produces a TGA G stop codon. In consideration of the use of these models for therapeutic

interventions, we chose to create a mouse model for each of the two PTC codons (**Fig. 1A**).

A modified Gateway R3-R4-destination vector was used to pull out the desired region of the mouse Atm

gene from a Bacterial Artificial Chromosome (BAC) and subsequently mutated to create either a TAG G

stop codon at codon 35 (M00001, position 103 (C>T)) or a TGA G stop codon (M00002, position 103

(CAG>TGA), replicating the human AT PTC). The genomic alleles were then cloned into a modified

version of the NorCOMM mammalian targeting vector using a 3-way Gateway Reaction (Bradley et al.

2012). The resulting targeting vectors were electroporated into C2 ES cells (C57Bl/6N, derived in A. Nagy

lab, Toronto, Canada) and successfully targeted clones were identified by selection with G418

(Gertsenstein et al. 2010). Integration of the mutated targeting cassette into the *Atm* gene locus was

confirmed by Southern blot, and by sequencing of PCR products to confirm the presence of the Atm PTC

mutation, error free targeting into the Atm locus and error free functional components of the vector (data

not shown). Positive ES clones were used for blastocyst injection to obtain the transgenic lines. The

transgenic allele contained a floxed human beta actin promoter - delta TK1- Neo cassette in the intron

upstream of the region containing the mutated exon. This floxed cassette was subsequently excised by

crossing with a Cre driver mouse (B6.C-Tg(CMV-cre)1Cgn/J) to generate *Atm^R35X/Wt* and *Atm^Q35X/Wt* (MGI

nomenclature: Atm^TM1(103CAG>TGA)MFGC and Atm^TM1(103C>T)MFGC, respectively) mouse lines (**Fig. 1A**).

Genotyping of the two *Atm* lines was performed by using the following primers at Tm 62°C: Atm gene

forward (F) primer: 5'-CCTTTGAGGCATAAGTTGCAACTTG-3'; and Atm gene reverse (R) primer: 5'-

GTACAGTGTATCAGGTTAGGCATGC-3', creating a Wild-type allele product of 151bp or targeted allele

product of 241bp (**Figs. 1A, 1B**).

Atm^{R35X} and Atm^{Q35X} were backcrossed with C57Bl/6J mice for 9 generations (99.2% isogenic) prior to

cryopreservation and subsequent rederivation using C57Bl/6J surrogate mothers. Atm^{R35X} and Atm^{Q35X}

breeders were obtained from F1 sibling $Atm^{R35X/Wt}$ and $Atm^{Q35X/Wt}$ mice. $Atm^{R35X/R35X}$ and $Atm^{Q35X/Q35X}$ were

both found to be fertile. Aptx knockout ($Aptx^{-/-}$) mice were created and provided to Dr. Mathews as

embryos from Dr. McKinnon (Ahel et al. 2006), and subsequently rederived via C57Bl/6J surrogate

mothers. $Aptx^{-/-}$ mice are on a C57Bl/6 and 129 mixed background. $Atm^{R35/R35XX}$; $Aptx^{-/-}$ mice of various

Wt, heterozygous, and homozygous combinations were created from $Atm^{R35X/Wt}$; $Aptx^{-/+}$ breeders

generated by crossing $Atm^{R35X/R35X}$ and $Aptx^{-/-}$ mice. One cohort of double mutant and corresponding

control mice were used in the longitudinal behavioral study for gait analyses and SHIRPA testing (**Figs.**

2, **3**). Multiple additional cohorts of age matched double mutant and control mice were used for

electrophysiological, immunohistological, and Vertical Pole test experiments (**Figs. 4**, **7**). Immunological

and protein expression experiments were carried out using mice bred from the original Atm^{R35X} and

Atm^{Q35X} rederived mice (**Figs**. **5**, **6**, and **8**).

Genotyping was performed from ear tissue samples of P8-11 mice. Real-time PCR methods conducted

by Transnetyx Inc. were used to determine each animals' genotype. Animals were made identifiable via

toe tattoos given at the same time as ear biopsy. Unique primers for Atm^{R35X} and Atm^{Q35X} were quantified

and used to identify Wt, hetero- and homo-zygous mice (listed above). $Aptx^{-/-}$ and $Aptx^{Wt/Wt}$ primers were

used to assess their genotypes.

4.3 Animal Health

Animals were weighed via a digital scale at P8, 45, 120, 210, 400. Animal death was recorded as the day

found dead, or on the day of euthanization when the animals reached a humane endpoint (animal unable

to right itself within 60s, significant hair matting indicating lack of self-grooming, or excessive distress as

noted by the veterinary staff). Animal carcasses were immediately frozen upon death, and postmortem

necropsies were carried out in batch. Probable cause of death was determined to the best of our ability

in collaboration with the staff veterinarian (Dr. Catalina Guerra) by visual inspection of the internal organs.

Some mice were cannibalized or accidentally disposed of by vivarium staff and were therefore labelled

as "missing." Mice with no discernable visual cause of death were labelled "indeterminable." Mice that

were found with thoracic masses near where the thymus would normally be in young mice were listed as

"thymic cancer." All other identified probable causes of death (e.g., enlarged livers, urinary blockage)

were labelled "other."

4.4 Behavior

Before performing any behavioral test, mice were acclimated to the behavioral suite for ~20-minutes.

Mice were tested at varying times of the day, in line with their day cycle. A battery of behavioral tests was

performed on naïve double mutant mice of the indicated genotypes at various time points depending on

the behavior but in the same cohort of mice. The battery of tests included Catwalk Gait assessment (P45,

120, 210, 400) and a subset of the SmithKline-Beecham Harwell Imperial-College and Royal-London-

Hospital Phenotype Assessment (SHIRPA) tests (P30 and 400). These tests were conducted by the

UCLA Behavioral Core. Double mutant and control mice were additionally examined on the Vertical Pole

test. All behavioral apparatus was wiped down with ethanol (70%) between each testing each subject.

Gait Analysis

We used a Noldus Catwalk Gait analysis system designed to semi-automatically measure and analyze

the gait of mice during normal ambulation. Briefly, the movement of mice across a glass bottom corridor

is video recorded from a ventral position. Paw prints are highlighted in the video due to light illumination

across the glass walking platform. Each mouse step within a video is subsequently detected using

Catwalk XT (Noldus) in a semi-automated fashion. A run for each mouse consists of 3 trials of consistent

ambulation across the monitored platform. Only consistent trials are accepted, and mice may take up to

10 attempts to complete 3 compliant trials in either direction across the corridor. Compliant trials were

defined as those with movement across the platform under 5 s-long and with no more than 60% speed

variation. Once placed onto the platform, mice generally ran back and forth without any need for

experimenter prompting.

Vertical Pole

Mice are placed at the top of an 80 cm tall bolt with their nose faced down and hind paws as close to the

top as possible. Mice are immediately released, and time started immediately upon placement. Time is

stopped when the first forepaw touches the surface below the pole. A mouse's natural predilection is to

immediately climb down the pole, and they are given up to 60 s to traverse the pole, otherwise they are

helped off the pole. A non-completed trial is automatically given a time of 30 s, as 95% of mice that did

not descend within 30 s were still on the pole at the 60 s mark.

<u>SHIRPA</u>

Behavioral tests were conducted by the University of California, Los Angeles Behavioral Core at P30 and

P400. All parameters are scored to provide a quantitative assessment, which enables comparison of

results both over time and between different laboratories. Each mouse was sequentially tested across all

behaviors within ~20-min time span before moving onto the next mouse. The experimenter was blinded

to animal genotype. The screen was performed as described previously (Rogers et al. 1997).

Behavioral Observation

The primary screen provides a behavioral observation profile and assessment of each animal begins by

observing undisturbed behavior in a viewing jar (10 cm diameter) for 5-min In addition to the scored

behaviors of **body position**, **spontaneous activity**, **respiration rate**, and **tremor**, the observer logs any

instances of bizarre or stereotyped behavior and convulsions, compulsive licking, self-destructive biting,

retropulsion (walking backwards) and indications of spatial disorientation.

Arena Behavior

Thereafter, the mouse is transferred to the arena ($30 \text{ cm} \times 50 \text{ cm}$) for testing of transfer arousal and

observation of normal behavior. The arena is marked into a grid of 10 cm^2 squares to measure locomotor

activity within a 30 s-period. While the mouse is active in the arena, measures of **startle response**, **gait**,

pelvic elevation, and **tail elevation** are recorded.

Supine Restraint

The animal is restrained in a supine position to record autonomic behaviors. During this assessment,

grip strength, **body tone**, **pinna reflex**, **corneal reflex**, **toe pinch**, **wire maneuver**, and **heart rate**,

were evaluated.

Balance and Orientation

Finally, several measures of vestibular system function were performed. The **righting reflex**, **contact**

righting reflex, and **negative geotaxis** tests were performed. Throughout this procedure vocalization,

urination and general fear, irritability, or aggression were recorded.

Equipment Used

1. Clear Plexiglas arena (approximate internal dimensions $55 \times 33 \times 18$ cm). On the floor of the arena

is a Plexiglas sheet marked with 15 squares (11 cm). A rigid horizontal wire (3 mm diameter) is

secured across the rear right corner such that the animals cannot touch the sides during the wire

maneuver. A grid (40×20 cm) with 12 mm mesh (approximate) is secured across the width of the

box for measuring tail suspension and grip strength behavior.

2. A clear Plexiglas cylinder (15×11 cm) was used as a viewing jar.

3. One grid floor (40×20 cm) with 12 mm meshes on which viewing jars stand.

4. Four cylindrical stainless-steel supports (3 cm high x 2.5 cm diameter) to raise grids off the bench.

5. One square (13 cm) stainless steel plate for transfer of animals to the arena.

6. Cut lengths of 3 / 0 Mersilk held in the forceps for corneal and pinna reflex tests

7. A plastic dowel rod sharpened to a pencil point to test salivation and biting.

8. A pair of dissecting equipment forceps, curved with fine points (125 mm forceps, Philip Harris

Scientific, Cat. No. D46-174), for the toe pinch.

9. A stopwatch.

10. An IHR Click box is used for testing the startle responses. The Click Box generates a brief 20 KHz

tone at 90dB SPL when held 30cm above the mouse. Contact Prof. K.P. Steel, MRC Institute of

Hearing Research, University Park, Nottingham NG7 2RD.

11. A ruler.

12. A 30 cm clear Plexiglas tube with an internal diameter of 2.5 cm for the contact righting reflex.

4.5 Electrophysiology

<u>Preparation of acute cerebellar slices</u>

Acute parasagittal slices of 300 μm thickness were prepared from the cerebellum of experimental and

control littermate mice by following published methods (Hansen et al., 2013). In brief, cerebella were

quickly removed and immersed in an ice-cold extracellular solution with composition of (mM): 119 NaCl,

26 NaHCO$_3$, 11 glucose, 2.5 KCl, 2.5 CaCl$_2$, 1.3 MgCl$_2$ and 1 NaH$_2$PO$_4$, pH 7.4 when gassed with 5%

CO$_2$/95% O$_2$. Cerebella were sectioned parasagittally using a vibratome (Leica VT-1000, Leica

Biosystems, Nussloch, Germany) and initially incubated at 35°C for ~30 min, and then equilibrated and

stored at room temperature until use.

<u>Extracellular Electrophysiology</u>

Extracellular and intracellular recordings were obtained from Purkinje neurons (PNs) in slices constantly

perfused with carbogen-bubbled extracellular solution and maintained at either 37° C (extracellular) or

32° C (intracellular) ± 1° C (see above). Cells were visualized with DIC optics and a water-immersion 40×

objective (NA 0.75) using a Zeiss Examiner microscope. Glass pipettes of ~3 MΩ resistance (Model P-

1000, Sutter instruments, Novato, CA) were filled with either extracellular solution and positioned near

PN axon hillocks in order to measure action potential-associated capacitive current transients in voltage

clamp mode with the pipette potential held at 0 mV. For whole-cell patch clamp recordings, pipettes were

filled with an intracellular solution (mM): 140 KMeth (CH3KO3S), 10 NaCl, 2 MgCl$_2$, 0.2 CaCl$_2$, 10 HEPES,

14 Phosphocreatine (tris salt), 1 EGTA, 4 Mg-ATP, 0.4 Na-GTP. 100 μM Picrotoxin (Sigma) was added

to block inhibitory GABAegeric synaptic inputs. Data was acquired using a MultiClamp 700B amplifier at

20 or 100 kHz in voltage or current clamp mode, Digidata 1440 with pClamp10 (Molecular Devices,

Sunnyvale, CA) and filtered at 2 to 4 kHz. The series resistance was usually between 10 and 15 MΩ.

Series resistance was compensated at 80% for short term plasticity experiments only.

For extracellular recordings, a total of 20 to 45 PNs were recorded from for each animal across all

genotypes, sexes, and age groups. Recordings were distributed across both the medial-lateral and

rostro-caudal axis of the cerebellum. Only cells with a "healthy" look (low contrast of cellular borders),

and regular, uninterrupted firing rate were examined. During analysis, a few cells were found to have

gaps in firing of greater than 2 second and these cells were eliminated from analysis, as this type of firing

is associated with being "unhealthy." Double mutant tissue did not qualitatively differ in appearance under

DIC microscopy prior to recordings, nor was the number of "unhealthy" cells greater than that of other

genotypes (7% vs 4 to 11% of all cells across control genotypes at P400). Spatial comparison of neural

activity was obtained by recording from serial sections in the flocculus, lateral (2nd or 3rd), intermediate

(6th or 7th), and medial (11th or 12th) slices. Lower number slices were used in the younger age groups

(P45 and 110) to roughly match the relative positioning of recordings across age groups. 0-3 recordings

were made from each lobule within each slice dependent on tissue quality and health. Each recording

lasted for 1-minute. 3 to 5 mice were used for each age group and the experimenter was blinded to the

genotype, age, and sex.

Intracellular recordings were obtained from PNs in either folia III or VIII of the vermis, no statistical

differences in properties were observed between folia.

<u>Analyses</u>

Spontaneous action potential interstimulus intervals detected and analyzed using standard and custom

routines in ClampFit (Molecular Device), IgorPro (Wavemetrics), and Excel (Microsoft). Specifically,

action potentials were threshold detected and spiking statistics (i.e., frequency and interval length),

determined using adapted IgorPro routines (Taro Tools; https://sites.google.com/site/tarotoolsregister/).

The coefficient of variation of the mean inter-spike interval (CV) and the median inter-spike interval

(CV2=2 |ISIn+1-ISIn|/(ISIn+1+ISIn)) were calculated in Excel using custom macros.

Standard membrane properties were analyzed using IgorPro. R_M was determined by averaging 3 voltage

trace responses to a -5 mV step pulse from a -80 mV holding potential and measuring the resulting

current deflection between 900 and 1000 ms after onset. The membrane time constant was measured

by fitting a single exponential to the initial decay phase from 90 to 10% of the peak. C_M was calculated

by dividing the membrane time constant by the R_M. mEPSC events were recorded over a 1-minute epoch

and detected and measured using Neuroexpress (v21.1.13). Parallel and climbing fiber axons were

stimulated using theta-glass electrodes (W.P.I.) and a TTL controlled stimulus isolator (ISO-Flex,

A.M.P.I.). Evoked EPSC amplitudes and decay time constants (1 exp. for parallel and 2 exp. for climbing

fibers) were analyzed using custom routines in IgorPro. Action potentials were examined as part of a set

of 1 s current injections between -500 and 2250 pA (250 pA steps) with a holding current adjusted to

maintain an ~70 mV potential. Action potential waveforms were measured using custom routines in

IgorPro. Action potential threshold was defined as the first membrane voltage in which the first derivative

exceeded 30 mV/ms (Zhu et al. 2006).

4.6 Examination of Cerebellar Atrophy

Cerebellar size

Immediately after brain removal from the skull, a dorsal, whole mount image was obtained. Images were

then processed using Fiji (NIH). The forebrain and cerebellar sizes were assessed by outlining their 2-

dimensional space and then calculating area. We normalized for possible differences in overall brain size

by dividing the results of the cerebellum by forebrain size to produce a relative cerebellum to forebrain

ratio. Experimenters were blind to the genotype of the animal.

Immunohistochemistry

At the respective study endpoints (P45, 120, 210, 400), male and female mice of all genotypes

represented in this study were anesthetized with isoflurane and underwent transcardial perfusion with

phosphate-buffered saline followed by 4% (w/v) buffered paraformaldehyde (PFA) and then dissected

to extract the brain. Images of the whole brain were taken immediately after removing the brain from

the skull and the brains were then submerged in 4% PFA for 24-hours, followed by 72-hours in 30%

sucrose in Tris-buffered saline (TBS) with 0.05% azide, and then cryoprotected in TBS-AF and stored

at 4°C until further use. The cerebellum was separated from the forebrain and parasagittally sectioned

using a sliding microtome (Microm HM 430, Thermo Scientific) set to section at 40 μm thickness.

Cerebellum sections were collected in a series of six and stored in TBS-AF at 4° C until further use. For

immunofluorescent visualization of Purkinje neurons, cerebellum sections of both $Atm^{Wt/Wt}$; $Aptx^{+/+}$ and

Atm^R35X/R35X; *Aptx^-/-* (n = 5 per genotype) were washed for $5 \min$ in TBS three times, and then blocked in

15% normal goat serum at room temperature for $30 \min$ followed by free floating incubation in rabbit

anti-calbindin D-28k $(1:1000)$ for 1 hour at room temperature on an orbital shaker, then washed for 5

$\min$ with TBS three times, followed by free floating incubation in goat anti-rabbit Alexa Fluor 488

$(1:1000)$ for $1 \mathrm{h}$ in the dark at room temperature on an orbital shaker. Following secondary antibody

incubation, sections were washed for $5 \min$ in TBS three times and stored in TBS until further use.

Sections were mounted and cover-slipped with Fluoromount-G with DAPI. Slides were scanned using

Stereo Investigator version 2020.1.3 64bit (MBF Bioscience) on a Zeiss Axio Imager.M2 microscope

(Carl Zeiss Microscopy) using a $20\mathrm{x}$ objective (NA 0.5) and images captured with a Hamamatsu ORCA

Flash 4.0 LT C11440 digital camera (Hamamatsu Photonics). To quantify the number of calbindin-

reactive cells in each folia in the resulting images, we randomly drew 2 lines between $300 \text{ to } 500 \mu m$

long in each folia and manually counted the total number of PNs along the length within the $40 \mu m$

thickness of the tissue slice. 2D density $(\# \text{ of PNs}/(\text{linear length} * 40 \text{ um thickness}))$ of the two samples

per folia were then averaged for further between folia and subject comparison. Experimenter was

blinded to mouse genotype.

Molecular (Calbindin) and granule cell (DAPI) layer widths were assessed by taking two width

measurements at two predefined locations for each folia, roughly halfway along the long extent of each

folia. Experimenter was blinded to the mouse genotype in which slices were examined and the order of

examination was interleaved.

4.7 Flow Cytometry Measurements

Flow cytometry analysis of blood and thymus cells was performed by staining with specific anti-mouse

antibodies: CD4, CD8, CD3, CD44, and CD25. Briefly, whole-blood samples $(50 \mu l)$ were stained using

fluorescent-labeled antibodies, then red-blood cells were lysed using BD lysing solution while live white-

blood cells were stained using a viability stain. Thymi were mechanically dissociated. $1 \text{ to } 2$ million

thymus cells were similarly stained using specific antibodies for CD4, CD8, CD44 and CD25. Analysis of

immuno-stained white blood cells or thymus samples was performed using FACS ARIA III and data

analyzed using FlowJo software as reported previously (Sanghez et al. 2017).

4.8 Western Blots

Protein extracts (cells/tissues) were homogenized in radioimmunoprecipitation assay (RIPA) lysis buffer

(150 mM NaCl, 1% Nonidet P-40 [NP-40], 0.5% deoxycholate, 0.1% SDS, 50 mM Tris, pH 8.0) with

protease inhibitors (10 ug/ml AEBSF, 10 ug/ml leupeptin, 5 ug/ml pepstatin, 5 ug/ml chymotrypsin, 10

ug/ml aprotinin). The protein extracts were sonicated then pelleted by centrifugation at 13,000 rpm for 15

min at 4°C. BCA protein assay was used to quantify protein concentrations. Samples containing equal

amounts of protein 50 to 100 μg per lane were separated using 4 to 12% gradient TGX precast gels

BioRad then transferred by TransBlot Semi-Dry BioRad system using Nitrocellulose transfer pack.

Transferred blots were stained by Ponceau S stain for equal protein loading then washed and blocked

with 5% nonfat dry milk in TBST for 60 min at room temp. Primary antibodies were incubated with shaking

overnight at 4°C. Blots were probed for the following antibodies: ATM (D2E2) Rabbit mAb Cell Signaling,

at 1:1000 dilution, β-Actin (D6A8) Rabbit mAb Cell Signaling, GAPDH (D16H11) Rabbit mAb Cell

Signaling followed by the appropriate horseradish peroxidase–conjugated (HRP) secondary Anti-rabbit,

Anti-mouse for 2 hours at room temperature. After multiple washes with TBST, Protein expression was

detected by Radiance Plus chemiluminescence substrate using the Azure c400 and the BioRad

ChemiDoc imaging systems. Densitometric analysis of the ATM was performed using ImageJ.

Experiments were performed with 2 technical and 2-3 biological replicates as indicated.

4.9 Statistical Assessment

The number of animals chosen for each group was based on a priori power analyses using GPower v3.1

based on an α size of 0.5, power of 0.8, and effect sizes estimated from preliminary data or prior studies.

We used both parametric (1- and 2-way ANOVA) for normally distributed and non-parametric (Kruskal

Wallace) statistical methods for interval data to test for differences between groups followed by pairwise

multiple comparisons tests as indicated in the text. Outliers for immune data in Figs. 6 and 7 were

excluded via the ROUT method (Q=2%). The specific analyses used for each data set is noted in each

figure legend. For all figures: * p≤0.05, ** p<0.01, *** p<0.001, **** p<0.0001. Data are reported as

$\mathrm{mean} \pm \mathrm{SEM}$ and box and whisker plots indicate the minimum, first quartile, median, third quartile, and

maximum data values. All figures and statistical analyses were completed using Excel (Microsoft 360) or

Prism v8 and 9 (Graphpad).

850 5.0 Acknowledgements

851 We would like to thank the UCLA Behavioral Core, especially Irina Zhuravka for her efforts assaying

852 behavioral deficits in the mice. We would also like to thank Dr. Jennifer Fogel for comments and edits to

853 the manuscript.

854 6.0 Competing interests

855 The authors declare that no competing interests exist.

856 7.0 Citations

Ady, Visou, Brenda Toscano-Márquez, Moushumi Nath, Philip K Chang, Jeanette Hui, Anna Cook,
François Charron, Roxanne Larivière, Bernard Brais, and R Anne McKinney. 2018. 'Altered
synaptic and firing properties of cerebellar Purkinje cells in a mouse model of ARSACS', *The*
Journal of physiology, 596: 4253-67.
Aguilar, M. J., S. Kamoshita, B. H. Landing, E. Boder, and R. P. Sedgwick. 1968a. 'Pathological
observations in ataxia-telangiectasia. A report of five cases', *Journal of Neuropathology and*
Experimental Neurology, 27: 659-76.
Aguilar, Mary Jane, Shigehiko Kamoshita, Benjamin H. Landing, Elena Boder, and Robert P. Sedgwick.
1968b. 'Pathological Observations in Ataxia-Telangiectasia: A Report on Five Cases*', *Journal of*
Neuropathology and Experimental Neurology, 27: 659-76.
Ahel, I., U. Rass, S. F. El-Khamisy, S. Katyal, P. M. Clements, P. J. McKinnon, K. W. Caldecott, and S.
C. West. 2006. 'The neurodegenerative disease protein aprataxin resolves abortive DNA ligation
intermediates', *Nature*, 443: 713-6.
Aicardi, J., C. Barbosa, E. Andermann, F. Andermann, R. Morcos, Q. Ghanem, Y. Fukuyama, Y. Awaya,
and P. Moe. 1988. 'Ataxia-ocular motor apraxia: a syndrome mimicking ataxia-telangiectasia',
Annals of Neurology, 24: 497-502.
Alviña, Karina, and Kamran Khodakhah. 2010. 'KCa channels as therapeutic targets in episodic ataxia
type-2', *Journal of Neuroscience*, 30: 7249-57.
Amromin, George D, Elena Boder, and Raymond Teplitz. 1979. 'Ataxia-telangiectasia with a 32 year
survival: a clinicopathological report', *Journal of Neuropathology and Experimental Neurology*, 38:
621-43.

Ando, K., J. L. Kernan, P. H. Liu, T. Sanda, E. Logette, J. Tschopp, A. T. Look, J. Wang, L. Bouchier-
Hayes, and S. Sidi. 2012. 'PIDD death-domain phosphorylation by ATM controls prodeath versus
prosurvival PIDDosome signaling', *Molecular Cell*, 47: 681-93.
Apps, R., and R. Hawkes. 2009. 'Cerebellar cortical organization: a one-map hypothesis', *Nature
Reviews: Neuroscience*, 10: 670-81.
Atluri, Pradeep P., and Wade G. Regehr. 1996. 'Determinants of the Time Course of Facilitation at the
Granule Cell to Purkinje Cell Synapse', *The Journal of Neuroscience*, 16: 5661-71.
Barlow, C., S. Hirotsune, R. Paylor, M. Liyanage, M. Eckhaus, F. Collins, Y. Shiloh, J. N. Crawley, T.
Ried, D. Tagle, and A. Wynshaw-Boris. 1996. 'Atm-deficient mice: a paradigm of ataxia
telangiectasia', *Cell*, 86: 159-71.
Beraldi, R., D. K. Meyerholz, A. Savinov, A. D. Kovacs, J. M. Weimer, J. A. Dykstra, R. D. Geraets, and
D. A. Pearce. 2017. 'Genetic ataxia telangiectasia porcine model phenocopies the multisystemic
features of the human disease', *Biochim Biophys Acta Mol Basis Dis*, 1863: 2862-70.
Bhatt, Jayesh M., and Andrew Bush. 2014. 'Microbiological surveillance in lung disease in ataxia
telangiectasia', *European Respiratory Journal*, 43: 1797-801.
Boder, Elena, and Robert P. Sedgwick. 1958. 'ATAXIA-TELANGIECTASIA', *A Familial Syndrome of
Progressive Cerebellar Ataxia, Oculocutaneous Telangiectasia and Frequent Pulmonary
Infection*, 21: 526-54.
Bogue, M. A., C. Wang, C. Zhu, and D. B. Roth. 1997. 'V(D)J recombination in Ku86-deficient mice:
distinct effects on coding, signal, and hybrid joint formation', *Immunity*, 7: 37-47.
Bosch, Marie K, Yarimar Carrasquillo, Joseph L Ransdell, Ajay Kanakamedala, David M Ornitz, and
Jeanne M Nerbonne. 2015. 'Intracellular FGF14 (iFGF14) is required for spontaneous and evoked
firing in cerebellar Purkinje neurons and for motor coordination and balance', *Journal of
Neuroscience*, 35: 6752-69.
Bradley, A., K. Anastassiadis, A. Ayadi, J. F. Battey, C. Bell, M. C. Birling, J. Bottomley, S. D. Brown, A.
Bürger, C. J. Bult, W. Bushell, F. S. Collins, C. Desaintes, B. Doe, A. Economides, J. T. Eppig, R.
H. Finnell, C. Fletcher, M. Fray, D. Frendewey, R. H. Friedel, F. G. Grosveld, J. Hansen, Y.
Hérault, G. Hicks, A. Hörlein, R. Houghton, M. Hrabé de Angelis, D. Huylebroeck, V. Iyer, P. J.
de Jong, J. A. Kadin, C. Kaloff, K. Kennedy, M. Koutsourakis, K. C. Lloyd, S. Marschall, J. Mason,
C. McKerlie, M. P. McLeod, H. von Melchner, M. Moore, A. O. Mujica, A. Nagy, M. Nefedov, L.
M. Nutter, G. Pavlovic, J. L. Peterson, J. Pollock, R. Ramirez-Solis, D. E. Rancourt, M. Raspa, J.
E. Remacle, M. Ringwald, B. Rosen, N. Rosenthal, J. Rossant, P. Ruiz Noppinger, E. Ryder, J.
Z. Schick, F. Schnütgen, P. Schofield, C. Seisenberger, M. Selloum, E. M. Simpson, W. C.
Skarnes, D. Smedley, W. L. Stanford, A. F. Stewart, K. Stone, K. Swan, H. Tadepally, L. Teboul,
G. P. Tocchini-Valentini, D. Valenzuela, A. P. West, K. Yamamura, Y. Yoshinaga, and W. Wurst.
2012. 'The mammalian gene function resource: the International Knockout Mouse Consortium',
Mammalian Genome, 23: 580-6.
Bredemeyer, Andrea L, Girdhar G Sharma, Ching-Yu Huang, Beth A Helmink, Laura M Walker, Katrina
C Khor, Beth Nuskey, Kathleen E Sullivan, Tej K Pandita, and Craig H Bassing. 2006. 'ATM
stabilizes DNA double-strand-break complexes during V (D) J recombination', *Nature*, 442: 466-
70.
Browne, Susan E., L. Jackson Roberts, Phyllis A. Dennery, Susan R. Doctrow, M. Flint Beal, Carrolee
Barlow, and Rodney L. Levine. 2004. 'Treatment with a catalytic antioxidant corrects the
neurobehavioral defect in ataxia–telangiectasia mice', *Free Radical Biology and Medicine*, 36:
938-42.
Çaglayan, Melike, Rajendra Prasad, Rachel Krasich, Matthew J. Longley, Kei Kadoda, Masataka Tsuda,
Hiroyuki Sasanuma, Shunichi Takeda, Keizo Tano, William C. Copeland, and Samuel H. Wilson.
2017. 'Complementation of aprataxin deficiency by base excision repair enzymes in mitochondrial
extracts', *Nucleic Acids Research*, 45: 10079-88.
Campbell, A., B. Krupp, J. Bushman, M. Noble, C. Pröschel, and M. Mayer-Pröschel. 2015. 'A novel
mouse model for ataxia-telangiectasia with a N-terminal mutation displays a behavioral defect
and a low incidence of lymphoma but no increased oxidative burden', *Human Molecular Genetics*,
24: 6331-49.

Campbell, Andrew, Jared Bushman, Joshua Munger, Mark Noble, Christoph Pröschel, and Margot
932    Mayer-Pröschel. 2016. 'Mutation of ataxia–telangiectasia mutated is associated with
933    dysfunctional glutathione homeostasis in cerebellar astroglia', *Glia*, 64: 227-39.
934 Chao, Connie, Eva Marie Yang, and Yang Xu. 2000. 'Rescue of Defective T Cell Development and
935    Function in Atm−/− Mice by a Functional TCRαβ Transgene', *The Journal of*
936    *Immunology*, 164: 345-49.
937 Chatterjee, Nimrat, and Graham C. Walker. 2017. 'Mechanisms of DNA damage, repair, and
938    mutagenesis', *Environmental and Molecular Mutagenesis*, 58: 235-63.
939 Chen, Philip, Cheng Peng, John Luff, Kevin Spring, Dianne Watters, Steven Bottle, Shigeki Furuya, and
940    Martin F. Lavin. 2003. 'Oxidative Stress Is Responsible for Deficient Survival and Dendritogenesis
941    in Purkinje Neurons from Ataxia-Telangiectasia Mutated Mutant Mice', *The Journal of*
942    *Neuroscience*, 23: 11453-60.
943 Chen, Xi, Tie-Shan Tang, Huiping Tu, Omar Nelson, Mark Pook, Robert Hammer, Nobuyuki Nukina, and
944    Ilya Bezprozvanny. 2008. 'Deranged calcium signaling and neurodegeneration in spinocerebellar
945    ataxia type 3', *Journal of Neuroscience*, 28: 12713-24.
946 Chou, W. C., L. Y. Hu, C. N. Hsiung, and C. Y. Shen. 2015. 'Initiation of the ATM-Chk2 DNA damage
947    response through the base excision repair pathway', *Carcinogenesis*, 36: 832-40.
948 Clark, H Brent, Eric N Burright, Wael S Yunis, Seth Larson, Claire Wilcox, Boyd Hartman, Antoni Matilla,
949    Huda Y Zoghbi, and Harry T Orr. 1997. 'Purkinje Cell Expression of a Mutant Allele of SCA1in
950    Transgenic Mice Leads to Disparate Effects on Motor Behaviors, Followed by a Progressive
951    Cerebellar Dysfunction and Histological Alterations', *Journal of Neuroscience*, 17: 7385-95.
952 Clements, P. M., C. Breslin, E. D. Deeks, P. J. Byrd, L. Ju, P. Bieganowski, C. Brenner, M. C. Moreira,
953    A. M. Taylor, and K. W. Caldecott. 2004. 'The ataxia-oculomotor apraxia 1 gene product has a
954    role distinct from ATM and interacts with the DNA strand break repair proteins XRCC1 and
955    XRCC4', *DNA Repair (Amst)*, 3: 1493-502.
956 Concannon, P., and R. A. Gatti. 1997. 'Diversity of ATM gene mutations detected in patients with ataxia-
957    telangiectasia', *Human Mutation*, 10: 100-7.
958 Cook, Anna A., Eviatar Fields, and Alanna J. Watt. 2020. 'Losing the Beat: Contribution of Purkinje Cell
959    Firing Dysfunction to Disease, and Its Reversal', *Neuroscience*.
960 Coutinho P, Barbot C, Coutinho P. 2002. 'Ataxia with Oculomotor Apraxia Type 1.' in Ardinger HH Adam
961    MP, Pagon RA, et al. (ed.), *GeneReviews®* (GeneReviews® [Internet]: University of Washington,
962    Seattle).
963 Crawford, T. O., R. L. Skolasky, R. Fernandez, K. J. Rosquist, and H. M. Lederman. 2006. 'Survival
964    probability in ataxia telangiectasia', *Archives of Disease in Childhood*, 91: 610-11.
965 De Leon, G. A., W. D. Grover, and D. S. Huff. 1976. 'Neuropathologic changes in ataxia-telangiectasia',
966    *Neurology*, 26: 947-51.
967 Dell'Orco, James M., Aaron H. Wasserman, Ravi Chopra, Melissa A. C. Ingram, Yuan-Shih Hu, Vikrant
968    Singh, Heike Wulff, Puneet Opal, Harry T. Orr, and Vikram G. Shakkottai. 2015. 'Neuronal Atrophy
969    Early in Degenerative Ataxia Is a Compensatory Mechanism to Regulate Membrane Excitability',
970    *The Journal of Neuroscience*, 35: 11292-307.
971 Dell'Orco, James M, Stefan M Pulst, and Vikram G Shakkottai. 2017. 'Potassium channel dysfunction
972    underlies Purkinje neuron spiking abnormalities in spinocerebellar ataxia type 2', *Human*
973    *Molecular Genetics*, 26: 3935-45.
974 Demaerel, PH, BE Kendall, and D Kingsley. 1992. 'Cranial CT and MRI in diseases with DNA repair
975    defects', *Neuroradiology*, 34: 117-21.
976 Dineen, Rob A., Felix Raschke, Hannah L. McGlashan, Stefan Pszczolkowski, Lorna Hack, Andrew D.
977    Cooper, Manish Prasad, Gabriel Chow, William P. Whitehouse, and Dorothee P. Auer. 2020.
978    'Multiparametric cerebellar imaging and clinical phenotype in childhood ataxia telangiectasia',
979    *NeuroImage: Clinical*, 25: 102110.
980 Du, L., M. E. Jung, R. Damoiseaux, G. Completo, F. Fike, J. M. Ku, S. Nahas, C. Piao, H. Hu, and R. A.
981    Gatti. 2013. 'A new series of small molecular weight compounds induce read through of all three
982    types of nonsense mutations in the ATM gene', *Molecular Therapy*, 21: 1653-60.

El-Brolosy, Mohamed A., and Didier Y. R. Stainier. 2017. 'Genetic compensation: A phenomenon in
search of mechanisms', *Plos Genetics*, 13: e1006780-e80.
Elson, A., Y. Wang, C. J. Daugherty, C. C. Morton, F. Zhou, J. Campos-Torres, and P. Leder. 1996.
'Pleiotropic defects in ataxia-telangiectasia protein-deficient mice', *Proceedings of the National
Academy of Sciences of the United States of America*, 93: 13084-9.
Gao, Zhenyu, Boeke J van Beugen, and Chris I De Zeeuw. 2012. 'Distributed synergistic plasticity and
cerebellar learning', *Nature Reviews: Neuroscience*, 13: 619-35.
Gatti, R. A., and H. V. Vinters. 1985. 'Cerebellar pathology in ataxia-telangiectasia: the significance of
basket cells', *Kroc Foundation Series*, 19: 225-32.
Genik, P. C., H. Bielefeldt-Ohmann, X. Liu, M. D. Story, L. Ding, J. M. Bush, C. M. Fallgren, and M. M.
Weil. 2014. 'Strain background determines lymphoma incidence in Atm knockout mice',
Neoplasia, 16: 129-36.
Germain, Ronald N. 2002. 'T-cell development and the CD4–CD8 lineage decision', *Nature Reviews
Immunology*, 2: 309-22.
Gertsenstein, Marina, Lauryl M. J. Nutter, Tammy Reid, Monica Pereira, William L. Stanford, Janet
Rossant, and Andras Nagy. 2010. 'Efficient Generation of Germ Line Transmitting Chimeras from
C57BL/6N ES Cells by Aggregation with Outbred Host Embryos', *PloS One*, 5: e11260.
Gilad, Shlomit, Rami Khosravi, Dganit Shkedy, Tamar Uziel, Yael Ziv, Kinneret Savitsky, Galit Rotman,
Sara Smith, Luciana Chessa, Timothy J. Jorgensen, Reli Harnik, Moshe Frydman, Ozden Sanal,
Sima Portnoi, Zipora Goldwicz, N. G. J. Jaspers, Richard A. Gatti, Gilbert Lenoir, Martin F. Lavin,
Kouichi Tatsumi, Rolf D. Wegner, Yosef Shiloh, and Anat Bar-Shira. 1996. 'Predominance of Null
Mutations in Ataxia-Telangiectasia', *Human Molecular Genetics*, 5: 433-39.
Gueven, N., O. J. Becherel, A. W. Kijas, P. Chen, O. Howe, J. H. Rudolph, R. Gatti, H. Date, O. Onodera,
G. Taucher-Scholz, and M. F. Lavin. 2004. 'Aprataxin, a novel protein that protects against
genotoxic stress', *Human Molecular Genetics*, 13: 1081-93.
Guo, Zhi, Sergei Kozlov, Martin F. Lavin, Maria D. Person, and Tanya T. Paull. 2010. 'ATM Activation by
Oxidative Stress', *Science*, 330: 517-21.
Hansel, Christian, and David J. Linden. 2000. 'Long-Term Depression of the Cerebellar Climbing Fiber–
Purkinje Neuron Synapse', *Neuron*, 26: 473-82.
Hansen, Stephen T, Pratap Meera, Thomas S Otis, and Stefan M Pulst. 2013. 'Changes in Purkinje cell
firing and gene expression precede behavioral pathology in a mouse model of SCA2', *Human
Molecular Genetics*, 22: 271-83.
Hathcock, K. S., S. Bowen, F. Livak, and R. J. Hodes. 2013. 'ATM influences the efficiency of TCRbeta
rearrangement, subsequent TCRbeta-dependent T cell development, and generation of the pre-
selection TCRbeta CDR3 repertoire', *PloS One*, 8: e62188.
Herzog, Karl-Heinz, Miriam J. Chong, Manuela Kapsetaki, James I. Morgan, and Peter J. McKinnon.
1998. 'Requirement for Atm in Ionizing Radiation-Induced Cell Death in the Developing Central
Nervous System', *Science*, 280: 1089-91.
Hoche, F., K. Seidel, M. Theis, S. Vlaho, R. Schubert, S. Zielen, and M. Kieslich. 2012.
'Neurodegeneration in ataxia telangiectasia: what is new? What is evident?', *Neuropediatrics*, 43:
119-29.
Hourez, Raphael, Laurent Servais, David Orduz, David Gall, Isabelle Millard, Alban de Kerchove
d'Exaerde, Guy Cheron, Harry T Orr, Massimo Pandolfo, and Serge N Schiffmann. 2011.
'Aminopyridines correct early dysfunction and delay neurodegeneration in a mouse model of
spinocerebellar ataxia type 1', *Journal of Neuroscience*, 31: 11795-807.
Hoxha, Eriola, Ilaria Balbo, Maria Concetta Miniaci, and Filippo Tempia. 2018. 'Purkinje Cell Signaling
Deficits in Animal Models of Ataxia', *Frontiers in Synaptic Neuroscience*, 10: 6-6.
Hurlock, Edward C, Anne McMahon, and Rolf H Joho. 2008. 'Purkinje-cell-restricted restoration of Kv3.
3 function restores complex spikes and rescues motor coordination in Kcnc3 mutants', *Journal of
Neuroscience*, 28: 4640-48.
Jackson, Thomas J, Gabriel Chow, Mohnish Suri, Philip Byrd, Malcolm R Taylor, and William P
Whitehouse. 2016. 'Longitudinal analysis of the neurological features of ataxia-telangiectasia',
Developmental Medicine and Child Neurology, 58: 690-97.

Jayabal, Sriram, Hui Ho Vanessa Chang, Kathleen E Cullen, and Alanna J Watt. 2016. '4-aminopyridine
reverses ataxia and cerebellar firing deficiency in a mouse model of spinocerebellar ataxia type
6', *Scientific Reports*, 6: 1-12.
Kaminsky, Natalie, Ofer Bihari, Sivan Kanner, and Ari Barzilai. 2016. 'Connecting Malfunctioning Glial
Cells and Brain Degenerative Disorders', *Genomics, Proteomics & Bioinformatics*, 14: 155-65.
Kano, Masanobu, and Takaki Watanabe. 2017. 'Type-1 metabotropic glutamate receptor signaling in
cerebellar Purkinje cells in health and disease', *F1000Research*, 6: 416-16.
Kastan, M. B., and J. Bartek. 2004. 'Cell-cycle checkpoints and cancer', *Nature*, 432: 316-23.
Kasumu, Adebimpe, and Ilya Bezprozvanny. 2012. 'Deranged calcium signaling in Purkinje cells and
pathogenesis in spinocerebellar ataxia 2 (SCA2) and other ataxias', *The Cerebellum*, 11: 630-39.
Kim, Jusik, Keeeun Kim, Jung-Soon Mo, and Youngsoo Lee. 2020. 'Atm deficiency in the DNA
polymerase β null cerebellum results in cerebellar ataxia and Itpr1 reduction associated with
alteration of cytosine methylation', *Nucleic Acids Research*, 48: 3678-91.
Krangel, Michael S. 2009. 'Mechanics of T cell receptor gene rearrangement', *Current Opinion in
Immunology*, 21: 133-39.
Larivière, Roxanne, Nicolas Sgarioto, Brenda Toscano Márquez, Rébecca Gaudet, Karine Choquet, R
Anne McKinney, Alanna J Watt, and Bernard Brais. 2019. 'Sacs R272C missense homozygous
mice develop an ataxia phenotype', *Molecular Brain*, 12: 1-15.
Lavin, M. F. 2013. 'The appropriateness of the mouse model for ataxia-telangiectasia: neurological
defects but no neurodegeneration', *DNA Repair (Amst)*, 12: 612-9.
Levy, Ariel, and Anthony E. Lang. 2018. 'Ataxia-telangiectasia: A review of movement disorders, clinical
features, and genotype correlations', *Movement Disorders*, 33: 1238-47.
Li, J., Y. R. Han, M. R. Plummer, and K. Herrup. 2009. 'Cytoplasmic ATM in neurons modulates synaptic
function', *Current Biology*, 19: 2091-6.
Li, Jiali, Jianmin Chen, Harry V. Vinters, Richard A. Gatti, and Karl Herrup. 2011. 'Stable Brain
ATM Message and Residual Kinase-Active ATM Protein in Ataxia-Telangiectasia',
The Journal of Neuroscience, 31: 7568-77.
Lindahl, T., and D. E. Barnes. 2000. 'Repair of endogenous DNA damage', *Cold Spring Harbor Symposia
on Quantitative Biology*, 65: 127-33.
Liu, Jing, Tie-Shan Tang, Huiping Tu, Omar Nelson, Emily Herndon, Duong P Huynh, Stefan M Pulst,
and Ilya Bezprozvanny. 2009. 'Deranged calcium signaling and neurodegeneration in
spinocerebellar ataxia type 2', *Journal of Neuroscience*, 29: 9148-62.
Livák, Ferenc, Michelle Tourigny, David G. Schatz, and Howard T. Petrie. 1999. 'Characterization of TCR
Gene Rearrangements During Adult Murine T Cell Development', *The Journal of Immunology*,
162: 2575-80.
Lorenzetti, Diego, Kei Watase, Bisong Xu, Martin M Matzuk, Harry T Orr, and Huda Y Zoghbi. 2000.
'Repeat instability and motor incoordination in mice with a targeted expanded CAG repeat in the
Sca1 locus', *Human Molecular Genetics*, 9: 779-85.
Marshall, Jean S., Richard Warrington, Wade Watson, and Harold L. Kim. 2018. 'An introduction to
immunology and immunopathology', *Allergy, asthma, and clinical immunology : official journal of
the Canadian Society of Allergy and Clinical Immunology*, 14: 49-49.
Matei, I. R., C. J. Guidos, and J. S. Danska. 2006. 'ATM-dependent DNA damage surveillance in T-cell
development and leukemogenesis: the DSB connection', *Immunological Reviews*, 209: 142-58.
Matsuoka, Shuhei, Bryan A. Ballif, Agata Smogorzewska, E. Robert McDonald, Kristen E. Hurov, Ji Luo,
Corey E. Bakalarski, Zhenming Zhao, Nicole Solimini, Yaniv Lerenthal, Yosef Shiloh, Steven P.
Gygi, and Stephen J. Elledge. 2007. 'ATM and ATR Substrate Analysis Reveals Extensive Protein
Networks Responsive to DNA Damage', *Science*, 316: 1160-66.
McKinnon, P. J. 2009. 'DNA repair deficiency and neurological disease', *Nature Reviews: Neuroscience*,
10: 100-12.
Meagher, M., and R. N. Lightowlers. 2014. 'The role of TDP1 and APTX in mitochondrial DNA repair',
Biochimie, 100: 121-4.
Micol, R., L. Ben Slama, F. Suarez, L. Le Mignot, J. Beauté, N. Mahlaoui, C. Dubois d'Enghien, A. Laugé,
J. Hall, J. Couturier, L. Vallée, B. Delobel, F. Rivier, K. Nguyen, T. Billette de Villemeur, J. L.

Stephan, P. Bordigoni, Y. Bertrand, N. Aladjidi, J. M. Pedespan, C. Thomas, I. Pellier, M. Koenig, O. Hermine, C. Picard, D. Moshous, B. Neven, F. Lanternier, S. Blanche, M. Tardieu, M. Debré, A. Fischer, and D. Stoppa-Lyonnet. 2011. 'Morbidity and mortality from ataxia-telangiectasia are associated with ATM genotype', *Journal of Allergy and Clinical Immunology*, 128: 382-9.e1.

Monaco, S., E. Nardelli, G. Moretto, T. Cavallaro, and N. Rizzuto. 1988. 'Cytoskeletal pathology in ataxia-telangiectasia', *Clinical Neuropathology*, 7: 44-6.

Moreira, M. C., S. Klur, M. Watanabe, A. H. Németh, I. Le Ber, J. C. Moniz, C. Tranchant, P. Aubourg, M. Tazir, L. Schöls, M. Pandolfo, J. B. Schulz, J. Pouget, P. Calvas, M. Shizuka-Ikeda, M. Shoji, M. Tanaka, L. Izatt, C. E. Shaw, A. M'Zahem, E. Dunne, P. Bomont, T. Benhassine, N. Bouslam, G. Stevanin, A. Brice, J. Guimarães, P. Mendonça, C. Barbot, P. Coutinho, J. Sequeiros, A. Dürr, J. M. Warter, and M. Koenig. 2004. 'Senataxin, the ortholog of a yeast RNA helicase, is mutant in ataxia-ocular apraxia 2', *Nature Genetics*, 36: 225-7.

Morrell, Daphne, Elizabeth Cromartie, and Michael Swift. 1986. 'Mortality and cancer incidence in 263 patients with ataxia-telangiectasia', *Journal of the National Cancer Institute*, 77: 89-92.

Paula-Barbosa, M. M., C. Ruela, M. A. Tavares, C. Pontes, A. Saraiva, and C. Cruz. 1983. 'Cerebellar cortex ultrastructure in ataxia-telangiectasia', *Annals of Neurology*, 13: 297-302.

Perkins, Emma M, Yvonne L Clarkson, Nancy Sabatier, David M Longhurst, Christopher P Millward, Jennifer Jack, Junko Toraiwa, Mitsunori Watanabe, Jeffrey D Rothstein, and Alastair R Lyndon. 2010. 'Loss of β-III spectrin leads to Purkinje cell dysfunction recapitulating the behavior and neuropathology of spinocerebellar ataxia type 5 in humans', *Journal of Neuroscience*, 30: 4857-67.

Petersen, Andrew J., Stacey A. Rimkus, and David A. Wassarman. 2012. 'ATM kinase inhibition in glial cells activates the innate immune response and causes neurodegeneration in Drosophila', *Proceedings of the National Academy of Sciences*, 109: E656-E64.

Pizzamiglio, L., E. Focchi, and F. Antonucci. 2020. 'ATM Protein Kinase: Old and New Implications in Neuronal Pathways and Brain Circuitry', *Cells*, 9.

Quarantelli, Mario, Giuliana Giardino, Anna Prinster, Giuseppina Aloj, Barbara Carotenuto, Emilia Cirillo, Angela Marsili, Elena Salvatore, Ennio Del Giudice, and Claudio Pignata. 2013. 'Steroid treatment in Ataxia-Telangiectasia induces alterations of functional magnetic resonance imaging during prono-supination task', *European Journal of Paediatric Neurology*, 17: 135-40.

Quek, Hazel, John Luff, KaGeen Cheung, Sergei Kozlov, Magtouf Gatei, C. Soon Lee, Mark C. Bellingham, Peter G. Noakes, Yi Chieh Lim, Nigel L. Barnett, Steven Dingwall, Ernst Wolvetang, Tomoji Mashimo, Tara L. Roberts, and Martin F. Lavin. 2016. 'A rat model of ataxia-telangiectasia: evidence for a neurodegenerative phenotype', *Human Molecular Genetics*, 26: 109-23.

Rashi-Elkeles, S., R. Elkon, N. Weizman, C. Linhart, N. Amariglio, G. Sternberg, G. Rechavi, A. Barzilai, R. Shamir, and Y. Shiloh. 2006. 'Parallel induction of ATM-dependent pro- and antiapoptotic signals in response to ionizing radiation in murine lymphoid tissue', *Oncogene*, 25: 1584-92.

Rass, Ulrich, Ivan Ahel, and Stephen C. West. 2007. 'Defective DNA Repair and Neurodegenerative Disease', *Cell*, 130: 991-1004.

Rogers, D. C., E. M. Fisher, S. D. Brown, J. Peters, A. J. Hunter, and J. E. Martin. 1997. 'Behavioral and functional analysis of mouse phenotype: SHIRPA, a proposed protocol for comprehensive phenotype assessment', *Mammalian Genome*, 8: 711-3.

Rothblum-Oviatt, Cynthia, Jennifer Wright, Maureen A. Lefton-Greif, Sharon A. McGrath-Morrow, Thomas O. Crawford, and Howard M. Lederman. 2016. 'Ataxia telangiectasia: a review', *Orphanet Journal of Rare Diseases*, 11: 159.

Sahama, Ishani, Kate Sinclair, Simona Fiori, James Doecke, Kerstin Pannek, Lee Reid, Martin Lavin, and Stephen Rose. 2015. 'Motor pathway degeneration in young ataxia telangiectasia patients: A diffusion tractography study', *NeuroImage: Clinical*, 9: 206-15.

Sahama, Ishani, Kate Sinclair, Kerstin Pannek, Martin Lavin, and Stephen Rose. 2014. 'Radiological imaging in ataxia telangiectasia: a review', *The Cerebellum*, 13: 521-30.

Sandoval, N., M. Platzer, A. Rosenthal, T. Dork, R. Bendix, B. Skawran, M. Stuhrmann, R. D. Wegner, K. Sperling, S. Banin, Y. Shiloh, A. Baumer, U. Bernthaler, H. Sennefelder, M. Brohm, B. H.

Weber, and D. Schindler. 1999. 'Characterization of ATM gene mutations in 66 ataxia
telangiectasia families', *Human Molecular Genetics*, 8: 69-79.
Sanghez, Valentina, Anna Luzzi, Don Clarke, Dustin Kee, Steven Beuder, Danielle Rux, Mitsujiro Osawa,
Joaquín Madrenas, Tsui-Fen Chou, Michael Kyba, and Michelina Iacovino. 2017. 'Notch
activation is required for downregulation of HoxA3-dependent endothelial cell phenotype during
blood formation', *PloS One*, 12: e0186818.
Savitsky, K., A. Bar-Shira, S. Gilad, G. Rotman, Y. Ziv, L. Vanagaite, D. A. Tagle, S. Smith, T. Uziel, S.
Sfez, M. Ashkenazi, I. Pecker, M. Frydman, R. Harnik, S. R. Patanjali, A. Simmons, G. A. Clines,
A. Sartiel, R. A. Gatti, L. Chessa, O. Sanal, M. F. Lavin, N. G. Jaspers, A. M. Taylor, C. F. Arlett,
T. Miki, S. M. Weissman, M. Lovett, F. S. Collins, and Y. Shiloh. 1995. 'A single ataxia
telangiectasia gene with a product similar to PI-3 kinase', *Science*, 268: 1749-53.
Schellenberg, M. J., P. P. Tumbale, and R. S. Williams. 2015. 'Molecular underpinnings of Aprataxin
RNA/DNA deadenylase function and dysfunction in neurological disease', *Progress in Biophysics*
and Molecular Biology, 117: 157-65.
Schubert, R., J. Reichenbach, and S. Zielen. 2002. 'Deficiencies in CD4+ and CD8+ T cell subsets in
ataxia telangiectasia', *Clinical and Experimental Immunology*, 129: 125-32.
Sedghi, Maryam, Mehri Salari, Ali-Reza Moslemi, Ariana Kariminejad, Mark Davis, Hayley Goullée, Björn
Olsson, Nigel Laing, and Homa Tajsharghi. 2018. 'Ataxia-telangiectasia-like disorder in a family
deficient for MRE11A, caused by a *MRE11* variant', *Neurology Genetics*, 4: e295.
Shakkottai, Vikram G, Maria do Carmo Costa, James M Dell'Orco, Ananthakrishnan Sankaranarayanan,
Heike Wulff, and Henry L Paulson. 2011a. 'Early changes in cerebellar physiology accompany
motor dysfunction in the polyglutamine disease spinocerebellar ataxia type 3', *Journal of*
Neuroscience, 31: 13002-14.
Shakkottai, Vikram G, Maolei Xiao, Lin Xu, Michael Wong, Jeanne M Nerbonne, David M Ornitz, and
Kelvin A Yamada. 2009. 'FGF14 regulates the intrinsic excitability of cerebellar Purkinje neurons',
Neurobiology of Disease, 33: 81-88.
Shakkottai, Vikram G., Maria do Carmo Costa, James M. Dell'Orco, Ananthakrishnan
Sankaranarayanan, Heike Wulff, and Henry L. Paulson. 2011b. 'Early Changes in Cerebellar
Physiology Accompany Motor Dysfunction in the Polyglutamine Disease Spinocerebellar Ataxia
Type 3', *The Journal of Neuroscience*, 31: 13002-14.
Shiloh, Yosef. 2020. 'The cerebellar degeneration in ataxia-telangiectasia: A case for genome instability',
DNA Repair, 95: 102950.
Shiloh, Yosef, and Yael Ziv. 2013. 'The ATM protein kinase: regulating the cellular response to genotoxic
stress, and more', *Nature Reviews Molecular Cell Biology*, 14: 197-210.
Solitare, G. B. 1968. 'Louis-Bar's syndrome (ataxia-telangiectasia). Anatomic considerations with
emphasis on neuropathologic observations', *Neurology*, 18: 1180-6.
Solitare, G. B., and V. F. Lopez. 1967. 'Louis-bar's syndrome (ataxia-telangiectasia). Neuropathologic
observations', *Neurology*, 17: 23-31.
Spring, K., S. Cross, C. Li, D. Watters, L. Ben-Senior, P. Waring, F. Ahangari, S. L. Lu, P. Chen, I. Misko,
C. Paterson, G. Kay, N. I. Smorodinsky, Y. Shiloh, and M. F. Lavin. 2001. 'Atm knock-in mice
harboring an in-frame deletion corresponding to the human ATM 7636del9 common mutation
exhibit a variant phenotype', *Cancer Research*, 61: 4561-8.
Staples, E. R., E. M. McDermott, A. Reiman, P. J. Byrd, S. Ritchie, A. M. Taylor, and E. G. Davies. 2008.
'Immunodeficiency in ataxia telangiectasia is correlated strongly with the presence of two null
mutations in the ataxia telangiectasia mutated gene', *Clinical and Experimental Immunology*, 153:
214-20.
Stoyas, Colleen A, David D Bushart, Pawel M Switonski, Jacquelyn M Ward, Akshay Alaghatta, Mi-bo
Tang, Chenchen Niu, Mandheer Wadhwa, Haoran Huang, and Alex Savchenko. 2020.
'Nicotinamide pathway-dependent Sirt1 activation restores calcium homeostasis to achieve
neuroprotection in spinocerebellar ataxia type 7', *Neuron*, 105: 630-44. e9.
Strich, Sabina J. 1966. 'Pathological findings in three cases of ataxia-telangiectasia', *Journal of*
Neurology, Neurosurgery, and Psychiatry, 29: 489.

Swift, M., D. Morrell, E. Cromartie, A. R. Chamberlin, M. H. Skolnick, and D. T. Bishop. 1986. 'The incidence and gene frequency of ataxia-telangiectasia in the United States', *American Journal of Human Genetics*, 39: 573-83.

Sykora, Peter, Deborah L. Croteau, Vilhelm A. Bohr, and David M. Wilson, 3rd. 2011. 'Aprataxin localizes to mitochondria and preserves mitochondrial function', *Proceedings of the National Academy of Sciences of the United States of America*, 108: 7437-42.

Takashima, H., C. F. Boerkoel, J. John, G. M. Saifi, M. A. Salih, D. Armstrong, Y. Mao, F. A. Quiocho, B. B. Roa, M. Nakagawa, D. W. Stockton, and J. R. Lupski. 2002. 'Mutation of TDP1, encoding a topoisomerase I-dependent DNA damage repair enzyme, in spinocerebellar ataxia with axonal neuropathy', *Nature Genetics*, 32: 267-72.

Tal, Efrat, Marina Alfo, Shan Zha, Ari Barzilai, Chris I. De Zeeuw, Yael Ziv, and Yosef Shiloh. 2018. 'Inactive Atm abrogates DSB repair in mouse cerebellum more than does Atm loss, without causing a neurological phenotype', *DNA Repair*, 72: 10-17.

Tavani, F., R. A. Zimmerman, G. T. Berry, K. Sullivan, R. Gatti, and P. Bingham. 2003. 'Ataxia-telangiectasia: the pattern of cerebellar atrophy on MRI', *Neuroradiology*, 45: 315-19.

Taylor, A. M., Z. Lam, J. I. Last, and P. J. Byrd. 2015. 'Ataxia telangiectasia: more variation at clinical and cellular levels', *Clinical Genetics*, 87: 199-208.

Terplan, K. L., and R. F. Krauss. 1969. 'Histopathologic brain changes in association with ataxia-telangiectasia', *Neurology*, 19: 446-54.

Tsutsumi, S., M. Yamazaki, T. Miyazaki, M. Watanabe, K. Sakimura, M. Kano, and K. Kitamura. 2015. 'Structure-function relationships between aldolase C/zebrin II expression and complex spike synchrony in the cerebellum', *Journal of Neuroscience*, 35: 843-52.

Tumbale, Percy, Matthew J Schellenberg, Geoffrey A Mueller, Emma Fairweather, Mandy Watson, Jessica N Little, Juno Krahn, Ian Waddell, Robert E London, and R Scott Williams. 2018. 'Mechanism of APTX nicked DNA sensing and pleiotropic inactivation in neurodegenerative disease', *The EMBO Journal*, 37: e98875.

Vacchio, Melanie S., Alexandru Olaru, Ferenc Livak, and Richard J. Hodes. 2007. 'ATM deficiency impairs thymocyte maturation because of defective resolution of T cell receptor α locus coding end breaks', *Proceedings of the National Academy of Sciences*, 104: 6323-28.

Vail, Graham, Aifang Cheng, Yu Ray Han, Teng Zhao, Shengwang Du, Michael M. T. Loy, Karl Herrup, and Mark R. Plummer. 2016. 'ATM protein is located on presynaptic vesicles and its deficit leads to failures in synaptic plasticity', *Journal of Neurophysiology*, 116: 201-09.

Valentin-Vega, Y. A., and M. B. Kastan. 2012. 'A new role for ATM: regulating mitochondrial function and mitophagy', *Autophagy*, 8: 840-1.

van der Burgt, I, K H Chrzanowska, D Smeets, and C Weemaes. 1996. 'Nijmegen breakage syndrome', *Journal of Medical Genetics*, 33: 153-56.

Verhagen, Mijke M.M., Jean-Jacques Martin, Marcel van Deuren, Chantal Ceuterick-de Groote, Corry M.R. Weemaes, Berry H.P.H. Kremer, Malcolm A.R. Taylor, Michèl A.A.P. Willemsen, and Martin Lammens. 2012. 'Neuropathology in classical and variant ataxia-telangiectasia', *Neuropathology*, 32: 234-44.

Voogd, Jan, and Mitchell Glickstein. 1998. 'The anatomy of the cerebellum', *Trends in Neurosciences*, 21: 370-75.

Wakasugi, M., T. Sasaki, M. Matsumoto, M. Nagaoka, K. Inoue, M. Inobe, K. Horibata, K. Tanaka, and T. Matsunaga. 2014. 'Nucleotide excision repair-dependent DNA double-strand break formation and ATM signaling activation in mammalian quiescent cells', *Journal of Biological Chemistry*, 289: 28730-7.

Wallis, LI, PD Griffiths, SJ Ritchie, CAJ Romanowski, G Darwent, and ID Wilkinson. 2007. 'Proton spectroscopy and imaging at 3T in ataxia-telangiectasia', *American journal of neuroradiology*, 28: 79-83.

Walter, Joy T, Karina Alvina, Mary D Womack, Carolyn Chevez, and Kamran Khodakhah. 2006. 'Decreases in the precision of Purkinje cell pacemaking cause cerebellar dysfunction and ataxia', *Nature Neuroscience*, 9: 389-97.

Weyemi, Urbain, Christophe E. Redon, Towqir Aziz, Rohini Choudhuri, Daisuke Maeda, Palak R. Parekh,
Michael Y. Bonner, Jack L. Arbiser, and William M. Bonner. 2015. 'NADPH oxidase 4 is a critical
mediator in Ataxia telangiectasia disease', *Proceedings of the National Academy of Sciences*,
112: 2121-26.
Xu, Y., and D. Baltimore. 1996. 'Dual roles of ATM in the cellular response to radiation and in cell growth
control', *Genes and Development*, 10: 2401-10.
Yamasaki, Miwako, Kouichi Hashimoto, and Masanobu Kano. 2006. 'Miniature Synaptic Events Elicited
by Presynaptic Ca^{2}+ Rise Are Selectively Suppressed by Cannabinoid Receptor
Activation in Cerebellar Purkinje Cells', *The Journal of Neuroscience*, 26: 86-95.
Zhou, Haibo, Zhanmin Lin, Kai Voges, Chiheng Ju, Zhenyu Gao, Laurens W. J. Bosman, Tom J. H.
Ruigrok, Freek E. Hoebeek, Chris I. De Zeeuw, and Martijn Schonewille. 2014. 'Cerebellar
modules operate at different frequencies', *Elife*, 3: e02536.
Zhu, L., B. Scelfo, F. Tempia, B. Sacchetti, and P. Strata. 2006. 'Membrane excitability and fear
conditioning in cerebellar Purkinje cell', *Neuroscience*, 140: 801-10.

1259

8.0 Figure Legends

Figure 1. New A-T mouse models expressing clinically related PTCs. A) The *Atm* gene locus was

targeted by homologous recombination of a targeting vector containing a modified NorCOMM cassette

in intron one and the corresponding A-T PTC mutation in exon 3 to create the targeted *Atm^{R35X}* and

Atm^{Q35X} ES cell lines. Following germline transmission of these alleles in mice, the floxed NorCOMM

cassette was removed by Cre excision *in vivo* to produce the final *Atm^{R35X}* and *Atm^{Q35X}* mouse lines. **B)**

Genotyping of A-T mouse models. PCR agarose gel of mouse DNA shows 151 bp wildtype (Wt) allele

band and 241 bp Cre-excised targeted allele band. **C)** ATM levels were examined using immunoblot

analyses of the spleen due to its high expression density in this tissue. Exemplar blots illustrate a gene

dose effect of ATM protein expression in samples harvested from wildtype (Wt), heterozygous (het), and

homozygous *Atm^{R35x}* and *Atm^{Q35X}* mice as indicated. **D)** Breeding scheme schematic for double mutant

and control mice for this study. **E)** *Atm^{R35X/R35X}; Aptx^{-/-}* mice develop an ataxia that at late stages results

in a severe loss of motor coordination and ability to ambulate (see Videos S1-S4). Abbreviations for

panel **1**: **hβA**-human beta Actin promotor; Δ**TK1**-delta TK1, inactivated Thymidine Kinase 1; **T2A**-self-

cleaving peptide sequence; **Neo**-Neomycin gene; **PGKpA**-Phosphoglycerate kinase poly A tail; **loxP**-

recombination elements are show as a blue triangle; orientation of the Gateway **attB** recombination

elements by an orange arrow; orientation of the genotyping **F** and **R** primers is shown by green and blue

arrows respectively; and engineered PTC sites are shown in exon 3 by a red circle.

Figure 2. Health and survivability of single and double mutant mice. A) Left: The line color and

symbol for each genotype is denoted and is consistent across all figures (1-8). **Right:** *Atm^{R35X/R35X}; Aptx^{-}*

^{/-} mice weighed significantly less than all control genotypes as indicated by the growth curves ($\pm$ 95%

confidence interval; dotted lines). Growth curve (*Atm^{R35X/R35X}; Aptx^{-/-} vs.* controls): Male k = 0.024 *vs.*

0.011-0.019, Y_{max} = 21.8 *vs.* 32.9-41.0 g, (n = 3 to 18); Female k = 0.030 *vs.* 0.017-0.022, Y_{max} = 16.9 *vs.*

23.3-31.3, (n = 2 to 19). Sum of squares F-test run across all curves: Male $F_{(12,\,364)}$ = 30.5, ****p<0.0001,

Female $F_{(12,\,339)}$ = 28.3, ****p<0.0001. **B)** ATM deficient mice, regardless of APTX expression, displayed

significantly lower survivability with ~55% of mice deceased by P400. No statistical differences between

ATM deficient mice were detected. Moreover, a single wildtype copy of the *Atm* gene was sufficient to

prevent premature death (no statistical difference detected between *Atm^R35X/Wt; Aptx^-/-* and *Atm^Wt/Wt;*

Aptx^+/+ mice). Log-rank (Mantel-Cox) tests across all ($\chi^2_{(6, 217)}$ = 48.4, ****p<0.0001), just the ATM

deficient ($\chi^2_{(2, 217)}$ = 1.06, p=0.6), and single comparisons to wildtype (see figure) were conducted. Total

number of animals indicated in panel **C**. **C)** Pie charts illustrating that ATM deficient mice displayed a

high prevalence of thymomas based on postmortem necropsies. "Other" probable causes of death

included enlarged livers, and obstructed kidneys. "Missing" mice were presumed dead and cannibalized

by cage mates, cause of death unknown. **Figure 2-figure supplement 1, Figure 2-source data**

Figure 3. *Atm^R35X/R35X; Aptx^-/-* mice develop a progressive loss in motor coordination. A)

Atm^R35X/R35X; Aptx^-/- mice take a similar amount of time to descend a vertical pole at P45, 120, and 210,

but significantly longer at P400. These overall results were found to be similar for both male (left, n = 2

to 12) and female (right, n=4 to 12) mice. **B)** Consistent with the vertical pole test, the gait of *Atm^R35X/R35X;*

Aptx^-/- mice measured during ambulation on a Catwalk gait analysis system was significantly different to

controls by P400, but not before P210. This includes the percent of time a mouse spends with 3 *vs.* 1, 2,

or 4 paws on the ground and the speed and cadence during each run across the platform. The effects of

the two null mutations were generally similar between males (left, n = 4 to 21) and females (right, n = 3

to 18). **C)** Behavioral data for male (blue) and female (pink) *Atm^R35X/R35X; Aptx^-/-* (dark purple, left) and

Atm^Wt/Wt; Aptx^+/+ (orange, right) mice are plotted at P30 (left 2 columns) and 400 (right 2 columns). Left-

right asymmetries of the horizontal bars indicate a difference in performance between genotypes for the

behavioral test listed in a column on the far left. A significant difference in the time to right during the

righting reflex at P8 was observed in both Male and Female mice (bottom). **A** and **B** were examined via

two-way ANOVA with age and genotype as factors followed by *potshot* Tukey's multiple comparison tests

between *Atm^R35X/R35X; Aptx^-/-* and each of the control genotypes. Behavioral tests in **C** were examined

using a non-parametric Kruskal Wallace followed by *postdocs* Dunn's multiple comparisons tests.

Symbol/color key: *Atm^Wt/Wt; Aptx^+/+* (purple circle), *Atm^Wt/Wt; Aptx^-/-* (blue diamond), *Atm^R35X/R35X; Aptx^+/+*

(green triangle), $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (orange square), $Atm^{Wt/R35X}$; $Aptx^{-/-}$ (red inverted triangle) **Figure 3-**

figure supplement 1, Figure 3-source data

Figure 4. The biophysical properties of PNs are significantly perturbed in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice.

A) Schematic diagram of intracellular recording from a single Purkinje neuron (PN) in an acute cerebellar

tissue slice preparation used to examine their biophysical properties. **B) Left:** Voltage clamp

measurements of PN neuron membrane properties were made from a $1\,s$, $-5\,mV$ step pulse as illustrated.

Right: The membrane input resistance (R_m), time constant (τ), and capacitance (C_m) were perturbed in

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ compared to wildtype mice. **C)** Current clamp recordings of PN action potentials (AP)

after $2\,nA$ step pulses from a $-70\,mV$ holding potential. PN action potentials recorded from $Atm^{R35X/R35X}$;

$Aptx^{-/-}$ fail to maintain constant firing and summary plots show that they have lower 1^{st} AP amplitudes,

firing threshold, and area under the curve. **D) Top:** Example mEPSC recordings taken from a PN under

voltage clamp at a $-80\,mV$ holding potential. **Bottom:** Median frequency and amplitude data, along with

the overall probability distribution function are plotted for both $Atm^{Wt/Wt}$; $Aptx^{+/+}$ ($n = 11$) and $Atm^{R35X/R35X}$;

$Aptx^{-/-}$ ($n = 11$) mice. The frequency, but not amplitude of PNs recorded in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice was

found to be perturbed. **E and F) Left:** Example traces of evoked EPSCs recorded from PNs as a result

of a 2-pulse stimulation ($50\,ms$ interval) of either parallel (**G**) or climbing (**H**) axon fibers. Traces illustrate

the first (A_1) and second (A_2) amplitude (normalized) and time course of first decay (blue fitted line) of

each synaptic response. **Right:** Summary plots of the paired pulse ratio. While parallel fiber paired pulse

facilitation was normal in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice, climbing fiber paired pulse depression and halfwidth

was significantly perturbed compared to $Atm^{Wt/Wt}$; $Aptx^{+/+}$ mice. Data in **B** were compared using an

ANOVA (Kruskal-Wallis) followed by Dunn's multiple comparisons test while data in **D to F** were

compared via Welch's t-test. Symbol/color key: $Atm^{Wt/Wt}$; $Aptx^{+/+}$ (purple circle), $Atm^{R35X/R35X}$; $Aptx^{-/-}$

(orange square) **Figure 4-figure supplement 1**

Figure 5. Cerebellar atrophy is associated with a progressive reduction in PN neural action

potential firing frequency and PN dendritic length. A) Schematic diagram of extracellular recording

from a single Purkinje neuron (PN) in an acute cerebellar tissue slice preparation. Example electrophysiological traces for $Atm^{Wt/Wt}$; $Aptx^{+/+}$ (purple, top) and $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (orange, bottom) PNs in the medial (vermis) area of the cerebellum. **B)** $Atm^{R35X/R35X}$; $Aptx^{-/-}$ PN action potential firing frequency progressively decreased with age and was significantly slower in comparison to all control genotypes expressing at least one copy of the Atm or $Aptx$ gene. [$Atm^{Wt/Wt}$; $Aptx^{+/+}$ (n=52 to 59), $Atm^{Wt/Wt}$; $Aptx^{-/-}$ (n = 51 to 64), $Atm^{R35X/R35X}$; $Aptx^{+/+}$ (n = 39 to 52), $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (n = 24 to 71), $Atm^{Wt/R35X}$; $Aptx^{-/-}$ (n = 69)] **C)** Cartoon image of the brain highlighting the dorsal forebrain and cerebellar surface. 2-dimensional area estimates from dorsal images of the brain were used to determine the cerebellum to forebrain ratio allowing us to control for any differences in overall size of the brain. We found the cerebellum decreased in size over age in $Atm^{R35X/R35X}$; $Aptx^{-/-}$(n = 5 to 10), but not control mice [$Atm^{Wt/Wt}$; $Aptx^{+/+}$ (n = 4 to 20), $Atm^{Wt/Wt}$; $Aptx^{-/-}$ (n = 4 to 12), $Atm^{R35X/R35X}$; $Aptx^{+/+}$ (n = 6 to 16), $Atm^{Wt/R35X}$; $Aptx^{-/-}$ (n = 6)]. **D)** Immunofluorescent images of $Atm^{Wt/Wt}$; $Aptx^{+/+}$ (top) and $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (bottom) at P400 (medial cerebellar lobule VIII). Scale bar = 50 μm **E)** The width of the molecular layer, but not the granule cell layer progressively declined over age (n = 5 to 7). Statistical significances were assessed via 2-way ANOVA with age and genotype as factors followed by *posthoc* Holm-Sidak (**B**, **C**, and **E (top)**) or Sidak (**E bottom**) pairwise multiple comparisons test. **Figure 5-figure supplements 1-5, Figure 5-source data**

Figure 6. T-cell deficits are found in the blood of $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice. A) Representative flow cytometric profiles of T-cell glycoprotein marker CD3 and summary plots indicate ATM and/or APTX deficient mice have decreased proportions of CD3+ T-cells in the blood. **B)** Representative flow cytometric profiles of T-cell glycoprotein markers CD4 and CD8 gated on CD3+ cells and summary plots for CD8 and CD4 single positive cell proportions. ATM deficient mice had reduced CD4+ proportions compared to mice with at least one copy of the Atm gene. Statistical significances were assessed via 1-way ANOVA followed by *posthoc* Tukey's pairwise multiple comparisons tests. Number of animals denoted at bottom of bar. Symbol/color key: $Atm^{Wt/Wt}$; $Aptx^{+/+}$ (purple circle), $Atm^{R35X/R35X}$; $Aptx^{+/+}$ (green triangle), $Atm^{Wt/R35X}$; $Aptx^{-/-}$ (red inverted triangle), $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (orange square) **Figure 6/7-source data**

Figure 7. ATM and APTX deficiency confer deficits in T-cell expression, but at different

developmental stages. A) Representative flow cytometric profiles of T-cell glycoprotein markers CD44

and CD25 gated on CD4$^-$CD8$^-$ double negative (DN) cells. Summary plots show proportions of

thymocytes at DN stages 1-4 (left to right). APTX deficient mice display increased proportions for DN1-3

and decreased proportion at DN4 consistent with a deficit in ontogeny from DN3 to DN4. **B)**

Representative flow cytometric profiles of T-cell glycoprotein markers CD4 and CD8 gated. ATM deficient

mice display decreased proportions for CD4 and CD8 single positive cells consistent with a deficit in

ontogeny from CD4$^+$CD8$^+$ double positive to CD4$^+$ and CD8$^+$ single positive fates. Statistical significances

were assessed via 1-way ANOVA followed by *posthoc* Tukey's pairwise multiple comparisons tests.

Number of animals denoted at bottom of bars. Symbol/color key: *Atm$^{Wt/Wt}$; Aptx$^{+/+}$* (purple circle),

Atm$^{R35X/R35X}$; Aptx$^{+/+}$ (green triangle), *Atm$^{Wt/R35X}$; Aptx$^{-/-}$* (red inverted triangle), *Atm$^{R35X/R35X}$; Aptx$^{-/-}$* (orange

square)

Figure 8. ATM protein expression is restored after readthrough compound exposure in explant

tissues from *Atm$^{R35X/R35X}$* and *Atm$^{Q35X/Q35X}$.* Spleen and cerebellar explant tissue from *Atm$^{R35X/R35X}$* and

Atm$^{Wt/Wt}$ mice were treated with vehicle, the readthrough compounds G418 ($100\ \mu M$), or GJ103 (100

μM) for $72\ hrs$. ATM immunoblots show recovery of ATM (MW $350\ kDa$) production in both the spleen (n

$= 2$) and cerebellum (n $= 3$). Equal loading was assessed via housekeeping genes (Actin or GAPDH)

and ponceau staining. **Figure 8-source data**

9.0 Figures

Fig. 1

Fig. 2

1382

1383

Fig. 3

1384

1385

Fig. 4

1386

1387

Fig. 5

Fig. 6

1389

Fig. 7

1390

1391

Fig. 8

10.0 Figure Supplements

Figure 2-figure supplement 1

Figure 1-figure supplement 1. Animal weight for each time point and genotype. A) The average weights are plotted for each genotype at each of the indicated time points. Growth curves without experimental, 2-way ANOVA with age and genotype as factors. Male: $F_{(10, 226)}= 5.6$, p<0.0001; Female: $F_{(10, 197)}= 7.3$, p<0.0001. **B)** The survivability of each genotype of mice are plotted for male and female individually.

1401

Figure 3-figure supplement 1. $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice develop progressive ataxia. A) $Atm^{R35X/R35X}$;

$Aptx^{-/-}$ mice develop an ataxia that at late stages results in a severe loss of coordination and ability to

ambulate. **B)** Additional gait analyses measuring stride length and time of overlap of the hindlimbs.

Examined via two-way ANOVA with age and genotype as factors followed by *potshot* Tukey's multiple

comparison tests between $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and each of the control genotypes. **C)** Behavioral deficits

are seen only in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice across all behavioral tests and sexes. Behavioral tests were

examined using a non-parametric Kruskal Wallace followed by *posthoc* Dunn's multiple comparisons

tests.

1410

Figure 4-figure supplement 1. Current *vs.* voltage responses significantly differ between *Atm$^{Wt/Wt}$;*

***Aptx$^{+/+}$ and Atm$^{R35X/R35X}$; Aptx$^{-/-}$ mice. A)** PN voltage responses to various current steps between -500

and 2250 pA (250 pA steps) from a -70 mV holding current in *Atm$^{Wt/Wt}$; Aptx$^{+/+}$* (top, purple) and

Atm$^{R35X/R35X}$; Aptx$^{-/-}$ (bottom, orange) mice. **(B)** I–V curves calculated from either max deflection (V$_{m\ max}$)

or steady state (V$_{m\ end}$) for *Atm$^{Wt/Wt}$; Aptx$^{+/+}$* (purple) and *Atm$^{R35X/R35X}$; Aptx$^{-/-}$* (orange) mice. **C)** Various

measurements of the voltage response to -500 pA step pulse (blue box in **B**) in *Atm$^{Wt/Wt}$; Aptx$^{+/+}$* (purple)

and *Atm$^{R35X/R35X}$; Aptx$^{-/-}$* (orange) mice. Significance was tested using a non-parametric Mann Whitney

test.

Figure 5-figure supplement 1. Mean variation between PN firing intervals across the cerebellum.

Average CV2 of PN firing frequency is plotted across the indicated locations at P45, 120, 210, and 400.

1424

Figure 5-figure supplement 2. Mean PN firing frequency across genotype and sex. Average PN firing frequency for all cells recorded from male and female mice is plotted for the indicated genotype. No significant differences were observed between sex. 2-Way ANOVA with age and sex as factors, $Atm^{Wt/Wt}$; $Aptx^{+/+}$ ($F_{(1, 751)} = 1.15$, p=0.3), $Atm^{Wt/Wt}$; $Aptx^{-/-}$ ($F_{(1, 797)} = 1.10$, p=0.3), $Atm^{R35X/R35X}$; $Aptx^{+/+}$ ($F_{(1, 630)} = 0.17$, p=0.7), $Atm^{R35X/R35X}$; $Aptx^{-/-}$ ($F_{(1, 666)} = 1.10$, p=0.4), t-test for P400 $Atm^{Wt/R35X}$; $Aptx^{-/-}$ (p=0.9)

Figure 5-figure supplement 3. Coefficient of Variation of PN firing frequency across the cerebellum. Average CV of PN firing frequency is plotted across the indicated locations at P45, 120, 210, and 400. No significant differences ($p<0.5$) were detected across all areas using 2-way ANOVA with age and genotype as factors.

1435

Figure 5-figure supplement 4. Mean variation between PN firing intervals across the cerebellum.

Average CV2 of PN firing frequency is plotted across the indicated locations at P45, 120, 210, and 400.

No significant differences ($p<0.5$) were detected across all areas using 2-way ANOVA with age and

genotype as factors.

Figure 5-figure supplement 5. Histopathological effects of ATM and ATPX deficiency. A) Width

measurements of the molecular and granule cell layer (ML and GCL respectively) for each folia across

the medial intermediate and lateral areas of the cerebellum. **B) (Top)** Summary plot indicates no

significant differences in the 2D linear density of PNs across each folia in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ vs. $Atm^{Wt/Wt}$;

$Aptx^{+/+}$. **(Bottom)** Images of parasagittal cerebellar slices ($2.5X$) illustrating the lack of significant gaps in

PN (green, Calbindin) or granule cell layers (magenta, DAPI). Scale bar = 0.5 mm **(C)** Images of

parasagittal sections from $Atm^{Wt/Wt}$; $Aptx^{+/+}$ (top row) and $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (bottom rows) mice ($63X$).

PNs (green, Calbindin) were found to have a larger diameter than those found in $Atm^{Wt/Wt}$; $Aptx^{+/+}$ cells

(red arrows). PN axonal swellings were also observed in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ sections (magenta arrows).

Scale bar = 50 μm

11.0 Source files
Figure 2- source data. Weight, age of death, and probable cause of death

Figure 3-source data. Raw behavior data

Figure 4-source data. Electrophysiological data

Figure 5- source data. Individual average firing frequencies for each recorded cell; Individual CV
for each recorded cell; Brain area

Figure 6/7- source data. Tables of FACs data

Figure 8-source data. Western blot measurements

12.0 Rich Media
Video S1. Pole test, $Atm^{Wt/Wt}$ vs. $Atm^{R35X/R35X}$. $Atm^{R35X/R35X}$ do not display an ataxic phenotype at
P460.

Video S2. Pole test, $Aptx^{+/+}$ vs. $Aptx^{-/-}$. $Aptx^{-/-}$ mice do not display an ataxic phenotype at P460.

Video S3. Pole test, $Atm^{Wt/Wt}$; $Aptx^{+/+}$ vs. $Atm^{R35X/R35X}$; $Aptx^{-/-}$. $Atm^{R35X/R35X}$; $Aptx^{-/-}$ have considerable
motor disability at P460.

Video S4. Open field, $Atm^{Wt/Wt}$; $Aptx^{+/+}$ vs. $Atm^{R35X/R35X}$; $Aptx^{-/-}$. $Atm^{R35X/R35X}$; $Aptx^{-/-}$ display a clear
inability to ambulate in the open field at P460.

1467

1468

1469

1470