A novel, ataxic mouse model of Ataxia Telangiectasia caused by a clinically relevant nonsense mutation

Perez, Harvey [¥,1]; Abdallah, May, F.[¥,1]; Chavira, Jose, I. [¥,1]; Norris, Angelina, S.[1]; Egeland, Martin, T. [1];

Vo, Karen, L. [1]; Buechsenschuetz, Callan, L. [1]; Sanghez, Valentina [1]; Kim, Jeannie, L. [1]; Pind, Molly [2],

Nakamura, Kotoka [3], Hicks, Geoffrey, G. [2]; Gatti, Richard, A. [3]; Madrenas, Joaquin [1,5]; Iacovino,

Michelina [1,4]; McKinnon, Peter, J. [6]; Mathews, Paul, J. [*,1,7]

1. *The Lundquist Institute for Biomedical Innovation, Harbor-UCLA Medical Center, Torrance, CA*
2. *Department of Biochemistry & Medical Genetics, Max Rady College of Medicine, University of Manitoba*
3. *Department of Pathology & Laboratory Medicine, David Geffen School of Medicine, University of California, Los Angeles, CA*
4. *Department of Pediatrics, Harbor-UCLA Medical Center, Torrance, CA*
5. *Department of Medicine, Harbor-UCLA Medical Center, Torrance, CA*
6. *Center for Pediatric Neurological Disease Research, St. Jude Pediatric Translational Neuroscience Initiative, St. Jude Children's Research Hospital, Memphis, TN, USA*
7. *Department of Neurology, Harbor-UCLA Medical Center, Torrance, CA*

[¥]*Contributed equally*

[*]*Corresponding Author*

Abstract

Ataxia Telangiectasia (A-T) and Ataxia with Ocular Apraxia Type 1 (AOA1) are devastating neurological disorders caused by null mutations in the genome stability genes, A-T mutated (*ATM*) and Aprataxin (*APTX*), respectively. Our mechanistic understanding and therapeutic repertoire for treating these disorders is severely lacking, in large part due to the failure of prior animal models with similar null mutations to recapitulate the characteristic loss of motor coordination (i.e., ataxia) and associated cerebellar defects. By increasing genotoxic stress through the insertion of null mutations in both the *Atm* (nonsense) and *Aptx* (knockout) genes in the same animal, we have generated a novel mouse model that for the first time develops a progressively severe ataxic phenotype associated with atrophy of the cerebellar molecular layer. We find biophysical properties of cerebellar Purkinje neurons are significantly perturbed (e.g., reduced membrane capacitance, lower action potential thresholds, etc.), while properties of synaptic inputs remain largely unchanged. These perturbations significantly alter

Purkinje neuron neural activity, including a progressive reduction in spontaneous action potential firing

frequency that correlates with both cerebellar atrophy and ataxia over the animal's first year of life.

Double mutant mice also exhibit a high predisposition to developing cancer (thymomas) and immune

abnormalities (impaired early thymocyte development and T-cell maturation), symptoms characteristic

of A-T. Lastly, by inserting a clinically relevant nonsense-type null mutation in *Atm*, we demonstrate that

Small **M**olecule **R**ead-**T**hrough (SMRT) compounds can restore ATM production, indicating their

potential as a future A-T therapeutic.

1.0 Introduction

Ataxia Telangiectasia (A-T) is a rare (1 in ~100,000) (Swift et al. 1986), autosomal recessive genetic

disorder characterized by cancer predisposition, immune deficiency, and a highly penetrant progressive

and severe ataxia linked to cerebellar atrophy (Rothblum-Oviatt et al. 2016; Boder and Sedgwick 1958;

Levy and Lang 2018). A-T patients typically die in their second or third decade of life (Crawford et al.

2006) from lymphatic cancers, respiratory infections, or complications of ataxia—unfortunately,

survivability has not dramatically changed since the 1950s (Micol et al. 2011; Rothblum-Oviatt et al.

2016). While disease progression and cause of death vary widely across patients, the highly penetrant

progressive decline in motor coordination is reported as having the greatest negative impact on a

patient's quality of life (Jackson et al. 2016). Care is generally palliative, directed at reducing, limiting, or

eliminating cancers or infections. No long-term therapies are available for treating the ataxia and

associated cerebellar dysfunction and atrophy.

A-T is caused by deficiency or dysfunction of the ATM (A-T mutated) protein (Savitsky et al. 1995).

Premature termination codon (PTC) causing nonsense mutations account for up to a half of known

cases, with missense and deletion mutations also contributing (Concannon and Gatti 1997; Sandoval et

al. 1999). ATM is a serine/threonine PIKK family kinase that plays a key role in the DNA damage

response (DDR), protecting cells from the tens of thousands of DNA lesions incurred each day (Lindahl

and Barnes 2000; Kastan and Bartek 2004; Shiloh and Ziv 2013). In the active monomeric form, ATM

phosphorylates several key proteins halting the production of new DNA (cell cycle arrest) (Ando et al.

2012), and then, depending on severity of the damage, initiates DNA repair or programmed cell death

(i.e., apoptosis) (Ando et al. 2012; Rashi-Elkeles et al. 2006). Several downstream DDR pathway

targets of ATM have been identified, including p53, CHK2, BRCA1, SMC1, and NBS1 (Matsuoka et al.

2007). ATM's role in DNA repair is also implicated in normal immune system development, where it is

proposed to contribute to the recombination of natural DNA splicing that occurs during gene

rearrangement in T- and B-lymphocyte maturation (Chao, Yang, and Xu 2000; Matei, Guidos, and

Danska 2006; Vacchio et al. 2007; Schubert, Reichenbach, and Zielen 2002). Although its roles are still

emerging, ATM has also been implicated in oxidative stress homeostasis (Guo et al. 2010) and

mitophagy (Valentin-Vega and Kastan 2012; Pizzamiglio, Focchi, and Antonucci 2020).

It is unclear why ATM deficiency causes ataxia, but it is far from the only DDR protein linked to ataxia,

as Aprataxin (APTX) (Aicardi et al. 1988), Meiotic recombination 11 homolog 1 (MRE11) (Sedghi et al.

2018), Nibrin (NBS1) (van der Burgt et al. 1996), Senataxin (SETX) (Moreira et al. 2004), and Tyrosyl-

DNA Phosphodiesterase 1 (TDP1) (Takashima et al. 2002) when absent or dysfunctional can cause

cerebellar-related ataxia. This suggests that the neurological features of genome instability syndromes

have a common underlying cause, although this has yet to be mechanistically demonstrated (McKinnon

2009; Rass, Ahel, and West 2007).

A major factor limiting our ability to define why loss of DDR proteins, like ATM, selectively impacts the

cerebellum and causes progressive ataxia is the lack of an animal model that recapitulates these

neurological symptoms (Lavin 2013). Several A-T rodent models have been created over the past

several years by inserting gene mutations that cause protein dysfunction (lack kinase activity) or

complete deficiency (Herzog et al. 1998; Xu and Baltimore 1996; Elson et al. 1996; Spring et al. 2001;

Campbell et al. 2015; Quek et al. 2016; Tal et al. 2018; Lavin 2013); a minipig was also recently

reported (Beraldi et al. 2017). However, none develop an overt, progressive ataxia with cerebellar

dysfunction and atrophy that recapitulates the human disease, even though other aspects of the

disorder like thyroid cancers, infertility, and immune abnormalities do develop. It remains unclear why

these prior animal models fail to display the progressive ataxic phenotype (Lavin 2013). It is possible

that species-specific molecular compensations in mice provide redundancies or alternative pathways

minimizing the effects of ATM deficiency in the brain (El-Brolosy and Stainier 2017). It is also possible

that the shortened lifespan of prior models (Barlow et al. 1996) is too brief for the stochastic

mechanisms driving cerebellar dysfunction and atrophy to accumulate and impact motor behavior.

Other challenges include potentially leaky genetic manipulations that result in low levels of ATM protein

or active fragments with residual kinase activity, thus limiting neuropathology (Li et al. 2011). The

impact of missing such a crucial animal model has been significant, severely limiting experimental

studies from identifying the cellular and molecular mechanisms and hampering pre-clinical development

and testing of much needed therapeutics.

We test here whether increasing genotoxic stress, by placing null mutations in not just the *Atm* gene,

but also the related *Aptx* gene, leads to a more representative mouse model that displays cerebellar

dysfunction, atrophy, and the development of progressive ataxia. We chose to additionally knock-out

Aptx because its deficiency causes an A-T like disorder in humans called ataxia with ocular apraxia

type 1 (AOA1), which does not feature A-T's other system defects that could increase the potential for

prenatal lethality or early death (e.g., immunodeficiency and cancer predisposition) (Coutinho P 2002).

Moreover, APTX is a phosphodiesterase involved in DNA reassembly after double- and single-stranded

repair, and has a function downstream of—but not directly regulated by or related to—ATM (Gueven et

al. 2004; Schellenberg, Tumbale, and Williams 2015; Ahel et al. 2006). We hypothesized that functional

expression of both proteins would have an additive effect and induce neurological dysfunction. Our

results indeed demonstrate that mice deficient in ATM and APTX develop cerebellar dysfunction,

atrophy, and a progressive and profound ataxia, while mice deficient in either protein alone do not.

Additionally, double mutants displayed other characteristic symptoms of A-T, including defects in

immune maturation and a high incidence of cancer (thymomas), making it the most representative

model, from a phenotypic standpoint, to date.

Finally, we designed this new mouse model to test our recently developed **S**mall **M**olecule **R**ead-

Through Compounds (SMRT) that enable translation through premature termination codons (Du et al.

2013). Thus, we inserted a premature termination-causing nonsense mutation (103C>T) in the *Atm*

gene common to a large family of North African A-T patients (Gilad, Bar-Shira, et al. 1996). This

mutation results in a premature termination codon (PTC) at what would normally be amino acid 35 and

the loss of ATM translation. Here, we report proof-of-principle experiments demonstrating that clinically

relevant genetic mutations incorporated into the A-T mouse model are amenable to read-through

compounds and thus appropriate for preclinical testing of SMRT compounds.

115   2.0 Results

2.1 Creation of a new A-T mutant mouse model expressing a clinically relevant nonsense

mutation

To create a more clinically relevant mouse model of A-T we used a gateway recombination cloning and

site-directed mutagenesis method to recapitulate a c.103C>T (p.R35X) mutation in the *ATM* gene

found in a large population of North African A-T patients (**Fig. 1A and Methods**) (Gilad, Khosravi, et al.

1996). The insertion of thymine in place of cytosine at this site in exon 3 results in a premature

termination codon (PTC)-causing nonsense mutation in the ATM gene. Since the c.103C>T mutation

results in different PTCs in the human gene compared to the mouse *Atm* gene—TGA *vs.* TAG,

respectively—we created two different mice by exchanging the mouse *Atm* exon 3 with either a human

or mouse exon 3 variant with the c.103C>T mutation (**Fig. 1B**). In the human variant, a 103C>T

mutation of the mouse codon, where the arginine (R) encoding codon (CGA) becomes a TGA stop

codon, results in a mouse we denote as Atm^{R35X} (officially $Atm^{Tm1.1(103CAG)TGA)Mfgc}$). In the mouse variant,

the c.103C>T mutation transforms a glutamine (Q)-encoding CAG codon into a TAG stop codon and is

denoted Atm^{Q35X} (officially $Atm^{Tm1.1(103C)T)Mfgc}$). The presence of the PTC results in a loss of ATM

expression, either reduced by about half in the heterozygote expressing one normal mouse copy of the

Atm gene ($Atm^{R35X/+}$ or $Atm^{Q35X/+}$), or completely in the homozygote ($Atm^{R35X/R35X}$ or $Atm^{Q35X/Q35X}$) (**Fig.**

1C).

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ (double mutant) mice were created by first crossing single mutant $Atm^{R35X/R35X}$

(congenic on the C57BL/6J background) and $Aptx^{-/-}$ (mixed C57BL/6J and 129 background) mice to

135 generate heterozygote $Atm^{R35X/+}$; $Aptx^{+/-}$ mice. F1-5 littermate $Atm^{R35X/+}$; $Aptx^{+/-}$ mice were then crossed

136 within litters to create sufficient numbers of the desired experimental and control genotypes to

137 determine how loss of different amounts of ATM and APTX affects the animal's phenotype (**Fig. 1D**).

138 Like prior ATM-deficient A-T mouse models, ATM or APTX deficiency alone did not result in mice with

139 ataxia (**Video 1 and 2**). However, deficiency in both proteins ($Atm^{R35X/R35X}$; $Aptx^{-/-}$) results in the

140 development of a severe and progressively ataxic phenotype (**Fig. 1E**, **Video 3 and 4**).

141 **2.2 ATM-deficient mice have lowered survivability and a high incidence of thymomas**

142 We assessed the general health and development of control and experimental mice expressing

143 different levels of ATM and APTX (**Fig. 2**). We found that $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice grew ~55% slower

144 and reached estimated plateau weights that were ~35% less than control genotypes (log-rank, n = 21 to

145 40, p<0.0001; **Fig. 2A**). These differences in weight were a postnatal phenomenon, as no significant

146 weight differences were detected just after birth (P8) across all genotypes (1-way ANOVA, n = 5 to 23,

147 p>0.23). Adolescent double mutant mice at postnatal day 45 (P45) weighed on average 30% less in

148 male mice [double mutant: 14.4 ± 1.0 g (n = 13) *vs.* wildtype: 20.2 ± 0.5 g (n = 16), *t*-test, p<0.0001] and

149 25% less in females [double mutant: 12.7 ± 0.6 g (n = 17) *vs.* wildtype: 17.0 ± 0.2 g (n = 15), *t*-test,

150 p<0.0001; **Fig. 1A**]. Differences across the control genotypes were observed, but they were small and

151 not consistent across time points or sex and therefore judged to not be physiologically relevant (**Fig.**

152 **2A**). Survivability of the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice was significantly reduced compared to $Atm^{+/+}$; $Aptx^{+/+}$

153 mice, with 53% of mice still alive at 400 days of age, compared to 97% of $Atm^{+/+}$; $Aptx^{+/+}$ mice at the

154 same time point (**Fig. 2B**). ATM deficiency alone was sufficient to reduce survivability; as compared to

155 $Atm^{+/+t}$; $Aptx^{+/+}$ mice, both $Atm^{R35X/R35X}$; $Aptx^{+/+}$ and $Atm^{R35X/R35X}$; $Aptx^{+/-}$ mice had significantly reduced

156 survivability [42%, log-rank, $\chi^2_{(1, 56)} = 13.49$, p=0.0002 and 52%, log-rank, $\chi^2_{(1, 53)} = 19.54$, p<0.0001,

157 respectively]. No significant difference in survivability between ATM deficient mice with partial or

158 complete APTX deficiency was detected [log-rank, $\chi^2_{(2, 85)} = 1.01$, p=0.6]. Conversely, mice harboring at

159 least one functional copy of the *Atm* gene had normal survivability, regardless of whether they

160 expressed APTX or not [log-rank, $\chi^2_{(3, 131)} = 3.08$, p=0.4]. No significant difference between male and

female mice was observed, and thus data were pooled [log-rank, $p > 0.4$ for all pairwise comparisons;

Fig. 2-fig. S1B]. Generally, a third of mice with ATM deficiency died from complications related to large

thymic cancers (thymoma) found in the thoracic cavity (**Fig. 2C**). The presence or absence of APTX did

not impact cancer prevalence, and mice with at least one *Atm* transcript were cancer free up until at

least P400. Overall, ATM but not APTX deficiency had severe effects on the health and survivability of

mice.

2.3 Both ATM and APTX deficiency are necessary to produce progressive motor dysfunction

The progressive development of severe ataxia is a hallmark characteristic of A-T that is recapitulated in

the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice but none of the other control genotypes we tested. Overall, we find motor

coordination deficits emerge between 210 and 400 days after birth in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice and find

no evidence of ataxia in mice with at least one copy of the *Atm* or *Aptx* gene (**Fig. 3A, B**). For the

vertical pole test, $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice took twice as long to descend at P400 compared to $Atm^{+/+}$;

$Aptx^{+/+}$, $Atm^{+/+}$; $Aptx^{-/-}$, $Atm^{R35X/R35X}$; $Aptx^{+/+}$, or $Atm^{R35X/+}$; $Aptx^{-/-}$ mice [Male: 29.1 ± 0.9 s (n = 3) *vs.* $7.5 \pm$

0.4 s (n = 12), 12.5 ± 2.5 s (n = 9), 9.2 ± 0.9 s (n = 10), 8.6 ± 0.9 s (n = 11), 1-way ANOVA, $F_{(4, 40)} = 19.9$,

$p < 0.0001$; Female: 19.0 ± 4.0 s (n = 4) *vs.* 7.5 ± 0.4 s (n = 12), 7.8 ± 0.4 s (n = 10), 10.5 ± 1.2 s (n = 6),

8.2 ± 0.5 s (n = 8), 1-way ANOVA, $F_{(4, 35)} = 13.9$, $p < 0.0001$]. An examination of gait indicated that

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice at P400, but not P210 need additional stabilization during ambulation, as they

spend twice as much time with 3 paws, rather than the normal 2, in contact with the ground as they

walk across the gait analysis platform [Male: 56.2 *vs.* 26.4 to 32.2 %, 1-way ANOVA, $F_{(4, 54)} = 14.3$,

$p < 0.0001$; Female: 58.4 *vs.* 18.9 to 28.8 %, 1-way ANOVA, $F_{(3, 178)} = 95.5$, $p < 0.0001$; **Fig. 3B**].

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ also display a slower cadence and average speed across the platform compared to

all other genotypes at P400 [cadence, Male: 9.5 *vs.* 13.3 to 15.9 steps/s, 1-way ANOVA, $F_{(3, 204)} = 36.8$,

$p < 0.0001$; Female: 9.1 *vs.* 14.2 to 15.9 steps/s, 1-way ANOVA, $F_{(3, 204)} = 39.7$, $p < 0.0001$; speed, Male:

8.8 *vs.* 22 to 26 cm/s, 1-way ANOVA, $F_{(4, 50)} = 28.3$, $p < 0.0001$; Female: 58.4 *vs.* 18.9 to 28.8 cm/s, 1-way

ANOVA, $F_{(3, 178)} = 39.7$, $p < 0.0001$; **Fig. 3B; Fig. 3-fig. S1**]. This difference in speed and cadence is

unlikely to be caused by animal size, as there are no significant differences in these parameters at

earlier time points when the difference in size is significant (**Fig. 2A**). These observations across the

two behavioral tests were found in both male and female mice at each of their respective time points,

consistent with the lack of sex differences observed in A-T patients.

We further examined behavioral differences between the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and $Atm^{+/+}$; $Aptx^{+/+}$ mice

using a standardized set of experimental procedures used to phenotype genetically modified mice (i.e.,

SHIRPA; **Fig. 3C**; **Fig. 3-fig. S1**) (Rogers et al. 1997). We first detected differences in motor function at

P8, where $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice took 3-4 times longer on average to right themselves compared to

$Atm^{+/+}$; $Aptx^{+/+}$ mice [Male: 6.4 ± 1.1 s (n = 24) *vs.* 1.5 ± 0.1 s (n = 23), *t*-test, p<0.0002; Female: $11.1 \pm$

1.9 s (n = 21) *vs.* 2.4 ± 0.3 s (n = 17), *t*-test, p<0.0002; **Fig. 3C bottom**]. At 30 days of age, we detected

significant differences between $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and $Atm^{+/+}$; $Aptx^{+/+}$ mice in behavioral tests that

qualitatively measure body position and spontaneous activity (**Fig. 3C**). Striking differences in

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ compared to $Atm^{+/+}$; $Aptx^{+/+}$ mice were observed at P400, especially for behaviors

related to movement, including locomotor activity, body position, and gait (**Fig. 3C**). The results from

this battery of tests demonstrates that $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice develop a severe change in behavior by

P400, consistent with purely visual observations of significant motor coordination deficits in the mice up

to this time point. Importantly, we do not find any significant differences between the other control

genotypes, including $Atm^{R35X/+}$; $Aptx^{-/-}$ mice that express at least some ATM but no APTX protein (**Fig.**

3-fig. S1).

2.4 Membrane and synaptic properties are perturbed in ATM- and APTX-deficient Purkinje

neurons

Purkinje neurons (PN) are a key neuronal subtype located in the cerebellar cortex. They display

considerable intrinsic excitability, firing action potentials spontaneously at rates significantly higher than

most other neurons in the brain (50 to 100 Hz more in many cases). Their activity shapes cerebellar

output via tonic inhibition of neurons of the cerebellar nuclei, which project to motor coordination

centers in the forebrain, brainstem, and spinal cord. Cerebellar PN dysfunction is associated with

several forms of ataxia and implicated in A-T (Hoxha et al. 2018; Cook, Fields, and Watt 2021; Shiloh

2020). We therefore examined if the electrophysiological properties of PNs in the $Atm^{R35X/R35X}$; $Aptx^{-/-}$

cerebellum were abnormal.

Since PN baseline activity and responsivity to input is mediated by a baseline set of passive and active

membrane properties (**Fig. 4**), we directly recorded from and compared the membrane properties of

PNs in acute cerebellar sections harvested from $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and $Atm^{+/+}$; $Aptx^{+/+}$ mice (P350 to

400). PNs recorded from $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice had significantly "tighter" membranes, displaying

higher membrane input resistances (R_m) than those from $Atm^{+/+}$; $Aptx^{+/+}$ mice [47.7 ± 5.6 (n = 15) *vs.*

30.2 ± 1.47 (n = 23) MΩ, *t*-test, p=0.008; **Fig. 4B**] and displayed a faster membrane time constant (τ)

[3.6 ± 0.4 (n = 15) *vs.* 5.1 ± 0.3 (n = 23) ms, t-test, p=0.009; **Fig. 4B**]. These results indicate that the total

membrane capacitance ($C_m = \tau/R_m$) of the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ PNs is significantly reduced [98.25 ±

19.23 (n = 15) *vs.* 175.6 ± 12.67 (n = 23) pF, *t*-test, p=0.0025; **Fig. 4B**]. At the cellular level, this

suggests that the ATM- and APTX-deficient PNs have less (i.e., decreased area) or thinner membranes

than that of wildtype PNs, a result suggestive of a developmental deficit or neurodegenerative process

(Dell'Orco et al. 2015). We next assessed the intrinsic excitability of PNs in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and

$Atm^{+/+}$; $Aptx^{+/+}$ mice by examining PN action potential (AP) generation and dynamics. Significant deficits

in the ability of PNs to fire continuously in response to current injection were observed in $Atm^{R35X/R35X}$;

$Aptx^{-/-}$ mice (**Fig. 4C**). These deficits were associated with significant perturbations in the amplitude,

threshold, and area of evoked action potentials [amplitude: 66.2 ± 0.7 (n = 14) *vs.* 72.1 ± 1.4 (n = 13)

ΔmV, *t*-test, p = 0.003; threshold: -55.2 ± 1.5 *vs.* -48.61 ± 1.9 mV, *t*-test, p=0.0196; area: 17.96 ± 0.6 *vs.*

20.63 ± 1.0 mV*ms, *t*-test, p=0.048; **Fig. 4C**]. Together, these experiments demonstrate significant

perturbations of PN physiological properties that likely disrupt their ability to function normally in the

cerebellum of $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice.

We next tested whether extrinsic and/or synaptic PN properties were also impacted in $Atm^{R35X/R35X}$;

$Aptx^{-/-}$ mice. We first examined spontaneous excitatory post synaptic currents (sEPSC) generated by

granule cell-to-PN synapses (i.e., parallel fiber inputs). No difference in sEPSC size was detected,

indicating the function of granule cell axon terminals (i.e. parallel fibers) was relatively normal in the

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ cerebellum [18.92 ± 1.3 (n = 11) *vs.* 23.4 ± 3.3 (n = 11) pA, *t*-test, p=0.477; **Fig. 4D**]

(Yamasaki, Hashimoto, and Kano 2006). sEPSC frequency, however, was found to be significantly

increased, a phenomenon that could be attributed to either an increase in the total number of synapses,

an increase in the size of the readily releasable pool of synaptic vesicles, or an increase in the

probability of neurotransmitter release in PNs of $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice [18.75 ± 2.8 Hz (n = 11) *vs.*

11.4 ± 1.0 Hz (n = 11), *t*-test, p=0.047; **Fig. 4D**]. We next explored evoked synaptic release and short-

term plasticity by simultaneously recording from PNs and electrically stimulating either granule cell (i.e.,

parallel fibers) or inferior olivary (i.e., climbing fiber) axons with a paired-pulse burst (2-pulses, 50 ms

apart). The synaptic properties of parallel fibers were found to be normal, displaying no significant

differences in the expected short-term facilitation (Atluri and Regehr 1996) or halfwidth and decay time

constant of the evoked EPSC [PPR: 1.3 ± 0.03 (n = 10) *vs.* 1.4 ± 0.05 (n = 13), *t*-test, p=0.162; halfwidth:

3.9 ± 0.6 *vs.* 4.9 ± 0.4 ms, *t*-test, p=0.175; time constant: 3.5 ± 0.5 *vs.* 4.7 ± 0.4 ms, *t*-test, p=0.054; **Fig.**

4E]. In comparison, we found climbing fiber-to-PN synaptic responses, which normally display pair-

pulse depression (Hansel and Linden 2000), to depress at significantly greater magnitudes in

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice [PPR: 0.6 ± 0.03 (n = 6) *vs.* 0.7 ± 0.02 (n = 9), *t*-test, p=0.03; **Fig. 4F**]. The

overall width and decay time constant of the evoked currents were also smaller [halfwidth: 2.3 ± 0.6 (n =

6) *vs.* 3.0 ± 0.2 (n = 9) ms, *t*-test, p=0.004; time constant (fast): 1.1 ± 0.14 *vs.* 2.9 ± 0.4 ms, t-test,

p=0.001]. While these results could be caused by a presynaptic deficit, such as reduced vesicle stores

in the climbing fiber axon terminal, the unaffected initial magnitude of the EPSC [2.4 ± 0.4 (n = 6) *vs.* 1.9

± 0.2 nA (n = 9), *t*-test, p=0.3] points to a more intrinsic deficit, such as a reduced Ca^{2+} influx from the

endoplasmic reticulum, which could significantly impact long-term synaptic plasticity critical to

cerebellar function (Hoxha et al. 2018; Kano and Watanabe 2017). Overall, the perturbations to the

passive and active PN properties that we observe here likely give rise to the significant cerebellar

dysfunction in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice.

2.5 ATM and APTX deficiency causes a progressive perturbation of PN neural activity that is

associated with dendritic shrinking and overall cerebellar atrophy

Decreased rates of spontaneous PN action potential firing, which can be indicative of PN dysfunction,

have been observed in several mouse models of ataxia, including spinocerebellar ataxias (SCA) 2, 3, 5,

6, 13, 27, several models of episodic ataxia (e.g., leaner, ducky, and tottering), and autosomal-

recessive spastic ataxia of the Charlevoix-Saguenay (Hourez et al. 2011; Hansen et al. 2013;

Dell'Orco, Pulst, and Shakkottai 2017; Kasumu and Bezprozvanny 2012; Liu et al. 2009; Perkins et al.

2010; Shakkottai et al. 2011; Jayabal et al. 2016; Stoyas et al. 2020; Hurlock, McMahon, and Joho

2008; Shakkottai et al. 2009; Bosch et al. 2015; Walter et al. 2006; Alviña and Khodakhah 2010; Ady et

al. 2018; Larivière et al. 2019; Cook, Fields, and Watt 2021). We therefore used this biomarker to

characterize the progression of PN perturbation in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice and assess whether deficits

were restricted to ATM- and APTX-deficient mice, consistent with the behavioral results (**Fig. 2,3**). We

additionally examined whether decreased PN activity differed across the cerebellum, as anecdotal

clinical pathology reports suggest degeneration may occur asymmetrically across the cerebellum, with

the anterior and posterior vermis and middle cerebellar hemispheres affected the most, although no

systematic analysis has been performed, and the consistency of results across patients is highly

variable (Verhagen et al. 2012; De Leon, Grover, and Huff 1976; Amromin, Boder, and Teplitz 1979;

Monaco et al. 1988; Terplan and Krauss 1969; Strich 1966; Solitare 1968; Solitare and Lopez 1967;

Aguilar et al. 1968a; Paula-Barbosa et al. 1983).

Using extracellular recording methods in the acute slice, we recorded spontaneous action potentials

from 3,300 PNs (**Fig. 4G**) across 188 animals, encompassing $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and 3 other

genotypes at four different time points (P45, 120, 210, and 400). We visually selected "healthy" cells

(see Methods) located deeper in the slice, that consistently fired during the extent of the 60 second

recording period. Qualitatively, tissue and cell quality did not visually differ across genotypes under DIC

microscopy. Cells were sampled in a distributed fashion across the lateral, intermediate, and medial

(vermis) cerebellum of each mouse to assess whether changes in PN firing activity was ubiquitous or

anatomically restricted. Regions were segregated based on gross anatomical domains in the mouse

defined by natural anatomical boundaries (e.g., foliation) and their general connectivity with different

regions of the nervous system (e.g., forebrain, brainstem, etc.) (Voogd and Glickstein 1998).

We found that complete deficiency of both ATM and APTX, consistent with the behavioral results, was

necessary to produce a significantly reduced spontaneous PN firing frequency (**Fig. 4G, H**). Although

the trend of slower PN firing rates was observed across most regions of the cerebellum, some

subregions appeared to be less or minimally impacted, including several areas of the lateral

cerebellum, including the paraflocculus, paramedian, and crus I and II (**Fig. 4-fig. S2**). Significant age

dependent changes in firing frequency were also only observed in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice (**Fig. 4H**),

with the most significant decline occurring between P120 and 210 [medial: 50.3 ± 2.4 Hz (n = 61) vs.

36.9 ± 2.2 Hz (n = 31), t-test, p=0.0006]. No significant difference in PN firing frequency was detected

between male and female mice within each genotype, thus the data were pooled (2-way ANOVA, p>0.3

across all pairwise comparisons; **Fig. 4-fig. S3**). Previous studies across several mouse models of

heritable ataxia, including episodic ataxia and several variants of spinocerebellar ataxia find that

physiological disruption in PN firing not only changes its frequency, but also its regularity (Kasumu and

Bezprozvanny 2012; Jayabal et al. 2016; Stoyas et al. 2020; Cook, Fields, and Watt 2021). We

compared both the coefficient of variation (CV) and variability in adjacent intervals (CV2) between

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ and control mice (**Fig. 4-figs. S4, S5**). No difference in these parameters across

sex, age, or genotype was detected. Consistent with the behavioral results, cerebellar dysfunction was

found only in the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice that developed ataxia and not in mice with partial or full

expression of ATM or APTX.

2.5 ATM and APTX deficiency induces cerebellar atrophy

In A-T patients, ataxia is usually detected between 1 to 2-years of age and is associated with little to no

cerebellar atrophy (Tavani et al. 2003; Taylor et al. 2015). Significant structural changes and atrophy

are usually first detected via neuroimaging between 5 to 10-years of age (Demaerel, Kendall, and

Kingsley 1992; Tavani et al. 2003). Postmortem clinical histopathology in A-T patients points to

significant changes in PN morphology and density, however, these reports primarily detail patients at

late stages of the disorder, and the relationship between the severity of PN pathology and ataxia is not

clear (Verhagen et al. 2012; De Leon, Grover, and Huff 1976; Amromin, Boder, and Teplitz 1979;

Monaco et al. 1988; Terplan and Krauss 1969; Strich 1966; Solitare 1968; Solitare and Lopez 1967;

Aguilar et al. 1968a; Paula-Barbosa et al. 1983; Gatti and Vinters 1985).

In the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice, we found the gross size of the cerebellum to be normal early in life, but

significant atrophy developed as the severity of ataxia increased (**Fig. 5A**). At early stages (P45-P210),

the size of the cerebellum in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice did not differ from mice with at least one copy of

the Atm gene (2-way ANOVA, $F_{(3,52)}= 1.0$, p=0.4). However, by P210, the overall size of the cerebellum

in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice was significantly reduced (1-way ANOVA, $F_{(3,37)}= 1.4$, p=0.3), with the

degenerative process continuing to at least the last time point investigated (P460). To rule out the

possibility that reduced cerebellar size was related to the smaller stature of $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice, we

examined, animal weight and cerebellar size and found no correlation [Pearson's correlation, p>0.3 for

all 4 genotypes at P460, n = 10 to 20]. Furthermore, we found that cerebellar size did not differ between

male and the on average 22% smaller female mice across genotypes at this age [2-way ANOVA, $F_{(2,}$

$_{153)}= 1.9$, p=0.2]. Therefore, cerebellar neurodegeneration in the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice, which begins

after P120, is correlated with ATM and APTX deficiency.

We found cerebellar atrophy to be associated with a selective reduction in the width of the molecular

layer where PN dendrites reside (**Fig. 5B**). Consistent with the temporal changes in gross cerebellar

size, PN firing frequency, and behavior, ML width in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice was normal in younger

mice but progressively decreased in width as the severity of ataxia increased [P400: 120.2 ± 2.1 μm (n

= 5) $vs.$ 140.2 ± 4.8 μm (n = 5), 2-way ANOVA, $F_{(1, 42)}= 45.04$, p<0.0001; **Fig. 5B, Fig. 5-figs. S1A**]. In

contrast, no difference in the width of the granular cell layer (GCL) across age was observed in the

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice [P400: 135.5 ± 2.4 μm (n = 6) $vs.$ 127.5 ± 4.3 μm (n = 5), 2-way ANOVA, $F_{(1,}$

$_{42)}= 3.3$, p=0.08; **Fig. 5B, Fig. 5-figs. S1A**]. Cerebellar atrophy, however, was not due to PN cell death,

as PN density did not significantly differ between $Atm^{R35X/R35X}$; $Aptx^{-/-}$ and $Atm^{+/+}$; $Aptx^{+/+}$ mice [P400: 4.5

± 0.3 (n = 9) $vs.$ 4.5 ± 0.2 (n = 7) PNs/4000 μm^2, Welch's t-test, p=0.9; **Fig. 5C,D Fig. 5-figs. S1B**].

Moreover, we found no evidence from immunohistolological experiments for increased levels of

programmed cell death or microglial activation in the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice (**Fig. 5-figs. S2**). At the

anatomical level, we found pathological changes in PN morphology, as somatic size was reduced (15.3

$\pm$ 0.3 (n = 5) *vs. 17.8 $\pm$ 0.1* (n = 5) μm, Welch's *t*-test, p=0.0004; **Fig. 5E**) and the primary and secondary

dendrites were abnormally large in caliber in the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice (3.1 $\pm$ 0.2 (n = 6) *vs.* 2.8 $\pm$ 0.1

(n = 6) μm, Welch's *t*-test, p=0.003; **Fig. 5F, Fig. 5-figs. S2C**). Overall, we found a good correlation

between the abnormal structural and electrophysiological properties of the cerebellum and the

progression of motor and deficits.

2.6 Differential disruption of thymocyte development in ATM-deficient vs. APTX-deficient mice

Chronic sinopulmonary infections associated with immunodeficiency are one of the leading causes of

death in A-T patients (Morrell, Cromartie, and Swift 1986; Bhatt and Bush 2014). Immunodeficiency is

linked to deficits in the generation of B- and T-lymphocytes that have been linked to defects in the

antigen receptor gene rearrangement processes during the generation of these cells in the bone

marrow and thymus, respectively (Staples et al. 2008). The resulting defects in mature lymphocyte

numbers include decreases in CD4[+] helper T-cells and killer CD8[+] T-cells (Schubert, Reichenbach, and

Zielen 2002). We therefore examined the percentages of T-cells in peripheral blood and of different

subpopulations in the thymus of Atm[R35X/R35X]; Aptx[-/-] mice using T-cell antigen receptor (TCR) and

CD4/CD8 co-receptor expression.

In the peripheral blood, we observed a significant reduction in the total fraction of CD3[+] T-cells in mice

with reduced or absent ATM expression compared to wildtype mice (**Fig. 6**). This reduction was further

compounded by concomitant deficiency of APTX. ATM and APTX deficiencies reduced T-cells in

peripheral blood by over 65% compared to wild type controls. The effect of APTX deficiency was

additive to that of ATM deficiency, suggesting a different mechanism of action for each of these two

proteins on T-cell generation. The reduction in the percentage of T-cells in peripheral blood was mostly

associated with reduction in the CD4+ helper T-cell population (**Fig. 6B**). Of interest, the proportion of

CD8+ T-cells was increased only in Atm[R35X/R35X]; Aptx[-/-] mice (**Fig. 6B**). Again, we observed a

differential effect of ATM and APTX deficiencies as seen for the effects of these mutations on the total

T-cell fraction.

Given the reduction in T-cell populations in the blood, we next assessed T-cell development in the

thymus. In this organ, bone marrow-derived T-cell progenitors undergo TCR gene rearrangement

followed by positive selection for MHC restriction and negative selection of autoreactive clones. The

phases of thymocyte development can be followed by monitoring expression of CD4 and CD8

expression in thymocytes. The progression of this developmental program goes from double negative

(CD4$^-$CD8$^-$) thymocytes to double positive (CD4$^+$CD8$^+$) thymocytes and then to single positive (CD4$^+$ or

CD8$^+$) thymocytes. In addition, within the double negative stage, four different subpopulations can be

identified, based on the expression of CD25 and CD44, known as DN1 (CD44$^+$CD25$^-$), DN2

(CD44$^+$CD25$^+$), DN3 (CD44$^-$ CD25$^+$) and DN4 (CD44$^-$CD25$^-$) (Germain 2002).

Gene rearrangement during thymocyte development occurs twice—once at the double negative

thymocyte stage in the CD25$^+$CD44$^-$ stage (Krangel 2009) and then again in double positive thymocyte

stage before progressing into separate CD4$^+$ and CD8$^+$ single positive populations (Livák et al. 1999).

ATM deficiency has been linked to defects in both bouts of rearrangement in mice (Vachio 2007,

Hathcock 2013). Therefore, we compared the proportion of cells in the thymus expressing these

different developmental cell surface markers in our ATM-deficient and control mice (**Fig. 7**).

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ and $Atm^{R35X/+}$; $Aptx^{-/-}$ but not $Atm^{R35X/R35X}$; $Aptx^{+/+}$ mice had significantly elevated

proportions of CD44$^+$CD25$^-$, CD44$^+$CD25$^+$, and CD44$^-$CD25$^+$ cells compared to wildtype (**Fig. 7A**).

These increased proportions appear to be due in part to an impediment of CD44$^-$CD25$^+$ cells maturing

into CD44$^-$CD25$^-$ double negative cells, as the fraction of CD44$^-$CD25$^-$ cells from $Atm^{R35X/R35X}$; $Aptx^{-/-}$

and $Atm^{R35X/+}$; $Aptx^{-/-}$ mice is significantly lower than wildtype (**Fig. 7A**). Of interest, APTX deficiency by

itself had the greatest effect on the loss of DN4 cells, suggesting that APTX deficiency, rather than ATM

deficiency, is responsible for this effect. To our knowledge, this finding implicates for the first time

APTX in gene rearrangement during the process of TCRβ recombination.

Next, we looked at the proportions of CD4$^+$CD8$^+$ thymocytes compared to CD4$^+$CD8$^-$ and CD4$^-$CD8$^+$

single positive thymocytes in these four different strains. In agreement with our results in the blood and

prior studies, we found that ATM-deficient mice but not control mice displayed decreased expression of

CD4$^+$CD8$^-$ and CD4$^-$CD8$^+$ single positive thymocytes (**Fig. 7B**). These results support the role of ATM

in TCR α/δ gene rearrangement during thymocyte development (Bredemeyer et al. 2006), a role that is

independent of the role played by APTX in early thymocyte maturation.

2.7 Read-through molecules overcome PTC to restore ATM expression

Our primary rationale for inserting a clinically relevant nonsense mutation in the *Atm* gene was to

generate a mouse amenable to critical pre-clinical testing of a novel set of SMRT compounds. We

previously demonstrated SMRT compounds recover production of ATM protein in A-T patient derived

lymphoblastoid cell lines by overcoming premature termination codons (PTC) caused by nonsense

mutations (Du et al. 2013). To demonstrate suitability of this new A-T animal model for SMRT

compound testing we chose to directly examine their ability to restore ATM expression using an explant

approach that circumvents challenges related to *in vivo* delivery (e.g., bioavailability, route of delivery,

etc.). ATM expression was measured in samples from the spleen, where ATM is normally expressed at

high levels, and the cerebellum, a key target tissue for the disorder. We exposed these explant tissues,

harvested from homozygous *Atm^{R35X}* and *Atm^{Q35X}* mice with either a candidate SMRT compound

(GJ103), or an aminoglycoside previously known to have read-through properties (G418) and then

measured ATM expression by immunoblot to assess restoration. In both types of ATM deficient mice,

ATM expression was consistently restored in the spleen and cerebellum by both G418 and GJ103 (**Fig.**

8). These results demonstrate that our SMRT compounds can enable read-through of at least 2 of the 3

possible nonsense mutations causing PTCs and provide the rationale for *in vivo* efficacy testing in

follow-on studies.

3.0 Discussion

By increasing genotoxic stress through the addition of a secondary hit to the DDR pathway, we

generated a novel mouse model that displays the most comprehensive set of A-T symptoms of any

model to date. This includes a severe and progressive ataxia associated with cerebellar atrophy and

perturbations of PN properties along with a high incidence of cancer and defects in immune cell

development. Together, these comorbidities encompass the three leading causes of premature death in

A-T—each contributing to roughly a third of deaths. Of these, the incapacitating effect of ataxia is the

most penetrant and is reported by patients and caregivers as having the greatest impact on their quality

of life. For this reason, the presence of ataxia and cerebellar atrophy in this new mouse model is of

great significance, as it provides for the very first time a resource to not only elucidate the mechanisms

of neurological dysfunction, but also a critically needed *in vivo* model to test severely needed A-T

therapeutics, such as the read-through compounds we describe here.

We found several similarities between the overall progression of ataxia in the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice

and A-T patients. In clinical A-T, motor deficits are observable by roughly 2 years of age, when parents

and doctors detect a lowered ability to transition from toddling to a smooth, reflexively coordinated

gate—unfortunately, little is known about motor defects at earlier stages due to the diseases low

prevalence and current lack of early diagnostic testing (Rothblum-Oviatt et al. 2016). Patients usually

learn to walk without assistance and neurological symptoms tend to remain stable through the first 4 to

5 years of life (Rothblum-Oviatt et al. 2016). We found a similar early progression of motor deficits in

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice, detecting early mild motor deficits at P8 (righting reflex deficit), followed by a

period of relative stability, prior to onset of a progressive and severe ataxia developing after p210 that

included changes in gait, startle reflex, tremor, and locomotor activity. Several important questions arise

out of these findings, including whether ATM and/or APTX have a neurodevelopmental role in the

cerebellum. Future studies focused on the early phase of the disorder will be critical in understanding if

the cerebellum develops normally prior to dysfunction or whether developmental defects are an initial

cause. We also found, similar to A-T patients, that the severity of the late-developing ataxia was

variable, with some mice ambulating with a clumsy, high-stepping rear gate (**Video 3**) and others

moving almost entirely via contortion of the rear trunk (**Fig. 1E** and **Video 4**) (Rothblum-Oviatt et al.

2016; Levy and Lang 2018; Boder and Sedgwick 1958). Overall, we found that $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice

developed a visually profound and measurable progressive loss in motor coordination similar to that

observed in A-T patients, which was rescued by expression of at least one copy of the *Atm* or *Aptx*

gene.

The loss of motor coordination in A-T has been attributed to cerebellar degeneration due to its relatively

selective neuropathology across the brain and its causal role in several different forms of ataxia (Hoche

et al. 2012). Consistent with A-T patient neuroimaging studies (Wallis et al. 2007; Sahama et al. 2015;

Sahama et al. 2014; Dineen et al. 2020; Tavani et al. 2003; Quarantelli et al. 2013), we find that

cerebellar size in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice is initially normal, but progressively atrophies concurrently

with changes in neurological function. While loss of cerebellar tissue has been considered a main

cause of ataxia in humans, it is unclear from clinical data if ataxia severity is a good predictor of the

extent of cerebellar degeneration found postmortem (Aguilar et al. 1968b; Crawford et al. 2006; Dineen

et al. 2020). In the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice, we find clear atrophy associated with thinning of the

Purkinje neuron dendrite layer that precedes the late, severe behavioral deficits. Our histological

observations in the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice suggest that changes in cerebellar function itself, rather

than profound loss of cerebellar cells, are sufficient to cause the ataxic phenotype, consistent with the

observation of behavioral defects prior to significant PN loss in several SCAs (Shakkottai et al. 2011;

Lorenzetti et al. 2000; Clark et al. 1997; Jayabal et al. 2016).

The reason why ATM and APTX deficiency is required to generate ataxia in mice, when loss of either is

sufficient to cause ataxia in humans, remains unclear. One possibility is that the rodent brain may more

flexibly utilize compensatory pathways or redundant proteins while responding to the 10-20k DNA

lesions that impact cells each day (Lindahl and Barnes 2000). Several forms of DNA repair exist to

potentially meet this challenge, including base excision repair (BER), nucleotide excision repair (NER),

as well as homologous and non-homologous end joining (HEJ and NHEJ, respectively), all of which

ATM and APTX have been implicated in (Chou et al. 2015; Çaglayan et al. 2017; Wakasugi et al. 2014;

Tumbale et al. 2018; Chatterjee and Walker 2017). Alternatively, it may be the case that deficiency in

ATM or APTX alone does not adequately impact cell health during the mouse's comparatively short

lifespan, and thus eliminating both proteins is necessary to achieve sufficient accumulation of DNA

damage to manifest over this time period. This possibility is strengthened by the fact that ATM and

APTX have distinct biochemical properties and functional roles in the DNA damage response, and

therefore deficiency in both would be predicted to cause a broader hit to genome stability (i.e.,

increased genotoxic stress).

Our finding that two genome stability pathway proteins are required to induce neurological defects in

mice strongly suggests that it is the loss of ATM's role in DNA repair, rather than potential functions in

oxidative stress signaling, mitophagy, or mitochondrial function that cause the cerebellar defects

(Shiloh 2020). Alternatives, however, cannot be completely ruled out, as APTX, like ATM, has been

observed within the mitochondria of brain cells, where it is thought to support the processing of

mitochondrial DNA (Meagher and Lightowlers 2014; Sykora et al. 2011). This new mouse model

provides a new tool to explore these possibilities and mechanistically define how loss of ATM and

APTX ultimately causes cerebellar dysfunction.

The biophysical perturbations observed in PNs recorded from the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice are similarly

found in several other mouse models of ataxia. This includes changes we observed in PN input

resistance, membrane capacitance, and AP threshold and width, which have also been described in

mouse models of SCA like 1, 3, and 7 (Stoyas et al. 2020; Shakkottai et al. 2011; Dell'Orco et al. 2015).

Moreover, the progressive reduction in PN action potential firing frequency we report, which positively

correlates with the development of ataxia in the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice, is reported in a large number

of ataxic mouse models, including SCAs 1, 2, 3, 5, 6, and 13 as well as a few episodic forms (see

review (Cook, Fields, and Watt 2021)).

Given the significant overlap in PN perturbations observed across many different ataxias caused by

distinct cellular defects, restoring PN AP firing frequencies has been considered as a broad-based

therapeutic approach. However, it remains unclear whether reduced PN firing is a causal factor of

ataxia. Moreover, experimental evidence suggests changes in PN activity may in fact be a generalized

response to maintain homeostasis during ongoing disease-related impairment of PN physiology

(Dell'Orco et al. 2015). Thus, continued efforts across all cerebellar ataxias are needed to link the

genetic, molecular, and cellular disruptions caused by disease to the specific changes in cerebellar

neural signaling that ultimately generates the ataxia. Of significant importance in this effort will be

determining whether disease-causing cerebellar defects commonly or differentially cause ataxia

through a loss of cerebellar function (e.g., loss of coordinating signals during movement), or from a

dominant negative effect (e.g., disrupting downstream neural circuits with abnormal neural output

patterns). Ultimately, while a common therapeutic strategy to address cerebellar ataxias would have the

greatest impact, a directed approach that addresses the distinct genetic and molecular causes of

cellular dysfunction may ultimately be necessary to successfully develop an efficacious therapeutic.

The mechanistic link between deficiency in DNA stability proteins like ATM and APTX and PN

dysfunction is far from clear. Our results suggest the effect of ATM and APTX loss on PNs is intrinsic,

as we do not find changes in the presynaptic properties of granule cells or evidence of their cellular loss

(no change in GCL thickness). Moreover, while we observed differences in short term plasticity of

inferior olivary inputs in ATM- and APTX-deficient PNs and wildtype, these results likely point to a

disruption in Ca^{2+} homeostasis potentially via reductions in Inositol 1,4,5-triphosphate receptor 1 (*Itpr1*)

expression, similar to those observed in SCAs 1, 2, and 3 mouse models as well as ATM-deficient mice

(Kim et al. 2020; Chen et al. 2008; Demirci et al. 2009; Shakkottai et al. 2011). While this provides a

promising avenue for future examination and comparison, it is as of yet unclear, even for the SCAs,

whether changes in Ca^{2+} homeostasis, is the causal factor or just another symptom or even

compensatory response of diseased or disturbed PNs (Dell'Orco et al. 2015).

In the immune system, ATM is implicated in the repair of DNA breaks that naturally occur during gene

rearrangement of antigen receptor genes in B- and T-cell precursors, a phenomenon critical for antigen

receptor (Ig and TCR) diversity of these cells. Our finding that T-cell proportions in the blood are

significantly reduced is consistent with prior studies in humans and A-T knockout mice (Schubert,

Reichenbach, and Zielen 2002; Hathcock et al. 2013; Chao, Yang, and Xu 2000; Barlow et al. 1996).

This reduction of T-cells in the periphery likely correlates with a defect in both cellular and humoral

immunity. Importantly, we found that expression of one copy of the ATM gene is enough to restore

CD4+ deficits in the blood indicating that therapies able to restore partial ATM expression would have

therapeutic efficacy. Although we have not assessed B-cell development in this paper, it is likely that

similar conclusions would apply to that process given their mechanistic similarities (Marshall et al.

2018).

As expected, the reduction of T-cells in peripheral blood is correlated with defective thymocyte

development. In the thymus, we found two main defects. One, induced primarily by APTX deficiency,

manifests as a defect in the DN3 to DN4 transition coinciding with early rearrangement of TCR β locus.

The other defect, primarily caused by ATM deficiency, correlates with decreased progression of double

positive CD4$^+$CD8$^+$ to single positive cells, primarily CD4$^+$ thymocytes. While the APTX finding was

surprising, as its deficiency (AOA 1) is not associated with immune deficits, APTX is known to interact

with TCR β gene rearrangement proteins, including XRCC4 (Clements et al. 2004). Future studies

aimed at defining APTX's role in end-joining mechanisms during TCR gene rearrangement will be

important, and the possibility that alternative end-joining mechanisms, like the use of microhomologies

account for the lack of an immune deficit in its absence needs further investigation (Bogue et al. 1997).

The survivability of $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice is considerably longer than prior A-T mouse models. In

comparison, the first A-T KO mouse model reported by Barlow et al. died from thymomas usually within

2-4 months after birth (Barlow et al. 1996). The decreased cancer survivability in this and many other

knockout A-T mouse models is likely genetic, as the background strain harboring the mutation has

been shown to have significant effects on cancer prevalence and survivability, with A/J and C57BL/6

backgrounds having significantly increased survivability over the BALBC and 129S strains (Genik et al.

2014). The fact that our ATM deficient mice were created on a C57BL/6 background likely underlies

their comparatively long lifespan. Given that the $Atm^{R35X/R35X}$; $Aptx^{+/+}$ mice do not develop ataxia, it is

unlikely that the early death in A-T KO mice prevents observation of an ataxic phenotype that would

otherwise develop in these mice. However, it is unknown whether the C57BL/6 background confers a

resilience to developing ataxia, as it does for cancer. Defining the genetic or possibly epigenetic factors

that influence the severity of the disease could provide avenues for future therapeutic development.

Given the global nature of the ATM and APTX null mutation in our mouse model, we cannot entirely

rule out that extra-cerebellar defects may also contribute to the severe ataxic phenotype, and thus

future examination outside the cerebellum in the forebrain, brainstem, spinal cord, and even muscle will

need to be conducted. Within the cerebellum, while we found some anatomical differences in the PN

firing properties within different regions of the cerebellum, we did not detect regional differences in ML

width or PN density. However, there are challenges in using regional anatomy as a grouping factor in

the cerebellum, as the physical folds of the tissue do not necessarily correlate with the boundaries of

functional, molecular expression, or physiological property domains that have been described (Apps

and Hawkes 2009; Tsutsumi et al. 2015; Gao, van Beugen, and De Zeeuw 2012; Zhou et al. 2014).

Experiments focused on examining the extent of cerebellar defects within these domains will be

important in future studies and compared to the anecdotal reports of anatomical differences in A-T

patients (Verhagen et al. 2012; De Leon, Grover, and Huff 1976; Amromin, Boder, and Teplitz 1979;

Monaco et al. 1988; Terplan and Krauss 1969; Strich 1966; Solitare 1968; Solitare and Lopez 1967;

Aguilar et al. 1968a; Paula-Barbosa et al. 1983).

While we detect two potential stages in the progression of ataxia in the $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice, the

later stage of severe ataxia develops in adulthood in mice, as compared to the childhood onset in

humans. This may limit its use in some neurodevelopmental studies. Also, the interpretation of future

experiments must carefully factor in the fact that this new model expresses null mutations in two

genome stability genes at the same time, a situation that has not been detected in human patients with

either A-T or AOA1.

Finally, pinpointing where, when, and how ATM deficiency causes cerebellar pathology and ataxia has

been a challenge, as prior ATM-deficient mice generally lack the characteristic features needed to

causally link cellular and molecular deficits to the ataxic phenotype. Multiple promising avenues of

investigation have been defined, including those focused at the neuronal level, where ATM is implicated

in oxidative stress signaling (Chen et al. 2003) and synaptic function (Li et al. 2009; Vail et al. 2016), as

well as glial function, where recent evidence suggests glial pathology may be a leading factor in

cerebellar pathology (Kaminsky et al. 2016; Campbell et al. 2016; Petersen, Rimkus, and Wassarman

2012; Weyemi et al. 2015). This novel animal model provides a new tool to test mechanistic

hypotheses regarding how ATM deficiency causes cerebellar pathology and ataxia. Additionally, this

model may serve most importantly as a critical preclinical tool for testing previously proposed

therapeutic candidates (Browne et al. 2004; Chen et al. 2003) and our own SMRT compounds (Du et

al. 2013). The severe limitations of not having a suitable preclinical model for therapeutic testing,

especially for a rare disorder like A-T and AOA1, cannot be overstated.

4.0 Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Strain, strain background (*Mus musculus*)	Atm^{R35X}, $Atm^{Tm1.1(103CAG)TGA)Mfgc}$	This paper	103C>T mutation, human exon replacement	Generated by Hicks laboratory. Has been backcrossed into C57b/6 9 times. Contact pmathews@lundquist.org
Strain, strain background (*Mus musculus*)	Atm^{Q35X}, $Atm^{Tm1.1(103C)T)Mfgc}$	This paper	103C>T mutation, targeted premature termination signal in the mouse codon	Generated by Hicks laboratory. Has been backcrossed into C57b/6 9 times. Contact pmathews@lundquist.org

Strain, strain background (*Mus musculus*)	*Aptx*[-/-]	Ahel et al. 2006	MGI Cat# 3687171, RRID:MGI:3687171	Contact peter.mckinnon@stjude.org
Gene (*Mus musculus*)	*Atm*	MGI	MGI:107202; C030026E19Rik; ENSMUSG00000034218	
Gene (*Homo Sapien*)	*ATM*	OMIM	OMIM: 607585 MGI: 107202 HomoloGene: 30952; ENSG00000149311	
Sequence-based reagent	Atm gene	Transnetyx	PCR primers	F-5'-CCTTTGAGGCATAAGTTGCAACTTG-3'
Sequence-based reagent	Atm gene	Transnetyx	PCR primers	R- 5'-GTACAGTGTATCAGGTTAGGCATGC-3'
Chemical compound/ drugs	GJ103 salt Formula: $C_{16}H_{14}N_4O_3S$	TargetMol	T3448; CAS No. : 1459687-96-7	100 μM in media
Antibody	Anti-mouse CD68 (Rat monoclonal)	Bio-Rad	Cat# MCA1957, RRID: AB_322219	IF (1:400)
Antibody	ATM (D2E2) (Rabbit-monoclonal)	Cell Signaling Technology	Cell Signaling Technology Cat# 2873, RRID:AB_2062659	WB(1:500) WB(1:1000)
Antibody	GAPDH (14C10) (Rabbit-monoclonal)	Cell Signaling Technology	Cell Signaling Technology Cat# 2118, RRID:AB_561	WB(1:4000)

				053
Antibody	β-Actin (D6A8) (Rabbit-monoclonal)	Cell Signaling Technology	Cell Signaling Technology Cat# 8457, RRID:AB_109 50489	WB(1:5000)
Antibody	Anti-Rabbit IgG,HRP-linked (Goat-monoclonal-polyclonal)	Cell Signaling Technology	Cell Signaling Technology Cat# 7074, RRID:AB_209 9233	WB(1:5000)
Antibody	Anti-Calbindin (D-28k) (Rabbit, polyclonal)	Swant Inc.	Swant Cat# CB 38, RRID:AB_100 00340	IF (1:1000)
Antibody	Anti-Mouse Alexa Fluor 488 (Goat polyclonal)	ThermoFisher Invitrogen	Cat# A11001, RRID: AB_2534069	IF (1:500)
Antibody	Anti-mouse Cleaved Caspase-3, Asp-175 (rabbit)	Cell Signaling Technology	Cat# 9961, RRID: AB_2341188	IF (1:200)
Antibody	Anti-Rat Alexa Fluor 555 (Goat polyclonal)	ThermoFisher Invitrogen	Cat# A21244, RRID: AB_2535812	IF (1:1000)
Antibody	Anti-Rabbit Alexa Fluor 647 (Goat polyclonal)	ThermoFisher Invitrogen	Cat# A-21434, RRID: AB_141733	IF (1:500)
Antibody	Anti-Calbindin D-28k (mouse-monoclonal)	Swant Inc.	Cat# CB300	IF (1:500)
Antibody	Anti-Rabbit Alexa Fluor 488 (Goat-polyclonal)	ThermoFisher Invitrogen	Thermo Fisher Scientific Cat# A-11034, RRID:AB_257 6217	IF(1:1000)
Antibody	CD4 (GK1.5) (Rat-monoclonal)	ThermoFisher Invitrogen	Thermo Fisher Scientific Cat# 50-0041-82,	FACS (5 ul per test)

			RRID:AB_106 09337	
Antibody	CD8 (53-6.7) (Rat-monoclonal)	ThermoFisher Invitrogen	Thermo Fisher Scientific Cat# 53-0081-82, RRID:AB_469 897	FACS (5 ul per test)
Antibody	CD3 (145-2C11) (Hamster-monoclonal)	ThermoFisher Invitrogen	Thermo Fisher Scientific Cat# 12-0031-83, RRID:AB_465 497	FACS (5 ul per test)
Antibody	CD44 (IM7) (Rat-monoclonal)	ThermoFisher Invitrogen	Thermo Fisher Scientific Cat# 25-0441-82, RRID:AB_469 623	FACS (5 ul per test)
Antibody	CD25 (PC61.5) (Rat-monoclonal)	ThermoFisher Invitrogen	Thermo Fisher Scientific Cat# 47-0251-82, RRID:AB_127 2179	FACS (5 ul per test)
Other	Eosin Y (Certified Biological Stain)	Thermofisher (Fisher Chemical)	Cat# E511-100	
Other	Hematoxylin Stain Solution, Modified Harris Formulation, Mercury Free Nuclear Stain	RICCA Chemical Company	Cat# 3530-16	
Other	Permount Mounting Medium	ThermoFisher (Fisher Chemical)	Cat# SP15-100	
Other	Fluoromount-G with DAPI	Southern Biotech	Cat# 0100-20, RRID: SCR_021261	
Commercial assay or kit	BCA Protein Assay Kit	ThermoFisher Pierce	Cat# 23225	Protein Assay

Commercial assay or kit	SuperSignal West Pico Chemiluminescent Substrate	ThermoFisher Pierce	Cat# 34580	Chemiluminescent Substrate
Commercial assay or kit	Radiance plus	Azure Biosystems	Cat# AC2103	Chemiluminescent Substrate
Software, algorithm	FlowJo	https://www.flowjo.com/solutions/flowjo	RRID:SCR_008520	
Software, algorithm	ImageJ software	ImageJ (http://imagej.nih.gov/ij/)	RRID:SCR_003070	Version 1.53
Software, algorithm	IgorPro	http://www.wavemetrics.com/products/igorpro/igorpro.htm	RRID:SCR_000325	Version 7; Tarotools procedures
Software, algorithm	Neuroexpress	https://www.researchgate.net/project/NeuroExpress-Analysis-software-for-whole-cell-electrophysiological-data	https://www.researchgate.net/project/NeuroExpress-Analysis-software-for-whole-cell-electrophysiological-data	Version 21.1.13; Used for sEPSC analyses
Software, algorithm	GraphPad, Prism	GraphPad Prism (https://graphpad.com)	RRID:SCR_015807	Versions 8 and 9
Software, algorithm	MBF, Stereo investigator	https://www.mbfbioscience.com/stereology	RRID:SCR_017667	Version 2021
Software, algorithm	Microsoft Excel	https://www.microsoft.com/en-us/microsoft-365/excel	RRID:SCR_016137	Version 365
Software, algorithm	Catwalk XT	https://www.noldus.com/catwalk-xt	RRID: SCR_021262	

4.1 Ethics Statement

This study was performed in strict accordance with the recommendations in the Guide for the Care and

Use of Laboratory Animals of the National Institutes of Health. All the animals were handled according

to approved Institutional Animal Care and Use Committee (IACUC) protocols at The Lundquist Institute

(31374-03, 31773-02) and UCLA (ARC-2007-082, ARC-2013-068). The protocol was approved by the

Committee on the Ethics of Animal Experiments of the Lundquist Institute (Assurance Number: D16-

00213). Every effort was made to minimize pain and suffering by providing support when necessary

and choosing ethical endpoints.

4.2 Mice

All mice were group housed and kept under a 12-h day/night cycle with food and water available *ad*

libitum. Animals were housed within the general mouse house population, and not in specialized

pathogen-free rooms. Older animals were made available wetted food or food gel packs on the ground

of the cages as ataxia developed. *Atm^{R35X}* and *Atm^{Q35X}* mice were created and provided by Dr. Hicks

and colleagues at the University of Manitoba.

These mice were created to contain the c.103C>T mutation found in a large population of North African

AT patients using recombineering Gateway technology and site-directed mutagenesis. A C>T mutation

at this position in the mouse *Atm* gene creates a TAG G stop codon. The same mutation in the human

ATM gene produces a TGA G stop codon. In consideration of the use of these models for therapeutic

interventions, we chose to create a mouse model for each of the two PTC codons (**Fig. 1A**).

A modified Gateway R3-R4-destination vector was used to pull out the desired region of the mouse *Atm*

gene from a Bacterial Artificial Chromosome (BAC) and subsequently mutated to create either a TAG G

stop codon at codon 35 (M00001, position 103 (C>T)) or a TGA G stop codon (M00002, position 103

(CAG>TGA), replicating the human AT PTC). The genomic alleles were then cloned into a modified

version of the NorCOMM mammalian targeting vector using a 3-way Gateway Reaction (Bradley et al.

2012). The resulting targeting vectors were electroporated into C2 ES cells (C57Bl/6N, derived in A.

Nagy lab, Toronto, Canada) and successfully targeted clones were identified by selection with G418

(Gertsenstein et al. 2010). Integration of the mutated targeting cassette into the *Atm* gene locus was

confirmed by Southern blot, and by sequencing of PCR products to confirm the presence of the *Atm*

PTC mutation, error free targeting into the *Atm* locus, and error-free functional components of the

vector (data not shown). Positive ES clones were used for blastocyst injection to obtain the transgenic

lines. The transgenic allele contained a floxed human beta actin promoter - delta TK1- Neo cassette in

the intron upstream of the region containing the mutated exon. This floxed cassette was subsequently

excised by crossing with a Cre driver mouse (B6.C-Tg(CMV-cre)1Cgn/J) to generate *Atm*$^{R35X/+}$ and

Atm$^{Q35X/+}$ (MGI nomenclature: *Atm*$^{TM1(103CAG>TGA)MFGC}$ and *Atm*$^{TM1(103C>T)MFGC}$, respectively) mouse lines

(**Fig. 1A**). Genotyping of the two *Atm* lines was performed by using the following primers at Tm 62°C:

Atm gene forward (F) primer: 5'-CCTTTGAGGCATAAGTTGCAACTTG-3'; and *Atm* gene reverse (R)

primer: 5'-GTACAGTGTATCAGGTTAGGCATGC-3', creating a wild-type allele product of 151bp or

targeted allele product of 241bp (**Figs. 1A, 1B**).

Atm^{R35X} and *Atm*^{Q35X} were back-crossed with C57Bl/6J mice for 9 generations (99.2% isogenic) prior to

cryopreservation and subsequent rederivation using C57Bl/6J surrogate mothers. *Atm*^{R35X} and *Atm*^{Q35X}

breeders were obtained from F1 sibling *Atm*$^{R35X/+}$ and *Atm*$^{Q35X/+}$ mice. *Atm*$^{R35X/R35X}$ and *Atm*$^{Q35X/Q35X}$ were

both found to be fertile. *Aptx* knockout (*Aptx*$^{-/-}$) mice were created and provided to Dr. Mathews as

embryos from Dr. McKinnon (Ahel et al. 2006), and subsequently rederived via C57Bl/6J surrogate

mothers. *Aptx*$^{-/-}$ mice are on a C57Bl/6 and 129 mixed background. *Atm*$^{R35/R35XX}$; *Aptx*$^{-/-}$ mice of various

wildtype, heterozygous, and homozygous combinations were created from *Atm*$^{R35X/+}$; *Aptx*$^{+/-}$ breeders

generated by crossing *Atm*$^{R35X/R35X}$ and *Aptx*$^{-/-}$ mice. One cohort of double mutant and corresponding

control mice were used in the longitudinal behavioral study for gait analyses and SHIRPA testing (**Figs.**

2, **3**). Multiple additional cohorts of age-matched double mutant and control mice were used for

electrophysiological, immunohistological, and Vertical Pole test experiments (**Figs. 4**, **7**). Immunological

and protein expression experiments were carried out using mice bred from the original *Atm*^{R35X} and

Atm^{Q35X} rederived mice (**Figs**. **5**, **6**, and **8**).

Genotyping was performed from ear tissue samples of P8-11 mice. Real-time PCR methods conducted

by Transnetyx Inc. were used to determine each animals' genotype. Animals were made identifiable via

toe tattoos given at the same time as ear biopsy. Unique primers for Atm^{R35X} and Atm^{Q35X} were

quantified and used to identify wildtype, heterozygous and homozygous mice (listed above). $Aptx^{-/-}$ and

$Aptx^{+/+}$ primers were used to assess their genotypes.

4.3 Animal Health

Animals were weighed via a digital scale at P8, 45, 120, 210, 400. Animal death was recorded as the

day found dead, or on the day of euthanization when the animals reached a humane endpoint (animal

unable to right itself within 60s, significant hair matting indicating lack of self-grooming, or excessive

distress as noted by the veterinary staff). Animal carcasses were immediately frozen upon death, and

postmortem necropsies were carried out in batch. Probable cause of death was determined to the best

of our ability in collaboration with the staff veterinarian (Dr. Catalina Guerra) by visual inspection of the

internal organs. Some mice were cannibalized or accidentally disposed of by vivarium staff and were

therefore labelled as "missing." Mice with no discernable visual cause of death were labelled

"indeterminable." Mice that were found with thoracic masses near where the thymus would normally be

in young mice were listed as "thymic cancer." All other identified probable causes of death (e.g.,

enlarged livers, urinary blockage) were labelled "other."

4.4 Behavior

Before performing any behavioral test, mice were acclimated to the behavioral suite for ~20 minutes.

Mice were tested at varying times of the day, in line with their day cycle. A battery of behavioral tests

was performed on naïve double mutant mice of the indicated genotypes at various time points

depending on the behavior but in the same cohort of mice. The battery of tests included Catwalk Gait

assessment (P45, 120, 210, 400) and a subset of the SmithKline-Beecham Harwell Imperial-College

and Royal-London-Hospital Phenotype Assessment (SHIRPA) tests (P30 and 400). These tests were

conducted by the UCLA Behavioral Core. Double mutant and control mice were additionally examined

on the Vertical Pole test. All behavioral apparatuses were wiped down with ethanol (70%) between

each testing each subject.

<u>Gait Analysis</u>

We used a Noldus Catwalk Gait analysis system designed to semi-automatically measure and analyze

the gait of mice during normal ambulation. Briefly, the movement of mice across a glass bottom corridor

is video recorded from a ventral position. Paw prints are highlighted in the video due to light illumination

across the glass walking platform. Each mouse step within a video is subsequently detected using

Catwalk XT (Noldus) in a semi-automated fashion. A run for each mouse consists of 3 trials of

consistent ambulation across the monitored platform. Only consistent trials are accepted, and mice may

take up to 10 attempts to complete 3 compliant trials in either direction across the corridor. Compliant

trials were defined as those with movement across the platform under 5 s-long and with no more than

60% speed variation. Once placed onto the platform, mice generally ran back and forth without any

need for experimenter prompting.

<u>Vertical Pole</u>

Mice are placed at the top of an 80 cm tall bolt with their nose facing down and hind paws as close to

the top as possible. Mice are immediately released, and time started immediately upon placement.

Time is stopped when the first forepaw touches the surface below the pole. A mouse's natural

predilection is to immediately climb down the pole, and they are given up to 60 s to traverse the pole,

otherwise they are helped off the pole. A non-completed trial is automatically given a time of 30 s, as

95% of mice that did not descend within 30 s were still on the pole at the 60 s mark.

<u>SHIRPA</u>

Behavioral tests were conducted by the University of California, Los Angeles Behavioral Core at P30

and P400. All parameters are scored to provide a quantitative assessment, which enables comparison

of results both over time and between different laboratories. Each mouse was sequentially tested

across all behaviors within ~20-min time span before moving onto the next mouse. The experimenter

was blinded to animal genotype. The screen was performed as described previously (Rogers et al.

1997).

Behavioral Observation

689   The primary screen provides a behavioral observation profile and assessment of each animal begins by

690   observing undisturbed behavior in a viewing jar (10 cm diameter) for 5 min. In addition to the scored

691   behaviors of **body position**, **spontaneous activity**, **respiration rate**, and **tremor**, the observer logs

692   any instances of bizarre or stereotyped behavior and convulsions, compulsive licking, self-destructive

693   biting, retropulsion (walking backwards) and indications of spatial disorientation.

Arena Behavior

695   Thereafter, the mouse is transferred to the arena (30 cm x 50 cm) for testing of transfer arousal and

696   observation of normal behavior. The arena is marked into a grid of 10×10 cm^2 squares to measure

697   locomotor activity within a 30 s-period. While the mouse is active in the arena, measures of **startle**

698   **response**, **gait**, **pelvic elevation**, and **tail elevation** are recorded.

Supine Restraint

700   The animal is restrained in a supine position to record autonomic behaviors. During this assessment,

701   **grip strength**, **body tone**, **pinna reflex**, **corneal reflex**, **toe pinch**, **wire maneuver**, and **heart rate**

702   were evaluated.

Balance and Orientation

704   Finally, several measures of vestibular system function were performed. The **righting reflex**, **contact**

705   **righting reflex**, and **negative geotaxis** tests were performed. Throughout this procedure vocalization,

706   urination and general fear, irritability, or aggression were recorded.

707   Equipment Used

708   1. Clear Plexiglas arena (approximate internal dimensions 55 x 33 x18 cm). On the floor of the arena

709   is a Plexiglas sheet marked with 15 squares (11 cm). A rigid horizontal wire (3 mm diameter) is

710   secured across the rear right corner such that the animals cannot touch the sides during the wire

maneuver. A grid (40×20 cm) with 12 mm mesh (approximate) is secured across the width of the

box for measuring tail suspension and grip strength behavior.

2. A clear Plexiglas cylinder (15×11 cm) was used as a viewing jar.

3. One grid floor (40×20 cm) with 12 mm meshes on which viewing jars stand.

4. Four cylindrical stainless-steel supports (3 cm high x 2.5 cm diameter) to raise grids off the bench.

5. One square (13 cm) stainless steel plate for transfer of animals to the arena.

6. Cut lengths of 3 / 0 Mersilk held in the forceps for corneal and pinna reflex tests

7. A plastic dowel rod sharpened to a pencil point to test salivation and biting.

8. A pair of dissecting equipment forceps, curved with fine points (125 mm forceps, Philip Harris

Scientific, Cat. No. D46-174), for the toe pinch.

9. A stopwatch.

10. An IHR Click box is used for testing the startle responses. The Click Box generates a brief 20

KHz tone at 90dB SPL when held 30cm above the mouse. Contact Prof. K.P. Steel, MRC Institute

of Hearing Research, University Park, Nottingham NG7 2RD.

11. A ruler.

12. A 30 cm clear Plexiglas tube with an internal diameter of 2.5 cm for the contact righting reflex.

4.5 Electrophysiology

<u>Preparation of acute cerebellar slices</u>

Acute parasagittal slices of 300 μm thickness were prepared from the cerebellum of experimental and

control littermate mice by following published methods (Hansen et al., 2013). In brief, cerebella were

quickly removed and immersed in an ice-cold extracellular solution with composition of (mM): 119 NaCl,

26 $NaHCO_3$, 11 glucose, 2.5 KCl, 2.5 $CaCl_2$, 1.3 $MgCl_2$ and 1 NaH_2PO_4, pH 7.4 when gassed with 5%

$CO_2/95\%$ O_2. Cerebella were sectioned parasagittally using a vibratome (Leica VT-1000, Leica

Biosystems, Nussloch, Germany) and initially incubated at 35°C for ~30 min, and then equilibrated and

stored at room temperature until use.

<u>Extracellular Electrophysiology</u>

Extracellular and intracellular recordings were obtained from Purkinje neurons (PNs) in slices

constantly perfused with carbogen-bubbled extracellular solution and maintained at either 37° C

(extracellular) or 32° C (intracellular) ± 1° C (see above). Cells were visualized with DIC optics and a

water-immersion 40✕ objective (NA 0.75) using a Zeiss Examiner microscope. Glass pipettes of ~ 3 MΩ

resistance (Model P-1000, Sutter instruments, Novato, CA) were filled with extracellular solution and

positioned near PN axon hillocks in order to measure action potential-associated capacitive current

transients in voltage clamp mode with the pipette potential held at 0 mV. For whole-cell patch clamp

recordings, pipettes were filled with an intracellular solution (mM): 140 KMeth (CH3KO3S), 10 NaCl, 2

$MgCl_2$, 0.2 $CaCl_2$, 10 HEPES, 14 Phosphocreatine (tris salt), 1 EGTA, 4 Mg-ATP, 0.4 Na-GTP. 100 µM

Picrotoxin (Sigma) was added to block inhibitory GABAegeric synaptic inputs. Data was acquired using

a MultiClamp 700B amplifier at 20 or 100 kHz in voltage or current clamp mode, Digidata 1440 with

pClamp10 (Molecular Devices, Sunnyvale, CA) and filtered at 2 to 4 kHz. The series resistance was

usually between 10 and 15 MΩ. Series resistance was compensated at 80% for short term plasticity

experiments only.

For extracellular recordings, a total of 20 to 45 PNs were recorded from for each animal across all

genotypes, sexes, and age groups. Recordings were distributed across both the medial-lateral and

rostro-caudal axis of the cerebellum. Only cells with a "healthy" look (low contrast of cellular borders)

and regular, uninterrupted firing rate were examined. During analysis, a few cells were found to have

gaps in firing of greater than 2 seconds, and these cells were eliminated from analysis, as this type of

firing is associated with being "unhealthy." Double mutant tissue did not qualitatively differ in

appearance under DIC microscopy prior to recordings, nor was the number of "unhealthy" cells greater

than that of other genotypes (7% vs 4 to 11% of all cells across control genotypes at P400). Spatial

comparison of neural activity was obtained by recording from serial sections in the flocculus, lateral (2[nd]

or 3[rd]), intermediate (6[th] or 7[th]), and medial (11[th] or 12[th]) slices. Lower number slices were used in the

younger age groups (P45 and 110) to roughly match the relative positioning of recordings across age

groups. 0-3 recordings were made from each lobule within each slice dependent on tissue quality and

health. Each recording lasted for 1-minute. 3 to 5 mice were used for each age group, and the

experimenter was blinded to the genotype, age, and sex.

Intracellular recordings were obtained from PNs in either lobule III or VIII of the medial cerebellum (i.e.,

vermis); no statistical differences in properties were observed between lobules.

<u>Analyses</u>

Spontaneous action potential interstimulus intervals were detected and analyzed using standard and

custom routines in ClampFit (Molecular Device), IgorPro (Wavemetrics), and Excel (Microsoft).

Specifically, action potentials were threshold detected and spiking statistics (i.e., frequency and interval

length) were determined using adapted IgorPro routines (Taro Tools;

https://sites.google.com/site/tarotoolsregister/). The coefficient of variation of the mean inter-spike

interval (CV) and the median inter-spike interval ($CV2 = 2 \ |ISI_{n+1}-ISI_n|/(ISI_{n+1}+ISI_n)$)) were

calculated in Excel using custom macros.

Standard membrane properties were analyzed using IgorPro. R_M was determined by averaging 3

voltage trace responses to a -5 mV step pulse from a -80 mV holding potential and measuring the

resulting current deflection between 900 and 1000 ms after onset. The membrane time constant was

measured by fitting a single exponential to the initial decay phase from 90% to 10% of the peak. C_M

was calculated by dividing the membrane time constant by the R_M. sEPSC events were recorded over

a 1-minute epoch and detected and measured using Neuroexpress (v21.1.13). Parallel and climbing

fiber axons were stimulated using theta-glass electrodes (W.P.I.) and a TTL-controlled stimulus isolator

(ISO-Flex, A.M.P.I.). Evoked EPSC amplitudes and decay time constants (1 exp. for parallel and 2 exp.

for climbing fibers) were analyzed using custom routines in IgorPro. Action potentials were examined as

part of a set of 1 s current injections between -500 and 2250 pA (250 pA steps) with a holding current

adjusted to maintain an ~70 mV potential. Action potential waveforms were measured using custom

routines in IgorPro. Action potential threshold was defined as the first membrane voltage in which the

first derivative exceeded 30 mV/ms (Zhu et al. 2006).

4.6 Examination of Cerebellar Atrophy

Cerebellar size

Immediately after brain removal from the skull, a dorsal, whole mount image was obtained. Images

were then processed using Fiji (NIH). The forebrain and cerebellar sizes were assessed by outlining

their 2-dimensional space and then calculating area. We normalized for possible differences in overall

brain size by dividing the results of the cerebellum by forebrain size to produce a relative cerebellum-to-

forebrain ratio. Experimenters were blind to the genotype of the animal.

Immunohistochemistry

At the respective study endpoints (P45, 120, 210, 400), male and female mice of all genotypes

represented in this study were anesthetized with isoflurane and underwent transcardial perfusion with

phosphate-buffered saline followed by 4% (w/v) buffered paraformaldehyde (PFA) and then dissected

to extract the brain. Images of the whole brain were taken immediately after removing the brain from

the skull and the brains were then submerged in 4% PFA for 24 hours, and then cryoprotected in Tris-

buffered saline (TBS) with 0.05% azide and 30% sucrose for 72 hours and stored at 4°C until further

use. The cerebellum was separated from the forebrain and parasagittally sectioned using a sliding

microtome (Microm HM 430, Thermo Scientific) set to section at $40\,\mu m$ thickness. Cerebellum sections

were collected in a series of six and stored in TBS-AF (TBS with 30% sucrose, 0.05% sodium azide,

and 30% ethylene glycol) at 4°C or -20°C until further use . For immunofluorescent visualization of

Purkinje neurons, cerebellum sections of both $Atm^{+/+}$; $Aptx^{+/+}$ and $Atm^{R35X/R35X}$; $Aptx^{-/-}$ ($n = 5$ per

genotype) were washed for $5\,min$ in TBS three times, and then blocked in 15% normal goat serum at

room temperature for $30\,min$ followed by free-floating incubation in rabbit or mouse anti-calbindin D-28k

(1:1000) for 1 hour at room temperature on an orbital shaker. Sections were then washed for $5\,min$ with

TBS three times, followed by free floating incubation in goat anti-rabbit or mouse Alexa Fluor 488

(1:1000) for $1\,h$ in the dark at room temperature on an orbital shaker. Following secondary antibody

incubation, sections were washed for $5\,min$ in TBS three times, then mounted and cover-slipped with

Fluoromount-G with DAPI. For some sections, anti-cleaved Caspase-3 (1:200) and anti-CD68 (1:400)

antibodies were additionally probed in parallel with Calbindin using an Alexa Fluor 647 (1:500)

secondary antibody. Slides were scanned using Stereo Investigator (MBF Bioscience, ver. 2020) on a

Zeiss microscope equipped with an ApoTome 2 (Carl Zeiss Microscopy, Axio Imager.M2) using either a

2.5, 10, 20, 40, or 63x objective and images captured with a Hamamatsu CMOS camera (Hamamatsu

Photonics, ORCA Flash 4.0 LT+).

To quantify the number of calbindin-reactive cells in each lobule in the resulting images, we used

Stereo Investigator to randomly draw 2 lines between 300 to 500 μm long in each lobule and manually

counted the total number of PNs along the length within the 40 μm thickness of the tissue slice under

40x magnification. 2D density (# of PNs/(linear length * 40 um thickness)) of the two samples per lobule

were then averaged for further comparison between lobules and animals.

Calbindin positive PN dendrite widths were measured at a predefined location in lobule VI from each

animal in 25 or 40 μm thick tissue sections under 20x magnification. Dendritic widths of the primary and

secondary branches were measured at the midline between the PN cell bodies and edge of the

molecular layer. Between 7 and 13 dendrites were measured per section, one section per animal.

For PN somatic measurements, Stereo Investigator was used to randomly select PNs distributed

across the entire medial section under 20x magnification. The average PN width per animal was

determined by averaging results across 3 serial sections (16 to 37 PNs per section). PN widths were

measured perpendicular to the PN layer or to the exiting dendrite if askew by more than a few degrees.

Molecular layer and granule cell layer (visualized with Calbindin and DAPI stains, respectively) widths

were assessed in Stereo Investigator by averaging two width measurements at predefined locations for

each lobule, roughly halfway along the long extent of each lobule under 2.5x magnification.

CD68 positivity in the cerebellar sections was quantified by measuring the total percent area of CD68[+]

positive staining across the entire medial cerebellar section. 10x stitched images were thresholded to

the negative control and quantified using ImageJ, one section per animal.

To quantify the percent of Calbindin-positive PNs that were positive for cleaved Caspase-3 we counted

PNs across the entire cerebellum using Stereo Investigator. Three, 20x magnification stitched images

per animal were examined and the results averaged. The threshold for Caspase-3 positivity was

established from control sections stained with only the secondary antibodies.

For non-fluorescent histological analysis, 25-µm-thick, free-floating tissue sections onto positively

charged slides and air-dried overnight. The tissue was washed in Phosphate-buffered saline (PBS)

twice for 5 min, then stained sequentially with 0.1% Hematoxylin in 95% ethanol for ~25s and 0.5%

Eosin in 95% ethanol for ~3s and washed in double distilled water after each stain. The tissue was

subsequently dehydrated for 1 min in 95% ethanol, 100% ethanol, and 100% Xylene washes, then

cover slipped with Permount. Slides were imaged using a color camera (Q Imaging, MBF Biosciences)

on the same Zeiss microscope and MBF acquisition software.

Experimenters were blinded to the mouse genotype in which sections were examined, and the order of

examination was interleaved for all histological measurements.

4.7 Flow Cytometry Measurements

Flow cytometry analysis of blood and thymus cells was performed by staining with specific anti-mouse

antibodies: CD4, CD8, CD3, CD44, and CD25. Briefly, whole-blood samples ($50\ \mu$l) were stained using

fluorescent-labeled antibodies, then red-blood cells were lysed using BD lysing solution while live white-

blood cells were stained using a viability stain. Thymi were mechanically dissociated. $1\ \text{to}\ 2\ \text{million}$

thymus cells were similarly stained using specific antibodies for CD4, CD8, CD44 and CD25. Analysis

of immuno-stained white blood cells or thymus samples was performed using FACS ARIA III and data

analyzed using FlowJo software as reported previously (Sanghez et al. 2017).

4.8 Western Blots

Protein extracts (cells/tissues) were homogenized in radioimmunoprecipitation assay (RIPA) lysis buffer

(150 mM NaCl, 1% Nonidet P-40 [NP-40], 0.5% deoxycholate, 0.1% SDS, 50 mM Tris, pH 8.0) with

protease inhibitors (10 ug/ml AEBSF, 10 ug/ml leupeptin, 5 ug/ml pepstatin, 5 ug/ml chymotrypsin, 10

ug/ml aprotinin). The protein extracts were sonicated then pelleted by centrifugation at 13,000 rpm for

15 min at 4°C. BCA protein assay was used to quantify protein concentrations. Samples containing

equal amounts of protein 50 to 100 μg per lane were separated using 4 to 12% gradient TGX precast

gels BioRad then transferred by TransBlot Semi-Dry BioRad system using Nitrocellulose transfer pack.

Transferred blots were stained by Ponceau S stain for equal protein loading then washed and blocked

with 5% nonfat dry milk in TBST for 60 min at room temp. Primary antibodies were incubated with

shaking overnight at 4°C. Blots were probed for the following antibodies: ATM (D2E2) Rabbit mAb Cell

Signaling, at 1:1000 dilution, β-Actin (D6A8) Rabbit mAb Cell Signaling, GAPDH (D16H11) Rabbit mAb

Cell Signaling followed by the appropriate horseradish peroxidase–conjugated (HRP) secondary Anti-

rabbit, Anti-mouse for 2 hours at room temperature. After multiple washes with TBST, Protein

expression was detected by Radiance Plus chemiluminescence substrate using the Azure c400 and the

BioRad ChemiDoc imaging systems. Densitometric analysis of the ATM was performed using ImageJ.

Experiments were performed with 2 technical and 2-3 biological replicates as indicated.

4.9 Statistical Assessment

The number of animals chosen for each group was based on a priori power analyses using GPower

v3.1 based on an α size of 0.5, power of 0.8, and effect sizes estimated from preliminary data or prior

studies. We used both parametric (1- and 2-way ANOVA) for normally distributed and non-parametric

(Kruskal Wallace) statistical methods for interval data to test for differences between groups followed by

pairwise multiple comparisons tests as indicated in the text. Outliers for immune data in Figs. 6 and 7

were excluded via the ROUT method (Q=2%). The specific analyses used for each data set is noted in

each figure legend. For all figures: ns not significant, * $p \leq 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

Data are reported as mean $\pm$ SEM and box and whisker plots indicate the minimum, first quartile,

median, third quartile, and maximum data values. All figures and statistical analyses were completed

using Excel (Microsoft 360) or Prism v8 and 9 (Graphpad).

5.0 Acknowledgements

We would like to thank the UCLA Behavioral Core, especially Irina Zhuravka for her efforts assaying

behavioral deficits in the mice. We would also like to thank Dr. Jennifer Fogel for comments and edits to

the manuscript.

6.0 Competing interests

The authors declare that no competing interests exist.

7.0 Citations

Ady, Visou, Brenda Toscano-Márquez, Moushumi Nath, Philip K Chang, Jeanette Hui, Anna Cook,
François Charron, Roxanne Larivière, Bernard Brais, and R Anne McKinney. 2018. 'Altered
synaptic and firing properties of cerebellar Purkinje cells in a mouse model of ARSACS', *The*
Journal of physiology, 596: 4253-67.
Aguilar, M. J., S. Kamoshita, B. H. Landing, E. Boder, and R. P. Sedgwick. 1968a. 'Pathological
observations in ataxia-telangiectasia. A report of five cases', *Journal of Neuropathology and*
Experimental Neurology, 27: 659-76.
Aguilar, Mary Jane, Shigehiko Kamoshita, Benjamin H. Landing, Elena Boder, and Robert P. Sedgwick.
1968b. 'Pathological Observations in Ataxia-Telangiectasia: A Report on Five Cases*', *Journal*
of Neuropathology and Experimental Neurology, 27: 659-76.
Ahel, I., U. Rass, S. F. El-Khamisy, S. Katyal, P. M. Clements, P. J. McKinnon, K. W. Caldecott, and S.
C. West. 2006. 'The neurodegenerative disease protein aprataxin resolves abortive DNA
ligation intermediates', *Nature*, 443: 713-6.
Aicardi, J., C. Barbosa, E. Andermann, F. Andermann, R. Morcos, Q. Ghanem, Y. Fukuyama, Y.
Awaya, and P. Moe. 1988. 'Ataxia-ocular motor apraxia: a syndrome mimicking ataxia-
telangiectasia', *Annals of Neurology*, 24: 497-502.
Alviña, Karina, and Kamran Khodakhah. 2010. 'KCa channels as therapeutic targets in episodic ataxia
type-2', *Journal of Neuroscience*, 30: 7249-57.
Amromin, G. D., E. Boder, and R. Teplitz. 1979. 'Ataxia-telangiectasia with a 32 year survival. A
clinicopathological report', *Journal of Neuropathology and Experimental Neurology*, 38: 621-43.
Ando, K., J. L. Kernan, P. H. Liu, T. Sanda, E. Logette, J. Tschopp, A. T. Look, J. Wang, L. Bouchier-
Hayes, and S. Sidi. 2012. 'PIDD death-domain phosphorylation by ATM controls prodeath
versus prosurvival PIDDosome signaling', *Molecular Cell*, 47: 681-93.
Apps, R., and R. Hawkes. 2009. 'Cerebellar cortical organization: a one-map hypothesis', *Nature*
Reviews: Neuroscience, 10: 670-81.
Atluri, Pradeep P., and Wade G. Regehr. 1996. 'Determinants of the Time Course of Facilitation at the
Granule Cell to Purkinje Cell Synapse', *The Journal of Neuroscience*, 16: 5661-71.
Barlow, C., S. Hirotsune, R. Paylor, M. Liyanage, M. Eckhaus, F. Collins, Y. Shiloh, J. N. Crawley, T.
Ried, D. Tagle, and A. Wynshaw-Boris. 1996. 'Atm-deficient mice: a paradigm of ataxia
telangiectasia', *Cell*, 86: 159-71.
Beraldi, R., D. K. Meyerholz, A. Savinov, A. D. Kovacs, J. M. Weimer, J. A. Dykstra, R. D. Geraets, and
D. A. Pearce. 2017. 'Genetic ataxia telangiectasia porcine model phenocopies the multisystemic
features of the human disease', *Biochim Biophys Acta Mol Basis Dis*, 1863: 2862-70.
Bhatt, Jayesh M., and Andrew Bush. 2014. 'Microbiological surveillance in lung disease in ataxia
telangiectasia', *European Respiratory Journal*, 43: 1797-801.

Boder, Elena, and Robert P. Sedgwick. 1958. 'ATAXIA-TELANGIECTASIA', *A Familial Syndrome of*
Progressive Cerebellar Ataxia, Oculocutaneous Telangiectasia and Frequent Pulmonary
Infection, 21: 526-54.
Bogue, M. A., C. Wang, C. Zhu, and D. B. Roth. 1997. 'V(D)J recombination in Ku86-deficient mice:
distinct effects on coding, signal, and hybrid joint formation', *Immunity*, 7: 37-47.
Bosch, Marie K, Yarimar Carrasquillo, Joseph L Ransdell, Ajay Kanakamedala, David M Ornitz, and
Jeanne M Nerbonne. 2015. 'Intracellular FGF14 (iFGF14) is required for spontaneous and
evoked firing in cerebellar Purkinje neurons and for motor coordination and balance', *Journal of*
Neuroscience, 35: 6752-69.
Bradley, A., K. Anastassiadis, A. Ayadi, J. F. Battey, C. Bell, M. C. Birling, J. Bottomley, S. D. Brown, A.
Bürger, C. J. Bult, W. Bushell, F. S. Collins, C. Desaintes, B. Doe, A. Economides, J. T. Eppig,
R. H. Finnell, C. Fletcher, M. Fray, D. Frendewey, R. H. Friedel, F. G. Grosveld, J. Hansen, Y.
Hérault, G. Hicks, A. Hörlein, R. Houghton, M. Hrabé de Angelis, D. Huylebroeck, V. Iyer, P. J.
de Jong, J. A. Kadin, C. Kaloff, K. Kennedy, M. Koutsourakis, K. C. Lloyd, S. Marschall, J.
Mason, C. McKerlie, M. P. McLeod, H. von Melchner, M. Moore, A. O. Mujica, A. Nagy, M.
Nefedov, L. M. Nutter, G. Pavlovic, J. L. Peterson, J. Pollock, R. Ramirez-Solis, D. E. Rancourt,
M. Raspa, J. E. Remacle, M. Ringwald, B. Rosen, N. Rosenthal, J. Rossant, P. Ruiz Noppinger,
E. Ryder, J. Z. Schick, F. Schnütgen, P. Schofield, C. Seisenberger, M. Selloum, E. M.
Simpson, W. C. Skarnes, D. Smedley, W. L. Stanford, A. F. Stewart, K. Stone, K. Swan, H.
Tadepally, L. Teboul, G. P. Tocchini-Valentini, D. Valenzuela, A. P. West, K. Yamamura, Y.
Yoshinaga, and W. Wurst. 2012. 'The mammalian gene function resource: the International
Knockout Mouse Consortium', *Mammalian Genome*, 23: 580-6.
Bredemeyer, Andrea L, Girdhar G Sharma, Ching-Yu Huang, Beth A Helmink, Laura M Walker, Katrina
C Khor, Beth Nuskey, Kathleen E Sullivan, Tej K Pandita, and Craig H Bassing. 2006. 'ATM
stabilizes DNA double-strand-break complexes during V (D) J recombination', *Nature*, 442: 466-
70.
Browne, Susan E., L. Jackson Roberts, Phyllis A. Dennery, Susan R. Doctrow, M. Flint Beal, Carrolee
Barlow, and Rodney L. Levine. 2004. 'Treatment with a catalytic antioxidant corrects the
neurobehavioral defect in ataxia–telangiectasia mice', *Free Radical Biology and Medicine*, 36:
938-42.
Çaglayan, Melike, Rajendra Prasad, Rachel Krasich, Matthew J. Longley, Kei Kadoda, Masataka
Tsuda, Hiroyuki Sasanuma, Shunichi Takeda, Keizo Tano, William C. Copeland, and Samuel H.
Wilson. 2017. 'Complementation of aprataxin deficiency by base excision repair enzymes in
mitochondrial extracts', *Nucleic Acids Research*, 45: 10079-88.
Campbell, A., B. Krupp, J. Bushman, M. Noble, C. Pröschel, and M. Mayer-Pröschel. 2015. 'A novel
mouse model for ataxia-telangiectasia with a N-terminal mutation displays a behavioral defect
and a low incidence of lymphoma but no increased oxidative burden', *Human Molecular*
Genetics, 24: 6331-49.
Campbell, Andrew, Jared Bushman, Joshua Munger, Mark Noble, Christoph Pröschel, and Margot
Mayer-Pröschel. 2016. 'Mutation of ataxia–telangiectasia mutated is associated with
dysfunctional glutathione homeostasis in cerebellar astroglia', *Glia*, 64: 227-39.
Chao, C., E. M. Yang, and Y. Xu. 2000. 'Rescue of defective T cell development and function in Atm-/-
mice by a functional TCR alpha beta transgene', *Journal of Immunology*, 164: 345-9.
Chatterjee, Nimrat, and Graham C. Walker. 2017. 'Mechanisms of DNA damage, repair, and
mutagenesis', *Environmental and Molecular Mutagenesis*, 58: 235-63.
Chen, P., C. Peng, J. Luff, K. Spring, D. Watters, S. Bottle, S. Furuya, and M. F. Lavin. 2003. 'Oxidative
stress is responsible for deficient survival and dendritogenesis in Purkinje neurons from ataxia-
telangiectasia mutated mutant mice', *Journal of Neuroscience*, 23: 11453-60.
Chen, Xi, Tie-Shan Tang, Huiping Tu, Omar Nelson, Mark Pook, Robert Hammer, Nobuyuki Nukina,
and Ilya Bezprozvanny. 2008. 'Deranged calcium signaling and neurodegeneration in
spinocerebellar ataxia type 3', *Journal of Neuroscience*, 28: 12713-24.
Chou, W. C., L. Y. Hu, C. N. Hsiung, and C. Y. Shen. 2015. 'Initiation of the ATM-Chk2 DNA damage
response through the base excision repair pathway', *Carcinogenesis*, 36: 832-40.

Clark, H Brent, Eric N Burright, Wael S Yunis, Seth Larson, Claire Wilcox, Boyd Hartman, Antoni
Matilla, Huda Y Zoghbi, and Harry T Orr. 1997. 'Purkinje Cell Expression of a Mutant Allele of
SCA1in Transgenic Mice Leads to Disparate Effects on Motor Behaviors, Followed by a
Progressive Cerebellar Dysfunction and Histological Alterations', *Journal of Neuroscience*, 17:
7385-95.
Clements, P. M., C. Breslin, E. D. Deeks, P. J. Byrd, L. Ju, P. Bieganowski, C. Brenner, M. C. Moreira,
A. M. Taylor, and K. W. Caldecott. 2004. 'The ataxia-oculomotor apraxia 1 gene product has a
role distinct from ATM and interacts with the DNA strand break repair proteins XRCC1 and
XRCC4', *DNA Repair (Amst)*, 3: 1493-502.
Concannon, P., and R. A. Gatti. 1997. 'Diversity of ATM gene mutations detected in patients with
ataxia-telangiectasia', *Human Mutation*, 10: 100-7.
Cook, A. A., E. Fields, and A. J. Watt. 2021. 'Losing the Beat: Contribution of Purkinje Cell Firing
Dysfunction to Disease, and Its Reversal', *Neuroscience*, 462: 247-61.
Coutinho P, Barbot C, Coutinho P. 2002. 'Ataxia with Oculomotor Apraxia Type 1.' in Ardinger HH
Adam MP, Pagon RA, et al. (ed.), *GeneReviews®* (GeneReviews® [Internet]: University of
Washington, Seattle).
Crawford, T. O., R. L. Skolasky, R. Fernandez, K. J. Rosquist, and H. M. Lederman. 2006. 'Survival
probability in ataxia telangiectasia', *Archives of Disease in Childhood*, 91: 610-11.
De Leon, G. A., W. D. Grover, and D. S. Huff. 1976. 'Neuropathologic changes in ataxia-telangiectasia',
Neurology, 26: 947-51.
Dell'Orco, James M., Aaron H. Wasserman, Ravi Chopra, Melissa A. C. Ingram, Yuan-Shih Hu, Vikrant
Singh, Heike Wulff, Puneet Opal, Harry T. Orr, and Vikram G. Shakkottai. 2015. 'Neuronal
Atrophy Early in Degenerative Ataxia Is a Compensatory Mechanism to Regulate Membrane
Excitability', *The Journal of Neuroscience*, 35: 11292-307.
Dell'Orco, James M, Stefan M Pulst, and Vikram G Shakkottai. 2017. 'Potassium channel dysfunction
underlies Purkinje neuron spiking abnormalities in spinocerebellar ataxia type 2', *Human
Molecular Genetics*, 26: 3935-45.
Demaerel, PH, BE Kendall, and D Kingsley. 1992. 'Cranial CT and MRI in diseases with DNA repair
defects', *Neuroradiology*, 34: 117-21.
Demirci, O., M. C. Stevens, N. C. Andreasen, A. Michael, J. Liu, T. White, G. D. Pearlson, V. P. Clark,
and V. D. Calhoun. 2009. 'Investigation of relationships between fMRI brain networks in the
spectral domain using ICA and Granger causality reveals distinct differences between
schizophrenia patients and healthy controls', *Neuroimage*, 46: 419-31.
Dineen, Rob A., Felix Raschke, Hannah L. McGlashan, Stefan Pszczolkowski, Lorna Hack, Andrew D.
Cooper, Manish Prasad, Gabriel Chow, William P. Whitehouse, and Dorothee P. Auer. 2020.
'Multiparametric cerebellar imaging and clinical phenotype in childhood ataxia telangiectasia',
NeuroImage: Clinical, 25: 102110.
Du, L., M. E. Jung, R. Damoiseaux, G. Completo, F. Fike, J. M. Ku, S. Nahas, C. Piao, H. Hu, and R. A.
Gatti. 2013. 'A new series of small molecular weight compounds induce read through of all three
types of nonsense mutations in the ATM gene', *Molecular Therapy*, 21: 1653-60.
El-Brolosy, Mohamed A., and Didier Y. R. Stainier. 2017. 'Genetic compensation: A phenomenon in
search of mechanisms', *Plos Genetics*, 13: e1006780-e80.
Elson, A., Y. Wang, C. J. Daugherty, C. C. Morton, F. Zhou, J. Campos-Torres, and P. Leder. 1996.
'Pleiotropic defects in ataxia-telangiectasia protein-deficient mice', *Proceedings of the National
Academy of Sciences of the United States of America*, 93: 13084-9.
Gao, Zhenyu, Boeke J van Beugen, and Chris I De Zeeuw. 2012. 'Distributed synergistic plasticity and
cerebellar learning', *Nature Reviews: Neuroscience*, 13: 619-35.
Gatti, R. A., and H. V. Vinters. 1985. 'Cerebellar pathology in ataxia-telangiectasia: the significance of
basket cells', *Kroc Foundation Series*, 19: 225-32.
Genik, P. C., H. Bielefeldt-Ohmann, X. Liu, M. D. Story, L. Ding, J. M. Bush, C. M. Fallgren, and M. M.
Weil. 2014. 'Strain background determines lymphoma incidence in Atm knockout mice',
Neoplasia, 16: 129-36.

Germain, Ronald N. 2002. 'T-cell development and the CD4–CD8 lineage decision', *Nature Reviews Immunology*, 2: 309-22.

Gertsenstein, Marina, Lauryl M. J. Nutter, Tammy Reid, Monica Pereira, William L. Stanford, Janet Rossant, and Andras Nagy. 2010. 'Efficient Generation of Germ Line Transmitting Chimeras from C57BL/6N ES Cells by Aggregation with Outbred Host Embryos', *PloS One*, 5: e11260.

Gilad, S., A. Bar-Shira, R. Harnik, D. Shkedy, Y. Ziv, R. Khosravi, K. Brown, L. Vanagaite, G. Xu, M. Frydman, M. F. Lavin, D. Hill, D. A. Tagle, and Y. Shiloh. 1996. 'Ataxia-telangiectasia: founder effect among north African Jews', *Human Molecular Genetics*, 5: 2033-7.

Gilad, Shlomit, Rami Khosravi, Dganit Shkedy, Tamar Uziel, Yael Ziv, Kinneret Savitsky, Galit Rotman, Sara Smith, Luciana Chessa, Timothy J. Jorgensen, Reli Harnik, Moshe Frydman, Ozden Sanal, Sima Portnoi, Zipora Goldwicz, N. G. J. Jaspers, Richard A. Gatti, Gilbert Lenoir, Martin F. Lavin, Kouichi Tatsumi, Rolf D. Wegner, Yosef Shiloh, and Anat Bar-Shira. 1996. 'Predominance of Null Mutations in Ataxia-Telangiectasia', *Human Molecular Genetics*, 5: 433-39.

Gueven, N., O. J. Becherel, A. W. Kijas, P. Chen, O. Howe, J. H. Rudolph, R. Gatti, H. Date, O. Onodera, G. Taucher-Scholz, and M. F. Lavin. 2004. 'Aprataxin, a novel protein that protects against genotoxic stress', *Human Molecular Genetics*, 13: 1081-93.

Guo, Z., S. Kozlov, M. F. Lavin, M. D. Person, and T. T. Paull. 2010. 'ATM activation by oxidative stress', *Science*, 330: 517-21.

Hansel, Christian, and David J. Linden. 2000. 'Long-Term Depression of the Cerebellar Climbing Fiber–Purkinje Neuron Synapse', *Neuron*, 26: 473-82.

Hansen, S. T., P. Meera, T. S. Otis, and S. M. Pulst. 2013. 'Changes in Purkinje cell firing and gene expression precede behavioral pathology in a mouse model of SCA2', *Human Molecular Genetics*, 22: 271-83.

Hathcock, K. S., S. Bowen, F. Livak, and R. J. Hodes. 2013. 'ATM influences the efficiency of TCRbeta rearrangement, subsequent TCRbeta-dependent T cell development, and generation of the pre-selection TCRbeta CDR3 repertoire', *PloS One*, 8: e62188.

Herzog, Karl-Heinz, Miriam J. Chong, Manuela Kapsetaki, James I. Morgan, and Peter J. McKinnon. 1998. 'Requirement for Atm in Ionizing Radiation-Induced Cell Death in the Developing Central Nervous System', *Science*, 280: 1089-91.

Hoche, F., K. Seidel, M. Theis, S. Vlaho, R. Schubert, S. Zielen, and M. Kieslich. 2012. 'Neurodegeneration in ataxia telangiectasia: what is new? What is evident?', *Neuropediatrics*, 43: 119-29.

Hourez, Raphael, Laurent Servais, David Orduz, David Gall, Isabelle Millard, Alban de Kerchove d'Exaerde, Guy Cheron, Harry T Orr, Massimo Pandolfo, and Serge N Schiffmann. 2011. 'Aminopyridines correct early dysfunction and delay neurodegeneration in a mouse model of spinocerebellar ataxia type 1', *Journal of Neuroscience*, 31: 11795-807.

Hoxha, Eriola, Ilaria Balbo, Maria Concetta Miniaci, and Filippo Tempia. 2018. 'Purkinje Cell Signaling Deficits in Animal Models of Ataxia', *Frontiers in Synaptic Neuroscience*, 10: 6-6.

Hurlock, Edward C, Anne McMahon, and Rolf H Joho. 2008. 'Purkinje-cell-restricted restoration of Kv3. 3 function restores complex spikes and rescues motor coordination in Kcnc3 mutants', *Journal of Neuroscience*, 28: 4640-48.

Jackson, Thomas J, Gabriel Chow, Mohnish Suri, Philip Byrd, Malcolm R Taylor, and William P Whitehouse. 2016. 'Longitudinal analysis of the neurological features of ataxia-telangiectasia', *Developmental Medicine and Child Neurology*, 58: 690-97.

Jayabal, Sriram, Hui Ho Vanessa Chang, Kathleen E Cullen, and Alanna J Watt. 2016. '4-aminopyridine reverses ataxia and cerebellar firing deficiency in a mouse model of spinocerebellar ataxia type 6', *Scientific Reports*, 6: 1-12.

Kaminsky, Natalie, Ofer Bihari, Sivan Kanner, and Ari Barzilai. 2016. 'Connecting Malfunctioning Glial Cells and Brain Degenerative Disorders', *Genomics, Proteomics & Bioinformatics*, 14: 155-65.

Kano, Masanobu, and Takaki Watanabe. 2017. 'Type-1 metabotropic glutamate receptor signaling in cerebellar Purkinje cells in health and disease', *F1000Research*, 6: 416-16.

Kastan, M. B., and J. Bartek. 2004. 'Cell-cycle checkpoints and cancer', *Nature*, 432: 316-23.

Kasumu, A., and I. Bezprozvanny. 2012. 'Deranged calcium signaling in Purkinje cells and
pathogenesis in spinocerebellar ataxia 2 (SCA2) and other ataxias', *Cerebellum*, 11: 630-9.
Kim, Jusik, Keeeun Kim, Jung-Soon Mo, and Youngsoo Lee. 2020. 'Atm deficiency in the DNA
polymerase β null cerebellum results in cerebellar ataxia and Itpr1 reduction associated with
alteration of cytosine methylation', *Nucleic Acids Research*, 48: 3678-91.
Krangel, Michael S. 2009. 'Mechanics of T cell receptor gene rearrangement', *Current Opinion in
Immunology*, 21: 133-39.
Larivière, Roxanne, Nicolas Sgarioto, Brenda Toscano Márquez, Rébecca Gaudet, Karine Choquet, R
Anne McKinney, Alanna J Watt, and Bernard Brais. 2019. 'Sacs R272C missense homozygous
mice develop an ataxia phenotype', *Molecular Brain*, 12: 1-15.
Lavin, M. F. 2013. 'The appropriateness of the mouse model for ataxia-telangiectasia: neurological
defects but no neurodegeneration', *DNA Repair (Amst)*, 12: 612-9.
Levy, Ariel, and Anthony E. Lang. 2018. 'Ataxia-telangiectasia: A review of movement disorders, clinical
features, and genotype correlations', *Movement Disorders*, 33: 1238-47.
Li, H. H., W. H. Yu, N. Rozengurt, H. Z. Zhao, K. M. Lyons, S. Anagnostaras, M. S. Fanselow, K.
Suzuki, M. T. Vanier, and E. F. Neufeld. 1999. 'Mouse model of Sanfilippo syndrome type B
produced by targeted disruption of the gene encoding alpha-N-acetylglucosaminidase',
Proceedings of the National Academy of Sciences of the United States of America, 96: 14505-
10.
Li, J., Y. R. Han, M. R. Plummer, and K. Herrup. 2009. 'Cytoplasmic ATM in neurons modulates
synaptic function', *Current Biology*, 19: 2091-6.
Li, Jiali, Jianmin Chen, Harry V. Vinters, Richard A. Gatti, and Karl Herrup. 2011. 'Stable Brain
ATM Message and Residual Kinase-Active ATM Protein in Ataxia-Telangiectasia',
The Journal of Neuroscience, 31: 7568-77.
Lindahl, T., and D. E. Barnes. 2000. 'Repair of endogenous DNA damage', *Cold Spring Harbor
Symposia on Quantitative Biology*, 65: 127-33.
Liu, Jing, Tie-Shan Tang, Huiping Tu, Omar Nelson, Emily Herndon, Duong P Huynh, Stefan M Pulst,
and Ilya Bezprozvanny. 2009. 'Deranged calcium signaling and neurodegeneration in
spinocerebellar ataxia type 2', *Journal of Neuroscience*, 29: 9148-62.
Livák, Ferenc, Michelle Tourigny, David G. Schatz, and Howard T. Petrie. 1999. 'Characterization of
TCR Gene Rearrangements During Adult Murine T Cell Development', *The Journal of
Immunology*, 162: 2575-80.
Lorenzetti, Diego, Kei Watase, Bisong Xu, Martin M Matzuk, Harry T Orr, and Huda Y Zoghbi. 2000.
'Repeat instability and motor incoordination in mice with a targeted expanded CAG repeat in the
Sca1 locus', *Human Molecular Genetics*, 9: 779-85.
Marshall, Jean S., Richard Warrington, Wade Watson, and Harold L. Kim. 2018. 'An introduction to
immunology and immunopathology', *Allergy, asthma, and clinical immunology : official journal of
the Canadian Society of Allergy and Clinical Immunology*, 14: 49-49.
Matei, I. R., C. J. Guidos, and J. S. Danska. 2006. 'ATM-dependent DNA damage surveillance in T-cell
development and leukemogenesis: the DSB connection', *Immunological Reviews*, 209: 142-58.
Matsuoka, Shuhei, Bryan A. Ballif, Agata Smogorzewska, E. Robert McDonald, Kristen E. Hurov, Ji
Luo, Corey E. Bakalarski, Zhenming Zhao, Nicole Solimini, Yaniv Lerenthal, Yosef Shiloh,
Steven P. Gygi, and Stephen J. Elledge. 2007. 'ATM and ATR Substrate Analysis Reveals
Extensive Protein Networks Responsive to DNA Damage', *Science*, 316: 1160-66.
McKinnon, P. J. 2009. 'DNA repair deficiency and neurological disease', *Nature Reviews:
Neuroscience*, 10: 100-12.
Meagher, M., and R. N. Lightowlers. 2014. 'The role of TDP1 and APTX in mitochondrial DNA repair',
Biochimie, 100: 121-4.
Micol, R., L. Ben Slama, F. Suarez, L. Le Mignot, J. Beauté, N. Mahlaoui, C. Dubois d'Enghien, A.
Laugé, J. Hall, J. Couturier, L. Vallée, B. Delobel, F. Rivier, K. Nguyen, T. Billette de Villemeur,
J. L. Stephan, P. Bordigoni, Y. Bertrand, N. Aladjidi, J. M. Pedespan, C. Thomas, I. Pellier, M.
Koenig, O. Hermine, C. Picard, D. Moshous, B. Neven, F. Lanternier, S. Blanche, M. Tardieu,
M. Debré, A. Fischer, and D. Stoppa-Lyonnet. 2011. 'Morbidity and mortality from ataxia-

telangiectasia are associated with ATM genotype', *Journal of Allergy and Clinical Immunology*, 128: 382-9.e1.

Monaco, S., E. Nardelli, G. Moretto, T. Cavallaro, and N. Rizzuto. 1988. 'Cytoskeletal pathology in ataxia-telangiectasia', *Clinical Neuropathology*, 7: 44-6.

Moreira, M. C., S. Klur, M. Watanabe, A. H. Nemeth, I. Le Ber, J. C. Moniz, C. Tranchant, P. Aubourg, M. Tazir, L. Schols, M. Pandolfo, J. B. Schulz, J. Pouget, P. Calvas, M. Shizuka-Ikeda, M. Shoji, M. Tanaka, L. Izatt, C. E. Shaw, A. M'Zahem, E. Dunne, P. Bomont, T. Benhassine, N. Bouslam, G. Stevanin, A. Brice, J. Guimaraes, P. Mendonca, C. Barbot, P. Coutinho, J. Sequeiros, A. Durr, J. M. Warter, and M. Koenig. 2004. 'Senataxin, the ortholog of a yeast RNA helicase, is mutant in ataxia-ocular apraxia 2', *Nature Genetics*, 36: 225-7.

Morrell, Daphne, Elizabeth Cromartie, and Michael Swift. 1986. 'Mortality and cancer incidence in 263 patients with ataxia-telangiectasia', *Journal of the National Cancer Institute*, 77: 89-92.

Paula-Barbosa, M. M., C. Ruela, M. A. Tavares, C. Pontes, A. Saraiva, and C. Cruz. 1983. 'Cerebellar cortex ultrastructure in ataxia-telangiectasia', *Annals of Neurology*, 13: 297-302.

Perkins, Emma M, Yvonne L Clarkson, Nancy Sabatier, David M Longhurst, Christopher P Millward, Jennifer Jack, Junko Toraiwa, Mitsunori Watanabe, Jeffrey D Rothstein, and Alastair R Lyndon. 2010. 'Loss of β-III spectrin leads to Purkinje cell dysfunction recapitulating the behavior and neuropathology of spinocerebellar ataxia type 5 in humans', *Journal of Neuroscience*, 30: 4857-67.

Petersen, Andrew J., Stacey A. Rimkus, and David A. Wassarman. 2012. 'ATM kinase inhibition in glial cells activates the innate immune response and causes neurodegeneration in Drosophila', *Proceedings of the National Academy of Sciences*, 109: E656-E64.

Pizzamiglio, L., E. Focchi, and F. Antonucci. 2020. 'ATM Protein Kinase: Old and New Implications in Neuronal Pathways and Brain Circuitry', *Cells*, 9.

Quarantelli, Mario, Giuliana Giardino, Anna Prinster, Giuseppina Aloj, Barbara Carotenuto, Emilia Cirillo, Angela Marsili, Elena Salvatore, Ennio Del Giudice, and Claudio Pignata. 2013. 'Steroid treatment in Ataxia-Telangiectasia induces alterations of functional magnetic resonance imaging during prono-supination task', *European Journal of Paediatric Neurology*, 17: 135-40.

Quek, Hazel, John Luff, KaGeen Cheung, Sergei Kozlov, Magtouf Gatei, C. Soon Lee, Mark C. Bellingham, Peter G. Noakes, Yi Chieh Lim, Nigel L. Barnett, Steven Dingwall, Ernst Wolvetang, Tomoji Mashimo, Tara L. Roberts, and Martin F. Lavin. 2016. 'A rat model of ataxia-telangiectasia: evidence for a neurodegenerative phenotype', *Human Molecular Genetics*, 26: 109-23.

Rashi-Elkeles, S., R. Elkon, N. Weizman, C. Linhart, N. Amariglio, G. Sternberg, G. Rechavi, A. Barzilai, R. Shamir, and Y. Shiloh. 2006. 'Parallel induction of ATM-dependent pro- and antiapoptotic signals in response to ionizing radiation in murine lymphoid tissue', *Oncogene*, 25: 1584-92.

Rass, Ulrich, Ivan Ahel, and Stephen C. West. 2007. 'Defective DNA Repair and Neurodegenerative Disease', *Cell*, 130: 991-1004.

Rogers, D. C., E. M. Fisher, S. D. Brown, J. Peters, A. J. Hunter, and J. E. Martin. 1997. 'Behavioral and functional analysis of mouse phenotype: SHIRPA, a proposed protocol for comprehensive phenotype assessment', *Mammalian Genome*, 8: 711-3.

Rothblum-Oviatt, C., J. Wright, M. A. Lefton-Greif, S. A. McGrath-Morrow, T. O. Crawford, and H. M. Lederman. 2016. 'Ataxia telangiectasia: a review', *Orphanet Journal of Rare Diseases*, 11: 159.

Sahama, Ishani, Kate Sinclair, Simona Fiori, James Doecke, Kerstin Pannek, Lee Reid, Martin Lavin, and Stephen Rose. 2015. 'Motor pathway degeneration in young ataxia telangiectasia patients: A diffusion tractography study', *NeuroImage: Clinical*, 9: 206-15.

Sahama, Ishani, Kate Sinclair, Kerstin Pannek, Martin Lavin, and Stephen Rose. 2014. 'Radiological imaging in ataxia telangiectasia: a review', *The Cerebellum*, 13: 521-30.

Sandoval, N., M. Platzer, A. Rosenthal, T. Dork, R. Bendix, B. Skawran, M. Stuhrmann, R. D. Wegner, K. Sperling, S. Banin, Y. Shiloh, A. Baumer, U. Bernthaler, H. Sennefelder, M. Brohm, B. H. Weber, and D. Schindler. 1999. 'Characterization of ATM gene mutations in 66 ataxia telangiectasia families', *Human Molecular Genetics*, 8: 69-79.

Sanghez, Valentina, Anna Luzzi, Don Clarke, Dustin Kee, Steven Beuder, Danielle Rux, Mitsujiro
Osawa, Joaquín Madrenas, Tsui-Fen Chou, Michael Kyba, and Michelina Iacovino. 2017.
'Notch activation is required for downregulation of HoxA3-dependent endothelial cell phenotype
during blood formation', *PloS One*, 12: e0186818.
Savitsky, K., A. Bar-Shira, S. Gilad, G. Rotman, Y. Ziv, L. Vanagaite, D. A. Tagle, S. Smith, T. Uziel, S.
Sfez, M. Ashkenazi, I. Pecker, M. Frydman, R. Harnik, S. R. Patanjali, A. Simmons, G. A.
Clines, A. Sartiel, R. A. Gatti, L. Chessa, O. Sanal, M. F. Lavin, N. G. Jaspers, A. M. Taylor, C.
F. Arlett, T. Miki, S. M. Weissman, M. Lovett, F. S. Collins, and Y. Shiloh. 1995. 'A single ataxia
telangiectasia gene with a product similar to PI-3 kinase', *Science*, 268: 1749-53.
Schellenberg, M. J., P. P. Tumbale, and R. S. Williams. 2015. 'Molecular underpinnings of Aprataxin
RNA/DNA deadenylase function and dysfunction in neurological disease', *Progress in*
Biophysics and Molecular Biology, 117: 157-65.
Schubert, R., J. Reichenbach, and S. Zielen. 2002. 'Deficiencies in CD4+ and CD8+ T cell subsets in
ataxia telangiectasia', *Clinical and Experimental Immunology*, 129: 125-32.
Sedghi, Maryam, Mehri Salari, Ali-Reza Moslemi, Ariana Kariminejad, Mark Davis, Hayley Goullée,
Björn Olsson, Nigel Laing, and Homa Tajsharghi. 2018. 'Ataxia-telangiectasia-like disorder in a
family deficient for MRE11A, caused by a -MRE11- variant', *Neurology Genetics*, 4: e295.
Shakkottai, Vikram G, Maria do Carmo Costa, James M Dell'Orco, Ananthakrishnan
Sankaranarayanan, Heike Wulff, and Henry L Paulson. 2011. 'Early changes in cerebellar
physiology accompany motor dysfunction in the polyglutamine disease spinocerebellar ataxia
type 3', *Journal of Neuroscience*, 31: 13002-14.
Shakkottai, Vikram G, Maolei Xiao, Lin Xu, Michael Wong, Jeanne M Nerbonne, David M Ornitz, and
Kelvin A Yamada. 2009. 'FGF14 regulates the intrinsic excitability of cerebellar Purkinje
neurons', *Neurobiology of Disease*, 33: 81-88.
Shiloh, Y., and Y. Ziv. 2013. 'The ATM protein kinase: regulating the cellular response to genotoxic
stress, and more', *Nature Reviews: Molecular Cell Biology*, 14: 197-210.
Shiloh, Yosef. 2020. 'The cerebellar degeneration in ataxia-telangiectasia: A case for genome
instability', *DNA Repair*, 95: 102950.
Solitare, G. B. 1968. 'Louis-Bar's syndrome (ataxia-telangiectasia). Anatomic considerations with
emphasis on neuropathologic observations', *Neurology*, 18: 1180-6.
Solitare, G. B., and V. F. Lopez. 1967. 'Louis-bar's syndrome (ataxia-telangiectasia). Neuropathologic
observations', *Neurology*, 17: 23-31.
Spring, K., S. Cross, C. Li, D. Watters, L. Ben-Senior, P. Waring, F. Ahangari, S. L. Lu, P. Chen, I.
Misko, C. Paterson, G. Kay, N. I. Smorodinsky, Y. Shiloh, and M. F. Lavin. 2001. 'Atm knock-in
mice harboring an in-frame deletion corresponding to the human ATM 7636del9 common
mutation exhibit a variant phenotype', *Cancer Research*, 61: 4561-8.
Staples, E. R., E. M. McDermott, A. Reiman, P. J. Byrd, S. Ritchie, A. M. Taylor, and E. G. Davies.
2008. 'Immunodeficiency in ataxia telangiectasia is correlated strongly with the presence of two
null mutations in the ataxia telangiectasia mutated gene', *Clinical and Experimental*
Immunology, 153: 214-20.
Stoyas, Colleen A, David D Bushart, Pawel M Switonski, Jacqueline M Ward, Akshay Alaghatta, Mi-bo
Tang, Chenchen Niu, Mandheer Wadhwa, Haoran Huang, and Alex Savchenko. 2020.
'Nicotinamide pathway-dependent Sirt1 activation restores calcium homeostasis to achieve
neuroprotection in spinocerebellar ataxia type 7', *Neuron*, 105: 630-44. e9.
Strich, Sabina J. 1966. 'Pathological findings in three cases of ataxia-telangiectasia', *Journal of*
Neurology, Neurosurgery, and Psychiatry, 29: 489.
Swift, M., D. Morrell, E. Cromartie, A. R. Chamberlin, M. H. Skolnick, and D. T. Bishop. 1986. 'The
incidence and gene frequency of ataxia-telangiectasia in the United States', *American Journal of*
Human Genetics, 39: 573-83.
Sykora, Peter, Deborah L. Croteau, Vilhelm A. Bohr, and David M. Wilson, 3rd. 2011. 'Aprataxin
localizes to mitochondria and preserves mitochondrial function', *Proceedings of the National*
Academy of Sciences of the United States of America, 108: 7437-42.

Takashima, H., C. F. Boerkoel, J. John, G. M. Saifi, M. A. Salih, D. Armstrong, Y. Mao, F. A. Quiocho,
B. B. Roa, M. Nakagawa, D. W. Stockton, and J. R. Lupski. 2002. 'Mutation of TDP1, encoding
a topoisomerase I-dependent DNA damage repair enzyme, in spinocerebellar ataxia with axonal
neuropathy', *Nature Genetics*, 32: 267-72.
Tal, E., M. Alfo, S. Zha, A. Barzilai, C. I. De Zeeuw, Y. Ziv, and Y. Shiloh. 2018. 'Inactive Atm abrogates
DSB repair in mouse cerebellum more than does Atm loss, without causing a neurological
phenotype', *DNA Repair (Amst)*, 72: 10-17.
Tavani, F., R. A. Zimmerman, G. T. Berry, K. Sullivan, R. Gatti, and P. Bingham. 2003. 'Ataxia-
telangiectasia: the pattern of cerebellar atrophy on MRI', *Neuroradiology*, 45: 315-9.
Taylor, A. M., Z. Lam, J. I. Last, and P. J. Byrd. 2015. 'Ataxia telangiectasia: more variation at clinical
and cellular levels', *Clinical Genetics*, 87: 199-208.
Terplan, K. L., and R. F. Krauss. 1969. 'Histopathologic brain changes in association with ataxia-
telangiectasia', *Neurology*, 19: 446-54.
Tsutsumi, S., M. Yamazaki, T. Miyazaki, M. Watanabe, K. Sakimura, M. Kano, and K. Kitamura. 2015.
'Structure-function relationships between aldolase C/zebrin II expression and complex spike
synchrony in the cerebellum', *Journal of Neuroscience*, 35: 843-52.
Tumbale, P., M. J. Schellenberg, G. A. Mueller, E. Fairweather, M. Watson, J. N. Little, J. Krahn, I.
Waddell, R. E. London, and R. S. Williams. 2018. 'Mechanism of APTX nicked DNA sensing
and pleiotropic inactivation in neurodegenerative disease', *EMBO Journal*, 37: e98875.
Vacchio, M. S., A. Olaru, F. Livak, and R. J. Hodes. 2007. 'ATM deficiency impairs thymocyte
maturation because of defective resolution of T cell receptor alpha locus coding end breaks',
Proceedings of the National Academy of Sciences of the United States of America, 104: 6323-8.
Vail, Graham, Aifang Cheng, Yu Ray Han, Teng Zhao, Shengwang Du, Michael M. T. Loy, Karl Herrup,
and Mark R. Plummer. 2016. 'ATM protein is located on presynaptic vesicles and its deficit
leads to failures in synaptic plasticity', *Journal of Neurophysiology*, 116: 201-09.
Valentin-Vega, Y. A., and M. B. Kastan. 2012. 'A new role for ATM: regulating mitochondrial function
and mitophagy', *Autophagy*, 8: 840-1.
van der Burgt, I, K H Chrzanowska, D Smeets, and C Weemaes. 1996. 'Nijmegen breakage syndrome',
Journal of Medical Genetics, 33: 153-56.
Verhagen, Mijke M.M., Jean-Jacques Martin, Marcel van Deuren, Chantal Ceuterick-de Groote, Corry
M.R. Weemaes, Berry H.P.H. Kremer, Malcolm A.R. Taylor, Michèl A.A.P. Willemsen, and
Martin Lammens. 2012. 'Neuropathology in classical and variant ataxia-telangiectasia',
Neuropathology, 32: 234-44.
Voogd, Jan, and Mitchell Glickstein. 1998. 'The anatomy of the cerebellum', *Trends in Neurosciences*,
21: 370-75.
Wakasugi, M., T. Sasaki, M. Matsumoto, M. Nagaoka, K. Inoue, M. Inobe, K. Horibata, K. Tanaka, and
T. Matsunaga. 2014. 'Nucleotide excision repair-dependent DNA double-strand break formation
and ATM signaling activation in mammalian quiescent cells', *Journal of Biological Chemistry*,
289: 28730-7.
Wallis, LI, PD Griffiths, SJ Ritchie, CAJ Romanowski, G Darwent, and ID Wilkinson. 2007. 'Proton
spectroscopy and imaging at 3T in ataxia-telangiectasia', *American journal of neuroradiology*,
28: 79-83.
Walter, Joy T, Karina Alvina, Mary D Womack, Carolyn Chevez, and Kamran Khodakhah. 2006.
'Decreases in the precision of Purkinje cell pacemaking cause cerebellar dysfunction and
ataxia', *Nature Neuroscience*, 9: 389-97.
Weyemi, Urbain, Christophe E. Redon, Towqir Aziz, Rohini Choudhuri, Daisuke Maeda, Palak R.
Parekh, Michael Y. Bonner, Jack L. Arbiser, and William M. Bonner. 2015. 'NADPH oxidase 4 is
a critical mediator in Ataxia telangiectasia disease', *Proceedings of the National Academy of
Sciences*, 112: 2121-26.
Xu, Y., and D. Baltimore. 1996. 'Dual roles of ATM in the cellular response to radiation and in cell
growth control', *Genes and Development*, 10: 2401-10.

Yamasaki, Miwako, Kouichi Hashimoto, and Masanobu Kano. 2006. 'Miniature Synaptic Events Elicited
by Presynaptic Ca^2+ Rise Are Selectively Suppressed by Cannabinoid Receptor Activation in
Cerebellar Purkinje Cells', *The Journal of Neuroscience*, 26: 86-95.
Zhou, Haibo, Zhanmin Lin, Kai Voges, Chiheng Ju, Zhenyu Gao, Laurens W. J. Bosman, Tom J. H.
Ruigrok, Freek E. Hoebeek, Chris I. De Zeeuw, and Martijn Schonewille. 2014. 'Cerebellar
modules operate at different frequencies', *Elife*, 3: e02536.
Zhu, L., B. Scelfo, F. Tempia, B. Sacchetti, and P. Strata. 2006. 'Membrane excitability and fear
conditioning in cerebellar Purkinje cell', *Neuroscience*, 140: 801-10.

8.0 Figure Legends

Figure 1. New A-T mouse models expressing clinically related PTCs. A) The *Atm* gene locus was targeted by homologous recombination of a targeting vector containing a modified NorCOMM cassette in intron one and the corresponding A-T PTC mutation in exon 3 to create the targeted *Atm*[R35X] and *Atm*[Q35X] ES cell lines. Following germline transmission of these alleles in mice, the floxed NorCOMM cassette was removed by Cre excision *in vivo* to produce the final *Atm*[R35X] and *Atm*[Q35X] mouse lines. **B)** Genotyping of A-T mouse models. PCR agarose gel of mouse DNA shows 151 bp wildtype (+) allele band and 241 bp Cre-excised targeted allele band. **C)** ATM levels were examined using immunoblot analyses of the spleen due to its high expression density in this tissue. Exemplar blots illustrate a gene dose effect of ATM protein expression in samples harvested from wildtype (+), heterozygous (R35X/+, Q35X/+), and homozygous *Atm*[R35X/R35X] (R35X) and *Atm*[Q35X/Q35X] (Q35X) mice as indicated. **D)** Breeding scheme schematic for double mutant and control mice for this study. **E)** *Atm*[R35X/R35X]*; Aptx*[-/-] mice develop an ataxia that at late stages results in a severe loss of motor coordination and ability to ambulate (see **Videos 1-4**). Abbreviations for panel **1**: **hβA**-human beta Actin promotor; **ΔTK1-**delta TK1, inactivated Thymidine Kinase 1; **T2A**-self-cleaving peptide sequence; **Neo**-Neomycin gene; **PGKpA**-Phosphoglycerate kinase poly A tail; **loxP**-recombination elements are show as a blue triangle; orientation of the Gateway **attB** recombination elements by an orange arrow; orientation of the genotyping **F** and **R** primers is shown by green and blue arrows respectively; and engineered PTC sites are shown in exon 3 by a red circle.

Figure 2. Health and survivability of single and double mutant mice. A) Left: The line color and symbol for each genotype is denoted and is consistent across all figures. **Right:** *Atm*[R35X/R35X]*; Aptx*[-/-] mice weighed significantly less than all control genotypes as indicated by the growth curves (± 95%

confidence interval; dotted lines). Growth curve ($Atm^{R35X/R35X}$; $Aptx^{-/-}$ vs. controls): Male k = 0.024 vs.

0.011-0.019, Y_{max} = 21.8 vs. 32.9-41.0 g, (n = 3 to 18); Female k = 0.030 vs. 0.017-0.022, Y_{max} = 16.9

vs. 23.3-31.3, (n = 2 to 19). Sum of squares F-test run across all curves: Male $F_{(12, 364)}$ = 30.5,

****p<0.0001, Female $F_{(12, 339)}$ = 28.3, ****p<0.0001. **B)** ATM-deficient mice, regardless of APTX

expression, displayed significantly lower survivability with ~55% of mice deceased by P400. No

statistical differences between ATM-deficient mice were detected. Moreover, a single wildtype copy of

the *Atm* gene was sufficient to prevent premature death (no statistical difference detected between

$Atm^{R35X/+}$; $Aptx^{-/-}$ and $Atm^{+/+}$; $Aptx^{+/+}$ mice). Log-rank (Mantel-Cox) tests across all ($\chi^2_{(6, 217)}$ = 48.4,

****p<0.0001), just the ATM deficient ($\chi^2_{(2, 217)}$ = 1.06, p=0.6), and single comparisons to wildtype (see

figure) were conducted. Total number of animals indicated in panel **C**. **C)** Pie charts illustrating that

ATM-deficient mice displayed a high prevalence of thymomas based on postmortem necropsies.

"Other" probable causes of death included enlarged livers and obstructed kidneys. "Missing" mice

were presumed dead and cannibalized by cage mates, cause of death unknown. **Figure 2-figure**

supplement 1, Figure 2-source data 1

Figure 2-figure supplement 1. Animal weight for each time point and genotype. A) The average

weights are plotted for each genotype at each of the indicated time points. 2-way ANOVA with age and

genotype as factors excluding the double mutant mice data. Male: $F_{(10, 226)}$= 5.6, p<0.0001; Female: $F_{(10, 197)}$= 7.3, p<0.0001. **B)** The survivability of each genotype of mice is plotted for male and female

individually.

Figure 3. $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice develop a progressive loss in motor coordination. A)

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice take a similar amount of time to descend a vertical pole at P45, 120, and 210,

but significantly longer at P400. These overall results were found to be similar for both male (left, n = 2

to 12) and female (right, n=4 to 12) mice. **B)** Consistent with the vertical pole test, the gait of

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice measured during ambulation on a Catwalk gait analysis system was

significantly different to controls by P400, but not before P210. This includes the percent of time a

mouse spends with 3 *vs.* 1, 2, or 4 paws on the ground and the speed and cadence during each run

across the platform. The effects of the two null mutations were generally similar between males (left, n

= 4 to 21) and females (right, n = 3 to 18). **C)** Behavioral data for male (blue) and female (pink)

Atm$^{R35X/R35X}$*; Aptx*$^{-/-}$ (dark purple, left) and *Atm*$^{+/+}$*; Aptx*$^{+/+}$ (orange, right) mice are plotted at P30 (left 2

columns) and P400 (right 2 columns). Left-right asymmetries of the horizontal bars indicate a difference

in performance between genotypes for the behavioral test listed in a column on the far left. A significant

difference in the time to right during the righting reflex at P8 was observed in both Male and Female

mice (bottom). **A** and **B** were examined via 2-way ANOVA with age and genotype as factors followed

by *potshot* Tukey's multiple comparison tests between *Atm*$^{R35X/R35X}$*; Aptx*$^{-/-}$ and each of the control

genotypes. Behavioral tests in **C** were examined using a non-parametric Kruskal Wallace followed by

postdocs Dunn's multiple comparisons tests. Symbol/color key: *Atm*$^{+/+}$*; Aptx*$^{+/+}$ (purple circle), *Atm*$^{+/+}$*;*

Aptx$^{-/-}$ (blue diamond), *Atm*$^{R35X/R35X}$*; Aptx*$^{+/+}$ (green triangle), *Atm*$^{R35X/R35X}$*; Aptx*$^{-/-}$ (orange square),

Atm$^{R35X/+}$*; Aptx*$^{-/-}$ (red inverted triangle) **Figure 3-figure supplement 1, Figure 3-source data 1**

Figure 3-figure supplement 1. *Atm*$^{R35X/R35X}$*; Aptx*$^{-/-}$ mice develop progressive ataxia. A)

Atm$^{R35X/R35X}$*; Aptx*$^{-/-}$ mice develop an ataxia that at late stages results in a severe loss of coordination

and ability to ambulate. **B)** Additional gait analyses measuring stride length and time of overlap of the

hindlimbs. Examined via 2-way ANOVA with age and genotype as factors followed by Tukey's multiple

comparison tests between *Atm*$^{R35X/R35X}$*; Aptx*$^{-/-}$ and each of the control genotypes. **C)** Behavioral deficits

are seen only in *Atm*$^{R35X/R35X}$*; Aptx*$^{-/-}$ mice across all behavioral tests and sexes. Behavioral tests were

examined using a non-parametric Kruskal Wallace followed by Dunn's multiple comparisons test.

Figure 4. The biophysical properties of PNs are significantly perturbed in *Atm*$^{R35X/R35X}$*; Aptx*$^{-/-}$

mice. A) Schematic diagram of intracellular recording from a single Purkinje neuron (PN) in an acute

cerebellar tissue slice preparation used to examine their biophysical properties. **B) Left:** Voltage-clamp

measurements of PN neuron membrane properties were made from a 1 s, -5 mV step pulse as

illustrated. **Right:** The membrane input resistance (R_m), time constant (τ), and capacitance (C_m) were

perturbed in *Atm*$^{R35X/R35X}$*; Aptx*$^{-/-}$ compared to *Atm*$^{+/+}$*; Aptx*$^{+/+}$ mice. **C)** Current clamp recordings of PN

action potentials (AP) after 2 nA step pulses from a -70 mV holding potential. PN action potentials

recorded from $Atm^{R35X/R35X}$; $Aptx^{-/-}$ fail to maintain constant firing and summary plots show that they

have lower 1st AP amplitudes, firing threshold, and area under the curve. **D) Top:** Example sEPSC

traces taken from a PN under voltage clamp at a -80 mV holding potential. **Bottom:** Median frequency

and amplitude data, along with the overall probability distribution function are plotted for both $Atm^{+/+}$;

$Aptx^{+/+}$ (n = 11) and $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (n = 11) mice. The frequency but not amplitude of PNs recorded

in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice was found to be perturbed. **E and F) Left:** Example traces of evoked

EPSCs recorded from PNs as a result of a 2-pulse stimulation (50 ms interval) of either parallel (**E**) or

climbing (**F**) fiber axons. Traces illustrate the first (A$_1$) and second (A$_2$) amplitude (normalized) and time

course of first decay (blue fitted line) of each synaptic response. **Right:** Summary plots of the paired-

pulse ratio. While parallel fiber paired-pulse facilitation was normal in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice, climbing

fiber paired-pulse depression and halfwidth was significantly perturbed compared to $Atm^{+/+}$; $Aptx^{+/+}$

mice. **G)** Schematic diagram of extracellular recording from a single Purkinje neuron (PN) in an acute

cerebellar tissue slice preparation. Example electrophysiological traces for $Atm^{+/+}$; $Aptx^{+/+}$ (purple, top)

and $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (orange, bottom) PNs in the medial area (i.e., vermis) of the cerebellum. **H)**

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ PN action potential firing frequency progressively decreased with age and was

significantly slower in comparison to all control genotypes expressing at least one copy of the Atm or

$Aptx$ gene. [$Atm^{+/+}$; $Aptx^{+/+}$ (n=52 to 59), $Atm^{+/+}$; $Aptx^{-/-}$ (n = 51 to 64), $Atm^{R35X/R35X}$; $Aptx^{+/+}$ (n = 39 to 52),

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ (n = 24 to 71), $Atm^{R35X/+}$; $Aptx^{-/-}$ (n = 69)] Data in **B** were compared using an ANOVA

(Kruskal-Wallis) followed by Dunn's multiple comparisons test, data in **D to F** were compared via

Welch's t-test, and data in **H** using a 2-way ANOVA followed by Holm-Šídák's multiple comparisons

test. Symbol/color key: $Atm^{+/+}$; $Aptx^{+/+}$ (purple circle), $Atm^{+/+}$; $Aptx^{-/-}$ (blue diamond), $Atm^{R35X/R35X}$; $Aptx^{+/+}$

(green triangle), $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (orange square), $Atm^{R35X/+}$; $Aptx^{-/-}$ (red inverted triangle) **Figure 4-**

figure supplement 1-5, Figure 4 source data 1

Figure 4-figure supplement 1. Current *vs.* voltage responses significantly differ between $Atm^{+/+}$;

$Aptx^{+/+}$ and $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice. A) PN voltage responses to various current steps between -500

and 2250 pA (250 pA steps) from a -70 mV holding current in $Atm^{+/+}$; $Aptx^{+/+}$ (top, purple) and

$Atm^{R35X/R35X}$; $Aptx^{-/-}$ (bottom, orange) mice. **(B)** I□V curves calculated from either max deflection ($V_{m\,max}$)

or steady state ($V_{m\,end}$) for $Atm^{+/+}$; $Aptx^{+/+}$ (purple) and $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (orange) mice. **C)** Various

measurements of the voltage response to -500 pA step pulse (blue box in **B**) in $Atm^{+/+}$; $Aptx^{+/+}$ (purple)

and $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (orange) mice. Significance was tested using a non-parametric Mann Whitney

test.

Figure 4-figure supplement 2. Mean PN firing frequency across the cerebellum. Average PN firing

frequency is plotted across the indicated locations at P45, 120, 210, and 400. Significance tested via 2-

way ANOVA with age and genotype as factors.

Figure 4-figure supplement 3. Mean PN firing frequency across genotype and sex. Average PN

firing frequency for all cells recorded from male and female mice is plotted for the indicated genotype.

No significant differences were observed between sex. 2-way ANOVA with age and sex as factors,

$Atm^{+/+}$; $Aptx^{+/+}$ ($F_{(1,\,751}$ = 1.15, p=0.3), $Atm^{+/+}$; $Aptx^{-/-}$ ($F_{(1,\,797)}$ = 1.10, p=0.3), $Atm^{R35X/R35X}$; $Aptx^{+/+}$ ($F_{(1,\,630)}$ =

0.17, p=0.7), $Atm^{R35X/R35X}$; $Aptx^{-/-}$ ($F_{(1,\,666)}$ = 1.10, p=0.4), t-test for P400 $Atm^{R35X/+}$; $Aptx^{-/-}$ (p=0.9)

Figure 4-figure supplement 4. Coefficient of Variation of PN firing frequency across the

cerebellum. Average CV of PN firing frequency is plotted across the indicated locations at P45, 120,

210, and 400. No significant differences (p<0.5) were detected across all areas using 2-way ANOVA

with age and genotype as factors.

Figure 4-figure supplement 5. Mean variation between PN firing intervals across the cerebellum.

Average CV2 of PN firing frequency is plotted across the indicated locations at P45, 120, 210, and 400.

No significant differences (p<0.5) were detected across all areas using 2-way ANOVA with age and

genotype as factors.

Figure 5. Cerebellar atrophy is associated with a progressive reduction in molecular layer width

and pathological changes in PN morphology but not PN cell death. A) Cartoon image of the brain

highlighting the dorsal forebrain and cerebellar surface. Area estimates from dorsal images of the brain

were used to determine the cerebellum to forebrain ratio allowing us to control for any differences in

overall size of the brain. We found the cerebellum decreased in size over age in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (n

= 5 to 10), but not control mice [$Atm^{+/+}$; $Aptx^{+/+}$ (n = 4 to 20), $Atm^{+/+}$; $Aptx^{-/-}$ (n = 4 to 12), $Atm^{R35X/R35X}$;

$Aptx^{+/+}$ (n = 6 to 16), $Atm^{R35X/+}$; $Aptx^{-/-}$ (n = 6)]. **B) Left:** Immunofluorescent images of parasagittal

cerebellar sections illustrating the length (blue arrows) of PN dendrites in the molecular layer (ML;

green) and width of the granule cell layer (GCL; magenta) in $Atm^{+/+}$; $Aptx^{+/+}$ and $Atm^{R35X/R35X}$; $Aptx^{-/-}$

mice (P400; medial cerebellar lobule VIII). scale bar = 50 μm. **Middle:** Summary graphs of molecular

(ML) and granule cell (GCL) layer width measurements averaged across all lobules in sections from the

medial, intermediate, and lateral parts of the cerebellum (P400). **Right:** Average ML and GCL widths at

different ages in the medial cerebellum (n = 5 to 7). **C) Left:** H&E stained, parasagittal cerebellar

sections from P400 $Atm^{+/+}$; $Aptx^{+/+}$ (top) and $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (bottom) mice. scale bar = 500 μm

Right: Magnified view (from white box) of PNs (white triangles) in cerebellar lobules 4/5 and 6. scale

bar = 50 μm, inset 10 μm **Right:** The average density of PNs across all lobules in the medial

cerebellum of $Atm^{+/+}$; $Aptx^{+/+}$ (n = 9) and $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (n = 7) P400 mice do not differ. **D) Left:**

Whole cerebellar, fluorescent images and magnified view of folia VIII (from white box). scale bar = 500

μm left, 75 μm right **Right:** Average density of Calbindin positive PNs across the whole cerebellum (n =

7 to 9). **E) Left:** Optically sectioned fluorescent images illustrate the smaller size of PNs in $Atm^{R35X/R35X}$;

$Aptx^{-/-}$ mice. **Right:** Plot of the average width of PN somas randomly sampled from across the

cerebellum (n = 5). scale bar = 10 μm **F) Left:** Images like in **E**. Abnormally large caliber PN dendrites

(inset) were observed in $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice (P400; medial cerebellar lobule VI). scale bar = 25 μm

main, 5 μm inset **Right:** Average width of primary and secondary PN dendrites measured at the midline

between the PN cell bodies and ML edge in $Atm^{+/+}$; $Aptx^{+/+}$ and $Atm^{R35X/R35X}$; $Aptx^{-/-}$ mice (n = 6).

Statistical significances were assessed via 2-way ANOVA with age and genotype as factors followed by

Holm-Šídák (**A** and **B right**) or Šídák (**B middle**) pairwise multiple comparisons test. Welch's *t*-test

used in **D**, **E,** and **F**. **Figure 5-figure supplements 1-5, Figure 5-source data 1**

Figure 5-figure supplement 1. Decreased molecular layer width but not cell death is a key

feature of the A-T model. A) Width measurements of the molecular and granule cell layer—ML and

GCL respectively—for each lobule across the medial intermediate and lateral areas of the cerebellum.

B) Left: Summary plot indicates no significant differences in PN density were observed across each

lobule in *Atm^{+/+}; Aptx^{+/+}* (n = 9) and *Atm^{R35X/R35X}; Aptx^{-/-}* (n = 7) mice**. Right:** Images of parasagittal

cerebellar sections from medial (top), intermediate (middle), and lateral (bottom) cerebellum. Scale bar

= 500 μm Statistical significances were assessed via 2-way ANOVA with age and genotype as factors

followed by Šídák post comparison test.

Figure 5-figure supplement 2. A) Fluorescent images of anti-microglial activation (CD68) staining in

Atm^{+/+}; Aptx^{+/+}, *Atm^{R35X/R35X}; Aptx^{-/-}*, and a positive control for cerebellar degeneration (mouse model of

MPS III; deficient in N-acetyl-alpha-glucosaminidase (*Naglu*) (Li et al. 1999). No positive staining for

CD68 was observed in the A-T mouse model, but extensive microglial activation (yellow) was seen

throughout the cerebellum of the *Naglu* knockout (*Naglu^{-/-}*). scale bar = 500 μm top, 50 μm bottom **B)**

Anti-cleaved Caspase-3 staining (yellow) in the cerebellum of *Atm^{+/+}; Aptx^{+/+}*, *Atm^{R35X/R35X}; Aptx^{-/-}*,

Naglu^{+/-}, and *Naglu^{-/-}* indicates there is no significant programmed cell death in the A-T mouse model as

compared to the *Naglu^{-/-}* which displays extensive PN loss. scale bar = 10 μm **C)** Additional fluorescent

images of large caliber dendrites (white arrows) in the A-T mouse model (bottom four images). scale

bar = 25 μm Significance was tested using a 1-Way ANOVA followed by Tukey's comparison test.

Figure 6. T-cell deficits are found in the blood of *Atm^{R35X/R35X}; Aptx^{-/-}* mice. A) Representative flow

cytometric profiles of T-cell glycoprotein marker CD3 and summary plots indicate ATM and/or APTX

deficient mice have decreased proportions of CD3^{+} T-cells in the blood. **B)** Representative flow

cytometric profiles of T-cell glycoprotein markers CD4 and CD8 gated on CD3^{+} cells and summary plots

for CD8 and CD4 single positive cell proportions. ATM deficient mice had reduced CD4^{+} proportions

compared to mice with at least one copy of the *Atm* gene. Statistical significances were assessed via 1-

way ANOVA followed by Tukey's pairwise multiple comparisons test. Number of animals denoted at

bottom of bar. Symbol/color key: $Atm^{+/+}$; $Aptx^{+/+}$ (purple circle), $Atm^{R35X/R35X}$; $Aptx^{+/+}$ (green triangle),

$Atm^{R35X/+}$; $Aptx^{-/-}$ (red inverted triangle), $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (orange square) **Figure 6-source data 1**

Figure 7. ATM and APTX deficiency confer deficits in T-cell expression, but at different

developmental stages. A) Representative flow cytometric profiles of T-cell glycoprotein markers CD44

and CD25 gated on CD4⁻CD8⁻ double negative (DN) cells. Summary plots show proportions of

thymocytes at DN stages 1-4 (left to right). APTX deficient mice display increased proportions for DN1-

3 and decreased proportion at DN4 consistent with a deficit in ontogeny from DN3 to DN4. **B)**

Representative flow cytometric profiles of T-cell glycoprotein markers CD4 and CD8 gated. ATM

deficient mice display decreased proportions for CD4 and CD8 single positive cells consistent with a

deficit in ontogeny from CD4⁺CD8⁺ double positive to CD4⁺ and CD8⁺ single positive fates. Statistical

significances were assessed via 1-way ANOVA followed by Tukey's pairwise multiple comparisons test.

Number of animals denoted at bottom of bars. Symbol/color key: $Atm^{+/+}$; $Aptx^{+/+}$ (purple circle),

$Atm^{R35X/R35X}$; $Aptx^{+/+}$ (green triangle), $At^{/R35X/+}$; $Aptx^{-/-}$ (red inverted triangle), $Atm^{R35X/R35X}$; $Aptx^{-/-}$ (orange

square) **Figure 7-source data 1**

Figure 8. ATM protein expression is restored after read-through compound exposure in explant

tissues from $Atm^{R35X/R35X}$ and $Atm^{Q35X/Q35X}$. Spleen and cerebellar explant tissue from $Atm^{R35X/R35X}$ and

$Atm^{+/+}$ mice were treated with vehicle, the read-through compounds G418 ($100 \,\mu M$) or GJ103 ($100 \,\mu M$)

for $72 \,hrs$. ATM immunoblots show recovery of ATM (MW $350 \,kDa$) production in both the spleen ($n = 3$)

and cerebellum ($n = 3$). Equal loading was assessed via housekeeping genes (Actin or GAPDH) and

ponceau staining. **Figure 8-source data**

11.0 Source files
Figure 1-source data 1. Original blots.
Figure 2-source data 1. Numerical data of weight, age of death, and probable cause of death
Figure 3-source data 1. Numerical data for animal behavioral assessments
Figure 4-source data 1. Numerical data of electrophysiological recordings for each panel
Figure 5-source data 1. Numerical data for histology
Figure 6-source data 1. Table of numerical FACs data for blood samples

Figure 7-source data 1. Table of numerical FACs data for thymus samples

Figure 8-source data 1. Original blots and numerical data

12.0 Rich Media

Video 1. Pole test, *Atm$^{+/+}$ vs. Atm$^{R35X/R35X}$*. *Atm$^{R35X/R35X}$* do not display an ataxic phenotype at P460.

Video 2. Pole test, *Aptx$^{+/+}$ vs. Aptx$^{-/-}$*. *Aptx$^{-/-}$* mice do not display an ataxic phenotype at P460.

Video 3. Pole test, *Atm$^{+/+}$; Aptx$^{+/+}$ vs. Atm$^{R35X/R35X}$; Aptx$^{-/-}$*. *Atm$^{R35X/R35X}$; Aptx$^{-/-}$* have considerable
motor disability at P460.

Video 4. Open field, *Atm$^{+/+}$; Aptx$^{+/+}$ vs. Atm$^{R35X/R35X}$; Aptx$^{-/-}$*. *Atm$^{R35X/R35X}$; Aptx$^{-/-}$* display a clear
inability to ambulate in the open field at P460.

1514

1515

1516

1517

1518

1519

Fig. 1

A Insertion of PTC mutation in *Atm* gene

B ATM gene

C ATM protein
Spleen

D

$Atm^{R35X/R35X}$ X $Aptx^{-/-}$ → $Atm^{R35X/+}; Aptx^{+/-}$ X →

$Atm^{+/+}; Aptx^{+/+}$

$Atm^{+/+}; Aptx^{-/-}$

$Atm^{R35X/R35X}; Aptx^{+/+}$

$Atm^{R35X/R35X}; Aptx^{-/-}$

$Atm^{R35X/+}; Aptx^{-/-}$

$Atm^{R35X/R35X}; Aptx^{+/-}$

E

$Atm^{+/+}; Aptx^{-/-}$

$Atm^{R35X/R35X}; Aptx^{-/-}$

Fig. 2

A

- *Atm*$^{+/+}$; *Aptx*$^{+/+}$
- *Atm*$^{+/+}$; *Aptx*$^{-/-}$
- *Atm*$^{R35X/R35X}$; *Aptx*$^{+/+}$
- *Atm*$^{R35X/R35X}$; *Aptx*$^{-/-}$
- *Atm*$^{R35X/+}$; *Aptx*$^{-/-}$
- *Atm*$^{R35X/R35X}$; *Aptx*$^{+/-}$
- *Atm*$^{R35X/+}$; *Aptx*$^{+/+}$

Male

Female

B Survival proportions

C Cause of death
- thymoma
- indeterminable
- other
- missing
- alive at 400d

Atm$^{+/+}$; *Aptx*$^{+/+}$ n=32

Atm$^{+/+}$; *Aptx*$^{-/-}$ n=33

Atm$^{R35X/+}$; *Aptx*$^{+/+}$ n=31

Atm$^{R35X/+}$; *Aptx*$^{-/-}$ n=37

Atm$^{R35X/R35X}$; *Aptx*$^{+/+}$ n=24

Atm$^{R35X/R35X}$; *Aptx*$^{+/-}$ n=20

Atm$^{R35X/R35X}$; *Aptx*$^{-/-}$ n=40

Fig. 3

A

Legend:
- $Atm^{+/+}$; $Aptx^{+/+}$
- $Atm^{+/+}$; $Aptx^{-/-}$
- $Atm^{R35X/+}$; $Aptx^{-/-}$
- $Atm^{R35X/R35X}$; $Aptx^{-/-}$
- $Atm^{R35X/R35X}$; $Aptx^{+/+}$

Pole test

Male / Female
Time to descend (s)

B

Gait Analyses

% standing on 3
Cadence (steps/s)
Average speed (cm/s)
Days Old

C

SHIRPA

30 Days old / 400 Days old

$Atm^{+/+}$ $Aptx^{+/+}$ / $Atm^{R35X/R35X}$ $Aptx^{-/-}$

Male / Female

Righting Reflex
Wire Manoeuvre
Grip Strength
Hindlimb Clasping
Ledge Test
Touch Escape
Tail Elevation
Pelvic Elevation
Gait
Startle Response
Transfer Arousal
Tremor
Respiration Rate
Spontaneous Activity
Body Position

score

Locomotor Activity
# of squares

Postnatal (P8) Righting Reflex
time to right (s)

A Acute slice 350-400 Day

PN   patch electrode

B
$Atm^{+/+}; Aptx^{+/+}$ $Atm^{R35X/R35X}; Aptx^{-/-}$

200 pA / 200 ms

-80 mV   - 5mV step pulse

Membrane properties

R_m (MΩ)   **

τ (ms)   **

C_m (pF)   **

C
-70 mV   2 nA step pulse   10 mV / 100 ms

APs/cell   Time from step pulse (ms)

1st AP Ampl. (ΔmV)   **

1st AP Thresh. (abs. mV)   *

1st AP area (mV*ms)   *

D sEPSC

10 pA / 100 ms

Amplitude (pA)

Probability / Amplitude (pA)
$Atm^{+/+}; Aptx^{+/+}$
$Atm^{R35X/R35X}; Aptx^{-/-}$

Frequency (Hz)   *

Probability / Frequency (Hz)

E Parallel fiber stimulation

A_1 A_2 5 ms

Pair Pulse Ratio (A_2/A_1)

F Climbing fiber stimulation

A_1 A_2 5 ms

Paired pulse ratio (A_2/A_1)   *

Halfwidth (ms)   **

G Acute slice

PN   extracellular electrode

$Atm^{+/+}; APTX^{+/+}$

$Atm^{R35X/R35X}; APTX^{-/-}$

100 ms

H
$Atm^{+/+}; Aptx^{+/+}$ $Atm^{R35X/R35X}; Aptx^{+/+}$ $Atm^{+/+}; Aptx^{-/-}$
$Atm^{R35X/+}; Aptx^{-/-}$ $Atm^{R35X/R35X}; Aptx^{-/-}$

Spont. AP freq. (Hz)

Lateral   Intermediate   Medial

Days old   Days old   Days old

*   ****   **

****   ****   ****   ****

Fig. 5

Fig. 6

Fig. 7

Fig. 8

Figure 2-figure supplement 1

Figure 3-figure supplement 1

Figure 4-figure supplement 1

Figure 4-figure supplement 2

Figure 4-figure supplement 3

Figure 4-figure supplement 4

Figure 4-figure supplement 5

Figure 5-figure supplement 1

Figure 5-figure supplement 2