Asymmetric requirement of Dpp/BMP morphogen dispersal in the Drosophila wing disc

Shinya Matsuda[*1], Jonas V. Schaefer[2], Yusuke Mii[3], Yutaro Hori[4], Dimitri Bieli[1], Masanori Taira[5], Andreas Plückthun[2], and Markus Affolter[*1]

Affiliations:

1. Biozentrum, University of Basel, Switzerland
2. Department of Biochemistry, University of Zurich, Switzerland
3. National Institute for Basic Biology and Exploratory Research Center on Life and Living Systems (ExCELLS), National Institutes of Natural Sciences, Japan
    JST, PRESTO, Japan
4. Institute for Quantitative Biosciences, The University of Tokyo, Japan
5. Department of Biological Sciences, Graduate School of Science, The University of Tokyo, Japan
    Present address: Department of Biological Sciences, Faculty of Science and Engineering, Chuo University, Japan

Corresponding authors

Correspondence to: markus.affolter@unibas.ch or shinya.matsuda@unibas.ch

Summary
How morphogen gradients control patterning and growth in developing tissues remains largely unknown due to lack of tools manipulating morphogen gradients. Here, we generate two membrane-tethered protein binders that manipulate different aspects of Decapentaplegic (Dpp), a morphogen required for overall patterning and growth of the Drosophila wing. "HA trap" is based on a single-chain variable fragment (scFv) against the HA tag that traps HA-Dpp to mainly block its dispersal, while "Dpp trap" is based on a Designed Ankyrin Repeat Protein (DARPin) against Dpp that traps Dpp to block both its dispersal and signaling. Using these tools, we found that, while posterior patterning and growth require Dpp dispersal, anterior patterning and growth largely proceed without Dpp dispersal. We show that *dpp* transcriptional refinement from an initially uniform to a localized expression and persistent signaling in transient *dpp* source cells render the anterior compartment robust against the absence of Dpp dispersal. Furthermore, despite a critical requirement of *dpp* for the overall wing growth, neither Dpp dispersal nor direct signaling is critical for lateral wing growth after wing pouch specification. These results challenge the long-standing dogma that Dpp dispersal is strictly required to control and coordinate overall wing patterning and growth.

Introduction

A fundamental question in developmental biology is how proteins work together to orchestrate developmental processes. Forward and reverse genetic approaches based on mutants and RNAi, together with biochemical analyses, provide insights into how proteins function. However, interpretational gaps often remain between the mutant phenotypes and the underlying mechanisms.

Recently, small, high affinity protein binders such as nanobodies, single-chain variable fragments (scFvs), Designed Ankyrin Repeat Proteins (DARPins) and others have emerged as versatile tools to fill this gap. By fusing these protein binders to well-characterized protein domains and expressing the fusion proteins *in vivo*, protein function can be directly manipulated in a predicted manner[1-5]. For example, when a protein functions with multiple parameters, protein binder tools targeting each or a subset of these parameters could help to dissect the requirement of each parameter. However, it remains challenging to design and customize such protein binder tools.

A class of molecules that exert its function with multiple parameters are morphogens, secreted molecules that disperse from a localized source and regulate target gene expression in a concentration-dependent manner[6-9]. A morphogen gradient is characterized by its parameters such as rates of secretion, diffusion, and degradation[10]. Temporal dynamics of a morphogen gradient also impact cell fates decisions[11]. Despite a variety of parameters involved, morphogen dispersal is generally thought to be critical for morphogen function based on severe morphogen mutant phenotypes and long-range action of morphogens to reorganize patterning and growth.

However, a recent study challenged this basic assumption for the case of the Wingless (Wg) morphogen, the main Wnt in Drosophila, by showing that a membrane-tethered non-diffusible form of Wg can replace the endogenous Wg without strongly affecting appendage development[12]. Although the precise contribution of Wg dispersal requires further investigations[13-16], the study raises the question of how important morphogen dispersal is for tissue patterning and growth in general.

In contrast to Wg, *Decapentaplegic (dpp)*, the vertebrate BMP2/4 homologue, is thought to act as a *bona fide* morphogen in the Drosophila prospective wing. Dpp disperses from a narrow anterior stripe of cells along the anterior-posterior (A-P) compartment boundary to establish a characteristic morphogen gradient in both compartments (Fig. 1a)[17,18]. How the Dpp dispersal-mediated morphogen gradient achieves and coordinates overall wing patterning and growth has served as a paradigm to study morphogens[19]. However, despite intensive studies, it remains controversial how Dpp/BMP disperses[17,20-23], controls growth[24-33], and coordinates patterning and growth (i.e. scaling)[32,34-36]. Regardless of the actual mechanisms, all the studies are based on the assumption that Dpp dispersal from the anterior stripe of cells controls overall wing patterning and growth, in line with the severe *dpp* mutant phenotypes.

To directly manipulate dispersal of Dpp, we recently generated morphotrap, a membrane-tethered anti-GFP nanobody, to trap GFP-tagged Dpp and thereby manipulate its dispersal[37]. Using morphotrap, the authors showed that a substantial amount of GFP-Dpp secreted from the anterior stripe of cells can reach the peripheral wing disc and that blocking GFP-Dpp dispersal from the source cells causes severe adult wing patterning and growth defects[37]. These results support the critical role of Dpp dispersal for overall wing patterning and growth[37]. However, application of morphotrap was limited to rescue conditions by overexpression of GFP-Dpp, due to the lack of an endogenous *GFP-dpp* allele.

In this study, we first generated a platform to manipulate the *dpp* locus and inserted a tag into *dpp* in order to investigate the precise requirement of the endogenous Dpp morphogen gradient for wing patterning and growth. We found that while a *HA-dpp* allele was functional, a *GFP-dpp* allele was not, thus limiting morphotrap application. To manipulate the endogenous Dpp morphogen gradient, we then generated two protein binder tools analogous to morphotrap. One is "HA trap" based on anti-HA scFv that traps HA-Dpp through the HA tag to mainly block Dpp dispersal, the other is "Dpp trap" based on anti-Dpp DARPin that directly binds to Dpp to block Dpp dispersal and signaling in the source cells. Thus, these tools allowed us to distinguish the requirements of Dpp dispersal and cell-autonomous signaling in the source cells for wing pouch growth and patterning.

Using these tools, we found that while posterior patterning and growth require Dpp dispersal,
anterior patterning and growth largely proceed without Dpp dispersal but require cell-autonomous Dpp
signaling in the source cells. We show that *dpp* transcriptional refinement from an initially uniform to
a localized expression and persistent signaling in transient *dpp* source cells allow relatively normal
anterior patterning and growth despite the absence of Dpp dispersal. Furthermore, despite a critical
requirement of *dpp* for overall wing growth, we also found that neither Dpp dispersal nor direct
signaling is critical for the lateral wing pouch to grow once the wing pouch is defined. These results
challenge the long-standing dogma that Dpp dispersal controls overall wing patterning and growth and
call for a revision of how Dpp controls and coordinates wing patterning and growth.
98
Results
A platform to manipulate the endogenous *dpp* locus
To manipulate the endogenous Dpp morphogen gradient, we utilized a MiMIC transposon inserted in
the *dpp* locus (*dpp^MI03752*), which allows to replace the sequence between the two *attP* sites in the
transposon with any sequence inserted between two inverted *attB* sites upon integrase expression[38]. A
genomic fragment containing sequences encoding a tagged version of *dpp* followed by an FRT and a
marker was first inserted into the locus (Fig. 1b), then the endogenous *dpp* exon was removed upon
FLP/FRT recombination to keep only the tagged *dpp* exon (Fig. 1b). Using this strategy, we inserted
different tags into the *dpp* locus and found that while a *GFP-dpp* allele was homozygous lethal during
early embryogenesis (data not shown), a *HA-dpp* allele was functional without obvious phenotypes[39]
(Fig. 1c). Immunostainings for the HA-tag including permeabilization steps showed HA-Dpp
expression in an anterior stripe of cells along the A-P compartment boundary in the late third instar
wing disc (Fig. 1d). In contrast, immunostainings for the HA-tag without permeabilization, which
allows antibodies to access only the extracellular antigens, revealed that a shallow extracellular HA-
Dpp gradient overlapped with the gradient of phosphorylated Mad (pMad), a downstream transcription
factor of Dpp signaling (Fig. 1e). Similar HA tag knock-in *dpp* alleles have recently been generated
by a CRISPR approach[25].
116
Generation and characterization of HA trap
Since we could not apply morphotrap due to the lethality of the *GFP-dpp* allele, we generated a "HA
trap", analogous to morphotrap. HA trap consists of an anti-HA scFv[40] fused to the transmembrane
domain of CD8 and mCherry (Fig. 2a). HA trap expression in the anterior stripe of cells of wild type
wing discs using *ptc-Gal4* did not interfere with Dpp signaling in the wing disc or patterning and
growth of the adult wing (Extended Data Fig. 1). Thus, HA trap is inert in the absence of a HA-tagged
protein. While we attempted to visualize extracellular HA-Dpp distribution upon HA trap expression,
we noticed that the HA tag can no longer be used for immunostaining when bound to HA trap. We
therefore additionally inserted an Ollas tag to generate a functional *Ollas-HA-dpp* allele in order to
visualize the extracellular Dpp distribution using the antibody against the Ollas tag. The extracellular
Ollas-HA-Dpp gradient was similar to the extracellular HA-Dpp gradient (Fig. 2b, c).
128
To test if HA trap can efficiently trap Ollas-HA-Dpp in the Dpp producing cells, HA trap was
expressed in the anterior stripe of cells using *ptc-Gal4* in Ollas-HA-Dpp heterozygous wing discs,
since *ptc*-Gal4 expression largely overlaps with *dpp* producing cells[41]. Under this condition,
extracellular immunostainings for the Ollas-tag revealed that Ollas-HA-Dpp accumulated on the
anterior stripe of cells, and that the extracellular gradient was abolished (Fig. 2d, e). To test if HA trap
can trap Ollas-HA-Dpp outside the anterior stripe of cells, clones of cells expressing *Gal4* were
randomly induced by heat-shock inducible FLP to express HA trap under UAS control. We found that
Ollas-HA-Dpp accumulated in clones of cells expressing HA trap induced outside the main *dpp* source
cells in both compartments (Fig. 2f-i, arrow). If HA trap can efficiently trap Ollas-HA-Dpp in the
source cells, the clonal Ollas-HA-Dpp accumulation should be blocked upon HA trap expression in

the source cells. Indeed, we found that clonal Ollas-HA-Dpp accumulation in both compartments was
drastically reduced upon HA trap expression using *ptc-Gal4* (Fig. 2j-k, arrow), indicating that the HA
trap can block HA-Dpp dispersal efficiently.
142
143   It has been shown that overexpression of GFP-Dpp from the anterior stripe cells leads to accumulation
144   of GFP-Dpp in clones of cells expressing morphotrap in the peripheral regions[37]. In contrast, we found
145   that Ollas-HA-Dpp accumulated in clones of cells expressing HA trap near the source cells but not in
146   the peripheral regions (Fig. 2l arrowhead). This raises a question whether Dpp can act in the peripheral
147   regions at physiological levels.
148

Asymmetric patterning and growth defects by HA trap

After we validated that HA trap can efficiently block Dpp dispersal, we then expressed HA trap using
different Gal4 driver lines in *HA-dpp* homozygous wing discs to address the requirement of Dpp
dispersal. Normally, Dpp binds to the Dpp receptors Thickveins (Tkv) and Punt, inducing a pMad
gradient and an inverse gradient of Brk, a transcription repressor repressed by Dpp signaling. The two
opposite gradients regulate growth and patterning (nested target gene expression, such as *sal*, and *omb*)
to define adult wing vein positions (such as L2 and L5)(Fig. 3a)[10,19,42-44]. Upon HA trap expression in
the anterior stripe of cells using *ptc-Gal4*, pMad, Sal, and Omb expression were undetectable in the P
compartment and Brk was also upregulated in the P compartment (Fig. 3b, e, f, g, h), indicating that
HA trap efficiently blocked HA-Dpp dispersal from source cells and interfered with patterning. The
posterior wing pouch growth was also affected as revealed by the expression of an intervein marker
DSRF and a wing pouch marker *5xQE.DsRed*[45] (Fig. 3b arrow, 3i). Interestingly, although
5xQE.DsRed contains five copies of the 806 bp Quadrant Enhancer (QE) of the wing master gene *vg*
containing a Mad binding site and is therefore thought to be directly regulated by Dpp signaling[46,47],
5xQE.DsRed remained expressed in the P compartment without detectable Dpp signaling (Fig. 3b
arrow). In the A compartment, pMad was slightly reduced in the anterior medial region (Fig. 3b, e),
probably because HA trap partially blocked Dpp signaling upon binding to HA-Dpp (Fig. 2l arrow).
Nevertheless, the anterior Brk gradient was not strongly affected (Fig. 3b, f). Although maximum
intensity of Sal or Omb was reduced, nested expression of Sal and Omb, and growth were maintained
in the A compartment (Fig. 3b, g, h, i). Consistent with these phenotypes in the wing discs, while
posterior patterning and growth were severely affected, anterior patterning and growth were relatively
normal in the resulting adult wings (Fig. 3c, d, j). Similar defects were observed upon HA trap
expression in the region covering the entire wing pouch using *nub-Gal4* (Fig. 3k-t) or in the entire
anterior compartment using *ci-Gal4* (Extended Data Fig. 2a-j). Even when HA trap was expressed
using both *nub-Gal4* and *ptc-Gal4*, the resulting phenotypes were not enhanced (Extended Data Fig.
3). To test whether the posterior growth defects upon HA trap expression is caused by cell death,
Caspase-3 was analyzed. We found that Caspase-3 was not upregulated (Extended Data Fig. 4a, b, d),
and blocking apoptosis by apoptosis inhibitor p35 did not rescue these growth defects upon HA trap
expression (Extended Data Fig. 4e-g). Thus, the posterior growth defects upon HA trap expression is
not caused by cell death. Taken together, these results suggest that, while critical for posterior
patterning and growth, Dpp dispersal is largely dispensable for anterior patterning and growth.
180

Lateral wing pouch growth without Dpp signaling

A critical role of Dpp dispersal for posterior patterning and growth is consistent with a role of Dpp as
a morphogen. However, the overall phenotypes caused by HA trap was surprisingly mild when
compared to the phenotypes seen in *dpp* mutants (Fig. 6a, c). Given the requirement of Dpp signaling
for cell proliferation and survival in the entire wing pouch[48], it was surprising that about 40% of the
posterior wing pouch was able to grow and differentiate into adult wing tissue without detectable Dpp
signaling (Fig. 3d, i, j).
188

We therefore tested whether the posterior growth and *5xQE.DsRed* expression seen upon HA trap
expression is caused by low levels of HA-Dpp leaking from the HA trap expressed in the source. In
this case, the posterior growth and *5xQE.DsRed* expression seen upon HA trap expression should be
dependent on *tkv*, an essential receptor for Dpp signaling. To test this, mutant clones of *tkv^a12*
(characterized as a null allele[49,50]) were induced in wing discs expressing HA trap with *ptc-Gal4*
between mid-second and beginning of third instar stages and analyzed in the late third instar stage. We
found that *tkv^a12* clones survived and expressed the *5xQE.DsRed* reporter in the anterior lateral regions
as well as in the entire posterior region. We also noticed that *tkv^a12* clones survived and expressed the
5xQE.DsRed reporter even next to the source cells in the P compartment (Fig. 4a). These results
indicate that the lateral growth and *5xQE.DsRed* expression seen upon HA trap expression is
independent of Dpp signaling, and not caused by a leakage of HA-Dpp from the HA trap, even if such
leakage would occur.
201
To test whether Dpp signaling-independent growth occurs also during normal development, *tkv^a12*
clones were induced in the wild type wing disc during mid-second and early third instar stage. We
found that *tkv^a12* clones were eliminated from the medial regions but often survived and expressed the
5xQE.DsRed reporter in the lateral wing pouch (Fig. 4b). Since *tkv^a12* may not be a complete null allele,
we then inserted an FRT cassette in the *tkv* locus and generated a new *tkv* flip-out allele (*tkvHA^FO*) to
induce FLP/FRT mediated excision of *tkv*. By generating *tkv* null clones upon heat-shock inducible
FLP expression, we confirmed that *tkv* null clones often survived and expressed the *5xQE.DsRed*
reporter in the lateral wing pouch (Fig. 4c, d arrow). We also found that, while most often eliminated,
medial *tkv* null clones survived and expressed *5xQE.DsRed* in rare cases (Fig. 4d), indicating that Dpp
signaling is dispensable for *5xQE.DsRed* expression also in the medial region, but medial cells lacking
Dpp signaling are normally eliminated[48].
213
How can Dpp signaling-independent wing pouch growth and *5xQE.DsRed* expression be reconciled
with a critical role of Dpp signaling for the entire wing pouch growth[48]? First, *tkv* clones generated in
the developing wing pouch have been shown to be eliminated by apoptosis or extrusion and do not
survive in the adult wing[48,51]. However, *tkv* clones survive better in the P compartment where Dpp
signaling is blocked by HA trap (Fig. 4a) and in the lateral region of wild type wing disc where Dpp
signaling is generally low (Fig. 4b-d). This raises a possibility that *tkv* clones are eliminated when
surrounded by wild type cells, even if *tkv* clones could grow and survive to a certain extent. Second,
wing pouch and *5xQE.DsRed* expression were completely lost in *dpp* mutants (Fig. 6a, Extended Data
Fig. 7a). It has been shown that initial wing pouch specification is mediated by Dpp derived from the
peripodial membrane, which covers the developing wing pouch, and this early *dpp* expression in the
peripodial membrane is lost in *dpp* disc alleles[52]. Thus, wing pouch and *5xQE.DsRed* expression could
be lost in *dpp* disc alleles due to failure of initial specification of the wing disc and subsequent
elimination of cells.
227
To minimize these potential problems, we applied Gal80ts to conditionally remove *dpp* from the entire
A compartment using *ci-Gal4*. At the permissive temperature of 18°C, Gal80ts actively represses Gal4
activity. At restrictive temperature of 29°C, Gal80ts can no longer block Gal4 activity, thus Gal4 can
be temporally activated using temperature shifts. Upon FLP expression, *dpp* was removed by FLP/FRT
mediated excision via *dpp^FO* allele[24], in which a FRT cassette was inserted into the *dpp* locus. To
remove *dpp* from the beginning of second instar stage when the wing pouch is specified, the larvae
were raised at 18°C for 4 days and then shifted to 29°C. By removing *dpp* from the entire A
compartment using *ci-Gal4* under this condition, we found that *5xQE.DsRed* remained expressed
despite severe growth defects in the late third instar stage (Fig. 4e-h). Similarly, genetic removal of *tkv*
via *tkvHA^FO* from the A compartment using *ci-Gal4* or from the P compartment using *hh-Gal4* from
the second instar stage revealed that, despite severe growth defects, *5xQE.DsRed* remained expressed

in each compartment lacking *tkv* (Extended Data Fig. 5). Surprisingly, similar results were obtained
even when *tkv* was removed from the entire P compartment using *hh-Gal4* from the embryonic stages
without Gal80ts (Extended Data Fig. 6). These results further support the presence of Dpp signaling-
independent *5xQE.DsRed* expression and wing pouch growth.

How is *5xQE.DsRed* expression regulated if QE is not directly regulated by Dpp signaling? While
5xQE.DsRed expression is completely lost in *dpp* mutants, we found that *5xQE.DsRed* reporter
expression was rescued in *dpp*, *brk* double mutant wing discs (Extended Data Fig. 7), indicating that
5xQE.DsRed expression is largely induced by repressing *brk*, similar to the regulation of other *dpp*
target genes. Indeed, QE has also been shown to be activated in *brk* mutant clones in the wing disc[53].
However, this notion appears inconsistent with the fact that *5xQE.DsRed* expression was not repressed
in the region where Brk is high in various conditions, in which Dpp signaling is compromised (Fig.
3b, 4h, Extended Data Fig. 5d', 5h'). We noticed that the observed high Brk levels upon Dpp trapping
were comparable to the Brk level in the lateral region of the control wing disc (Fig. 3f, 3p), and Brk
and *5xQE.DsRed* were co-expressed in the lateral region of the control wing disc (Fig. 4f, Extended
Data Fig. 5b', 5f'). Thus, we speculate that Brk is not sufficient to repress *5xQE.DsRed* expression at
physiological levels in lateral regions and that there are additional inductive inputs. Consistent with
this idea, QE has also been shown to be regulated by Wg[45,54].

Severe patterning and growth defects by Dpp trap
Even if the lateral wing pouch region can grow independent of Dpp signaling after wing pouch
specification (Fig. 4e-h), this growth cannot account for the overall minor growth phenotypes caused
by HA trap (Fig. 3, Extended Data Fig. 2a-j). How can relatively normal patterning and growth be
achieved without Dpp dispersal? Since pMad was completely lost in *dpp* mutants (Fig. 4g) but
remained active in the source cells upon HA trap expression (Fig. 3, Extended Data Fig. 2), we asked
whether Dpp signaling in the source cells could account for the minor phenotypes caused by HA trap.

To test this, we selected DARPins, protein binders based on ankyrin repeats[55-57], that bind to the mature
Dpp ligand and block Dpp signaling. For each of the 36 candidates obtained from the *in vitro* screening,
we generated a "Dpp trap" by fusing the anti-Dpp DARPin to the transmembrane domain of CD8 and
mCherry (Fig. 5a). By expressing each trap in the wing disc, we identified one Dpp trap (containing
DARPin 1242_F1), which efficiently blocked Dpp dispersal (Fig. 5b) and signaling (Fig. 5d, 5l,
Extended Data Fig. 2k). We found that the expression of the Dpp trap using *ptc-Gal4* (Fig. 5c-i), *nub-*
Gal4 (Fig. 5k-t), and *ci-Gal4* (Extended Data Fig. 2k-t) caused severe patterning and growth defects,
similar to *dpp* mutants (Fig. 4g, h). Adult wings expressing Dpp trap using *nub-Gal4* were recovered
and also showed severe patterning and growth defects comparable to *dpp* mutants (Fig. 5n). Although
Caspase-3 was upregulated upon Dpp trap expression (Extended Data Fig. 4a, c, d), the growth defects
were not rescued by p35 (Extended Data Fig. 4h-j), indicating that apoptosis was not the main cause
of growth defects caused by Dpp trap. These severe phenotypes were not due to a common scaffold
effects of DARPins, since one of the traps (containing DARPin 1240_C9) that failed to trap Dpp did
not interfere with pMad accumulation in the wing disc or patterning and growth of the adult wing when
expressed using *ptc-Gal4* (Extended Data Fig. 8).

We note that upon Dpp trap expression, Sal expression was lost from the medial region but appeared
to be upregulated in the lateral region (Fig. 5d, l, Extended Data Fig. 2o, arrow). It has previously been
shown that Sal is expressed not only in the medial region but also in the lateral region[58]. The same
study also showed that the medial Sal expression is Dpp signaling-dependent but lateral Sal expression
is Brk-dependent. Thus, upregulation of Brk upon Dpp trap expression could cause the lateral Sal
upregulation. However, when we focused on the peripheral region of the control wing disc (basal
confocal section), we noticed that the lateral Sal expression of the control wing disc was actually

comparable to that of the wing disc expressing Dpp trap (Extended Data Fig. 9a, b). Thus, when we
focused on the medial Sal expression (apical confocal section), the lateral Sal expression of the control
wing disc was simply missed due to the tissue architecture. Consistently, when *dpp* was removed from
the entire A compartment using *ci-Gal4* from mid-second instar stage, Sal expression was lost from
the medial region but not significantly upregulated in the lateral region (Extended Data Fig. 9c-e).

By comparing the phenotypes caused by Dpp trap and HA trap, we noticed that the phenotypes caused
by Dpp trap were much stronger than those caused by HA trap (Fig. 3, Fig. 5, Extended Data Fig. 2).
Comparison of each compartment size when expressing HA trap and Dpp trap using different Gal4
driver lines also showed that each compartment size was smaller upon Dpp trap expression than upon
HA trap expression (Fig. 5j, u, Extended Data Fig. 2u). To test if the difference could be due to more
efficient blocking of Dpp dispersal by Dpp trap than by HA trap, each trap was expressed in the anterior
stripe of cells using *ptc-Gal4* and posterior pMad signal was analyzed. We found that HA trap blocked
posterior pMad signal more efficiently than Dpp trap (Extended Data Fig. 10), indicating that Dpp trap
actually blocks Dpp dispersal less efficiently than HA trap. Thus, the severe phenotypes caused by
Dpp trap are likely because Dpp trap blocks Dpp signaling more efficiently than HA trap. Interestingly,
despite the slight leakage of Dpp from the Dpp trap (Extended Data Fig. 10), anterior Dpp trap
expression caused more severe posterior growth defects than HA trap (Fig. 5j, Extended Data Fig. 2u),
indicating that anterior Dpp signaling is non-autonomously required for the posterior growth. We note
that, even though the anterior Dpp signaling was eliminated, genetic removal of *tkv* from the A
compartment using *ci-Gal4* did not interfere with posterior growth as severe as Dpp trap (Extended
Data Fig. 5c, d), probably because Dpp secreted from the A compartment can disperse to control
posterior growth. Taken together, these results suggest that Dpp signaling in the source cells is required
for a majority of patterning and growth seen upon HA trap expression.

Rescue of *dpp* mutants by cell-autonomous Dpp signaling
Our results so far suggest that, while the requirement for Dpp dispersal is relatively minor and
asymmetric, Dpp signaling in the source cells is critically required for the majority of patterning and
growth seen upon blocking Dpp dispersal. This raises the question of how cell-autonomous Dpp
signaling in the source cells can control patterning and growth outside the anterior stripe of cells, the
main *dpp* source cells. There are a couple of possible scenarios; if *dpp* expression is successively
restricted to the anterior stripe of cells during development, the anterior stripe of cells may retain and
deliver the earlier Dpp signaling to the peripheral region after they leave from the anterior stripe of
cells via proliferation[59], or downstream factor(s) of Dpp signaling in the anterior stripe of cells may
act non-autonomously to control patterning and growth outside the stripe of cells. Alternatively, *dpp*
expression may not be restricted to the anterior stripe of cells in the early stages, similar to what has
been shown in the case of *wg*[12].

Before we address these possibilities, we first asked how important cell-autonomous Dpp signaling in
the source cells is for wing pouch patterning and growth. If the relatively mild phenotypes caused by
HA trap are due to cell-autonomous Dpp signaling in the source cells, a constitutively active version
of Tkv (TkvQD)[50] expressed in the anterior stripe of cells using *dpp-Gal4* should rescue *dpp* mutants
to an extent mimicking the phenotypes caused by HA trap (Fig. 3). Indeed, under this condition, pMad
activation in the anterior stripe of cells rescued nested Sal and Omb expression in the A compartment
(Fig. 6a, b). Interestingly, growth, but not patterning, was also partially rescued in the P compartment
as indicated by DSRF and *5xQE.DsRed* expression (Fig. 6b arrow), thus indeed mimicking phenotypes
caused by HA trap (Fig. 3). Unfortunately, the resulting adult flies were not recovered at 25 °C.
However, when the temperature was shifted from 25 °C to 18 °C during mid- to late-third instar stages
in order to reduce Gal4 activity during pupal stages, rare survivors were recovered from the pupal
cases or managed to hatch although they died shortly after hatching. In such survivors, although the

anterior wing veins tends to be affected, probably due to continuous TkvQD expression during pupal
stages, the anterior growth was largely rescued and the posterior growth was also partially rescued,
similar to phenotypes caused by HA trap (Fig. 6c, d). These results suggest that the phenotypes caused
by HA trap expression largely depend on cell-autonomous Dpp signaling in the source cells.
343
How can cell-autonomous Dpp signaling in the source cells control posterior growth if *dpp* expression
is restricted to the A compartment? One trivial possibility is that the posterior growth was induced by
non-specific *dpp-Gal4* expression in the P compartment. To test this, *dpp-Gal4* was converted into a
ubiquitous *LexA* driver to express TkvQD permanently in lineage of *dpp-Gal4* (Fig. 6e). In this setup,
pMad was constitutively activated in all the cells where *dpp-Gal4* has been expressed, including cells
expressing *dpp-Gal4* non-specifically. Under this condition, pMad, Sal, and Omb were uniformly
upregulated in the A compartment, and Brk was completely lost in the entire A compartment (Fig. 6f),
indicating that *dpp-Gal4* has been expressed in the entire A compartment[60]. In contrast, pMad was not
activated in the P compartment, but *5xQE.DsRed* was still induced in the P compartment (Fig. 6f
arrow), indicating that non-autonomous posterior growth control by anterior Dpp signaling is
permissive rather than instructive (see discussion).
355

Initial uniform *dpp* transcription in the anterior compartment
Next, we asked how cell-autonomous Dpp signaling in the source cells can control anterior patterning
and growth. The uniform lineage of Dpp producing cells in the A compartment[60] (Fig. 6f) suggests
two possibilities; either *dpp* expression is always restricted to the anterior stripe of cells but the lineage
of these cells can cover the lateral region via proliferation[59], or earlier *dpp* expression covers the entire
A compartment as in the case of *wg*[12]. Since the existing *dpp-Gal4* line is derived from a fragment of
the *dpp* disc enhancer inserted outside the *dpp* locus, we first generated an endogenous *dpp-Gal4* line
using our platform (Fig. 7a). We traced its lineage with G-TRACE analysis, in which RFP expression
labels the real-time Gal4-expressing cells and GFP expression labels the entire lineage of the Gal4-
expressing cells (Fig. 7b). We found that the lineage of Dpp producing cells indeed covers the entire
A compartment (Fig. 7c).
367
To distinguish between the two possibilities mentioned above, we then generated a *dpp* transcription
reporter line by inserting a destabilized GFP (half-life <2 hrs) into the *dpp* locus (Fig. 7d). Consistent
with the latter possibility, we found that this *dpp* transcription reporter was uniformly expressed in the
entire A compartment until the early third instar stage (Fig. 7e, e', f, f') and refined to an anterior stripe
expression during much of the third instar stage (Fig. 7g, g', h, h'). To directly follow *dpp* transcription,
we also performed smFISH using RNAscope technology to visualize *dpp* transcripts *in situ*. Consistent
with the dynamic expression of the *dpp* transcription reporter, we found uniform anterior *dpp*
transcription until the early third instar stage (Fig. 7i, i', j, j') and an anterior stripe of *dpp* transcription
in the later stages (Fig. 7k, k', l, l'). Despite the initial broad anterior *dpp* expression, we found that
pMad signal is low in the middle of the wing disc and graded toward the lateral regions, similar to the
pMad gradient in the later stages (Extended Data Fig. 11).
379

Transient *dpp* soure outside Sal domain is required for anterior patterning and growth
The earlier anterior *dpp* source outside the anterior stripe of cells could provide a local *dpp* source to
control anterior patterning and growth when Dpp dispersal is blocked. However, since *ptc-Gal4* is also
initially expressed in the entire A compartment[41], the relatively minor defects by HA trap could be due
to the perdurance of Dpp signaling via artificially stabilized Dpp by HA trap. To avoid HA trap
expression in the entire A compartment, we applied *tubGal80ts* to express HA trap using *ptc-Gal4* at
defined time points in the A compartment. To do so, the larvae were raised at 18 °C until a temperature
shift to 29 °C to induce *Gal4* expression (Extended Data Fig. 12a). Upon HA trap expression from the
mid-second instar stage, the lineage of *ptc-Gal4* covered at most the anterior Sal domain (Fig. 8b),

which corresponds to the region between L2 and L4 in the adult wing. We found that the later the
temperature shift, the milder the posterior growth defects (Extended Data Fig. 12a, b). In contrast, the
A compartment size (between L1 and L4) remained rather normal, independent of the timing of the
temperature shift (Extended Data Fig. 12a, c). Interestingly, the size of peripheral regions (between L1
and L2) and the specification of L2 were not affected, independent of the timing of the temperature
shift (Extended Data Fig. 12a, d). These results are consistent with a role of an anterior lateral *dpp*
source for patterning and/or growth in the anterior lateral regions.
396
To directly test this, we applied Gal80ts to genetically remove *dpp* via *dpp*[FO] allele upon FLP
expression using *ptc-Gal4* from the mid-second instar stage. To do so, the larvae were raised at 18 °C
for 5 days before a temperature shift to 29 °C and were then dissected 48 hr later. In this setup, *dpp*
was removed approximately from the anterior Sal region, where cells in which the FRT cassette was
removed were marked by lacZ staining (Fig. 8a-f). Given that it takes about 20 hr to eliminate the
majority of Dpp protein under this condition[24,41], wing pouches are devoid of the majority of the Dpp
protein derived from the anterior stripe of cells for 28 hr at 29°C until they reach the late third instar
stage, which corresponds to a lack of Dpp protein secreted from the main source from early third instar
stages onward.
406
Under this setup, we found that pMad, Sal, and Omb were significantly reduced in the P compartment,
consistent with the removal of the main *dpp* source (Fig. 8d-f). In contrast, in the A compartment, low
levels of pMad persisted and Brk remained graded with lowest expression outside the lacZ positive
region (Fig. 8d arrow). As a consequence, while Sal was completely lost (Fig. 8e), weak Omb remained
expressed in the A compartment (Fig. 8f). Consistent with a critical role of the *dpp* stripe for wing
pouch growth[25,41,61], both anterior and posterior growth were affected (Fig. 8k).
413
To test if the remaining anterior Dpp signaling activity is due to Dpp produced outside the anterior Sal
domain, we then compared removal of *dpp* from the anterior Sal domain using *ptc-Gal4* (the same
setup above) with removal of *dpp* from the entire A compartment using *ci-Gal4* from the mid-second
instar stage. We found that the anterior weak Dpp signaling and Omb expression seen upon removal
of *dpp* from the anterior Sal domain using *ptc-Gal4* (Fig. 8g, h) was completely lost by removing *dpp*
from the entire A compartment using *ci-Gal4* (Fig. 8i, j). Furthermore, anterior growth defects upon
removal of *dpp* from the anterior Sal domain was further enhanced upon removal of *dpp* from the
entire A compartment (Fig. 8k), indicating that the anterior *dpp* source outside the anterior Sal domain
is locally required for anterior Dpp signaling and growth.
423
How can the transient *dpp* transcription sustain Dpp target gene expression? One possibility is that
persistent low pMad levels are continuously required to repress Brk. Alternatively, Brk repression by
early Dpp signaling persists in the later stages without continuous Dpp signaling, for example, by
epigenetic regulation or via autoregulation. To distinguish between these possibilities, we applied
tubGal80ts to genetically remove *tkv* via *tkvHA*[FO] allele upon FLP expression from the entire A
compartment using *ci-Gal4* at different time points. To do so, the larvae were raised at 18 °C until a
temperature shift to 29 °C to induce Gal4 expression. Consistent with a role of *tkv* in wing pouch
growth, the earlier *tkv* was removed, the smaller the A compartment was (Extended Data Fig. 13). We
found that Brk is largely derepressed as early as 16 hr after *tkv* was removed (Extended Data Fig. 13).
By considering perdurance activity of Gal80ts for 6 hr after temperature shift[62], we can estimate that
Brk is derepressed within 10 hr at 29 °C after Dpp signaling was lost. Given that the transient *dpp*
transcription, which terminated in the early third instar stage, can sustain the anterior pMad signaling
and Brk repression at least 28 hr at 29 °C after Dpp protein from the main source is eliminated, these
results suggest that persistent weak pMad signaling is continuously required to repress Brk.
438

439   Taken together, Dpp dispersal-independent anterior patterning and growth can therefore be achieved
440   by a combination of a persistent weak signaling by transient *dpp* transcription outside the stripe and a
441   stronger signaling by continuous *dpp* transcription in the anterior stripe of cells.
442
Discussion
It has long been thought that Dpp dispersal from the anterior stripe of cells generates the morphogen
gradient in both compartments to control overall wing patterning and growth mainly based on the
complete loss of wing tissue in *dpp* disc mutant alleles (Fig. 9a). Here, we generated two novel protein
binder tools, namely HA trap and Dpp trap, to manipulate distinct parameters of the Dpp morphogen
to determine the requirement of Dpp dispersal and cell-autonomous signaling in the source cells. We
show that, although endogenous Dpp generates a gradient in both compartments, requirement of Dpp
dispersal is relatively minor and asymmetric along the A-P axis.
451
New protein binder tools manipulating distinct aspects of Dpp
Although nanobodies against GFP have been used most intensively in the field[63,64], fusion to GFP
could affect protein functions, as is the case for GFP-Dpp. To bypass this, we generated HA trap,
which is analogous to morphotrap and provides an alternative way to trap secreted proteins. Although
HA trap can trap HA-Dpp as efficient as morphotrap, we found several differences between the two.
First, while trapping GFP-Dpp by morphotrap in the source cells activates Dpp signaling in at least
one cell row in the P compartment[37], trapping HA-Dpp by HA trap did not (Fig. 3b). We think that the
difference is not because GFP-Dpp trapped by morphotrap activates Dpp signaling *in trans*, since
clonal accumulation of GFP-Dpp by morphotrap in the P compartment failed to do so[37]. Second, while
morphotrap could trap GFP-Dpp even in the peripheral regions[37], HA trap did not (Fig. 2l). Third,
while trapping GFP-Dpp by morphotrap in the source cells induced excessive Dpp signaling in the
source cells and caused severe defects in the adult wing[37], trapping HA-Dpp by HA trap in the source
cells slightly reduced Dpp signaling in the source cells and caused relatively minor defects in the adult
wing (Fig. 3, Extended Data Fig. 2a-j). We speculate that these differences are in part due to
overexpression of GFP-Dpp. Due to excess amount of GFP-Dpp, some GFP-Dpp may leak from the
morphotrap to activate Dpp signaling in the P compartment, or may reach to the peripheral regions.
Excessive Dpp signaling may cause cell death during pupal stages to cause severe adult wing defects
as previously shown[65]. These differences highlight the importance of investigating endogenous protein
functions.
471
In addition to HA trap, we isolated DARPins against Dpp, and generated a Dpp trap analogous to HA
trap. To our knowledge, this is the first DARPin isolated for an endogenous protein in Drosophila and
applied in the field of developmental biology. Interestingly, while HA trap blocks mainly Dpp
dispersal (Fig. 3), we found that Dpp trap blocks Dpp dispersal and cell-autonomous signaling in the
source cells (Fig. 5). We speculate that HA trap binds to the HA tag and thereby allows Dpp to bind
to its receptors, while Dpp trap directly binds to Dpp to block its interaction with the receptors.
Regardless of the actual mechanisms underlying this difference, these tools allowed us to dissect the
requirements of dispersal and cell-autonomous signaling in the source cells for wing pouch growth
and patterning. Relatively mild phenotypes by HA trap and severe phenotypes by Dpp trap indicate a
minor and asymmetric role of Dpp dispersal and a critical role of cell-autonomous Dpp signaling in
the source cells, respectively. Furthermore, these results also suggest that the severe *dpp* mutant
phenotypes do not reflect the role of Dpp dispersal alone, but reflect the role of both Dpp dispersal and
Dpp signaling in the source cells, with more contribution from the latter parameter (Fig. 9).
485
Asymmetric requirement of Dpp dispersal
The relatively minor requirement of Dpp dispersal for the anterior compartment is reminiscent of the
minor requirement of dispersal of Wg[12]. Wg dispersal is largely dispensable for the wing growth, likely

due to the early uniform *wg* transcription in the entire wing pouch and a memory of the earlier
signaling[12]. In contrast, the requirement of Dpp dispersal is asymmetric along the A-P axis due to the
early uniform *dpp* transcription in the A compartment. In both cases, transcriptional refinement of each
morphogen and persistent signaling by transient morphogen expression appear to be a key for
robustness against the absence of dispersal of each morphogen.
494
Although not identified, a *dpp* source outside the anterior stripe of cells has previously been implicated
to control the entire wing pouch growth based on the minor growth defects by removal of *dpp* from
the anterior stripe of cells using *dpp-Gal4* and severe growth defects by removal of *dpp* from the A
compartment using *ci-Gal4*[24]. However, we and others showed that removal of *dpp* using *dpp-Gal4*
was imprecise and inefficient, and that *dpp* derived from the anterior stripe of cells is indeed critical
for wing growth[25,41,61]. Thus, the presence of a *dpp* source outside the anterior stripe of cells has been
questioned. Our results suggest that such a *dpp* source indeed exists and contributes to the anterior
patterning and growth, but not to the growth of the entire wing pouch (Fig. 7, 8).
503
It remains unknown how transient *dpp* expression can maintain Dpp signaling in the anterior lateral
region (Fig. 8). One possibility is that feedback factors control the duration of Dpp signaling. Various
feedback factors have been shown to regulate Dpp signaling. For example, the Drosophila tumor
necrosis factor α homolog Eiger and a secreted BMP-binding protein Crossveinless-2 (Cv-2) are
positively regulated by Dpp signaling to either positively and negatively influence Dpp signaling in
the early embryo, respectively[66]. While JNK is not activated by Eiger in the wild type wing pouch[67],
Cv-2 acts as a positive feedback factor to specify the posterior crossvein during pupal stages[68]. Whether
these two factors act during wing pouch patterning and growth remains to be addressed. Pentagon is a
secreted feedback factor repressed by Dpp signaling to positively regulate Dpp signaling in the wing
disc[69]. Pent is produced in the lateral region of the wing pouch and regulates Dpp signaling and
proliferation there. Thus, in addition to its role for scaling, Pent may control the duration of Dpp
signaling in the lateral region. Another possibility is that, since it takes quite some time (20-24 hr at
29 °C) to eliminate Dpp protein upon excision of *dpp*[24,41], relatively stable *dpp* mRNA and/or protein
after the termination of transient *dpp* transcription may contribute to Brk repression (~28 hr at 29 °C).
518
Growth without Dpp dispersal and signaling
Despite a critical requirement of *dpp* for the entire wing pouch growth, our results uncover Dpp
dispersal- and signaling-independent lateral wing pouch growth (Fig. 3, 4, 6, Extended Data Fig. 5,
6)? The presence of Dpp signaling-independent growth appears inconsistent with the fact that the wing
pouch is completely lost in *dpp* mutant (Fig. 6a, c, Extended Data Fig. 7b). This could be in part due
to a failure of the initial specification of the wing pouch in *dpp* disc alleles[52]. Indeed, despite severe
growth defects, part of the wing pouch could still grow upon removal of *dpp* after wing pouch
specification (Fig. 4e-h) and upon removal of *tkv* after wing pouch specification (Extended Data Fig.
5).
528
Consistent with the presence of Dpp signaling-independent lateral growth, it has been shown that
lateral wing fates are less sensitive than medial wing fates in various *dpp* mutant alleles[70]. However,
this appears counter-intuitive since the lateral region, where morphogen level are low, is expected to
be more sensitive than the medial region to a reduction of morphogen levels. It has been proposed that
another BMP-type ligand, Glass bottom boat (Gbb), which is expressed ubiquitously in the wing pouch,
contributes to lateral cell fates[70]. However, since Gbb signaling is also mediated by Tkv[71] but the lateral
growth is independent of Tkv (Fig. 3, 4, 6, Extended Data Fig. 5, 6), we think that the lateral region
develops independent of direct Dpp and Gbb signaling. Thus, the lateral region is less sensitive than
the medial region in various *dpp* mutant alleles, probably because the lateral region can develop
independent of Dpp signaling. What regulates the Dpp signaling-independent lateral wing pouch

growth? Given that *5xQE.DsRed* is also dependent on Wg[45,54], Wg may regulate *5xQE.DsRed*
expression and growth in the absence of Dpp dispersal.
541
In addition to the Dpp signaling-independent lateral growth, requirement of anterior Dpp signaling for
posterior growth (Fig. 5j, Extended Data Fig. S2u) and rescue of posterior growth in *dpp* mutants by
anterior Dpp signaling (Fig. 6) indicates that anterior Dpp signaling is non-autonomously involved in
posterior lateral growth. We note that similar rescue of posterior growth in *dpp* mutant by anterior Dpp
signaling has previously been recognized, but the rescued posterior growth was interpreted as growth
of the hinge region, without immunostainings for relevant markers[59]. It remains unknown how anterior
Dpp signaling contributes to posterior growth. One possibility is that factors from the A compartment
may act non-autonomously to promote posterior growth. Such factors include, but are not limited to,
direct downstream factors of Dpp signaling. Given that *5xQE.DsRed* is dependent on Wg[45,54], Wg
derived from the rescued A compartment may regulate *5xQE.DsRed* expression and growth in the P
compartment. Alternatively, mechanical forces may be involved in the non-autonomous growth. It has
been proposed that growth factors such as Dpp induce medial growth and subsequently stretch the
peripheral regions to induce lateral growth. As the wing disc grows, the peripheral regions in turn
compresses the medial region of the wing disc to inhibit the growth of the medial region[72-76]. Thus, the
growth of the A compartment may stretch the P compartment cells to stimulate their proliferation. It
has also been shown that juxtaposition of cells with different Dpp signaling level can induce
proliferation non-autonomously but the growth is transient[26,27]. Therefore, we think it unlikely that the
difference of Dpp signaling levels between two compartments can induce sustained growth.
560
A refined growth model
The presence of Dpp signaling-independent lateral wing pouch growth is at odds with all the growth
models assuming that Dpp dispersal directly controls overall wing patterning and growth based on the
complete loss of wing tissue in *dpp* mutants[24-33] (Fig. 9a). For example, no wing pouch growth is
expected without Dpp signaling due to a lack of either a temporal increase of Dpp signaling (temporal
model)[32], a detectable Dpp signal (threshold model)[25,61,77], or a slope of Dpp signaling activity
(gradient model)[26,27] (Fig. 9a).
568
It has recently been proposed that Wg and Dpp control wing pouch size by two distinct
mechanisms[33,45,54,78]. One is an intracellular mechanism, in which Vg controls its own expression
through QE in response to Wg and Dpp; the other one is an intercellular mechanism, in which Vg-
positive wing pouch cells send a feed-forward signal to induce QE-dependent *vg* expression in the
neighboring pre-wing cells to recruit them into the wing pouch in response to Dpp and Wg. The
dispersal of each morphogen is critical for the two mechanisms in the model[33,45,54,78]. In the genetic
setup used, in which morphogens and *vg* expression are eliminated, the two mechanisms appear to
recapitulate the dynamics of *vg* expression seen during normal wing disc development, in which *vg*
expression is gradually expanded in the wing pouch area. However, although sufficient, these
mechanisms do not necessarily account for *vg* expression under physiological conditions. Indeed, we
found Dpp dispersal and/or signaling-independent *vg* expression in various conditions.
580
Among the models, the presence of Dpp signaling-independent lateral wing pouch growth appears
most consistent with the growth equalization model, in which Dpp signaling removes Brk to allow
medial regions to grow, while Brk suppresses Dpp signaling-independent lateral growth with higher
proliferation nature to equalize the non-uniform growth[22,28]. However, the identity of medial and lateral
regions remained undefined in this model. Given the complete lack of wing pouch in *dpp* mutants, it
is tempting to speculate that the Dpp signaling-dependent medial region corresponds to the entire wing
pouch region, and the Dpp signaling-independent lateral region corresponds to the hinge region located
next to the wing pouch region (Fig. 9a). Indeed, using morphotrap, such a Dpp signaling-insensitive

posterior growth has previously been interpreted as the growth of the hinge region due to severe adult
wing defects[37]. However, in contradiction to this interpretation, which would predict overgrowth of
the hinge region in *brk* mutant, a massive overgrowth of wing pouch region was observed in *brk* mutant
(Extended Data Fig. 7). Based on our results that the lateral wing pouch regions can grow independent
of direct Dpp signaling (Fig. 3, 4, 6, Extended Data Fig. 5, 6), we therefore suggest to refine the growth
equalization model and propose that both Dpp signaling-dependent medial and Dpp signaling-
independent lateral regions are located within the wing pouch (Fig. 9b). The permissive role of Dpp
signaling in modulating a non-uniform growth potential within the wing pouch raises questions about
what kind of instructive signals control proliferation and growth, and how the non-uniform growth
potential emerges within the wing pouch independent of the Dpp/Brk system[44].
599

Conclusion
Our approach applying customized protein binder tools to manipulate distinct parameters of Dpp
challenges the long-standing dogma that Dpp dispersal controls overall wing patterning and growth.
Given that the tools developed in this study are easily applicable in other tissues, it would be interesting
to investigate the precise requirement of Dpp dispersal and signaling in other tissues or between
different organs.
606

 Experimental procedures
Data reporting

No statistical methods were used to predetermine sample size. The experiments were not randomized,
and investigators were not blinded to allocation during experiments and outcome assessment.

Fly stocks
Flies were kept in standard fly vials (containing polenta and yeast) in a 25°C incubator. The following
fly lines were used: *dpp^FO*, *dpp-Gal4*, *UAS-FLP* (Matthew Gibson), *ptc-Gal4* (BL2017),
P{act5C(FRT.polyA)lacZ.nls1}3, ry506 (BL6355), *w[*]; P{w[+mC]=UAS-RedStinger}6,*
P{w[+mC]=UAS-FLP.Exel}3, P{w[+mC]=Ubi-p63E(FRT.STOP)Stinger}15F2 (G-
TRACE)(BL28281), *brk^XA* (BL58792), *dpp^MI03752* (BL36399), *PBac{RB}e00178, Dp(2;2)DTD48*
(Bloomington stock center). *omb*-LacZ (Kyoto101157). *act>Stop, y+>LexA^LHG, tkv^a12, UAS-TkvQD,*
pLexAop-TkvQD (Konrad Basler), *5xQE.DsRed* (Gary Struhl), *UAS/LexAop-HAtrap* (this study),
UAS/LexAop-Dpp trap (F1) (this study), *dpp^d8, dpp^d12, nub-Gal4* (II), *ci-Gal4* (II), *hh-Gal4* (III), *UAS-*
p35(III), *tub-Gal80ts* (III) are described from Flybase. *tub>CD2, Stop>Gal4, UAS-nlacZ* (Francesca
Pignoni). *TkvHA* (Giorgos Pyrowolakis).

622
Genotypes by figures
Fig. 1c-e: *yw; HA-dpp/HA-dpp*
Fig. 2b: *yw; ptc-Gal4, Ollas-HA-dpp/+*
Fig. 2d: *yw; ptc-Gal4, Ollas-HA-dpp/+; UAS/LexAop-HAtrap/+*
Fig. 2f, h: *hsFLP; Ollas-HA-dpp/tub>CD2, Stop>Gal4, UAS-nlacZ; UAS/LexAop-HAtrap/+*
Fig. 2j: *hsFLP; ptc-Gal4, Ollas-HA-dpp/tub>CD2, Stop>Gal4, UAS-nlacZ; UAS/LexAop-HAtrap/+*
Fig. 2l: *hsFLP; Ollas-HA-dpp/tub>CD2, Stop>Gal4, UAS-nlacZ; UAS/LexAop-HAtrap/+*
Fig. 3a, c: *(5xQE.DsRed); ptc-Gal4, HA-dpp/HA-dpp*
Fig. 3b, d: *(5xQE.DsRed); ptc-Gal4, HA-dpp/HA-dpp; UAS/LexAop-HAtrap/+*
Fig. 3k, m: *yw; nub-Gal4, HA-dpp/HA-dpp*
Fig. 3l, n: *yw; nub-Gal4, HA-dpp/HA-dpp; UAS/LexAop-HAtrap/+*
Fig. 4a: *hsFLP/5xQE.DsRed; HA-dpp, tkv^a12 FRT40/HA-Dpp, UbiGFP, FRT40, ptc-Gal4;*
UAS/LexAop-HAtrap/+
Fig. 4b: *hsFLP/5xQE.DsRed; tkv^a12 FRT40/UbiGFP, FRT40*
Fig. 4c, d: *hsFLP/5xQE.DsRed; tkvHA^FO/tkvHA^FO*
Fig. 4e, f: (internal control within a cross) *5xQE.DsRed/+; (dpp^FO, ci-Gal4)/(dpp^FO); (UAS-*
FLP)/tubGal80ts,
Fig. 4g, h: *5xQE.DsRed/+; dpp^FO, ci-Gal4/dpp^FO; UAS-FLP/tubGal80ts*
Fig. 5b: (left) *yw; ptc-Gal4, Ollas-HA-dpp/+*, (right) *yw; ptc-Gal4, Ollas-HA-dpp/+; UAS/LexAop-*
Dpptrap/+
Fig. 5c: *yw; ptc-Gal4, HA-dpp/+*
Fig. 5d: *yw; ptc-Gal4, HA-dpp/+; UAS/LexAop-Dpptrap/+*
Fig. 5k, m: *yw; nub-Gal4, HA-dpp/+*
Fig. 5l, n: *yw; nub-Gal4, HA-dpp/+; UAS/LexAop-Dpptrap/+*
Fig. 6a, c: *(y)w; (5xQE.DsRed); dpp^d8/dpp^d12*
Fig. 6b, d: *(y)w; (5xQE.DsRed); dpp^d8/dpp^d12; dpp-Gal4/UAS-tkvQD*
Fig. 6f: *(5xQE.DsRed); dpp^d8, UAS-FLP/dpp^d12, act>Stop, y+>LexA^LHG; dpp-Gal4/LexAop-tkvQD*
Fig. 7c: *yw; dpp-T2A-Gal4, Dp(2;2)DTD48(dpp+)/+; P{w[+mC]=UAS-RedStinger}6,*
P{w[+mC]=UAS-FLP.Exel}3, P{w[+mC]=Ubi-p63E(FRT.STOP)Stinger}15F2/+
Fig. 7e-h: *yw M{vas-int.Dm}zh-2A; dpp-T2A-d2GFP-NLS/Cyo, P23*
Fig. 7i-l: *yw*
Fig. 8a-c: *ptc-Gal4, dpp^FO/+; tubGal80ts/UAS-FLP, act5C(FRT.polyA)lacZ.nls*

Fig. 8d-h: *ptc-Gal4, dpp^{FO}/dpp^{FO}; tubGal80ts/UAS-FLP, act5C(FRT.polyA)lacZ.nls*
Fig. 8i-j: *ci-Gal4, dpp^{FO}/dpp^{FO}; tubGal80ts/UAS-FLP, act5C(FRT.polyA)lacZ.nls*
657
Extended Data Fig. 1a, e: *ptc-Gal4/+*
Extended Data Fig. 1b, f: *ptc-Gal4/+; UAS/LexAop-HAtrap/+*
Extended Data Fig. 2a, c, e, g, i: *HA-dpp/HA-dpp, ci>+* (left) and *HA-dpp/HA-dpp, ci>HAtrap* (right)
Extended Data Fig. 2k, m, o, q, s: *HA-dpp/+, ci>+* (left) and *HA-dpp/+, ci>Dpptrap* (right)
Extended Data Fig. 3a: *nub-Gal4, ptc-Gal4, HAdpp/HA-dpp* (control),
Extended Data Fig. 3b: *nub-Gal4, ptc-Gal4, HA-dpp/HA-dpp; UAS/LexAop-HAtrap/+*
Extended Data Fig. 4a: *ptc-Gal4, HA-dpp/HA-dpp*
Extended Data Fig. 4b: *ptc-Gal4, HA-dpp/HA-dpp; UAS/LexAop-HAtrap/+*
Extended Data Fig. 4c: *nub-Gal4, HA-dpp/+; UAS/LexAop-Dpptrap/+,*
Extended Data Fig. 4e: *nub-Gal4, HA-dpp/HA-dpp; UAS/LexAop-HAtrap/+,*
Extended Data Fig. 4f: *nub-Gal4, HA-dpp/HA-dpp; UAS/LexAop-HAtrap/UAS-p35*
Extended Data Fig. 4h: *nub-Gal4, HA-dpp/+; UAS/LexAop-Dpptrap/+*
Extended Data Fig. 4i: *nub-Gal4, HA-dpp/+; UAS/LexAop-Dpptrap/UAS-p35*
Extended Data Fig. 5a, b: (control within the cross) *5xQE.DsRed/+; (tkvHA^{FO}, ci-Gal4)/(tkvHA^{FO});*
(UAS-FLP)/tubGal80ts
Extended Data Fig. 5c, d: *5xQE.DsRed/+; tkvHA^{FO}, ci-Gal4/tkvHA^{FO}; UAS-FLP/tubGal80ts*
Extended Data Fig. 5e, f: (control within the cross) *5xQE.DsRed/+; (tkvHA^{FO})/(tkvHA^{FO}); +/Hh-Gal4,*
tubGal80ts
Extended Data Fig. 5g, h: *5xQE.DsRed/+; tkvHA^{FO}/tkvHA^{FO}; UAS-FLP/Hh-Gal4, tubGal80ts*
Extended Data Fig. 6a: (control within the cross) *5xQE.DsRed/+; (tkvHA^{FO})/tkvHA^{FO}; (Hh-Gal4)/+*
Extended Data Fig. 6b: (experiment) *5xQE.DsRed/+; tkvHA^{FO}/tkvHA^{FO}; Hh-Gal4/UAS-FLP*
Extended Data Fig. 7a: *5xQE.DsRed/+, dpp^{d8} or dpp^{d12}/+*
Extended Data Fig. 7b: *5xQE.DsRed/+; dpp^{d8}/dpp^{d12}*
Extended Data Fig. 7c: *5xQE.DsRed, brk^{XA}/Y, dpp^{d8} or dpp^{d12}/+*
Extended Data Fig. 7d: *5xQE.DsRed, brk^{XA}/Y, dpp^{d8}/dpp^{d12}*
Extended Data Fig. 8a: *Ollas-HA-dpp, ptc-Gal4/+*
Extended Data Fig. 8b: *Ollas-HA-dpp, ptc-Gal4/+; UAS/LexAop-C9/+*
Extended Data Fig. 8d: *ptc-Gal4/+*
Extended Data Fig. 8e: *ptc-Gal4/+; UAS/LexAop-C9/+*
Extended Data Fig. 9a: *yw; ptc-Gal4, HA-dpp/+* (identical disc as Fig. 5c)
Extended Data Fig. 9b: *yw; ptc-Gal4, HA-dpp/+; UAS/LexAop-Dpptrap/+* (identical disc as Fig. 5d)
Extended Data Fig. 9c, d: *dpp^{FO}/+; UAS-FLP/tubGal80ts*
Extended Data Fig. 9e: *dpp^{FO}, ci-Gal4/dpp^{FO}; UAS-FLP/tubGal80ts*
Extended Data Fig. 10a: (control) *ptc-Gal4, HA-dpp/+,*
Extended Data Fig. 10b: (Dpptrap) *ptc-Gal4, HA-dpp/+; UAS/LexAop-Dpptrap/+*
Extended Data Fig. 10c: (HAtrap) *ptc-Gal4, HA-dpp/HA-dpp; UAS/LexAop-HAtrap/+*
Extended Data Fig. 11: *yw M{vas-int.Dm}zh-2A; dpp-T2A-d2GFP-NLS/Cyo, P23*
Extended Data Fig. 12: (control) *ptc-Gal4, HA-dpp/HA-dpp; tubGal80ts/+*, (experiment) *ptc-Gal4,*
HA-dpp/HA-dpp; UAS/LexAop-HAtrap/tubGal80ts
Extended Data Fig. 13: (control) *tkvHA^{FO}/+; UAS-FLP/tubGal80ts*
Extended Data Fig. 13: (experiment) *tkvHA^{FO}, ci-Gal4/tkvHA^{FO}; UAS-FLP/tubGal80ts*
699

Immunostainings and antibodies
Protocol as described previously[37]. For extracellular staining, dissected larvae were incubated with
primary antibodies in M3 medium for 1 hr on ice before fixation to allow antibodies to access only the
extracellular antigens. Each fly cross was set up together with a proper control and genotypes were

processed in parallel. If the genotype could be distinguished, experimental and control samples were
processed in the same tube. To minimize variations, embryos were staged by collecting eggs for 2-
4hrs. An average intensity image from 3 sequential images from a representative wing disc is shown
for all the experiments. The following primary antibodies were used; anti-HA (3F10, Roche; 1:300 for
conventional staining, 1:20 for extracellular staining), anti-Ollas (Novus Biologicals, 1:300 for
conventional staining, 1:20 for extracellular staining), anti-phospho-Smad1/5 (1:200; Cell Signaling),
anti-Brk (1:1.000; Gines Morata), anti-Sal (1:500; Rosa Barrio), anti-Omb (1:500; Gert Pflugfelder),
anti-Wg (1:120; DSHB, University of Iowa), anti-Ptc (1:40; DSHB, University of Iowa), anti-β-
Galactosidase (1:1.000; Promega, 1:1000; abcam), anti-Cleaved Caspase-3 (1:500; Cell Signaling).
All the primary and secondary antibodies were diluted in 5% normal goat serum (NGS) (Sigma) in
PBT (0.03% Triton X-100/PBS). All secondary antibodies from the AlexaFluor series were used at
1:500 dilutions. Wing discs were mounted in Vectashield (H-1000, Vector Laboratories). Images of
wing discs were obtained using a Leica TCS SP5 confocal microscope (section thickness 1 μm).
717
RNAscope
smFISH using RNAscope technology with the probe against *dpp* (Cat No. 896761) has previously
been successful to visualize *dpp* mRNA in the germline stem cell niche[79]. The probes against *dpp*
target 682-1673 of MM_057963.5 (accession number from NCBI). RNAscope was performed in an
Eppendorf tube. Larvae were dissected in PBS and fixed in 4% paraformaldehyde in PBS for 30 min.
Fixed larvae were washed with PBS and then with PBT (PBS containing 0.03% Triton X-100). Fixed
larvae were dehydrated at RT (room temperature) in a series of 25%, 50%, 75% and 100% methanol
in PBT, and stored overnight in 100% methanol at −20 °C. Larvae were rehydrated at RT in a series
of 75%, 50%, 25%, 0% methanol in PBT. Larvae were treated with protease using Pretreat 3
(RNAscope H2O2 & Protease Plus Reagents; ACD, 322330) at RT for 5 min. After washing with PBT,
hybridization using probes against *dpp* was performed overnight at 40 °C. The following day, larvae
were washed with in RNAscope wash buffer and re-fixed with 4% paraformaldehyde in PBS for 10
min. After washing with PBS, fluorescent signal was developed using RNAscope Fluorescent
Multiplex Reagent Kit according to the manufacturer's instructions. Wing discs were mounted in
Vectashield (H-1000, Vector Laboratories) and dissected for imaging. Images of wing discs were
obtained using a Leica TCS SP5 confocal microscope (section thickness 1 μm).
734
Quantification
Quantification of pMad, Brk, Sal, and Omb
From each z-stack image, signal intensity profile along A/P axis was extracted from average projection
of 3 sequential images using ImageJ. Each signal intensity profile was aligned along A/P compartment
boundary (based on anti-Ptc staining) and average signal intensity profile from different samples was
generated and plotted by the script (wing_disc-alignment.py). The average intensity profile from
control and experimental samples were then compared by the script (wingdisc_comparison.py). Both
scripts can be obtained from (https://etiennees.github.io/Wing_disc-alignment/). The resulting signal
intensity profiles (mean with SD) were generated by Prism.
Quantification of wing pouch size and adult wing size
The A and P compartment of the wing pouches were approximated by Ptc/Wg staining and positions
of folds, and the A/P compartment boundary of the adult wings were approximated by L4 position.
The size of each compartment was measured using ImageJ. Scatter dot plots (mean with SD) were
generated by Prism.
Statistics
Statistical significance was assessed by Prism based on the normality tests using a two-sided Mann-
Whitney test (Fig. 3t, for P compartment, Fig. 5s, for A compartment, Fig. 5u, for A compartment) and
a two-sided Student's *t*-test with unequal variance for the rest of the comparisons (***p<0.0002 ****
p<0.0001).

Generation of *HA-dpp* and *GFP-dpp* knock-in allele

Cloning of plasmids for injection.

A fragment containing multi-cloning sites (MCS) between two inverted attB sites was synthesized and inserted in the pBS (BamHI) vector (from Mario Metzler). A genomic fragment of *dpp* between *dpp^{MI03752}* and *PBac{RB}e00178* (about 4.4kb), as well as an *FRT* and *3xP3mCherry* were inserted in this MCS by standard cloning procedures. A fragment encoding HA tag or GFP was inserted between the XhoI and NheI sites inserted after the last Furin processing site[18].

Inserting *dpp* genomic fragments in the *dpp* locus

The resulting plasmids were injected in *yw M{vas-int.Dm}zh-2A; dpp^{MI03752}/Cyo, P23*. *P23* is a transgene containing a *dpp* genomic fragment to rescue *dpp* haploinsufficiency. After the hatched flies were backcrossed, flies that lost *y* inserted between inverted attP sites in the mimic transposon lines were individually backcrossed to establish stocks. The orientation of inserted fragments was determined by PCR.

Removal of the endogenous *dpp* exon by FLP/FRT recombination

Males from the above stock were crossed with females of genotype *hsFLP; al, PBac{RB}e00178 /SM6, al, sp* and subjected to heat-shock at 37°C for 1hr/day. *PBac{RB}e00178* contains *FRT* sequence and *w+* and upon recombination, the *dppHA* genomic fragment are followed by FRT and w+. Hatched males of *hsFLP;dppHA/al,PBac{RB}e00178* were crossed with *yw; al, b, c, sp/ SM6, al, sp*. From this cross, flies *yw; dppHA(w+)/SM6, al, sp* were individually crossed with *yw; al, b, c, sp/ SM6, al, sp* to establish the stock.

Construction of α-HAscFv

cDNA of HAscFv was constructed by combining coding sequences of variable regions of the heavy chain (V_H: 1-423 of LC522514) and of the light chain (V_L: 67-420 of LC522515) cloned from anti-HA hybridoma (clone 12CA5)[40] with a linker sequence (5'-accggtGGCGGAGGCTCTGGCGGAGGAGGTTCCGGCGGAGGTGGAAGCgatatc-3') in the order of V_H-linker-V_L. The coding sequence of HAscFv was cloned into pCS2+mcs-2FT-T for FLAG-tagging. Requests for HAscFv should be addressed to YM (mii@nibb.ac.jp). To generate HA trap, the region encoding morphotrap (VHH-GFP4) was replaced with KpnI and SphI sites in pLOTattB-VHH-GFP4:CD8-mChery[37]. A fragment encoding HAscFv was amplified by PCR and then inserted via KpnI and SphI sites by standard cloning procedures.

Selection of Dpp-binding DARPins and generation of Dpp trap

Streptavidin-binding peptide (SBP)-tagged mature C-terminal domain of Dpp was cloned into pRSFDuet vector by a standard cloning. Dpp was overexpressed in *E. coli*, extracted from inclusion bodies, refolded, and purified by heparin affinity chromatography followed by reverse phase HPLC[80]. To isolate suitable DARPins, SBP-tagged Dpp was immobilized on streptavidin magnetic beads and used as a target for DARPin selections by employing multiple rounds of Ribosome Display[81,82]. Due to the aggregation and precipitation propensity of the purified SBP-Dpp, the refolded dimers previously stored in 6 M urea buffer (6 M urea, 50 mM Tris-HCl, 2 mM EDTA pH8.0, 0.25 M NaCl) were diluted to a concentration of 100-120 µg/ml in the same buffer and subsequently dialyzed against 4 mM HCl at 4°C overnight. To ensure binding of correctly folded Dpp to the beads, this solution was diluted five times in the used selection buffer just prior to bead loading and the start of the ribosome display selection. In each panning round, the target concentration presented on magnetic beads was reduced, while the washing stringency was simultaneously increased to enrich for binders with high affinities[81]. In addition, from the second round onward, a pre-panning against Streptavidin beads was performed prior to the real selection to reduce the amounts of matrix binders. After four rounds of selection, the enriched pool was cloned into an *E. coli* expression vector, enabling the production of

both N-terminally His$_8$- and C-terminally FLAG-tagged DARPins. Nearly 400 colonies of transformed
E. coli were picked and the encoded binders expressed in small scale. Bacterial crude extracts were
subsequently used in enzyme-linked immunosorbent assay (ELISA) screenings, detecting the binding
of candidate DARPins to streptavidin-immobilized Dpp, or just streptavidin (indicating background
binding) by using a FLAG-tag based detection system (data not shown). Of those 127 candidate
DARPins interacting with streptavidin-immobilized Dpp, 73 (or 57%) specifically bound to Dpp (i.e.,
having at least threefold higher signal for streptavidin-immobilized Dpp than to streptavidin alone). 36
of these (50%) revealed unique and full-length sequences. To generate Dpp trap, the region encoding
morphotrap (VHH-GFP4) was replaced with KpnI and SphI sites in pLOTattB-VHH-GFP4:CD8-
mChery[37]. Each fragment encoding a DARPin was amplified by PCR and then inserted via KpnI and
SphI sites by standard cloning procedures.
814

815   **Generation of *tkvHA^{FO}* (Flip-out) allele**
816   The *tkvHA* allele was previously described[83]. An FRT cassette was inserted in the re-insertion vector
817   for *tkvHA* (Genewiz) and re-inserted into the attP site in the *tkv* locus.
818

819   **Generation of endogenous *dpp-Gal4***
820   *pBS-KS-attB2-SA(1)-T2A-Gal4-Hsp70* (addgene 62897) was injected in the *yw M{vas-int.Dm}zh-2A;*
821   *dpp^{MI03752}/Cyo, P23* stock. Since the *Gal4* insertion causes haploinsufficiency, the *dpp-Gal4* was
822   recombined with *Dp(2;2)DTD48* (duplication of *dpp*) for G-TRACE analysis.
823

824   **Generation of an endogenous *dpp* reporter**
825   A DNA fragment containing T2A-d2GFP-NLS was synthesized and used to replace the region
826   containing *T2A-Gal4* in *pBS-KS-attB2-SA(1)-T2A-Gal4-Hsp70* via BamHI to generate *pBS-KS-attB2-*
827   *SA(1)-T2A-d2GFP-NLS-Hsp70* (Genewiz). The resulting plasmid was injected in the *yw M{vas-*
828   *int.Dm}zh-2A; dpp^{MI03752}/ Cyo, P23* stock.
829
830

Acknowledgments:
We thank Konrad Basler, Giorgos Pyrowolakis, Gustavo Aguilar and Sheida Hadji Rasouliha for
comments on the manuscript. Stocks obtained from the Bloomington Drosophila Stock Center (NIH
P40OD018537) were used in this study. We thank Konrad Basler, Gary Struhl, Matthew Gibson,
Giorgos Pyrowolakis, and Kyoto stock center for flies. We thank Gines Morata, Rosa Barrio, Gert
Pflugfelder and the Developmental Studies Hybridoma Bank at The University of Iowa for antibodies.
We thank the Biozentrum Imaging Core Facility for maintenance of microscopes and support. We
thank Etienne Schmelzer for scripts for quantification and Oguz Kanca for the idea to manipulate the
dpp locus. We thank Dietmar Schreiner and Caroline Bornmann for introducing RNAscope and
sharing reagents, and Minkyoung Lee for the advice for smFISH and sharing reagents. We would like
to thank Bernadette Bruno, Gina Evora and Karin Mauro for constant and reliable supply with world's
best fly food. We further acknowledge all current and former members of the High-Throughput Binder
Selection facility at the Department of Biochemistry of the University of Zurich for their contribution
to the establishment of the semi-automated ribosome display that resulted in the generation of the used
anti-Dpp DARPin binders, particularly Thomas Reinberg. **Funding:** SM has been supported by a JSPS
Postdoctoral Fellowship for Research Abroad and the Research Fund Junior Researchers University
of Basel, and is currently supported by a SNSF Ambizione grant (PZ00P3_180019). YM has been
supported by JST PRESTO (JPMJPR194B) and JSPS KAKENHI (18K14720). The work in the
Affolter laboratory was supported by grants from Kantons Basel-Stadt and Basel-Land and from the
SNSF (MA).
851
852 **Author contributions:** This project was conceived by SM and MA. SM designed, performed, and
853 analyzed all the experiments except isolation of α-HA scFv and DARPins. JVS and AP isolated
854 DARPins against Dpp. YM, YH, and MT cloned α-HA scFv. DB helped with Dpp purification. The
855 main text was written by SM and MA with comments from all the authors.
856
857 **Competing interests:** Authors declare no competing interests.
858

859
860
861
862 **Figure 1. Generation of HA trap**
863 **a**, A schematic view of the wing disc and the adult wing. **b**, A schematic view of a platform
864 manipulating endogenous *dpp* locus. **c**, Adult wing of a homozygous *HA-dpp* fly. **d**, Conventional α-
865 HA staining of *HA-dpp* homozygous wing disc. **e**, Extracellular α-HA staining (Ex HA) and α-pMad
866 staining of *HA-dpp* homozygous wing disc. Quantification of each staining.
867

868
869
870
Figure 2. Characterization of HA trap
a, A schematic view of HA trap. **b-e**, Extracellular α-Ollas staining (Ex Ollas), HA trap (mCherry), and merge of control
Ollas-HA-dpp/+ disc (**b**), and of *Ollas-HA-dpp/+ ptc>HA trap* disc (**d**). **c, e,** Quantification of extracellular α-Ollas
staining of (**b**) and (**d**), respectively. **f-k**, Extracellular α-Ollas staining, HA trap (mCherry), and merge of *Ollas-HA-dpp/+*
disc with an anterior clone of cells expressing HA trap (**f**), of *Ollas-HA-dpp/+* disc with a posterior clone of cells expressing
HA trap (**h**), and of *Ollas-HA-dpp/+* disc with HA trap expression using *ptc-Gal4* and clones of cells expressing HA trap
in both compartments (**j**). **g, i, k,** Quantification of extracellular α-Ollas staining and HA trap (mCherry) of (**f**), (**h**), (**j**),
respectively. Arrows indicate clones of cells expressing HA trap where quantification was performed. **l**, Extracellular α-
Ollas staining, HA trap (mCherry), pMad and merge of *Ollas-HA-dpp/+* wing disc with clones of cells expressing HA trap.
Arrows indicate clones of cells expressing HA trap where pMad signal is reduced upon trapping Ollas-HA-Dpp. Arrow
heads indicate a clone of cells expressing HA trap that accumulates Ollas-HA-Dpp near the source cells and a clone of
cells expressing HA trap that does not accumulate Ollas-HA-Dpp far from the source cells. Dashed white lines mark the
A-P compartment border. Scale bar 50 μm.

884
885
Figure 3. Blocking Dpp dispersal by HA trap causes minor and asymmetric patterning and
growth defects
a-b, α-pMad, α-Brk, α-Sal, α-Omb, 5xQE.DsRed, DSRF, and Dpp trap (mCherry) (inset) of control
wing disc (**a**) and wing disc expressing HA trap using *ptc-Gal4* (**b**). **c-d**, Control adult wing (**c**), and
adult wing expressing HA trap using *ptc-Gal4* (**d**). **e-h**, Quantification of α-pMad (**e**), α-Brk (**f**), α-Sal
(**g**), α-Omb (**h**) staining in (**a-b**). **i-j**, Quantification of compartment size of wing pouch (**i**) and adult
wing (**j**) upon HA trap expression using *ptc-Gal4*. **k-l**, α-pMad, α-Brk, α-Sal, α-Omb, 5xQE.DsRed,
DSRF, and Dpp trap (mCherry) (inset) of control wing disc (**k**) and wing disc expressing HA trap
using *nub-Gal4* (**l**). **m-n**, Control adult wing (**m**) and adult wing expressing HA trap using *nub-Gal4*
(**n**). **o-r**, Quantification of α-pMad (**o**), α-Brk (**p**), α-Sal (**q**), α-Omb (**r**) staining in (**k-l**). **s-t**,
Quantification of compartment size of wing pouch (**s**) and adult wing (**t**) upon HA trap expression
using *nub-Gal4*. Dashed white lines mark the A-P compartment border. Scale bar 50 μm.
898

Figure 4. Lateral wing pouch growth without Dpp signaling

a-b, *tkv^{a12}* clones (indicated by the absence of GFP signal) induced in *HA-dpp/HA-dpp, ptc>HA* trap wing discs (**a**) and in wild type wing discs (**b**). **c-d**, *tkvHA^{FO}* clones (indicated by the absence of α-HA staining) in wild type wing discs. Clones were induced at 60-72 hr AEL (after egg laying) during mid-second to early third instar stages. **e-h**, α-pMad and 5xQE.DsRed (**e**, **g**) and α-Brk and 5xQE.DsRed (**f**, **h**) of control wing disc (**e**, **f**) and wing disc removing *dpp* using *ci-Gal4* with *tubGal80s* (**g**, **h**). Crosses were shifted from 18°C to 29°C at 4day AEL (early second instar). Scale bar 50 μm.

909
910
Figure 5. Blocking Dpp dispersal and signaling by Dpp trap causes severe defects similar to *dpp* mutant
a, a schematic view of Dpp trap based on DARPins [2] against Dpp. **b**, α-ExOllas staining and Dpp trap expression
(mCherry) (inset) of control wing disc (left) and wing disc expressing Dpp trap using *ptc-Gal4* (right). Quantification
of ExOllas signal. **c-d**, α-pMad, α-Brk, α-Sal, α-Omb staining and Dpp trap (mCherry) expression (inset) of control
wing disc (**c**) and wing disc expressing Dpp trap using *ptc-Gal4* (**d**). **e-h**, Quantification of α-pMad (**e**), α-Brk (**f**), α-
Sal (**g**), α-Omb (**h**) staining in (**c-d**). **i**, Quantification of compartment size of wing pouch upon Dpp trap expression
using *ptc-Gal4*. **j**, Comparison of wing pouch compartment size upon HA trap and Dpp trap expression using *ptc-*
Gal4 (comparison of Fig. 3i and Fig. 5i). **k-l**, α-pMad, α-Brk, α-Sal, α-Omb staining and Dpp trap (mCherry)
expression (inset) of control wing disc (**k**) and wing disc expressing Dpp trap using *nub-Gal4* (**l**). **m-n**, Control adult
wing (**m**), and adult wing expressing Dpp trap using *nub-Gal4* (**n**). **o-r**, Quantification of α-pMad (**o**), α-Brk (**p**), α-
Sal (**q**), α-Omb (**r**) staining in (**k-l**). **s-t**, Quantification of compartment size of wing pouch (**s**) and adult wing (**t**)
upon Dpp trap expression using *nub-Gal4*. **u**, Comparison of compartment size of wing pouch upon HA trap and
Dpp trap expression using *nub-Gal4* (comparison of Fig. 3s and Fig. 5s). Scale bar 50 μm.

924
925
Figure. 6 Rescue of *dpp* mutant by cell-autonomous Dpp signaling
a-b, α-pMad, α-Brk, α-Sal, α-Omb, 5xQE.DsRed, and DSRF staining of (**a**) *dpp^d8^/dpp^d12^* wing disc
and (**b**) *dpp^d8^/dpp^d12^, dpp>tkvQD* wing disc. Arrows indicate rescued posterior wing pouch identified
by 5xQE.DsRed and DSRF staining. **c**, Adult wing of *dpp^d8^/dpp^d12^*. **d**, Adult wing of *dpp^d8^/dpp^d12^,*
dpp>tkvQD. **e**, A schematic view of converting *dpp-Gal4* into a *LexA* driver, which is permanently
expressed in lineage of *dpp-Gal4* expressing cells. In this experimental setup, the lineage of *dpp-Gal4*
(including lineage of non-specific *dpp-Gal4* expression) will permanently activate TkvQD and thus
pMad signaling. **f**, α-pMad, α-Brk, α-Sal, α-Omb staining, and *5xQE.DsRed* signal of *dpp^d8^/dpp^d12^,*
dpp-Gal4>UAS-FLP, act>y+>LexA-LHG, LOP-tkvQD wing disc. Arrow indicates rescued posterior
wing pouch identified by 5xQE.DsRed staining. Scale bar 50 μm.
936

937
938
Figure 7. Initial uniform *dpp* transcription in the anterior compartment
a-c, Lineage of endogenous *dpp-Gal4* during wing development. (**a**) a schematic view of a *Gal4*
insertion into the *dpp* locus. (**b**) A schematic view of G-TRACE analysis. While RFP expression labels
the real-time *Gal4*-expressing cells, GFP expression labels the lineage of the *Gal4*-expressing cells (**c**),
G-TRACE analysis of the endogenous *dpp-Gal4*. Scale bar 50 μm. **d**, A schematic view of *d2GFP*
insertion into the *dpp* locus. **e-h**, α-GFP and α-Ptc/Wg staining of wing disc expressing the *d2GFP*
reporter at mid-second instar stage (60hr AEL) (**e**), at early third instar stage (72hr AEL) (**f**), at mid-
third instar stage (84hr AEL) (**g**), and at mid- to late- third instar stage (96hr AEL) (**h**). Arrows indicate
dpp transcription outside the stripe of cells. **e'-h'**, Magnified wing discs from (**e-h**). **i-l**, smFISH against
dpp using RNA scope technology. *yw* wing disc at 60 hr AEL (**i**), 72 hr AEL (**j**), 84 hr AEL (**k**), 96 hr
AEL (**l**). **i'-l'**, Magnified wing discs from (**i-l**).

Figure 8. Transient anterior *dpp* source outside Sal domain is required for anterior patterning and growth

a-f, α-pMad, α-Brk, and α-LacZ (**a**, **d**), α-Sal and α-LacZ (**b**, **e**), α-Omb and α-LacZ (**c**, **f**) staining of control wing disc (**a-c**) and wing discs removing *dpp* using *ptc-Gal4* from mid-second instar (**d-f**). **g-j**, α-pMad, α-Brk, and α-LacZ (**g**, **i**), α-Omb and α-LacZ (**h**, **j**) staining of wing disc removing *dpp* using *ptc-Gal4* from mid-second instar (same condition as (d-f)) (**g**, **h**) and using *ci-Gal4* from mid-second instar (**i**, **j**). The genotypes of the wing discs in (d-f) and in (g, h) are identical. **k**, Quantification of each compartment size of wing discs removing *dpp* using *ptc-Gal4* or *ci-Gal4*. Crosses were shifted from 18°C to 29°C at 5day AEL (mid-second instar). α-LacZ staining marks the region where *dpp* is removed upon FLP expression. Dashed white lines mark the A-P compartment border. Scale bar 50 μm.

967
968
Figure 9. Requirements of Dpp dispersal and signaling for wing patterning and growth
a, Based on complete lack of wing pouch by *dpp* disc alleles, it has long been thought that Dpp
dispersal from the anterior stripe of cells and Dpp dispersal-mediated signaling control overall
patterning and growth of the wing pouch.
b, In contrast, the present study shows that while critical for the posterior patterning and growth, Dpp
dispersal is largely dispensable for anterior patterning and growth, and that while critical for the medial
growth, Dpp signaling is dispensable for the lateral wing pouch growth after the wing pouch
specification. In a refined growth equalization model, growth rate is not uniform but higher in the
lateral wing pouch region without Dpp/Brk system. Dpp removes a growth repressor Brk to control
medial wing pouch growth, and Brk represses the lateral wing pouch growth with higher growth rate
to equalize growth rates. Both the Dpp signaling-dependent medial region and -independent lateral
region exist within the wing pouch in the refined model.
981

982
983
1   Bieli, D. *et al.* Development and Application of Functionalized Protein Binders in Multicellular
Organisms. *Int Rev Cell Mol Biol* **325**, 181-213, doi:10.1016/bs.ircmb.2016.02.006 (2016).
2   Harmansa, S. & Affolter, M. Protein binders and their applications in developmental biology.
Development (Cambridge, England) **145**, 10.1242/dev.148874, doi:dev148874 [pii] (2018).
3   Helma, J., Cardoso, M. C., Muyldermans, S. & Leonhardt, H. Nanobodies and recombinant
binders in cell biology. *The Journal of cell biology* **209**, 633-644, doi:10.1083/jcb.201409074
[doi] (2015).
4   Kaiser, P. D., Maier, J., Traenkle, B., Emele, F. & Rothbauer, U. Recent progress in generating
intracellular functional antibody fragments to target and trace cellular components in living
cells. *Biochimica et biophysica acta* **1844**, 1933-1942, doi:S1570-9639(14)00111-3 [pii]
(2014).
5   Vigano, M. A. *et al.* Protein manipulation using single copies of short peptide tags in cultured
cells and in Drosophila melanogaster. *Development* **148**, doi:10.1242/dev.191700 (2021).
6   Ashe, H. L. & Briscoe, J. The interpretation of morphogen gradients. *Development (Cambridge,*
England) **133**, 385-394, doi:133/3/385 [pii] (2006).
7   Rogers, K. W. & Schier, A. F. Morphogen gradients: from generation to interpretation. *Annual*
Review of Cell and Developmental Biology **27**, 377-407, doi:10.1146/annurev-cellbio-092910-
154148 [doi] (2011).
8   Tabata, T. Genetics of morphogen gradients. *Nature reviews.Genetics* **2**, 620-630,
doi:10.1038/35084577 [doi] (2001).
9   Turing, A. The chemical basis of morphogenesis. . *Philosophical Transactions of the Royal*
Society of London B **237** (1952).
10   Kicheva, A. & Gonzalez-Gaitan, M. The Decapentaplegic morphogen gradient: a precise
definition. *Current opinion in cell biology* **20**, 137-143, doi:10.1016/j.ceb.2008.01.008 [doi]
(2008).
11   Sagner, A. & Briscoe, J. Morphogen interpretation: concentration, time, competence, and
signaling dynamics. *Wiley interdisciplinary reviews.Developmental biology* **6**,
10.1002/wdev.1271. Epub 2017 Mar 1020, doi:10.1002/wdev.271 [doi] (2017).
12   Alexandre, C., Baena-Lopez, A. & Vincent, J. P. Patterning and growth control by membrane-
tethered Wingless. *Nature* **505**, 180-185, doi:10.1038/nature12879 [doi] (2014).
13   Apitz, H. & Salecker, I. Spatio-temporal relays control layer identity of direction-selective
neuron subtypes in Drosophila. *Nature communications* **9**, 2295-2018-04592-z,
doi:10.1038/s41467-018-04592-z [doi] (2018).
14   Beaven, R. & Denholm, B. Release and spread of Wingless is required to pattern the proximo-
distal axis of Drosophila renal tubules. *eLife* **7**, 10.7554/eLife.35373, doi:10.7554/eLife.35373
[doi] (2018).
15   Chaudhary, V. *et al.* Robust Wnt signaling is maintained by a Wg protein gradient and Fz2
receptor activity in the developing Drosophila wing. *Development (Cambridge, England)* **146**,
10.1242/dev.174789, doi:dev174789 [pii] (2019).
16   Tian, A., Duwadi, D., Benchabane, H. & Ahmed, Y. Essential long-range action of
Wingless/Wnt in adult intestinal compartmentalization. *PLoS genetics* **15**, e1008111,
doi:10.1371/journal.pgen.1008111 [doi] (2019).
17   Entchev, E. V., Schwabedissen, A. & Gonzalez-Gaitan, M. Gradient formation of the TGF-
beta homolog Dpp. *Cell* **103**, 981-991, doi:10.1016/s0092-8674(00)00200-2 (2000).
18   Teleman, A. A. & Cohen, S. M. Dpp gradient formation in the Drosophila wing imaginal disc.
Cell **103**, 971-980 (2000).
19   Affolter, M. & Basler, K. The Decapentaplegic morphogen gradient: from pattern formation to
growth regulation. *Nature reviews.Genetics* **8**, 663-674, doi:10.1038/nrg2166 (2007).

20 Belenkaya, T. Y. *et al.* Drosophila Dpp morphogen movement is independent of dynamin-
mediated endocytosis but regulated by the glypican members of heparan sulfate proteoglycans.
Cell **119**, 231-244, doi:10.1016/j.cell.2004.09.031 (2004).
21 Hsiung, F., Ramirez-Weber, F. A., Iwaki, D. D. & Kornberg, T. B. Dependence of Drosophila
wing imaginal disc cytonemes on Decapentaplegic. *Nature* **437**, 560-563, doi:nature03951 [pii]
(2005).
22 Schwank, G. *et al.* Formation of the long range Dpp morphogen gradient. *PLoS Biol* **9**,
e1001111, doi:10.1371/journal.pbio.100111
10.1371/journal.pbio.1001111 (2011).
23 Zhou, S. *et al.* Free extracellular diffusion creates the Dpp morphogen gradient of the
Drosophila wing disc. *Curr Biol* **22**, 668-675, doi:10.1016/j.cub.2012.02.065 (2012).
24 Akiyama, T. & Gibson, M. C. Decapentaplegic and growth control in the developing
Drosophila wing. *Nature* **527**, 375-378, doi:10.1038/nature15730 [doi] (2015).
25 Bosch, P. S., Ziukaite, R., Alexandre, C., Basler, K. & Vincent, J. P. Dpp controls growth and
patterning in Drosophila wing precursors through distinct modes of action. *eLife* **6**,
10.7554/eLife.22546, doi:10.7554/eLife.22546 [doi] (2017).
26 Rogulja, D. & Irvine, K. D. Regulation of cell proliferation by a morphogen gradient. *Cell* **123**,
449-461, doi:10.1016/j.cell.2005.08.030 (2005).
27 Rogulja, D., Rauskolb, C. & Irvine, K. D. Morphogen control of wing growth through the Fat
signaling pathway. *Developmental cell* **15**, 309-321, doi:10.1016/j.devcel.2008.06.003 [doi]
(2008).
28 Schwank, G., Restrepo, S. & Basler, K. Growth regulation by Dpp: an essential role for Brinker
and a non-essential role for graded signaling levels. *Development (Cambridge, England)* **135**,
4003-4013, doi:10.1242/dev.025635; 10.1242/dev.025635 (2008).
29 Schwank, G. *et al.* Antagonistic growth regulation by Dpp and Fat drives uniform cell
proliferation. *Developmental cell* **20**, 123-130, doi:10.1016/j.devcel.2010.11.007 [doi] (2011).
30 Schwank, G., Yang, S. F., Restrepo, S. & Basler, K. Comment on "Dynamics of dpp signaling
and proliferation control". *Science (New York, N.Y.)* **335**, 401; author reply 401,
doi:10.1126/science.1210997 [doi] (2012).
31 Wartlick, O., Mumcu, P., Jülicher, F. & Gonzalez-Gaitan, M. Response to Comment on
"Dynamics of Dpp Signaling and Proliferation Control". *Science* **335**, 401,
doi:10.1126/science.1211373 (2012).
32 Wartlick, O. *et al.* Dynamics of Dpp signaling and proliferation control. *Science (New York,*
N.Y.) **331**, 1154-1159, doi:10.1126/science.1200037 [doi] (2011).
33 Zecca, M. & Struhl, G. A unified mechanism for the control of Drosophila wing growth by the
morphogens Decapentaplegic and Wingless. *PLoS biology* **19**, e3001111,
doi:10.1371/journal.pbio.3001111 [doi] (2021).
34 Ben-Zvi, D., Pyrowolakis, G., Barkai, N. & Shilo, B. Z. Expansion-repression mechanism for
scaling the Dpp activation gradient in Drosophila wing imaginal discs. *Current biology : CB*
21, 1391-1396, doi:10.1016/j.cub.2011.07.015 [doi] (2011).
35 Hamaratoglu, F., de Lachapelle, A. M., Pyrowolakis, G., Bergmann, S. & Affolter, M. Dpp
signaling activity requires Pentagone to scale with tissue size in the growing Drosophila wing
imaginal disc. *PLoS biology* **9**, e1001182, doi:10.1371/journal.pbio.1001182 [doi] (2011).
36 Zhu, Y., Qiu, Y., Chen, W., Nie, Q. & Lander, A. D. Scaling a Dpp Morphogen Gradient
through Feedback Control of Receptors and Co-receptors. *Developmental cell* **53**, 724-
739.e714, doi:S1534-5807(20)30418-4 [pii] (2020).
37 Harmansa, S., Hamaratoglu, F., Affolter, M. & Caussinus, E. Dpp spreading is required for
medial but not for lateral wing disc growth. *Nature* **527**, 317-322,
doi:http://dx.doi.org/10.1038/nature15712 (2015).

1081	38	Venken, K. J. *et al.* MiMIC: a highly versatile transposon insertion resource for engineering Drosophila melanogaster genes. *Nature methods* **8**, 737-743, doi:10.1038/nmeth.1662 [doi] (2011).
1084	39	Galeone, A. *et al.* Regulation of BMP4/Dpp retrotranslocation and signaling by deglycosylation. *Elife* **9**, doi:10.7554/eLife.55596 (2020).
1086	40	Mii, Y. *et al.* Quantitative analyses reveal extracellular dynamics of Wnt ligands in Xenopus embryos. *eLife* **10**, 10.7554/eLife.55108, doi:10.7554/eLife.55108 [doi] (2021).
1088	41	Matsuda, S. & Affolter, M. Dpp from the anterior stripe of cells is crucial for the growth of the Drosophila wing disc. *eLife* **6**, 10.7554/eLife.22319, doi:10.7554/eLife.22319 [doi] (2017).
1090	42	Akiyama, T. & Gibson, M. C. Morphogen transport: theoretical and experimental controversies. *Wiley interdisciplinary reviews.Developmental biology* **4**, 99-112, doi:10.1002/wdev.167 [doi] (2015).
1093	43	Matsuda, S., Harmansa, S. & Affolter, M. BMP morphogen gradients in flies. *Cytokine & growth factor reviews* **27**, 119-127, doi:10.1016/j.cytogfr.2015.11.003 [doi] (2016).
1095	44	Restrepo, S., Zartman, J. J. & Basler, K. Coordination of patterning and growth by the morphogen DPP. *Current biology : CB* **24**, R245-255, doi:10.1016/j.cub.2014.01.055 [doi] (2014).
1098	45	Zecca, M. & Struhl, G. Recruitment of cells into the Drosophila wing primordium by a feed-forward circuit of vestigial autoregulation. *Development (Cambridge, England)* **134**, 3001-3010, doi:dev.006411 [pii] (2007).
1101	46	Kim, J., Johnson, K., Chen, H. J., Carroll, S. & Laughon, A. Drosophila Mad binds to DNA and directly mediates activation of vestigial by Decapentaplegic. *Nature* **388**, 304-308, doi:10.1038/40906 [doi] (1997).
1104	47	Kim, J. *et al.* Integration of positional signals and regulation of wing formation and identity by Drosophila vestigial gene. *Nature* **382**, 133-138, doi:10.1038/382133a0 [doi] (1996).
1106	48	Burke, R. & Basler, K. Dpp receptors are autonomously required for cell proliferation in the entire developing Drosophila wing. *Development (Cambridge, England)* **122**, 2261-2269 (1996).
1109	49	Nellen, D., Affolter, M. & Basler, K. Receptor serine/threonine kinases implicated in the control of Drosophila body pattern by decapentaplegic. *Cell* **78**, 225-237, doi:0092-8674(94)90293-3 [pii] (1994).
1112	50	Nellen, D., Burke, R., Struhl, G. & Basler, K. Direct and long-range action of a DPP morphogen gradient. *Cell* **85**, 357-368 (1996).
1114	51	Gibson, M. C. & Perrimon, N. Extrusion and death of DPP/BMP-compromised epithelial cells in the developing Drosophila wing. *Science (New York, N.Y.)* **307**, 1785-1789, doi:10.1126/science.1104751 (2005).
1117	52	Paul, L. *et al.* Dpp-induced Egfr signaling triggers postembryonic wing development in Drosophila. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 5058-5063, doi:10.1073/pnas.1217538110 [doi] (2013).
1120	53	Campbell, G. & Tomlinson, A. Transducing the Dpp morphogen gradient in the wing of Drosophila: regulation of Dpp targets by brinker. *Cell* **96**, 553-562, doi:10.1016/s0092-8674(00)80659-5 (1999).
1123	54	Zecca, M. & Struhl, G. Control of Drosophila wing growth by the vestigial quadrant enhancer. *Development (Cambridge, England)* **134**, 3011-3020, doi:dev.006445 [pii] (2007).
1125	55	Pluckthun, A. Designed ankyrin repeat proteins (DARPins): binding proteins for research, diagnostics, and therapy. *Annual Review of Pharmacology and Toxicology* **55**, 489-511, doi:10.1146/annurev-pharmtox-010611-134654 [doi] (2015).
1128	56	Brauchle, M. *et al.* Protein interference applications in cellular and developmental biology using DARPins that recognize GFP and mCherry. *Biology open* **3**, 1252-1261, doi:10.1242/bio.201410041 [doi] (2014).

| 1131 | 57 | Vigano, M. A. *et al.* DARPins recognizing mTFP1 as novel reagents for in vitro and in vivo |
| 1132 | | protein manipulations. *Biology open* **7**, 10.1242/bio.036749, doi:bio036749 [pii] (2018). |

58 Ziv, O. *et al.* Inverse regulation of target genes at the brink of the BMP morphogen activity
gradient. *J Cell Sci* **125**, 5811-5818, doi:10.1242/jcs.110569 (2012).

59 Lecuit, T. *et al.* Two distinct mechanisms for long-range patterning by Decapentaplegic in the
Drosophila wing. *Nature* **381**, 387-393, doi:10.1038/381387a0 [doi] (1996).

60 Evans, C. J. *et al.* G-TRACE: rapid Gal4-based cell lineage analysis in Drosophila. *Nature*
methods **6**, 603-605, doi:10.1038/nmeth.1356 [doi] (2009).

61 Barrio, L. & Milan, M. Boundary Dpp promotes growth of medial and lateral regions of the
Drosophila wing. *eLife* **6**, 10.7554/eLife.22013, doi:10.7554/eLife.22013 [doi] (2017).

62 Bergantinos, C., Corominas, M. & Serras, F. Cell death-induced regeneration in wing imaginal
discs requires JNK signalling. *Development (Cambridge, England)* **137**, 1169-1179,
doi:10.1242/dev.045559 [doi] (2010).

63 Aguilar, G., Matsuda, S., Vigano, M. A. & Affolter, M. Using Nanobodies to Study Protein
Function in Developing Organisms. *Antibodies (Basel)* **8**, doi:10.3390/antib8010016 (2019).

64 Aguilar, G., Vigano, M. A., Affolter, M. & Matsuda, S. Reflections on the use of protein
binders to study protein function in developmental biology. *Wiley Interdiscip Rev Dev Biol* **8**,
e356, doi:10.1002/wdev.356 (2019).

65 Akiyama, T., User, S. D. & Gibson, M. C. Somatic clones heterozygous for recessive disease
alleles of BMPR1A exhibit unexpected phenotypes in Drosophila. *eLife* **7**,
10.7554/eLife.35258, doi:10.7554/eLife.35258 [doi] (2018).

66 Gavin-Smyth, J., Wang, Y. C., Butler, I. & Ferguson, E. L. A genetic network conferring
canalization to a bistable patterning system in Drosophila. *Current biology : CB* **23**, 2296-2302,
doi:S0960-9822(13)01242-6 [pii] (2013).

67 Adachi-Yamada, T., Fujimura-Kamada, K., Nishida, Y. & Matsumoto, K. Distortion of
proximodistal information causes JNK-dependent apoptosis in Drosophila wing. *Nature* **400**,
166-169, doi:10.1038/22112 [doi] (1999).

68 Conley, C. A. *et al.* Crossveinless 2 contains cysteine-rich domains and is required for high
levels of BMP-like activity during the formation of the cross veins in Drosophila. *Development*
(Cambridge, England) **127**, 3947-3959 (2000).

69 Vuilleumier, R. *et al.* Control of Dpp morphogen signalling by a secreted feedback regulator.
Nat Cell Biol **12**, 611-617, doi:10.1038/ncb2064 (2010).

70 Bangi, E. & Wharton, K. Dpp and Gbb exhibit different effective ranges in the establishment
of the BMP activity gradient critical for Drosophila wing patterning. *Developmental biology*
295, 178-193, doi:S0012-1606(06)00210-7 [pii] (2006).

71 Shimmi, O., Ralston, A., Blair, S. S. & O'Connor, M. B. The crossveinless gene encodes a new
member of the Twisted gastrulation family of BMP-binding proteins which, with Short
gastrulation, promotes BMP signaling in the crossveins of the Drosophila wing. *Dev Biol* **282**,
70-83, doi:10.1016/j.ydbio.2005.02.029 (2005).

72 Aegerter-Wilmsen, T., Aegerter, C. M., Hafen, E. & Basler, K. Model for the regulation of size
in the wing imaginal disc of Drosophila. *Mechanisms of development* **124**, 318-326, doi:S0925-
4773(06)00220-6 [pii] (2007).

73 Aegerter-Wilmsen, T. *et al.* Integrating force-sensing and signaling pathways in a model for
the regulation of wing imaginal disc size. *Development (Cambridge, England)* **139**, 3221-3231,
doi:10.1242/dev.082800 [doi] (2012).

74 Aegerter-Wilmsen, T. *et al.* Exploring the effects of mechanical feedback on epithelial
topology. *Development (Cambridge, England)* **137**, 499-506, doi:10.1242/dev.041731 [doi]
(2010).

75   Nienhaus, U., Aegerter-Wilmsen, T. & Aegerter, C. M. Determination of mechanical stress
1180   distribution in Drosophila wing discs using photoelasticity. *Mechanisms of development* **126**,
1181   942-949, doi:10.1016/j.mod.2009.09.002 [doi] (2009).
1182   76   Schluck, T., Nienhaus, U., Aegerter-Wilmsen, T. & Aegerter, C. M. Mechanical control of
1183   organ size in the development of the Drosophila wing disc. *PloS one* **8**, e76171,
1184   doi:10.1371/journal.pone.0076171 [doi] (2013).
1185   77   Barrio, L. & Milan, M. Regulation of Anisotropic Tissue Growth by Two Orthogonal Signaling
1186   Centers. *Developmental cell* **52**, 659-672.e653, doi:S1534-5807(20)30049-6 [pii] (2020).
1187   78   Zecca, M. & Struhl, G. A feed-forward circuit linking wingless, fat-dachsous signaling, and
1188   the warts-hippo pathway to Drosophila wing growth. *PLoS biology* **8**, e1000386,
1189   doi:10.1371/journal.pbio.1000386 [doi] (2010).
1190   79   Wang, X. & Page-McCaw, A. Wnt6 maintains anterior escort cells as an integral component
1191   of the germline stem cell niche. *Development (Cambridge, England)* **145**, 10.1242/dev.158527,
1192   doi:dev158527 [pii] (2018).
1193   80   Groppe, J. *et al.* Biochemical and biophysical characterization of refolded Drosophila DPP, a
1194   homolog of bone morphogenetic proteins 2 and 4. *The Journal of biological chemistry* **273**,
1195   29052-29065, doi:10.1074/jbc.273.44.29052 [doi] (1998).
1196   81   Dreier, B. & Pluckthun, A. Rapid selection of high-affinity binders using ribosome display.
1197   *Methods in molecular biology (Clifton, N.J.)* **805**, 261-286, doi:10.1007/978-1-61779-379-
1198   0_15 [doi] (2012).
1199   82   Pluckthun, A. Ribosome display: a perspective. *Methods in molecular biology (Clifton, N.J.)*
1200   **805**, 3-28, doi:10.1007/978-1-61779-379-0_1 [doi] (2012).
1201   83   Norman, M., Vuilleumier, R., Springhorn, A., Gawlik, J. & Pyrowolakis, G. Pentagone
1202   internalises glypicans to fine-tune multiple signalling pathways. *eLife* **5**, 10.7554/eLife.13301,
1203   doi:10.7554/eLife.13301 [doi] (2016).
1204

Extended Data Fig 1. Expression of HA trap using *ptc*-Gal4 did not affect Dpp signaling or patterning and growth of the adult wing in the absence of a HA-tagged protein.
a-b, pMad staining of *ptc>+* wing disc (control) (**a**), and *ptc>HA trap* disc (**b**). **c**, Quantification of pMad staining in (**a-b**). **d**, Quantification of wing pouch size in (**a-b**). **e-f**, Adult wing of *ptc>+* (**e**), and *ptc>HA trap* (**f**).

9
10
Extended Data Fig 2. Patterning and growth defects by HA trap and Dpp trap expression
using *ci-Gal4*
a–j, Patterning and growth defects by HA trap. (**a, c, e, g, i**) α-pMad (**a**), α-Brk (**c**), α-Sal (**e**), α-Omb
(**g**), α-Ptc/Wg staining and HA trap (mCherry) (**i**) of control *HA-dpp/HA-dpp, ci>+* (left) and *HA-*
dpp/HA-dpp, ci>HA trap (right). (**b, d, f, h**) Quantification of (**a, c, e, g**), respectively. (**j**)
Quantification of compartment size of wing pouch expressing HA trap using *ci-Gal4*. **k-t**, Patterning
and growth defects by Dpp trap. (**k, m, o, q, s**) α-pMad (**k**), α-Brk (**m**), α-Sal (**o**), α-Omb (**q**), α-
Ptc/Wg staining and HA trap (mCherry) (**s**) of control *HA-dpp/+, ci>+* (left) and *HA-dpp/+, ci>Dpp*
trap (right). (**l, n, p, r, t**) Quantification of (**k, m, o, q, s**) respectively. **t**, Quantification of compartment
size of wing pouch expressing Dpp trap using *ci-Gal4*. **u**, Comparison of compartment size of wing
pouch upon HA trap and Dpp trap expression using *ci-Gal4* (comparison of Extended Data Fig 2j and
Extended Data Fig 2t). Dashed white lines mark the A-P compartment border. Scale bar 50 μm.
23
24

HAdpp/HAdpp, nub, ptc>+

HAdpp/HAdpp, nub, ptc>HAtrap

25
26

Extended Data Fig 3. Patterning and growth defects by concomitant HA trap expression using *ptc-Gal4* and *nub-Gal4*

a-b, Adult wing of control *HA-dpp/HA-dpp, nub, ptc>+* (**a**), and *HA-dpp/HA-dpp, nub, ptc>HA trap* (**b**). **c**, Quantification of compartment size of (**a**) and (**b**). Note that patterning and growth defects were not enhanced by concomitant HA trap expression using *ptc-Gal4* and *nub-Gal4*.

33
34
Extended Data Fig 4. Blocking cell death does not rescue growth defects caused by HA trap or
Dpp trap
a-d, α-Caspase-3 and α-Ptc/Wg staining of control wing disc (**a**), *HA-dpp/HA-dpp, ptc>HA trap* disc
(**b**), and *HA-dpp/+, nub>Dpp trap disc* (**c**). The insets show HA trap (mCherry) expression. Scale bar
50 μm. Note that wing discs expressing HA trap using *ptc*-Gal4 and wing discs expressing Dpp trap
using *nub*-Gal4 were analyzed since each condition showed the most severe phenotypes among Gal4
lines used. (**d**) Quantification of the number of α-Caspase-3 positive cells in (**a-c**). **e-g**, Adult wing of
HA-dpp/HA-dpp, nub>HAtrap (control) (**e**) and *HA-dpp/HA-dpp, nub>HAtrap, p35* (**f**). **g**,
Quantification of compartment size of (**e, f**). **h-j**, Adult wing of *HA-Dpp/+, nub>Dpptrap* (control)
(**h**) and *HA-dpp/+, nub>Dpptrap, p35* (**i**). **j**, Quantification of compartment size of (**h**, **i**).
45

Extended Data Fig 5. *5xQE.DsRed* remains expressed in each compartment where *tkv* is genetically removed from the beginning of second instar stage.

a-d, α-HA (TkvHA[FO]) staining (**a-d**) and α-Brk staining and 5xQE.DsRed expression (**a'-d'**) of control wing disc (**a, b**) and *5xQE.DsRed/+; tkvHA[FO], ci-Gal4/tkvHA[FO]; UAS-FLP/tubGal80ts* (**c, d**). **e-h**, α-HA (TkvHA[FO]) staining (**e-h**) and α-Brk staining and 5xQE.DsRed expression (**e'-h'**) of control wing disc (**e, f**) and *5xQE.DsRed/+; tkvHA[FO]/tkvHA[FO]; UAS-FLP/Hh-Gal4, tubGal80ts* (**g, h**). Scale bar 50 μm. Arrows indicate 5xQE.DsRed expression in the compartment where *tkv* is genetically removed.

Extended Data Fig 6. A part of posterior wing pouch can grow without *tkv*

a-b, α-HA (TkvHA^FO), α-pMad, 5xQE.DsRed, and merge of control wing disc (**a**), and wing disc where *tkv* is genetically removed from the entire P compartment using *Hh*-Gal4 (**b**). Upon removal of *tkv* from the P compartment, the 5xQE.DsRed reporter remained expressed in the P compartment (arrow) despite complete loss of pMad signal and severe growth defects in the P compartment. Note that anterior pMad signal was also affected probably because Hh target *dpp* expression is affected by the reduced number of Hh producing posterior cells.

67
68
69 **Extended Data Fig 7. *5xQE.DsRed* reporter expression is largely independent of Dpp signaling**
70 **a-d**, α-pMad, 5xQE.DsRed, and merge of control (**a**), *dpp^{d8}/dpp^{d12}* (**b**), *brk^{XA}* (**c**), and *brk^{XA}; dpp^{d8}/*
71 *dpp^{d12}* (**d**) wing discs. Scale bar 50 μm.
72
73

74
75
Extended Data Fig 8. Expression of a trap (containing DARPin 1240_C9) using *ptc*-Gal4 did not
affect extracellular distribution of Dpp, pMad signaling, or patterning and growth of the adult
wing
a-b, Extracellular α-Ollas staining (ExOllas) of *Ollas-HA-dpp/+, ptc>+* wing disc (control) disc (**a**),
and *Ollas-HA-dpp/+, ptc>C9* disc (**b**). **c**, Quantification of extracellular α-Ollas staining in (**a-b**). **d-**
e, α-pMad of *Ollas-HA-dpp/+, ptc>+* wing disc (control) disc (**d**), and *Ollas-HA-dpp/+, ptc>C9* disc
(**e**). **f**, Quantification of α-pMad staining in (**d-e**). **g**, Quantification of wing pouch size in (**d-e**). **h-i**,
Adult wing of *Ollas-HA-dpp/+, ptc>+* wing disc (control) (**h**), and *Ollas-HA-dpp/+, ptc>C9* (**f**). Scale
bar 50 μm.
85
86
87

Extended Data Fig 9. Lateral Sal expression is not affected by loss of Dpp signaling

a-b, α-Sal staining of *HA-dpp/+, ptc>+* disc (control) (**a**), and *HA-dpp/+, ptc>Dpp trap* disc (**b**). Each wing disc is from Fig. 5c and Fig. 5d, respectively. In an apical confocal section of control wing disc (z=-4μm), the lateral Sal expression is hidden due to the tissue architecture but in a basal confocal section of control wing disc (z=-15μm), the lateral Sal expression is easily detected (**a**). **c-e**, α-Brk, α-Sal, and merge of *dpp^{FO}/+, tubGal80ts, FLP>+* disc (control) (**c-d**, same wing disc), and *dpp^{FO}/dpp^{FO}, tubGal80ts, ci>FLP* disc (**e**). *dpp* was genetically removed from the mid-second instar. The lateral Sal expression is found in a basal confocal section of control wing disc (z=-12μm) (**c-d**). The lateral Sal expression is not significantly upregulated, although Brk is uniformly upregulated upon generic removal of *dpp* from the entire A compartment using *ci*-Gal4 (**e**). Scale bar 50 μm.

101
102
103 **Extended Data Fig. 10. HA trap can trap Dpp more efficiently than Dpp trap**
104 **a-c**, α-pMad and mCherry (Dpp trap or HA trap) in the P compartment of control wing disc (**a**), wing
105 disc expressing Dpp trap using *ptc*-Gal4 (**b**), wing disc expressing HA trap using *ptc*-Gal4 (**c**). **d**,
106 Quantification of α-pMad of **a-c**. Arrow indicates pMad signal by leaked Dpp from Dpp trap.
107

108
109
110 **Extended Data Fig. 11. pMad signaling at mid-second instar stage**
111 **a-a'**, α-GFP (a) and α-pMad (a') staining of wing disc expressing the *d2GFP* reporter at mid-second
112 instar stage (60hr AEL). Dotted line indicates A-P compartment boundary. **b**, Quantification of
113 pMad signaling.
114
115
116

117
118
Extended Data Fig. 12 Relatively normal anterior patterning and growth by blocking Dpp
dispersal at different time points
a, Control adult wings and adult wings expressing HA trap using *ptc-Gal4* at different time points.
Crosses were shifted from 18 °C to 29 °C at indicated time point. **b-d**, A compartment size (**b**), P
compartment size (**c**), and the peripheral region between L1 and L2 (**d**) were normalized against each
counter part of control wings.
125
126
127

128
129
Extended Data Fig 13. De-repression of Brk by genetic removal of *tkv*

α-HA (TkvHA[FO]) and α-Brk staining of wing discs in which *tkv* was genetically removed from A
compartment using *ci*-Gal4 from different time points. The larvae were raised at 18 °C until a
temperature shift to 29 °C to induce Gal4 expression. Time shown in each figure indicates the time of
dissection after temperature shift. All the discs are the same age and only difference is how long tkv
had been removed. Scale bar 50 μm.