Title: Sex differences and similarities in the neuroimmune response to central administration of

poly I:C

Authors: Caitlin K. Posillico, M.S.[1], Rosa E. Garcia-Hernandez, B.S.[2], & Natalie C. Tronson,

Ph.D.[1]

Affiliations: [1]Psychology Department, [2]University of Michigan, Ann Arbor 48109

Corresponding Author: Natalie C. Tronson

Address: 530 Church St

12   Ann Arbor, MI 48109

13   USA

14 Phone: (734) 936-1495

15 Email: ntronson@umich.edu

16 Keywords: poly I:C; sex differences; neuroimmune; cytokines; hippocampus;

17   chemokines; male; female; interferon; mouse

ABSTRACT

Background: The neuroimmune system is required for normal neural processes, including

modulation of cognition, emotion, and adaptive behaviors. Aberrant neuroimmune activation is

associated with dysregulation of memory and emotion, though the precise mechanisms at play

are complex and highly context dependent. Sex differences neuroimmune activation and

function further complicate our understanding of its roles in cognitive and affective regulation.

Methods: Here, we characterized the physiological sickness and inflammatory response of the

hippocampus following intracerebroventricular (ICV) administration of a synthetic viral mimic,

polyinosinic:polycytidylic acid (poly I:C), in both male and female C57Bl/6 mice.

Results: We observed that poly I:C induced weight loss, fever, and elevations of cytokine and

chemokines in the hippocampus of both sexes. Specifically, we found transient increases in

gene expression and protein levels of IL-1α, IL-1β, IL-4, IL-6, TNFα, CCL2, and CXCL10, where

males showed a greater magnitude of response compared with females. Only males showed

increased IFNα and IFNγ in response to poly I:C, whereas both males and females exhibited

elevations of IFNβ, demonstrating a specific sex difference in the anti-viral response in the

hippocampus.

Conclusion: Our data suggest that type I interferons are one potential node mediating sex-

specific cytokine responses and neuroimmune effects on cognition. Together, these findings

highlight the importance of using both males and females and analyzing a broad set of

inflammatory markers in order to identify the precise, sex-specific roles for neuroimmune

dysregulation in neurological diseases and disorders.

1. BACKGROUND

The neuroimmune system is responsible for surveying the microenvironment and responding to illness, injury, and infection. Importantly, it is also required to carry out normal neural processes (1–3), thus making it well-placed to modify or disrupt learning and memory. In fact, dysregulated immune signaling has been implicated in disorders of affect and cognition, many of which show sex-biases in prevalence and outcomes (4–6). There is significant evidence for sex differences in immune responses in the periphery (7,8), but there is limited literature on similar findings in adult brains. Knowing whether such sex differences occur in similar magnitudes and directions in the neuroimmune system is important for understanding exactly how neuroimmune dysregulation impacts cognition, and contributes to psychiatric and neurological disorders, in both sexes.

Illness, injury, or aseptic triggers of the innate immune system – either bacterial endotoxins (e.g., lipopolysaccharide, LPS) or viral mimics (e.g., polyinosinic:polycytidylic acid, poly I:C) – cause activation of neuroimmune cells, including microglia and astrocytes, and rapid production of cytokines in the brain (9,10). Due to key roles in peripheral inflammation, the cytokines IL-1β (11–13), IL-6 (14–16), and TNFα (17–19) have been the focus of much of the research of neuroimmune function (20). More recently, other cytokines, including interferons (4,21), CCL2 (22,23), and CXCL10 (21,24,25) also play critical roles in modulation of behavior, cognition, and affective states, suggesting that many cytokines play important roles in these processes.

Sex differences in immune and neuroimmune activity have also been reported in various contexts. Females have a greater peripheral immune response compared with males (7). In contrast, neuroimmune cells *in vitro*, including astrocytes derived from male cortical tissue, have a significantly greater reaction to inflammatory insults compared with female-derived cells (26,27). We have identified sex differences in the magnitude, time course, and pattern of cytokines activated in the hippocampus following peripheral LPS (28), and in the long-lasting

impact of LPS on hippocampal function (29). Thus, sex differences in neuroimmune responses

specifically may be a contributing factor to sex differences in neural and cognitive processes

and disorders.

Despite incredible advances in psychoneuroimmunology over the past decade, there are

critical gaps in our knowledge that preclude a holistic understanding of neuroimmune function

and its impacts on cognition and disease. First, with some notable exceptions (22,30,31),

studies have typically focused on a few inflammatory cytokines (e.g., IL-1β, IL-6, and TNFα)

critical for neuroimmune activation and its effects on cognition. Yet more recently, it has become

clear that the massive, coordinated cytokine response observed in the periphery also occurs in

the CNS (28,32). The roles played by other cytokines, and the similarities and differences of this

response in males versus females are yet to be defined (20). Second, the bulk of studies aimed

at understanding neuroimmune activation and its behavioral sequelae have used the gram-

negative bacterial shell and toll-like receptor 4 (TLR4) agonist LPS. Nevertheless, viral illnesses

also trigger changes in behavior, cognition, and emotional states, and significant sex differences

have been observed in the context of viral infections (7,8,33) – an issue that has been propelled

to the forefront during the current COVID-19 pandemic (34,35). Given that viruses act through

distinct toll-like receptors, their impact is likely mediated by a different, albeit overlapping,

pattern of cytokine activation compared with LPS or bacterial triggers. Third, due to its relevance

for disease states, many *in vivo* studies of neuroimmune function use a peripheral immune

challenge. Here, neuroimmune activation is primarily driven by peripheral immune signals that

infiltrate the brain (36). This complicates the interpretation of whether sex differences in cytokine

levels observed in the brain are due to indirect effects based on sex differences in peripheral

immune response, or to direct effect of sex differences in neuroimmune function.

In this study, we aimed to identify a broad set of inflammatory cytokines induced in the

hippocampus by direct neuroimmune stimulation *via* central administration of poly I:C in both

males and females. Elevation of hippocampal cytokines, in particular, is associated with both

disruption of memory processes (1,11,13,37–40) and increased depression-like behaviors

(41,42). Here, we demonstrate that poly I:C induces fever, weight loss, and changes in mRNA

expression and protein levels of cytokines, chemokines, and markers of glial activation across a

24-hour period in both sexes. Notably, only IFNα and IFNɣ showed male-specific patterns of

activation after central poly I:C administration, and many cytokines and chemokines showed a

greater magnitude increase in males compared with females. Whether these sex differences in

neuroimmune activation contribute to sex differences in modulation of cognition and affect and

subsequent prevalence of memory- and mood-related diseases and disorders is an important

area of research for our ongoing studies.

2. METHODS

2.1. Animals

A total of 99 male and female 8-9 week-old C57BL/6N mice from Envigo (Indianapolis,

IN) were used in these experiments. For all experiments, mice were individually housed in

standard mouse cages with *ad libitum* access to food and water in a room with maintained

temperature and pressure under a 12:12-hour light:dark cycle. All mice had at least one week of

acclimation to the colony room prior to any manipulations. All protocols were approved by the

University of Michigan Institutional Animal Care and Use Committee (IACUC).

2.2. Stereotaxic Surgeries

Bilateral guide cannulae (PlasticsOne, Roanoke, VA) targeting the lateral ventricles were

implanted using standard stereotaxic methods (KOPF, Tujunga, CA) at the following

coordinates relative to Bregma: medial-lateral: +/- 1.00mm, anterior-posterior: 0.30mm, dorsal-

ventral: -2.50mm. Animals were administered a pre-surgical analgesic (5mg/kg Carprofen,

subcutaneous) and anesthetized for surgery using an intraperitoneal injection of 250mg/kg of

Avertin (2,2,2-tribromoethanol) which maintained a surgical plane of anesthesia for the duration

of the craniotomy. Bilateral holes were drilled into the skull at the above coordinates, and guide

cannulae were implanted using dental cement. Animals were given a second dose of Carprofen

(5mg/kg, subcutaneous) 24 hours after surgery to maintain a total of 48 hours of analgesia.

Mice were monitored daily for 10 days post-operative and were given at least 2 weeks to

recover from surgery prior to use in experiments.

2.3. Poly I:C Administration

Polyinosinic:polycytidylic acid (poly I:C; Cat. No. P1530; Sigma-Aldrich, St. Louis, MO)

was prepared according to the manufacturer's instructions and sterile-filtered using a 0.22μm

filter prior to administration. For intracerebroventricular (ICV) administration, we infused 20μg of

poly I:C (2μL of 10μg/μL poly I:C) (43) or an equal volume of 0.9% sterile saline *via* the

implanted guide cannula under brief isoflurane anesthesia.

2.4. Sickness Behavior Assessment

To determine whether poly I:C increased physiological measures of sickness in males

and females, body weights and rectal temperatures (RET-3; Physitemp, Clifton, NJ) were

assessed at 2, 4, 6, 12, 24, and 48 hours following ICV administration of poly I:C (n = 10 male; n

= 9 female) or sterile saline (n = 10 male; n = 8 female; Figure 2A). Visual and behavioral

measures of sickness (piloerections, squinted eyes, hunched posture, and low responsivity)

were assessed throughout (44). No changes in overt sickness behaviors were observed for any

experiment (data not shown).

2.4.1. Statistical Analysis of Sickness Behaviors

Analysis of body weight and temperature changes in response to poly I:C was

completed using a mixed repeated-measures ANOVA, using time post-infusion as the within-

subjects factor and treatment and sex as the between-subjects factors with Greenhouse-

Geisser corrections for sphericity. Significant main effects and interactions ($p < 0.05$) were

followed up using post-hoc tests with Bonferroni corrections for multiple comparisons, and effect

sizes were calculated using the partial eta squared method. Any outliers were identified as

samples outside the range of 2 standard deviations from the group mean.

2.5. Characterization of the Acute Neuroimmune Response

We used RNA and protein endpoints to examine induction of cytokines and glial

activation markers in the hippocampus. Males and females were treated with either poly I:C (n =

22 male; n = 24 female) or sterile saline (n = 8/sex) and brains were collected 0.5 hours (n = 5

male; n = 6 female), 2 hours (n = 6/sex), 4 hours (n = 5 male; n = 6 female), and 24 (n = 6/sex)

hours later. All animals were transcardially perfused with 0.1M phosphate buffer to remove

circulating blood from the brain. Both hemispheres of dorsal hippocampus tissue were collected

in separate RNase-/DNase-free, sterile microcentrifuge tubes and immediately flash frozen. All

samples were stored at -80°C before tissue processing.

2.5.1. Quantitative Real-Time PCR

One hemisphere of dorsal hippocampal tissue per mouse was processed for gene

expression analysis using quantitative real-time PCR (qPCR). Frozen samples were

homogenized, and messenger RNA (mRNA) was extracted (PureLink RNA Mini Kit; Cat. No.

12183020; Invitrogen, Carlsbad, CA) under sterile, RNase-free conditions. RNA quality was

assessed using gel electrophoresis, and UV spectroscopy was used to assess RNA purity

(A260/280 > 1.80) and quantity (BioSpectrometer Basic; Eppendorf, Hamburg, Germany). Any

genomic DNA in the sample was removed using DNase treatment, and 800ng of cDNA was

synthesized from each mRNA sample (QuantiTect Reverse Transcriptase Kit; Cat. No. 205314;

Qiagen, Hilden, Germany). Any samples that did not have a high enough concentration of RNA

to make 800ng of cDNA were removed from further analyses (*n* = 3 male; *n* = 5 female).

Relative gene expression was measured using Power SYBR Green PCR Master Mix (Cat. No.

4368702; Applied Biosystems, Foster City, CA) in 10μL reactions (ABI 7500 real-time PCR

system; Cat. No. 4351105; Applied Biosystems).

We measured expression of four commonly-used housekeeping genes: *18s*, *gapdh*,

hprt1, and *rplp0* (all QuantiTect Primer Assays: *18s* Cat. No. QT02448082, *gapdh* Cat. No.

QT01658692, *hprt1* Cat. No. QT00166768, *rplp0* Cat. No. QT00249375; Qiagen). We analyzed

the relative expression of the following genes of interest: *ccl2*, *cd11b*, *cxcl10*, *gfap*, *ifnα*, *ifnβ*,

ifnγ, *il-1α*, *il-1β*, *il-6*, *il-10*, and *tnfα*. The gene primer for *il-1α* was a QuantiTect Primer Assay

(Cat. No. QT00113505; Qiagen). The sequences for the remaining gene primers can be found

in Table 1 and were ordered through Integrated DNA Technologies and diluted to 0.13 μM to be

used for PCR. All Qiagen primers were diluted as per the manufacturer's instruction.

Table 1.

Primer sequences used for real-time PCR.

Gene Target	Forward Primer Sequence (5' to 3')	Reverse Primer Sequence (5' to 3')	NCBI Reference Sequence
ccl2	CCACAACCACCTCAAGCACT	AAGGCATCACAGTCCGAGT	NM_011333.3
cd11b	CGTGAATGGGGACAAACTGAC	GCACTGAGGCTGGCTATTGA	NM_008401.2
cxcl10	TCCATCACTCCCCTTTACCCA	TGGCTTGACCATCATCCTGC	NM_021274.2
gfap	AAACCGCATCACCATTCCTG	CCCGCATCTCCACAGTCTTTA	NM_010277.3
ifnα	AGAGAAGAAACACAGCCCCT	AGCACATTGGCAGAGGAAGA	NM_010502.2
ifnβ	GCTCCAAGAAAGGACGAACAT	GGATGGCAAAGGCAGTGTAA	NM_010510.1
ifnγ	GTCAACAACCCACAGGTCCA	CGACTCCTTTTCCGCTTCCT	NM_008337.4
il-1β	TGCCACCTTTTGACAGTGATG	GCTCTTGTTGATGTGCTGCT	NM_008361.4
il-6	GAGACTTCCATCCAGTTGCCT	TCATTTCCACGATTTCCCAGAG	NM_001314054.1
il-10	CTGGACAACATACTGCTAACCG	AATGCTCCTTGATTTCTGGGC	NM_010548.2
tnfα	ACCCCTTTACTCTGACCCCTT	ACTGTCCCAGCATCTTGTGT	NM_001278601.1

2.5.1.1. Housekeeping Gene Stability Analysis

To control for the transcriptional activity of the samples being analyzed, we confirmed the stability of four housekeeping genes (*18s*, *gapdh*, *hprt1*, and *rplp0*). While many studies use common housekeeping genes such as GAPDH or HPRT1, it is less common for authors to report that their chosen housekeeping gene is indeed stable across experimental groups or tissues prior to use in analyses. Thus, we confirmed the stability of our housekeeping genes using a combination of four techniques to ensure the most reliable quantification of gene expression in our studies. First, we assessed the variability of the candidate genes by measuring the standard deviation of the raw quantification cycle (Cq) values from all samples (Figure 1A). We found that *18s* had the largest standard deviation of Cq values (1.540), followed by *gapdh* (0.527), *rplp0* (0.225), and *hprt1* (0.151; Figure 1B). By this approach, *rplp0* and *hprt1* showed the greatest stability compared to *18s* and *gapdh*, with *hprt1* exhibiting the lowest variability.

Second, we employed a comparative ∆Cq approach in which the standard deviations of the differences in Cq values (∆Cqs) between all possible pairs of candidate genes were compared (45) (Figure 1C). From highest to lowest variability, the genes ranked as follows: *18s* (1.609 average standard deviation), *gapdh* (0.911), *rplp0* (0.764), and *hprt1* (0.753). Again, this method indicated that the most variable genes were *18s* and *gapdh* while the most stable genes were *rplp0* and *hprt1*, and this is most apparent when considering the lowest ∆Cq standard deviation from this method was from the *rplp0* and *hprt1* comparison at 0.206 (Figure 1D).

The third method we employed was that developed by Vandesompele and colleagues, which calculated the average pairwise variation of one candidate gene with all other candidate genes (46). We used R packages ReadqPCR and NormqPCR (47) to calculate *M* stability

values, as depicted in Figure 1E. Consistent with the previous methods, *hprt1* and *rplp0* were

the most stable of the candidate genes, with the lowest pairwise variability, *M* value, of 0.206.

Fourth, and last, we used a model-based stability analysis approach developed by

Andersen et al., an algorithm called NormFinder (v5) (48). This method protects against

identifying two genes via the pairwise approach that might be misinterpreted as being the most

stable if they are coregulated. Using this method, again, *hprt1* and *rplp0* were found to be the

most stable genes with the lowest expression stability values (Figure 1F). However, NormFinder

resulted in *rplp0* having the lowest stability value of 0.326, indicating that the model-based

approach identified *rplp0* as the most stable gene.

Together, these methods identified the two most stable candidate housekeeping genes

as *hprt1* and *rplp0*. Vandesompele et al. (46) posits that using the geometric mean of multiple

housekeeping genes results in more accurate expression levels of genes of interest. We

calculated the geometric mean of the Cq values from *hprt1* and *rplp0* to be used in the $2^{-\Delta\Delta Cq}$

method for calculations of relative expression for our target genes.

Figure 1. Housekeeping gene stability analysis. (A) Distribution of the quantification cycles (Cq) for housekeeping genes *18s*, *gapdh*, *hprt1*, and *rplp0,* with **(B)** associated standard deviations. **(C)** Distribution of the difference of Cq values (ΔCq) between pairs of housekeeping genes, and **(D)** the associated standard deviations. **(E)** Stability values calculated using gene ratio method by Vandesompele et al., 2002, which uses stepwise elimination of lowest stability

240 (highest *M* value) to rank gene stability. **(F)** Stability values calculated using a model-based
241 approach by Andersen et al., 2004 which measures expression variation such that highest
242 stability results in the lowest Rho value. **(G)** Summary of results from each of the four methods
243 of housekeeping gene stability are shown.
244

2.5.1.2. Statistical Analysis of mRNA Gene Expression

For each PCR reaction, the quantification cycle (Cq) was determined, and the $2^{-\Delta\Delta Cq}$

method was used to calculate the relative gene expression of each gene. Any samples with

abnormal amplification curves, melt curves, and/or melt peaks across replicates were removed

from analyses (*n* = 1/sex). Any outliers were identified as samples outside the range of 2

standard deviations from the group mean and excluded from analyses.

Baseline sex differences in relative gene expression (qPCR) were assessed by

evaluating the male and female saline-treated groups. To directly and meaningfully compare

these two groups in the PCR analysis, the male saline-treated group was normalized to the

female saline-treated group and analyzed using independent, two-sample t-tests.

To appropriately analyze sex differences in relative gene expression (qPCR) across the

24-hour time course, we normalized each group to its respective same-sex saline-treated group

to control for any sex differences in gene expression at baseline and used two-way ANOVA

tests using treatment and sex as factors. Significant main effects and interactions ($p < 0.05$)

were followed up using post-hoc tests with Bonferroni corrections for multiple comparisons, and

effect sizes were calculated using the partial eta squared method.

2.5.2. Multiplex Assays

The second hemispheres of dorsal hippocampal tissue were processed as previously

described using low-detergent RIPA buffer sonication (28). Milliplex magnetic bead panel

assays (CCL2, CXCL10, IFNγ, IL-1α, IL-1β, IL-2, IL-4, IL-6, and IL-10; Millipore Sigma,

Burlington, MA) were used as per manufacturer's instructions. Cytokine concentrations were

calculated as pg/mg of hippocampal tissue via Luminex software. Only samples that showed

readable bead counts according to the Luminex software were included in the analyses.

2.5.2.1. Statistical Analysis of Protein Levels

Baseline sex differences in protein levels from multiplex assays were analyzed with

independent, two-sample t-tests comparing the saline-treated groups. To analyze changes in

protein levels from poly I:C across the 24-hour time frame, we used two-way ANOVA tests using

treatment and sex as factors. Significant main effects and interactions ($p < 0.05$) were followed

up using post-hoc tests with Bonferroni corrections for multiple comparisons, and effect sizes

were calculated using the partial eta squared method. Any outliers were identified as samples

outside the range of 2 standard deviations from the group mean and excluded from analyses.

2.6. Data Visualization and Statistical Software

Data visualization and statistical analyses were completed using R 3.6.2 (R Core Team,

2019) with the following packages: dplyr (v0.8.5; (49)), tidyr (v1.0.2; (50)), rstatix (v0.5.0; (51)),

DescTools (v0.99.34; (52)), sjstats (v0.17.9; (53)), ReadqPCR and NormqPCR (47), ggplot2

(54), gridExtra (v2.3; (55)), pheatmap (v1.0.12; (56)), and viridis (v0.5.1; (57)).

3. RESULTS

3.1. Central administration of poly I:C induces physiological sickness responses

Both females and males showed physiological responses to poly I:C. Whereas both

saline- and poly I:C-treated animals showed changes in weight across the 48-hour period

(Figure 2B, main effect of Time: $F(3.13, 96.92) = 28.899$, $p < 0.001$, $\eta^2_p = 0.482$), poly I:C

caused weight loss in both sexes (main effect of Treatment: $F(1, 31) = 8.781$, $p = 0.006$, $\eta^2_p = $

0.221; trend towards a Time x Treatment interaction: $F(3.13, 96.92) = 2.476$, $p = 0.064$, $\eta^2_p = $

0.074). Specifically, males and females treated with poly I:C lost significantly more weight than

the saline-treated animals at the 12- (p = 0.004) and 24-hour (p = 0.022) time points. By 48

hours post-treatment, the weights of poly I:C-treated animals had recovered and were no longer

different from those of saline-treated animals (p = 1.00).

Figure 2. Analysis of sickness behaviors following poly I:C administration. (A) Timeline of
body weight and temperature measurements following poly I:C or sterile saline administration.
(B) Average weight change from baseline (Time = 0 hour) prior to treatment. **(C)** Average body
temperature as measured via rectal thermometer. Analyzed using mixed repeated-measures
ANOVA. *****: $p < 0.05$ poly I:C- vs saline-treated groups.

In both males and females, poly I:C caused significant increases in body temperature

relative to the saline-treated group (Figure 2C; main effect of Treatment: $F_{(1, 31)}$ = 23.759, $p <$

0.001, η^2_p = 0.434; Time x Treatment interaction: $F_{(4.6, 142.62)}$ = 11.635, $p < 0.001$, η^2_p =

0.273). Post-hoc tests revealed that body temperature began to increase 2 hours following poly

I:C (p = 0.068), remained elevated at the 4- (p < 0.001) and 6-hour (p < 0.001) time points, and

recovered to saline-treated body temperatures by 12 hours post treatment (all p = 1.00).

3.2. Baseline sex differences in mRNA expression and protein levels of select

hippocampal immune molecules

3.2.1. Interleukins

Select immune signaling markers showed baseline sex differences. mRNA expression of

interleukin *(il)-1α* exhibited a trend towards a sex difference in baseline expression (Figure 3C1;

t(12) = 2.006, p = 0.068), and *il-6* showed a significant sex difference (Figure 3E1; t(11) = 3.079,

p = 0.01, 95% CI [0.182, 1.062]) where females showed greater mRNA expression than males.

These results did not hold true for protein levels of IL-1α or IL-6. Rather, multiplex data showed

that protein levels of IL-1β and IL-10 were significantly different between saline-treated males

and females (Figures 4D and 4F, respectively; IL-1β: t(13) = 4.275, p = 0.001, 95% CI [5.682,

17.291]; IL-10: t(13) = 2.236, p = 0.044, 95% CI [0.672, 39.314]). Again, for these cytokines,

females showed greater protein levels at baseline compared with males.

3.2.2. Interferons

Two interferons (IFN), *ifnα* and *ifnγ*, also showed baseline sex differences in relative

mRNA expression (Figures 3G1 and 3I1: IFNα: t(12) = 5.546, p = 0.0001, 95% CI [0.441, 1.01];

IFNγ: t(11) = 2.995, p = 0.012, 95% CI [0.259, 1.694]). Likewise, protein levels of IFNγ exhibited

a sex difference in saline-treated groups (Figure 4G; t(14) = 6.475, p < 0.001, 95% CI [6.534,

13.006]). For both mRNA expression and protein levels, female levels were greater than males.

3.2.3. Chemokines

Expression of CC chemokine *ccl2* similarly followed this pattern at baseline (Figure 3K1;

t(12) = 3.287, p = 0.006, 95% CI [0.259, 1.279]), and this held true for protein levels of CCL2 as

well (Figure 4H; t(12) = 2.751, p = 0.018, 95% CI [12.798, 110.318]). Interestingly, all genes with

sex differences in saline-treated animals showed that both female baseline mRNA expression

and protein levels were at least twice as high as that of males.

Figure 3. mRNA gene expression of cytokines, chemokines, and markers of glial activation in the hippocampus. Baseline gene expression was measured by normalizing the male saline-treated group to the female saline-treated group and analyzed using independent, two-sample t-tests. Baseline expression of **(A1)** CD11b, **(B1)** GFAP, **(C1)** IL-1α, **(D1)** IL-1β, **(E1)** IL-6, **(F1)** IL-10, **(G1)** IFNα, **(H1)** IFNβ, **(I1)** IFNγ, **(J1)** TNFα, **(K1)** CCL2, and **(L1)** CXCL10 are shown. Gene expression changes following poly I:C treatment were calculated by normalizing time points after poly I:C treatment to the saline-treated groups within sex to eliminate confounding variables of baseline sex differences. Gene expression was analyzed using two-way ANOVA tests for **(A2)** CD11b, **(B2)** GFAP, **(C2)** IL-1α, **(D2)** IL-1β, **(E2)** IL-6, **(F2)** IL-10, **(G2)** IFNα, **(H2)** IFNβ, **(I2)** IFNγ, **(J2)** TNFα, **(K2)** CCL2, and **(L2)** CXCL10. ***** above a bracket covering both sexes indicates a significant main effect of sex ($p < 0.05$); * above a horizontal line covering just one sex indicates a significant main effect of treatment ($p < 0.0.5$); ***** above a single bar indicates a significant post-hoc test ($p < 0.05$) vs the saline-treated group within sex; **%** above a single bar indicates a significant post-hoc test ($p < 0.05$) vs females at the same time point.

351
Figure 4. Protein levels of cytokines and chemokines in the hippocampus. Protein
concentration (pg/mg) of **(A)** IL-1α, **(B)** IL-2, **(C)** IL-4, **(D)** IL-1β, **(E)** IL-6, **(F)** IL-10, **(G)** IFNγ, **(H)**
CCL2, and **(I)** CXCL10 are shown. Two-way ANOVA tests were used to analyze these data. *
above a bracket covering both sexes indicates a significant main effect of sex (*p* < 0.05); *
above a horizontal line covering just one sex indicates a significant main effect of treatment (*p* <
0.0.5); * above a single bar indicates a significant post-hoc test (*p* < 0.05) vs the saline-treated
group within sex; **%** above a single bar indicates a significant post-hoc test (*p* < 0.05) vs
females at the same time point.

3.2.4. Glial activation markers and other cytokines did not show baseline sex differences

It is equally important to note the markers that did not show sex differences at baseline.

Both the microglial activation marker *cd11b* and the astrocyte activation marker *gfap* showed

equivalent levels of gene expression in the saline-treated groups in both sexes (Figures 3A1

and 3B1, respectively; *cd11b*: $t(12) = 0.723$, $p = 0.483$; *gfap*: $t(12) = 1.603$, $p = 0.135$). Other

commonly examined inflammatory cytokines, including IL-2, IL-4, and tumor necrosis factor

(TNF)α similarly showed no sex difference at baseline (IL-2 $p = 0.420$; IL-4 $p = 0.633$; TNFα =

0.139). No other sex differences in mRNA gene expression or protein levels were observed

under baseline conditions.

3.3. Central poly I:C causes greater increases in male mRNA expression of hippocampal

cytokines compared with females

3.3.1. Glial Activation Markers

Poly I:C treatment significantly increased expression of both *cd11b* and *gfap*, although

this appeared to be true only at the 24-hour time point (Figures 3A2 and 3B2; *cd11b* main effect

of Treatment: $F(4, 42) = 12.96$, $p < 0.001$, $\eta^2_p = 0.552$; *gfap* main effect of Treatment: $F(4, 42) =$

12.992, $p < 0.001$, $\eta^2_p = 0.553$). Sex did not affect the response of either *cd11b* or *gfap* to poly

I:C (insignificant Sex x Treatment interactions: *cd11b*: $F(4, 42) = 0.684$, $p = 0.607$; *gfap*: $F(4, 42)$

$= 0.923$, $p = 0.460$).

3.3.2. Interleukins

Poly I:C caused increased expression of *il-1α*, *il-1β*, and *il-6* in both males and females

(Figures 3C2, 3D2, and 3E2; main effects of Treatment: *il-1α*: $F(4, 42) = 9.784$, $p < 0.001$, $\eta^2_p =$

0.482; *il-1β*: $F(4, 42) = 9.512$, $p < 0.001$, $\eta^2_p = 0.475$; *il-6*: $F(4, 42) = 22.28$, $p < 0.001$, $\eta^2_p =$

0.680). In males, expression began to increase at the 2-hour time point following poly I:C

treatment for *il-1α* ($p = 0.015$), *il-1β* ($p = 0.057$), and *il-6* ($p = 0.029$), showed peaks at the 4-

hour time point ($p < 0.001$ for all), and decreased to saline-treated levels by 24 hours ($p = 1.00$

for all). Each of these genes also showed an overall greater expression in males than females

(main effects of Sex: *il-1α*: $F(1, 42) = 6.398$, $p = 0.015$, $\eta^2_p = 0.132$; *il-1β*: $F(1,42) = 6.695$, $p =$

0.013, $\eta^2_p = 0.137$; *il-6*: $F(1, 42) = 21.1$, $p < 0.001$, $\eta^2_p = 0.334$), and a significantly greater

magnitude of response in males compared with females (Sex x Treatment interactions: *il-1α*:

$F(4, 42) = 3.103$, $p = 0.025$, $\eta^2_p = 0.228$; *il-1β*: $F(4, 42) = 4.288$, $p = 0.005$, $\eta^2_p = 0.290$; *il-6*: $F(4,$

$42) = 15$, $p < 0.001$, $\eta^2_p = 0.588$). Post-hoc tests revealed for all three genes, males exhibited an

even greater response at only the 4-hour time point compared with females ($p < 0.05$ for all).

Notably, the peak *il-1α* and *il-1β* expression in males was roughly 3-fold higher than that of the

peak female expression for these cytokines, and the *il-6* peak expression in males was more

than 10-fold higher than that of females (Figures 3C2, 3D2, and 3E2).

397   Males showed greater *il-10* gene expression across all time points compared with

females (Figure 3F2; main effect of Sex: $F(1, 39) = 25.642$, $p < 0.001$, $\eta^2_p = 0.397$). Additionally,

poly I:C significantly increased gene expression of *il-10* in males, but not females (Sex x

Treatment interaction: $F(4, 39) = 3.304$, $p = 0.02$, $\eta^2_p = 0.253$). Specifically, male expression of

il-10 at the 4-hour time point following poly I:C was significantly greater than that of saline-

treated controls ($p = 0.001$), and this was also greater than the 4-hour expression in females (p

$= 0.001$).

3.3.3. Tumor Necrosis Factor Alpha

405   Gene expression of *tnfα* increased in response to poly I:C, males had significantly higher

expression than females overall, and males showed a greater magnitude of response compared

with females (Figure 3J2; main effect of Treatment: $F(4, 42) = 6.407$, $p = 0.0004$, $\eta^2_p = 0.379$;

main effect of Sex: $F(1, 42) = 10.1$, $p = 0.003$, $\eta^2_p = 0.194$; Sex x Treatment interaction: $F(4, 42)$

$= 4.117$, $p = 0.007$, $\eta^2_p = 0.282$). Post-hoc tests showed that males 4 hours post-treatment had

significantly greater expression than those treated with saline ($p < 0.001$), and this was again

greater than the 4-hour peak expression in females ($p = 0.001$).

3.3.4. Interferons

Both *ifnα* and *ifnγ* showed a similar response pattern to poly I:C, whereby males treated

with poly I:C exhibited a significant acute increase in gene expression of both cytokines, but

females did not show the same response (Figures 3G2 and 3I2; *ifnα*: main effect of Treatment:

$F(4, 42) = 5.007$, $p = 0.002$, $\eta^2_p = 0.323$; Sex x Treatment interaction: $F(4, 42) = 3.35$, $p = 0.018$,

$\eta^2_p = 0.242$; *ifnγ*: main effect of Treatment: $F(4, 40) = 4.698$, $p = 0.003$, $\eta^2_p = 0.32$; Sex x

Treatment interaction: $F(4, 40) = 4.178$, $p = 0.006$, $\eta^2_p = 0.295$). Specifically, 4 hours after poly

I:C treatment, males showed significantly elevated expression compared to the saline-treated

controls ($p = 0.0001$), and this was greater in magnitude than the 4-hour time point in females (p

$= 0.001$). Female *ifnα* and *ifnγ* did not seem to respond to poly I:C treatment at any time point.

In contrast, poly I:C transiently increased expression of *ifnβ* in both males and females,

and there were no sex differences seen in magnitude of expression increase (Figure 3H2; main

effect of Treatment: $F(4, 42) = 4.855$, $p = 0.003$, $\eta^2_p = 0.316$; insignificant Sex x Treatment

interaction: $F(4, 42) = 1.297$, $p = 0.287$). Unlike all other cytokines examined in this study, peak

expression appeared to be at the 2-hour time point, and expression began decreasing again by

4 hours post-treatment. The magnitude increase was also notable, with a 100-fold increase in

females and a 300-fold increase in males.

3.3.5. Chemokines

Poly I:C significantly increased the expression of both *ccl2* and *cxcl10* in males and

females, with a peak increase in expression at 4-hours post-infusion (Figures 3K2 and 3L2;

main effects of Treatment: *ccl2*: $F(4, 41) = 25.47$, $p < 0.001$, $\eta^2_p = 0.713$; *cxcl10*: $F(4, 42) =$

87.37, $p < 0.001$, $\eta^2_p = 0.893$). Also consistent with many other cytokines, expression of both

ccl2 and *cxcl10* was greater overall in males compared with females (main effects of Sex: *ccl2*:

$F(1, 41) = 44.55$, $p < 0.001$, $\eta^2_p = 0.521$; *cxcl10*: $F(1, 42) = 92.79$, $p < 0.001$, $\eta^2_p = 0.688$). Finally,

males showed a markedly greater magnitude of response than did females for both chemokines

(Sex x Treatment interactions: *ccl2*: $F(4, 41) = 20.96$, $p < 0.001$, $\eta^2_p = 0.672$; *cxcl10*: $F(4, 42) =$

60.51, $p < 0.001$, $\eta^2_p = 0.852$). Remarkably, male *ccl2* expression peaked at nearly 450-fold

greater than the expression of saline-treated males compared to a roughly 20-fold increased

peak in females (Figure 3K2). Similarly, *cxcl10* expression in males peaked at nearly 500-times

that of saline-treated males while female *cxcl10* expression peaked at just over 40-times greater

than saline-treated females (Figure 3L2). These massive increases in gene expression are

reflected in the strong effect sizes noted for the interaction effect above. Post-hoc tests

confirmed that the male 2- and 4-hour time points post-treatment showed significantly greater

gene expression of both *ccl2* and *cxcl10* than saline-treated males ($p < 0.001$). Additionally,

both the male 2- and 4-hour time points of both genes proved to be significantly greater than the

2- and 4-hour time points in females, respectively ($p < 0.01$).

3.4. Central poly I:C increases select hippocampal cytokine protein levels in males and

females

3.4.1. Interleukins

We analyzed several members of the interleukin family of cytokines, including classically

inflammatory IL-1α, IL-1β, IL-2, IL-4, and IL-6, and classically anti-inflammatory IL-10. Of these,

IL-1α, IL-1β, IL-4, and IL-6 significantly increased following ICV poly I:C administration in both

males and females (Figures 4A, 4C, 4D, 4E; main effects of Treatment: IL-1α: $F(4, 51) = 3.523$,

$p = 0.013$, $\eta^2_p = 0.216$; IL-1β: $F(4, 51) = 5.721$, $p = 0.001$, $\eta^2_p = 0.31$; IL-4: $F(4, 51) = 5.146$, $p =$

0.001, $\eta^2_p = 0.288$; IL-6: $F(4,51) = 10.298$, $p < 0.001$, $\eta^2_p = 0.447$). In all cases, protein levels

appear to increase to a peak 4 hours following poly I:C, similar to the effects seen in mRNA

expression. In addition, both IL-1β and IL-4 also exhibited a main effect of Sex such that protein

levels of these cytokines, regardless of time point, were significantly higher in females

compared with males (Figures 4C and 4D; IL-1β: $F(1, 51) = 114.226$, $p < 0.001$, $\eta^2_p = 0.691$; IL-

4: $F(1, 51) = 11.03$, $p = 0.002$, $\eta^2_p = 0.178$). No interactions of sex and treatment were found for

any of the interleukin cytokines examined here (IL-1α: $F(4, 51) = 0.446$, $p = 0.775$; IL-1β: $F(4,$

51$) = 0.513$, $p = 0.726$; IL-4: $F(4, 51) = 0.982$, $p = 0.426$; IL-6: $F(4, 51) = 1.779$, $p = 0.148$).

Neither IL-2 nor IL-10 showed any effects of poly I:C treatment in either sex (Figures 4B and 4F;

insignificant main effects of Treatment: IL-2: $F(4, 51) = 1.498$, $p = 0.217$; IL-10: $F(4, 51) = 1.122$,

$p = 0.357$). Similar to IL-1β and IL-4, though, we found that females had overall higher levels of

IL-10 than did males (main effect of Sex; $F(1, 51) = 20.27$, $p < 0.001$, $\eta^2_p = 0.284$).

3.4.2. Interferon

We also measured IFNγ levels in response to poly I:C. Unlike mRNA expression, IFNγ

protein levels did not change following poly I:C administration in either sex (Figure 4G;

insignificant main effect of Treatment: $F(4, 52) = 1.93$, $p = 0.119$). However, IFNγ levels were

higher in females relative to males (main effect of Sex: $F(1, 52) = 150.64$, $p < 0.001$; $\eta^2_p = $

0.743). This was consistent with mRNA expression data where saline-treated females also

showed significantly higher expression of *ifnγ* at baseline than did males (see Figure 3Q).

3.4.3. Chemokines

Finally, we measured protein levels of the chemokines CCL2 and CXCL10 in both

sexes. Both chemokines were significantly affected by central administration of poly I:C and in

different ways in males and females (Figures 4H and 4I; CCL2: main effect of Treatment: $F(4,$

46$) = 18.517$, $p < 0.001$, $\eta^2_p = 0.617$; Sex x Treatment interaction: $F(4, 46) = 3.381$, $p = 0.017$,

$\eta^2_p = 0.227$; CXCL10: main effect of Treatment $F(4, 52) = 14.54$, $p < 0.001$, $\eta^2_p = 0.528$; Sex x

Treatment interaction: $F(4, 52) = 2.796$, $p = 0.035$, $\eta^2_p = 0.177$). In males, CCL2 levels increased

earlier (at 2 hours) post-infusion relative to females. For CXCL10, females took longer for

protein levels to begin to decrease as compared to the time course in males, with females still

showing the massive elevation at 24 hours post-infusion as they did at 4 hours.

Notably, both CCL2 and CXCL10 levels showed the most substantial increases out of all

cytokines measured in protein analysis in the hippocampus. Specifically, female CCL2 levels

induced by poly I:C peaked at approximately 4 times that of the saline-treated animals, and

male CCL2 levels peaked nearly 8 times that of saline-treated males (Figure 4H). More

remarkably, female CXCL10 levels rose roughly 16-fold by 4 hours after poly I:C administration,

and males showed a 12-fold increase by 4-hours (Figure 4I).

3.5. Summary of mRNA and protein data

Overall, hippocampal mRNA expression and protein levels of most of the cytokines and

chemokines examined in this experiment responded to central administration of poly I:C in both

males and females. We found significant sex differences in baseline mRNA expression and

protein levels of several cytokines, where females showed greater basal levels than males. In

addition, we found the magnitude of mRNA expression increases was greater in males than

females. Protein data showed this to be true only for 2 chemokines, CCL2 and CXCL10.

The heatmaps shown in Figure 5 indicate that most of the immune signaling molecules

affected in the immediate phase following poly I:C treatment peaked at 4 hours for both mRNA

expression (Figure 5A) and protein levels (Figure 5B) and returned to levels of saline-treated

animals by 24 hours post-infusion.

Figure 5. Heatmaps of gene expression and protein levels in the hippocampus. Changes in **(A)** mRNA gene expression and **(B)** protein levels for cytokines, chemokines, and markers of glial activation are shown. Values are centered and scaled across rows to highlight changes across the time course for each gene of interest; thus, differences in magnitude between genes are not depicted.

4. DISCUSSION

Here we demonstrated that central administration of poly I:C induced both physiological measures of sickness and concomitant neuroinflammatory activation in both male and female mice. Both males and females treated with poly I:C mounted an acute fever response and showed weight loss that recovered to saline-treated levels 48 hours later. Additionally, both sexes showed acute cytokine and chemokine responses in the hippocampus, as measured by both mRNA expression and protein levels, that followed the time course of fever. Interestingly, mRNA gene expression of *il-1α*, *il-1β*, *il-6*, *il-10*, *ifnα*, *tnfα*, *ccl2*, and *cxcl10* and protein levels of CCL2 and CXCL10 showed a stronger response in males compared with females. Further, gene expression of *il-10*, *ifnα*, and *ifnγ* increased in males only. Given the specific role of type I interferons in viral response (58–60), this sex difference may have important implications for how neuroimmune activation differentially disrupts cognition in males and females.

Poly I:C treatment in both sexes resulted in a significant and transient increase in gene expression and protein levels of most, but not all, cytokines and chemokines in the hippocampus, including IFNβ, IL-1α, IL-1β, IL-6, TNFα, CCL2, and CXCL10. That administration of an immune stimulant, including viral mimics such as poly I:C, induces a neuroimmune response is not new; however, most of the previous studies on poly I:C used peripheral administration (61–64). As such, multiple, indirect mechanisms are likely involved in causing inflammation in the brain (36). Peripheral administration of poly I:C, specifically, was found to induce neuroinflammation through a separate and distinct pathway than central administration (43). Additionally, these studies examined a subset of cytokines or chemokines, including the most commonly examined IL-1β, IL-6, and TNFα, and type I interferons that typically respond to viruses. Given evidence of mechanistic complexities governing neuroinflammation, particularly from stimulants such as poly I:C, and given that there are over 300 cytokines with important roles in the immune system and neural function, it is critical to begin looking beyond IL-1β, IL-6,

and TNFα and more strongly consider implications of such limits in experimental design for the

field of psychoneuroimmunology.

In addition to the intricacies of how immune stimulants such as poly I:C activate the

neuroimmune system, males and females differ in immune and neuroimmune responses. The

direction of these differences depends on whether you are looking in the periphery (7) or the

brain (26,27). We found that mRNA gene expression of *il-1*α, *il-1*β, *il-6*, *il-10*, *ifn*α, *tnf*α, *ccl2*, and

cxcl10 and protein levels of CCL2 and CXCL10 in the hippocampus showed a stronger

response in males compared with females. A greater magnitude of cytokine and chemokine

response in males is consistent with previous findings that male-derived astrocytes have a

greater reaction to inflammatory insults compared with females (26,27,65,66). Poly I:C is

recognized by microglia, astrocytes, and neurons via toll-like receptor 3 (TLR3) (67–69). The

interaction of these 3 cell types is crucial in mediating inflammatory responses (70,71). Given

that TLR3 shows much greater expression in astrocytes relative to microglia (72), the reaction of

astrocytes in males may be driving the sex differences in magnitude gene expression response

of cytokines following poly I:C. The astrocyte activation marker, GFAP, and the microglial

activation marker, CD11b, did not increase until 24 hours after poly I:C treatment and did not

show sex differences. However, this does not absolve astrocytes or microglia from the acute

response to poly I:C. Specifically, Norden and colleagues found that cytokine gene expression

from both astrocytes and microglia preceded increases in astrocyte and microglial activation

markers, (GFAP and Iba1, respectively), and that these activation markers similarly did not

show reliable increases until the 24-hour time point (73). It is likely, therefore, that a greater

astrocytic reaction in males is driving the greater magnitude of response to poly I:C by males.

Viral infections are not immune to observations of sex differences. Specifically, males

and females show sex differences in prevalence, intensity, and survival outcomes of viruses

(7,8,33). Many report that when females show better survival outcomes or seem more protected

against certain viruses, it is due to the female immune system having a greater anti-viral

response that allows the virus to be cleared faster (33,74–76). Data on SARS-CoV-2, the virus

that causes the coronavirus disease 2019 (COVID-19), has shown a similar sex difference in

intensity of infection, with more males having worse symptoms, including neurological

complications, and survival outcomes relative to females (77–81). However, in contrast to the

data that suggests a greater immune response results in faster recovery in females, recent

studies indicate that higher levels of inflammatory cytokines including IL-6, IL-1β, CXCL10, and

CCL2 are actually correlated with worse COVID-19 outcomes (35,82). Unequivocally, sex as a

biological factor matters in the context of immune and neuroimmune responses as well as in the

acute and chronic implications of neuroimmune modulation of neural function, cognition, and

emotion in males and females. Limiting studies by only using one sex and extrapolating the data

to the other is not only inappropriate given the intricacies of sex differences in the literature

discussed here, but it is also potentially dangerous for both development and administration of

treatments in inflammatory diseases and disorders.

We observed that most cytokines examined here showed a sex difference in response to

poly I:C. Whereas others have reported increases in select inflammatory markers following poly

I:C treatment, these studies used either only used male (62,63) or female rodents (61,64). To

the best of our knowledge, this is the first direct comparison of hippocampal cytokines in males

and females as a consequence of poly I:C. Whether the greater magnitude in male response to

poly I:C indicates greater neuroprotection or vulnerability is yet to be determined. This is an

especially intriguing question given that we also directly compared mRNA expression and

protein levels of cytokines and chemokines at baseline and found that females had greater

basal immune activity relative to males.

Perhaps it is not only that females are able to mount a greater immune response to help

clear viral loads faster and aid in a faster recovery as previously discussed (33,74–76), but also

that females start out with greater immune activity that allows them to reach necessary

activation states faster than males. Specifically, maybe females do not need to have as great of

a magnitude increase in cytokines and chemokines when the neuroimmune system is activated

because they already have "more players in the game". Future work will need to address

whether and how sex differences in the cytokine and chemokine basal levels or activation in

response to immune challenge result in functional differences in modulation of neural function

and contribute to sex-biases in neurological and psychiatric disease.

Of particular note, we observed a sex-specific pattern of expression of the interferon

family of cytokines in the hippocampus. Specifically, males showed increases in IFNα, IFNβ,

and IFNγ, but females only showed a significant response in IFNβ. This is consistent with

previous findings that showed increased gene expression of IFNβ, but not IFNα, in females in

response to peripheral poly I:C, though this study did not measure these effects in males for

comparison (64). Type I interferons, IFNα and IFNβ, are key to the anti-viral response of the

immune system and, as such, are known to respond to viral stimulants including poly I:C (58–

60,83,84). Consistent with our data, in which IFNβ showed an early peak expression levels, type

I interferon activity is responsible for inducing inflammatory cytokines such as IL-6 and TNFα

(62,64). Additionally, interferon signaling from poly I:C treatment also results in altered

glutamatergic signaling (83,84), which is known to be critical for hippocampal memory formation

(85). Given that males induce both IFNα and IFNβ in the hippocampus following poly I:C

whereas females only induce IFNβ, interferons have important implications for sex differences in

neuroimmune disruption of cognition from virus or viral-like stimulants.

5. CONCLUSION

This study characterizes the sickness and neuroimmune responses to central

administration of poly I:C, and we observed sex-specific patterns of hippocampal cytokine

responses. Specifically, we identified type I interferons as one potential node mediating sex-

specific cytokine responses and neuroimmune effects on synaptic plasticity and cognition.

Additionally, the magnitude of response of cytokines such as CCL2 and CXCL10 highlight the

importance of future work incorporating a more comprehensive set of inflammatory markers

using multiple endpoints. Neuroimmune activation is known to play a role in cognitive deficits

and affective dysregulation in diseases such as Alzheimer's Disease and other dementias (86),

Post-Traumatic Stress Disorder (87,88), depression (4,89), and now also COVID-19 (90). Given

the sex/gender biases in prevalence, severity, and/or survival outcomes, identifying sex-specific

neuroimmune responses will provide novel targets for personalized prevention and treatment of

these diseases.

LIST OF ABBREVIATIONS

ANOVA = analysis of variance

CCL2 = C-C motif chemokine ligand 2

Cd11b = cluster of differentiation molecule 11B

cDNA = complementary deoxyribonucleic acid

COVID-19 = coronavirus disease-19

CXCL10 = C-X-C motif chemokine ligand 10

DNA = deoxyribonucleic acid

GAPDH = glyceraldehyde-3-phosphate dehydrogenase

GFAP = glial fibrillary acidic protein

HPRT1 = hypoxanthine phosphoribosyltransferase 1

Iba1 = ionized calcium-binding adaptor molecule 1

ICV = intracerebroventricular

IFNα = interferon α

IFNβ = interferon β

IFNγ = interferon γ

IL-1α = interleukin-1α

IL-1β = interleukin-1β

IL-2 = interleukin-2

IL-4 = interleukin-4

IL-6 = interleukin-6

IL-10 = interleukin-10

kg = kilogram

LPS = lipopolysaccharide

mg = milligram

mm = millimeter

mRNA = messenger ribonucleic acid

PCR = polymerase chain reaction

Poly I:C = polyinosinic:polycytidylic acid

qPCR = quantitative real-time polymerase chain reaction

RNA = ribonucleic acid

RPLP0 = ribosomal protein lateral stalk subunit P0

SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2

TLR = toll-like receptor

TNFα = tumor necrosis factor α

μg = microgram

μL = microliter

μm = micrometer

DECLARATIONS

Ethics Approval

All animal protocols used in these experiments were approved by the Institutional Animal

Care and Use Committee (IACUC) at the University of Michigan.

Consent for Publication

Not applicable.

Availability of Data and Materials

The data used and analyzed for the current study are available from the corresponding

author upon reasonable request.

Competing Interests

The authors have no competing interests to declare.

Funding

These experiments were supported by a University of Michigan Office of Research

Award to NCT. This funding body did not provide input for the design of the study, collection,

analysis, data interpretation, or writing of this manuscript. The award was used solely to fund

the cost of animals, reagents, and equipment used in these experiments.

Author's Contributions

CKP designed the experiments, performed all animal surgeries and treatments, collected

physiological measurements (body temperature and weights) and tissue, processed tissue

samples and conducted all molecular biology assays (qPCR and multiplex assays), ran

statistical analyses, and wrote the manuscript. RGH aided in the collection of physiological

measurements (body temperature and weights) and monitored animals for sickness behaviors

and surgical recovery throughout the experiments. NCT provided intellectual contributions

regarding experimental design and data interpretation, supplied financial support and laboratory

space and equipment to complete the experiments, and contributed significantly to the editing of

this manuscript. All authors read and approved the final version of this manuscript.

Acknowledgements

We would like to thank Dr. Ashley Keiser, Dr. Daria Tchessalova, Brynne Raines, and

Sarah Jacob for assistance with the techniques, reagents, and data collection for these

experiments.

Author's Information

CKP is trained in behavioral neuroscience with specific interests in behavioral

neuroimmunology. Since 2013, CKP has worked on models of neuroimmune activation in both

rats and mice with the goal of understanding how the neuroimmune system functions to

modulate cognition and affect and both whether and how biological sex interacts with these

processes.

NCT studies the mechanisms of learning and memory in males and females, the modulation of

memory and affective processes by neuroimmune activation, and long-term impact of illness on

the hippocampus, cognitive decline, Alzheimer's Disease, and neuropsychiatric disorders.

REFERENCES

1.   Marin I, Kipnis J. Learning and memory ... and the immune system. *Learn Mem*.
2013;20(10):601-606; doi:10.1101/lm.028357.112

2.   Rizzo FR, Musella A, De Vito F, et al. Tumor Necrosis Factor and Interleukin-1β
Modulate Synaptic Plasticity during Neuroinflammation. *Neural Plast*.
2018;2018:8430123; doi:10.1155/2018/8430123

3.   Yirmiya R, Goshen I. Immune modulation of learning, memory, neural plasticity and
neurogenesis. *Brain Behav Immun*. 2011;25(2):181-213; doi:10.1016/J.BBI.2010.10.015

4.   Bekhbat M, Neigh GN. Sex differences in the neuro-immune consequences of stress:
Focus on depression and anxiety. *Brain Behav Immun*. 2018;67:1-12;
doi:10.1016/j.bbi.2017.02.006

5.   Dantzer R, O'Connor JC, Freund GG, Johnson RW, Kelley KW. From inflammation to
sickness and depression: when the immune system subjugates the brain. *Nat Rev*
Neurosci. 2008;9(1):46-56; doi:10.1038/nrn2297

6.   Neigh GN, Bekhbat M, Rowson SA. Neuroimmunology: Behavioral Effects. January
2018; doi:10.1093/ACREFORE/9780190264086.013.7

7.   Klein SL, Flanagan KL. Sex differences in immune responses. *Nat Rev Immunol*.
2016;16(10):626-638; doi:10.1038/nri.2016.90

8.   Klein SL, Huber S. Sex differences in susceptibility to viral infection. In: *Sex Hormones*
and Immunity to Infection. Springer-Verlag Berlin Heidelberg; 2010:93-122;
doi:10.1007/978-3-642-02155-8_4

9.   Ransohoff RM. Chemokines and chemokine receptors: standing at the crossroads of
immunobiology and neurobiology. *Immunity*. 2009;31(5):711-721;
doi:10.1016/j.immuni.2009.09.010

10. Ransohoff RM, Brown MA. Innate immunity in the central nervous system. *J Clin Invest*.
2012;122(4):1164-1171; doi:10.1172/JCI58644

741 11. Barrientos RM, O'Reilly RC, Rudy JW. Memory for context is impaired by a post context

742   exposure injection of interleukin-1 beta into dorsal hippocampus. *Behav Brain Res*.

743   2002;134(1-2):291-298; doi:10.1016/S0166-4328(02)00043-8

744 12. Goshen I, Kreisel T, Ounallah-Saad H, et al. A dual role for interleukin-1 in hippocampal-

745   dependent memory processes. *Psychoneuroendocrinology*. 2007;32(8-10):1106-1115;

746   doi:10.1016/J.PSYNEUEN.2007.09.004

747 13. Yirmiya R, Winocur G, Goshen I. Brain interleukin-1 is involved in spatial memory and

748   passive avoidance conditioning. *Neurobiol Learn Mem*. 2002;78(2):379-389;

749   doi:10.1006/nlme.2002.4072

750 14. Sparkman NL, Buchanan JB, Heyen JRR, Chen J, Beverly JL, Johnson RW. Interleukin-6

751   facilitates lipopolysaccharide-induced disruption in working memory and expression of

752   other proinflammatory cytokines in hippocampal neuronal cell layers. *J Neurosci*.

753   2006;26(42):10709-10716; doi:10.1523/JNEUROSCI.3376-06.2006

754 15. Trapero I, Cauli O. Interleukin 6 and cognitive dysfunction. *Metab Brain Dis*.

755   2014;29(3):593-608; doi:10.1007/s11011-014-9551-2

756 16. Weaver JD, Huang MH, Albert M, Harris T, Rowe JW, Seeman TE. Interleukin-6 and risk

757   of cognitive decline: Macarthur studies of successful aging. *Neurology*. 2002;59(3):371-

758   378; doi:10.1212/WNL.59.3.371

759 17. Feuerstein GZ, Liu T, Barone FC. Cytokines, inflammation, and brain injury: Role of

760   tumor necrosis factor-alpha. *Cerebrovascbrain Metabrev*. 1994;6(4):341-360.

761 18. Lindbergh CA, Casaletto KB, Staffaroni AM, et al. Systemic Tumor Necrosis Factor-Alpha

762   Trajectories Relate to Brain Health in Typically Aging Older Adults. *J Gerontol A Biol Sci*

763   *Med Sci*. 2020;75(8):1558-1565; doi:10.1093/gerona/glz209

764 19. Strieter RM, Kunkel SL, Bone RC. Role of tumor necrosis factor-α in disease states and

765   inflammation. *Crit Care Med*. 1993;21(10 SUPPL.); doi:10.1097/00003246-199310001-

766   00006

767   20.   Donzis EJ, Tronson NC. Modulation of learning and memory by cytokines: Signaling

768   mechanisms and long term consequences. *Neurobiol Learn Mem*. 2014;115:68-77;

769   doi:10.1016/J.NLM.2014.08.008

770   21.   Blank T, Detje CN, Spieß A, et al. Brain Endothelial- and Epithelial-Specific Interferon

771   Receptor Chain 1 Drives Virus-Induced Sickness Behavior and Cognitive Impairment.

772   *Immunity*. 2016;44(4):901-912; doi:10.1016/j.immuni.2016.04.005

773   22.   Westin K, Buchhave P, Nielsen H, Minthon L, Janciauskiene S, Hansson O. CCL2 Is

774   Associated with a Faster Rate of Cognitive Decline during Early Stages of Alzheimer's

775   Disease. Ginsberg SD, ed. *PLoS One*. 2012;7(1):e30525;

776   doi:10.1371/journal.pone.0030525

777   23.   Xu J, Dong H, Qian Q, et al. Astrocyte-derived CCL2 participates in surgery-induced

778   cognitive dysfunction and neuroinflammation via evoking microglia activation. *Behav*

779   *Brain Res*. 2017;332:145-153; doi:10.1016/j.bbr.2017.05.066

780   24.   Bajova H, Nelson TE, Gruol DL. Chronic CXCL10 alters the level of activated ERK1/2

781   and transcriptional factors CREB and NF-κB in hippocampal neuronal cell culture. *J*

782   *Neuroimmunol*. 2008;195(1-2):36-46; doi:10.1016/j.jneuroim.2008.01.003

783   25.   Bradburn S, McPhee J, Bagley L, et al. Dysregulation of C-X-C motif ligand 10 during

784   aging and association with cognitive performance. *Neurobiol Aging*. 2018;63:54-64;

785   doi:10.1016/j.neurobiolaging.2017.11.009

786   26.   Loram LC, Sholar PW, Taylor FR, et al. Sex and estradiol influence glial pro-inflammatory

787   responses to lipopolysaccharide in rats. *Psychoneuroendocrinology*. 2012;37:1688-1699;

788   doi:10.1016/j.psyneuen.2012.02.018

789   27.   Santos-Galindo M, Acaz-Fonseca E, Bellini MJ, Garcia-Segura LM. Sex differences in the

790   inflammatory response of primary astrocytes to lipopolysaccharide. *Biol Sex Differ*.

791   2011;2:7; doi:10.1186/2042-6410-2-7

28.   Speirs IC, Tronson NC. Sex differences in hippocampal cytokines after systemic immune

challenge. *bioRxiv*. July 2018:378257; doi:10.1101/378257

29.   Tchessalova D, Tronson NC. Enduring and Sex-specific Changes in Hippocampal Gene

Expression after a Subchronic Immune Challenge. *Neuroscience*. 2020;428:76-89;

doi:10.1016/j.neuroscience.2019.12.019

30.   Bettcher BM, Neuhaus J, Wynn MJ, et al. Increases in a Pro-inflammatory Chemokine,

MCP-1, Are Related to Decreases in Memory Over Time. *Front Aging Neurosci*.

2019;11(FEB):25; doi:10.3389/fnagi.2019.00025

31.   Richwine AF, Sparkman NL, Dilger RN, Buchanan JB, Johnson RW. Cognitive deficits in

interleukin-10-deficient mice after peripheral injection of lipopolysaccharide. *Brain Behav*

Immun. 2009;23(6):794-802; doi:10.1016/j.bbi.2009.02.020

32.   Kronfol Z, Remick DG. Cytokines and the brain: Implications for clinical psychiatry. *Am J*

Psychiatry. 2000;157(5):683-694; doi:10.1176/appi.ajp.157.5.683

33.   Barna M, Komatsu T, Bi Z, Reiss CS. Sex differences in susceptibility to viral infection of

the central nervous system. *J Neuroimmunol*. 1996;67(1):31-39; doi:10.1016/0165-

5728(96)00022-7

34.   Klein SL, Dhakal S, Ursin RL, Deshpande S, Sandberg K, Mauvais-Jarvis F. Biological

sex impacts COVID-19 outcomes. *PLoS Pathog*. 2020;16(6);

doi:10.1371/journal.ppat.1008570

35.   Takahashi T, Ellingson MK, Wong P, et al. Sex differences in immune responses that

underlie COVID-19 disease outcomes. *Nature*. 2020; doi:10.1038/s41586-020-2700-3

36.   Watkins LR, Maier SF, Goehler LE. Cytokine-to-brain communication: A review &

analysis of alternative mechanisms. *Life Sci*. 1995;57(11):1011-1026; doi:10.1016/0024-

3205(95)02047-M

37.   Cunningham C, Campion S, Lunnon K, et al. Systemic Inflammation Induces Acute

Behavioral and Cognitive Changes and Accelerates Neurodegenerative Disease. *Biol*

Psychiatry. 2009;65(4):304-312; doi:10.1016/j.biopsych.2008.07.024

38.   Czerniawski J, Guzowski JF. Acute neuroinflammation impairs context discrimination

820   memory and disrupts pattern separation processes in hippocampus. *J Neurosci*.

821   2014;34(37):12470-12480; doi:10.1523/JNEUROSCI.0542-14.2014

822   39.   McAfoose J, Baune BT. Evidence for a cytokine model of cognitive function. *Neurosci*

823   *Biobehav Rev*. 2009;33(3):355-366; doi:10.1016/j.neubiorev.2008.10.005

824   40.   Vasek MJ, Garber C, Dorsey D, et al. A complement–microglial axis drives synapse loss

825   during virus-induced memory impairment. *Nature*. 2016;534(7608):538-543;

826   doi:10.1038/nature18283

827   41.   Koo JW, Duman RS. Evidence for IL-1 receptor blockade as a therapeutic strategy for the

828   treatment of depression. *Curr Opin Investig Drugs*. 2009;10(7):664-671.

829   42.   Tang M ming, Lin W juan, Pan Y qin, Guan X ting, Li Y cong. Hippocampal neurogenesis

830   dysfunction linked to depressive-like behaviors in a neuroinflammation induced model of

831   depression. *Physiol Behav*. 2016;161:166-173; doi:10.1016/j.physbeh.2016.04.034

832   43.   Zhu X, Levasseur PR, Michaelis KA, Burfeind KG, Marks DL. A distinct brain pathway

833   links viral RNA exposure to sickness behavior. *Sci Rep*. 2016;6(1):29885;

834   doi:10.1038/srep29885

835   44.   Hart BL. The behavior of sick animals. *Vet Clin North Am Small Anim Pract*.

836   1991;21(2):225-237; doi:10.1016/S0195-5616(91)50028-0

837   45.   Silver N, Best S, Jiang J, Thein SL. Selection of housekeeping genes for gene expression

838   studies in human reticulocytes using real-time PCR. *BMC Mol Biol*. 2006;7(1):33;

839   doi:10.1186/1471-2199-7-33

840   46.   Vandesompele J, De Preter K, Pattyn F, et al. Accurate normalization of real-time

841   quantitative RT-PCR data by geometric averaging of multiple internal control genes.

842   *Genome Biol*. 2002;3(7):research0034.1; doi:10.1186/gb-2002-3-7-research0034

843   47.   Perkins JR, Dawes JM, McMahon SB, Bennett DLH, Orengo C, Kohl M. ReadqPCR and

844   NormqPCR: R packages for the reading, quality checking and normalisation of RT-qPCR

quantification cycle (Cq) data. *BMC Genomics*. 2012;13(1):296; doi:10.1186/1471-2164-

13-296

48.   Andersen CL, Jensen JL, Ørntoft TF. Normalization of real-time quantitative reverse

transcription-PCR data: A model-based variance estimation approach to identify genes

suited for normalization, applied to bladder and colon cancer data sets. *Cancer Res*.

2004;64(15):5245-5250; doi:10.1158/0008-5472.CAN-04-0496

49.   Wickham H, Francois R, Henry L, Muller K. dplyr: A Grammar of Data Manipulation.

2020. https://cran.r-project.org/package=dplyr

50.   Wickham H, Henry L. tidyr: Tidy Messy Data. 2020. https://cran.r-

project.org/web/packages/tidyr/

51.   Kassambara A. rstatix: Pipe-Friendly Framework for Basic Statistical Tests. 2020.

https://cran.r-project.org/package=rstatix.

52.   Signorell A, Aho K, Alfons A, et al. DescTools: Tools for descriptive statistics. 2020.

53.   Ludecke D. sjstats: Statistical Functions for Regression Models. 2020. https://cran.r-

project.org/web/packages/sjstats/index.html

54.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. 2016.

https://doi.org/10.18637/jss.v035.b01

55.   Auguie B. gridExtra: Miscellaneous Functions for "Grid" Graphics. 2017. https://CRAN.R-

project.org/package=gridExtra

56.   Kolde R. pheatmap: Pretty Heatmaps. 2019. https://cran.r-

project.org/web/packages/pheatmap

57.   Garnier S. viridis: Default Color Maps from "matplotlib." 2018. https://cran.r-

project.org/package=viridis

58.   Seth RB, Sun L, Chen ZJ. Antiviral innate immunity pathways. In: *Cell Research*. Vol 16.

Nature Publishing Group; 2006:141-147; doi:10.1038/sj.cr.7310019

59.   Stetson DB, Medzhitov R. Type I Interferons in Host Defense. *Immunity*. 2006;25(3):373-

381; doi:10.1016/j.immuni.2006.08.007

60.   Teijaro JR. Type i interferons in viral control and immune regulation. *Curr Opin Virol*.

2016;16:31-40; doi:10.1016/j.coviro.2016.01.001

61.   Cunningham C, Campion S, Teeling J, Felton L, Perry VH. The sickness behaviour and

CNS inflammatory mediator profile induced by systemic challenge of mice with synthetic

double-stranded RNA (poly I:C). *Brain Behav Immun*. 2007;21(4):490-502;

doi:10.1016/J.BBI.2006.12.007

62.   Fortier ME, Kent S, Ashdown H, Poole S, Boksa P, Luheshi GN. The viral mimic,

polyinosinic:polycytidylic acid, induces fever in rats via an interleukin-1-dependent

mechanism. *Am J Physiol - Regul Integr Comp Physiol*. 2004;287(4 56-4):759-766;

doi:10.1152/ajpregu.00293.2004

63.   Kamerman P, Skosana M, Loram L, Mitchell B, Weber J. Fever and inflammatory

cytokine response in rats injected subcutaneously with viral double-stranded RNA analog,

polyinosinic:polycytidylic acid (Poly-I:C). *J Therm Biol*. 2011;36(7):397-402;

doi:10.1016/J.JTHERBIO.2011.07.003

64.   Murray C, Griffin ÉW, O'Loughlin E, et al. Interdependent and independent roles of type I

interferons and IL-6 in innate immune, neuroinflammatory and sickness behaviour

responses to systemic poly I: C. *Brain Behav Immun*. 2015;48:274-286;

doi:10.1016/j.bbi.2015.04.009

65.   Astiz M, Acaz-Fonseca E, Garcia-Segura LM. Sex differences and effects of estrogenic

compounds on the expression of inflammatory molecules by astrocytes exposed to the

insecticide dimethoate. *Neurotox Res*. 2014;25(3):271-285; doi:10.1007/s12640-013-

9417-0

66.   Chistyakov D V., Azbukina N V., Astakhova AA, Goriainov S V., Chistyakov V V.,

Sergeeva MG. Sex-mediated differences in lps induced alterations of TNFα, IL-10

expression, and prostaglandin synthesis in primary astrocytes. *Int J Mol Sci*. 2018;19(9);

897   doi:10.3390/ijms19092793

898 67. Alexopoulou L, Holt AC, Medzhitov R, Flavell RA. Recognition of double-stranded RNA

899   and activation of NF-κB by Toll-like receptor 3. *Nature*. 2001;413(6857):732-738;

900   doi:10.1038/35099560

901 68. Kielian T. Toll-like receptors in central nervous system glial inflammation and

902   homeostasis. *J Neurosci Res*. 2006;83(5):711-730; doi:10.1002/jnr.20767

903 69. Matsumoto M, Seya T. TLR3: Interferon induction by double-stranded RNA including

904   poly(I:C). *Adv Drug Deliv Rev*. 2008;60(7):805-812; doi:10.1016/J.ADDR.2007.11.005

905 70. Cerbai F, Lana D, Nosi D, et al. The neuron-astrocyte-microglia triad in normal brain

906   ageing and in a model of neuroinflammation in the rat hippocampus. *PLoS One*.

907   2012;7(9):e45250; doi:10.1371/journal.pone.0045250

908 71. Scholz J, Woolf CJ. The neuropathic pain triad: neurons, immune cells and glia. *Nat*

909   *Neurosci*. 2007;10(11):1361-1368; doi:10.1038/nn1992

910 72. Bsibsi M, Ravid R, Gveric D, Van Noort JM. Broad expression of Toll-like receptors in the

911   human central nervous system. *J Neuropathol Exp Neurol*. 2002;61(11):1013-1021;

912   doi:10.1093/jnen/61.11.1013

913 73. Norden DM, Trojanowski PJ, Villanueva E, Navarro E, Godbout JP. Sequential activation

914   of microglia and astrocyte cytokine expression precedes increased iba-1 or GFAP

915   immunoreactivity following systemic immune challenge. *Glia*. 2016;64(2):300-316;

916   doi:10.1002/glia.22930

917 74. Channappanavar R, Fett C, Mack M, Ten Eyck PP, Meyerholz DK, Perlman S. Sex-

918   Based Differences in Susceptibility to Severe Acute Respiratory Syndrome Coronavirus

919   Infection. *J Immunol*. 2017;198(10):4046-4053; doi:10.4049/jimmunol.1601896

920 75. Klein SL. Sex influences immune responses to viruses, and efficacy of prophylaxis and

921   treatments for viral diseases. *BioEssays*. 2012;34(12):1050-1059;

922   doi:10.1002/bies.201200099

76.   vom Steeg LG, Klein SL. SeXX Matters in Infectious Disease Pathogenesis. *PLoS*

Pathog. 2016;12(2); doi:10.1371/journal.ppat.1005374

77.   Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of

2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet.*

2020;395(10223):507-513; doi:10.1016/S0140-6736(20)30211-7

78.   Gebhard C, Regitz-Zagrosek V, Neuhauser HK, Morgan R, Klein SL. Impact of sex and

gender on COVID-19 outcomes in Europe. *Biol Sex Differ.* 2020;11(1);

doi:10.1186/s13293-020-00304-9

79.   Guan W, Ni Z, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China.

N Engl J Med. 2020;382(18):1708-1720; doi:10.1056/nejmoa2002032

80.   Klein SL, Dhakal S, Ursin RL, Deshpande S, Sandberg K, Mauvais-Jarvis F. Biological

sex impacts COVID-19 outcomes. Coyne CB, ed. *PLOS Pathog.* 2020;16(6):e1008570;

doi:10.1371/journal.ppat.1008570

81.   Onder G, Rezza G, Brusaferro S. Case-Fatality Rate and Characteristics of Patients

Dying in Relation to COVID-19 in Italy. *JAMA - J Am Med Assoc.* 2020;323(18):1775-

1776; doi:10.1001/jama.2020.4683

82.   Liu B, Han J, Cheng X, et al. Persistent SARS-CoV-2 presence is companied with defects

in adaptive immune system in non-severe COVID-19 patients. *medRxiv.* March

2020:2020.03.26.20044768; doi:10.1101/2020.03.26.20044768

83.   Costello DA, Lynch MA. Toll-like receptor 3 activation modulates hippocampal network

excitability, via glial production of interferon-β. *Hippocampus.* 2013;23(8):696-707;

doi:10.1002/hipo.22129

84.   Scumpia PO, Kelly KM, Reeves WH, Stevens BR. Double-stranded RNA signals antiviral

and inflammatory programs and dysfunctional glutamate transport in TLR3-expressing

astrocytes. *Glia.* 2005;52(2):153-162; doi:10.1002/glia.20234

85.   Katagiri H, Tanaka K, Manabe T. Requirement of appropriate glutamate concentrations in

the synaptic cleft for hippocampal LTP induction. *Eur J Neurosci*. 2001;14(3):547-553;

doi:10.1046/j.0953-816X.2001.01664.x

86.   Krstic D, Knuesel I. Deciphering the mechanism underlying late-onset Alzheimer disease.

Nat Rev Neurol. 2013;9(1):25-34; doi:10.1038/nrneurol.2012.236

87.   Pace TWW, Heim CM. A short review on the psychoneuroimmunology of posttraumatic

stress disorder: From risk factors to medical comorbidities. *Brain Behav Immun*.

2011;25(1):6-13; doi:10.1016/J.BBI.2010.10.003

88.   Wang Z, Young MRI. PTSD, a Disorder with an Immunological Component. *Front

Immunol*. 2016;7:219; doi:10.3389/fimmu.2016.00219

89.   Hodes GE, Kana V, Menard C, Merad M, Russo SJ. Neuroimmune mechanisms of

depression. *Nat Neurosci*. 2015;18(10):1386-1393; doi:10.1038/nn.4113

90.   Zhou H, Lu S, Chen J, et al. The landscape of cognitive function in recovered COVID-19

patients. *J Psychiatr Res*. 2020;129:98-102; doi:10.1016/j.jpsychires.2020.06.022

G.

	Cq Value Standard Deviations	Avg Standard Deviations of ΔCq Values	M Stability Values	Rho Stability Values
18s	1.5402440	1.609	1.609	1.1459049
gapdh	0.5266292	0.911	0.510	0.6235678
hprt1	0.1507111	0.753	0.206	0.3610396
rplp0	0.2245596	0.764	0.206	0.3256827