Bidirectional redox cycling of phenazine-1-carboxylic acid by *Citrobacter portucalensis* MBL

drives increased nitrate reduction

Lev M. Tsypin[a] and Dianne K. Newman[a,b#]

[a] Division of Biology and Biological Engineering, California Institute of Technology, Pasadena,

CA, USA

[b]Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena,

CA, USA

Running Head: *C. portucalensis* MBL links PCA and nitrate redox cycles

Address correspondence to Dianne K. Newman, dkn@caltech.edu

ABSTRACT

Phenazines are secreted metabolites that microbes use in diverse ways, from quorum sensing to

antimicrobial warfare to energy conservation. Phenazines are able to contribute to these activities

due to their redox activity. The physiological consequences of cellular phenazine reduction have

been extensively studied, but the counterpart phenazine oxidation has been largely overlooked.

Phenazine-1-carboxylic acid (PCA) is common in the environment and readily reduced by its

producers. Here, we describe its anaerobic oxidation by *Citrobacter portucalensis* strain MBL,

which was isolated from topsoil in Falmouth, MA, and which does not produce phenazines itself.

This activity depends on the availability of a suitable terminal electron acceptor, specifically

nitrate or fumarate. When *C. portucalensis* MBL is provided reduced PCA and either nitrate or

fumarate, it continuously oxidizes the PCA. We compared this terminal electron acceptor-

dependent PCA-oxidizing activity of *C. portucalensis* MBL to that of several other γ-

proteobacteria with varying capacities to respire nitrate and/or fumarate. We found that PCA

oxidation by these strains in a fumarate- or nitrate-dependent manner is decoupled from growth

and correlated with their possession of the fumarate or periplasmic nitrate reductases,

respectively. We infer that bacterial PCA oxidation is widespread and genetically determined.

Notably, reduced PCA enhances the rate of nitrate reduction to nitrite by *C. portucalensis* MBL

beyond the stoichiometric prediction, which we attribute to *C. portucalensis* MBL's ability to

also reduce oxidized PCA, thereby catalyzing a complete PCA redox cycle. This bidirectionality

highlights the versatility of PCA as a biological redox agent.

IMPORTANCE Phenazines are increasingly appreciated for their roles in structuring microbial

communities. These tricyclic aromatic molecules have been found to regulate gene expression,

be toxic, promote antibiotic tolerance, and promote survival under oxygen starvation. In all of

these contexts, however, phenazines are studied as electron acceptors. Even if their utility arises

primarily from being readily reduced, they would need to be oxidized in order to be recycled.

While oxygen and ferric iron can oxidize phenazines abiotically, biotic oxidation of phenazines

has not been studied previously. We observed bacteria that readily oxidize phenazine-1-

carboxylic acid (PCA) in a nitrate-dependent fashion, concomitantly increasing the rate of nitrate

reduction to nitrite. Because nitrate is a prevalent terminal electron acceptor in diverse anoxic

environments, including soils, and phenazine-producers are widespread, this observation of

linked phenazine and nitrogen redox cycling suggests an underappreciated role for redox-active

secreted metabolites in the environment.

OBSERVATION

Physiological studies of phenazines have focused on cellular reduction of these secreted

molecules for over 120 years. Reduction of phenazines by bacteria was first proposed in the 19th

century as an indicator for the presence of enteric bacteria in water supplies (1). Several decades

later, pyocyanin, one of the phenazines produced by *Pseudomonas aeruginosa*, was described as

an "accessory respiratory pigment" that increased the rate of oxygen consumption by

Staphylococcus, *Pneumococcus*, and erythrocytes by shuttling electrons from the cells to oxygen

(2). Once it became apparent that phenazines can have cytotoxic effects for cells that do not

produce them, they were characterized as antimicrobial compounds that destructively abstract

electrons from the transport chain (3). It was then discovered that phenazine reduction can

greatly benefit *P. aeruginosa* by: 1) regulating gene expression in *P. aeruginosa* during quorum

sensing by oxidizing a transcription factor; 2) acting as alternative terminal electron acceptors to

promote anoxic survival; and 3) facilitating iron acquisition (4–8). These reports paint a complex

picture of the multifarious effects phenazines can have, but in each case the conceptual model

ends with the cell reducing the phenazine. Reduced phenazines can be re-oxidized by inorganic

electron acceptors like oxygen and ferric iron, and this abiotic process has been invoked to

explain redox cycling of phenazines in biofilms (9, 10). However, when these electron acceptors

are unavailable, biotic oxidation of reduced phenazines could close the redox cycle by

regenerating oxidized phenazines. This process has not been shown to exist for secreted redox-

active metabolites.

In parallel to these physiological studies, phenazines have been used as generic electron shuttles

in bioelectrochemical reactor research, selected according to their chemical properties and

suitability for a given application (11, 12). Electrochemically reduced neutral red (NR), a

phenazine, has been successfully used as an electron donor to cells, chosen for its standard

midpoint potential (very near to that of NADH/NAD$^+$, $E^{o\prime}_{\frac{1}{2}} = -320$ mV vs. NHE) and

hydrophobicity (13–15). Anaerobic NR oxidation in BERs has been coupled to the reduction of

several terminal electron acceptors, including nitrate and fumarate (16). A limitation of these

studies is that NR is not found in nature. Therefore, despite NR oxidation being useful in

regulating electrosynthesis, the existence of natural bacterially driven phenazine oxidation

remains unexplored.

PCA is one of the mostly widely synthesized phenazines in the microbial world, from which

other phenazines are derived (17, 18). PCA is known to be reduced by its producers, driving

current generation in bioelectrochemical systems, in which it is re-oxidized by the anode (5, 19).

These facts make PCA a fitting candidate for microbial oxidation during anaerobic metabolism.

In previous work, we enriched for PCA oxidizers from topsoil by incubating them with reduced

PCA, acetate (a non-fermentable carbon source), and nitrate as the only terminal electron

acceptor, and successfully isolated the PCA-oxidizing *Citrobacter portucalensis* MBL (20).

***C. portucalensis* MBL oxidizes PCA in a nitrate- and fumarate-dependent manner.** We did

not observe reduced PCA (PCA$_{red}$/PCA$_{ox}$ $E^{o\prime}_{\frac{1}{2}} = -116$ mV vs. NHE (9)) oxidation in the

absence of a terminal electron acceptor in both the abiotic and biotic regimes (Fig 1A, top left).

When 10 mM fumarate (fumarate/succinate $E^{o\prime}_{\frac{1}{2}} = +33$ mV vs NHE (15)) or nitrate (NO$_3^-$/NO$_2^-$

$E^{o\prime}_{\frac{1}{2}} = +433$ mV vs NHE (15)) were added (right panels), *C. portucalensis* MBL readily

oxidized PCA$_{red}$. Neither fumarate nor nitrate oxidize PCA abiotically (grey curves). During both

dissimilatory and assimilatory nitrate reduction, nitrate is first reduced to nitrite (21). Nitrite

(NO$_2^-$/NO $E^{o\prime}_{\frac{1}{2}} = +350$ mV vs NHE (15)) is a more reactive nitrogen oxide than nitrate.

Accordingly, we observed slow abiotic oxidation of PCA$_{red}$ by nitrite occurs (Fig 1A, bottom left

in grey).

When *C. portucalensis* MBL is incubated with PCA$_{red}$ and nitrate or fumarate, PCA$_{red}$ oxidation

proceeds much faster than it does abiotically by nitrite. Furthermore, when *C. portucalensis*

MBL is incubated with PCA$_{red}$ and nitrite, there is no observable PCA$_{red}$ oxidation. We interpret

this to mean that *C. portucalensis* catalyzes the oxidation of PCA$_{red}$ when an appropriate terminal

electron acceptor is available but reduces oxidized PCA (PCA$_{ox}$) when such an electron acceptor

is absent. We observed PCA$_{ox}$ reduction when the cells started with 200 μM PCA$_{ox}$, but the rate

of reduction decreased according to the provided terminal electron acceptor (Fig 1B). Adding

fumarate (Fig. 1B, top right) decreased the rate of PCA_{ox} reduction by *C. portucalensis* MBL by

2.38 ± 0.12 µM/hr; adding nitrate caused reduction to be non-detectable; and the presence of

nitrite, which abiotically oxidized PCA_{red}, caused a slight dampening in the reduction rate by

1.25 ± 0.09 µM/hr (Fig. 1C and Supp. Fig. 4). The asymptotic steady state that *C. portucalensis*

MBL approaches with fumarate in Fig. 1A and B indicates that the oxidation and reduction of

PCA proceed simultaneously. This is further supported by the fact that PCA_{ox} reduction by *C.*

portucalensis MBL in the presence of fumarate is significantly slower than in the absence of a

terminal electron acceptor (Fig. 1C).

Comparative study of nitrate- and fumarate-dependent PCA oxidation by several γ-

proteobacteria. We assayed whether other γ-proteobacteria can also oxidize PCA in a terminal

electron acceptor-dependent manner (Fig. 1). Using the same assay as above, *E. coli MG1655*, *P.*

aeruginosa UCBPP-PA14 Δ*phz**, *P. chlororaphis phzB::TnluxAB*, and *P. aureofaciens*

phzB::lacZ (all of which cannot synthesize PCA either naturally or due to the specified

mutations) were incubated with PCA and 10 mM nitrate, nitrite, or fumarate. With the exception

of *P. chlororaphis* and *aureofaciens*, all strains exhibited PCA oxidation with nitrate, with *C.*

portucalensis MBL being the fastest with an initial rate of -25.23 ± 0.84 µM PCA_{red}/hr (Fig. 1C).

The assayed *P. aureofaciens* and *chlororaphis* oxidized PCA_{red} with nitrite faster than the abiotic

control: -2.75 ± 0.80 and -3.10 ± 0.58, respectively, vs. -1.48 ± 0.29 µM PCA_{red}/hr abiotically

(Fig. 1C). Only *C. portucalensis* MBL obviously oxidized PCA in the presence of fumarate. *E.*

coli, like *C. portucalensis* MBL, reduces PCA_{ox} more slowly with fumarate than without it,

indicating that it can also oxidize PCA_{red} with fumarate (Fig 1C, light blue), even if not apparent

in Fig. 1A. When the reduction rate by cells was not determined (N.D.), we interpret that the

concurrent oxidation rate overwhelmed the reduction rate. None of the strains exhibited

significant growth in these assays (Supp. Figs. 1 and 2).

The effect of PCA oxidation on the initial rate of nitrate reduction by *C. portucalensis*

MBL. In Fig 1A it is clear that *C. portucalensis* MBL completes its oxidation of ~200 μM

PCA_{red} within 10 hours when nitrate is available. We repeated this experiment in anaerobic

culture tubes inoculated with *C. portucalensis* MBL, 10 mM NO_3^-, and either 200 μM PCA_{red},

200 μM PCA_{ox}, or no PCA and measured nitrate and nitrite concentrations over time via ion

chromatography. We observed that PCA_{red} oxidation significantly increased the rate of nitrate

reduction to nitrite (Fig 2A). Nitrate consumption was stoichiometrically matched by nitrite

production. We did not observe the production of any other nitrogen oxides or ammonium (data

not shown). The no-PCA control did not show any nitrate reduction over the first eight hours (3

$\pm$ 55 μM/hr; 95% confidence interval reported for all rate measurements, calculated from a linear

regression of the data from the first two timepoints for each curve in Fig. 2A). During this time,

131 $\pm$ 49 μM/hr nitrate was reduced in the PCA_{red} condition. In contrast, the PCA_{ox} control

exhibited only a small increase in nitrate reduction (35 $\pm$ 35 μM/hr). To verify that nitrate was

reduced to nitrite, we tracked nitrite's production. We observed that over the first eight hours the

no-PCA control produced nitrite at the rate of 22 $\pm$ 3 μM/hr, the PCA_{ox} control at 58 $\pm$ 2 μM/hr,

and the PCA_{red} condition at 147 $\pm$ 44 μM/hr of nitrite over the first eight hours. Thus, we

estimate the effect of PCA_{red} vs PCA_{ox} to be 96 $\pm$ 60 μM/hr of increased nitrate reduction or 89 $\pm$

44 μM/hr of increased nitrite production. The increase in nitrate reduction due to PCA_{red} was not

stoichiometric: PCA redox and nitrate reduction are both two-electron processes (9, 21), and a

process without a redox cycle would predict that oxidizing PCA_{red} at a rate of 25 µM/hr (Fig. 1C)

would reduce at most 25 µM/hr nitrate to nitrite. However, the lowest range of the confidence

intervals suggests that at least an additional 36 µM/hr nitrate was reduced by the cells when

PCA_{red} was provided, implying that the cells are redox-cycling the PCA.

Conclusions. The effect of PCA oxidation by *C. portucalensis* MBL on its rate of nitrate

reduction was not stoichiometric (Fig 2A). This is consistent with two explanations: 1) a prior

report argues that neutral red (a synthetic phenazine) oxidation affects electrosynthesis during

anaerobic respiration primarily by changing gene regulation via menaquinone reduction (16), and

so it is plausible that PCA oxidation may increase transcription of a rate-limiting factor in the

electron transport chain to nitrate; 2) we observed that a PCA redox cycle by *C. portucalensis*

MBL is possible (Fig 1, most evident in the case of fumarate, top right panels in A and B, where

oxidation and reduction proceed simultaneously in the steady state); it is likely that PCA redox

cycling may also directly stimulate nitrate reduction. We cannot determine whether reduced PCA

(PCA_{red}) serves as an effective electron donor to the cells' metabolism from our above

observations. Regardless, nitrate reduction driven by PCA redox-cycling has environmental

ramifications far beyond the cell.

Despite all being capable of nitrate reduction, not all the γ-proteobacteria that we tested were

capable of oxidizing PCA_{red} with nitrate (Fig. 1C), suggesting a genetic basis for this process.

The only strains that did not readily oxidize PCA_{red} with nitrate were *P. chlororaphis* and

aureofaciens, and, unlike the other species, they do not possess the periplasmic nitrate reductase

operon *nap*. Curiously, it has been previously noted that wildtype *P. aeruginosa* UCBPP-PA14

oxidizes another phenazine, pyocyanin, in the presence of nitrate, while a *napA* transposon

knockout strain does not (22). We propose that this operon may be necessary for rapid PCA_{red}

oxidation and are developing genetic tools in *C. portucalensis* MBL to test this idea directly.

Nitrate-dependent PCA_{red} oxidation is likely to be common in anoxic environments, such as

soils, as it is not species-specific and occurs readily. If the outsized effect of PCA_{red} oxidation on

nitrate reduction that we observe generalizes to other species, the production and reduction of

phenazines by organisms like *Pseudomonads* likely affects the rate of nitrate consumption in

their environs, adding another function to the phenazine arsenal (18). We propose that cells may

catalyze a PCA redox cycle whenever they have internal stores of reducing equivalents and a

usable terminal electron acceptor (Fig 2B). While both intracellular (e.g., sulfur) and

extracellular (e.g., humics) bacterial redox cycles have been described (23, 24), to our knowledge

this has not been appreciated for secreted metabolites, such as phenazines. Our observation

implies that these molecules may act as "electron buffers", enabling cells to reduce and oxidize

them according to whether they are lacking a terminal electron acceptor or an electron donor,

respectively, and in so doing significantly impact other biogeochemical cycles.

MATERIALS AND METHODS

Strains and media. *Citrobacter portucalensis* MBL was isolated in our previous work (20). In

the comparative PCA oxidation and reduction experiments, we used strains of γ-proteobacteria

that cannot synthesize phenazines, either natively or due to mutations. We used *E. coli* MG1655,

P. aeruginosa UCBPP-PA14 Δ*phz**, *P. chlororaphis phzB::TnLuxAB*, and *P. aureofaciens*

phzB::lacZ. The wildtype *Pseudomonads* can synthesize PCA, but the used mutants cannot. All

strains were grown and incubated under the same conditions. The basal medium for the

experiments contained 20 mM potassium phosphate buffer (final pH 7), 1 mM sodium sulfate, 10

mM ammonium chloride, 1x SL-10 trace elements, 1x freshwater salt solution (17.1 mM sodium

chloride, 1.97 mM magnesium chloride, 0.68 mM calcium chloride, and 6.71 mM potassium

chloride), and 1x 13-vitamin solution (10 μM MOPS pH 7.2, 0.1 μg/mL riboflavin, 0.03 μg/mL

biotin, 0.1 μg/mL thiamine HCl, 0.1 μg/mL L-ascorbic acid, 0.1 μg/mL d-Ca-pantothenate, 0.1

μg/mL folic acid, 0.1 μg/mL nicotinic acid, 0.1 μg/mL 4-aminobenzoic acid, 0.1 μg/mL

pyridoxine HCl, 0.1 μg/mL lipoic acid, 0.1 μg/mL NAD, 0.1 μg/mL thiamine pyrophosphate,

and 0.01 μg/mL cyanocobalamin). Depending on the experimental condition, as indicated in the

figure legends, a terminal electron acceptor would be added (10 mM of fumarate, nitrate, or

nitrite) or omitted. For oxidized PCA, a 10 mM stock in 20 mM NaOH was prepared. For

reduced PCA, an 800 μM stock in the basal medium was reduced by electrolysis. Both stocks

were diluted into plates to a final target concentration of 200 μM.

Cell preparation. All cell incubations and experiments were performed at 30 C. Cells were

preserved in 35% glycerol stocks at -80 C. Two days prior to the experiments, frozen cells for

each strain assayed were struck out on lysogeny broth (LB) agar plates and incubated overnight.

The evening prior to the experiment, a patch from the streaks was inoculated into liquid LB in a

respective culture tube and incubated slanted, shaking at 250 rpm, overnight. The morning of the

experiment, 1 mL of each cell culture was washed three times into the basal medium by spinning

for two minutes at 6000 x g, aspirating the supernatant, and gently resuspending with a pipette.

The OD_{600} of each washed culture was measured. The cultures were brought into a Coy glove

box, where they were washed three times into the same basal medium that had been made

anoxic, following the same procedure as above. After being left to stand for 1-2 hours, the cells

were inoculated into the different experimental conditions at a target starting OD_{600} of 0.1.

Measurement of PCA redox and nitrogen oxide concentrations. All PCA redox

measurements were performed in a Coy chamber using a BioTek Synergy 5 plate reader.

Reduced PCA concentration was measured by fluorescence (excitation 360 nm and emission 528

nm) (25). In these cultures, 1 mM acetate was provided. Plates were incubated shaking on the

"medium" setting. For nitrogen oxide concentration measurements, *C. portucalensis* MBL cells

were prepared as above, but incubated in culture tubes in the Coy chamber to allow for sampling.

In these cultures, no acetate was provided. The tubes were kept at 30 C, but not shaking. Samples

were filtered through a 0.2 µm cellulose-acetate spin filter and stored at -80 C prior to analysis.

Nitrate and nitrite concentrations were measured by ion chromatography using a Dionex ICS-

2000 instrument.

Data analysis. Initial redox rates in the conditions with supplied nitrite did not include the first

1.5 hours of the assay because that period showed a systematic artifact due to fluorescence

quenching (Fig. 1A, lower left panel). The linear fits were calculated over the first five hours

after the first detectable PCA_{red} measurement, which was determined to be appropriate based on

scanning for R^2 values over increasing time frames (Supp. Figs. 4 and 5). 95% confidence

intervals are calculated as the estimated value ± 1.96 * (standard error). When comparing rates,

the reported error is the geometric mean of the intervals for the two measurements. All plots

were generated using Bokeh and the legends and titles were adjusted using Inkscape. All the raw

data and the Jupyter notebook used for their analysis are available at

https://github.com/ltsypin/Cportucalensis_observation.

ACKNOWLEDGEMENTS

We would like to thank the members of the Newman lab, and especially Scott Saunders, Darcy

McRose, Avi Flamholz, John Ciemniecki, Chelsey VanDrisse, and Justin Bois for their insight

and helpful discussions throughout this work. We are grateful to Nathan Dalleska at the

Environmental Analysis Center at Caltech for training LMT on the Dionex instrument and

providing a facility for analytical chemistry. LMT was supported by the Rosen Endowment

Fellowship at Caltech and the National Science Foundation Graduate Research Fellowship

(DGE-1745301). Additional support to DKN came from NIH (1R01AI127850-01A1 and

1R01HL152190-01) and ARO (W911NF-17-1-0024) grants.

REFERENCES

1. Rothberger CJ. 1898. Differential diagnostische Untersuchungen mit gefarbten Nahrboden. Centralblatt für Bakteriologie, Parasitenkunde und Infektionskrankheiten 24:513–518.

2. Friedheim EA. 1931. Pyocyanine, an accessory respiratory enzyme. J Exp Med 54:207–221.

3. Baron SS, Terranova G, Rowe JJ. 1989. Molecular mechanism of the antimicrobial action of pyocyanin. Curr Microbiol 18:223–230.

4. Dietrich LEP, Price-Whelan A, Petersen A, Whiteley M, Newman DK. 2006. The phenazine pyocyanin is a terminal signalling factor in the quorum sensing network of *Pseudomonas aeruginosa*. Mol Microbiol 61:1308–1321.

5. Glasser NR, Kern SE, Newman DK. 2014. Phenazine redox cycling enhances anaerobic survival in *Pseudomonas aeruginosa* by facilitating generation of ATP and a proton-motive force. Mol Microbiol 92:399–412.

6. Gu M, Imlay JA. 2011. The SoxRS response of *Escherichia coli* is directly activated by redox-cycling drugs rather than by superoxide. Mol Microbiol 79:1136–1150.

7. Dietrich LEP, Kiley PJ. 2011. A shared mechanism of SoxR activation by redox-cycling compounds. Mol Microbiol 79:1119–1122.

8. Wang Y, Wilks JC, Danhorn T, Ramos I, Croal L, Newman DK. 2011. Phenazine-1-carboxylic acid promotes bacterial biofilm development via ferrous iron acquisition. J Bacteriol 193:3606–3617.

9. Wang Y, Newman DK. 2008. Redox reactions of phenazine antibiotics with ferric (hydr)oxides and molecular oxygen. Environ Sci Technol 42:2380–2386.

10. Saunders SH, Tse ECM, Yates MD, Otero FJ, Trammell SA, Stemp EDA, Barton JK,

Tender LM, Newman DK. 2020. Extracellular DNA promotes efficient extracellular

electron transfer by pyocyanin in *Pseudomonas aeruginosa* biofilms. Cell 182:919–

932.e19.

11. Rabaey K, Boon N, Höfte M, Verstraete W. 2005. Microbial phenazine production

enhances electron transfer in biofuel cells. Environ Sci Technol 39:3401–3408.

12. Pham TH, Boon N, Aelterman P, Clauwaert P, De Schamphelaire L, Vanhaecke L, De

Maeyer K, Höfte M, Verstraete W, Rabaey K. 2008. Metabolites produced by

Pseudomonas sp. enable a Gram-positive bacterium to achieve extracellular electron

transfer. Appl Microbiol Biotechnol 77:1119–1129.

13. Park DH, Zeikus JG. 1999. Utilization of electrically reduced neutral red by *Actinobacillus*

succinogenes: physiological function of neutral red in membrane-driven fumarate

reduction and energy conservation. J Bacteriol 181:2403–2410.

14. Hongo M, Iwahara M. 1979. Application of electro-energizing method to L-glutamic acid

fermentation. Agric Biol Chem 43:2075–2081.

15. Thauer RK, Jungermann K, Decker K. 1977. Energy conservation in chemotrophic

anaerobic bacteria. Bacteriol Rev 41:100–180.

16. Harrington TD, Tran VN, Mohamed A, Renslow R, Biria S, Orfe L, Call DR, Beyenal H.

2015. The mechanism of neutral red-mediated microbial electrosynthesis in *Escherichia*

coli: menaquinone reduction. Bioresour Technol 192:689–695.

17. Blankenfeldt W, Parsons JF. 2014. The structural biology of phenazine biosynthesis. Curr

Opin Struct Biol 29:26–33.

18. Dar D, Thomashow LS, Weller DM, Newman DK. 2020. Global landscape of phenazine

biosynthesis and biodegradation reveals species-specific colonization patterns in

301   agricultural soils and crop microbiomes. eLife 9.

302 19. Feng J, Qian Y, Wang Z, Wang X, Xu S, Chen K, Ouyang P. 2018. Enhancing the

303   performance of *Escherichia coli*-inoculated microbial fuel cells by introduction of the

304   phenazine-1-carboxylic acid pathway. J Biotechnol 275:1–6.

305 20. Tsypin LM, Saunders SH, Bar-On Y, Leadbetter JR, Newman DK. 2020. Draft genome

306   sequence of the redox-active enteric bacterium *Citrobacter portucalensis* strain MBL.

307   Microbiol Resour Announc 9.

308 21. Sparacino-Watkins C, Stolz JF, Basu P. 2014. Nitrate and periplasmic nitrate reductases.

309   Chem Soc Rev 43:676–706.

310 22. Price-Whelan AM. 2009. PhD Thesis: Physiology and mechanisms of pyocyanin reduction

311   in *Pseudomonas aeruginosa*. California Institute of Technology.

312 23. Schmidt TM, Arieli B, Cohen Y, Padan E, Strohl WR. 1987. Sulfur metabolism in

313   *Beggiatoa alba*. J Bacteriol 169:5466–5472.

314 24. Coates JD, Cole KA, Chakraborty R, O'Connor SM, Achenbach LA. 2002. Diversity and

315   ubiquity of bacteria capable of utilizing humic substances as electron donors for anaerobic

316   respiration. Appl Environ Microbiol 68:2445–2452.

317 25. Sullivan NL, Tzeranis DS, Wang Y, So PTC, Newman D. 2011. Quantifying the dynamics

318   of bacterial secondary metabolites by spectral multiphoton microscopy. ACS Chem Biol

319   6:893–899.

Main text figure legends

Figure 1. Oxidation and reduction of PCA by bacteria provided different terminal electron acceptors (TEAs). (A-B) The light circles correspond to three independent biological replicates and the dark lines to their respective means. Cells only oxidize PCA when an appropriate TEA is available. Fumarate, nitrite, or nitrate (top right, bottom left, and bottom right panels, respectively) stimulate different strains to oxidize PCA. When no TEA is provided (top left panels), no strains oxidize PCA and all reduce it at different rates (enumerated in C). With nitrate, only *P. chlororaphis* and *aureofaciens* (blue and umber curves) appear to not oxidize PCA but reduce it. Nitrite abiotically oxidizes PCA (grey curve, lower left in A), but *P. aureofaciens and chlororaphis* catalyze an even faster biological oxidation. In contrast, the *Enterics* (yellow and orange curves) reduce PCA faster than the abiotic reaction. Given fumarate, *C. portucalensis* MBL (orange) clearly oxidizes PCA, and the *Pseudomonads* reduce it. (C) This table reports the estimated initial rates of oxidation or reduction according to a linear fit over the first five hours (Supp. Figs. 3 and 4). This timeframe was determined by tracking the R^2 for the linear fit over increasing time windows (Supp. Figs. 5 and 6). N.D. signifies rates that were not detectable. We can only observe the relative magnitude of the rates of oxidation and reduction in a given condition. However, whether the oxidation of PCA overpowers its reduction depends on both the provided TEA and is associated with the presence of the periplasmic nitrate reductase (NAP) or fumarate reductase (FRD). PCA reduction is calculated as a positive rate (shades of blue); PCA oxidation is calculated as a negative rate (shades of red).

Figure 2. PCA oxidation by *C. portucalensis* MBL increases its initial rate of nitrate reduction. (A) Either 200 μM reduced PCA (PCA$_{red}$), 200 μM oxidized PCA (PCA$_{ox}$), or no

PCA was added to each condition. Ion chromatography shows that over the 10 hours that the

cells are oxidizing PCA_{red} (Fig. 1A), their initial rate of nitrate reduction is substantially

increased. The nitrate is stoichiometrically reduced to nitrite. Error bars are 95% confidence

intervals around the mean of three independent biological replicates. When not visible, they are

smaller than the circles (nitrate) or squares (nitrite) denoting the measurements. (B) Any cell that

has internal stores of reducing equivalents and an appropriate terminal electron acceptor (TEA_1)

may catalyze an internal PCA redox cycle (bolded arrows). We observe this for *E. coli* and *C.*

portucalensis MBL with nitrate or fumarate, *P. aeruginosa* with nitrate, and *P. aureofaciens* and

chlororaphis with nitrite. In addition to this cellularly-catalyzed reaction, the product of TEA_1's

reduction (TEA_2) may abiotically oxidize PCA, which we observe when nitrate is reduced to

nitrite (Fig. 1A, bottom left).

356
357

Supplementary Figure Legends

Supplementary Figure 1. OD_{600} measurements for all strains in all conditions during the oxidation assays (Fig. 1A). Light circles represent all measurements from three independent biological replicates corresponding to the PCA_{red} measurements in Figure 1A. The solid lines are the means of the replicates. The curves are colored by condition (i.e., which, if any, terminal electron acceptor was added). While there are some OD_{600} trends in the first five hours, there is no evidence of growth that would indicate a doubling. In the abiotic graph, all the data points overlie each other.

Supplementary Figure 2. OD_{600} measurements for all strains in all conditions during the reduction assays (Fig. 1B). Light circles represent all measurements from three independent biological replicates corresponding to the PCA_{red} measurements in Figure 1B. The solid lines are the means of the replicates. The curves are colored by condition (i.e., which, if any, terminal electron acceptor was added). While there are some OD_{600} trends in the first five hours, there is no evidence of growth that would indicate a doubling. In the abiotic graph, all the data points overlie each other.

Supplementary Figure 3. Oxidation assay results grouped by strain rather than by condition. This is a different representation of the same data as in Fig. 1A, including the linear regressions (dashed lines) that generated the initial rate estimates as reported in Fig. 1Cα.

Supplementary Figure 4. Reduction assay results grouped by strain rather than by condition. This is a different representation of the same data as in Fig. 1B, including the linear regressions

(dashed lines) that generated the initial rate estimates as reported in Fig. 1Cβ. In the abiotic

conditions, no reduction was detected (N.D.).

Supplementary Figure 5. Evaluation of the appropriate window for performing the linear

regressions to estimate initial redox rates in the oxidation assays (Fig. 1Cα). Starting from the

first valid PCA$_{red}$ measurement in each condition, a linear regression was performed over

increasing lengths of time and the corresponding R^2 value was plotted. A time window of five

hours (red line) was selected as a systematic compromise for all conditions.

Supplementary Figure 6. Evaluation of the appropriate window for performing the linear

regressions to estimate initial redox rates in the reduction assays (Fig. 1Cβ). Starting from the

first valid PCA$_{red}$ measurement in each condition, a linear regression was performed over

increasing lengths of time and the corresponding R^2 value was plotted. A time window of five

hours (red line) was selected as a systematic compromise for all conditions.

Figure 1. Oxidation and reduction of PCA by bacteria provided different terminal electron acceptors (TEAs). (A-B) The light circles correspond to three independent biological replicates and the dark lines to their respective means. Cells only oxidize PCA when an appropriate TEA is available. Fumarate, nitrite, or nitrate (top right, bottom left, and bottom right panels, respectively) stimulate different strains to oxidize PCA. When no TEA is provided (top left panels), no strains oxidize PCA and all reduce it at different rates (enumerated in C). With nitrate, only P. chlororaphis and aureofaciens (blue and umber curves) appear to not oxidize PCA but reduce it. Nitrite abiotically oxidizes PCA (grey curve, lower left in A), but P. aureofaciens and chlororaphis catalyze an even faster biological oxidation. In contrast, the Enterics (yellow and orange curves) reduce PCA faster than the abiotic reaction. Given fumarate, C. portucalensis MBL (orange) clearly oxidizes PCA, and the Pseudomonads reduce it. (C) This table reports the estimated initial rates of oxidation or reduction according to a linear fit over the first five hours (Supp. Figs. 3 and 4). This timeframe was determined by tracking the R^2 for the linear fit over increasing time windows (Supp. Figs. 5 and 6). N.D. signifies rates that were not detectable. We can only observe the relative magnitude of the rates of oxidation and reduction in a given condition. However, whether the oxidation of PCA overpowers its reduction depends on both the provided TEA and is associated with the presence of the periplasmic nitrate reductase (NAP) or fumarate reductase (FRD). PCA reduction is calculated as a positive rate (shades of blue); PCA oxidation is calculated as a negative rate (shades of red).

Figure 2. PCA oxidation by C. portucalensis MBL increases its initial rate of nitrate reduction. (A) Either 200 µM reduced PCA (PCA$_{red}$), 200 µM oxidized PCA (PCA$_{ox}$), or no PCA was added to each condition. Ion chromatography shows that over the 10 hours that the cells are oxidizing PCAred (Fig. 1A), their initial rate of nitrate reduction is substantially increased. The nitrate is stoichiometrically reduced to nitrite. Error bars are 95% confidence intervals around the mean of three independent biological replicates. When not visible, they are smaller than the circles (nitrate) or squares (nitrite) denoting the measurements. (B) Any cell that has internal stores of reducing equivalents and an appropriate terminal electron acceptor (TEA$_1$) may catalyze an internal PCA redox cycle (bolded arrows). We observe this for E. coli and C. portucalensis MBL with nitrate or fumarate, P. aeruginosa with nitrate, and P. aureofaciens and chlororaphis with nitrite. In addition to this cellularly-catalyzed reaction, the product of TEA$_1$'s reduction (TEA$_2$) may abiotically oxidize PCA, which we observe when nitrate is reduced to nitrite (Fig. 1A, bottom left).