Ipomoeassin-F inhibits the *in vitro* biogenesis of the SARS-CoV-2

spike protein and its host cell membrane receptor

Sarah O'Keefe[1,4], Peristera Roboti[1], Kwabena B. Duah[2], Guanghui Zong[3], Hayden

Schneider[2], Wei Q. Shi[2] and Stephen High[1,4]

[1]School of Biological Sciences, Faculty of Biology, Medicine and Health, University of

Manchester, Manchester, M13 9PT, United Kingdom

[2]Department of Chemistry, Ball State University, Muncie, Indiana 47306, USA

[3]Department of Chemistry and Biochemistry, University of Maryland, College Park,

Maryland 20742, USA

[4]Lead   Contacts   for   correspondence: sarah.okeefe@manchester.ac.uk;

stephen.high@manchester.ac.uk

Keywords

Cell-free translation, Endoplasmic reticulum (ER), ER membrane complex (EMC),

Sec61 translocon, viral protein biogenesis.

Abstract

In order to produce proteins essential for their propagation, many pathogenic human viruses, including SARS-CoV-2 the causative agent of COVID-19 respiratory disease, commandeer host biosynthetic machineries and mechanisms. Three major structural proteins, the spike, envelope and membrane proteins, are amongst several SARS-CoV-2 components synthesised at the endoplasmic reticulum (ER) of infected human cells prior to the assembly of new viral particles. Hence, the inhibition of membrane protein synthesis at the ER is an attractive strategy for reducing the pathogenicity of SARS-CoV-2 and other obligate viral pathogens. Using an *in vitro* system, we demonstrate that the small molecule inhibitor ipomoeassin F (Ipom-F) potently blocks the Sec61-mediated ER membrane translocation/insertion of three therapeutic protein targets for SARS-CoV-2 infection; the viral spike and ORF8 proteins together with angiotensin-converting enzyme 2, the host cell plasma membrane receptor. Our findings highlight the potential for using ER protein translocation inhibitors such as Ipom-F as host-targeting, broad-spectrum, antiviral agents.

Introduction

Viruses are obligate parasites that require host cell machineries to support their cellular invasion, genomic replication and the synthesis of viral progeny. Inhibition of the host factors that are co-opted during viral infection thus provides a powerful strategy for the development of broad-spectrum antiviral agents; a current global need given the paucity of clinically effective antiviral drugs and the emergence of highly pathogenic human viruses (Adalja and Inglesby, 2019).

Many of these viruses, including the causative agent of COVID-19 respiratory disease, SARS-CoV-2 (Zhou et al., 2020; Zhu et al., 2020) (Fig. 1A), hijack the mechanisms of host secretory pathway-mediated protein synthesis in order to correctly synthesise, fold and assemble important viral proteins (Bojkova et al., 2020; Gordon et al., 2020; Sicari et al., 2020). Hence, small molecule inhibitors of Sec61-mediated co-translational protein entry into the endoplasmic reticulum (ER) (Luesch and Paavilainen, 2020) have recently attracted attention as potentially broad-spectrum antiviral agents (Heaton et al., 2016; Shah et al., 2018). Such inhibitors offer a dual approach to reduce viral pathogenicity; first, by directly inhibiting the production of key viral proteins and, second, by reducing the levels of host proteins that are co-opted during viral infection. In the latter case, human angiotensin-converting enzyme 2 (ACE2) is one of the host cell receptors crucial for SARS-CoV-2 viral entry and infectivity (Cantuti-Castelvetri et al., 2020; Daly et al., 2020; Walls et al, 2020). As an integral membrane protein, ACE2 is synthesised at the ER prior to its trafficking to the plasma membrane (Warner et al., 2005), and is thus a potential candidate for down-regulation using inhibitors of ER entry.

Our recent studies show that ipomoeassin-F (Ipom-F) (Fig. 1B) is a potent and selective inhibitor of Sec61-mediated protein translocation into and across the ER membrane (Zong et al., 2019; O'Keefe et al., 2020 submitted). Given that SARS-CoV-2 membrane proteins likely co-opt host mechanisms of ER entry during their biogenesis (cf. Gordon et al., 2020; Sicari et al., 2020), we concluded that their sensitivity to Ipom-F would likely be comparable to that of endogenous Sec61 clients (Fig. 1C; see also Zong et al., 2019; O'Keefe et al., 2020 submitted). We, therefore, evaluated the effects of Ipom-F on a small panel of SARS-CoV-2 proteins that each contain hydrophobic ER targeting signals (Fig. 1D). The *in vitro* membrane insertion

of the viral spike (S) protein and membrane translocation of the ORF8 protein are both

strongly inhibited by Ipom-F, whilst several other viral membrane proteins are

unaffected (Fig. 2). Likewise, the ER membrane integration of ACE2, an important

host receptor for SARS-CoV-2 (Walls et al., 2020), is highly sensitive to Ipom-F (Fig.

2).

We show that the principle molecular basis for the Ipom-F sensitivity of these SARS-

CoV-2 proteins is driven by their dependence on the Sec61 translocation complex, as

dictated by their individual structural features and membrane topologies (Fig. 3).

Taken together, our *in vitro* study of SARS-CoV-2 protein synthesis at the ER

highlights Ipom-F as a promising candidate for the development of a broad-spectrum,

host-targeting, antiviral agent.

Fig. 1. Ipom-F as a potential inhibitor of SARS-CoV-2 viral protein synthesis. (A) A
schematic of the (+) ssRNA genome architecture of SARS-CoV-2 (29903 nt) which contains
a 5' capped mRNA with a leader sequence (LS), a 3' end poly-A tail, a 5' and 3' UTR and the
open reading frames (ORFs) ORF1a, ORF1b, spike (S), ORF3a, envelope (E), membrane
(M), ORF6, ORF7, ORF8, nucleoprotein (N) and ORF10 (Firth, 2020; Naqvi et al., 2020). An
important mode of SARS-CoV-2 host entry proceeds via interaction of the viral S protein with
human angiotensin-converting enzyme 2 (ACE2) (Walls et al., 2020). **(B)** The chemical
structure of Ipomoeassin-F (Ipom-F), a small molecule inhibitor of Sec61-mediated protein
translocation into the endoplasmic reticulum (ER). **(C)** Ipom-F efficiently blocks the ER
membrane translocation of secretory proteins and the ER insertion of single-pass type I and

type II TMPs, but not the insertion of type III TMPs or tail-anchored (TA) proteins. In the
schematic, SA denotes a signal anchor. (**D**) Based on the known/predicted membrane
topology of candidate SARS-CoV-2 proteins and the sensitivity of comparable host cell
proteins (Zong et al., 2019; O'Keefe et al., 2020 submitted), we anticipated their likely
sensitivity to an Ipom-F-induced blockade of ER protein translocation.

Results and Discussion
<u>Ipom-F selectively inhibits ER translocation of the viral ORF8 and S proteins</u>
In order to explore the ability of Ipom-F to inhibit the ER translocation of a small, yet

structurally diverse, panel of SARS-CoV-2 membrane and secretory-like proteins, we

first used a well-established *in vitro* translation system supplemented with canine

pancreatic microsomes (Fig. 2A). To facilitate the detection of ER translocation, we

modified the viral ORF8, S, E, M and ORF6 proteins by adding an OPG2-tag; an

epitope that supports efficient ER lumenal N-glycosylation and enables product

recovery via immunoprecipitation, without affecting Ipom-F sensitivity of the substrate

(Fig. S1A) (O'Keefe et al., 2020 submitted). The identity of the resulting N-glycosylated

species for each of these OPG2-tagged viral proteins was confirmed by

endoglycosidase H (Endo H) treatment of the radiolabelled products associated with

the membrane fraction prior to SDS-PAGE (Fig. 2B, cf. lanes 1 and 2 in each panel).

Using ER lumenal modification of either endogenous N-glycosylation sites (viral S and

M proteins) or the appended OPG2-tag (viral E and ORF8 proteins) as a robust

reporter for ER membrane translocation, we found that 1 µM Ipom-F strongly inhibited

both the translocation of the soluble, secretory-like protein ORF8-OPG2 and the

integration of the type I transmembrane proteins (TMP) S-OPG2, and various

truncated derivatives thereof (Fig. 2B, Fig. 2C, Fig. S1C). Furthermore, membrane

insertion of the human type I TMP, ACE2, was inhibited to a similar extent (Fig. 2B,

Fig. 2C, ~70 to ~90% inhibition for these three proteins). These results mirror previous

findings showing that precursor proteins that bear N-terminal signal peptides, and are

thus obligate clients for the Sec61-translocon, are typically very sensitive to Ipom-F-

mediated inhibition (Zong et al., 2019; O'Keefe et al., 2020 submitted). In the context

of SARS-CoV-2 infection, wherein ACE2 acts as an important host cell receptor for

the SARS-CoV-2 virus via its interaction with the viral S protein (Walls et al., 2020),

these data suggest that an Ipom-F-induced antiviral effect might be achieved via

selective reductions in the biogenesis of both host and viral proteins alike (cf. Fig. 1A).

In contrast to the viral S and ORF8 proteins, insertion of the viral E protein was
unaffected by Ipom-F (Fig. 2B-C), consistent with its recently reported classification as
a type III TMP (Duart et al., 2020). Hence, the integration of type III TMPs is highly
resistant to Ipom-F (Zong et al., 2019), most likely because they exploit a novel
pathway for ER membrane insertion (cf. Fig. 3; O'Keefe et al., 2020 submitted). We
therefore conclude that the known substrate-selective inhibitory action of Ipom-F at
the Sec61 translocon is directly applicable to viral membrane proteins; whereby the
ER translocation of secretory proteins and type I TMPs, but not type III TMPs, is
efficiently blocked by Ipom-F. It seems likely that viral type II TMPs will also be inhibited
by Ipom-F (cf. Morel et al., 2018; Zong et al., 2019; Gérard et al., 2020) but, since
SARS-CoV-2 has no such proteins, we have not addressed this issue.

The viral M protein is a multi-pass TMP with its first TMD oriented so the N-terminus
is exoplasmic (N_{exo}) and hence can be considered "type III-like". Although human
multi-pass TMPs of this type typically require both the ER membrane complex (EMC)
and Sec61 translocon for their authentic ER insertion (Chitwood et al., 2018), Ipom-F
had no significant effect on the ER translocation of the N-terminal domain of the M
protein *in vitro* (Fig. 2C). We conclude that the integration of its first TMD is unaffected
by Ipom-F, consistent with its use of the EMC (Chitwood et al., 2018; O'Keefe et al.,
2020 submitted). Although we cannot rule out any Ipom-F-induced defects in the
Sec61-dependent integration of the second/third TM-spans of the M protein (cf.
Chitwood et al., 2018), previous studies have shown that similar host cell multi-pass
TMPs are resistant to comparable Sec61 inhibitors such as mycolactone (Morel et al.,
2018). In practice, the potential resistance of this highly abundant and functionally
diverse class of endogenous membrane proteins (von Heijne, 2007) may limit any
Ipom-F-induced cytotoxicity towards host cells.

Fig. 2. Ipom-F selectively inhibits the ER membrane translocation of SARS-CoV-2 proteins. (A) Schematic of the *in vitro* ER import assay in which canine pancreatic microsomes are used as a source of ER membrane. Following translation, fully translocated or membrane inserted radiolabelled precursor proteins are recovered with the ER membrane fraction and analysed by SDS-PAGE and phosphorimaging. The identity of N-glycosylated species was confirmed by treatment with endoglycosidase H (Endo H). **(B)** Protein precursors of the human angiotensin-converting enzyme 2 (ACE2) and OPG2-tagged versions of the SARS-CoV-2 ORF8 (ORF8-OPG2), spike (S-OPG2), envelope (OPG2-E), membrane (M-OPG2) and ORF6 (a doubly-OPG2 tagged version, OPG2-ORF6-OPG2, and two singly-OPG2 tagged forms, OPG2-ORF6 and ORF6-OPG2, with their respective predominant N-glycan species highlighted in bold) were synthesised in rabbit reticulocyte lysate

supplemented with ER-derived microsomes in the absence or presence of Ipom-F (lanes 1 and 3). Phosphorimages of membrane-associated products resolved by SDS-PAGE, together with representative substrate outlines, are shown. N-glycosylation was used as a read-out for the efficiency of membrane translocation/insertion and N-glycosylated (X-Gly) versus non-N-glycosylated (0Gly) species were identified by treatment with Endo H (see lane 2). **(C)** The relative efficiency of membrane translocation/insertion in the presence of Ipom-F was calculated using the ratio of N-glycosylated protein to non-glycosylated protein, relative to the DMSO treated control (set to 100% efficiency). Quantifications are given as means ±SEM for independent translation reactions performed in triplicate (*n*=3) and statistical significance (one-way ANOVA, DF and F values shown in the figure) was determined using Dunnett's multiple comparisons test. Statistical significance is given as n.s., non-significant >0.1; ****, P < 0.0001.

ORF6 assumes a lumenal-facing hairpin topology in ER-derived microsomes

Cell-based studies of the ORF6 protein from SARS-CoV-1 suggest it has an unusual hairpin topology with both its N- and C-termini located on the exoplasmic side of the host cell membrane, to which it binds via an N-terminal amphipathic helix (Netland et al., 2007). To independently determine the membrane topology of SARS-CoV-2 ORF6, we prepared versions with OPG2 tags at both its N- and C-termini, or single tagged equivalents (see Fig. 2B, schematics, OPG2-ORF6-OPG2, OPG2-ORF6 and ORF6-OPG2). Following membrane insertion, doubly tagged OPG2-ORF6-OPG2 shows significant amounts of species with 3- and 4-N-linked glycans (Fig. 2B). This pattern confirms that the SARS-CoV-2 ORF6 protein assumes a 'hairpin' conformation in the ER membrane with both its N- and C-termini in the lumen (Fig. 2B, OPG2-ORF6-OPG2). These 3- and 4-N-glycan bearing OPG2-ORF6-OPG2 species are also resistant to extraction with alkaline sodium carbonate buffer (Fig. S1D) and protected from added protease (Fig. S1E), further indicating that the majority of the ORF6 protein is stably associated with the ER membrane in a 'hairpin' (N_{exo}/C_{exo}) topology.

Consistent with this unusual membrane topology, we find no indication that the membrane insertion of any of our OPG2-tagged ORF6 variants is reduced by Ipom-F, strongly suggesting that its association with the inner leaflet of the ER membrane does not require protein translocation via the central channel of the Sec61 translocon (Gérard et al., 2020; O'Keefe et al., 2020 submitted). We noted a sub-set of OPG2-ORF6-OPG2 species bearing only a single N-glycan was also clearly present in the membrane-associated fraction with or without Ipom-F treatment (Fig. 2B, OPG2-ORF6-OPG2, see 1Gly). Based on comparison to singly OPG2-tagged variants (Fig. 2B), we conclude that OPG2-ORF6-OPG2-1Gly has its N-terminus in the ER lumen,

where only one of its two consensus sites is efficiently N-glycosylated (cf. Nilsson and von Heijne, 1993), whilst its C-terminus is either ER luminal but non-glycosylated or remains on the cytosolic side of the membrane. In the latter case, it may be that, in addition to its hairpin topology, some fraction of ORF6 may be integrated into ER-derived microsomes as a type III TMP (cf. Fig. S2E; see also Netland et al., 2007) that is resistant to Ipom-F inhibition (this study; Zong et al., 2019; O'Keefe et al., 2020 submitted).

The molecular basis for SARS-CoV-2 protein sensitivity to Ipom-F

Having ascertained that Ipom-F inhibits the ER membrane translocation/insertion of the viral ORF8 and S proteins, but not of the ORF6, E or M proteins (Fig. 2C), we next investigated the molecular basis for this selectivity. For these studies we employed semi-permeabilised (SP) mammalian cells, depleted of specific membrane components via siRNA-mediated knockdown, as our source of ER membrane (Fig. 3A; Wilson et al., 2007). Consistent with our recent work (O'Keefe et al., 2020 submitted), and based on the quantitative immunoblotting of target and control gene products (Fig. S2A-C), we selectively depleted HeLa cells for core components of the Sec61 translocon (Sec61α-kd, ~65% reduction), the EMC (EMC5-kd, ~73% reduction) and both together (Sec61α+EMC5-kd, ~68% and ~78% reduction) prior to semi-permeabilisation with digitonin and use for *in vitro* ER translocation assays.

Following the analysis of total OPG2-tagged translation products recovered by immunoprecipitation, we found that: i) the S protein and a truncated derivative were both more strongly affected by the depletion of Sec61α than of EMC5 (Fig. 3B, Fig. S2D); ii) the ORF8 protein was likewise strongly affected by Sec61α depletion but also sensitive to EMC5 depletion (Fig. 3C); iii) the E protein showed diminished insertion efficiency after knock-down of Sec61α and EMC5, although the latter had a more pronounced effect (Fig. 3D). In each case, the combined knockdown of Sec61α and EMC5 resulted in a reduction of membrane insertion that was either comparable to, or greater than, that achieved following the knock-down of Sec61α alone (Figs. 3B to 3D). For the ORF6 protein, the total level of N-glycosylated OPG2-ORF6-OPG2 species was unaffected by any knockdown condition tested (Fig. 3E). However, we note a marked increase in the proportion of potentially mis-inserted OPG2-ORF6-

OPG2-1Gly species, particularly after co-depletion of EMC5 and Sec61α (see Fig. 3E;

Fig. S2E). We speculate that the unusual hairpin topology of the ORF6 protein may

be attributed to the EMC and Sec61 complex acting in concert to provide an Ipom-F

insensitive pathway for protein translocation across the ER membrane (O'Keefe et al.,

2020, submitted). Perturbation of this pathway seemingly increases the potential for

ORF6 to mis-insert (cf. Chitwood et al., 2018), perhaps as a consequence of disruption

to the translocation of its C-terminus (Fig. S2E).

Taken together, our data establish that, analogous to human membrane and secretory

proteins, the principal molecular basis for the Ipom-F-sensitivity of the SARS-CoV-2

ORF8 and S proteins is their dependence on Sec61-mediated protein translocation

into and across the ER membrane. In contrast, the E, M, and ORF6 proteins appear

capable of exploiting one or more alternative membrane insertion/translocation

pathways that can bypass the translocase activity of the Sec61 complex. These

alternatives most likely include a recently described route for type III TMP insertion

that requires the insertase function of the EMC (O'Keefe et al., 2020 submitted), which

our data suggest is also sufficient to confer Ipom-F-resistance to the viral E protein

and at least the first TM-span of the viral M protein.

<u>Concluding Remarks</u>

We conclude, that Sec61-selective protein translocation inhibitors like Ipom-F hold

promise as broad-spectrum antivirals that may exert a therapeutic effect by selectively

inhibiting the ER translocation of viral and/or host proteins which are crucial to viral

infection and propagation (Mast et al., 2020). In the context of SARS-CoV-2,

integration of the viral S protein and its host cell receptor, ACE2, into the ER

membrane is significantly reduced by Ipom-F (Fig. 2C, 3B). Likewise, translocation of

the viral ORF8 protein across the ER membrane and into its lumen is substantially

diminished (Fig. 2C, 3C). The binding of the viral S protein to cell surface ACE2 is a

key step in host cell infection (Drew and Janes, 2020), whilst ORF8 may protect SARS-

CoV-2 infected cells against host cytotoxic T lymphocytes (Zhang et al., 2020), making

all three of these proteins viable therapeutic targets (Drew and Janes, 2020; Li et al.,

2020; Young et al., 2020). Given that Ipom-F is a potent inhibitor of Sec61-mediated

protein translocation in cell culture models (Zong et al., 2019) and appears well

tolerated in mice (Zong et al., in press), we propose that future studies investigating

its effect on SARS-CoV-2 infection and propagation in cellular models are clearly
warranted.

Fig. 3: SARS-CoV-2 proteins are variably dependent on the Sec61 complex and/or the
EMC for ER membrane translocation/insertion. (A) Schematic of the *in vitro* ER import
assay in which control SP cells, or those depleted of a subunit of the Sec61 complex and/or
the EMC via siRNA, are used as a source of ER membrane. Following translation, OPG2-
tagged translation products (i.e. membrane-associated and non-targeted nascent chains) are
recovered by immunoprecipitation, resolved by SDS-PAGE and analysed by
phosphorimaging. OPG2-tagged variants of the SARS-CoV-2 **(B)** spike (S-OPG2), **(C)** ORF8
(ORF8-OPG2), **(D)** envelope (OPG2-E) and **(E)** ORF6 (OPG2-ORF6-OPG2, different Gly
species labelled as described in the legend to Fig. 2) proteins were synthesised in rabbit
reticulocyte lysate supplemented with control SP cells (lanes 1-2) or those with impaired
Sec61 complex and/or EMC function (lanes 3-6). Radiolabelled products were recovered and

analysed as described in **(A)**. Membrane translocation/insertion efficiency was determined
using the ratio of the N-glycosylation of lumenal domains, identified by treatment with Endo H
(EH, lane 1), relative to the NT control (set to 100% translocation/insertion efficiency).
Quantifications are given as means ±SEM for independent translation reactions from separate
siRNA treatments performed in triplicate (*n*=3) and statistical significance (two-way ANOVA,
DF and F values shown in the figure) was determined using Dunnett's multiple comparisons
test. Statistical significance is given as: n.s., non-significant >0.1; *, $P < 0.05$; **, $P < 0.01$; ***,
$P < 0.001$; ****, $P < 0.0001$.

Materials and Methods

Ipom-F and Antibodies

Ipom-F was synthesised as previously described (Zong et al., in press). Antibodies used to validate Sec61 and/or EMC subunit depletions in SP cells (Fig. S2) were purchased from Santa Cruz Biotechnology (goat polyclonal anti-LMNB1 (clone M-20, sc-6217), Bethyl Laboratories (rabbit polyclonal anti-EMC5 (A305-832-A)), Abcam (rabbit polyclonal anti-EMC6, (ab84902)), gifted by Sven Lang and Richard Zimmerman (University of Saarland, Homburg, Germany, rabbit anti-Sec61α) or as previously described (mouse monoclonal anti-OPG2 tag (McKenna et al., 2016) and rabbit polyclonal anti-OST48 (Wilson et al., 2007).

DNA constructs

The cDNA for human ACE2 (Uniprot: Q9BYF1) was purchased from Sino Biological (HG10108-M). cDNAs encoding the SARS-CoV-2 genes for ORF6, ORF8 and the E M and S proteins (Uniprot: P0DTC6, P0DTC8, P0DTC4, P0DTC5, P0DTC2 respectively) were kindly provided by Nevan Krogan (UCSF, US) (Gordon et al. 2020), amplified by PCR, subcloned into the pcDNA5 vector and constructs validated by DNA sequencing (GATC, Eurofins Genomics). ORF6-OPG2, ORF8-OPG2, M-OPG2 and S-OPG2 were generated by inserting the respective cDNAs in frame between NheI and AflII sites of a pcDNA5/FRT/V5-His vector (Invitrogen) containing a C-terminal OPG2 tag (MNGTEGPNFYVPFSNKTG). OPG2-E was generated by cloning the cDNA encoding the E-protein into the same pcDNA5-OPG2 vector using the KpnI and BamHI sites and deleting the stop codon after the OPG2 tag by site-directed mutagenesis (Stratagene QuikChange, Agilent Technologies). The N-terminal OPG2-tag of OPG2-ORF6-OPG2 was inserted by site-directed mutagenesis of ORF6-OPG2 using the relevant forward and reverse primers (Integrated DNA Technologies). Linear DNA templates were generated by PCR and mRNA transcribed using T7 polymerase.

siRNA-mediated knockdown and SP cell preparation

HeLa cells (human epithelial cervix carcinoma cells, ATCC, female) were cultured in DMEM supplemented with 10% (v/v) FBS and maintained in a 5% CO_2 humidified incubator at 37°C. Knockdown of target genes were performed as previously described (O'Keefe et al., 2020 submitted) using 20 nM (final concentration) of either control siRNA (ON-TARGETplus Non-targeting control pool; Dharmacon), *SEC61A1*

siRNA (Sec61α-kd, GE Healthcare, sequence AACACUGAAAUGUCUACGUUUUU), *MMGT1* siRNA (EMC5-kd, ThermoFisher Scientific, s41129) and INTERFERin (Polyplus, 409-10) as described by the manufacturer. 96 h post-initial transfection, cells were semi-permeabilsed using 80 µg/mL high purity digitonin (Calbiochem) and treated with 0.2 U Nuclease S7 Micrococcal nuclease from *Staphylococcus aureus* (Sigma-Aldrich, 10107921001) as previously described (O'Keefe et al., 2020 submitted; Wilson et al., 2007). SP cells lacking endogenous mRNA were resuspended (3×10^6 SP cells/mL as determined by trypan blue (Sigma-Aldrich, T8154) staining) in KHM buffer (110 mM KOAc, 2 mM Mg(OAc)$_2$, 20 mM HEPES-KOH pH 7.2) prior to analysis by western blot, or inclusion in translation master mixes such that each translation reaction contained 2×10^5 cells/mL.

In vitro ER import assays

Standard translation and membrane translocation/insertion assays, supplemented with nuclease-treated canine pancreatic microsomes (from stock with OD$_{280}$ = 44/mL) or siRNA-treated SP HeLa cells, were performed in nuclease-treated rabbit reticulocyte lysate (Promega) as previously described (Zong et al., 2019; O'Keefe et al., 2020 submitted): namely in the presence of EasyTag EXPRESS ^{35}S Protein Labelling Mix containing [^{35}S] methionine (Perkin Elmer) (0.533 MBq; 30.15 TBq/mmol), 25 µM amino acids minus methionine (Promega), 1 µM Ipom-F, or an equivalent volume of DMSO, 6.5% (v/v) ER-derived microsomes or SP cells and ~10% (v/v) of *in vitro* transcribed mRNA (~500 ng/µL) encoding the relevant precursor protein. Microsomal translation reactions (20 µL) were performed for 30 min at 30°C whereas those using SP HeLa cells were performed on a 1.5X scale (30 µL translation reactions) for 1 h at 30°C. As the S protein was most efficiently synthesised using the TNT® Coupled system (Fig. S1B), import assays of the comparatively higher molecular weight ACE2 and S proteins (50 µL reactions) were both performed using the TNT® Coupled Transcription/ Translation system (Promega) for 90 min at 30°C as described by the manufacturer (~50 ng/µL cDNA, 1 µM Ipom-F or an equivalent volume of DMSO, 12% (v/v) ER-derived microsomes or SP cells). All translation reactions were finished by incubating with 0.1 mM puromycin for 10 min at 30°C to ensure translation termination and ribosome release of newly synthesised proteins prior to analysis.

Recovery and analysis of radiolabelled products

Following puromycin treatment, microsomal membrane-associated fractions were recovered by centrifugation through an 80 µL high-salt cushion (0.75 M sucrose, 0.5 M KOAc, 5 mM Mg(OAc)$_2$, 50 mM Hepes-KOH, pH 7.9) at 100,000 g for 10 min at 4°C and the pellet suspended directly in SDS sample buffer. To confirm the topology of ORF6 (Fig. S2), the membrane-associated fraction of the doubly-OPG2-tagged form (OPG2-ORF6-OPG2) was resuspended in KHM buffer (20 µL) and subjected to either carbonate extraction (0.1 M Na$_2$CO$_3$) (McKenna et al., 2016) or a protease protection assay using trypsin (1 µg/mL) with or without 0.1% Triton X-100 (Ray-Sinha et al., 2009) prior to suspension in SDS sample buffer. For translation reactions using SP cells, the total reaction material was diluted with nine volumes of Triton immunoprecipitation buffer (10 mM Tris-HCl, 140 mM NaCl, 1 mM EDTA, 1% Triton X-100, 5 mM PMSF, 1 mM methionine, pH 7.5). Samples were incubated under constant agitation with an antibody recognising the OPG2 epitope (1:200 dilution) for 16 h at 4°C to recover both the membrane-associated and non-targeted nascent chains. Samples were next incubated under constant agitation with 10% (v/v) Protein-A-Sepharose beads (Genscript) for a further 2 h at 4°C before recovery by centrifugation at 13,000 g for 1 min. Protein-A-Sepharose beads were washed twice with Triton immunoprecipitation buffer prior to suspension in SDS sample buffer. Where indicated, samples were treated with 1000 U of Endoglycosidase Hf (~10-50 kDa protein substrates; New England Biolabs, P0703S) or Endoglycosidase H (~50-150 kDa protein substrates; New England Biolabs, P0702S), and all samples were solubilised for 12 h at 37°C and then sonicated prior to resolution by SDS-PAGE (10% or 16% PAGE, 120V, 120-180 min). Gels were fixed for 5 min (20% MeOH, 10% AcOH), dried for 2 h at 65°C and radiolabelled products visualised using a Typhoon FLA-700 (GE Healthcare) following exposure to a phosphorimaging plate for 24-72 h.

Western Blotting

Following semi-permeabilisation, aliquots of siRNA-treated HeLa cells were suspended in SDS sample buffer, denatured for 12 h at 37°C and sonicated prior to resolution by SDS-PAGE (16% or 10% PAGE, 120V, 120-150 min). Following transfer to a PVDF membrane in transfer buffer (0.06 M Tris, 0.60 M glycine, 20% MeOH) at 300 mA for 2.5 h, PVDF membranes were incubated in 1X Casein blocking buffer (10X

stock from Sigma-Aldrich, B6429) made up in TBS, incubated with appropriate primary
antibodies (1:500 or 1:1000 dilution) and processed for fluorescence-based detection
as described by LI-COR Biosciences using appropriate secondary antibodies (IRDye
680RD Donkey anti-Goat, IRDye 680RD Donkey anti-Rabbit, IRDye 800CW Donkey
anti-Mouse) at 1:10,000 dilution. Signals were visualised using an Odyssey CLx
Imaging System (LI-COR Biosciences).
391

<u>Quantification and Statistical Analysis</u>
Bar graphs depict either the efficiency of membrane translocation/insertion calculated
as the ratio of N-glycosylated protein relative to the amount of non-N-glycosylated
protein (Fig. 2-3), or the efficiencies of siRNA-mediated knockdown in SP cells
calculated as a proportion of the protein content when compared to the NT control
(Fig. S2), with all control samples set to 100%. Normalised values were used for
statistical comparison (one-way or two-way ANOVA; DF and F values are shown in
each figure as appropriate and the multiple comparisons test used are indicated in the
appropriate figure legend). Statistical significance is given as n.s., non-significant >0.1;
*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$.
402

Acknowledgements
We thank Quentin Roebuck for technical assistance, Nevan Krogan (UCSF, US) for
providing the original SARS-CoV-2 plasmids, Sven Lang (University of Saarland,
Homburg, Germany) for Sec61α antisera, Belinda Hall (University of Surrey, UK) and
Rachel Simmonds (University of Surrey, UK) for passing on the SARS-CoV-2 plasmids
and useful discussions. We are indebted to Richard Zimmermann (University of
Saarland) for catalyzing SARS-CoV-2 related discussions amongst the ER research
community and for critical feedback during the preparation of our manuscript.

Competing interests
The authors declare no competing interests.

Author Contributions

K.B.D., G.Z. and H.S. participated in the chemical synthesis of Ipom-F and W.Q.S supervised the synthesis; P.R. subcloned and validated the SARS-CoV-2 plasmids used in this study; S.O'K. performed the site-directed mutagenesis and experiments; S.O'K. and S.H. designed the study, analysed the data and wrote the manuscript.

Funding

This work was supported by a Welcome Trust Investigator Award in Science 204957/Z/16/Z (S.H.), an AREA grant 2R15GM116032-02A1 from the National Institute of General Medical Sciences of the National Institutes of Health (NIH) and a Ball State University (BSU) Provost Startup Award (W.Q.S.).

Supplementary Information

Supplementary information accompanies this short report and includes Fig. S1 and Fig. S2.

References

429
430 **Adalja, A., and Inglesby, T.** (2019). Broad-spectrum antiviral agents: A crucial
431 pandemic tool. *Exp. Rev. Anti-Infect. Ther.* **17**, 467-470.

432
433 **Bojkova, D., Klann, K., Koch, B., Widera, M., Krause, D., Ciesek, S., Cinatl, J.,**
434 **and Münch, C.** (2020). Proteomics of SARS-CoV-2 infected host cells reveals therapy
435 targets. Nature. 583, 469-472.

436
437 **Cantuti-Castelvetri, L., Ojha, R., Pedro, L. D., Djannatian, M., Franz, J., Kuivanen,**
438 **S., van der Meer, F., Kallio, K., Kaya, T., Anastasina, M., et al.** (2020). Neuropilin-
439 1 facilitates SARS-CoV-2 cell entry and infectivity. *Science.* **370**, 856-860.

440
441 **Chitwood, P. J., Juszkiewicz, S., Guna, A., Shao, S., and Hegde, R. S.** (2018).
442 EMC is required to initiate accurate membrane protein topogenesis. *Cell.* **175**, 1-13.

443
444 **Daly, J. L., Simonetti, B., Klein, K., Chen, K.-E., Kavanagh Williamson, M., Antón-**
445 **Plágaro, C., Shoemark, D. K., Simón-Gracia, L., Bauer, M. et al.** (2020). Neuropilin-
446 1 is a host factor for SARS-CoV-2 infection. *Science*. **370**, 861-865.

447
448 **Drew, E. D., and Janes, R. W.** (2020). Identification of a druggable binding pocket in
449 the spike protein reveals a key site for existing drugs potentially capable of combating
450 Covid-19 infectivity. *BMC Mol. Cell Biol.* **21**, 49.

451
452 **Duart, G., García-Murria, M. J., Grau, B., Acosta-Cáceres, J. M., Martínez-Gil, L.,**
453 **and Mingarro I.** (2020) SARS-CoV-2 envelope protein topology in eukaryotic
454 membranes. *Open Biol.* **10**, 200209.

455
456 **Firth, A. E.** (2020). A putative new SARS-CoV protein, 3c, encoded in an ORF
457 overlapping ORF3a. *J. Gen. Virol.* **101**, 1085-1089**.**

458
Gérard, S. F., Hall, B. S., Zaki, A. M., Corfield, K. A., Mayerhofer, P. U., Costa, C.,
Whelligan, D. K., Biggin P. C., Simmonds, R. E., and Higgins, M. K. (2020).
Structure of the inhibited state of the Sec translocon. *Mol. Cell.* **79**, 406-415.e7.

Gordon, D. E., Jang, G. M., Bouhaddou, M., Xu, J., Obernier, K., White, K. M.,
O'Meara, M. J., Rezelj, V. V., Guo, J. Z., Swaney, D. L., Tummino, T. A. et al. (2020)
A SARS-CoV-2 protein interaction map reveals targets for drug repurposing. *Nature.*
583, 459-468.

Heaton, N. S., Moshkina, N., Fenouil, R., Gardner, T. J., Aguirre, S., Shah, P. S.,
Zhao, N., Manganaro, L., Hultquist, J. F., Noel, J. et al. (2016). Targeting viral
proteostasis limits influenza virus, HIV, and Dengue virus infection. *Immunity.* **44**, 46-
58.

Li, J.-Y., Liao, C.-H., Wang, Q., Tan, Y.-., Luo, R., Qiu, Y., and Ge, X.-Y. (2020). The
ORF6, ORF8 and nucleocapsid proteins of SARS-CoV-2 inhibit type I interferon
signalling pathway. *Virus Res.* **286**, 198074.

Luesch, H., and Paavilainen, V. O. (2020). Natural products as modulators of
eukaryotic protein secretion. *Nat. Prod. Rep.* **37**, 717-736.

Mast, F. D., Navare, A. T., van der Sloot, A. M., Coulombe-Huntington, J. Rout,
M. P., Baliga, N. S., Kaushansky, A., Chait, B. T., Aderem, A., Rice, C. M. et al.
(2020). Crippling life support for SARS-CoV-2 and other viruses through synthetic
lethality. J. Cell. Biol. **219**, e202006159.

McKenna, M., Simmonds, R.E., and High, S. (2016). Mechanistic insights into the
inhibition of Sec61-dependent co- and post-translational translocation by
mycolactone. *J. Cell Sci.* **129**, 1404-1415.

Morel, J. D., Paatero, A. O., Wei, J., Yewdell, J. W., Guenin-Macé, L., Van Haver,
D., Impens, F., Pietrosemoli, N., Paavilainen, V. O., and Demangel, C. (2018).
Proteomics reveals scope of mycolactone-mediated Sec61 blockade and distinctive
stress signature. *Mol. Cell Prot.* **17**, 1750-1765.

Naqvi, A. A. T., Fatima, K., Mohammad, T., Fatima, U., Singh, I. K., Singh, A., Atif,
A. M., Hariprasad, G., Hasan, G. M., and Hassan, M. I. (2020) Insights into SARS-

CoV-2 genome, structure, evolution, pathogenesis and therapies: structural genomics approach. *Biochim. Biophys. Acta. Mol. Basis Dis.* **1866**, 165878.

Netland, J., Ferraro, D., Pewe, L., Olivares, H., Gallagher, T., and Perlman, S. (2007). Enhancement of murine coronavirus replication by severe acute respiratory syndrome coronavirus protein 6 requires the N-terminal hydrophobic region but not C-terminal sorting motifs. *J. Virol.* **81**, 11520-11525.

Nilsson, I. M., and von Heijne, G. (1993). Determination of the distance between the oligosaccharyltransferase active site and the endoplasmic reticulum membrane. *J. Biol. Chem.* **268**, 5798-5801.

Ray-Sinha, A., Cross, B. C. S., Mironov, A., Wiertz, E., and High, S. (2009). Endoplasmic reticulum-associated degradation of a degron-containing polytopic membrane protein. *Mol. Membr. Biol.* **26**, 448-464.

O'Keefe, S., Zong, G., Duah, K. B., Andrews, L. E., Shi, W. Q., and High, S. (2020). Type III transmembrane protein integration requires both the EMC and Sec61 complex. Submitted.

Sicari, D., Chatziioannou, A., Koutsandreas, T., Sitia, R., and Chevet, E. (2020). Role of the early secretory pathway in SARS-CoV-2 infection. *J. Cell. Biol.* **219**, e202006005.

Shah, P. S., Link, N., Jang, G. M., Sharp, P. P., Zhu, T., Swaney, D. L., Johnson, J. R., Von Dollen, J., Ramage, H. R., Satkamp, L. et al. (2018). Comparative flavivirus-host protein interaction mapping reveals mechanisms of Dengue and Zika virus pathogenesis. *Cell.* **175**, 1931-1945.e18.

Von Heijne, G. (2007). The membrane protein universe: what's out there and why bother? *J. Intern. Med.* **261**, 543-547.

Walls, A. C., Park, Y.-J., Tortorici, M. A., Wall, A., McGuire, A. T., and Veesler, D.
(2020). Structure, function and antigenicity of the SARS-CoV-2 spike glycoprotein.
Cell. **181**, P281-292.E6.

Warner, F. J., Lew, R. A., Smith, A. I., Lambert, D. W., Hooper, N. M., and Turner,
A. T. (2005). Angiotensin-converting enzyme 2 (ACE2), but not ACE, is preferentially
localised to the apical surface of polarised kidney cells. *J. Biol. Chem.* **280**, 39353-
39362.

Wilson, C. M., and High, S. (2007). Ribophorin I acts as a substrate-specific facilitator
of N-glycosylation. *J. Cell. Sci.* **120**, 648-657.

Young, B. E., Fong, S.-W., Chan, Y. H., Mak, T.-M., Ang, L. W., Anderson, D. E.,
Yi-Pin Lee, C., Naqiah Amrun, S., Lee, B., Shan Goh, Y. et al. (2020). Effects of a
major deletion in the SARS-CoV-2 genome on the severity of infection and
inflammatory response: an observational cohort study. *Lancet.* **396**, 603-611.

Zhang, Y., Zhang, J., Chen. Y., Luo, B., Yuan, Y., Huang, F., Yang, T., Yu, F., Liu,
J., Song, Z. et al. (2020). The ORF8 protein of SARS-CoV-2 mediates immune
evasion through potently downregulating MHC-1. bioRxiv.
doi: 10.1101/2020.05.24.111823

Zhou, P., Yang, X.-L., Wang, X.-G., Hu, B., Zhang, L., Zhang, W., Si, H.-R., Zhu,
Y., Li, B., Huang, C.-L. et al. (2020). A pneumonia outbreak associated with a new
coronavirus of probable bat origin. *Nature.* **579**, 270-273.

Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi,
W., Lu, R., et al. (2020). A novel coronavirus from patients with pneumonia in China,
2019. *N. Eng. J. Med.* **382**, 727-733.

Zong, G., Hu, Z., O'Keefe, S., Tranter, D., Iannotti, M. J., Baron, L., Hall, B.,
Corfield, K., Paatero, A., Henderson M. et al. (2019). Ipomoeassin F binds Sec61α
to inhibit protein translocation. *J. Am. Chem. Soc.* **141**, 8450-8461.

Zong, G., Hu, Z., Duah, K., B., Andrews, L. E., Zhou, J., O'Keefe, S., Whisenhunt,

L., Shim, J. S., Du, Y., High, S., et al. (2020) Ring-expansion leads to a more potent

analogue of ipomoeassin F. *J. Org. Chem.* In press.

Supplementary information for

Ipomoeassin-F inhibits the *in vitro* biogenesis of the SARS-CoV-2 spike protein and its host cell membrane receptor

Sarah O'Keefe, Peristera Roboti, Kwabena B. Duah, Guanghui Zong, Hayden Scheider, Wei Q. Shi and Stephen High

Fig. S1. Additional studies using ER microsomes, Related to Figures 2 and 3.
(**A**) Non-tagged (lanes 1-3) and OPG2-tagged (lanes 4-6) versions of the SARS-CoV-2 spike protein (S, S-OPG2), ORF8 (ORF8, ORF8-OPG2) and membrane protein (M, M-OPG2) were synthesised in rabbit reticulocyte lysate supplemented with ER-derived canine pancreatic microsomes in the absence and presence of Ipom-F (lanes 1 and 3). Phosphorimages of membrane-associated products resolved by SDS-PAGE together with representative substrate outlines are shown. N-glycosylated (X-Gly) versus non-N-glycosylated (0Gly) species were identified by treatment with endoglycosidase H (Endo H, lanes 2 and 5). (**B**) The S protein was synthesised in a Flexi® rabbit reticulocyte system with varying concentrations of magnesium acetate (lanes 1-5) and a TNT® Coupled system (lane 6) in the absence of ER-derived

microsomes. 5% of the total reaction material was resolved by SDS-PAGE and visualised by phosphorimaging. (**C**) The ER import of truncated variants of the S protein (S-short, S-s.s.-TMD, S-half-OPG2) was analysed as described for (**A**). (**D**) The membrane-associated products of the doubly tagged form of ORF6 (OPG2-ORF6-OPG2) were synthesised as in (**A**) and, following treatment with sodium carbonate buffer and centrifugation, the pellet, enriched for membrane-integrated material, and supernatant, largely containing peripherally membrane-associated material, were analysed for OPG2-ORF6-OPG2. (**E**) The membrane-associated products of OPG2-ORF6-OPG2 were treated with trypsin in the absence or presence of Triton-100 (TX-100, lanes 2-3).

Fig. S2. Validation of Sec61 and/or EMC subunit depletions in SP cells, Related to Figure 3.

(A) The effects of transfecting HeLa cells with non-targeting (NT; lane 1), Sec61α-targeting (lane 2), EMC5-targeting (lane 3) and Sec61α+EMC5-targeting (lane 4) siRNAs were determined after semi-permeabilisation by immunoblotting for target genes (Sec61α, EMC5). Controls to assess destabilisation of the wider EMC complex (EMC2 and EMC6), any effect on the N-glycosylation machinery (the ER-resident 48 kDa subunit of the oligosaccharyl-transferase complex (OST48) and the quantity of SP cells used in each experiment (the nuclear protein Lamin-B1 (LMNB1)), are also shown. **(B)** The efficiencies of siRNA-mediated knockdown (bold) were calculated as a proportion of the signal intensity obtained with the NT control (set as 100%).

Quantifications are given as means ±SEM for three separate siRNA treatments (*n*=3) with statistical significance of siRNA-mediated knockdowns (two-way ANOVA, DF and F values shown in the figure) determined using Dunnett's multiple comparisons test. Statistical significance is given as n.s., non-significant >0.1; *, $P < 0.05$; ****, $P < 0.0001$. (**C**) Mean knockdown efficiencies ($\pm$SEM) for each of the target genes. (**D**) A truncated variant of the S protein (S-half-OPG2) was synthesised in rabbit reticulocyte lysate supplemented with SP cells with impaired Sec61 complex and/or EMC function and recovered by immunoprecipitation via the OPG2 tag. Radiolabelled products resolved by SDS-PAGE and analysed by phosphorimaging. N-glycosylated (14-Gly) versus non-N-glycosylated (0Gly) species were identified by treatment with endoglycosidase H (Endo H, lane 1). (**E**) Further analysis of the data presented in Fig. 3E of the main text. Here, the ratio of 3Gly and 4Gly bearing OPG2-ORF6-OPG2 N-glycosylated species relative to the 1Gly species present in the same sample was used as a proxy to estimate potential mis-insertion of the ORF6 protein in SP cells with impaired Sec61 complex and/or EMC function relative to the NT control (set to 100% efficiency). Quantifications are given as means ±SEM for independent translation reactions from separate siRNA treatments performed in triplicate (*n*=3) and statistical significance (two-way ANOVA, DF and F values shown in the figure) was determined using Dunnett's multiple comparisons test. Statistical significance is given as n.s., non-significant >0.1; *, $P < 0.05$; ***, $P < 0.001$.