Snowball Earths, population bottlenecks, and the evolution of marine photosynthetic bacteria

Hao Zhang[1,^], Ying Sun[1,^], Qinglu Zeng[2], Sean A. Crowe[3], Haiwei Luo[1,*]

[1]Simon F. S. Li Marine Science Laboratory, School of Life Sciences and State Key Laboratory of Agrobiotechnology, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[2]Department of Ocean Science, The Hong Kong University of Science and Technology, Clear Water Bay, Hong Kong SAR

[3]Department of Earth Sciences, School of Biological Sciences, and Swire Institute for Marine Science (SWIMS), University of Hong Kong, Pokfulam Road, Hong Kong SAR

^These authors contributed equally to this study.

*Corresponding author:
Haiwei Luo
The Chinese University of Hong Kong
Shatin, Hong Kong SAR
Phone: (+852) 39436121
E-mail: hluo2006@gmail.com

Running Title: *Prochlorococcus* and Snowball Earths

Keywords: *Prochlorococcus*, Neoproterozoic Snowball Earth, genome reduction, molecular dating.

Abstract

Prochlorococcus are the most abundant photosynthetic organisms in the modern ocean. A massive DNA loss event occurred in their early evolutionary history, leading to highly reduced genomes in nearly all lineages, as well as enhanced efficiency in both nutrient uptake and light absorption. The environmental landscape that shaped this ancient genome reduction, however, remained unknown. Through careful molecular clock analyses, we established that this *Prochlorococcus* genome reduction occurred during the Neoproterozoic Snowball Earth climate catastrophe. The lethally low temperature and exceedingly dim light during the Snowball Earth event would have inhibited *Prochlorococcus* growth and proliferation and caused severe population bottlenecks. These bottlenecks are recorded as an excess of deleterious mutations that accumulated across genomic regions in the descendant lineages. *Prochlorococcus* adaptation to extreme environmental conditions during Snowball Earth intervals can be inferred by tracing the evolutionary paths of genes that encode key metabolic potential. This metabolic potential includes modified lipopolysaccharide structure, strengthened peptidoglycan biosynthesis, the replacement of a sophisticated circadian clock with an hourglass-like mechanism that resets daily for dim light adaption, and the adoption of ammonia diffusion as an efficient membrane transporter-independent mode of nitrogen acquisition. In this way, the Neoproterozoic Snowball Earth event altered the physiological characters of *Prochlorococcus*, shaping their ecologically vital role as the most abundant primary producers in the modern oceans.

Introduction

Prochlorococcus are the smallest and most abundant photosynthetic organisms on Earth [1]. They are prevalent throughout the photic zone of the oligotrophic oceans between 40 °N and 40 °S[1], where they account for more than 40% of the biomass and contribute almost half of the net primary production[2]. *Prochlorococcus* have diversified into two major phylogenetic groups with distinct ecology (ecotypes), with the high-light (HL) adapted monophyletic group imbedded in the low-light (LL) adapted paraphyletic group[3]. The distinct ecotypes of *Prochlorococcus* evolved different pigments, light-harvesting systems, and the phycobiliproteins, which allowed for efficient light absorption in the water column[4], and thus increased growth rates and primary production[5].

Prochlorococcus genomes have been shaped by stepwise streamlining, including a major genome reduction in their early evolution and a few minor modifications that followed[6,7]. Modern marine *Prochlorococcus* lineages, in particular those with small genomes, show very low ratios of nonsynonymous (d_N) to synonymous (d_S) nucleotide substitution rates, suggesting that natural selection is a highly efficient throttle on the accumulation of deleterious mutations (i.e., nonsynonymous mutations) in *Prochlorococcus*[6,8]. On long time scales, however, nucleotide substitutions at synonymous sites become saturated, invalidating the use of d_N/d_S to infer selection efficiency in deep time[9]. Thus, an alternative approach focuses instead on different types of nonsynonymous substitutions leading to radical versus conservative changes in amino acid sequences, with the former more likely to be deleterious[10,11]. Excess radical mutations accumulate from random fixation of deleterious mutations by genetic drift (i.e., reduced efficiency of selection). Using this approach reveals that the major genome

reduction in *Prochlorococcus* took place under reduced selection efficiency[9] and implies

that the ancient population went through severe bottlenecks as the likely result of

environmental catastrophe.

The environmental context underlying *Prochlorococcus* genome reduction remains

unknown, however, and precise molecular dating is needed to link this important

evolutionary event to its possible environmental drivers. Implementing comprehensive

molecular clock analyses, we now link the early major genome reduction event of

Prochlorococcus to the Neoproterozoic Snowball Earth events. These catastrophic

disruptions to the Earth system would likely have challenged warm-water-loving

photosynthetic *Prochlorococcus*, with strong potential to cause the population

bottlenecks inferred from the genome sequences described above. *Prochlorococcus*

survived this catastrophe through likely gains and losses of key metabolic functions

reconstructed from the same genome sequences, which have far-reaching impact on their

success in today's oceans. *Prochlorococcus* are thus vital "guardians of metabolism"[12],

shepherding genes critical to the functioning of the biosphere across environmental

catastrophes, including global glaciations.

Results and Discussion

Prochlorococcus experienced a massive gene loss event on the ancestral branch

leading to the last common ancestor (LCA) of clades HL, LLI and LLII/III[6,7,13]. This is

confirmed by our analysis, which reconstructed 366 and 107 gene family losses and gains

on this branch, respectively (Fig. 1A & S1), among which 163 and 84 encode unknown

proteins. On the same ancestral branch, it was shown that d_R/d_C is significantly elevated

compared to the sister ancestral branch leading to the LCA of the LLIV[9], which was

validated here (both sign test and paired *t*-test, $p < 0.001$; Fig. 1B). These results confirm

that the major genome reduction event occurring on this branch was likely driven by

genetic drift as a result of one or recurrent population bottlenecks[9]. Given the global

distribution and abundance of *Prochlorococcus*, and cyanobacteria more generally, such

a bottleneck would likely require a global-scale event, like an environmental or climate

catastrophe (e.g. meteorite impact, large igneous province emplacement, or glaciation).

To establish the environmental context for the large, ancient genome reduction, we

estimated the timeline of *Prochlorococcus* evolution by implementing molecular clock

analyses based on essential calibrations available in the cyanobacterial lineage. We

recognize that the use of calibration sets adapted from previous studies (under

calibrations C1-C6 in Table S1 with related references included there) results in up to

~320 Ma disparity (Fig. S2) in the estimated time for the LCA of *Prochlorococcus* HL,

LLI and LLII/III clades that emerged with the major genome reduction. We note that the

calibrations in previous studies were not properly used. For example, the akinete fossil

identified to 2,100 Mya was used as either the maximum bound or the minimum bound to

calibrate the crown group of *Nostocales*[14,15]. However, given the fact that apomorphic

character must evolve earlier than the divergence of crown group, morphological fossils

can only serve as the minimum bounds on total groups of assigned lineages[16] (see Section

2.3 in Supplemental Methods for details). Thus, in the present study, we modified the

calibration sets by constraining the lower bounds of the *Nostocales* (and also

Pleurocapsales) total groups with morphological fossils and by leaving their upper

bounds open (C9-C14; Table S1). Intriguingly, the variation is reduced to less than 10 Ma

when these modified calibration sets are used (Fig. S2). By comparing the width of the

95% highest posterior density (HPD) derived from each molecular clock analysis (Fig.

S3), we inferred the most precise timeline of *Prochlorococcus* evolution (corresponding

to calibration set C14 in Table S1; see Section 2.3 in Supplemental Methods).

Our time estimates revealed that the LCA of *Prochlorococcus* HL, LLI and LLII/III

clades diversified at 682 Mya [95% HPD (=highest posterior density) 632-732 Mya],

precisely dating the genome reduction event to this time. A 682 Mya date for the

emergence of the LCA of *Prochlorococcus* HL, LLI and LLII/III clades places the large

genome reduction that took place in this lineage firmly within the Cryogenian Period

(~720 to 635 Mya; Fig. 1A) and implicates the Snowball Earth icehouse climate

conditions eponymous with the Period in the corresponding *Prochlorococcus* population

bottleneck. We therefore refer to this ancestor as SBE-LCA (see Fig. 1A), short for

"Snowball Earth" LCA. The Neoproterozoic climate catastrophe culminated in the

Sturtian (~717 to 659 Mya) and Marinoan (~645 to 635 Mya) glaciations (Fig. 1A),

which stretched from the poles to sea level near the equators, possibly wrapping the

entire Earth under a frozen skin[17]. This "Snowball Earth" persisted with the freezing

temperature of seawater below the ice sheet lowered to -3.5 ℃[18]. Since all assayed

Prochlorococcus strains, including those affiliated with the basal LL ecotypes, reach

maximum growth rates at approximately 25 ℃ and rarely survive when temperature drops

to ~10 ℃ (Fig. 1C)[19], we propose that extreme climate cooling during the Neoproterozoic

Snowball Earth events was likely the major driver of severe bottlenecks in early

Prochlorococcus populations.

Survival of *Prochlorococcus* populations through the Cryogenian would have

required refugia, the nature of which would have shaped continued *Prochlorococcus*

evolution. A variety of biotic refugia have been identified during Snowball Earth

intervals, including the sea-ice brine channels within ice grounding-line crack systems[20]

and cryoconite holes/ponds on the surface of the sublimation zone, which may have

represented ~12% of the global sea glacier surface[21] (Fig. 1D). Despite providing the

essential space for *Prochlorococcus* survival, these refugia would have presented a

number of environmental stresses to *Prochlorococcus* populations including: low

temperature, low light, high and variable osmotic pressure, and limited nutrients[20-22].

Prochlorococcus thus evolved a number of adaptive mechanisms to cope with these

stresses via gene gains and losses, which we assessed by reconstructing the evolutionary

paths of imprints that the Snowball Earth climate left in extant *Prochlorococcus*

genomes.

Among these stresses, the most prominent was likely lethally low temperature.

Maintaining membrane fluidity is of paramount importance under low temperature

conditions, which is largely achieved by the activities of fatty acid desaturase encoded by

desA and *desC*. As a result, we inferred that these genes were retained in SBE-LCA (Fig.

1A). Lipopolysaccharide (LPS) in the outer membrane is known to provide the first line

of defense against harsh environments[23], which contains the O-specific polysaccharide,

the glycolipid anchor lipid A, and the polysaccharide core region. Based on our analyses,

genes encoding the polysaccharide core region (*kdsABCD* for 3-deoxy-d-manno-

octulosonate biosynthesis; Fig. 1A) were likely lost at SBE-LCA, while those encoding

the other components were retained (*lpxABCD* and *rfbABC* for Lipid A precursor and O-

specific LPS precursor biosynthesis; Fig. 1A). This inference is consistent with a

previous conclusion that the loss of the LPS core region would increase the

hydrophobicity and permeability of the cell envelope[24] to protect against the cold

conditions[25]. The amino sugar N-acetylglucosamine (GlcNAc) is used by bacteria such as

Corynebacterium glutamicum as a carbon, energy and nitrogen source[26]. GlcNAc enters

bacteria in the form of GlcNAc-6-phosphate (GlcNAc-6-P). However, instead of being

metabolized, the loss of *nagB* for GlcNAc-6-P deamination at SBE-LCA suggests that

GlcN6P is more likely to be involved in peptidoglycan (PG) recycling through the

cascade catalysis by GlmM and GlmU (Fig. 1A) to generate UDP-GlcNAc, which is an

essential precursor of cell wall PG and LPS[27]. During cell turnovers, PG is continuously

broken down and reused through the PG recycling pathway to produce new PG, and in

some bacteria, PG recycling is critical for their long-term survival when growth is stalled

under nutrient limitation[28]. Thus, such a recycling mechanism seems to be key for

maintenance of cell integrity in SBE-LCA under oligotrophic and lethally freezing

conditions[21]. Another metabolic modification in SBE-LCA was related to heat shock

proteins (HSPs), which play crucial roles in tolerating environmental stresses including

thermal shocks. Typically, HSPs are tightly regulated, as they respond quickly to stress

and turn off rapidly once the stress disappears[29]. However, the HSP repressor protein

encoded by *hrcA* was inferred to be lost at SBE-LCA, which thus likely allowed the

organism to continuously express HSPs to cope with prolonged lethally low temperature.

In fact, constitutive expression of HSPs occurs in polar organisms such as the Antarctic

ciliate *Euplotes focardii*[30] and the polar insect *Belgica antarctica* larvae[31]. Extremely low

temperature also made substrate acquisition difficult due to increased lipid stiffness and

decreased efficiency and affinity of membrane transporters[32]. Under such conditions,

bacteria may increasingly rely on substrates whose uptake shows lower dependence on

temperature. In sea-ice brines where less CO_2 is dissolved[33], elevated pH promotes the

conversion of ammonium to ammonia, which diffuses directly into cells without the aid

of transporters in the membrane. Accordingly, species of bacteria and microalgae show a

greater dependence on ammonium and ammonia at low temperatures and high pH than

nitrate[34,35], thereby reducing reliance on membrane transporters. In SBE-LCA, the

potentially efficient utilization of ammonia made other N acquisition genes dispensable,

leading to the neutral loss of nitrite transporter (*nitM*), whereas glutamine synthetase

(*glnA*) and glutamate synthase (*gltS*) responsible for the utilization of ammonia after its

assimilation were conserved (Fig. 1A).

An additional stress to *Prochlorococcus* during Snowball Earth would have been

variable osmotic pressure. Although multiple refugia might have supported bacterial

survival during Snowball Earth, the schizohaline nature of these refugia must have

imposed strong osmotic pressure on ice-trapped bacteria[21]. As temperature dropped, salts

would have become increasingly concentrated in brine channels[20], whereas cryoconite

ponds would have remained hyposaline, similar to modern Arctic and Antarctic

cryoconite ecosystems[36]. Glycine betaine (GB) is among the most important organic

osmolytes in halophilic cyanobacteria, and is used as the major osmolyte in

Synechococcus sp. WH8102[37]. However, genes involved in glycine betaine biosynthesis

and transport were lost at SBE-LCA, including *bsmB* for dimethylglycine N-

methyltransferase, *gsmt* for glycine/sarcosine N-methyltransferase, and *proVWXP* for

glycine betaine/proline transport system (Fig. 1A). Instead, several other organic

osmolytes might have been used during the Snowball Earth, as their biosynthetic genes

were retained at SBE-LCA. The first examples are the *ggpS* gene encoding

glucosylglycerol phosphate synthase for glucosylglycerol (GG) synthesis and the *gpgS*

encoding glucosyl-phosphoglycerate synthase for glucosylglycerate (GGA) synthesis

(Fig. 1A). GG and GGA may be sufficient to provide osmotic tolerance under moderately

saline conditions[38]. Interestingly, the biosynthesis of GG and GGA requires less N

compared to that of GB[38,39], and thus the potential use of GG/GGA instead of GB

appeared favorable to SBE-LCA, which had a reduced efficiency of membrane

transporters and a low affinity for external nutrients at exceedingly low temperature.

Low light intensity during Snowball Earth was another formidable challenge to

phototrophs including *Prochlorococcus*. In the Neoproterozoic, the Sun was still at least

6% dimmer than that at present[40]. Moreover, sea ice, especially when covered with snow,

is an effective barrier to light transmission[41]. This is in analogy to the deeper layers of

today's polar snow and glacier ice where irradiation is reduced and photosynthetic

organisms and activities are scarcely detectable[42]. Consequently, photosynthetic

organisms trapped in the brine channels or inhabiting waters below ice need to be

physiologically geared to cope with low light. It was proposed that modification of the

photosystem structure enables adaptation to the low light condition[43]. We inferred a few

changes in photosystem I and II (PSI/PSII) that occurred in SBE-LCA, including the gain

of RC1 subunit PsaM, RC2 subunit PsbY, and an extra copy of the RC2 subunit PsbF, the

loss of RC2 protein PsbU, and the replacement of RC2 subunit PsbX (Fig. 1A), but the

molecular mechanism of these changes underlying low light adaptation is poorly

understood. We also inferred an expansion of the *Prochlorococcus* antenna Pcb from two

to six copies during the Snowball Earth (Fig. 1A), which may boost the light-harvesting

capacity under low-light conditions[44].

Many cyanobacteria have a sophisticated circadian clock, which is essential in

controlling global diel transcriptional activities of the cells. This circadian oscillator

system requires only three components: KaiA, KaiB and KaiC[45]. While all marine

Synechococcus possess the three *kai* genes, most *Prochlorococcus* lack *kaiA* and, as a

consequence, their circadian clocks rather behave like an "hourglass" which is reset every

morning[46-48]. Our analysis indicated that *kaiA* was lost at SBE-LCA (Fig. 1A). This is

likely due to the prolonged darkness or low light conditions during the Snowball Earth,

rendering the sophisticated circadian clock dispensable.

We argue that the genome reduction and metabolic adaptation events discussed

above not only enabled *Prochlorococcus* to survive the Snowball Earth climate

catastrophe, but also shaped the physiological characters and the biogeographic

distribution of their descendants in the modern ocean. For example, the genome reduction

that occurred in the early evolution of *Prochlorococcus* likely resulted in the reduced cell

size and increased surface-to-volume ratio in their descendants, which may have

enhanced their efficiency in nutrient acquisition[49] and eventually led them to dominate

the photosynthetic communities in the most oligotrophic regions of today's oceans[2].

Likewise, new metabolic strategies that *Prochlorococcus* evolved to overcome the

nutrient stresses during Snowball Earth, such as the recycling of cell wall components

and the use of GG and GGA instead of nitrogen-rich GB as the organic osmolytes,

decreased the nutrient requirements of the descendants' cells and thus contributed to their

success in the modern oligotrophic nitrogen-limited oceans. On the other hand,

modifications of some important metabolic pathways may also have imposed deleterious

effects on *Prochlorococcus* descendants. For example, whereas the replacement of

circadian clock with an hourglass-like mechanism might have facilitated the ancestral

lineage to adapt to the prolonged dim light condition during the Snowball Earth

catastrophe, it likely prevents the dispersal of *Prochlorococcus* to high latitude regions in

the modern ocean, where the day length varies substantially across seasons. Normally,

organisms with circadian rhythms deal with these changes by anticipating the changes of

light intensity and promptly regulating cellular processes such as DNA transcription and

recombination via chromosome compaction, a known mechanism to protect DNA from

UV radiation[50,51]. In the absence of the circadian clock, however, species such as

Prochlorococcus cannot synchronize the endogenous oscillation with the environmental

cycles and thus are under high risks of cell damages[52].

Concluding Remarks

The Neoproterozoic Snowball Earth hypothesis was proposed decades ago, which

claimed the entire extinction of the photosynthetic organisms[17]. In contrast to the original

"hard" version of the hypothesis, a modified "soft" version of the Snowball Earth

hypothesis was later proposed to include the likely persistence of refugia across the

Cryogenian Period, which allowed for the survival of bacterial and simple eukaryotic

lineages[53,54]. Survivors of the Snowball Earth included photosynthetic

microorganisms[55,56], which enabled continuous primary production across the

interval[53,57]. Like other autotrophic organisms at the base of a food web, the survival of

Prochlorococcus was likely important in sustaining primary production, heterotrophy and

carbon cycling, as well as broader ecosystem functioning during the Snowball Earth

glaciations[53].

On the other extreme, a few studies have proposed that microbial communities might

have been only mildly affected by the Snowball Earth climate catastrophe[56,57]. These

inferences were based on the microfossil and biomarker records, which, due to the lack of

lineage-specificity, did not capture the nuances required to reconstruct effects on many

ecologically important lineages like the *Prochlorococcus* studied here. Instead, we find

that substantial disruptions to the Earth system, like the Neoproterozoic Snowball Earth,

leave indelible signatures in microbial genomes, such that these heritable changes allow

us to reconstruct interactions between environmental change and biological evolution

deep in Earth's history. By employing the accelerated genome-wide accumulation of the

deleterious type mutations as a proxy for a rapid decrease in the population size of

ancient lineages, we uncovered severe bottlenecks that shaped the early evolution of

Prochlorococcus lineages. The precise molecular clock analyses as well as the ancestral

genome reconstruction, furthermore, enabled us to link dynamics in ancestral population

sizes to changes in metabolic potential and adaptation to icehouse climates through

natural selection. Collectively, our findings demonstrate how paleomicrontological

approaches can be used to connect large-scale dynamics in the Earth System to the

genomic imprints left on extant microorganisms, which shape their ecological role and

biogeographic distribution in the world today. They also illustrate how *Prochlorococci*

acted as important "guardians of metabolism"[12], safeguarding photosynthetic metabolic

potential across the Snowball Earth climate catastrophe.

Materials and Methods

Genomic sequences of Cyanobacteria were downloaded from public databases and

manually annotated (Table S2; see Section 1 in Supplemental Methods). Divergence time

of *Prochlorococcus* was estimated with MCMCTree v4.9e[58] on top of 27 genes (Table

S3) previously proposed to be valuable to date bacterial divergence[59] and *Cyanobacteria*

phylogenomic trees. In previous studies, the LPP (*Leptolyngbya*, *Plectonema* and

Phormidium) group of *Cyanobacteria* located either at the basal of the

Microcyanobacteria group[14,60] or at the basal of the Macrocyanobacteria group[61,62]. Our

analysis showed that this controversy is likely caused by the inclusion of composition-

heterogeneous proteins, and that using composition-homogeneous proteins led to

consistent support for the former hypothesis (Fig. S4; Table S4; see Section 2.2 in

Supplemental Methods). Since molecular dating analysis is known to be intrinsically

associated with calibration points[63], we summarized the calibrations of Cyanobacteria

used in previous studies, and modified them for our own analyses with caution (Table S1;

see Section 2.3 in Supplemental Methods for justification). We further assessed the

fitness of different molecular clock models implemented in MCMCTree by using the

package "mcmc3r" v0.3.2, based on which we decided to use the independent rates

model for further molecular clock analyses. For each molecular clock analysis, the

software ran twice with a burn-in of 50,000 and a total of 500,000 generations. The

convergence was assessed based on the correlations of posterior mean time of all

ancestral nodes between independent runs (Fig. S5). By implementing statistical tests

based on the "infinite-site" theory (Fig. S3; see Section 2.6 in Supplemental Methods) we

were able to select the most precise estimates of *Prochlorococcus* evolutionary timeline

for illustration (Fig. S6) and further discussion.

Evolution of genome content via gene gains and losses was inferred using two

independent methods, AnGST[64] and BadiRate v1.35[65]. The former assumes that the

statistically supported topological differences between a gene tree and the species tree

result from evolutionary events (gene loss, gene duplication, HGT, gene birth and

speciation), and infers these evolutionary events by reconciling the topological

incongruences under a generalized parsimony framework by achieving a minimum

number of the evolutionary events along the species tree, with penalties of an

evolutionary event determined by the genome flux analysis (Fig. S7)[64]. The latter does

not rely on the tree topological incongruence information, but instead uses a full

maximum-likelihood approach to determine the gene family turnover rates that

maximizes the probability of observing the gene count patterns provided by the family

size table. The BadiRate analyses were run using nine strategies each with a distinct

turnover rate model and a distinct branch model. The likelihoods of different runs were

compared (Fig. S8), and three strategies with highest likelihood values were used (Fig.

S9). Further, results derived from AnGST (Fig. S10) and BadiRate were compared and

summarized to determine the common patterns shared by the two software (Fig. S11),

and important functional genes discussed were consistently inferred by these two

methods. As the two methods inferred the qualitatively same pattern of genome size

reduction on the branches leading to SBE-LCA, the number of gene gains and losses

derived from the AnGST analysis was presented.

The inference of a potential change of selection efficiency on a given branch was

performed by comparing the genome-wide d_R/d_C value across single-copy orthologous

genes of the branch to that of the closest sister branch. The d_R/d_C value was calculated

using RCCalculator (http://www.geneorder.org/RCCalculator/; see Section 4 in

Supplemental Methods) based on two independent amino acid classification schemes

(Table S5).

References

Biller, S. J., Berube, P. M., Lindell, D. & Chisholm, S. W. Prochlorococcus: the structure and function of collective diversity. *Nat Rev Microbiol* **13**, 13 (2015).

Johnson, Z. I. *et al.* Niche partitioning among Prochlorococcus ecotypes along ocean-scale environmental gradients. *Science (80-)* **311**, 1737-1740 (2006).

West, N. J. & Scanlan, D. J. Niche-partitioning of Prochlorococcus populations in a stratified water column in the eastern North Atlantic Ocean. *Appl Environ Microbiol* **65**, 2585-2591, doi:10.1128/aem.65.6.2585-2591.1999 (1999).

Hess, W. R. *et al.* The photosynthetic apparatus of Prochlorococcus: insights through comparative genomics. *Photosynth Res* **70**, 53-71 (2001).

Moore, L. R., Rocap, G. & Chisholm, S. W. Physiology and molecular phylogeny of coexisting Prochlorococcus ecotypes. *Nature* **393**, 464 (1998).

Batut, B., Knibbe, C., Marais, G. & Daubin, V. Reductive genome evolution at both ends of the bacterial population size spectrum. *Nat Rev Microbiol* **12**, 841 (2014).

Luo, H., Friedman, R., Tang, J. & Hughes, A. L. Genome reduction by deletion of paralogs in the marine cyanobacterium Prochlorococcus. *Mol Biol Evol* **28**, 2751-2760 (2011).

Hu, J. & Blanchard, J. L. Environmental sequence data from the Sargasso Sea reveal that the characteristics of genome reduction in Prochlorococcus are not a harbinger for an escalation in genetic drift. *Mol Biol Evol* **26**, 5-13 (2008).

Luo, H., Huang, Y., Stepanauskas, R. & Tang, J. Excess of non-conservative amino acid changes in marine bacterioplankton lineages with reduced genomes. *Nat Microbiol* **2**, 17091 (2017).

Zuckerkandl, E. & Pauling, L. in *Evolving genes and proteins* 97-166 (Elsevier, 1965).

Dayhoff, M. O. A model of evolutionary change in proteins. *Atlas of protein sequence and structure* **5**, 89-99 (1972).

Falkowski, P. G., Fenchel, T. & Delong, E. F. The microbial engines that drive Earth's biogeochemical cycles. *Science (80-)* **320**, 1034-1039 (2008).

Kettler, G. C. *et al.* Patterns and implications of gene gain and loss in the evolution of Prochlorococcus. *PLoS Genet* **3**, e231 (2007).

Sánchez-Baracaldo, P. Origin of marine planktonic cyanobacteria. *Sci Rep* **5**, 17418 (2015).

Sánchez-Baracaldo, P., Ridgwell, A. & Raven, J. A. A neoproterozoic transition in the marine nitrogen cycle. *Current Biology* **24**, 652-657 (2014).

Marshall, C. R. Using the Fossil Record to Evaluate Timetree Timescales. *Front Genet* **10**, 1049, doi:10.3389/fgene.2019.01049 (2019).

Hoffman, P. F., Kaufman, A. J., Halverson, G. P. & Schrag, D. P. A Neoproterozoic snowball earth. *Science (80-)* **281**, 1342-1346 (1998).

Ashkenazy, Y. *et al.* Dynamics of a Snowball Earth ocean. *Nature* **495**, 90 (2013).

Zinser, E. R. *et al.* Influence of light and temperature on Prochlorococcus ecotype distributions in the Atlantic Ocean. *Limnol Oceanogr* **52**, 2205-2220 (2007).

Thomas, D. & Dieckmann, G. Antarctic sea ice--a habitat for extremophiles.

391   *Science (80-)* **295**, 641-644 (2002).

392 21 Hoffman, P. F. *et al.* Snowball Earth climate dynamics and Cryogenian geology-
393   geobiology. *Sci Adv* **3**, e1600983 (2017).

394 22 Takeuchi, N. Optical characteristics of cryoconite (surface dust) on glaciers: the
395   relationship between light absorbency and the property of organic matter
396   contained in the cryoconite. *Annals of Glaciology* **34**, 409-414 (2002).

397 23 Benforte, F. C. *et al.* Novel role of the LPS core glycosyltransferase WapH for
398   cold adaptation in the Antarctic bacterium Pseudomonas extremaustralis. *PLoS*
399   *One* **13**, e0192559 (2018).

400 24 Wang, Z., Wang, J., Ren, G., Li, Y. & Wang, X. Influence of core oligosaccharide
401   of lipopolysaccharide to outer membrane behavior of Escherichia coli. *Mar Drugs*
402   **13**, 3325-3339 (2015).

403 25 Feller, G. & Gerday, C. Psychrophilic enzymes: hot topics in cold adaptation. *Nat*
404   *Rev Microbiol* **1**, 200 (2003).

405 26 Uhde, A. *et al.* Glucosamine as carbon source for amino acid-producing
406   Corynebacterium glutamicum. *Appl Microbiol Biotechnol* **97**, 1679-1687 (2013).

407 27 Park, J. T. & Uehara, T. How Bacteria Consume Their Own Exoskeletons
408   (Turnover and Recycling of Cell Wall Peptidoglycan). *Microbiology and*
409   *Molecular Biology Reviews* **72**, 211-227, doi:10.1128/mmbr.00027-07 (2008).

410 28 Borisova, M. *et al.* Peptidoglycan Recycling in Gram-Positive Bacteria Is Crucial
411   for Survival in Stationary Phase. *MBio* **7**, doi:10.1128/mBio.00923-16 (2016).

412 29 Schumann, W. Regulation of bacterial heat shock stimulons. *Cell Stress and*
413   *Chaperones* **21**, 959-968 (2016).

414 30 La Terza, A., Papa, G., Miceli, C. & Luporini, P. Divergence between two
415   Antarctic species of the ciliate Euplotes, E. focardii and E. nobilii, in the
416   expression of heat-shock protein 70 genes. *Mol Ecol* **10**, 1061-1067,
417   doi:10.1046/j.1365-294x.2001.01242.x (2001).

418 31 Rinehart, J. P. *et al.* Continuous up-regulation of heat shock proteins in larvae, but
419   not adults, of a polar insect. *Proc Natl Acad Sci U S A* **103**, 14223-14227,
420   doi:10.1073/pnas.0606840103 (2006).

421 32 Lawrence, R. P. & William, J. W. Temperature and substrates as interactive
422   limiting factors for marine heterotrophic bacteria. *Aquatic Microbial Ecology* **23**,
423   187-204 (2001).

424 33 Gleitz, M., v.d. Loeff, M. R., Thomas, D. N., Dieckmann, G. S. & Millero, F. J.
425   Comparison of summer and winter inorganic carbon, oxygen and nutrient
426   concentrations in Antarctic sea ice brine. *Mar Chem* **51**, 81-91, (1995).

427 34 Reay, D. S., Nedwell, D. B., Priddle, J. & Ellis-Evans, J. C. Temperature
428   dependence of inorganic nitrogen uptake: reduced affinity for nitrate at
429   suboptimal temperatures in both algae and bacteria. *Appl. Environ. Microbiol.* **65**,
430   2577-2584 (1999).

431 35 Raven, J. A., Wollenweber, B. & Handley, L. L. A comparison of ammonium and
432   nitrate as nitrogen sources for photolithotrophs. *New Phytologist* **121**, 19-32
433   (1992).

434 36 Webster-Brown, J. G., Hawes, I., Jungblut, A. D., Wood, S. A. & Christenson, H.
435   K. The effects of entombment on water chemistry and bacterial assemblages in

436   closed cryoconite holes on Antarctic glaciers. *FEMS Microbiol Ecol* **91** (2015).

437   37   Mao, X. *et al.* Computational prediction of the osmoregulation network in
438   Synechococcus sp. WH8102. *BMC Genomics* **11**, 291, doi:10.1186/1471-2164-
439   11-291 (2010).

440   38   Scanlan, D. J. *et al.* Ecological genomics of marine picocyanobacteria. *Microbiol.*
441   *Mol. Biol. Rev.* **73**, 249-299 (2009).

442   39   Empadinhas, N. & da Costa, M. S. To be or not to be a compatible solute:
443   bioversatility of mannosylglycerate and glucosylglycerate. *Syst Appl Microbiol*
444   **31**, 159-168 (2008).

445   40   Carver, J. H. & Vardavas, I. M. Precambrian glaciations and the evolution of the
446   atmosphere. *Ann Geophys* **12**, 674-682, doi:10.1007/s00585-994-0674-3 (1994).

447   41   Raven, J. A., Kübler, J. & Beardall, J. Put out the light, and then put out the light.
448   *Journal of the Marine Biological Association of the United Kingdom* **80**, 1-25,
449   doi:10.1017/s0025315499001526 (2000).

450   42   Simon, C., Wiezer, A., Strittmatter, A. W. & Daniel, R. Phylogenetic diversity and
451   metabolic potential revealed in a glacier ice metagenome. *Appl Environ Microbiol*
452   **75**, 7519-7526, doi:10.1128/aem.00946-09 (2009).

453   43   Kouřil, R., Wientjes, E., Bultema, J. B., Croce, R. & Boekema, E. J. High-light vs.
454   low-light: effect of light acclimation on photosystem II composition and
455   organization in Arabidopsis thaliana. *Biochimica et Biophysica Acta (BBA)-*
456   *Bioenergetics* **1827**, 411-419 (2013).

457   44   Bibby, T., Mary, I., Nield, J., Partensky, F. & Barber, J. Low-light-adapted
458   Prochlorococcus species possess specific antennae for each photosystem. *Nature*
459   **424**, 1051 (2003).

460   45   Dong, G. & Golden, S. S. How a cyanobacterium tells time. *Curr Opin Microbiol*
461   **11**, 541-546 (2008).

462   46   Holtzendorff, J. *et al.* Genome streamlining results in loss of robustness of the
463   circadian clock in the marine cyanobacterium Prochlorococcus marinus PCC
464   9511. *J Biol Rhythms* **23**, 187-199, doi:10.1177/0748730408316040 (2008).

465   47   Axmann, I. M. *et al.* Biochemical evidence for a timing mechanism in
466   prochlorococcus. *J Bacteriol* **191**, 5342-5347, doi:10.1128/jb.00419-09 (2009).

467   48   Mella-Flores, D. *et al.* Prochlorococcus and Synechococcus have Evolved
468   Different Adaptive Mechanisms to Cope with Light and UV Stress. *Front*
469   *Microbiol* **3**, 285, doi:10.3389/fmicb.2012.00285 (2012).

470   49   Giovannoni, S. J., Thrash, J. C. & Temperton, B. Implications of streamlining
471   theory for microbial ecology. *ISME J* **8**, 1553 (2014).

472   50   Warters, R. L. & Lyons, B. W. Variation in radiation-induced formation of DNA
473   double-strand breaks as a function of chromatin structure. *Radiat Res* **130**, 309-
474   318 (1992).

475   51   Simons, M. J. The evolution of the cyanobacterial posttranslational clock from a
476   primitive "phoscillator". *J Biol Rhythms* **24**, 175-182 (2009).

477   52   Mullineaux, C. W. & Stanewsky, R. The rolex and the hourglass: a simplified
478   circadian clock in Prochlorococcus? *J Bacteriol* **191**, 5333-5335 (2009).

479   53   Moczydłowska, M. The Ediacaran microbiota and the survival of Snowball Earth
480   conditions. *Precambrian Res* **167**, 1-15 (2008).

54 Hoffman, P. F. & Schrag, D. P. The snowball Earth hypothesis: testing the limits
of global change. *Terra nova* **14**, 129-155 (2002).
55 Allison, C. W. & Awramik, S. M. Organic-walled microfossils from earliest
Cambrian or latest Proterozoic Tindir Group rocks, northwest Canada.
Precambrian Res **43**, 253-294 (1989).
56 Corsetti, F. A., Awramik, S. M. & Pierce, D. A complex microbiota from snowball
Earth times: microfossils from the Neoproterozoic Kingston Peak Formation,
Death Valley, USA. *Proc Natl Acad Sci USA* **100**, 4399-4404 (2003).
57 Olcott, A. N., Sessions, A. L., Corsetti, F. A., Kaufman, A. J. & De Oliviera, T. F.
Biomarker evidence for photosynthesis during Neoproterozoic glaciation. *Science
(80-)* **310**, 471-474 (2005).
58 Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol*
24, 1586-1591 (2007).
59 Battistuzzi, F. U. & Hedges, S. B. A major clade of prokaryotes with ancient
adaptations to life on land. *Mol Biol Evol* **26**, 335-343 (2008).
60 Shih, P. M. *et al.* Improving the coverage of the cyanobacterial phylum using
diversity-driven genome sequencing. *Proc Natl Acad Sci USA* **110**, 1053-1058
(2013).
61 Blank, C. & Sanchez-Baracaldo, P. Timing of morphological and ecological
innovations in the cyanobacteria–a key to understanding the rise in atmospheric
oxygen. *Geobiology* **8**, 1-23 (2010).
62 Uyeda, J. C., Harmon, L. J. & Blank, C. E. A comprehensive study of
cyanobacterial morphological and ecological evolutionary dynamics through deep
geologic time. *PLoS One* **11**, e0162539 (2016).
63 Schirrmeister, B. E., Sanchez-Baracaldo, P. & Wacey, D. Cyanobacterial evolution
during the Precambrian. *Int J Astrobiol* **15**, 187-204 (2016).
64 David, L. A. & Alm, E. J. Rapid evolutionary innovation during an Archaean
genetic expansion. *Nature* **469**, 93 (2011).
65 Librado, P., Vieira, F. & Rozas, J. BadiRate: estimating family turnover rates by
likelihood-based methods. *Bioinformatics* **28**, 279-281 (2011).
511
512

Acknowledgments

We thank Allison Coe, Erik Zinser, and Zackary Johnson for providing the data of

Prochlorococcus growth rates, Sishuo Wang, Tianhua Liao and Xiaoyuan Feng for their

suggestions on molecular dating analyses. This work is supported by the National Natural

Science Foundation of China (92051113), the Hong Kong Research Grants Council

General Research Fund (14110820), the Hong Kong Research Grants Council Area of

Excellence Scheme (AoE/M-403/16), HKU FoS funds to SAC, and the Direct Grant of

CUHK (4053257 and 3132809).

Author Contributions

522   H.L. conceived and directed the study. H.Z. and Y.S. performed the

523   bioinformatics. All authors contribute to the interpretation of the results. H.Z., Y.S.,

524   S.A.C and H.L. wrote the paper.

Conflict of interests

526   The author declares no competing interests.

Fig. 1

A full-page scientific figure (Fig. 1) comprising four panels (A–D).

A — A time-calibrated phylogenetic tree of *Prochlorococcus* and *Synechococcus* strains plotted against a geological time axis (Mya), with gene presence/absence dot-matrix for genes grouped under "Low temperature", "Osmotic pressure" and "Low light". Clades are labeled HLII, HLI, LLI, LLII/III, LLIV, Syne 5.1, Syne 5.2. A pie chart legend indicates Replacement, Loss, Duplication, Gain, Vertical inheritance. SBE-LCA 682 Mya (632-732 Mya). Geological periods: Meso (S), Neo (T, C, E), P (C, O, S, D, C, P), M (T, J, C), C. Sturtian glaciation and Marinoan glaciation are marked.

Gene labels (left to right): *desA, desC, kdsA, kdsB, kdsC, kdsD, lpxA, lpxB, lpxC, lpxD, rfbA, rfbB, rfbC, nagB, glmU, glmM, hrcA, nlrM, glnA, gltS, bsmB, gsmt, proV, proW, proX, proP, sspS, spgS, psaM, psbY, psbF, psbX, psbU, psbX, pcbD, kaiA, kaiB, kaiC*

Strains (top to bottom): MIT 9314, SB, AS9601, MIT 9301, GP2, MIT 0604, MIT 9311, MIT 9312, MIT 9302, MIT 9321, MIT 9401, MIT 9322, MIT 9201, MIT 9202, MIT 9215, MIT 9116, MIT 9123, MIT 9107, EQPAC1, MED4, MIT 9515, NATL2A, PAC1, NATL1A, MIT 0801, LG, SS51, SS2, SS52, SS120, SS35, MIT 0602, MIT 0603, MIT 9211, MIT 0601, MIT 0702, MIT 0703, MIT 0701, MIT 9303, MIT 9313, CC9605, WH 8109, KORDI-52, BL107, CC9902, WH 8102, KORDI-49, CC9616, KORDI-100, CC9311, WH 8016, WH 7803, WH 7805, RS9916, RS9917, PCC 7001, GFB01, CB0101, CB0205, PCC 6307, WH 5701

Time axis: 1000, 800, 600, 400, 200, 0 Mya

B — Cladogram (HLII, HLI, LLI, LLII/III, LLIV, Syne 5.1) with plots of "Charge" and "Volume and Polarity" for Target vs Control. Legend: Uncorrected (grey), GC-corrected (AA frequency) (red), GC-corrected (codon frequency) (cyan).

C — Average growth rates of *Prochlorococcus* at different temperatures. Y axis: Average growth rate (day⁻¹), 0.0 to 0.6. X axis: Temperature (°C), 10 to 30. Ecotype: HL (red), LL (blue). Clade symbols: HLI-1 (square), HLI-2 (circle), HLII-1 (triangle), HLII-2 (plus), HLII-3 (cross), HLII-4 (diamond), LLI (inverted triangle), LLIV (asterisk).

D — Schematic cross-section of an ice sheet environment showing Accumulation Zone, Sublimation Zone, Solar Radiation, Ice Sheet, Sea-ice brine channels, Cryoconite holes, Ground, Melt water, Brine, and a cryoconite hole detail with Refrozen layer, Melt water, Cryoconite, Solar Radiation. Legend: LL *Prochlorococcus*.

Fig. 1   Evolution of *Prochlorococcus* during the Neoproterozoic Snowball Earth events.

(A) (Left) Chronogram of the evolutionary history of *Prochlorococcus* estimated by

MCMCTree. The evolutionary tree shown here is part of the species tree constructed with

MrBayes based on 90 compositionally homogenous gene families shared by 159

cyanobacterial genomes (Fig. S4 D). Divergence time is estimated based on 27 gene

sequences under calibration set C9 (Table S1). The vertical bars represent the estimated

time of the Neoproterozoic glaciation events. The flanking horizontal blue bars on

ancestral nodes represent the posterior 95% highest probability density (HPD) interval of

the estimated divergence time. The pie chart on the ancestral branches leading to the node

SBE-LCA provides the proportion of reconstructed genomic events including gene gain,

gene loss, gene replacement, gene duplication and gene vertical inheritance. (Right)

Phyletic pattern of key gene families that potentially enabled *Prochlorococcus* to survive

harsh conditions during the Neoproterozoic Snowball Earth (at the ancestral node 'SBE-

LCA'). Solid square, solid circle and open circle next to each extant taxon represent

multi-copy gene family, single-copy gene family, and absence of the gene family,

respectively, in the genome. (B) (Left) The diagram helps understand how the d_R/d_C was

calculated. In this context, the 'Target' group includes all genomes of all HL clades, LLI

and LLII/III, the 'Control' group includes all genomes of LLIV, and the 'reference'

group includes all genomes of Syne 5.1. The d_R/d_C for the 'Target' group (shown in

Middle & Right) is calculated by comparing a genome from the 'Target' group to a

genome from the 'reference' group (marked with red), followed by averaging the value

across all possible genome pairs. Likewise, the d_R/d_C for the 'Control' group (shown in

Middle & Right) is calculated by comparing a genome from the 'Control' group to a

genome from the 'reference' group (marked with green) and then by averaging the value

across all possible genome pairs. (Middle & Right) The genome-wide means of d_R/d_C

values at the ancestral branch leading to SBE-LCA and that at its sister lineage. They

were classified based on the physicochemical classification of the amino acids by charge

or by volume and polarity, and were either GC-corrected by codon frequency (blue), GC-

corrected by amino acid (AA) frequency (red) or uncorrected (gray). Error bars of d_R/d_C

values represent the standard error of the mean. (C) Diagram of putative bacterial refugia

including cryoconite holes and sea ice brine channels in Neoproterozoic Snowball Earth,

which were featured with a number of stresses such as low temperature, low light, and

variable osmotic pressure. (D) The average growth rate of *Prochlorococcus* ecotypes at

different temperatures. Replicate cell cultures were grown in a 14:10 light: dark cycle at

66 ±1 μmol m^{-2}s^{-1}. The growth data used for plotting are collected from Johnson et al.

2006 and Zinser et al. 2007.