Snowball Earths, population bottlenecks, and *Prochlorococcus* evolution

Hao Zhang[1,^], Ying Sun[1,^], Qinglu Zeng[2], Sean A. Crowe[3], Haiwei Luo[1*]

[1]Simon F. S. Li Marine Science Laboratory, School of Life Sciences and State Key Laboratory of Agrobiotechnology, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[2]Department of Ocean Science, The Hong Kong University of Science and Technology, Clear Water Bay, Hong Kong SAR

[3]Department of Earth Sciences, School of Biological Sciences, and Swire Institute for Marine Science (SWIMS), University of Hong Kong, Pokfulam Road, Hong Kong SAR

[^]These authors contributed equally to this study.

*Corresponding author: Haiwei Luo

Email: hluo2006@gmail.com

Competing Interest Statement: The authors declare no competing interests.

Keywords: *Prochlorococcus*, Neoproterozoic Snowball Earth, genome reduction, molecular dating.

Abstract

Prochlorococcus are the most abundant photosynthetic organisms in the modern ocean. A massive DNA loss event occurred in their early evolutionary history, leading to highly reduced genomes in nearly all lineages, as well as enhanced efficiency in both nutrient uptake and light absorption. The environmental landscape that shaped this ancient genome reduction, however, remained unknown. Through careful molecular clock analyses, we established that this *Prochlorococcus* genome reduction occurred during the Neoproterozoic Snowball Earth climate catastrophe. The lethally low temperature and exceedingly dim light during the Snowball Earth event would have inhibited *Prochlorococcus* growth and proliferation and caused severe population bottlenecks. These bottlenecks are recorded as an excess of deleterious mutations that accumulated across genomic regions in the descendant lineages. *Prochlorococcus* adaptation to extreme environmental conditions during Snowball Earth intervals can be inferred by tracing the evolutionary paths of genes that encode key metabolic potential. This metabolic potential includes modified lipopolysaccharide structure, strengthened peptidoglycan biosynthesis, the replacement of a sophisticated circadian clock with an hourglass-like mechanism that resets daily for dim light adaption, and the adoption of ammonia diffusion as an efficient membrane transporter-independent mode of nitrogen acquisition. In this way, the Neoproterozoic Snowball Earth event altered the physiological characters of *Prochlorococcus*, shaping their ecologically vital role as the most abundant primary producers in the modern oceans.

Introduction

Prochlorococcus are the smallest and most abundant photosynthetic organisms on Earth (1). They are prevalent throughout the photic zone of the oligotrophic oceans between 40 °N and 40 °S (1), where they account for more than 40% of the biomass and contribute almost half of the net primary production (2). *Prochlorococcus* have diversified into two major phylogenetic groups with distinct ecology (ecotypes), with the high-light (HL) adapted monophyletic group imbedded in the low-light (LL) adapted paraphyletic group (3). The distinct ecotypes of *Prochlorococcus* evolved different pigments, light-harvesting systems, and phycobiliproteins, which allowed for efficient light absorption in the water column (4), and thus increased growth rates and primary production (5).

Prochlorococcus genomes have been shaped by stepwise streamlining, including a major genome reduction in their early evolution and a few minor modifications that followed (6, 7). Modern marine *Prochlorococcus* lineages, in particular those with small genomes, show very low ratios of nonsynonymous (d_N) to synonymous (d_S) nucleotide substitution rates, suggesting that natural selection is a highly efficient throttle on the accumulation of deleterious mutations (i.e., nonsynonymous mutations) in *Prochlorococcus* (6, 8). On long time scales, however, nucleotide substitutions at synonymous sites become saturated, invalidating the use of d_N/d_S to infer selection efficiency in deep time (9). Thus, an alternative approach focuses instead on different types of nonsynonymous substitutions leading to radical versus conservative changes in amino acid sequences, with the former more likely to be deleterious (10, 11). Excess radical mutations accumulate from random fixations of deleterious mutations by genetic drift (i.e., reduced efficiency of selection). Using this approach reveals that the major genome reduction in

Prochlorococcus took place under reduced selection efficiency (9) and implies that the ancient

population went through severe bottlenecks as the likely result of environmental catastrophe.

The environmental context underlying *Prochlorococcus* genome reduction remains

unknown, however, and precise molecular dating is needed to link this important evolutionary

event to its possible environmental drivers. By implementing comprehensive molecular clock

analyses, we now link the early major genome reduction event of *Prochlorococcus* to the

Neoproterozoic Snowball Earth events. These catastrophic disruptions to the Earth system would

likely have challenged warm-water-loving photosynthetic *Prochlorococcus*, with strong potential

to cause the population bottlenecks inferred from the genome sequences described above.

Prochlorococcus survived this catastrophe through likely gains and losses of key metabolic

functions reconstructed from the same genome sequences, which have a far-reaching impact on

their success in today's oceans. *Prochlorococcus* are thus vital "guardians of metabolism" (12),

shepherding genes critical to the functioning of the biosphere across environmental catastrophes,

including global glaciations.

Results and Discussion

Prochlorococcus experienced a massive gene loss event on the ancestral branch leading to

the last common ancestor (LCA) of clades HL, LLI, and LLII/III (6, 7, 13). This is confirmed by

our analysis, which reconstructed 366 and 107 gene family losses and gains on this branch,

respectively (Fig. 1A & S1). On the same ancestral branch, it was shown that d_R/d_C is

significantly elevated compared to the sister ancestral branch leading to the LCA of the LLIV

(9), which was validated here (both sign test and paired *t*-test, $p < 0.001$; Fig. 1B). These results

confirm that the major genome reduction event occurring on this branch was likely driven by

genetic drift as a result of one or recurrent population bottlenecks (9). Given the global

distribution and abundance of *Prochlorococcus*, and cyanobacteria more generally, such a

bottleneck would likely require a global-scale event, like an environmental or climate catastrophe

(e.g. meteorite impact, large igneous province emplacement, or glaciation).

To establish the environmental context for the large, ancient genome reduction, we estimated

the timeline of *Prochlorococcus* evolution by implementing molecular clock analyses based on

essential calibrations available in the cyanobacterial lineage. We recognize that the use of

calibration sets adapted from previous studies (under calibrations C1-C8 in Table S1 with related

references included there) results in up to ~320 Ma disparity (Fig. S2A) in the estimated time for

the LCA of *Prochlorococcus* HL, LLI and LLII/III clades that emerged with the major genome

reduction. We note that the calibrations in previous studies were not properly used. For example,

the akinete fossil identified to 2,100 Mya was used as either the maximum bound or the

minimum bound to calibrate the crown group of Nostocales (14, 15). However, given the fact

that apomorphic character must evolve earlier than the divergence of crown group,

morphological fossils can only serve as the minimum bounds on total groups of assigned

lineages (16) (see Section 2.3 in Supplemental Methods for details). Thus, in the present study,

we modified the calibration sets by constraining the lower bounds of the Nostocales (and the

Pleurocapsales) total groups with morphological fossils and by leaving their upper bounds open

(C9-C14; Table S1). Intriguingly, the variation is reduced to less than 10 Ma when these

modified calibration sets are used (Fig. S2A).

Recent identification of non-oxygenic Cyanobacteria lineages such as Melainabacteria and

Sericytochromatia as sister groups of oxygenic Cyanobacteria (17, 18) provides an alternative

way to constrain the evolution of oxygenic Cyanobacteria. Specifically, given that oxygenic

photosynthesis evolved at the stem lineage of oxygenic Cyanobacteria, we constrained the

minimum age of the total Cyanobacteria group at 3.0 Ga, which is supported by geochemical

evidence as the time when atmospheric oxygen became available (19, 20). To avoid the overly

precise and potentially misleading age estimates, we calibrated the upper limit of the

Cyanobacteria root using the ages when the planet Earth formed and became habitable (C15-C38

in Table S1; see Section 2.3 in Supplemental Methods for details). Using this strategy, we show

that the age of *Prochlorococcus* major genome reduction remains stable when non-oxygenic

Cyanobacteria outgroups were included (Fig. S2B). Since including the non-oxygenic

Cyanobacteria have consistently reduced the precision of posterior age estimates, manifested as

the higher slopes of the regression line between highest posterior density (HPD) width and the

posterior age estimates compared to those without including these lineages (C15-C38 versus C1-

C14 in Fig. S3; also see Section 2.6 in Supplemental Methods for an extended discussion), we

focus on the crown oxygenic Cyanobacteria group dating (C7-C14) in the following discussions.

By comparing the width of the 95% HPD derived from each molecular clock analysis (Fig.

S3), we inferred the most precise timeline of *Prochlorococcus* evolution (corresponding to the

calibration set C14 in Table S1; see Section 2.3 in Supplemental Methods). Our time estimates

revealed that the LCA of *Prochlorococcus* HL, LLI, and LLII/III clades diversified at 682 Mya

(95% HPD 632-732 Mya), precisely dating the genome reduction event to this time. A 682 Mya

date for the emergence of the LCA of *Prochlorococcus* HL, LLI, and LLII/III clades places the

large genome reduction that took place in this lineage firmly within the Cryogenian Period (~720

to 635 Mya; Fig. 1A) and implicates the Snowball Earth icehouse climate conditions eponymous

with the Period in the corresponding *Prochlorococcus* population bottleneck. We, therefore, refer

to this ancestor as SBE-LCA (see Fig. 1A), short for "Snowball Earth" LCA. The

Neoproterozoic climate catastrophe culminated in the Sturtian (~717 to 659 Mya) and Marinoan

(~645 to 635 Mya) glaciations (Fig. 1A), which stretched from the poles to sea level near the

equators, possibly wrapping the entire Earth under a frozen skin (21). This "Snowball Earth"

persisted with the freezing temperature of seawater below the ice sheet lowered to -3.5°C (22).

Since all assayed *Prochlorococcus* strains, including those affiliated with the basal LL ecotypes,

reach maximum growth rates at approximately 25°C and rarely survive when the temperature

drops to ~10°C (Fig. 1C) (23), we propose that extreme climate cooling during the

Neoproterozoic Snowball Earth events was likely the major driver of severe bottlenecks in early

Prochlorococcus populations.

Survival of *Prochlorococcus* populations through the Cryogenian would have required

refugia, the nature of which would have shaped continued *Prochlorococcus* evolution. A variety

of biotic refugia have been identified during Snowball Earth intervals, including the sea-ice brine

channels within ice grounding-line crack systems (24) and cryoconite holes/ponds on the surface

of the sublimation zone, which may have represented ~12% of the global sea glacier surface (25)

(Fig. 1D). Despite providing the essential space for *Prochlorococcus* survival, these refugia

would have presented a number of environmental stresses to *Prochlorococcus* populations such

as low temperature, dim light, and limited nutrients (24-26). *Prochlorococcus* thus evolved a

number of adaptive mechanisms to cope with these stresses via gene gains and losses, which we

assessed by reconstructing the evolutionary paths of imprints that the Snowball Earth climate left

in extant *Prochlorococcus* genomes.

Among these stresses, the most prominent was likely lethally low temperature. Maintaining

membrane fluidity is of paramount importance under low-temperature conditions, which is

largely achieved by the activities of fatty acid desaturase encoded by *desA* and *desC*. As a result,

we inferred that these genes were retained in SBE-LCA (Fig. 1A). Lipopolysaccharide (LPS) in

the outer membrane is known to provide the first line of defense against harsh environments

(27), which contains the O-specific polysaccharide, the glycolipid anchor lipid A, and the

polysaccharide core region. Based on our analyses, genes encoding the polysaccharide core

region (*kdsABCD* for 3-deoxy-d-manno-octulosonate biosynthesis; Fig. 1A) were likely lost at

SBE-LCA, while those encoding the other components were retained (*lpxABCD* and *rfbABC* for

Lipid A precursor and O-specific LPS precursor biosynthesis; Fig. 1A). This inference is

consistent with a previous conclusion that the loss of the LPS core region would increase the

hydrophobicity and permeability of the cell envelope (28) to protect against cold conditions (29).

Another metabolic modification in SBE-LCA was related to heat shock proteins (HSPs), which

play crucial roles in tolerating environmental stresses including thermal shocks. Typically, HSPs

are tightly regulated, as they respond quickly to stress and turn off rapidly once the stress

disappears (30). However, the HSP repressor protein encoded by *hrcA* was inferred to be lost at

SBE-LCA, which thus likely allowed the organism to continuously express HSPs to cope with

prolonged lethally low temperature. In fact, constitutive expression of HSPs occurs in polar

organisms such as the Antarctic ciliate *Euplotes focardii* (31) and the polar insect *Belgica*

antarctica larvae (32). Extremely low temperature also made substrate acquisition difficult due

to increased lipid stiffness and decreased efficiency and affinity of membrane transporters (33).

Under such conditions, bacteria may increasingly rely on substrates whose uptake shows lower

dependence on temperature. In sea-ice brines where less CO_2 is dissolved (34), elevated pH

promotes the conversion of ammonium to ammonia, which diffuses directly into cells without

the aid of transporters in the membrane. Accordingly, species of bacteria and microalgae show a

greater dependence on ammonium and ammonia at low temperatures and high pH than nitrate

(35, 36), thereby reducing reliance on membrane transporters. In SBE-LCA, the potentially

efficient utilization of ammonia made other N acquisition genes dispensable, leading to the

neutral loss of nitrite transporter (*nitM*), whereas glutamine synthetase (*glnA*) and glutamate

synthase (*gltS*) responsible for the utilization of ammonia after its assimilation were conserved

(Fig. 1A).

An additional stress to *Prochlorococcus* during Snowball Earths was likely the extremely

oligotrophic condition presented by bacteria refugia (37). Accordingly, SBE-LCA of

Prochlorococcus evolved a few metabolic strategies for their survival. The amino sugar N-

acetylglucosamine (GlcNAc) is used by bacteria such as *Corynebacterium glutamicum* as a

carbon, energy, and nitrogen source (38). GlcNAc enters bacteria in the form of GlcNAc-6-

phosphate (GlcNAc-6-P). However, instead of being metabolized, the loss of *nagB* for GlcNAc-

6-P deamination at SBE-LCA suggests that GlcN6P is more likely to be involved in

peptidoglycan (PG) recycling through the cascade catalysis by GlmM and GlmU (Fig. 1A) to

generate UDP-GlcNAc, which is an essential precursor of cell wall PG and LPS (39). During cell

turnovers, PG is continuously broken down and reused through the PG recycling pathway to

produce new PG, and in some bacteria, PG recycling is critical for their long-term survival when

growth is stalled under nutrient limitation (40). Thus, such a recycling mechanism seems to be

key for the maintenance of cell integrity in SBE-LCA under oligotrophic conditions (25).

Glycine betaine (GB) is known to be a ubiquitous protein-stabilizing osmolyte in bacteria, in

particular cyanobacteria (41). However, genes involved in glycine betaine biosynthesis and

transport were lost at SBE-LCA, including *bsmB* for dimethylglycine N-methyltransferase, *gsmt*

for glycine/sarcosine N-methyltransferase, and *proVWXP* for glycine betaine/proline transport

system (Fig. 1A). Instead, several other organic osmolytes might have been used during the

Snowball Earths, as their biosynthetic genes were retained at SBE-LCA. The first examples are

the *ggpS* gene encoding glucosylglycerol phosphate synthase for glucosylglycerol (GG)

synthesis and the *gpgS* encoding glucosyl-phosphoglycerate synthase for glucosylglycerate

(GGA) synthesis (Fig. 1A). Since the biosynthesis of GG and GGA requires less N compared to

that of GB (42, 43), the potential use of GG/GGA instead of GB appeared favorable to SBE-

LCA.

Low light intensity during Snowball Earth was another formidable challenge to phototrophs

including *Prochlorococcus*. In the Neoproterozoic, the Sun was still at least 6% dimmer than that

at present (44). Moreover, sea ice, especially when covered with snow, is an effective barrier to

light transmission (45). This is in analogy to the deeper layers of today's polar snow and glacier

ice where irradiation is reduced and photosynthetic organisms and activities are scarcely

detectable (46). *Prochlorococcus* may also need to compete with contemporary eukaryotic algae

for the limited amount of light, because the latter became increasingly abundant during the

Cryogenian glaciations and likely shared the same refugia with bacteria (37, 47). Consequently,

photosynthetic organisms trapped in bacterial refugia or inhabiting waters below ice need to be

physiologically geared to cope with low light. It was proposed that modification of the

photosystem structure enables adaptation to the low light condition (48). We inferred a few

changes in photosystem I and II (PSI/PSII) that occurred in SBE-LCA, including the gain of RC1

subunit PsaM, RC2 subunit PsbY, and an extra copy of the RC2 subunit PsbF, the loss of RC2

protein PsbU, and the replacement of RC2 subunit PsbX (Fig. 1A), but the molecular mechanism

of these changes underlying low light adaptation is poorly understood. We also inferred an

expansion of the *Prochlorococcus* antenna Pcb from two to six copies during the Snowball Earth

(Fig. 1A), which may boost the light-harvesting capacity under low-light conditions (49).

Many cyanobacteria have a sophisticated circadian clock, which is essential in controlling

the global diel transcriptional activities of the cells. This circadian oscillator system requires only

three components: KaiA, KaiB, and KaiC (50). While all marine *Synechococcus* possess the

three *kai* genes, most *Prochlorococcus* lack *kaiA* and, as a consequence, their circadian clocks

rather behave like an "hourglass" which is reset every morning (51-53). Our analysis indicated

that *kaiA* was lost at SBE-LCA (Fig. 1A). This is likely due to the prolonged darkness or low

light conditions during the Snowball Earth, rendering the sophisticated circadian clock

dispensable.

We argue that the genome reduction and metabolic adaptation events discussed above not

only enabled *Prochlorococcus* to survive the Snowball Earth climate catastrophe, but also shaped

the physiological characters and the biogeographic distribution of their descendants in the

modern ocean. For example, the genome reduction that occurred in the early evolution of

Prochlorococcus likely resulted in the reduced cell size and increased surface-to-volume ratio in

their descendants, which may have enhanced their efficiency in nutrient acquisition (54) and

eventually led them to dominate the photosynthetic communities in the most oligotrophic regions

of today's oceans (2). Likewise, new metabolic strategies that *Prochlorococcus* evolved to

overcome the nutrient stresses during Snowball Earth, such as the recycling of cell wall

components and the use of GG and GGA instead of nitrogen-rich GB as the organic osmolytes,

decreased the nutrient requirements of the descendants' cells and thus contributed to their

success in the modern oligotrophic nitrogen-limited oceans. On the other hand, modifications of

some important metabolic pathways may also have imposed deleterious effects on

Prochlorococcus descendants. For example, whereas the replacement of circadian clock with an

hourglass-like mechanism might have facilitated the ancestral lineage to adapt to the prolonged

dim light condition during the Snowball Earth catastrophe, it likely prevents the dispersal of

Prochlorococcus to high latitude regions in the modern ocean, where the day length varies

substantially across seasons. Normally, organisms with circadian rhythms deal with these

changes by anticipating the changes of light intensity and promptly regulating cellular processes

such as DNA transcription and recombination via chromosome compaction, a known mechanism

to protect DNA from UV radiation (55, 56). In the absence of the circadian clock, however,

species such as *Prochlorococcus* cannot synchronize the endogenous oscillation with the

environmental cycles and thus are at high risks of cell damages (57).

Caveats and Concluding Remarks

The relaxed clock model implemented in the present study takes into account the rate

variation among species and allows us to estimate a reliable timeline of *Prochlorococcus*

evolution. On this basis, we link an ancestral phylogenetic branch that supported population

bottlenecks and genome reduction of *Prochlorococcus* to the Neoproterozoic Snowball Earth.

Using ancestral gene gain and loss analysis, we further identify potentially important metabolic

strategies that the Cryogenian *Prochlorococcus* evolved to survive the glacial catastrophe.

Despite these fascinating results, there are important caveats. We postulated that the icehouse

conditions during the Cryogenian were lethal to the ancestral *Prochlorococcus* and thus inducing

population bottlenecks. Apparently, this key assumption derives from our knowledge on the

modern *Prochlorococcus* populations which do not grow below ~10°C, but this physiological

character is not necessarily transmissible to the ancestral population. We also postulated that the

metabolic traits we discussed earlier, which allowed *Prochlorococcus* to survive the lethally low

temperature, extremely oligotrophic condition, and low light intensity, each must have conferred

a strong fitness advantage (i.e., large selection coefficient s) to the Cryogenian *Prochlorococcus*

population. This is because for populations under powerful genetic drift and thus having small

effective population sizes (Ne), which is the case for the Cryogenian *Prochlorococcus*

populations, only those mutants that confer sufficiently large benefits ($s > 1/Ne$) can be promoted

by positive selection (58). While the metabolic traits we discussed fit the geochemical conditions

well, they need additional evidence to support the hypothesis that they were subjected to positive

selection and facilitated *Prochlorococcus* adaptation in those harsh conditions. From the

perspective of the dating methodology, the uncertainty of our analysis largely comes from the

use of calibrations. Molecular clock analysis requires at least one maximum age constraint (59).

However, available cyanobacterial fossils can only serve as the minimum bounds (16), and

therefore our analysis has to rely on the maximum bound provided by the root. We took a

conservative approach by successively increasing the maximum bound of the root from 3,800

Ma to 4,500 Ma, which showed that the posterior age of the ancestral node SBE-LCA increased

only slightly by ~7% (Fig. S2 and Table S1) and thus strengthened our conclusion of the

coincidence between SBE-LCA and Cryogenian Snowball Earth.

The Neoproterozoic Snowball Earth hypothesis was proposed decades ago, which claimed

the entire extinction of the photosynthetic organisms (21). In contrast to the original "hard"

version of the hypothesis, a modified "soft" version of the Snowball Earth hypothesis was later

proposed to include the likely persistence of refugia across the Cryogenian Period, which

allowed for the survival of bacterial and simple eukaryotic lineages (60, 61). Survivors of the

Snowball Earth included photosynthetic microorganisms (62, 63), which enabled continuous

primary production across the interval (60, 64). Like other autotrophic organisms at the base of a

food web, the survival of *Prochlorococcus* was likely important in sustaining primary

production, heterotrophy, and carbon cycling, as well as broader ecosystem functioning during

the Snowball Earth glaciations (60). On the other hand, the population bottlenecks we learned

from *Prochlorococcus* may not necessarily have occurred in other cooccurring photoautotrophic

lineages. Take *Synechococcus*, which evolutionarily most closely related to *Prochlorococcus*, as

an example. The modern *Synechococcus* have wider geographical distribution than

Prochlorococcus, and those inhabiting higher latitude regions are known to be more adaptive to

the lower seawater temperature (65, 66). Moreover, *Synechococcus* harbor more diverse

pigments than *Prochlorococcus*, which allow them to live with a wider range of light niches (67).

These unique traits may increase the survivorship of *Synechococcus* during the Snowball Earths

and thus reduce the chance to detect population bottlenecks, if any, in that difficult time.

On the other extreme, a few studies have proposed that microbial communities might have

been only mildly affected by the Snowball Earth climate catastrophe (63, 64). These inferences

were based on the microfossil and biomarker records, which, due to the lack of lineage-

specificity, did not capture the nuances required to reconstruct effects on many ecologically

important lineages as the *Prochlorococcus* studied here. Instead, we find that substantial

disruptions to the Earth system, like the Neoproterozoic Snowball Earth, leave indelible

signatures in microbial genomes, such that these heritable changes allow us to reconstruct

interactions between environmental change and biological evolution deep in Earth's history. By

employing the accelerated genome-wide accumulation of the deleterious type mutations as a

proxy for a rapid decrease in the population size of ancient lineages, we uncovered severe

bottlenecks that shaped the early evolution of *Prochlorococcus* lineages. The precise molecular

clock analyses as well as the ancestral genome reconstruction, furthermore, enabled us to link

dynamics in ancestral population sizes to changes in metabolic potential and adaptation to

icehouse climates through natural selection. Collectively, our findings demonstrate how

paleomicrontological approaches can be used to connect large-scale dynamics in the Earth

System to the genomic imprints left on extant microorganisms, which shape their ecological role

and biogeographic distribution in the world today. They also illustrate how *Prochlorococcus*

acted as important "guardians of metabolism" (12), safeguarding photosynthetic metabolic

potential across the Snowball Earth climate catastrophe.

Materials and Methods

Genomic sequences of Cyanobacteria were downloaded from public databases and manually

annotated (see "Dataset_1.tbl" in online GitHub repository and Section 1 in Supplemental

Methods). Divergence time of *Prochlorococcus* was estimated with MCMCTree v4.9e (68) on

top of 27 genes (see "Dataset_2.tbl" in online GitHub repository) previously proposed to be

valuable to date bacterial divergence (69) and *Cyanobacteria* phylogenomic trees. In previous

studies, the LPP (*Leptolyngbya*, *Plectonema*, and *Phormidium*) group of *Cyanobacteria* located

either at the basal of the Microcyanobacteria group (14, 70) or at the basal of the

Macrocyanobacteria group (71, 72). Our analysis showed that this controversy is likely caused

by the inclusion of composition-heterogeneous proteins, and that using composition-

homogeneous proteins led to consistent support for the former hypothesis (Fig. S4; see

"Dataset_3.tbl" in online GitHub repository and Section 2.2 in Supplemental Methods). Since

molecular dating analysis is known to be intrinsically associated with calibration points (73), we

summarized the calibrations of Cyanobacteria used in previous studies and modified them for our

analyses with caution. Moreover, we proposed a new strategy to use calibrations when non-

oxygenic Cyanobacteria were used as outgroups (Table S1; see Section 2.3 in Supplemental

Methods for justification). We further assessed the fitness of different molecular clock models

implemented in MCMCTree by using the package "mcmc3r" v0.3.2, based on which we decided

to use the independent rates model for further molecular clock analyses. For each molecular

clock analysis, the software ran twice with a burn-in of 50,000 and a total of 500,000

generations. The convergence was assessed based on the correlations of posterior mean time of

all ancestral nodes between independent runs (Fig. S5). By implementing statistical tests based

on the "infinite-site" theory (Fig. S3; see Section 2.6 in Supplemental Methods) we were able to

select the most precise estimates of *Prochlorococcus* evolutionary timeline for illustration (Fig.

S6) and further discussion.

Evolution of genome content via gene gains and losses was inferred using two independent

methods, AnGST (74) and BadiRate v1.35 (75). The former assumes that the statistically

supported topological differences between a gene tree and the species tree result from

evolutionary events (gene loss, gene duplication, HGT, gene birth, and speciation), and infers

these evolutionary events by reconciling the topological incongruences under a generalized

parsimony framework by achieving a minimum number of the evolutionary events along the

species tree, with penalties of an evolutionary event determined by the genome flux analysis

(74). The latter does not rely on the tree topological incongruence information, but instead uses a

full maximum-likelihood approach to determine the gene family turnover rates that maximize the

probability of observing the gene count patterns provided by the family size table. The BadiRate

analyses were run using nine strategies each with a distinct turnover rate model and a distinct

branch model. The likelihoods of different runs were compared, and three strategies with the

highest likelihood values were used (Fig. S7A). Further, results derived from AnGST (Fig. S7B)

and BadiRate were compared and summarized to determine the common patterns shared by the

two software (Fig. S7C), and important functional genes discussed were consistently inferred by

these two methods. As the two methods inferred the qualitatively same pattern of genome size

reduction on the branches leading to SBE-LCA, the number of gene gains and losses derived

from the AnGST analysis was presented.

The inference of a potential change of selection efficiency on a given branch was performed

by comparing the genome-wide d_R/d_C value across single-copy orthologous genes of the branch

to that of the closest sister branch. The d_R/d_C value was calculated using RCCalculator

(http://www.geneorder.org/RCCalculator/; see Section 4 in Supplemental Methods) based on two

independent amino acid classification schemes (Table S2).

Data Availability

The custom scripts as well as the sequence datasets are available in the online GitHub

repository (https://github.com/luolab-cuhk/Prochl-SBE).

Acknowledgments

We thank Allison Coe, Erik Zinser, and Zackary Johnson for providing the data of

Prochlorococcus growth rates, Sishuo Wang, Tianhua Liao and Xiaoyuan Feng for their

suggestions on molecular dating analyses. This work is supported by the National Natural

Science Foundation of China (92051113), the Hong Kong Research Grants Council General

Research Fund (14110820), the Hong Kong Research Grants Council Area of Excellence

Scheme (AoE/M-403/16), HKU FoS funds to SAC, and the Direct Grant of CUHK (4053257

and 3132809).

References

1. S. J. Biller, P. M. Berube, D. Lindell, S. W. Chisholm, Prochlorococcus: the structure and function of collective diversity. *Nat Rev Microbiol* **13**, 13 (2015).
2. Z. I. Johnson *et al.*, Niche partitioning among Prochlorococcus ecotypes along ocean-scale environmental gradients. *Science (80-)* **311**, 1737-1740 (2006).
3. N. J. West, D. J. Scanlan, Niche-partitioning of Prochlorococcus populations in a stratified water column in the eastern North Atlantic Ocean. *Appl Environ Microbiol* **65**, 2585-2591 (1999).
4. W. R. Hess *et al.*, The photosynthetic apparatus of Prochlorococcus: insights through comparative genomics. *Photosynth Res* **70**, 53-71 (2001).
5. L. R. Moore, G. Rocap, S. W. Chisholm, Physiology and molecular phylogeny of coexisting Prochlorococcus ecotypes. *Nature* **393**, 464 (1998).
6. B. Batut, C. Knibbe, G. Marais, V. Daubin, Reductive genome evolution at both ends of the bacterial population size spectrum. *Nat Rev Microbiol* **12**, 841 (2014).
7. H. Luo, R. Friedman, J. Tang, A. L. Hughes, Genome reduction by deletion of paralogs in the marine cyanobacterium Prochlorococcus. *Mol Biol Evol* **28**, 2751-2760 (2011).
8. J. Hu, J. L. Blanchard, Environmental sequence data from the Sargasso Sea reveal that the characteristics of genome reduction in Prochlorococcus are not a harbinger for an escalation in genetic drift. *Mol Biol Evol* **26**, 5-13 (2008).
9. H. Luo, Y. Huang, R. Stepanauskas, J. Tang, Excess of non-conservative amino acid changes in marine bacterioplankton lineages with reduced genomes. *Nat Microbiol* **2**, 17091 (2017).
10. E. P. Zuckerkandl, Linus, "Evolutionary divergence and convergence in proteins" in Evolving genes and proteins. (Elsevier, 1965), pp. 97-166.
11. M. O. Dayhoff, A model of evolutionary change in proteins. *Atlas of protein sequence and structure* **5**, 89-99 (1972).
12. P. G. Falkowski, T. Fenchel, E. F. Delong, The microbial engines that drive Earth's biogeochemical cycles. *Science (80-)* **320**, 1034-1039 (2008).
13. G. C. Kettler *et al.*, Patterns and implications of gene gain and loss in the evolution of Prochlorococcus. *PLoS Genet* **3**, e231 (2007).
14. P. Sánchez-Baracaldo, Origin of marine planktonic cyanobacteria. *Sci Rep* **5**, 17418 (2015).
15. P. Sánchez-Baracaldo, A. Ridgwell, J. A. Raven, A neoproterozoic transition in the marine nitrogen cycle. *Curr. Biol.* **24**, 652-657 (2014).
16. C. R. Marshall, Using the Fossil Record to Evaluate Timetree Timescales. *Front Genet* **10**, 1049 (2019).
17. S. C. Di Rienzi *et al.*, The human gut and groundwater harbor non-photosynthetic bacteria belonging to a new candidate phylum sibling to Cyanobacteria. *Elife* **2**, e01102 (2013).
18. R. M. Soo, J. Hemp, D. H. Parks, W. W. Fischer, P. Hugenholtz, On the origins of oxygenic photosynthesis and aerobic respiration in Cyanobacteria. *Science (80-)* **355**, 1436-1440 (2017).
19. S. A. Crowe *et al.*, Atmospheric oxygenation three billion years ago. *Nature* **501**, 535 (2013).
20. N. J. Planavsky *et al.*, Evidence for oxygenic photosynthesis half a billion years before the Great Oxidation Event. *Nat Geosci* **7**, 283 (2014).
21. P. F. Hoffman, A. J. Kaufman, G. P. Halverson, D. P. Schrag, A Neoproterozoic snowball earth. *Science (80-)* **281**, 1342-1346 (1998).
22. Y. Ashkenazy *et al.*, Dynamics of a Snowball Earth ocean. *Nature* **495**, 90 (2013).
23. E. R. Zinser *et al.*, Influence of light and temperature on Prochlorococcus ecotype distributions in the Atlantic Ocean. *Limnol Oceanogr* **52**, 2205-2220 (2007).
24. D. Thomas, G. Dieckmann, Antarctic sea ice--a habitat for extremophiles. *Science (80-)* **295**, 641-644 (2002).
25. P. F. Hoffman *et al.*, Snowball Earth climate dynamics and Cryogenian geology-geobiology. *Sci Adv* **3**, e1600983 (2017).

26.   N. Takeuchi, Optical characteristics of cryoconite (surface dust) on glaciers: the relationship
between light absorbency and the property of organic matter contained in the cryoconite. *Annals of*
Glaciology **34**, 409-414 (2002).
27.   F. C. Benforte *et al.*, Novel role of the LPS core glycosyltransferase WapH for cold adaptation in
the Antarctic bacterium Pseudomonas extremaustralis. *PLoS One* **13**, e0192559 (2018).
28.   Z. Wang, J. Wang, G. Ren, Y. Li, X. Wang, Influence of core oligosaccharide of lipopolysaccharide
to outer membrane behavior of Escherichia coli. *Mar Drugs* **13**, 3325-3339 (2015).
29.   G. Feller, C. Gerday, Psychrophilic enzymes: hot topics in cold adaptation. *Nat Rev Microbiol* **1**,
200 (2003).
30.   W. Schumann, Regulation of bacterial heat shock stimulons. *Cell Stress and Chaperones* **21**, 959-
968 (2016).
31.   A. La Terza, G. Papa, C. Miceli, P. Luporini, Divergence between two Antarctic species of the
ciliate Euplotes, E. focardii and E. nobilii, in the expression of heat-shock protein 70 genes. *Mol*
Ecol **10**, 1061-1067 (2001).
32.   J. P. Rinehart *et al.*, Continuous up-regulation of heat shock proteins in larvae, but not adults, of a
polar insect. *Proc Natl Acad Sci U S A* **103**, 14223-14227 (2006).
33.   R. P. Lawrence, J. W. William, Temperature and substrates as interactive limiting factors for marine
heterotrophic bacteria. *Aquat. Microb. Ecol.* **23**, 187-204 (2001).
34.   M. Gleitz, M. R. v.d. Loeff, D. N. Thomas, G. S. Dieckmann, F. J. Millero, Comparison of summer
and winter inorganic carbon, oxygen and nutrient concentrations in Antarctic sea ice brine. *Mar*
Chem **51**, 81-91 (1995).
35.   D. S. Reay, D. B. Nedwell, J. Priddle, J. C. Ellis-Evans, Temperature dependence of inorganic
nitrogen uptake: reduced affinity for nitrate at suboptimal temperatures in both algae and bacteria.
Appl. Environ. Microbiol. **65**, 2577-2584 (1999).
36.   J. A. Raven, B. Wollenweber, L. L. Handley, A comparison of ammonium and nitrate as nitrogen
sources for photolithotrophs. *New Phytol.* **121**, 19-32 (1992).
37.   P. Hoffman, Cryoconite pans on Snowball Earth: supraglacial oases for Cryogenian eukaryotes?
Geobiology **14**, 531-542 (2016).
38.   A. Uhde *et al.*, Glucosamine as carbon source for amino acid-producing Corynebacterium
glutamicum. *Appl Microbiol Biotechnol* **97**, 1679-1687 (2013).
39.   J. T. Park, T. Uehara, How Bacteria Consume Their Own Exoskeletons (Turnover and Recycling
of Cell Wall Peptidoglycan). *Microbiol. Mol. Biol. Rev.* **72**, 211-227 (2008).
40.   M. Borisova *et al.*, Peptidoglycan Recycling in Gram-Positive Bacteria Is Crucial for Survival in
Stationary Phase. *MBio* **7** (2016).
41.   G. C. Papageorgiou, N. Murata, The unusually strong stabilizing effects of glycine betaine on the
structure and function of the oxygen-evolving Photosystem II complex. *Photosynth Res* **44**, 243-
252 (1995).
42.   D. J. Scanlan *et al.*, Ecological genomics of marine picocyanobacteria. *Microbiol. Mol. Biol. Rev.*
73, 249-299 (2009).
43.   N. Empadinhas, M. S. da Costa, To be or not to be a compatible solute: bioversatility of
mannosylglycerate and glucosylglycerate. *Syst Appl Microbiol* **31**, 159-168 (2008).
44.   J. H. Carver, I. M. Vardavas, Precambrian glaciations and the evolution of the atmosphere. *Ann*
Geophys **12**, 674-682 (1994).
45.   J. A. Raven, J. Kübler, J. Beardall, Put out the light, and then put out the light. *J. Mar. Biol. Assoc.*
U.K. **80**, 1-25 (2000).
46.   C. Simon, A. Wiezer, A. W. Strittmatter, R. Daniel, Phylogenetic diversity and metabolic potential
revealed in a glacier ice metagenome. *Appl Environ Microbiol* **75**, 7519-7526 (2009).
47.   J. J. Brocks *et al.*, The rise of algae in Cryogenian oceans and the emergence of animals. *Nature*
548, 578 (2017).
48.   R. Kouřil, E. Wientjes, J. B. Bultema, R. Croce, E. J. Boekema, High-light vs. low-light: effect of

light acclimation on photosystem II composition and organization in Arabidopsis thaliana.
Biochimica et Biophysica Acta (BBA)-Bioenergetics **1827**, 411-419 (2013).

49.   T. Bibby, I. Mary, J. Nield, F. Partensky, J. Barber, Low-light-adapted Prochlorococcus species
possess specific antennae for each photosystem. *Nature* **424**, 1051 (2003).

50.   G. Dong, S. S. Golden, How a cyanobacterium tells time. *Curr Opin Microbiol* **11**, 541-546 (2008).

51.   J. Holtzendorff *et al.*, Genome streamlining results in loss of robustness of the circadian clock in
the marine cyanobacterium Prochlorococcus marinus PCC 9511. *J Biol Rhythms* **23**, 187-199
(2008).

52.   I. M. Axmann *et al.*, Biochemical evidence for a timing mechanism in prochlorococcus. *J Bacteriol*
191, 5342-5347 (2009).

53.   D. Mella-Flores *et al.*, Prochlorococcus and Synechococcus have Evolved Different Adaptive
Mechanisms to Cope with Light and UV Stress. *Front Microbiol* **3**, 285 (2012).

54.   S. J. Giovannoni, J. C. Thrash, B. Temperton, Implications of streamlining theory for microbial
ecology. *ISME J* **8**, 1553 (2014).

55.   R. L. Warters, B. W. Lyons, Variation in radiation-induced formation of DNA double-strand breaks
as a function of chromatin structure. *Radiat Res* **130**, 309-318 (1992).

56.   M. J. Simons, The evolution of the cyanobacterial posttranslational clock from a primitive
"phoscillator". *J Biol Rhythms* **24**, 175-182 (2009).

57.   C. W. Mullineaux, R. Stanewsky, The rolex and the hourglass: a simplified circadian clock in
Prochlorococcus? *J Bacteriol* **191**, 5333-5335 (2009).

58.   H. Luo, M. A. Moran, How do divergent ecological strategies emerge among marine
bacterioplankton lineages? *Trends Microbiol* **23**, 577-584 (2015).

59.   G. J. Szollosi *et al.*, Relative time constraints improve molecular dating. *bioRxiv* (2020).

60.   M. Moczydłowska, The Ediacaran microbiota and the survival of Snowball Earth conditions.
Precambrian Res **167**, 1-15 (2008).

61.   P. F. Hoffman, D. P. Schrag, The snowball Earth hypothesis: testing the limits of global change.
Terra nova **14**, 129-155 (2002).

62.   C. W. Allison, S. M. Awramik, Organic-walled microfossils from earliest Cambrian or latest
Proterozoic Tindir Group rocks, northwest Canada. *Precambrian Res* **43**, 253-294 (1989).

63.   F. A. Corsetti, S. M. Awramik, D. Pierce, A complex microbiota from snowball Earth times:
microfossils from the Neoproterozoic Kingston Peak Formation, Death Valley, USA. *Proc Natl*
Acad Sci USA **100**, 4399-4404 (2003).

64.   A. N. Olcott, A. L. Sessions, F. A. Corsetti, A. J. Kaufman, T. F. De Oliviera, Biomarker evidence
for photosynthesis during Neoproterozoic glaciation. *Science (80-)* **310**, 471-474 (2005).

65.   P. Flombaum *et al.*, Present and future global distributions of the marine Cyanobacteria
Prochlorococcus and Synechococcus. *Proc Natl Acad Sci USA* **110**, 9824-9829 (2013).

66.   K. Zwirglmaier *et al.*, Global phylogeography of marine Synechococcus and Prochlorococcus
reveals a distinct partitioning of lineages among oceanic biomes. *Environ Microbiol* **10**, 147-161
(2008).

67.   C. Six *et al.*, Diversity and evolution of phycobilisomes in marine Synechococcus spp.: a
comparative genomics study. *Genome Biol* **8**, R259 (2007).

68.   Z. Yang, PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**, 1586-1591
(2007).

69.   F. U. Battistuzzi, S. B. Hedges, A major clade of prokaryotes with ancient adaptations to life on
land. *Mol Biol Evol* **26**, 335-343 (2008).

70.   P. M. Shih *et al.*, Improving the coverage of the cyanobacterial phylum using diversity-driven
genome sequencing. *Proc Natl Acad Sci USA* **110**, 1053-1058 (2013).

71.   C. Blank, P. Sanchez-Baracaldo, Timing of morphological and ecological innovations in the
cyanobacteria–a key to understanding the rise in atmospheric oxygen. *Geobiology* **8**, 1-23 (2010).

72.   J. C. Uyeda, L. J. Harmon, C. E. Blank, A comprehensive study of cyanobacterial morphological

529    and ecological evolutionary dynamics through deep geologic time. *PLoS One* **11**, e0162539 (2016).

530 73. B. E. Schirrmeister, P. Sanchez-Baracaldo, D. Wacey, Cyanobacterial evolution during the
531    Precambrian. *Int J Astrobiol* **15**, 187-204 (2016).

532 74. L. A. David, E. J. Alm, Rapid evolutionary innovation during an Archaean genetic expansion.
533    *Nature* **469**, 93 (2011).

534 75. P. Librado, F. Vieira, J. Rozas, BadiRate: estimating family turnover rates by likelihood-based
535    methods. *Bioinformatics* **28**, 279-281 (2011).

Fig. 1   Evolution of *Prochlorococcus* during the Neoproterozoic Snowball Earth events. (A) (Left) Chronogram of the evolutionary history of *Prochlorococcus* estimated by MCMCTree. The evolutionary tree shown here is part of the species tree constructed with MrBayes based on 90 compositionally homogenous gene families shared by 159 cyanobacterial genomes (Fig. S4 D). Divergence time is estimated based on 27 gene sequences under calibration set C14 (Table S1). The vertical bars represent the estimated time of the Neoproterozoic glaciation events. The flanking horizontal blue bars on ancestral nodes represent the posterior 95% highest probability density (HPD) interval of the estimated divergence time. The pie chart on the ancestral branches leading to the node SBE-LCA provides the proportion of reconstructed genomic events including gene gain, gene loss, gene replacement, gene duplication and gene vertical inheritance. (Right) Phyletic pattern of key gene families that potentially enabled *Prochlorococcus* to survive harsh conditions during the Neoproterozoic Snowball Earth (at the ancestral node 'SBE-LCA'). Solid square, solid circle tend open circle next to each extant taxon represent multi-copy gene family, single-copy gene family, and absence of the gene family, respectively, in the genome. (B) (Left) The diagram helps understand how the d_R/d_C was calculated. In this context, the 'Target' group includes all genomes of all HL clades, LLI and LLII/III, the 'Control' group includes all genomes of LLIV, and the 'reference' group includes all genomes of Syne 5.1. The d_R/d_C for the 'Target' group (shown in Middle & Right) is calculated by comparing a genome from the 'Target' group to a genome from the 'reference' group (marked with red), followed by averaging the value across all possible genome pairs. Likewise, the d_R/d_C for the 'Control' group (shown in Middle & Right) is calculated by comparing a genome from the 'Control' group to a genome from the 'reference' group (marked with green) and then by averaging the value across all possible genome pairs. (Middle & Right) The genome-wide means of d_R/d_C values at the ancestral branch leading to SBE-LCA and that at its sister lineage. They were classified based on the physicochemical classification of the amino acids by charge or by volume and polarity, and were either GC-corrected by codon frequency (blue), GC-corrected by amino acid (AA) frequency (red) or uncorrected (gray). Error bars of d_R/d_C values represent the standard error of the mean. (C) The average growth rate of *Prochlorococcus* ecotypes at different temperatures. Replicate cell cultures were grown in a 14:10 light: dark cycle at 66 ±1 μmol m⁻²s⁻¹. The growth data used for plotting are collected from Johnson et al. 2006 and Zinser et al. 2007. (D) Diagram of putative bacterial refugia including cryoconite holes and sea ice brine channels in Neoproterozoic Snowball Earth.