A mixed model approach for estimating drivers of microbiota community composition and differential taxonomic abundance

Amy R Sweeny[1]*, Hannah E Lemon[1]*, Anan Ibrahim[1], Kathryn A. Watt[1], Kenneth Wilson[3], Dylan Z Childs[4], Daniel H Nussey[1], Andrew Free[2$], Luke McNally[1$]

* These authors contributed equally
$ Joint last authors

Affiliations

[1] Institute of Evolutionary Biology, University of Edinburgh, Edinburgh EH9 3FL UK

[2] Institute of Quantitative Biology, Biochemistry and Biotechnology, University of Edinburgh EH9 3FL UK

[3] Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK

[4] Department of Animal & Plant Sciences, University of Sheffield, Sheffield, UK

Corresponding author: Amy R. Sweeny, amyr.sweeny@gmail.com

Running title: GLMM approaches to metabarcoding data

Key words: microbiota, metabarcoding, 16S, amplicon sequence variants, generalised linear mixed effects model, community composition, differential abundance, Bayesian estimation

Abstract

1. Next-generation sequencing (NGS) and meta-barcoding approaches have revolutionized understanding of within-host communities, such as the gut microbiome, in humans and laboratory animals. The application of such approaches in wild animal populations is growing, but there is a disconnect between the widely-applied generalised linear mixed model (GLMM) approaches commonly used to study phenotypic variation and the statistical toolkit from community ecology which is typically applied to meta-barcoding data.

2. Here, we describe and illustrate a novel GLMM-based approach for analysing the taxon-specific sequence read counts derived from standard meta-barcoding data. This approach allows us to decompose the contribution of different drivers of variation in community structure (e.g. year, season, individual host), via interaction terms in the random effects structure of the model. We also show how these models can be used to determine the degree to which specific taxa or taxonomic groups are responsible for variance attributed to different drivers.

3. To illustrate this approach, we applied it to two cross-sectional meta-barcoding data sets from the Soay Sheep population of St. Kilda. The GLMM approach yielded results that were in agreement with more classical approaches from community ecology, showing that variation the gut microbiota community in these sheep was better explained by age group than by season. We were able to quantify the contributions of different sources of variation to community structure, and also to drill down into the model predictions to show that the age effects we observed were principally due to increases in taxa of the phyla Bacteroidetes and declines in taxa of the phyla Firmicutes.

4. Our proposed models offer a powerful new approach to understanding the drivers of variation in estimates of community structure derived from meta-barcoding data. We discuss how our approach could be readily adapted to allow researchers to estimate that contribution of host genotype, environment, and microbial/parasite phylogeny to observed community structure, and thus provide a powerful means to answer emerging questions surrounding the ecological and evolutionary roles of within-host communities.

1. Introduction

The ecological dynamics of within-host communities of parasites and commensal microbes can have dramatic effects on host health and fitness (Alberdi *et al.* 2016; Koskella *et al.* 2017). One increasingly well studied example of such a within-host community is the so-called gut microbiota: the often complex and diverse community of commensal bacteria resident to the gastrointestinal tracts of their animal hosts. As well as playing a crucial role in the digestion of food, studies from humans and model laboratory animals highlight impacts of the gut microbiota on host behaviour, metabolism, as well as endocrine and immune homeostasis (Sudo *et al.* 2004; Cox-Foster *et al.* 2007; Desbonnet *et al.* 2014; Wahlström *et al.* 2016). A growing number of studies within ecology and evolutionary biology are seeking to study the dynamics of the gut microbiota of natural systems using a combination of faecal sampling and next-generation sequencing (NGS) meta-barcoding approaches. Understanding the role of within-host communities in underpinning host phenotypic variation, as well as wider ecological and evolutionary dynamics, in the wild will require statistical approaches that allow us to robustly quantify the contribution of different environmental and host-related factors to such meta-barcoding data. Generalised linear mixed models (GLMM) are a well-established and widely used suite of statistical models within ecology and evolution which provide a flexible means for appropriately dealing with the complex data structures and relationships between predictors of interest that arise in ecological data (Bolker *et al.* 2009). Although they have yet to be widely applied in this context, GLMMs have huge potential to help dissect and understand the drivers of within-host community dynamics, as revealed by meta-barcoding data.

Standard methodologies for investigating hypotheses concerning gut microbiota dynamics in the wild typically include collection of faecal samples from selected study subjects and application of next-generation sequencing (NGS) techniques for meta-barcoding of informative bacterial genes for taxonomic assignment of sequenced reads (J *et al.* 2018). Microbiota community analysis commonly relies on transformation of OTU (operational taxonomic units) or ASV (amplicon sequence variant) counts into relative proportions per sample or rarefaction such that a set library size is randomly subsampled from all samples (McMurdie & Holmes 2014; Weiss *et al.* 2017; McMurdie 2018) Hypothesis testing using transformed counts from 16S taxonomic assignments typically is focused on community-level differences in taxonomic diversity and

composition between experimental groups or time points of interest. Statistical approaches to this

end include estimation of alpha diversity (the number of distinguishable taxa within a sample),

distance measures (e.g. Bray-Curtis dissimilarity (Bray and Curtis, 1957)), and ordination with

dimensionality reduction (e.g. principal coordinates analysis). Data transformations and

hypothesis tests in these approaches have several limitations. Standardisations of data based on

proportions ignore heteroscedasticity from different library sizes across samples, while those

relying on rarefaction restrict data such that the reads considered per each sample are limited to

the minimum number of reads across all samples (McMurdie & Holmes 2014). This in turn can

significantly elevate rates of false positives or reduce performance in microbiome clustering

approaches. In addition to statistical pitfalls, these traditional approaches for assessing

community-level differences differ philosophically from GLMM-based approaches which

partition complex sources of variance. Although traditional approaches have provided substantial

insights into microbiota community composition, they fall short of the flexibility and power

offered by GLMM-based approaches to dissect the manifold and complex contributors to

variation in measured phenotypes in natural populations. There has therefore been movement

among community ecologists toward such model-based methods (Niku *et al.* 2019). Here we

develop a GLMM-based approach to decompose the sources of variation in count data derived

from meta-barcoding approaches and discuss the advantages of this approach for analysis of

microbiota and other community data.

The application of mixed effects models to microbiota datasets is not new. The "Hierarchical

Modelling of Species Communities" (HMSC) approach uses latent variable modelling and

random effects to model community compositions 1 and has been applied to the composition of

the microbiota (refs). A similar approach developed for microbiota datasets has also shown

insights into microbiota composition in the wild (ref Bjork). Our suggested approach differs from

these in several ways. Firstly, these approaches have a focus on modelling correlations among

microbial taxa using latent variables to model residual correlation, which adds quite a lot of

complexity to the modelling process. Our approach does not attempt to model these correlations,

it focuses on variance decomposition of the sort familiar to ecologists and evolutionary biologists

working on wild systems. If correlations among microbial taxa is of primary interest we would

direct readers to these approaches. Secondly, our approach does not require the use of any

particular modelling package or a high degree of proficiency in coding. There are two central

ideas in our approach – using sample level random effects in Poisson models to account for

variability in library size, and using random effects of microbial taxonomy to allow for effects of

host and environment on microbiota composition – that can be implemented in almost any

random effects modelling software or packages with which the reader is familiar. Thus, at the

expense of modelling residual correlation among species, our approach offers a familiar method

to decompose sources of variance in the microbiota for field scientists. Below we outline the

motivation for this approach and illustrate this via an application to two 16S metabarcoding

datasets from a wild mammal.

2.   A GLMM approach

As gut microbes have such important effects on host physiology, behaviour and health much

research has sought to identify individual microbial taxa that are responsible for alterations of

host phenotype and state. This has in many ways mirrored the goals of many genome-wide

association study (GWAS) analyses, which have sought to identify particular genetic variants

associated with phenotypes of interest, often with a goal of developing diagnostics or drug

targets (Visscher *et al.* 2012). However, just as GWAS analyses have shown us that most

phenotypes are highly polygenic, being determined by a complex combination of genetic

variants of small effects (Goldstein 2009; Consortium 2009; Loh *et al.* 2015), the study of host-

associated microbiomes has often failed to find single taxa associated with host states (Clemente

et al. 2012; Vayssier-Taussat 2014). Instead many changes in host state are associated with

general shifts in microbiome composition, often termed dysbiosis (Clemente *et al.* 2012; Carding

et al. 2015). Phenomena such as dysbiosis shift the level at which we look for associations with

host phenotype move from a small number of microbial taxa to the whole microbiota. In

addition, the most pressing questions about host-associated microbiota in ecology and evolution

are very general and focused on the entire microbiota community (Koskella *et al.* 2017). For

example, what are the relative roles of host physiology and environment in shaping the

microbiota? How heritable is the microbiota? How much does microbiota composition impact

fitness? The shift in focus of these questions from individual taxa to complete community poses

an important conceptual and statistical challenge.

As previously discussed, host-associated microbiotas often constitute hundreds or thousands of

different taxa. Whenever we need to estimate a large ensemble of related parameters, a common

statistical approach is to treat them as random variables from some distribution (Gelman & Hill

2006). To understand how this approach applies to the microbiota let us consider the concrete

question of estimating how the composition of the gut microbiota might change with season in a

wild mammal. In traditional approaches to analysing microbiota datasets, it would be common to

visualise an ordination of the data, distinguishing points by season. Then, one would perform a

permutational ANOVA on a dissimilarity matrix to test if microbiotas from different seasons are

more dissimilar than those from the same season, and go on to test for differential abundance of

individual taxa across seasons to identify taxa with a major role in these changes (Segata *et al.*

2011; Mandal *et al.* 2015). In this approach, estimates for how individual taxa differ by season

are all independent of each other. Using a random effects model would approach this in a

fundamentally different manner, where a model would be fitted with a random effect for the

seasonal change for each microbial taxon. This approach has the advantage that all taxa inform

the estimate of the mean and variance of the distribution that the effects across taxa come from.

The estimates of parameters for individual taxa are then "shrunk" to this distribution. This

"shrinkage" is known to improve the accuracy of parameter estimation as long as there are large

numbers of groups for the random effects, which is generally true for most host-associated

microbiota owing to their large number of taxa.

While fitting such random effects models is known to improve parameter estimation owing to

shrinkage, its biggest advantage is in allowing us to shift the questions we ask to the whole

microbiota level, and partition complex & inter-related sources of variance. GLMM approaches

have been used across other ecological and evolutionary contexts to estimate repeatability,

relative levels of spatiotemporal variance (Albery *et al.* 2019), social and common environment

effects (Rushmore *et al.* 2013; Froy *et al.* 2018), as well as heritability and the role of host

genetics (Hayward *et al.* 2014). Answering such questions has proved hugely challenging in the

microbiota field as most analyses rely on tools which are not multi-level, from which it is

extremely difficult to decompose the relative contribution of simultaneous processes at the host

and environmental scale. However, mutli-level models have been shown to offer significant

advantages over many other compositional methods in community ecology for species

abundance data (Jackson *et al.* 2012). Here, we develop and illustrate on method to appropriately

structure random effects across microbial taxa within a community using a GLMM, and thus

partition the sources of variance driving microbiota composition.

To see how such a model can be structured let us again return to the example of estimating the

effect of season on the gut microbiota of a wild mammal (Figure 1). Consider a scenario with

two samples taken per host from a sample of hosts in a population, one in winter and one in

summer, and with samples appropriately sequenced and reads bioinformatically assigned to

ASVs. This will yield data in the form of a count of reads belonging to each ASV (the focal

taxonomic group) within each sample (Figure 1A), with two samples per individual host one

from each season. We can directly analyse such count data by fitting a Poisson family GLMM

with log-link. The predicted values on the link scale are given according to the following model

(Figure 1B).

$$log\big(y_{h,asv,s}\big) = \beta_0 + \beta_1 s + u_h + u_{asv} + u_{h:s} + u_{asv:h} + u_{asv:s} + u_{asv:h:s}$$

Here $y_{h,asv,s}$ is the read count, β_0 is a global intercept, and remaining terms account for technical

variation effects in read counts (abundance) as well as biological variation in taxonomic

composition. Fixed and random terms dealing with technical variation are as follows: β_1 is the

effect of season (s) on total read count, u_h is a random effect describing variation in total read

count among individual hosts (h, where there are multiple samples per host), u_{asv} is a random

effect describing variation in total read count of each ASV across samples and hosts , $u_{h:s}$ is a

random effect accounting for library size by describing variation in mean read count in each

sample (i.e. host by season), and $u_{asv:h:s}$ is an additional random effect accounting for row-level

variation (over-dispersion). In this example biological effects of interest are specified as follows:

$u_{asv:h}$ is a random effect describing the abundance (read count) of an ASV in host h, $u_{asv:s}$ is a

random effect describing how ASV abundances (read counts) ASV differ between seasons. By

apportioning the variance attributed to these different random effects we can assess the relative

contributions of these different factors to microbiota composition (Figure 1B-C). For example, a

high variance associated with $u_{h:s}$ would indicate a high degree of technical variation due to

library size variation across samples, and high variance associated with u_{asv} could indicate

variation in read counts across ASVs due to over-dispersion introduced by many rare taxa

(Figure 1C). With regard to biological inference, variance associated with $u_{asv:h}$ can be

interpreted as indicative of individual 'repeatability' of ASV community composition and $u_{asv:s}$

can be interpreted as reflecting variance associated with compositional shifts across seasons.

Continuing with the above example, we can further use Poisson model outputs to explore which

specific taxa are driving differential abundances between groups of interests (e.g. season), which

is commonly of great interest in microbiome studies, but for which many existing methods may

be affected by library size and normalisation methods (Weiss *et al.* 2015). Differential

abundance in this example can be estimated using each ASV-by- season level of the random

effect $u_{asv:s}$ and comparing posterior distributions for each ASV across factor levels. For

example, the mean of the posterior distribution for ASV1:summer – that for ASV1:spring can be

interpreted as the differential abundance of ASV1 between spring and summer, allowing

identification of ASVs which exhibit the largest deviations from the means.

While the Poisson model accounts for variation in library size across samples, there has also

been a shift in microbiota research towards explicitly compositional data analysis, which

removes any effects of library size (other than in quantifying uncertainty) prior to analysis. The

centered log-ratio (CLR) described by Aitchsion (1982) represents one such transformation. We

present full details of how to implement the above GLMM approach using CLRs, and then apply

this alternative parameterization to the example data described below, in our Supplementary

Files.

Figure 1. Overview of mixed model approach to wild microbiota analysis. Data processing (A) generates amplicon sequence variant (ASV)-level abundances for each sample. These raw abundances are used as the response for generalised linear mixed-effects models with Poisson error families. In the example illustrated, data includes sampling timepoints for a group of individuals taken during two seasons. Model syntax therefore specifies a fixed effect of age, and random effects for taxonomy (asv), sample id (host:season, h:s), individual differential abundance of asvs (asv:h), differential abundance of asvs across seasons (asv:s), and a residual variance at the row-level (asv:h:s). GLMM output can be used to partition the variance explained by each random effect term (B). These variance components can be interpreted as the relative contributions of both technical variation and host or environmental contributions to differential abundance as illustrated in (C). Created with BioRender.com

3. A worked example: age and season effects on gut microbiota in wild sheep

To test and illustrate our approach, we obtained faecal samples from Soay Sheep (*Ovis aries*) from the island of Hirta in the St. Kilda archipelago of the Outer Hebrides of Scotland. These animals are free-living and are part of a long-term study in which individuals have been marked and monitored longitudinally since 1985 (Clutton-Brock Pemberton 2004). All animals sampled had been caught and uniquely tagged within a few days of birth, so that their age and sex were known with certainty. Each year, fieldwork teams visit St Kilda in spring to monitor lambing and capture, mark and sample newborn lambs within a few days of birth. Subsequently, each August a larger field team visits to capture, mark and sample animals living in the study area using a series of corral traps (Clutton-Brock & Pemberton 2004).

Two sets of faecal samples were collected, in 2013 and 2016, to allow comparison of the gut microbiota of individuals of different ages (2013) and from the same individuals sampled in different season (2016). The 2013 samples were collected during the August catch and included 30 samples from lambs (around 4 months old) and 28 samples from older adults (ages 2-13). The 2016 samples were collected from a set of 36 females age 1-13 years who were sampled in both spring (around the time of parturition) and then 3-4 months later in August. Microbial DNA was extracted from samples, amplified using bacterial 16S rRNA V4 region primers, and sequenced using the Ilumina MiSeq platform to generate 250 base pair (bp) paired-end reads. Sequences were processed using the DADA2 pipeline in R (v1.12.1) to call amplicon sequence variants (ASVs) (Callahan *et al.* 2016). Full details of sampling, sequencing, and data processing methods are provided in the Electronic Supplementary Material (ESM 1.1-1.4). We conducted

standard dissimilarity analysis and PERMANOVA tests on effects of age and season for

comparison to the results of our GLMM approach (see ESM 1.4.1 for full details).

3.1. Specification of GLMMs:

We applied separate GLMMs to the 2013 and 2016 datasets. First, an aggregate dataset for each

year was created from the sample metadata, taxonomic classifications for each ASV, and an

ASV-by-sample abundance matrix. We fit GLMMS with Poisson errors and log links to each

dataset using the Bayesian package 'MCMCglmm' (Hadfield 2010) using the approach

introduced in Section 2. Models fit to 2013 data, which included samples from hosts of two age

classes from a single season (one sample per host) were specified as follows:

$$log(\bar{y}_{h,asv}) = \beta_0 + a\beta_a + u_h + u_{asv} + u_{asv:a} + u_{asv:h}$$

Where $y_{h,asv}$ is the read count per ASV within each host, β_0 is a global intercept, β_a is the effect

of age (a, binary factor: lamb versus adult) on total read count, u_h is a random effect random

effect describing variation in total read count among individual hosts (h, equivalent to sample

here where hosts are sampled once each), u_{asv} is a random effect describing variation in total

read count of each ASV across hosts/samples, $u_{asv:a}$ is a random effect a random effect

describing how ASV abundances (read counts) ASV differ between host age classes and $u_{asv:h}$

is an additional is an additional row-level random effect describing residual variation.

Models fit to 2016 data, which included samples from individual hosts of similar age sampled in

both spring and summer of the same year (two samples per host), were specified as follows:

$$log(\bar{y}_{h,asv,s}) = \beta_0 + s\beta_s + u_h + u_{asv} + u_{h:s} + u_{asv:h} + u_{asv:s} + u_{asv:h:s}$$

Here $y_{h,asv,s}$ is the read count, β_0 is a global intercept, β_s is the effect of season (s, binary factor:

spring versus summer) on total read count, u_h is a random effect describing variation in total

read count among individual hosts (h, where there are multiple samples per host), u_{asv} is a

random effect by describing variation in total read count of each ASV across samples and

hosts, $u_{h:s}$ is a random effect describing variation in total read count among individual hosts (h,

where there are multiple samples per host), $u_{asv:h}$ is a random effect describing the abundance

(read count) of an ASV in host *h,* $u_{asv:s}$ is a random effect describing how ASV abundances

(read counts) ASV differ between seasons, and $u_{asv:h:s}$ is an additional row-level random effect

describing residual variation.

β_0 is a global intercept, and remaining terms account for technical variation as well as biological

variation of interest. Fixed and random terms dealing with technical variation are as follows: β_1

is the effect of season (*s*) on total read count, u_h is a random effect describing variation in total

read count among individual hosts (h, where there are multiple samples per host), u_{asv} is a

random effect by describing variation in total read count of each ASV across samples and hosts

, $u_{h:s}$ is a random effect accounting for library size by describing variation in mean read count in

each sample (i.e. host by season), and $u_{asv:h:s}$ is an additional random effect accounting for row-

level variation (over-dispersion). In this example biological effects of interest are specified as

follows: $u_{asv:h}$ is a random effect describing the abundance (read count) of an ASV in host *h,*

$u_{asv:s}$ is a random effect describing how ASV abundances (read counts) ASV differ between

seasons.

Using this GLMM approach, we calculated the relative contributions to sources of variance in

the data from both technical and biological model components. We followed Nakagawa &

Schielzeth (Nakagawa & Schielzeth 2013) for calculation of r^2 from GLMMs with Poisson error

distributions. Using this formula, there is a portion of variance equal to 1 minus the sum variance

of the model components which represents variance arising from the Poisson distribution.

Where multiple samples are present per individual (2016), repeatability of the community

composition of ASVs can be estimated as the proportion of variance attributable to differential

taxonomic composition across individuals divided by the sum of the variance explained by all

other non-technical component terms estimating compositional effects $u_{asv:h}/(u_{asv:s} + u_{asv:h} +$

$u_{asv:h:s})$.

We investigated differential abundances as outlined above (2). To extract information of specific

bacterial taxa contributing to differential abundance across age groups or season, we used

Poisson model outputs and subtracted the posterior distributions for each ASV between group

levels (2013: Age; 2016: Season). We used the resultant distribution to calculate a mean

difference and highest posterior density interval (HPDI) to estimate differential abundance for

each ASV. For example, the mean of the posterior distribution for ASV1:summer – that for

ASV1:spring can be interpreted as the differential abundance of ASV between spring and

summer, where a difference can be considered significant when credible intervals do not span

zero. We calculated p-values of each differential abundance by calculating the

proportional overlap of each pairwise set of distributions divided by half the number of stored

iterations.

3.2. Results:

The microbiota communities of Soay Sheep were dominated by two phyla, Firmicutes and

Bacteroidetes (Figure S1), as has been previous observed in most vertebrates (Ley *et al.* 2008)

Principle coordinates analysis (PCoA) based on Bray-Curtis dissimilarity indicated clustering of

samples by age and by season (Figure 2). The result of a PERMANOVA test on the 2013 data

set showed a significant difference in group centroids for lambs and adults (pseudo-F = 7.161, p

< 0.001), with 11.34% of the variance in gut microbiota composition (R^2) is explained by

differences between lambs and adults. PERMANOVA results for the 2016 data showed that

group centroids for April and August are significantly distinct (pseudo-F = 2.026, p = 0.002), but

season only explains 2.81% (R^2) of the observed variance.

349
350
351   **Figure 2**. Soay Sheep gut microbiota beta diversity in adults and lambs from 2013 (A) and from April
352   and August of 2016 (B). Principle coordinates (PCoA) plots represent Bray-Curtis dissimilarity indicating
353   clustering of samples by group. Ellipsoids represent a 95% confidence interval surrounding each group.
354

Poisson GLMMs from 2013 data showed comparable results to ordination approaches, where

community composition differed substantially between age classes (Proportion variance $u_{asv:a}$:

19.88% CI 18.47-21.39%; Figure 3). Additional effects estimated by the model showed a

substantial proportion of variance explained by taxonomic variation in ASV abundance (u_{asv}:

17.35%), a small portion of variation explained by variation in mean library size across samples

(u_h 2013: 1.59 %), and considerable residual variance (estimated by the 'units' term in

MCMCglmm; $u_{asv:h}$ 2013: 49.42%; Figure 3).

Poisson GLMMs from 2016 data likewise showed comparable results to ordination approaches,

where community composition changed negligibly between seasons ($u_{asv:s}$: 1.24% CI 0.99 –

1.44%; Figure 3). The 2016 model showed a substantial proportion of variance explained by

taxonomic variation in ASV abundance (u_{asv} 2016: 34.74%), a small portion of variation

explained by variation in mean library size across samples ($u_{h:s}$ 2013: 1.99 %), and considerable

residual variance ($u_{asv:h:s}$ 2013: 46.67%; Figure 3). Repeated sampling of individuals in 2016

additionally showed moderate variance explained by inter-individual variation in community

composition ($u_{asv:h}$ 2016: 5.49 %). The equated to an individual repeatability of 11.1% for their

microbiota community composition across sampling time points.

As outlined in 3.1 we calculated differential abundances using the saved posterior distributions

for each random effect level of specific taxa across age classes (2013 data set; Figure 4A & B)

and seasons (2016 data set; Figure 4C & D). For the ASV-by-age effect in the 2013 data ($u_{asv:a}$,

Figure 3), the estimates of taxa-specific differential abundances suggest that ASVs

demonstrating significant shifts between lambs and adults belong primarily to two phyla,

Firmicutes and Bacteroidetes (Figure 4A-B). 683 out of 2,023 (33.76%) ASVs present in 2013

data showed significant shifts between lambs and adults (50.81% positive shifts, 49.19%

negative shifts). Bacteroidetes represented 52.16% of significant positive shifts into adulthood,

and Firmicutes represented 72.02% of the significant negative shifts into adulthood (Figure 4B;

Table S4). For the ASV-by-season effect in the 2016 data (2016 $u_{asv:s}$, Figure 3 & Table S4),

very few ASVs (24 of 2,364; 1.02%) showed significant differential abundance between spring

and summer sampling (Figure 4C-D; Table S4).

Our results indicate that there are substantive developmental shifts in the Soay sheep microbiota

between lambs and adults, and that the majority of taxa shifting in abundance belong to the

Bacteroidetes and Firmicutes. However, this raises the question of whether this shift is because

Bacteroidetes are *generally* more abundant in adults and Firmicutes more abundant in lambs, or

if the ASVs that show these patterns *just happen* to be in these phyla. To illustrate how GLMMs

can be used to address questions of this sort, we modified our models for the 2013 dataset to

include additional taxonomic effects of family and phylum, allowing us to identify taxonomic

levels most responsible for differential abundances. Details of these phylogenetically more

explicit GLMMs and their results and implications are presented in detail in the ESM (see

sections 1.4.3, Figure S4, Table S3). Results suggest considerably variation with respect to age

across families and ASVs within both the Bacteroidetes and Firmicutes phyla, and that most of

the age effects on microbiota community composition occurs at these lower taxonomic levels

(Figure S4; Table S3).

399
400
401

402
403 **Figure 3**. Proportion of variance in bacterial read counts from different ASVs explained by GLMM
404 component terms for two datasets. The 2013 dataset (A) compared gut microbiota across two age classes
405 from individuals sampled once at the same time point), while the 2016 dataset (B) compared samples
406 taken from the same individuals over two seasons.

407
408

Figure 4. Differential abundances across age classes (A-B) or season (C-D) for individual ASVs
calculated from GLMMs with Poisson error families and taxonomic level specified as ASV-only. A & C
represent all ASV-level effects. Violin plots represent distribution of effect estimates, size of point
represents the inverse variance of the estimate. Rectangles indicate the ASVs with the highest magnitude
(positive or negative) differential abundances in Forest plots (B & D). Forest plots represent point
estimate and HPDI for the ASVs involved in the 50 (age class) or 20 (season) strongest increases and
decreases of abundance.

417
418

4.   Discussion & wider applications

We have described a novel approach to analysing meta-taxonomic data derived from NGS using a GLMM framework, and have illustrated this method using data describing variation in the gut microbiota community in wild sheep. Our approach represents an important step forward for researchers interested in using meta-taxonomic approaches to understand variation in community structure in complex, non-experimental settings. It allows the well-established power and flexibility of GLMM-based approaches to be harnessed to decompose drivers of variation in NGS-derived data on the taxonomic composition of samples. Our example analyses provide very simple illustrations of how the approach can be used to estimate the contribution of host-related or environmental factors (specifically, age and season) to variation in the community structure of the gut microbiota. We show that results are comparable to those derived from widely-applied ordination-based approaches, and discuss the implications of observed variation in gut microbiota structure with age and season briefly below. However, these analyses are intended mainly as templates to help illustrate the approach, and barely scratch the surface of the types of important outstanding questions the method could be used to tackle with larger-scale data sets. Applying GLMMs to taxonomic-level sequence count data provides a rich toolkit from fields like quantitative genetics to dissect the contributions of different environmental and host factors to variation in community structure, with the potential to advance our understanding of community ecology, host-microbe or -pathogen interactions, and evolutionary dynamics.

In our illustrative analyses of wild Soay sheep, both GLMM and ordination-based approaches identified a stronger effect of age than season on the gut microbiota. Changes in the structure of the gut microbiota across development in early life and during senescence in later adulthood are well-established in human studies (Popkes & Valenzano 2020), but remain poorly understood in natural populations. Our data clearly show the gut microbiota community structure changes between recently weaned lambs and adults, and argues for further longitudinal studies in natural systems to test whether shifts in gut community structure could play a role in patterns of demographic ageing in the wild. A growing number of studies in the wild have documented seasonal gut microbiota changes (Amato *et al.* 2015; Maurice *et al.* 2015; Ren *et al.* 2017; Orkin *et al.* 2019). The absence of a strong seasonal different in our data may be related to the relative homogeneity of the herbivorous diet of Soay sheep, as most previous wild studies are of

omnivores with strong seasonal shifts in diet preference. Alternatively, it may be because the

spring and summer seasons we sampled in are both periods of relatively high food abundance

and quality, and more pronounced change would be observed in autumn and winter when habitat

quality and food availability changes more dramatically. In future studies, repeated sampling of

the same individuals over time will be crucial to understand the effects of age, environmental and

other variables on gut community structure. The application of our approach to such longitudinal

metataxonomic data sets will help researchers robustly estimate within-individual patterns of

change in community structures over time or space, whilst also estimating how repeatable

community structure across hosts.

Our novel GLMM approach allows estimation of key ecological and evolutionary parameters

from meta-taxonomic data sets which can advance our understanding of within host-microbe

evolutionary dynamics. Individual repeatability of measured phenotypes is an important and

widely estimated parameter in ecology and quantitative genetics (Wilson 2018). Estimating the

within-host repeatability of microbiota community structure over time can offer insight into the

extent to which host control and environmental selection determine species composition (Foster

et al. 2017). The GLMM structure presented here directly estimates this repeatability across two

seasons in our 2016 data set at around 11%, although the small sample size and temporal

proximity of the samples should mean we interpret this parameter estimate with caution.

However, the model is illustrative and it should be clear that it is readily extendible to mixed-

effect model of more a more complex nature to address emerging questions in the field. For

example, effects of host relatedness & inbreeding effects on microbiota composition in have

been explored previously in microbiome studies but via ordination methods using a small

number of genetic clusters as grouping units (Yuan *et al.* 2015). Including host genetic

relatedness matrices as a random effect in a mixed-effects model (so-called "animal models")

within our GLMM framework can offer insights into heritability and inbreeding effects on

community composition as it compares to other forces in the population (Wilson *et al.* 2010).

Factors which can be of both considerable ecological and evolutionary interest, and that can also

confound heritability estimates, such as maternal effects or shared nest or litter effects can

likewise be incorporated in these models (Kruuk & Hadfield 2007). Statistical advances to

address maternal and social effects as well spatial autocorrelation in ecological datasets can also

be incorporated in microbiota analyses (Hayward *et al.* 2010; Albery *et al.* 2020). Applied to

larger longitudinal data sets, our GLMM approach can allow researchers to directly estimate how

different aspects of host state, genotype and environment influence the structure of within-host

communities, and address many outstanding questions about the evolutionary and ecological

causes and consequences of host-microbe interactions.

Full realisation of the role of microbiota communities in the ecology and evolution of wild

organisms depends on both identifying factors with important effects on global microbiota

composition and on being able to test whether and how individual taxa or taxonomic groups

underpin those effects. Our GLMM approach readily lends itself to addressing both questions.

We have illustrated how the approach can be used to identify ASVs involved in community level

shifts with age identified in the random effects structure of the 2013 models (Figure 4), and to

further decompose the contribution of different taxonomic levels to community level effects

(ESM 1.4.3; Table S3). For example, our analysis highlights that analysis at the phylum level

could provide a misleading view of compositional shifts associated with age of Soay sheep, and

that there is substantial variation at the family level within each phylum (Figure S4). This

analysis approach should represent similar insights to linear discriminant analysis (LDA) used in

approaches such as LefSE (Segata *et al.* 2011), with the advantage of extraction of this

information for multiple factors of interest rather than requirement of a priori knowledge of

effects of interest to run differential abundance analysis. Our approach could be further

developed beyond the nested taxonomic levels used here to identify taxonomic levels associated

with the greatest variance (ESM 1.4.3) to explicitly include the microbiota phylogeny within the

GLMMs, allowing environment and host effects on community composition to be estimated

accounting more accurately for phylogenetic distances between ASVs (Hadfield & Nakagawa

2010) in a similar manner to UNIFRAC clustering approaches (Lozupone *et al.* 2006). A

GLMM-based approach capable of simultaneously dissecting the contributions of host

environment, state and genetics alongside microbial phylogeny to variation in microbial

community structure seems to us to represent a powerful step towards robustly address emerging

questions surrounding the role of the microbiome in ecology and evolution. However, we note

however that there are computational challenges associated with incorporating more complex

covariance structures into our GLMMs, and provide some discussion of this issue in the

supplementary methods with a worked example of an alternate approach (ESM Section 2).

Beyond applications to microbiota community analyses, approaches outlined in this manuscript

are applicable more broadly to different types of metataxonomic data being collected across

myriad systems and research disciplines. For example, there has been great interest in describing

the dynamics of the parasite community as an ecosystem and understanding its influence on host

health (Pedersen & Fenton 2007; Graham 2008; Rynkiewicz *et al.* 2015). A growing number of

studies apply meta-barcoding to faecal samples to estimate the community structure of gastro-

intestinal parasite communities (Avramenko *et al.* 2015; Aivelo & Medlar 2017), and our

GLMM approach could readily applied to such data sets to dissect the drivers of variation in

parasite community composition. Another area of interest within ecology and evolution is using

meta-barcoding of faecal samples to estimate diet composition and its relationship to host

phenotypes. Bayesian mixed model approaches have also recently been applied to analysis of

presence and absence of *Cyanistes caeruleus* (blue tit) diet components and align conceptually

with approaches presented in this manuscript (Shutt *et al.* 2020). GLMM approaches to

metabarcoding data maintain key similarities to other multivariate community ecology

approaches to abundance data (Wang *et al.* 2012) while integrating benefits of ecological and

evolutionary approaches to quantifying phenotypic variation. We therefore suggest that

approaches presented in this manuscript can be applied across a range of systems and data types

for powerful and flexible understanding of complex drivers of community dynamics.

Acknowledgements
This work was funded by a large NERC grant (NE/R016801/1), and the long-term study on St
Kilda was funded principally by responsive mode grants from NERC. We thank Adam Hayward
and Jill Pilkington for sample collection, the National Trust for Scotland for support of our work
on St Kilda, and QinetiQ and Kilda Cruises for logistical support. We also thank the Ecology
Within Team for input in the analysis and manuscript and Josephine Pemberton for support and
management of the field project. Figure 1 was created with Biorender.com. Photos on which
sheep icons in Figure 1 are based by Hannah Vallin & Martin Stoffel.

Author contributions.
LM and ARS conceived and developed the statistical methods; LM, ARS, AF, HL conducted the
analyses; AF, AI, KAW, KW and DHN oversaw and undertook sample collection and laboratory
work; ARS wrote the first draft of the manuscript and all authors contributed to writing of the
final manuscript.

Data Accessibility
Data and code for this manuscript will be available on Dryad Repository upon acceptance.
Sequences and metadata on which analysis is based can be found on the European Nucleotide
Archive (ENA): http://www.ebi.ac.uk/ena/data/view/PRJEB39322

References

Aitchison, J. (1982). The Statistical Analysis of Compositional Data. *Journal of the Royal Statistical Society: Series B (Methodological)*, **44**, 139–160.

Aivelo, T. & Medlar, A. (2017). Opportunities and challenges in metabarcoding approaches for helminth community identification in wild mammals. *Parasitology*, **145**, 608–621.

Alberdi, A., Aizpurua, O., Bohmann, K., Zepeda-Mendoza, M.L. & Gilbert, M.T.P. (2016). Do Vertebrate Gut Metagenomes Confer Rapid Ecological Adaptation? *Trends in Ecology & Evolution*, **31**, 689–699.

Albery, G.F., Becker, D.J., Kenyon, F., Nussey, D.H. & Pemberton, J.M. (2019). The fine-scale landscape of immunity and parasitism in a wild ungulate population. *Integrative and Comparative Biology*, **16**, e2003538.

Albery, G.F., Kirkpatrick, L., Firth, J.A. & Bansal, S. (2020). Unifying spatial and social network analysis in disease ecology (D. Farine, Ed.). *Journal of Animal Ecology*, **30**, 2009–18.

Amato, K.R., Leigh, S.R., Kent, A., Mackie, R.I., Yeoman, C.J., Stumpf, R.M., Wilson, B.A., Nelson, K.E., White, B.A. & Garber, P.A. (2015). The gut microbiota appears to compensate for seasonal diet variation in the wild black howler monkey (Alouatta pigra). *Microbial ecology*, **69**, 434–443.

Avramenko, R.W., Redman, E.M., Lewis, R., Yazwinski, T.A., Wasmuth, J.D. & Gilleard, J.S. (2015). Exploring the Gastrointestinal 'Nemabiome': Deep Amplicon Sequencing to Quantify the Species Composition of Parasitic Nematode Communities (E. Serrano Ferron, Ed.). *PLoS One*, **10**, e0143559–18.

Bolker, B.M., Brooks, M.E., Clark, C.J., Geange, S.W., Poulsen, J.R., Stevens, M.H.H. & White, J.-S.S. (2009). Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in Ecology & Evolution*, **24**, 127–135.

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A. & Holmes, S.P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature methods*, **13**, 581–583.

Carding, S., Verbeke, K., Vipond, D.T., Corfe, B.M. & Owen, L.J. (2015). Dysbiosis of the gut microbiota in disease. *Microbial Ecology in Health and Disease*, **26**, 1–10.

Clemente, J.C., Ursell, L.K., Parfrey, L.W. & Knight, R. (2012). The Impact of the Gut Microbiota on Human Health: An Integrative View. *Cell*, **148**, 1258–1270.

Consortium, T.I.S. (2009). Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature*, **460**, 748–752.

Cox-Foster, D.L., Conlan, S., Holmes, E.C., Palacios, G., Evans, J.D., Moran, N.A., Quan, P.-L., Briese, T., Hornig, M., Geiser, D.M., Martinson, V., vanEngelsdorp, D., Kalkstein, A.L., Drysdale, A., Hui, J., Zhai, J., Cui, L., Hutchison, S.K., Simons, J.F., Egholm, M., Pettis, J.S. & Lipkin, W.I. (2007). A metagenomic survey of microbes in honey bee colony collapse disorder. *Science*, **318**, 283–287.

Desbonnet, L., Clarke, G., Shanahan, F., Dinan, T.G. & Cryan, J.F. (2014). Microbiota is essential for social development in the mouse. *Molecular psychiatry*, **19**, 146–148.

Foster, K.R., Schluter, J., Coyte, K.Z. & Rakoff-Nahoum, S. (2017). The evolution of the host
microbiome as an ecosystem on a leash. *Nature*, **548**, 43–51.

Froy, H., Borger, L., Regan, C.E., Morris, A., Morris, S., Pilkington, J.G., Crawley, M.J., Clutton Brock,
T.H., Pemberton, J.M. & Nussey, D.H. (2018). Declining home range area predicts reduced late-life
survival in two wild ungulate populations. *Ecology letters*, **21**, 1001–1009.

Gelman, A. & Hill, J. (2006). *Data Analysis Using Regression and Multilevel/Hierarchical Models*.
Cambridge University Press.

Goldstein, D.B. (2009). Common Genetic Variation and Human Traits. *N. Engl. J. Med.*, 1696–1703.

Graham, A.L. (2008). Ecological rules governing helminth-microparasite coinfection. *Proceedings of the
National Academy of Sciences of the United States of America*, **105**, 566–570.

Hadfield, J.D. (2010). MCMC methods for multi-response generalized linear mixed models: The
MCMCglmm R Package. *Journal of Statistical Software*, **33**, 1–22.

Hadfield, J.D. & Nakagawa, S. (2010). General quantitative genetic methods for comparative biology:
phylogenies, taxonomies and multi-trait models for continuous and categorical characters. *Journal of
Evolutionary Biology*, **23**, 494–508.

Hayward, A.D., Garnier, R., Watt, K.A., Pilkington, J.G., Grenfell, B.T., Matthews, J.B., Pemberton,
J.M., Nussey, D.H. & Graham, A.L. (2014). Heritable, heterogeneous, and costly resistance of sheep
against nematodes and potential feedbacks to epidemiological dynamics. *American Naturalist*, **184**,
S58–S76.

Hayward, A.D., Pilkington, J.G., Pemberton, J.M. & Kruuk, L.E.B. (2010). Maternal effects and early-life
performance are associated with parasite resistance across life in free-living Soay sheep.
Parasitology, **137**, 1261–1273.

J, P., Glendinning, L., Wisedchanwet, T. & Watson, M. (2018). The Madness of Microbiome: Attempting
To Find Consensus 'Best Practice' for 16S Microbiome Studies. *Applied and environmental
microbiology*, **84**, e02627–17.

Jackson, M.M., Turner, M.G., Pearson, S.M. & Ives, A.R. (2012). Seeing the forest and the trees:
multilevel models reveal both species and community patterns. *Ecosphere*, **3**, 79.

Koskella, B., Hall, L.J. & Metcalf, C.J.E. (2017). The microbiome beyond the horizon of ecological and
evolutionary theory. *Nature Ecology & Evolution*, **1**, 1606–1615.

Kruuk, L.E.B. & Hadfield, J.D. (2007). How to separate genetic and environmental causes of similarity
between relatives. *Journal of Evolutionary Biology*, **20**, 1890–1903.

Ley, R.E., Hamady, M., Lozupone, C., Turnbaugh, P.J., Ramey, R.R., Bircher, J.S., Schlegel, M.L.,
Tucker, T.A., Schrenzel, M.D., Knight, R. & Gordon, J.I. (2008). Evolution of mammals and their
gut microbes. *Science*, **320**, 1647–1651.

Loh, P.-R., Bhatia, G., Gusev, A., Finucane, H.K., Bulik-Sullivan, B.K., Pollack, S.J., de Candia, T.R.,
Lee, S.H., Wray, N.R., Kendler, K.S., O'Donovan, M.C., Neale, B.M., Patterson, N. & Price, A.L.
(2015). Contrasting genetic architectures of schizophrenia and other complex diseases using fast
variance-components analysis. *Nature Genetics*, 1–11.

Lozupone, C., Hamady, M. & Knight, R. (2006). **UniFrac – An online tool for comparing microbial**
community diversity in a phylogenetic context. *BMC Bioinformatics*, **7**.

Mandal, S., Van Treuren, W., White, R.A., Eggesbø, M., Knight, R. & Peddada, S.D. (2015). Analysis of
composition of microbiomes: a novel method for studying microbial composition. *Microbial Ecology*
in Health and Disease, **26**, 1–8.

Maurice, C.F., Knowles, S.C.L., Ladau, J., Pollard, K.S., Fenton, A., Pedersen, A.B. & Turnbaugh, P.J.
(2015). Marked seasonal variation in the wild mouse gut microbiota. *ISME Journal*, **9**, 2423–2434.

McMurdie, P.J. (2018). Normalization of Microbiome Profiling Data. *Microbiome Analysis*, pp. 143–168.
Methods and Protocols. Humana Press, New York, NY, New York, NY.

McMurdie, P.J. & Holmes, S. (2014). Waste Not, Want Not: Why Rarefying Microbiome Data Is
Inadmissible (A.C. McHardy, Ed.). *PLoS computational biology*, **10**, e1003531.

Nakagawa, S. & Schielzeth, H. (2013). A general and simple method for obtaining R2 from generalized
linear mixed-effects models. *Methods in Ecology and Evolution*, **4**, 133–142.

Niku, J., Hui, F.K.C., Taskinen, S. & Warton, D.I. (2019). gllvm: Fast analysis of multivariate abundance
data with generalized linear latent variable models in. *Methods in Ecology and Evolution*, **10**, 2173–
2182.

Orkin, J.D., Campos, F.A., Myers, M.S., Cheves Hernandez, S.E., Guadamuz, A. & Melin, A.D. (2019).
Seasonality of the gut microbiota of free-ranging white-faced capuchins in a tropical dry forest. *Isme*
Journal, **13**, 183–196.

Pedersen, A.B. & Fenton, A. (2007). Emphasizing the ecology in parasite community ecology. *Trends in*
Ecology & Evolution, **22**, 133–139.

Popkes, M. & Valenzano, D.R. (2020). Microbiota–host interactions shape ageing dynamics.
Philosophical Transactions of the Royal Society of London B: Biological Sciences, **375**, 20190596–9.

Ren, T., Boutin, S., Humphries, M.M., Dantzer, B., Gorrell, J.C., Coltman, D.W., McAdam, A.G. & Wu,
M. (2017). Seasonal, spatial, and maternal effects on gut microbiome in wild red squirrels.
Microbiome, **5**, 163–14.

Rushmore, J., Caillaud, D., Matamba, L., Stumpf, R.M., Borgattti, S.P. & Altizer, S. (2013). Social
network analysis of wild chimpanzees provides insights for predicting infectious disease risk.
Journal of Animal Ecology, **82**, 976–986.

Rynkiewicz, E.C., Pedersen, A.B. & Fenton, A. (2015). An ecosystem approach to understanding and
managing within-host parasite community dynamics. *Trends in Parasitology*, **31**, 212–221.

Segata, N., Izaard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W.S. & Huttenhower, C. (2011).
Metagenomic biomarker discovery and explanation. *Genome biology*, **12**, R60.

Shutt, J.D., Nicholls, J.A., Trivedi, U.H., Burgess, M.D., Stone, G.N., HADFIELD, J.D. & Phillimore,
A.B. (2020). Gradients in richness and turnover of a forest passerine's diet prior to breeding: A mixed
model approach applied to faecal metabarcoding data. *Molecular Ecology*, **29**, 1199–1213.

Sudo, N., Chida, Y., Aiba, Y., Sonoda, J., Oyama, N., Yu, X.-N., Kubo, C. & Koga, Y. (2004). Postnatal
665   microbial colonization programs the hypothalamic-pituitary-adrenal system for stress response in
666   mice. *The Journal of physiology*, **558**, 263–275.

667 Vayssier-Taussat, M. (2014). Shifting the paradigm from pathogens to pathobiome: new concepts in the
668   light of meta-omics. 1–7.

669 Visscher, P.M., Brown, M.A., McCarthy, M.I. & Yang, J. (2012). REVIEW Five Years of GWAS
670   Discovery. *The American Journal of Human Genetics*, **90**, 7–24.

671 Wahlström, A., Sayin, S.I., Marschall, H.-U. & Bäckhed, F. (2016). Intestinal Crosstalk between Bile
672   Acids and Microbiota and Its Impact on Host Metabolism. *Cell Metabolism*, **24**, 41–50.

673 Wang, Y., Naumann, U., Wright, S.T. & Warton, D.I. (2012). mvabund- an Rpackage for model-based
674   analysis of multivariate abundance data. *Methods in Ecology and Evolution*, **3**, 471–474.

675 Weiss, S.J., Xu, Z., Amir, A., Peddada, S., Bittinger, K., Gonzalez, A., Lozupone, C., Zaneveld, J.R.,
676   Vazquez-Baeza, Y., Birmingham, A. & Knight, R. (2015). Effects of library size variance, sparsity,
677   and compositionality on the analysis of microbiome data. *PeerJ PrePrints*, 3:e1157v1.

678 Weiss, S., Xu, Z.Z., Peddada, S., Amir, A., Bittinger, K., González, A., Lozupone, C., Zaneveld, J.R.,
679   Vázquez-Baeza, Y., Birmingham, A., Hyde, E.R. & Knight, R. (2017). Normalization and microbial
680   differential abundance strategies depend upon data characteristics. *Microbiome*, **5**, 1–18.

681 Wilson, A.J. (2018). How should we interpret estimates of individual repeatability? *Evolution Letters*, **2**,
682   4–8.

683 Wilson, A.J., Réale, D., Clements, M.N., Morrissey, M.M., Postma, E., WALLING, C.A., Kruuk, L.E.B.
684   & Nussey, D.H. (2010). An ecologist's guide to the animal model. *Journal of Animal Ecology*, **79**,
685   13–26.

686 Yuan, M.L., Dean, S.H., Longo, A.V., Rothermel, B.B., Tuberville, T.D. & Zamudio, K.R. (2015).
687   Kinship, inbreeding and fine-scale spatial structure influence gut microbiota in a hindgut-fermenting
688   tortoise. *Molecular Ecology*, **24**, 2521–2536.