Heterochiasmy facilitated the establishment of *gsdf* as a novel sex determining gene in

Atlantic halibut

Edvardsen, R.B*[1]., Wallerman, O[3]., Furmanek, T[1]., Kleppe, L[1]., Jern, P[3], Wallberg, A[3].,

Kjærner-Semb, E[1]., Mæhle, S[1]., Olausson, S. K[2]., Sundström, E[3]., Harboe, T[2]., Mangor-Jensen,

R[2]., Møgster, M[2]., Perrichon, P[2]., Norberg, B.[2], Rubin, C-J*[3]

*Corresponding authors

rolf.brudvik.edvardsen@hi.no

carl-johan.rubin@imbim.uu.se

Affiliations:

[1] Institute of Marine Research, P.O. Box 1870, Nordnes, NO-5817, Bergen, Norway.

[2] Institute of Marine Research, Austevoll Research Station, NO-5392, Storebø, Norway.

[3] Department of Medical Biochemistry and Microbiology, Uppsala University, 75123 Uppsala,

Sweden.

Keywords: aquaculture, nucleotide diversity, transposable elements, genome assembly, meiotic

recombination, sex chromosome

Abstract

Atlantic Halibut (Hippoglossus hippoglossus) has a X/Y genetic sex determination system, but

the sex determining factor is not known. We produced a high-quality genome assembly and

identified parts of chromosome 13 as the Y chromosome due to sequence divergence between

sexes and segregation of sex genotypes in pedigrees. Linkage analysis revealed that all

chromosomes exhibit heterochiasmy, i.e. male- and female restricted meiotic recombination

intervals (MRR/FRR). We show that FRR/MRR intervals differ in nucleotide diversity and

repeat class content and that this is true also for other Pleuronectidae species. We further show

that remnants of a Gypsy-like transposable element insertion on chr13 promotes early male

specific expression of gonadal somatic cell derived factor (gsdf). Less than 4 MYA, this male-

determining element evolved on an autosomal FRR segment featuring pre-existing male

meiotic recombination barriers, thereby creating a Y chromosome. We propose that

heterochiasmy may facilitate the evolution of genetic sex determination systems.

Introduction

The Atlantic halibut (*Hippoglossus hippoglossus*) is the largest teleost species in the North

Atlantic and can reach more than 50 years of age. In recent years, the demand for Atlantic

halibut has been larger than the supply of wild fish. In some areas the stocks have been defined

as vulnerable and sensitive to overfishing, due to relatively slow growth and high age at

maturity. In combination with a high market price, this has led to an increased interest in halibut

aquaculture. Female halibuts can weigh more than 350 kg and become more than 3.6 m long,

while males rarely weigh more than 50 kg. A problem encountered in Atlantic halibut

aquaculture is that males mature earlier and at a smaller size than females[1], and efficient

elimination of male halibut in production of market fish would therefore constitute a

particularly great advantage in halibut compared to other aquaculture fish species. Atlantic

halibut has an XX/XY, male heterogametic, sex determination system[2]. The most common and

practical method to achieve production of all female populations is sex reversal of F1 female

juveniles, to create phenotypic and functional males with an XX karyotype[3,4]. The offspring of

these sex-reversed genetic females will be 100% female, but identifying and separating XX

males from XY males involves raising offspring sibling groups until their sex can be visually

determined by histology or microscopy of gonads[4,5], a time- and resource-consuming process

that can take up to four years. A reliable and practical method for determination of genetic sex

based on identification of Y chromosome markers is instrumental for the development of

efficient protocols for all-female production.

In contrast to the highly conserved mammalian and avian sex determination systems, fish

generally display much more plasticity and turnover when it comes to sex chromosomes and

sex determination systems, with many already described systems being young in comparison

to their mammalian and avian counterparts[6]. Furthermore, teleosts display a rich diversity of

sex-determining mechanisms, both environmental and genetic[7]. As for genetic control of sex,

both XX/XY and ZZ/ZW systems have been described in teleosts, sometimes with closely

related species having differing systems, such as for sticklebacks[8], tilapias[9], swordtail fish[10] and

mosquitofish[11]. Righteye flounders (*Pleuronectidae*) have variable genetic sex determinations

systems with both XX/XY and ZZ/ZW systems represented[12]. The Pacific Halibut

(*Hippoglossus stenolepis*), which diverged from the Atlantic halibut about 3.7 million years

ago[2], has a ZZ/ZW system[13]. Several master sex determining (MSD) genes have been identified

in teleosts, most of which belong to one of three protein families (DMRT, SOX, and TGF-ß

and its signaling pathway)[14], and these have originated either from sub- or neo-functionalization

of duplicated genes, or by allelic diversification[6]. Exceptions to these protein families are found

for salmonids, where *sdy* is the MSD gene[15] and recently the *breast cancer anti-estrogen*

resistance protein 1 (*BCAR1*) and *bone morphogenetic protein receptor type-1B* (*bmpr1b*) have

been reported as MSD genes in catfish (*Ictalurus punctatus*)[16] and Atlantic herring (*Clupea*

harengus)[17], respectively. Thus, a wide range of systems have evolved for genetic control of

sex determination in fish and additional complexity is added by environmental variables

contributing to sex determination in some species.

Single molecule sequencing of long reads has transformed the way the scientific community

produces genome assemblies[18]. In relation to short-read sequencing technologies, methods

capable of sequencing long reads can greatly increase the contiguity of genome assemblies due

to long reads bridging repetitive parts of the genome. In this project we used an Oxford

Nanopore Technologies (ONT) MinION device to sequence long native DNA molecules from

a single male Atlantic halibut and used the assembled contigs in conjunction with Chromatin

Conformation Capture sequencing (HiC) data to produce a chromosome-scale scaffolded

genome assembly. To identify and characterize the sex chromosome we mapped short read data

from male and female DNA pools onto the assembly to screen for divergent genetic markers

between the sexes. The differing systems within the *hippoglossus* genus indicates that at least

one of Atlantic- and Pacific halibut must have evolved a novel sex determination system after

their 3.7 MYA split and that the mechanism for genetic sex determination in Atlantic halibut

may be recent on an evolutionary time scale. Thus, our high-quality Atlantic halibut genome

assembly offers great potential to study the emergence of a young genetic sex determination

system and evolution of sex chromosomes.

Results

Genome assembly and annotation

We used nanopore sequencing in combination with Illumina sequencing of 10X Genomics

Chromium-, paired-end- and HiC libraries to generate a chromosome-scale assembly of the

Atlantic halibut. The final scaffolded assembly (IMR_hipHip.v1) had a scaffold N50 of 27 Mb

(Supplementary Fig. 1a,b) and this assembly was evaluated using BUSCO using 4584 genes in

Actinopterygii odb9 data set, annotating 95.4% of analyzed genes as "Complete" in the genome

assembly. The assembly is organized into 24 larger scaffolds containing 97.6% of the

assembled sequence (Supplementary Fig. 2), corresponding to the chromosomes inferred from

karyotyping[19] and linkage analysis[20]. A number of smaller contigs and scaffolds, together

comprising only 2.4 % of the assembled sequence were left as unplaced. We named the

scaffolds by mapping microsatellite primer sequences from a published genetic linkage map for

Atlantic halibut[20] to IMR_hipHip.v1. Genome to genome alignment of IMR_hipHip.v1 to a

recently finished reference genome of the Pacific halibut (NCBI accession GCF_013339905.1)

revealed a largely co-linear genome organization, with few large-scale disagreements

(Supplementary Fig. 1c) as expected due to the recent split of the two species. To generate

RNA-seq data for genome annotation and to obtain transcriptomic data describing the early

stages of development, we used samples from different time points throughout embryonic

development from newly fertilized to 54 days post fertilization (dpf). For the first four stages

(1 hour post fertilization, 5 dpf, 8 dpf and 12 dpf), four pools of eight individuals were collected

and subjected to RNA extraction and Illumina HiSeq mRNA-sequencing. For the 28 dpf, 43

dpf and 54 dpf, four individual fish were sampled and sequenced for each stage. Genome

annotation based on the RNA-sequencing identified 18,314 genes, of which 18,092 (98.8%)

significantly matched with proteins in public databases.

Identification of the Atlantic halibut sex chromosome

In order to identify and characterize the sex chromosomes we sampled and extracted and pooled

DNA from ten full siblings of each sex and sequenced these using Illumina HiSeq sequencing.

The sequencing data was mapped against the IMR_hipHip.v1 assembly and the sequence

divergence between the male and female pools were determined for non-overlapping windows

of 100 SNPs along each chromosome using χ^2-tests comparing male- and female allele

frequencies. This screen revealed chromosome 13 as the most diverged between males and

females (Fig. 1a), in agreement with a previously published linkage study using RAD-

sequencing[21]. It should be noted that the haploid nature of the IMR_hipHip.v1 assembly makes

it possible that chr13 in this assembly is a hybrid between an X and a Y, which is what we find

when comparing the pool-seq genotypes of male vs. female divergent SNPs. The divergence

between phenotypic males and females on chr13 is not uniform, but overall, gradually declines

along the length of the chromosome, with the end being no more divergent than autosomes (Fig

1b and c).

a   **b**

c

Fig. 1: Genetic differentiation between phenotypic males and females indicates chr13 as the Y-chromosome. **a**

Average χ^2-values for 100 SNPs along each assembled chromosome in the Atlantic halibut genome. **b** Distribution

of average χ^2-values for chr13 compared to the rest of the genome. **c** Average χ^2-values for 500 kb windows along

chr13.

Identification of the Atlantic halibut sex determining gene

RNA-seq data was generated from 12 individuals covering three developmental stages with the

assumption that both genetic sexes would be represented. As we were unable to infer

phenotypic sex for these samples, an unbiased screen could involve searching for the most

variably expressed genes among the 12 individuals, but after conducting a Principal Component

Analysis we concluded that the samples clustered strongly by development stage

(Supplementary Fig. 3). An alternative strategy could involve screening for variable gene

expression between known phenotypic sexes during gonadal differentiation. As known

phenotypic sex was unavailable, we determined the genetic sex of the RNA-seq samples (XY

or XX). We first mapped the DNA-seq pools to the genome and extracted SNPs that had

approximately 50% allele frequency in the male pool-seq while being fixed for one allele in the

female pool. We next mapped the RNA-seq samples to the genome, extracted RNA-seq alleles

at the extracted divergent positions and performed hierarchical clustering (Supplementary Fig.

4) showing clearly that 7 samples were genetically male (XY) and 5 were genetically female

(XX). We then produced a Volcano plot showing the global genetic male vs. female expression

ratio plotted against the statistical significance of a differential expression (DE) between sexes

(Fig. 2a). The most significantly DE gene was *gsdf*, previously shown to be the male-

determining MSD gene in Luzon ricefish and sablefish[22,23]. Furthermore, among genes

previously identified as MSD genes in other fish species[14], only *gsdf* resides on chr13 in Atlantic

halibut (chr13:8506700-8510213). Among 12 RNA-seq individuals the 5 XX fish had almost

no *gsdf* expression, while the 7 XY fish all had high expression, consistent with Gsdf being a

male determining factor. To verify sex specific expression of *gsdf* during development and to

obtain better temporal resolution of expression levels, we performed quantitative PCR (qPCR)

on PCR sex-determined samples (Fig. 2b and c). RNA was extracted from individual embryos

from 11 development stages, ranging from 1 hour post fertilization (hpf) to 13 dpf. A PCR assay

based on SNPs in the 3'UTR of *brx* (chr13:9125004-9125007) was used to determine genetic

sex of samples from 48 hpf and onwards (Supplementary Fig. 5). The qPCR results show that

the expression of *gsdf* starts in males already at 48 hpf, and from 7 dpf onwards there is a clear

increase in expression compared to the earlier stages. For one individual sampled at 6 dpf

determined as a genetic male, we could not detect any expression, possibly due to non-genetic

factors having the chance to overrule the genotype in the determination of phenotypic sex in

the species. *in silico* searches for conserved genes located up- and downstream of *gsdf*

confirmed the *gsdf* identity. (Supplementary Fig. 6).

a

Fig. 2: Identification of *gsdf* as the Atlantic halibut sex determining gene. **a** Volcano plot representation of differential expression (DE) between individuals stratified by chr13 genotyping-based sex assignment. Dot colors indicate the chromosome from which genes are expressed and dot sizes were made proportional to the significance of DE (higher significance=larger dots). The most highly differentially expressed gene, *gsdf*, is indicated in the upper right corner. **b** qPCR result showing the sex specific *gsdf* expression (relative to *gtf3c6*) across developmental stages. **c** Pictures of fish representing the developmental stages sampled for gene expression analyses; 1 hpf to 13 dpf.

Most of the Atlantic halibut genome is subject to male or female restricted meiotic recombination

In order to validate chr13 as the Atlantic halibut sex chromosome we downloaded the pedigree RAD-sequencing data previously used to assign linkage group 13 as X/Y[21], aligned the data to our assembly IMR_hipHip.v1, called SNPs and investigated genotypes by sex phenotype in the pedigrees. The pedigree analysis verified our divergence-based assignment of IMR_hipHip.v1 chr13 as the sex chromosome and 90/94 F1 individuals were in agreement with regard to the

association between phenotypic sex and chr13 genotype. A similar observation was made for the embryonal qPCR analysis where one out of 34 genetic males (*brx*-assay) had no *gsdf* expression. The reason for the non-perfect association between genetic- and phenotypic sex could be erroneous annotation of the downloaded RAD-seq dataset or a logical consequence of non-perfect penetrance of young genetic sex determination systems. In our genome assembly-guided linkage analysis, we detected suppression of meiotic recombination along a large part of chr13 in male germ cells as would be expected in an X/Y system. Surprisingly, the majority of the sex chromosome pair exhibited strict male- or female restricted meiotic recombination (MRR/FRR) (Fig. 3a, Supplementary Fig. 7a). The male- and female genetic maps both sum up to 49 cM but the recombining intervals are nonoverlapping. None of the 90 F1 offspring used for analysis carried a male meiotic recombination in the first ~15 Mb, while all detected female meiotic recombination events on chr13 occurred in this region. Thus, the first ~15 Mb of chr13 behaves like a X/Y chromosome, with high male vs. female divergence (Fig. 1c) and nonexistent meiotic recombination in males. The later part of chr13 (~15 Mb until chromosome end) only recombines in males (X/Y recombination) and has lower male vs. female divergence, suggesting that sequence diversity in this part of the chromosome is shared between X and Y. Even more surprising than the meiotic recombination barrier observed for the of chr13 in females was the observation that all chromosomes clustered into more or less strict MRR/FRR regions (Fig. 3b, Supplementary Table 1 and Supplementary Fig. 7a). We limited our analysis to the placed chromosome part of the genome (chromosomes 1-24), spanning 596 Mb out of the total 611 Mb, and defined FRRs and MRRs according to our classification outlined in Supplementary Table 2. Genome-wide, this annotation defined 407 Mb (68% of the sequence) as FRRs and 154 Mb as MRRs (26% of the sequence) and only ~6% of the sequence fell outside of this classification, i.e. either both sexes or neither sex showed meiotic recombination.

Fig. 3: Cumulative proportion of all recombination events occurring during male and female meiosis in a two-pedigree intercross. **a** chr13 with *gsdf* location indicated. *Mapping interval of markers most highly associated with sex in a previous study[21]. **b** Male and female linkage maps differ strongly. All 24 chromosomes exhibit clear sex restricted recombination intervals.

Nucleotide diversity and repeat content is correlated with sex-restricted recombination

We investigated what distinguishes an MRR from an FRR and concluded that for all chromosomes except chr9 MRRs are much smaller than FRRs while retaining approximately the same number of recombination events (Supplementary Fig. 7b). Furthermore, MRRs, again with the exception of chr9, exhibit higher nucleotide diversity (π) than their FRR counterparts

(Fig. 4a). Recombination rate is correlated with π (Supplementary Fig. 8) and chr13 is an outlier

with the largest MRR over FRR π ratio among all chromosomes, presumably due to the limited

effective population size of its FRR, the herein identified Y-chromosome. Typically, MRRs

occupy one end of a chromosome and FRRs start from the other end. Atlantic halibut has been

shown to have subtelomeric or acrocentric centromeres[19] and for 20 out of 21 chromosomes

where we could infer centromere placement by mapping centromeric markers[20] to

IMR_Hiphip.v1, the female end coincided with the centromere (Supplementary Table 3). We

next sought to determine whether meiotic recombination is sex restricted also in species closely

related to Atlantic halibut but could not find any publicly available pedigree datasets for other

Righteye flounders. Since Atlantic halibut π was found to be higher for MRRs than FRRs, we

decided to investigate whether π is correlated with MRR/FRR classification also in other

Righteye flounder species. We could not find any publicly available genome sequencing data

for Pacific halibut and instead used RNA-sequencing data to estimate π for this species. For

Greenland halibut (*Reinhardtius hippoglossoides*) and European plaice (*Pleuronectes platessa*)

we estimated π from publicly available RAD-seq- and whole genome shotgun sequencing data,

respectively. The across-species comparison of π distribution in MRRs and FRRs revealed a

striking agreement among the four species analyzed, with regions corresponding to Atlantic

halibut MRR having higher π (Fig. 4a-c)*,* indicating that heterochiasmy has been conserved

among righteye flounders.

Fig. 4: Nucleotide diversity (π) and repeat content is correlated with sex-restricted recombination. **a** π distribution

in the Atlantic halibut genome, contrasting regions of male- vs. female restricted recombination, MRRs and FRRs,

respectively. **b** Distribution of Pacific Halibut π estimates for regions mapped back to the Atlantic halibut genome,

classified as either MRRs or FRRs. Chromosomal and MRR/FRR assignments were extracted from the Atlantic

halibut genome assembly coordinates. **c** Genome-wide distributions of estimated π for Atlantic-, Pacific- and

Greenland halibut and European plaice in regions corresponding to herein defined Atlantic halibut MRRs and

FRRs. Numbers above MRR distributions indicate, within species, how much higher π is in MRR regions

compared to FRR regions. * The time since divergence of Greenland halibut from *Hippoglossus* was estimated

from[24]. ** Divergence estimate for European plaice was obtained from TimeTree[25]. **d** Densities of repeat classes

and π for 500 kb windows along each Atlantic halibut chromosome. Class color indicates whether a region was

annotated as FRR, MRR or was unassigned. LINE and SINE represent Long- and short interspersed nucleotide

elements, respectively. LTR indicates Long Terminal Repeats. LCR and SR indicate low complexity- and simple

repeats, respectively. π indicates nucleotide diversity. The white to grey scale in heatmaps is proportional to the

min. and max. values within each class.

We investigated whether MRRs and FRRs differed for repeat content and annotated repeats in

IMR_Hiphip.v1 by means of RepeatMasker[26] using the *Actinopterygii* database. RepeatMasker

raw output was parsed by repeat subclass and the density per subclass was investigated for each

chromosome (Supplementary Table 4). Overall, Long- and Short Interspersed Nucleotide

Element (LINE and SINE) densities increase towards the FRR end, while simple repeats and

low complexity repeats show the opposite pattern, being enriched towards the MRR end (Fig.

4d). Chr13 and chr10 (Supplementary Fig. 9 and 10) are both typical with respect to clustering

of major repeat classes into MRRs/FRRs. Chr10 is the only example where the MRR end

overlaps the predicted centromere, suggesting that the clustering and accumulation of repeats

is shaped by evolutionary forces associated with sex restricted recombination rather than

centromere location.

The insertion on chromosome Y is of transposon origin and contains a *gsdf* promoter

In an effort to identify candidate Structural Variants (SVs) between Y and X that may influence

gsdf expression, we aligned all the ONT reads from the male individual used for genome

assembly to the IMR_hipHip.v1 assembly and used the software Sniffles[27] to call SVs along

the genome. This analysis revealed a large set of putative SVs (Supplementary Table 5), two of

which coincided with the *gsdf* locus, both detected as heterozygous deletions in relation to the

reference at chr13:8503901-8505128 (1,227 bp) and chr13:8509688-8509767 (79 bp). Manual

inspection of these hits revealed that both ONT reads and linked read pseudo-haplotypes from

the male assembly individual validate the assignments made by Sniffles. We used alignments

of the male and female pool sequencing data and the phases of ONT long reads to conclude that

the insertion allele at these two indels is present on chrY. The HiC data used for scaffolding the

assembly was used to screen for large-scale SVs that may differ between X and Y. This analysis

indicated that no large inversions are present on chr13 and that neither Y nor X are part of any

large translocations with other chromosomes, as such events would appear as off-diagonal

contacts in the contact map (Supplementary Fig. 11).

The 1.2 kb insertion upstream of *gsdf* (INS1) is not found in the corresponding Pacific halibut

locus, supporting that this region is derived on the Y chromosome in Atlantic halibut. INS1

(IMR_Hiphip.v1 chr13:8503902-8505127) is highly similar to 24 other loci in the genome

assembly. Sequence alignments showed high sequence similarity starting with TG and ending

with CA di-nucleotides in 19/24 loci. The remaining 5 loci showed strong, further extending

similarities to each other, and we therefore extracted additional sequence flanking these five

loci to perform a 7-8 kb alignment, which also terminated in conserved TG/CA di-nucleotides

in four out of the five sequences. The TG/CA-ends are hallmarks of retrotransposon long

terminal repeat (LTR) boundaries, and we could also identify 4-6 nucleotide sequence

duplications immediately flanking the insertions, indicating target site duplications (TSDs) that

confirm the transposable element (TE) insertions as retrotransposons. Although these TE

hallmarks were obscured at the INS1 site upstream if *gsdf*, the sequence similarity and

alignment to the related loci in the genome support its TE origin (Supplementary File 1). One

of the five longer sequences (IMR_Hiphip.v1 chr14:3028070-3045206) was used as query to

search the DFAM database[28]. This search revealed a 5254 nt long hit against the 5410 nt long

Danio rerio Gypsy transposable element (Gypsy6-I_DR, Bit score 2342), supporting by

sequence similarity association that INS1 was derived from a Gypsy retrotransposon that

inserted in antisense orientation relative to *gsdf*. Strong similarities between the full-length

Gypsy LTR sequences and INS1 revealed that INS1 is indeed an LTR remnant of an ancient

Gypsy insertion. This association is further supported by RepeatMasker analysis of the five

putative full-length hits partially matching INS1, as they all contained Gypsy sequence

annotations.

From the ONT data used for genome assembly we could assess DNA methylation at the *gsdf*

locus. Interestingly, we observed low DNA methylation within INS1, in a region coinciding

with a predicted CpG-island (Fig. 5 and Supplementary Fig. 12). Further inspection of the lowly

methylated CpG-island revealed predicted Sp1 binding sites indicating that this locus may act

as a promoter region for *gsdf*. It is clear from both ONT read alignment and the linked read

pseudo-haplotype alignments that the chrY haplotype indeed carries INS1. We developed a

PCR assay to amplify into INS1 and observed that this assay gave clear bands of the expected

size for 10 adult phenotypic males while no bands were observed for 10 phenotypic females

(Supplementary Fig. 13). Expression of embryonal *gsdf* (28-54 dpf) from the Y-allele (*gsdf*-Y)

extends further upstream of the transcription start site than gene models both in adult Atlantic

halibut testis and in male- and female reproductive tissues in adult Pacific halibut. The longer

gsdf-Y isoforms contain 5'UTRs extending into the Gypsy LTR INS1 (Fig. 5 and

Supplementary Fig. 12), which supports promoter activity of INS1.

Fig. 5: Chr. Y carries a transposon-derived LTR upstream of *gsdf*. Schematic representation of findings. Male and female chr13 haplotypes differ for a 1.2 kb transposable element derived segment 2 kb upstream of the *gsdf* transcription start site. Males carry the insertion allele which contains a lowly methylated CpG-island inferred to act as a derived promoter of *gsdf* uniquely in males. The shorter *gsdf* isoforms show the extent of *gsdf* expression observed in adult testis in Atlantic halibut as well as adult ovary and testis of Pacific halibut. The two *gsdf* gene models labeled chr13Y-specific indicate isoforms detected only in males from three separate developmental stages. An IGV-plot of this region is presented in Supplementary Fig. 12.

Analyzing chrY decay

In addition to the ONT genome assembly, we also sequenced the same male using a Chromium Genome linked read library and used the resulting data to generate two separate pseudo-haplotype assemblies which successfully phased the first 10 Mb of chr13 into separate scaffolds for the X and the Y chromosome. Loss of X/Y recombination is expected to lead to chrY decay, and we investigated whether we could find evidence of this using the phased chrX (X^{PH}) and chrY (Y^{PH}) along the first 10 Mb of chr13. This analysis, which involved quantification of possibly deleterious variants along the first 10 Mb of X^{PH} and Y^{PH}, revealed a 30% excess of variants predicted to be deleterious (HIGH classification in SNPEff) on Y^{PH} (n=35) compared to X^{PH} (n=27). That no large-scale loss of genetic material has occurred on chrY is supported by the observation that super-scaffolds of X^{PH} and Y^{PH} were both 20.4 Mb in size.

Discussion

In this study we show that *gsdf* is most likely the master sex determining gene. Our interpretation is that following an ancient Gypsy element insertion on the autosomal progenitor of chr13, this element degraded and left a remnant LTR, which now acts as a promoter driving the expression of *gsdf* ectopically during development of XY individuals, making carriers develop into phenotypic males. Interestingly, In Luzon ricefish, where *gsdf* is also the MSD gene, the region immediately upstream of *gsdf* was shown to be most important for modulating *gsdf* expression[22]. The region tested corresponds approximately to the site of TE insertion in Atlantic halibut. A role of *gsdf* in early gonadal development is supported by experiments in medaka where targeted homozygous deletion of *gsdf* in genetic (XY) males triggers an early development of ovaries, with the majority of these *gsdf* KO homozygote fish still developing testes later in life[29]. It would be interesting to use CRISPR/Cas9 to remove the Gypsy LTR

from XY Atlantic halibut (X/Y^{LTR-}) and insert the Gypsy LTR into XX individuals X/X^{LTR+}),

as we have done in Atlantic salmon[30,31]. Our predictions are that X/Y^{LTR-} individuals would

develop as females and X/X^{LTR+} individuals as males. Furthermore, in accordance with previous

studies[32], we observe *gsdf* expression in adult testes and ovaries in Pacific halibut and in

Atlantic halibut testis, indicating a function of Gsdf in mature gonads. In medaka ovaries and

testes, *gsdf* is expressed by somatic cells surrounding germ cells, with higher expression in

testis[29]. In Atlantic salmon *gsdf* expression has been detected in Granulosa and Sertoli in ovaries

and testes, respectively[33]. This supports a role for Gsdf in the mature gonad and suggests that

high expression of *gsdf* from Atlantic halibut chrY during development is expected to trigger

testis development. TEs are known to be involved in sex-chromosome evolution[34], e.g. TEs are

probably responsible for translocation of *sdY* in salmonides[35] and in Medaka a LINE insertion

at the *dmrt1b* locus has been implicated as the sex-determining variant[36]. It has been repeatedly

observed that TEs are repurposed for genomic innovation by domestication[37], as might be

expected for a dynamic part of the genome.

In this study we observed that meiotic recombination is dramatically different between male

and female gonads, and this can also be seen when revisiting a previous Atlantic halibut linkage

study[20], where this phenomenon was not strongly emphasized. Interestingly, the guppy

(*Poecilia reticulata*) has genome-wide heterochiasmy, while no significant difference was

observed in the closely related genus *Xiphophorus*[38], suggesting that the mechanism underlying

heterochiasmy may be easily modified over short evolutionary time-spans. A common feature

for observations of heterochiasmy in teleosts is enrichment of male crossovers towards

chromosome ends[38-40], which is also the case in Atlantic halibut. We found that four analyzed

righteye flounder species all had higher π in the regions orthologous to Atlantic halibut MRRs

than in corresponding FRRs orthologous regions (Fig. 4c). This likely reflects conservation of

sex restricted recombination across *Pleuronectidae*. The reason for this interpretation is that

few polymorphic sites are expected to be shared between species over the millions of years

separating the four analyzed species. Thus, the diverging π distributions must have been

established independently, presumably by selective forces associated with heterochiasmy.

In medaka, sex restricted recombination between sex chromosomes has been shown to follow

the XY male pattern also in XX neomales[41], which supports that a mechanism associated with

the gonadal environment governs recombination suppression in meiosis. We propose to study

the mechanisms of heterochiasmy in sex-reversed genetic female Atlantic halibut, which are

used as breeding males in aquaculture. Regions of TE-accumulation in genomes have been

implicated as putative heterochromatin sinks[42], and both TEs and heterochromatin are

positively correlated with suppressed recombination[43]. It is difficult to disentangle whether TEs

are involved in the establishment of meiotic recombination suppression or if their differing

distributions in MRRs/FRRs is merely a consequence of how the genome evolves due to

unknown selective pressures imposed by heterochiasmy already being in place. One possibility

could be that Atlantic halibut male and female gonads differ for which regions of the genome

become heterochromatin, perhaps as a defense against TE transposition, and that male and

female gonads direct recombination differently in response to variable TE content along the

chromosomes. Possibly, transposable elements are more likely to accumulate in FRRs, for

example if regions only recombining in males (MRRs) exist in a state less accessible for

transposition during female and/or male meiotic recombination. Accumulation of local TE

clusters in FRRs may also contribute to the further homogenization of this chromosome

compartment by gene conversion of homologous TE insertions. It should be noted that we have

analyzed repeats broadly, and future studies aiming to investigate repeat classifications in

relation to heterochiasmy may benefit from more detailed classifications. We show that π is

higher in MRRs than in FRRs and hypothesize that the higher recombination rate on the

physically smaller MRRs increase the probability of retaining diversity at linked sites during

purifying selection[44], or that differences in purifying selection due to the differing TE-

composition landscapes of FRR and MRRs has been involved in shaping π.

Sex-restricted meiotic recombination, as we describe here, suppresses recombination for large

segments along autosomes. Due to genome-wide, pre-existing sex-restricted recombination

being in place, this system is expected to facilitate the survival of novel sex determining factors

which in turn could give rise to new, and even competing genetic sex determination systems

(Supplementary Fig. 14). We propose that heterochiasmy has the potential to facilitate evolution

of sex determination systems, and presumably speciation as postulated in stickleback[45]. It has

been reported that female is the heterogametic sex (ZZ/ZW) in both Pacific halibut and

European plaice[12] as well as the half-smooth tongue sole[46], with the main candidate MSD gene

being *dmrt1*. This makes it conceivable, but not certain, that the Atlantic halibut X/Y system

replaced a Z/W system shared across different flatfish species.

Regardless of heterochiasmy being a facilitating factor in the evolution of sex determination

systems or not, we conclude that the TE insertion in the *gsdf* locus occurred after the split

between Atlantic halibut and Pacific halibut around 3.7 MYA. Sex chromosomes originate from

autosomes that acquire a sex determination locus and then lose recombination with the former

autosomal pair, which typically involves structural variation in the non-recombining region of

the sex chromosome pair, while other regions may still recombine and act as pseudo-autosomal

regions. For mammalian Y, as well as avian W, loss of purifying selection over millions of

years has led to pseudogenization, gradual loss of genes and a reduction in chromosome size[47],

where remaining genes and regulatory elements determine male and female phenotypes,

respectively. Our analysis of deleterious alleles identified a non-significant 30% increase in

predicted deleterious alleles on chrY compared to chrX, which may indicate an early stage of

chrY decay. We observe a 1 Mb segment (chr13:5.5-6.5 Mb) (Fig. 1c) showing differentiation

equal to the genome average, interpreted as X/Y recombination, although rare, not being fully

suppressed in the FRR of chr13. This is supported by previous fine mapping of the sex

determining locus to a smaller marker interval[21] (Fig. 3a) in a region corresponding to the herein

defined chrY. A retrotransposon insertion into the *gsdf* locus started the transformation of an

autosome into a sex chromosome in Atlantic halibut less than 4 million years ago. Incomplete

recombination barriers and little, if any, chrY decay both suggest that Atlantic halibut chrY is

in the early stages of the evolutionary trajectory of a sex-chromosome.

Methods

Ethics statement

For this study the halibut were reared under conditions similar to standard commercial fish

farming conditions. Such conditions are listed as an exception in The Norwegian Regulation on

Animal Experimentation, thus approval of the experimental protocol of this experiment by

NARA (the governmental Norwegian Animal Research Authority) was not required.

Halibut material

All samples were taken during routine production of Atlantic halibut at the Austevoll Research

Station, Institute of Marine Research, Storebø, Norway. Halibut breeding individuals were

stripped, eggs fertilized and incubated, and juvenile halibut produced according to standard

aquaculture protocols developed at the Austevoll Research Station[48-50]. Briefly, females were

stripped according to their individual ovulatory rhythms to ensure high (>90%) fertilization.

Fertilized eggs were incubated in 250 l conical incubators with upwelling water for 60 day

degrees (i.e. 10 days at 6ºC) and transferred to larger incubation units, silos (vol. 50,000 l)

before hatching. Yolk sac larvae were held in the silos until ca. 265 day degrees posthatch when

they were moved to first feeding tanks and fed *Artemia* for 70 days until metamorphosis was

complete. At this stage, the larvae were weaned onto a formulated diet. Juvenile halibut were

moved to ongrowing tanks when they had reached an average size of ca. 10 cm and held there

until they had reached a harvest size of 2-5 kg. One adult, three-year old male was euthanized

with metacain (100 mg/l) and muscle, spleen and blood samples were taken for reference

genome sequencing. For sequencing of pooled family samples, tissue samples were taken from

ten full siblings of mixed sex (ten males and ten females). All samples were snap frozen in

liquid nitrogen (N_2) and stored at -80ºC before DNA extraction. RNA-sequencing was made on

a series of samples taken through embryonic development. Before hatching, embryo samples

were snap frozen in liquid N_2 and stored at -80ºC. Larvae were euthanized in metacain (100

mg/l), transferred into RNA later, kept at 4ºC for 24 h and stored at -20ºC until RNA isolation.

The samples for qPCR were snap frozen in liquid N_2 and stored at -80ºC. Digital images of all

embryo-larval stages were captured using video cameras Moticam 1080 (Motic) mounted on

Olympus SZX10 stereomicroscope using Motic Live Imaging Module. Living embryos and

larvae were individually immobilized in a Petri dish containing 2% methylcellylose/98%

seawater. Images were then scaled and processed by ImageJ software[51].

DNA extraction and sequencing of male and female pools

DNA was extracted from muscle samples using Qiagen DNeasy Blood and Tissue Kit. Paired-

end libraries of equimolar male (10 individuals) and female (10 individuals) DNA pools and

from one single individual of each sex were constructed using the Genomic DNA Sample

Preparation Kit (Illumina) according to manufacturer's instructions and sequenced on the

Illumina HiSeq2000 platform (Illumina) at the Norwegian Sequencing center

(https://www.sequencing.uio.no, Oslo, Norway).

High Molecular Weight DNA extraction and sequencing

Several tissues were collected from a single male Atlantic Halibut. Muscle and spleen were

snap frozen in liquid nitrogen and stored at -80ºC until further use. Fresh blood was used to

isolate erythrocyte nuclei and then prepare High Molecular Weight (HMW) DNA from these

nuclei according to a Phenol/Chloroform protocol (Quick). Extracted DNA was either left

unfragmented or sheared using either MegaRuptor (Diagenode) or by passing DNA through a

100µl pipette tip 10-20 times. From these DNA samples we produced MinION sequencing

libraries with the standard adapter ligation protocol kit (LSK-108), 1D^2 sequencing kit (LSK-

308) and the RAD002 and RAD003 kits (Oxford Nanopore Technologies). For libraries

prepared with the RAD002 and RAD003 kits we used a protocol developed to produce ultra–

long reads (Quick, protocols.io). In total we used 9 flow cells and sequenced 18.8 Gbp in

2,963,108 reads with a read N50 of 17.8 kb. Sequenced fast5 raw signal data files were

basecalled using albacore v.2. The size distribution of sequenced reads is presented in

Supplementary Figure 1a. We also used the collected muscle samples to produce HiC libraries

(Dovetail), sequenced on one lane of HiSeq X (Illumina). One of the DNA preps inferred to

have HMW DNA was used to prepare a linked reads Chromium library (10x genomics) and

was sequenced on one lane of HiSeqX (Illumina).

Genome assembly, polishing and scaffolding

Nanopore reads passing filters were used in assembly by first constructing all-vs-all alignments

between all reads using minimap2[52]. These all-vs-all alignments were then used in the software

miniasm to produce a haploid contig assembly. The contig assembly was polished using racon[53]

with the nanopore data followed by pilon[54] using the Illumina data from the male pool.

Scaffolding of the polished miniasm assembly was performed using the HiRise pipeline

(Dovetail Genomics). The quality of the generated assembly was assessed using BUSCO[55].

Initial nanopore contigs assembled with miniasm[52] had a contig N50 of 3.2Mb and the resulting

final assembly had a scaffold N50 of 27.2 Mb.

Assembly of linked read data

The software Supernova2[56] (10X Genomics) was used to produce a genome assembly of the

sequencing data resulting from the Illumina sequencing of the Chromium library. For this

assembly, reads were subsampled to get close to the recommended 56x genome coverage

recommended in the Supernova2 user manual. The two pseudo-haplotype linked read

assemblies featured one scaffold each which covered the whole of IMR_Hiphip.v1 chr13. We

used SNP markers identified as divergent between the male and female short read data to

investigate the pseudo-haplotypes in relation to sex divergent markers, i.e. if one pseudo-

haplotype could represent the Y chromosome and the other the X-chromosome. This analysis

revealed that for the first 10 Mb of chr13, the allele being heterozygous in males and not present

in females (the perceived Y-allele) was observed on pseudo-haplotype number 2 (1,954 SNPs).

The perceived Y-allele was observed only for 2 SNPs in this region for pseudo-haplotype

number 1. The corresponding counts for the rest of chr13 was 125 and 213 alleles on pseudo-

haplotype numbers 2 and 1, respectively.

RNA-sequencing and gene annotation

Total RNA was extracted from whole larvae from seven different stages of development using

miRNeasy Mini Kit (Qiagen) and treated with TURBO DNase-Free kit (Ambion). TruSeq

cDNA libraries were sequenced on the HiSeq 4000 sequencing platform (Illumina). The

average number of reads was 50,320,639 per sample. Twenty-two samples (four samples from

54 dpf and three samples from each other stage) were assembled individually with Trinity using

the Agalma pipeline[57]. The transcriptome assemblies were annotated with BLAST against

Swissprot (download date 20191227) and the Ensembl gene models of zebrafish (GRCz11) and

stickleback (BROAD S1). The hits with the best scores were kept. Transcripts from all

assemblies were mapped to the halibut genome with BLAT. The thresholds for mapping were

set to 98% match for sequences with length >=401 nt and 95% match for sequences with length

<=400 nt. Long sequences were mapped before short sequences. The mapped sequences were

clustered on genome location, selecting the sequence with the highest score for each non-

overlapping cluster. Only clusters with an annotation (>300 in BLAST score threshold) were

saved. Neighboring clusters annotated with the same gene from either Swissprot, zebrafish or

stickleback were merged in cases where the annotation indicated that the gene was split in

several clusters (the neighboring clusters were only merged in cases where the annotation was

from different parts of the same gene). For the merged clusters, the sequence with the highest

BLAT score from all merged clusters were chosen. Additionally, the transcriptome assembly

sequence with the highest scoring BLAT match to the merged cluster was added. We found

15,273 annotated genes based on the Trinity assemblies. The zebrafish and stickleback

ENSEMBL gene models were mapped to the halibut genome using exonerate in order to add

genes without RNA-seq evidence. The exonerate hits were clustered by location, using the

exonerate hit with the highest score for each cluster. Sequences for each exonerate based gene

were extracted from the halibut genome sequence. 2,269 genes were added based on the

exonerate analysis. We added 222 genes that had no annotation (BLAST threshold score <300)

but had RNA-seq expression. After running BUSCO 3.1 with the Actinopterygii odb9 dataset

on the non-filtered Trinity assemblies that mapped to the genome[55], we added 550 genes from

the BUSCO analysis that were missing from our filtered list. This resulted in a total number of

18,314 genes and 20,579 transcripts giving a 80.9% complete, 8.7% fragmented and 10.4%

missing BUSCO coverage.

Nanopore sequencing of testis RNA

Testis tissue was homogenized at 4°C in 1ml QIAzol Lysis Reagent (Qiagen, Hilden, Germany)

together with RNase-Free Zirconium Oxide Beads (NextAdvance, Inc., Troy, NY, USA) during

1 min, maximum effect, in a BulletBlender (NextAdvance) and total RNA was prepared

according to the QIAzol (Qiagen) protocol. Polyadenylated RNA was isolated with the

Dynabeads mRNA direct purification kit (Thermo Fisher Scientific), following the kit protocol.

For full-length cDNA Oxford Nanopore sequencing, the cDNA-PCR Sequencing SQK-

PCS108 kit (Oxford Nanopore Technologies) was used according to the kit protocol with minor

modifications. The finished cDNA library was sequenced on a MinION device using a R9.4

flow cell (Oxford Nanopore Technologies).

Screening for male vs. female divergence

Illumina sequences from the male- and female DNA pools were filtered by trimmomatic[58], then

aligned to the assembly IMR_hipHip.v1 using BWA-MEM[59]. Aligned reads were subjected to

duplicate flagging ("Picard Toolkit." 2019. Broad Institute, GitHub Repository.

http://broadinstitute.github.io/picard/; Broad Institute) and were then used as input for calling

SNPs as well as small Insertions/Deletions using Genome Analysis Toolkit (GATK)[60]

UnifiedGenotyper. Raw variant vcf files were subjected to variant filtration using Best Practices

procedure with SNP-clusters being added as an additional filter (more than 5 SNPs in a window

of 20 bases were filtered out). SNPs passing filters were isolated from the vcf file and allele

frequencies from the male- and female pools were extracted and used in χ^2-analysis examining

the null hypothesis; that allele frequencies were not different between the male- and female

pools. Prior to the χ^2-test, in order not to inflate the significance at positions with high depths

of coverage, SNP read depths >12 for either pool were set to 12 while retaining the observed

allele frequency ratio. χ^2-tests were then performed SNP-by-SNP, and average χ^2-scores of

non-overlapping windows of 100 adjacent SNPs along the chromosomes were extracted.

Sex differentiating PCR assay

595 Genetic sex of each embryo was determined by PCR using sex-specific primers (Table 1). The

596 thermal cycling conditions were as follows: 94°C for 5 min, 32 cycles with 94°C for 30 sec,

597 65°C for 1 min and 72°C for 1 min, and finally 72°C for 5 min. Each 12 µl PCR reaction

598 contained 5X GOTaq Flexi Buffer, MgCl, dNTPs, GoTaq Polymerase, PCR primers and

599 template cDNA.

Differential gene expression between sexes

We generated RNA-sequencing data from 12 Atlantic halibut whole embryos representing

stages 28 (n=4), 43 (n=4) and 54 (n=4) dpf. Since one cannot determine the phenotypic sex at

this stage we aligned the RNA-seq data using Bowtie 2[61] to IMR_hipHip.v1 and extracted

expressed allele counts at positions identified as divergent between males and females in the

pool DNA-seq. Extracted RNA-seq allele ratios were used to cluster the individuals, revealing

two well-separated clusters of XY and XX individuals and this assignment was used to identify

genetic male and female individuals for the differential expression screen. The differential

expression screen involved contrasting across-sample normalized gene expression levels of the

seven inferred males to the five inferred females in a two-tailed t-test. Principal Component

Analysis (PCA) was performed, using the R function "prcomp", on the normalized gene

expression levels from the 12 embryos.

gsdf **quantitative PCR**

Genomic DNA and total RNA was extracted from single embryos using the Allprep

DNA/RNA/miRNA Universal Kit (Qiagen), according to the manufacturer's instructions. The

RNA was DNase treated as part of the extraction process, according to the manufacturer's

instructions. For each sample, cDNA was synthesized from 125 ng RNA using the Superscript

VILO cDNA synthesis Kit (ThermoFisher Scientific). The primers used for qPCR were

designed using BatchPrimer3 (https://wheat.pw.usda.gov/demos/BatchPrimer3/) (Table 2). The

gene *gtf3c6* was used as an internal reference gene for normalization of the qPCR data due to

its stable expression in the developmental stages analysed in this study (Supplementary Fig.

15). Using different RNA concentrations, the PCR efficiencies (E = -1+10(-1/slope)) for the

primers were calculated to be 108.5% (*gsdf*) and 92.4% (*gtf3c6*). A melt curve analysis was

performed to confirm that the primers did not amplify more than one PCR product. qPCR was

performed in duplicates in 384-well optical plates in a QuantStudio 5 Real-Time PCR system

(ThermoFisher Scientific) using default settings. 2 µl cDNA (diluted 1:20) was used in a 6 µl

Fast SYBR Green qPCR reaction (ThermoFisher Scientific). No-template controls for each

gene were run in all qPCR plates. The relative gene expression level was calculated using the

comparative Ct (or $2^{-\Delta\Delta Ct}$) method. All values were normalized to *gtf3c6* and calibrated to the

sample with the lowest ΔCt.

Detection of structural variants, small insertion/deletions and annotation of repeat

sequences

The long-read SV identification tool Sniffles[27] was used to identify SVs from the ONT data.

The ONT data from the single male individual used for genome assembly was aligned to

IMR_hipHip.v1 using minimap2[52] with the "-x map-ont" option and the resulting bam file was

used in sniffles, requiring >=5 observations of a variant allele to be retained as a potential SV.

Analysis of chrY decay

The two linked-read pseudo-haplotype assemblies of the male Atlantic halibut used for the

genome assembly were aligned against the Pacific halibut reference genome (nucmer) and

SNPs and small indels were called using the "show-snps" function of the MUMmer 4.0

package.　　Identified　　variants　　were　　converted　　to　　vcf-format

(https://github.com/MatteoSchiavinato/Utilities/blob/master/my-mummer-2-vcf.py) and were

then annotated using SNPEff, with gene models derived from the Pacific halibut

IPHC_HiSten_1.0 gtf-file "GCF_013339905.1_IPHC_HiSten_1.0_genomic.gtf". SNPs and

indels that were unique to either pseudo-haplotype were investigated for their annotations.

Annotation frequencies were compared between the two pseudohaplotypes for the interval

where they represent the X and Y (Atlantic halibut: chr13:0-10Mb, Pacific halibut:

NC_048944.1:0-11Mb).

Analysis of the evolutionary origin of the 1.2 kb insertion upstream of *gsdf*

The 1.2 kb insertion (IMR_Hiphip.v1 chr13:8503902-8505127) was used as a query sequence

in BLAT[62] searches against IMR_HipHip.v1. Full length hits were retained (n=30) and were

extended to include 2kb up- and downstream sequences and these approximately 5.3 kb

sequences were used as input for multiple sequence alignment using MUSCLE[63]. The resulting

alignments were visualized in AliView[64] and abrupt drops in sequence similarities were

observed for 24 of the sequences on each flank. The extent of highly similar sequences started

and ended at perfectly conserved TG and CA di-nucleotides, the hallmark motifs of LTR

boundaries, and the insertion was immediately flanked by 4-6 nucleotide duplications of loci

TE target sites (TSD) confirming the LTR-retrotransposon insertion.

DNA methylation calling from ONT data and CpG island prediction

Basecalled ONT reads previously used for the genome assembly were aligned back to the

IMR_Hiphip.v1 assembly using minimap2. The software nanopolish[65] was used to index the

raw ONT fast5-files to enable extraction of likelihoods of CpG site methylation along

individual mapped reads in genome assembly coordinates. Next, the nanopolish script

"calculate_methylation_frequency.py" was used to extract the methylation frequency at each

adequately covered CpG cluster in the assembly. CpG islands were predicted for the

IMR_Hiphip.v1 assembly using gcluster[66] employing default settings.

Estimation of nucleotide diversity

For Atlantic Halibut we estimated π by creating chromosome-specific samtools mpileup files

from the aligned data previously used for variant calling, using only positions with a mapping

quality >=20 and read depths ranging between 10 and 50. The script "WinNuclDivmPileup.py"

(https://github.com/cjrubinlab/python_scripts2) was used to estimate π in 500kb windows

along the genome for the male- and female pools. For Pacific halibut no DNA-sequencing data

was available in public sequence repositories, so we devised a way to estimate π from RNA-

sequencing data. Sequencing data from 12 RNA-seq experiments of various tissues and

individuals was downloaded from the SRA (SRR11826178-SRR11826189) and these data were

aligned to the Oxford nanopore derived Pacific Halibut genome assembly (GCF_013339905.1)

using the STAR aligner[67]. The resulting alignments were randomly subsampled 20 times and

then subjected to duplicate-pair removal using picard-tools MarkDuplicates, retaining 70% of

the reads after filtering. From the bam file containing filtered reads an mpileup file was

generated using samtools (mapping quality exclusion filter < 20). For the Greenland Halibut

(*Reinhandtius Hippoglossoides*) we downloaded the genome assembly from NCBI

(GCA_006182925.2_MU_Rhippo_v1) as well as RAD-sequencing data from 20 individuals

(SRA accessions: SRR11351808-SRR11351827), aligned the short-read data to the assembly

using minimap2 with the "-x sr" option. From mpileup files derived from samtools mpileup for

Pacific Halibut and Greenland halibut the estimated heterozygosity (He) was calculated for

positions having coverage>=10 and mapping quality >=20 using the WinNuclDivmPileup.py"

script. For European plaice (*Pleuronectes platessa*), a species lacking a genome assembly, we

downloaded paired genome resequencing Illumina reads (SRA accession: SRX8240416),

aligned these reads to the IMR_Hiphip.v1 assembly using bwa and proceeded with the same

pipeline as was used for estimation of Atlantic halibut π. In order to lift over genomic intervals

annotated as MRRs and FRRs in the Atlantic halibut to the Pacific- and Greenland halibut

assemblies we used the satsuma[68] commands "Chromosemble" to generate a detailed list of lift-

over intervals and "BlockDisplaySatsuma" to summarize the detailed list into coherent blocks.

We then converted these coordinates to the bed-format and used the bedtools[69] "intersectbed"

command to link coordinates in the other assemblies to the corresponding coordinates in the

Atlantic halibut assembly and MRRs and FRR classification.

Linkage analysis

RAD-sequencing data from an Atlantic halibut half-sib pedigree altogether comprising three

parents and 90 offspring were downloaded from the SRA (accession=SRX193134). The

downloaded individual files were split by barcodes and were then connected with annotation

information from the original publication regarding phenotypic sex, parent/offspring status and

family B/C[21]. The paired-end data from each individual in the pedigree was then aligned to

IMR_hipHip.v1 using BWA-MEM. Resulting bam-files were subjected to duplicate removal

using picard-tools "MarkDuplicates", then raw variant (SNP and indel) calling using GATK

UnifiedGenotyper. Raw SNPs were filtered using GATK "VariantFiltration" with the following

arguments, also with a list of raw indel positions given to the "InDel" mask; "--maskName

InDel", "--maskExtension=5", "--clusterSize=5", "--clusterWindowSize=15",

"BaseQRankSum<-7.0", "BaseQRankSum > 7.0", "MQRankSum < -7.0", "MQRankSum >

7.0", "QD < 5.0","FS > 50.0","ReadPosRankSum > 7.0", "ReadPosRankSum < -7.0" and "DP

> 40". SNPs remaining after filtration were saved in a new vcf file which was given as input to

Lep-MAP3[70] to generate recombination maps for the two families. The commands used in Lep-

MAP were: "ParentCall2", with the options (removeNonInformative=1 XLimit=2 halfSibs=1),

then "Filtering2", with the option data=- (dataTolerance=0.01), then "SeparateChromosomes2"

with the options (lodLimit=10 distortionLod=1) to identify linkage groups in the pedigrees.

SeparateChromosomes2 with lodLimit=10 generated a map of 31 linkage groups, out of which

20 had 964–3063 markers assigned to them (24 were expected based on karyotype). We

continued with the output from SeparateChromosomes2 (lodLimit=10) to run "OrderMarkers2"

to, for each identified linkage group, order the markers without having given the software access

to genome assembly coordinates. The output files from the first iteration of "OrderMarkers2"

were then subjected to a second round of "OrderMarkers2", this time giving the Lep-MAP3

access to genome coordinates of markers on each linkage group using the option

"evaluateOrder". The resulting linkage group specific files containing male- and female

centiMorgan (cM) positions of each marker were filtered to prune obviously inflated cM

positions in the very beginning and the very end of assembled chromosomes and by splitting

four linkage groups where two assembled chromosomes had been merged into one linkage

group, thereby increasing the observed linkage group count from 20 to the expected 24.

Resulting maps from this stage were retained for downstream analyses.

Pacific halibut RNA-seq contrasts

Sequencing data from 12 RNA-seq experiments of various tissues and individuals (SRR11826178-SRR11826189) was downloaded from SRA. Four of these tissues (testis, ovary, white muscle and brain) were used for characterization of *gsdf* gene expression levels and validation of gene models from the gff-file accompanying the Pacific halibut assembly on NCBI. Alignments were performed using STAR as described above for π estimation.

RepeatMasker analysis

RepeatMasker v4.1.0[26] was used to annotate both simple and interspersed repeats. We deployed the RMBLASTN (v2.6.0.+) search engine to scan for known repeats among ray-finned fishes ("-species actinopterygii") using a database that combined Dfam_3.1 and RepBase-20181026 repeat sequences. A custom Perl script was used to parse the main output table from RepeatMasker and compute the densities of major repeat classes (i.e. "Simple_repeat", "LINE", "LTR", "Low_complexity", "DNA", "RC", "SINE", "Satellite", "Unknown", "Retroposon") in windows of 50 kbp and 500 kbp, respectively.

Supplementary data

Supplementary File 1: Integration site analysis of the 1.2 kb insertion upstream of *gsdf*, which controls male development, and related insertions in the Atlantic halibut genome.

References

1   Jakupsstovu, S. H. & Haug, T. Growth, Sexual-Maturation, and Spawning Season of Atlantic Halibut, Hippoglossus-Hippoglossus, in Faroese Waters. *Fish Res* **6**, 201-215 (1988).

2   Tvedt, H. B., Benfey, T. J., Martin-Robichaud, D. J., McGowan, C. & Reith, M.

Gynogenesis and sex determination in Atlantic halibut (Hippoglossus hippoglossus).

Aquaculture **252**, 573-583, doi:10.1016/j.aquaculture.2005.06.042 (2006).

3   Hendry, C. I., Martin-Robichaud, D. J. & Benfey, T. J. Hormonal sex reversal of

Atlantic halibut (Hippoglossus hippoglossus L.). *Aquaculture* **219**, 769-781, doi:Pii

S0044-8486(02)00344-7

Doi 10.1016/S0044-8486(02)00344-7 (2003).

4   Babiak, J. *et al.* Induced sex reversal using an aromatase inhibitor, Fadrozole, in Atlantic

halibut (Hippoglossus hippoglossus L.). *Aquaculture* **324**, 276-280,

doi:10.1016/j.aquaculture.2011.11.001 (2012).

5   Hendry, C. I., Martin-Robichaud, D. J. & Benfey, T. J. Gonadal sex differentiation in

Atlantic halibut. *J Fish Biol* **60**, 1431-1442, doi:10.1006/jfbi.2002.1950 (2002).

6   Kikuchi, K. & Hamaguchi, S. Novel sex-determining genes in fish and sex chromosome

evolution. *Dev Dynam* **242**, 339-353, doi:10.1002/dvdy.23927 (2013).

7   Heule, C., Salzburger, W. & Bohne, A. Genetics of sexual development: an evolutionary

playground for fish. *Genetics* **196**, 579-591, doi:10.1534/genetics.114.161158 (2014).

8   Ross, J. A., Urton, J. R., Boland, J., Shapiro, M. D. & Peichel, C. L. Turnover of sex

chromosomes in the stickleback fishes (gasterosteidae). *Plos Genet* **5**, e1000391,

doi:10.1371/journal.pgen.1000391 (2009).

9   Cnaani, A. *et al.* Genetics of sex determination in tilapiine species. *Sexual development*

: genetics, molecular biology, evolution, endocrinology, embryology, and pathology of

sex determination and differentiation **2**, 43-54, doi:10.1159/000117718 (2008).

10 Franchini, P. *et al.* Long-term experimental hybridisation results in the evolution of a

new sex chromosome in swordtail fish. *Nat Commun* **9**, 5136, doi:10.1038/s41467-018-

07648-2 (2018).

11   Kottler, V. A. *et al.* Independent Origin of XY and ZW Sex Determination Mechanisms

in Mosquitofish Sister Species. *Genetics* **214**, 193-209,

doi:10.1534/genetics.119.302698 (2020).

12   Luckenbach, J. A., Borski, R. J., Daniels, H. V. & Godwin, J. Sex determination in

flatfishes: Mechanisms and environmental influences. *Semin Cell Dev Biol* **20**, 256-263,

doi:10.1016/j.semcdb.2008.12.002 (2009).

13   Drinan, D. P., Loher, T. & Hauser, L. Identification of Genomic Regions Associated

With Sex in Pacific Halibut. *J Hered* **109**, 326-332, doi:10.1093/jhered/esx102 (2018).

14   Herpin, A. & Schartl, M. Plasticity of gene-regulatory networks controlling sex

determination: of masters, slaves, usual suspects, newcomers, and usurpators. *EMBO*

reports **16**, 1260-1274, doi:10.15252/embr.201540667 (2015).

15   Yano, A. *et al.* An immune-related gene evolved into the master sex-determining gene

in rainbow trout, Oncorhynchus mykiss. *Current biology : CB* **22**, 1423-1428,

doi:10.1016/j.cub.2012.05.045 (2012).

16   Bao, L. *et al.* The Y chromosome sequence of the channel catfish suggests novel sex

determination mechanisms in teleost fish. *BMC biology* **17**, 6, doi:10.1186/s12915-019-

0627-7 (2019).

17   Rafati, N. *et al.* Reconstruction of the birth of a male sex chromosome present in

Atlantic herring. *Proc Natl Acad Sci U S A* **117**, 24359-24368,

doi:10.1073/pnas.2009925117 (2020).

18   Rhie, A. *et al.* Towards complete and error-free genome assemblies of all vertebrate

species. *bioRxiv*, doi:10.1101/2020.05.22.110833 (2020).

19   Brown, N. P., Bromage, N. R., Penman, D. J. & Shields, R. J. The karyotype of the

Atlantic halibut, Hippoglossus hippoglossus (Linnaeus). *Aquac Res* **28**, 489-491 (1997).

20 Reid, D. P. *et al.* A genetic linkage map of Atlantic halibut (Hippoglossus hippoglossus

L.). *Genetics* **177**, 1193-1205, doi:10.1534/genetics.107.075374 (2007).

21 Palaiokostas, C. *et al.* Mapping the sex determination locus in the Atlantic halibut

(Hippoglossus hippoglossus) using RAD sequencing. *BMC genomics* **14**, doi:Artn 566

10.1186/1471-2164-14-566 (2013).

22 Myosho, T. *et al.* Tracing the emergence of a novel sex-determining gene in medaka,

Oryzias luzonensis. *Genetics* **191**, 163-170, doi:10.1534/genetics.111.137497 (2012).

23 Rondeau, E. B. *et al.* Genomics of sablefish (Anoplopoma fimbria): expressed genes,

mitochondrial phylogeny, linkage map and identification of a putative sex gene. *BMC*

genomics **14**, 452, doi:10.1186/1471-2164-14-452 (2013).

24 Vinnikov, K. A., Thomson, R. C. & Munroe, T. A. Revised classification of the righteye

flounders (Teleostei: Pleuronectidae) based on multilocus phylogeny with complete

taxon sampling. *Mol Phylogenet Evol* **125**, 147-162, doi:10.1016/j.ympev.2018.03.014

(2018).

25 Kumar, S., Stecher, G., Suleski, M. & Hedges, S. B. TimeTree: A Resource for

Timelines, Timetrees, and Divergence Times. *Mol Biol Evol* **34**, 1812-1819,

doi:10.1093/molbev/msx116 (2017).

26 Tarailo-Graovac, M. & Chen, N. Using RepeatMasker to identify repetitive elements in

genomic sequences. *Curr Protoc Bioinformatics* **Chapter 4**, Unit 4 10,

doi:10.1002/0471250953.bi0410s25 (2009).

27 Sedlazeck, F. J. *et al.* Accurate detection of complex structural variations using single-

molecule sequencing. *Nat Methods* **15**, 461-+, doi:10.1038/s41592-018-0001-7 (2018).

28 Hubley, R. *et al.* The Dfam database of repetitive DNA families. *Nucleic Acids Res* **44**,

D81-D89, doi:10.1093/nar/gkv1272 (2016).

29   Imai, T., Saino, K. & Matsuda, M. Mutation of Gonadal soma-derived factor induces

medaka XY gonads to undergo ovarian development. *Biochem Bioph Res Co* **467**, 109-

114, doi:10.1016/j.bbrc.2015.09.112 (2015).

30   Edvardsen, R. B., Leininger, S., Kleppe, L., Skaftnesmo, K. O. & Wargelius, A.

Targeted mutagenesis in Atlantic salmon (Salmo salar L.) using the CRISPR/Cas9

system induces complete knockout individuals in the F0 generation. *PloS one* **9**,

e108622, doi:10.1371/journal.pone.0108622 (2014).

31   Straume, A. H. *et al.* Indel locations are determined by template polarity in highly

efficient in vivo CRISPR/Cas9-mediated HDR in Atlantic salmon. *Sci Rep* **10**, 409,

doi:10.1038/s41598-019-57295-w (2020).

32   Sawatari, E., Shikina, S., Takeuchi, T. & Yoshizaki, G. A novel transforming growth

factor-beta superfamily member expressed in gonadal somatic cells enhances primordial

germ cell and spermatogonial proliferation in rainbow trout (Oncorhynchus mykiss).

Dev Biol **301**, 266-275, doi:10.1016/j.ydbio.2006.10.001 (2007).

33   Kleppe, L. *et al.* Transcriptomic analysis of dead end knockout testis reveals germ cell

and gonadal somatic factors in Atlantic salmon. *BMC genomics* **21**, 99,

doi:10.1186/s12864-020-6513-4 (2020).

34   Dechaud, C., Volff, J. N., Schartl, M. & Naville, M. Sex and the TEs: transposable

elements in sexual development and function in animals. *Mob DNA* **10**, 42,

doi:10.1186/s13100-019-0185-0 (2019).

35   Faber-Hammond, J. J., Phillips, R. B. & Brown, K. H. Comparative Analysis of the

Shared Sex-Determination Region (SDR) among Salmonid Fishes. *Genome Biol Evol*

7, 1972-1987, doi:10.1093/gbe/evv123 (2015).

36   Herpin, A. *et al.* Transcriptional rewiring of the sex determining dmrt1 gene duplicate

857   by transposable elements. *Plos Genet* **6**, e1000844, doi:10.1371/journal.pgen.1000844

858   (2010).

859   37   Jangam, D., Feschotte, C. & Betran, E. Transposable Element Domestication As an

860   Adaptation   to   Evolutionary   Conflicts.   *Trends   Genet*   **33**,   817-831,

861   doi:10.1016/j.tig.2017.07.011 (2017).

862   38   Bergero, R., Gardner, J., Bader, B., Yong, L. & Charlesworth, D. Exaggerated

863   heterochiasmy in a fish with sex-linked male coloration polymorphisms. *Proc Natl Acad*

864   *Sci U S A* **116**, 6924-6931, doi:10.1073/pnas.1818486116 (2019).

865   39   Sardell, J. M. *et al.* Sex Differences in Recombination in Sticklebacks. *G3-Genes*

866   *Genom Genet* **8**, 1971-1983, doi:10.1534/g3.118.200166 (2018).

867   40   Sutherland, B. J. G., Rico, C., Audet, C. & Bernatchez, L. Sex Chromosome Evolution,

868   Heterochiasmy, and Physiological QTL in the Salmonid Brook Charr Salvelinus

869   fontinalis. *G3 (Bethesda)* **7**, 2749-2762, doi:10.1534/g3.117.040915 (2017).

870   41   Matsuda, M., Sotoyama, S., Hamaguchi, S. & Sakaizumi, M. Male-specific restriction

871   of recombination frequency in the sex chromosomes of the medaka, Oryzias latipes.

872   *Genet Res* **73**, 225-231, doi:Doi 10.1017/S0016672399003754 (1999).

873   42   Lemos, B., Branco, A. T. & Hartl, D. L. Epigenetic effects of polymorphic Y

874   chromosomes modulate chromatin components, immune response, and sexual conflict.

875   *Proc Natl Acad Sci U S A* **107**, 15826-15831, doi:10.1073/pnas.1010383107 (2010).

876   43   Kent, T. V., Uzunovic, J. & Wright, S. I. Coevolution between transposable elements

877   and recombination. *Philos T R Soc B* **372**, doi:ARTN 20160458

878   10.1098/rstb.2016.0458 (2017).

879   44   McGaugh, S. E. *et al.* Recombination Modulates How Selection Affects Linked Sites in

880   Drosophila. *Plos Biol* **10**, doi:ARTN e1001422

10.1371/journal.pbio.1001422 (2012).

45   Kitano, J. *et al.* A role for a neo-sex chromosome in stickleback speciation. *Nature* **461**,

1079-1083, doi:10.1038/nature08441 (2009).

46   Chen, S. L. *et al.* Whole-genome sequence of a flatfish provides insights into ZW sex

chromosome evolution and adaptation to a benthic lifestyle. *Nat Genet* **46**, 253-+,

doi:10.1038/ng.2890 (2014).

47   Bachtrog, D. Y-chromosome evolution: emerging insights into processes of Y-

chromosome degeneration. *Nat Rev Genet* **14**, 113-124, doi:10.1038/nrg3366 (2013).

48   Harboe, T., Mangor-Jensen, A., Naas, K. E. & Naess, T. A tank design for first feeding

of Atlantic halibut, Hippoglossus hippoglossus L., larvae. *Aquac Res* **29**, 919-923,

doi:DOI 10.1046/j.1365-2109.1998.29120919.x (1998).

49   Harboe, T., Tuene, S., Mangorjensen, A., Rabben, H. & Huse, I. Design and Operation

of an Incubator for Yolk-Sac Larvae of Atlantic Halibut. *Prog Fish Cult* **56**, 188-193,

doi:Doi 10.1577/1548-8640(1994)056<0188:Daooai>2.3.Co;2 (1994).

50   Norberg, B., Valkner, V., Huse, J., Karlsen, I. & Grung, G. L. Ovulatory Rhythms and

Egg Viability in the Atlantic Halibut (Hippoglossus-Hippoglossus). *Aquaculture* **97**,

365-371, doi:Doi 10.1016/0044-8486(91)90328-5 (1991).

51   Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of

image analysis. *Nat Methods* **9**, 671-675, doi:10.1038/nmeth.2089 (2012).

52   Li, H. Minimap and miniasm: fast mapping and de novo assembly for noisy long

sequences. *Bioinformatics* **32**, 2103-2110, doi:10.1093/bioinformatics/btw152 (2016).

53   Vaser, R., Sovic, I., Nagarajan, N. & Sikic, M. Fast and accurate de novo genome

assembly from long uncorrected reads. *Genome research* **27**, 737-746,

doi:10.1101/gr.214270.116 (2017).

54   Walker, B. J. *et al.* Pilon: An Integrated Tool for Comprehensive Microbial Variant

Detection and Genome Assembly Improvement. *PloS one* **9**, doi:ARTN e112963

10.1371/journal.pone.0112963 (2014).

55   Simao, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M.

BUSCO: assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics* **31**, 3210-3212, doi:10.1093/bioinformatics/btv351 (2015).

56   Weisenfeld, N. I., Kumar, V., Shah, P., Church, D. M. & Jaffe, D. B. Direct

determination of diploid genome sequences. *Genome research* **27**, 757-767,

doi:10.1101/gr.214874.116 (2017).

57   Dunn, C. W., Howison, M. & Zapata, F. Agalma: an automated phylogenomics

workflow. *BMC bioinformatics* **14**, 330, doi:10.1186/1471-2105-14-330 (2013).

58   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114-2120, doi:10.1093/bioinformatics/btu170

(2014).

59   Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. *arXiv* **1303.3997** (2013).

60   Van der Auwera, G. A. *et al.* From FastQ data to high confidence variant calls: the

Genome Analysis Toolkit best practices pipeline. *Curr Protoc Bioinformatics* **43**, 11 10

11-11 10 33, doi:10.1002/0471250953.bi1110s43 (2013).

61   Langmead, B. & Salzberg, S. L. Fast gapped-read alignment with Bowtie 2. *Nat*

Methods **9**, 357-359, doi:10.1038/nmeth.1923 (2012).

62   Kent, W. J. BLAT - The BLAST-like alignment tool. *Genome research* **12**, 656-664,

doi:10.1101/gr.229202 (2002).

63   Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. *Nucleic Acids Res* **32**, 1792-1797, doi:10.1093/nar/gkh340 (2004).

64   Larsson, A. AliView: a fast and lightweight alignment viewer and editor for large

datasets. *Bioinformatics* **30**, 3276-3278, doi:10.1093/bioinformatics/btu531 (2014).

65   Simpson, J. T. *et al.* Detecting DNA cytosine methylation using nanopore sequencing.

Nat Methods **14**, 407-+, doi:10.1038/nmeth.4184 (2017).

66   Li, X. Y., Chen, F. & Chen, Y. P. Gcluster: a simple-to-use tool for visualizing and

comparing genome contexts for numerous genomes. *Bioinformatics* **36**, 3871-3873,

doi:10.1093/bioinformatics/btaa212 (2020).

67   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,

doi:10.1093/bioinformatics/bts635 (2013).

68   Grabherr, M. G. *et al.* Genome-wide synteny through highly sensitive sequence

alignment: Satsuma. *Bioinformatics* **26**, 1145-1151, doi:10.1093/bioinformatics/btq102

(2010).

69   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing

genomic features. *Bioinformatics* **26**, 841-842, doi:10.1093/bioinformatics/btq033

(2010).

70   Rastas, P. Lep-MAP3: robust linkage mapping even for low-coverage whole genome

sequencing data. *Bioinformatics* **33**, 3726-3732, doi:10.1093/bioinformatics/btx494

(2017).

Table 1. Brx PCR primers

PCR primer	5'-3' sequence
brx_fwd	CAAGCCTGTTCACATCGAAA
brx_malerev	ATGGCAACAAACGCTTATCAA
brx_femalerev	TGGCAACAAACGCTTAACAC

952 **Table 2. Primers used for qPCR**

QPCR primer	5'-3' sequence
gsdf forward	TGCTGGACTCGTCCACATAG
gsdf reverse	GTCCATCATCCCACACCAAT
gtf3c6 forward	CACCGTGAAGAAGCTGATGA
gtf3c6 reverse	ACTTCCTGCTGTGCTGGTTT

Acknowledgements

We would like to thank the National Genomics Infrastructure/SNP&SEQ Technology Platform

in Uppsala for preparing and sequencing the 10X Genomics Chromium Genome library and

thank the Norwegian Sequencing Centre (Oslo) for Illumina sequencing of DNA and RNA.

Computations and data handling were enabled by resources provided by the Swedish National

Infrastructure for Computing (SNIC) at Uppmax, Uppsala, Sweden, with these resources being

partially funded by the Swedish Research Council through grant agreement no. 2018-05973.

We would like to thank Leif Andersson for comments on an earlier version of the manuscript.

We are grateful for financial support from Sterling White Halibut for the sex determining factor

part of this work. This study was co-funded by the Norwegian Ministry of Trade, Industry and

Fisheries. This research was supported by grants from the Swedish Research Councils Formas

(2012-00740 to C.J.R.; 2018-01008 to P.J.; 2017-00413 to A.W) and Vetenskapsrådet 2018-

03017 to P.J.; 2018-04444 to A.W.).

Author contributions

R.B.E., C.J.R and B.N. designed the project. T.H., R.M.J. and M.M. contributed with rearing

of fish. R.M.J., P.P. and B.N. collected the tissue samples. S.K.O. extracted the RNA. L.K.

performed the qPCR. E.K.S. and C.J.R performed the structural variant bioinformatic analysis.

C.J.R. and O.W. did HMW DNA isolation, nanopore sequencing and genome assembly. E.S.

generated RNA-sequencing data. A.W. and C.J.R. performed analysis of RepeatMasker data.

C.J.R performed and P.J. contributed to the TE analysis. C.J.R performed variant calling,

comparative genomics- and divergence analyses. R.B.E. T.F and O.W. performed annotation.

R.B.E. designed, and L.K. and S.M. tested the PCR sexing assay. R.B.E., C.J.R and B.N.

analyzed the results and wrote the paper. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Data Availability

The authors declare that the data supporting the findings of this study are available within the

article and its Supplementary Information. All datasets will be made publicly available under

bioproject PRJNA680301 upon publication.