How ancient forest fragmentation and riparian connectivity generate high levels of genetic diversity in a micro-endemic Malagasy tree

Jordi Salmona[1*], Axel Dresen[1], Anicet E. Ranaivoson[1,2], Sophie Manzi[1], Barbara Le Pors[3], Cynthia Hong-Wa[4], Jacqueline Razanatsoa[5], Nicole V. Andriaholinirina[2], Solofonirina Rasoloharijaona[2], Marie-Elodie Vavitsara[2], Guillaume Besnard[1*]

[1] CNRS-UPS-IRD, UMR5174, Laboratoire Évolution & Diversité Biologique, Université Paul Sabatier, 118 route de Narbonne, 31062 Toulouse, France

[2] Faculté des Sciences, Université de Mahajanga, BP 652 401, Mahajanga, Madagascar

[3] Instituto Gulbenkian de Ciênca, Rua da Quinta Grande, 6, P-2780-156 Oeiras, Portugal

[4] Claude E. Phillips Herbarium, Delaware State University, 1200 N. Dupont Hwy, Dover, DE 19901-2277, USA

[5] Herbier, Département Flore, Parc Botanique et Zoologique de Tsimbazaza, BP 4096, Antananarivo - 101, Madagascar

*Corresponding authors:

Jordi Salmona: jordi.salmona@gmail.com

Guillaume Besnard: guillaume.besnard@univ-tlse3.fr

Number of figures: 4, all figures should be published in color.

Number of tables: 2

The Supporting Information comprises 10 method and 3 result supporting paragraphs, 10 tables, 25 figures and associated references.

Abstract

- Understanding landscape changes is central to predicting evolutionary trajectories and defining conservation practices. While human-driven deforestation is intense throughout Madagascar, exception in areas like the Loky-Manambato region (North) raises questions. This region also harbors a rich and endemic flora, whose evolutionary origin remains poorly understood.

- We assessed the genetic diversity of an endangered micro-endemic Malagasy olive species (*Noronhia spinifolia*) to better understand the vegetation dynamic in the Loky-Manambato region and its influence on past evolutionary processes. We characterized 72 individuals sampled across eight forests through nuclear and mitochondrial restriction associated sequencing data (RADseq) and chloroplast microsatellites (cpSSR).

- Extremely high genetic diversity was revealed in the three genomic compartments (chloroplast $h = 0.99$, mitochondrial $h = 0.85$, and nuclear $H_O = 0.07$-0.20). Combined population and landscape genetics analyses indicate that *N. spinifolia* diversity is best explained by the current forest cover ($R^2 = 0.90$), highlighting a long-standing forest fragmentation in the region. Our results further suggest a predominant role of forest-dwelling organisms in mediating pollen and seed dispersals.

- This sustains a major and long-term role of riparian corridors in maintaining connectivity across those antique mosaic-habitats, calling for the study of organismal interactions that promote gene flow.

Key words: Habitat loss and fragmentation, Landscape genetics, Malagasy olive, Mitochondrial DNA, gene flow, connectivity, cpSSR, RADseq, Madagascar.

Introduction

Offsetting rapid anthropogenic habitat destruction and fragmentation, the primary causes of declines in global biodiversity (Fahrig, 2003; Lindenmayer & Fischer, 2013; Goudie, 2018), requires, among others, to urgently preserving connectivity (Haddad *et al.*, 2015). Although defining appropriate conservation programs largely depends on knowledge of species dispersal strategies (Sutherland *et al.*, 2004; LeBuhn *et al.*, 2015; Gardner *et al.*, 2018), these remain poorly understood, in particular in tropical biodiversity hotspots. This typically requires understanding species diversity, their dynamic, behavior and interactions across rapidly changing landscapes (Pressey *et al.*, 2007), which can be efficiently inferred from genetic data (Frankham, 2010; Salmona *et al.*, 2017a).

Madagascar's unique biodiversity (Goodman & Benstead, 2003; Myers *et al.*, 2000), constitutes an ideal model to study evolutionary processes of diversification (Vences, 2005; Wilmé *et al.*, 2006; Vences *et al.*, 2009). Drivers of evolution, such as riverine barriers (Craul *et al.*, 2008), refugia interconnection (Wilmé *et al.*, 2006), and habitat loss and fragmentation (Yoder *et al.*, 2016; Salmona *et al.*, 2017b), have been identified from taxonomic diversity and the genetic makeup of the Malagasy biota. However, assessing the relative and confounding effects of complex landscape dynamics (forest loss, fragmentation, barriers emergence, etc.) on population dynamics, is notoriously challenging (Nater *et al.*, 2015; Salmona *et al.*, 2017a,b; Beichman *et al.*, 2018).

Deforestation is among the greatest drivers of biodiversity and habitat loss, and fragmentation in Madagascar [~40-50% area since the 1950's (Harper *et al.*, 2007; Vieilledent *et al.*, 2018)]. However, the recent documentation of the Miocene origin of the Malagasy grassland endemics (Bond *et al.*, 2008; Vorontsova *et al.*, 2016; Hackel *et al.*, 2018; Solofondranohatra *et al.*, 2018; Salmona *et al.*, 2020) sparked a hot debate on the antiquity of open-canopy environments (Godfrey & Crowley, 2016; Joseph & Seymour, 2020, 2021). Since the genetic diversity of an organism, and its conservation implications, are the combined results of its distribution structure and history, it is crucial to assess the antiquity of landscapes, which can be questioned from genetic data [e.g. (Quéméré *et al.*, 2010; Yoder *et al.*, 2016; Salmona *et al.*, 2017b, 2020)].

The Loky-Manambato (LM) region in northern Madagascar rose as a small-scale model-
region to assess landscape antiquity and to study habitat loss and fragmentation, thanks to its
perplexingly mild deforestation (Quéméré *et al.*, 2012; Salmona *et al.*, 2017b), its well-
characterized matrix of forests and open-habitats, the diversity of its putative barriers to gene
flow, as well as its high levels of endemicity across living kingdoms (Goodman & Wilmé, 2006;
Goodman *et al.*, 2018). For instance, the forest-matrix was identified as the landscape feature
shaping genetic diversity across all species studied in the LM region, while the Manankolana
River, showed a strong effect on *Propithecus tattersalli*, not consistently recovered in other
species (Quéméré *et al.*, 2010; Rakotoarisoa *et al.*, 2013a; Sgarlata *et al.*, 2018; Aleixo-Pais *et
al.*, 2019; Tang *et al.*, 2020). Although multiple studies on mammals attempted to describe and
understand the processes that shaped its landscape and generated its diversity (Quéméré *et al.*,
2012; Rakotoarisoa *et al.*, 2013b; Salmona *et al.*, 2017b; Sgarlata *et al.*, 2018, 2019),
contributions on other taxa, such as plants, are crucial to draw taxonomically-broad generalities
regarding the antiquity of its landscape, its connectivity and conservation.

Despite their long generation time, native tree species are putatively good models for
landscape genetics studies in fragmented habitats, being the primary and immediate target of
deforestation and landscape changes. However, only a few studies have used the genetic
diversity of Malagasy plant populations (Andrianoelina *et al.*, 2009; Gardiner *et al.*, 2017;
Salmona *et al.*, 2020) to infer landscape dynamics and inform conservation. The Malagasy olives
(genus *Noronhia*), with a high number of taxa and a high micro-endemism rate, are among the
major components of Madagascar forests and of the LM region in particular (Hong-Wa &
Besnard, 2014; Hong-Wa, 2016). Among them, the Malagasy spiny olive (*Noronhia spinifolia*
Hong-Wa) is mostly endemic to the dry to sub-humid forests of the LM region; and although it is
relatively frequent there, it is of high conservation concern due to its narrow range. With such a
distribution, *N. spinifolia*'s genetic diversity holds the potential to have retained information
about the macro- and micro-evolutionary processes that have shaped the genus and species-level
diversity in the region. Furthermore, being narrowly distributed, it may hold relatively low
genetic diversity (Kimura, 1983) and suffer from inbreeding depression due to recent population
collapse. Although its pollen and seed dispersal have yet to be studied, *N. spinifolia*'s flower and
fruit morphology suggests insect pollination and animal-mediated dispersal of fruits (see below).
Noronhia spinifolia therefore represents an excellent model to better understand Malagasy

olives' ecology and offers a case study to define appropriate action for dry-forests plant
conservation in northern Madagascar.

In such sexually-reproducing plants, dispersal occurs by two means: via haploid male
gametes in pollen, and via diploid embryos in seeds. Without field data, population and
landscape genetics offer an alternative way to estimate effective dispersal (Holderegger *et al.*,
2010; Balkenhol *et al.*, 2016). In particular, the combined use of complementary maternally and
biparentally inherited genetic data [respectively from chloroplast or mitochondrial genomes
(cpDNA or mtDNA) and the nuclear genome (nDNA)] allows disentangling, to a certain level,
the relative contribution of seed and pollen dispersals in gene flow. For instance, the congeneric
N. lowryi exhibited contrasting strong chloroplast and near-panmixia nuclear genetic structure
suggesting a long and short distance dispersal of pollen and seed, respectively (Salmona *et al.*,
2020). While progresses in sequencing technologies facilitated the generation of such genetic
data for non-model organisms (Allendorf *et al.*, 2010), recent advances in spatially explicit
analyses also unlocked our ability to estimate the effect of numerous collinear landscape features
on genetic diversity (Balkenhol *et al.*, 2016; Prunier *et al.*, 2017). Furthermore, although the
limited number of tested alternative landscape hypotheses long relied on prior knowledge or
expert opinions, recent approaches iterating around a large panel of resistance values (Graves *et*
al., 2013) or searching for Bayesian optima (Peterman, 2018), widened the potential to
identifying relevant landscape components while optimizing their cost values from the genetic
data itself.

Here, we used genomic data from recently collected specimens of *N. spinifolia* across
most of its range, the LM region. We first tested whether its restricted geographic distribution
resulted in a low genetic diversity, as expected under a neutral model (Kimura, 1983), or
remained relatively high as for co-distributed primates [*P. tattersalli* and *Microcebus tavaratra*
(Quéméré *et al.*, 2010; Aleixo-Pais *et al.*, 2019)]. We then measured the effect of landscape
components on maternally and biparentally inherited genetic diversity, to investigate patterns of
seed and pollen dispersals, and assessed their congruence with those of a congeneric species
from the High Plateau [*N. lowryi* (Salmona *et al.*, 2020)], and of co-distributed mammal taxa
(abovementioned). From the latter, we expect open-canopy habitats and rivers to cause resistance
to *N. spinifolia*'s gene-flow. In contrast, congruence with its congener from the High Plateau
would imply near-panmixia on pollen-dispersed genes, but very short seed dispersal. The little

knowledge about its pollen and seed dispersal agents does not allow making strong predictions,
except that dispersal will depend on the vectors and on their use of the landscape. We also
examined whether the relative stability of the forest cover in the past 70 years (Quéméré *et al.*,
2012; Salmona *et al.*, 2017b) is reflected in *N. spinifolia* genetic makeup, comparing the effect of
recent and historical forest covers on gene flow, as a proxy for the temporality of its habitat
fragmentation. Finally, we present the application of our work to the conservation of the LM
region forest network.

Material and methods

Study region

The Loky-Manambato (LM) region (Daraina; Fig. 1) is a biogeographical transition zone
between dry deciduous and humid forests (Goodman & Wilmé, 2006), which is delimited by the
Loky and Manambato Rivers. This region is crossed by the relatively shallow Manankolana
River, bordered by riparian forests along most of its course, and by a national dirt road (Fig. 1). It
consists of an area of ~2,500 km² covered by ~360 km² of forests (Goodman *et al.*, 2018),
fragmented into a dozen major forest patches surrounded by human-altered grasslands, dry scrub
and agricultural lands. Most forests are situated at low- to mid-elevations and mostly consist of
dry deciduous vegetation. In contrast, some mountain forests (Binara and Antsahabe, plus
Bobankora to a lower extent) are covered by a gradient of dry deciduous, transition, humid and
ericoid vegetation (Gautier *et al.*, 2006). Despite sustained grassland fires, slash-and-burn
agriculture and charcoal production, as well as exploitation of wood, gold and sapphires
(Fanamby, 2010; Goodman *et al.*, 2018), deforestation rate in the LM region is still relatively
low (Quéméré *et al.*, 2012) compared with those of eastern and southwestern Madagascar
(Vieilledent *et al.*, 2018), likely stemming from its remoteness, difficult accessibility and
climate. However, to mitigate the threats, the LM region progressively became managed as a
protected area by the Malagasy NGO "Fanamby" since 2005 (Fanamby, 2010; Goodman *et al.*,
2018).

Study species

Noronhia spinifolia (Oleaceae) is a small-sized, understory tree that is easily distinguishable from other *Noronhia* species by its narrow linear leaves with a spiny tip. The plant has cream-white, urceolate, small (> 7 mm long), and hermaphroditic flowers, as well as small (> 10 mm long) and drupaceous fruits that have a thin mesocarp and a rather crustaceous endocarp (Hong-Wa, 2016). Flowering and fruiting typically occur from October to May, during the rainy season. Flower and fruit characteristics, along with observational accounts, suggest insect pollination (e.g. bees) and animal dispersal (e.g. birds, lemurs, rodents). This *Noronhia* species is micro-endemic to northern Madagascar, mainly found in the LM region except for one record from further north in Montagne des Français, and is reported mainly in semi-deciduous forests of low altitude, mostly on alkaline substrate (e.g. limestone, calc-alkaline rocks). *Noronhia spinifolia* has been assigned a preliminary conservation status of "Endangered" due to threats to its habitat (Hong-Wa, 2016).

Figure 1: Map of *Noronhia spinifolia* sampling in the Loky-Manambato (LM) region.
The small black points represent samples collected for all *Noronhia* species (ca. 30 distinct taxa) and illustrate the survey effort conducted in the region. The yellow and red dots represent *N. spinifolia* samples, with the red dots corresponding to samples included in our genomic analyses. The forest cover is adapted from Hansen *et al.* (2013). Pixels with less than 30% tree cover are represented in white. The remaining tree cover percentage values are represented from light green (30%) to dark green (100%). This forest cover representation also illustrates the presence of riparian forests along streams of the LM region.

Plant sampling

To sample *N. spinifolia* populations, we surveyed all major forests of the LM region (Fig. 1) in 2017 and 2018, during the dry season (July-September), and used topography (altitude and shape) as a sampling guide to maximize the representation of all landscape features. Most surveys started from the forest edge at low altitude towards the forest core at higher elevation. We identified *Noronhia* species based on tree characteristics, leaf morphology and tissue structure, and collected leaf samples of 220 *N. spinifolia* trees, preserved in silica gel for DNA conservation. We prioritized fully-grown mature tree sampling because much of the density-dependent mortality takes place before maturity in trees, and their effective population size contributing to the genetic diversity is thus closer to the actual adult census size than to the size of the entire population including young trees and seedlings (Dodd *et al.*, 1999; Petit & Hampe, 2006). Therefore, the regional patterns of diversity are expected to be better represented by adult samples. For each tree, we systematically recorded its height, diameter and reproductive state, as well as its geographical coordinates (GPS) and elevation. For all forests, at least one specimen voucher was prepared and deposited at the herbarium of the Parc Botanique et Zoologique de Tsimbazaza (TAN).

Laboratory procedures

DNA extraction, organellar and nuclear genotyping

We extracted DNA from 137 samples of *N. spinifolia* using a commercial protocol adapted to plants, followed by quality control procedures ensuring high quality genomic DNA. We subsequently genotyped 72 high DNA quality samples (Fig. 1, Methods S1); a cost-effective subsampling that nonetheless maximizes geographic and altitudinal representation, and also prioritizes reproductively mature and fully-grown trees with a targeted sequencing depth >15×. Using a two-pronged approach, we genotyped 15 chloroplast microsatellites (cpSSR) and one mitochondrial microsatellite (mtSSR), originally developed on *Olea europaea* (Table S1, Methods S2, S3; Besnard *et al.*, 2011), and also used restriction associated DNA sequencing (RADseq; generating data from the biparentally inherited nuclear genome and the mitogenome; Methods S4). RADseq consists in sequencing regions neighboring restriction sites, to obtain

220   homologous sequences across individuals, spread across the genome, at a decent coverage and a

221   reasonable cost (Baird *et al.*, 2008; Andrews *et al.*, 2016).

Data processing

Organellar RADseq loci and *de-novo* assembly of the nuclear loci catalog

After ad-hoc demultiplexing and cleaning of reads (Methods S4), we screened the organellar

genomes using bwa-mem sequence alignment (Li, 2013) to the *N. clarinerva* mitogenome and *N.*

spinifolia plastome (MW202230 and MT081057, respectively; Methods S5). We identified ten

mitochondrial *Sbf*I RAD loci *in silico*, from which haplotypes were called using ANGSD v0.92

(Nielsen *et al.*, 2012; Korneliussen *et al.*, 2014), based on their highest effective base depth

(Wang *et al.*, 2013). Conversely, no cpDNA RAD locus was recovered, confirming *in silico*

analyses (Methods S5).

A catalog of nuclear tags (loci) was *de-novo* optimized (Methods S6) by iterating around

the core parameters of Stacks (Rochette *et al.*, 2019) to maximize the amount of available

biological information (Paris *et al.*, 2017). The final catalog was further cleaned (Methods S6)

for exogenous contaminants using DeconSeq (Schmieder & Edwards, 2011) and endogenous

orthologs using MUMmer (Kurtz *et al.*, 2004).

RADseq genotyping

We used two fundamentally distinct genotyping approaches to ensure the robustness of our

results: single nucleotide polymorphism (SNPs) called in Stacks, and genotype likelihoods (GLs)

estimated with ANGSD (Methods S7). GLs retain information about uncertainty in base calls,

which alleviates some issues associated with RADseq data such as unevenness in sequencing

depth and allele drop-outs (Pedersen *et al.*, 2018; Warmuth & Ellegren, 2019; Heller *et al.*,

2021).

Landscape genetics

We conducted complementary analyses to assess the effect of landscape components on the

genetic diversity of *N. spinifolia*. We first investigated the raw patterns of genetic diversity and

structure without priors to describe the major trends and build hypotheses. Then, using univariate

approaches under an isolation-by-resistance model (IBR; McRae, 2006), we assessed the effect

of each landscape component, iterating through their cost and resolution. Finally, using a

multivariate model considering spatial autocorrelation and multicollinearity, we assessed the

contribution of selected landscape components.

Genetic diversity

252 We assessed the proportion of heterozygous genotypes (H_E) from nuclear genotype likelihoods

253 (GL) based on folded site frequency spectra estimated in ANGSD. We further estimated

254 organellar diversity (h), the probability that two haplotypes are different (Nei, 1987).

Population structure

We assessed the level of genetic differentiation among localities with Reynolds' weighted F_{ST}

(Reynolds *et al.*, 1983) from GL inferred in ANGSD. We explored the genetic structure of our

study system through naive clustering analyses (Methods S8), based on ANGSD GLs using

NgsAdmix v32 (Skotte *et al.*, 2013) and on Stacks called genotypes using ADMIXTURE v1.3.0

(Alexander *et al.*, 2009), and with a principal component analysis (PCA) from GLs with

PCAngsd. We estimated the level of organellar genetic differentiation among forests with Nei's

weighted F_{ST} (Nei, 1973) using the R package *hierfstat*. We also investigated the phylogenetic

structure of organellar DNA data using minimum spanning networks of genetic distances (see

below) constructed with the R package *poppr* (Kamvar *et al.*, 2015).

Genetic distances

We assessed the power of several individual pairwise estimates of genetic relationships

(distances or relatedness) from chloroplast, mitochondrial and nuclear data. For cpSSR data, we

used the Bruvo's and Prevosti's genetic distances (Prevosti *et al.*, 1975; Bruvo *et al.*, 2004).

From mtRAD SNPs, we inferred Euclidian and Manhattan distances. We estimated an overall

genetic distance for organellar genomes by combining weighted Manhattan mtDNA and Bruvo's

cpDNA distances (Methods S3).

We estimated the covariance of nuclear RADseq GLs (Meisner & Albrechtsen, 2018), as

well as Hall's and Vieira's metrics (Hall *et al.*, 2012; Vieira *et al.*, 2013) in PCAnsgd. Using

nuclear SNP data, we also computed Nei's genetic distance (Nei, 1972) and Yang's relatedness
(Yang *et al.*, 2010) in the *StAMPP* R package (Pembleton *et al.*, 2013).

Isolation by distance

We investigated patterns of isolation by distance (IBD) to assess how the geographic distance
alone explains the genetic diversity (Wright, 1943; Slatkin, 1993). We used Mantel tests (Mantel,
1967) between individual geographic and genetic distances (Methods S9). Since IBD may be
limited to a certain scale (e.g. Keller & Holderegger, 2013; Van Strien *et al.*, 2015; Cayuela *et*
al., 2019), we compared subsets of pairwise data defined by a maximum geographic distance (S)
between samples (Methods S9).

Isolation by resistance

284   Landscapes are rarely homogeneous, and gene flow may be limited or facilitated by its
285   components. We used an IBR approach (McRae, 2006) to assess the cost associated with
286   effective dispersal through each landscape feature.

Landscape variables, cost and resolution

As *N. spinifolia* was recently described and occurs in a remote area (Hong-Wa, 2016), we had
little prior knowledge on the landscape variables that may affect pollen and seed dispersal. We
therefore assessed the effect of most available landscape variables (Table 1; Methods S10). To
test if the genetic diversity of old trees may be better explained by past forest cover, we used
forest cover data from 1953, 1973, and 2000s (Hansen *et al.*, 2013; Vieilledent *et al.*, 2018).

Although strong priors associating a landscape component to a particular cost may be
available for well-studied species (e.g. Dellicour *et al.*, 2019; Quéméré *et al.*, 2010), landscape
variables and their associated cost are often chosen almost arbitrarily when little or no data are
available (Beier *et al.*, 2008, 2011). To identify the variable-cost associations that matter for our
study system, we iteratively tested 14 conductance-resistance values (Methods S10). Similarly,
organisms do not necessarily perceive each environmental component at the same resolution (or
granularity: Baguette & Van Dyck, 2007; Everson & Boucher, 1998; Laurance *et al.*, 2007;
Murcia, 1995). To identify the variable-cost-granularity relevant for *N. spinifolia*, we tested four
pixel resolutions (Methods S10).

Table 1: Landscape variables.

Variable	Abbreviation	Type	Univariate effect	Unique contribution
Geographic distance	IBD	Continuous	RES	NS*
Rivers	Rivers	Discrete	NA	NS*
Streams	Streams	Discrete	NA	NS
Roads	Roads	Discrete	NA	NS
Trails	Trails	Discrete	NA	NS
Slope	Slope	Continuous	NA	NS*
Wind	Wind_November	Continuous	NA	NS
% tree cover	%_tree_cov	Continuous	CON	CON*
% tree cover discrete	%_tree_cov_dis	Discrete	CON	CON
Forest cover ~2000	Veg_2000	Continuous	CON	CON
Forest cover ~1973	Veg_1973	Continuous	CON	CON
Forest cover ~1953	Veg_1953	Continuous	NA	NS

303
RES = variable exhibiting resistance; CON = variable exhibiting conductance; NA = no major
effect detected; NS = non-significant unique contribution; * variable included in the final model
presented in the main manuscript.

Movement models

To determine which dispersal model best applies to *N. spinifolia*, we used both the Least Cost
Path (LCP) and the Circuit Theory (CT). These two approaches, respectively, consider the least
cost trajectory and the cost of all possible trajectories (McRae & Beier, 2007). We computed
landscape distances using the R package *gdistance* (Van Etten, 2012).

Statistical procedures

We used a two-step procedure to first select landscape components, as well as their best fitting
cost, resolution, and movement model, and then, to assess their unique and common
contributions to the spatial structure of *N. spinifolia*'s genetic diversity.

We estimated the correlation between geographic or landscape distance and genetic
matrices (i.e. Landscape variables and Genetic distances as described above) using Mantel tests
(Mantel, 1967) in the R Package *vegan* (Dixon, 2003). We retained variables showing a better fit
(R^2) than IBD, exhibiting sensitivity to cost values (i.e. variables with a fixed fit across all cost
values were discarded), and selected their best fitting cost, movement model, and resolution. We
modeled the contribution of the retained landscape variables using logistic regressions on
distance matrices [LRDM] (Smouse *et al.*, 1986; Prunier *et al.*, 2015), a statistical procedure that

is similar to classical multiple ordinary least-square regressions, except that the significance of

model fit (multiple R^2) is assessed through permutations of the dependent matrix (Legendre *et*

al., 1994). We finally disentangled multicollinearity among variables and decomposed their

unique and common contributions using commonality analyses (CA; Prunier *et al.*, 2015).

Results

Species occurrence

We sampled *N. spinifolia* in eight of the 11 surveyed major forests of the LM region (Fig. 1). The species occurs from low to medium elevation, between 87 and 505 m, but with strong discrepancies among forests (Fig. S1). While it was mainly recorded in dry forests, it was surprisingly found in dry to wet transition forests at medium elevation (451-505 m) in Binara. Furthermore, the species was not found in three major forest patches of the LM region - namely Antsahabe, Ambohitsitondroina and Antsaharaingy - despite (*i*) large prospection efforts in these forests, and (*ii*) apparently similar habitat as the neighboring forests harboring the species (Fig. 1).

Organellar DNA genotyping and nuclear catalog construction

Of the 15 chloroplast microsatellites, 14 showed polymorphism (Table S2), and allowed distinguishing 55 chlorotype profiles among 72 trees (Results S1). The ten mitochondrial RAD loci (mtRAD) allowed identifying 11 SNPs (Results S1; Table S3). The combination of mtRADs and the mtSSR locus permits the identification of 15 mitotypes among 72 trees (Table 2). The cpSSR markers showed low to moderate linkage disequilibrium (LD; Fig. S2), a likely consequence of microsatellite-repeat-length homoplasy. Meanwhile, the mtDNA markers showed either high (among seven loci) or no LD (Fig. S3). Because SNPs are expected to be more stable (unlikely homoplasy) than SSRs, no LD between SNP loci was not expected, and could indicate recombination in the mitogenome. Finally, the overall LD among mtDNA and cpDNA markers (Fig. S4) suggests that they are both maternally inherited, although paternal leaks may occur occasionally.

The nuclear catalog parameter space exploration iterating around the core parameters for Stacks [i.e. m – the minimum number of reads required to build a stack, M – the maximum number of differences between stacks of an individual allowed when building a locus; and N – the maximum number of differences between loci of multiple individuals allowed when building a loci] allowed selecting values ($m = 4$, $M = 5$, $N = 8$) that offer a trade-off between the coverage,

loci number, and SNP number, while limiting the number of paralogs and the presence of contaminants (Figs S5-7; Results S2). The SNP-calling procedure showing low ability to recover the genetic makeup of *N. spinifolia* (when compared to the GL-based procedure; Figs S8-13), we therefore limited its use to preliminary analyses (ADMIXTURE & genetic distances) and proceeded with the GL-based procedure for downstream analyses.

Genetic diversity

Chloroplast microsatellites revealed a relatively high genetic diversity with only two chlorotypes shared by individuals from more than one forest, resulting in a high probability that two randomly sampled haplotypes are different ($h = 0.99$) and a mean allelic richness (A_r; estimated for five individuals) of 2.41 (Table 2). Consequently, most forests showed an extremely high cpSSR genetic diversity ($h > 0.92$) with the exception of Binara that appeared slightly less diverse ($h = 0.73$; Table 2). A relatively high mitotype diversity was also revealed [$h = 0.85$ (ranging from 0.66 to 0.97 per forest), $A_r = 2.12$]. Similarly, most sampled individuals exhibit relatively high levels of nuclear diversity with ~7 to ~20% of polymorphic sites and large discrepancies within and among forests (Table S1; Fig. S14). This diversity is not homogeneously distributed in space, and higher levels of genetic diversity seem to occur in the area from Solaniampilana to southern Bekaraoka (Fig. S15). Furthermore, genetic diversity does not seem influenced by altitude (Fig. S16).

Table 2: Chloroplast and mitochondrial summary statistics.

Forests	cpSSR				mtRAD			
	N	n_h	h	A_r	N	n_h	h	A_r
Ambilondamba	6	5	0.98	2.22	6	4	0.97	2.16
Ampondrabe	1	1	-	-	1	1	-	-
Antsiasia	6	4	0.92	2.67	6	3	0.81	2.14
Bekaraoka	25	19	0.99	2.38	22	5	0.66	2.04
Benanofy	11	8	0.94	2.39	11	4	0.78	2.26
Binara	5	2	0.73	2.36	5	2	0.73	2.05
Bobankora	11	10	0.99	2.45	11	3	0.73	2.04
Solaniampilana	10	8	0.97	2.37	10	5	0.87	2.17
Total / Mean	75	55	0.99	2.41	72	15	0.85	2.12

N = number of analyzed individuals; n_h = number of haplotypes; h = haplotype diversity; A_r: allelic richness (estimated for five individuals).

Population structure

The chloroplast and mitochondrial data both revealed substantial differentiation among forests (F_{ST} estimates ranging from 0.040 to 0.393 for cpSSRs; and 0.005 to 0.661 for mtRADs). As expected, a strong differentiation was also observed when combining cpDNA and mtDNA data (F_{ST} estimates ranging from 0.101 to 0.401; Table S4). The Solaniampilana-Benanofy forest cluster was clearly distinguished from other forests for both mtDNA and cpDNA (Figs S17-18), while Bekaraoka and Bobankora showed limited divergence with their neighboring forests. Haplotype networks based on cpSSR and/or mtRAD data also revealed that one maternal lineage is unique to Solaniampilana and Benanofy (Fig. 2). Furthermore, the geographic Euclidean distances showed low, but highly significant, power at explaining genetic distances among individuals (R^2 [cpSSR]: 11.7%; R^2 [mtRAD]: 20.7%; and R^2 [cpSSR + mtRAD]: 21.3%; Figs S13, S19; Results S3).

Figure 2: Organellar DNA haplotype network of *Noronhia spinifolia*.

Line length and grey scale are proportional to the Bruvo's cpDNA + Manhattan mtDNA
combined genetic distances between distinct organellar haplotypes. Pie chart size is proportional
to the occurrence number of a given haplotype. All edges of equal weight are represented.
Distances among haplotypes are represented both through longer edges and the grey scale. The
network highlights the huge organellar DNA diversity in *N. spinifolia*, with only one haplotype
shared by individuals from at least two forests. It further shows a limited spatial structure, with,
for instance, haplotypes from Solaniampilana and Benanofy grouping together at the bottom of
the network.

F_{ST} estimates based on nuclear markers (Table S5) ranged from 0.089 to 0.210, indicating

that most forests are differentiated from each other. However, we found no strong structure in

sub-populations, with no particular support for number of clusters >1, both for GL- and SNP-

based analyses (Figs S8-9). Instead, we found a clear northwest-southeast signal of continuous genetic differentiation across space, through GL-based PCA (First axis, ~15% of the variance explained; Fig. S20), clustering (Figs 3, S10-11), and IBD analyses (Figs S13, S19). The observed continuous structure is well illustrated by the clustering structure for $K = 3$ that shows admixed patterns at sampling sites (Fig. 3). We found a clear IBD signal explaining up to 56.6% of the among-individuals nuclear GL covariance (Fig. S19).

Figure 3: Spatial genetic structure of *Noronhia spinifolia* in the Loky-Manambato region. NgsAdmix ancestry proportions (for $K = 3$ genetic clusters) represented either (**A**) spatially by sampling site, or (**B**) per individual. Size of pie charts (in A) is proportional to the number of samples per site. Pie shares represent the sums of individual ancestry proportions that are shown in B. Results are arbitrarily represented for $K = 3$, according to the likelihood and delta$K$ results in Fig. S8, because this K value best illustrates the continuous pattern of structure inferred using ngsAdmix and other approaches.

Landscape genetics

The optimization of resistance surfaces through univariate comparison of genetic and landscape distances (IBR) showed lower fit for cpDNA (R^2 max ~0.14) than for mtDNA (R^2 max ~0.38) and nDNA (R^2 max ~0.90). Among the four vegetation layers, the continuous and discrete percent tree cover layer always exhibited the highest fit for conductance values at high resolution with cpDNA, mtDNA and nDNA (R^2 = 0.14; 0.38 and 0.90, respectively; Figs S21- 24). In other words, the percent tree cover data alone shows a strong conducting effect on gene flow and explains a very large portion of the genetic variation (R^2 = 0.90). Altogether the parameter space exploration reveals a strong effect of all forest cover layers, whereas some other variables (i.e., rivers, roads and slope) may have subtle lower effects too. To build multivariate models, we retained in priority landscape variables showing a better fit (R^2) than the null model considering IBD alone, and exhibiting sensitivity to cost values (e.g. % forest cover). Our results combining LRDM and CA confirmed that forest cover was the best landscape predictor of genetic differentiation, releasing other landscape components and IBD to account mostly for collinearity with the forest cover (Fig. 4; Table S6). This pattern was consistent across organellar and nuclear DNA (Table S6), and the high quality percent tree cover from Hansen *et al.* (2013) was always the best forest cover predictor (Table S6). The 2000's forest covers all better fit genetic diversity than the 1953 and 1973 forest covers, meaning we did not recover particular effect of the documented forest-cover changes on the genetic diversity of *N. spinifolia*.

Figure 4: Landscape contribution to nuclear gene flow in *Noronhia spinifolia*.

A) Unique and common contributions of four selected landscape variables to nuclear gene flow, estimated using commonality analysis. **B**) Geographic representation of the percent tree cover conductance (inverse of cost), which illustrates the landscape conductance. **C**) Graphic representation of the relationship between percent tree cover conductance and genetic distances (isolation by resistance). This figure illustrates a strong conducting effect of forest cover (percent tree cover) on the connectivity of *N. spinifolia*, and it further shows that Euclidean geographic distance (IBD), the Manankolana River (Rivers) and the topology (Slope) have very low unique contribution, if any, to *N. spinifolia* nuclear gene flow. Cover_continue: Percent tree cover, conductance = 5; IBD: Isolation by distance, resistance = 1; Rivers: resistance = 5; Slope: conductance = 5.

Discussion

From a comprehensive and extensive sampling of *Noronhia spinifolia* in its core distribution area, and leveraging the rare combination of nuclear and mitochondrial RADseq data with cpDNA microsatellites, this study allowed us to reveal a strong effect of forest cover on gene flow in a fragmented habitat in northern Madagascar. We not only report a surprisingly high organellar and nuclear genetic diversity unevenly distributed in space, but also found that GL-based approaches were able to recover much more information than SNP-calling approaches in our model species. Moreover, the iterative optimization of resistance surface allowed identifying outstanding landscape variables with a strong effect on the connectivity of *N. spinifolia*. Finally, we show that recent forest cover better explains the genetic structure of *N. spinifolia* than more ancient ones.

Noronhia spinifolia, a highly diverse Malagasy micro-endemic

Our analyses exhibit unexpectedly high chloroplast ($h = 0.99$; 55 chlorotypes for 72 individuals), mitochondrial ($h = 0.85$; 15 mitotypes), and nuclear (~7-20% polymorphic sites) genetic diversity in a micro-endemic Malagasy tree species.

Firstly, the cpDNA diversity is tremendously higher than that of another micro-endemic congener of the High Plateau (*N. lowryi*) when using the same 15 cpSSR loci [6 haplotypes in 77 individuals; $h = 0.58$ (Salmona *et al.*, 2020)]. More surprisingly, more cpDNA haplotypes and diversity were revealed in 72 *N. spinifolia* individuals than in 1263 wild olive trees from the whole Mediterranean basin [47 chlorotypes; $h_{cp} = 0.35$ (Besnard *et al.*, 2013)] and thus across very different geographic scales (LM region = 900 km² *vs* Mediterranean basin = ~2.5 Million km²) and despite the use of more polymorphic cpSSRs (n = 35) in olive. Similarly, the *N. spinifolia* mtDNA diversity is also higher than in the Mediterranean olive [4 mitotypes; $h_{mt} = 0.58$; (Besnard *et al.*, 2002)], although comparable diversity levels have been revealed in other plant groups exhibiting large mitogenomes with high mutation rates as *Silene vulgaris* in Central Europe [30 mitotypes; $h = 0.94$; (Štorchová & Olson, 2004)]. Finally, the nuclear genomic diversity is ~20-40 times higher than that estimated in poplar populations across all Eurasia (Ma *et al.*, 2018). For the sake of an approach-based comparison, the diversity is also twice as large

as in five eastern-Madagascar mouse lemurs and two orders of magnitude higher than in African
plains zebra both estimated using RADseq data and a GL-based analytical procedure (Pedersen
et al., 2018; Poelstra *et al.*, 2021). This high genetic diversity is particularly unexpected for a
narrowly distributed micro-endemic, and thus threatened, species.

Although high standing genetic diversity is common in forest trees, the relative
importance of the multiple mechanisms generating and maintaining this diversity are still
debated (Petit & Hampe, 2006; Scotti *et al.*, 2016; Isabel *et al.*, 2020). In *N. spinifolia*, several
non-exclusive evolutionary mechanisms may explain such an exceptionally high intraspecific
genetic diversity. Firstly, it suggests that a long-term maintenance of a large effective population
size precluded significant genetic drift. Persistent connectivity between forest patches may have
been key in this process, particularly during climatic fluctuations of the Late Quaternary that
may have contributed to fragmenting habitat, as suggested for other species of the LM region
(Quéméré *et al.*, 2012; Salmona *et al.*, 2017b). Secondly, the genus *Noronhia* has extremely
diversified in northern Madagascar (Hong-Wa, 2016), and about 30 taxa have been recently
recorded and sampled in the LM region (JS & GB, unpublished data). What caused such
diversification remains unknown. The co-occurrence of closely related taxa may offer some
opportunities for hybridization events, which could have contributed to the increased genetic
diversity in *N. spinifolia*. However, the cpSSR characterization of four sympatric/parapatric LM
Noronhia (i.e. *N. candicans*, *N. clarinerva*, *N. crassinodis* and *N. intermedia*; > 200 individuals),
closely related to *N. spinifolia* (according to cpDNA and nrDNA data; Salmona *et al.*, 2020),
shows that these species have no shared chlorotype with our study model (GB, unpubl. data),
thus suggesting that maternal introgression events to *N. spinifolia*, if any, may not be recent.
Lastly, high mutation rate may also contribute to the high genetic diversity in *N. spinifolia*. An
obvious acceleration of the mitogenome evolutionary rate has been recently documented in the
closely related species *N. candicans*, *N. clarinerva*, *N. intermedia* and *N. spinifolia,* with a high
number of di- or tri-nucleotide mutations possibly reflecting frequent mtDNA recombination in
this clade (Van de Paer, 2017), as also suggested by a lack of LD between some SNPs. While
accelerated mutation rate was missing on the plastome (Salmona *et al.*, 2020), we are still
lacking any evidence for the nuclear genome. Such accelerated evolutionary rate could result
from relatively frequent and recurrent hybridization events in this group, promoting genomic
instability (Fontdevila, 2005; Payseur & Rieseberg, 2016). Moreover, the strong linear

relationship between geographic and genetic distance could preclude cryptic radiation (Pillon *et*
al., 2014) and microgeographic adaptation (Scotti *et al.*, 2016) as major drivers of the observed
diversity. In conclusion, the surprisingly high genetic diversity calls for the identification of the
evolutionary, ecological and/or molecular mechanisms underlying this peculiar pattern.

513 **Landscape effects on the genetic diversity of *Noronhia spinifolia***

514 A strong continuous spatial structure

Beyond revealing surprisingly high levels of diversity, our results also show complementary
signals of a strong continuous structure in space (PCA, clustering and IBD), from both organelles
and the nucleus, in contrast to generally expected incongruent patterns among genomes
(Olofsson *et al.*, 2019; Bianconi *et al.*, 2020). While the northwest-southeast differentiation cline
represented as much as ~15% of the variance of the PCA, the geographic Euclidean distance
alone explained up to ~55% of the nuclear genetic variance using IBD tests. This strong pattern
of nuclear genetic structure sharply contrasts with the absence of nuclear spatial structure in the
savanna olive tree, *N. lowryi* (Salmona *et al.*, 2020). However, reported IBD patterns in trees
show a wide range from low values in *Dalbergia monticola* across eastern Madagascar humid
forests [R^2 = 0.18; (Andrianoelina *et al.*, 2009)], or *Coffea mauritiana* in the Reunion Island [R^2
= 0.21; (Garot *et al.*, 2019)], to high values in *Swietenia macrophylla* in Central America [R^2 =
0.62; (McRae & Beier, 2007)]. Unexpectedly, this genetic structure was here extremely well
explained by the vegetation cover (percent tree cover; mtDNA R^2 = 0.38; nDNA R^2 = 0.90),
releasing IBD to account mostly for collinearity with the forest cover. Although strong landscape
effects were also found in *S. macrophylla* (McRae & Beier, 2007), we report a unique evidence
of a strong habitat fragmentation effect explained mostly by one landscape variable.

531 On seed-mediated gene flow: the organellar DNA testimony

532 Although organellar IBR patterns (Figs S19, S21-24) suggest that seed-mediated gene flow is
533 driven by forest cover, the recovered pattern was of lower intensity than for pollen-mediated
534 gene flow (nDNA). Despite slope and watershed networks being candidates for barochory and
535 hydrochory, we could not recover any landscape variable (other than forest cover) with
536 noticeable effect on seed dispersal. Similarly, the overall structures of organellar haplotype

networks (Figs 2, S17-18) are coherent with the geographic repartition of forests, and in line with the effect of the forest cover. These prevailing effects of forest cover suggest that seed dispersal may be primarily performed by forest-dwelling animals (zoochory), especially those with limited and/or rare across-forest movements, such as lemurs, rodents and territorial birds (Quéméré *et al.*, 2010; Rakotoarisoa *et al.*, 2013a; Sgarlata *et al.*, 2018; Aleixo-Pais *et al.*, 2019). However, the networks also show multiple potential fluxes among forests, hence supporting the network complementarity to the IBR approach. Several non-exclusive interpretations can be invoked for explaining these patterns: (*i*) relevant landscape variables are not included or of low resolution (e.g. forest type and climatic variables); (*ii*) the cpDNA and mtDNA diversities are confounded by homoplasy, recombination, strong drift, long-term phylogenetic or demographic history; and (*iii*) seed dispersal also result from infrequent seed ingestion by wide-ranging birds (or other vertebrates).

A deep forest cover effect on gene flow

Unlike organellar DNAs, nDNA diversity is deeply explained by the LM region forest cover (Fig. 4). While this partially confirms the effect of forest cover on seed dispersal since nDNA diversity is influenced by both seed and pollen movement, wind-mediated pollen dispersal favored in open-canopy environments is not supported here. It thus further sustains that pollen dispersal is mediated by forest-dwelling organisms with movements limited by open-canopy environments. Insect-mediated pollen dispersal in *N. spinifolia* is also strongly suggested by its flower morphology and color (Hong-Wa, 2016). However, the currently limited knowledge of the Malagasy entomofauna and plant-pollinator networks prevents us from clearly identifying this species' forest-dwelling pollinators.

The antiquity of forest fragmentation in northern Madagascar

Our results further support a long-standing forest fragmentation in the LM region. First, the better fit of all recent forest cover (2000's), compared to older vegetation cover (1953, 1973), suggests that the small forest changes that have occurred through this period (Quéméré *et al.*, 2012) are unable to explain the genetic diversity of *N. spinifolia*. These mild landscape changes in the LM region contrast with the high deforestation rates observed throughout Madagascar since the fifties (Hansen *et al.*, 2013; Vieilledent *et al.*, 2018). Under such high recent

deforestation rates, a better fit of the recent forest cover layer would be very unlikely, even
considering that its better resolution could positively bias its fit. Second, because we mostly
genotyped fully-grown mature trees, and since the generation time of *Noronhia* is potentially
long [>20-50 years; (Salmona *et al.*, 2020)], the genetic diversity is expected to reflect ancient
forest cover. The time lag for a particular landscape feature to imprint its effects in the genetic
diversity of a species, has been little studied (Landguth *et al.*, 2010; Mona *et al.*, 2014).
However, in *N. spinifolia*, based on the strength of the signal, the high level of diversity and of
gene-flow, the re-shuffling of allele frequencies after fragmentation can be roughly expected to
last at least 40 generations, before harboring the signature of the new geographical pattern. This
suggests that the landscape changes leading to the current forest cover are at least ~800 years (40
generations x 20 years), i.e. long pre-dating the most ancient available layer (1953). The strong
genetic correlation with the recent forest cover is, therefore, sound evidence that the landscape of
the LM region was relatively stable at least for the last century (i.e. when most of Madagascar's
deforestation occurred), and possibly the last millennium. This result concurs with those of
recent studies (Quéméré *et al.*, 2012; Salmona *et al.*, 2020) supporting the a relative antiquity of
forest fragmentation in Northern Madagascar. Furthermore, both the high diversity of *Noronhia*
spinifolia, and its predominant distribution in low-elevation dry forest suggests that this habitat
type may have been spatially, topographically, and temporally extensive in northern Madagascar,
albeit frequently fragmented, as seemingly evidenced by a rare and likely relictual occurrence of
the species in contemporary high-elevation humid forest (e.g. Binara) and similarly peculiar
presence further north (e.g. Montagne des Français). To assess forest-cover changes over a larger
timeframe (e.g. the last ten or so millennia), inferences of *N. spinifolia*'s demography over time
would be relevant (Salmona *et al.*, 2017a; Beichman *et al.*, 2018). Coupling these inferences,
with that of short-generation grassland organisms, would also help clarifying the dynamics of
fire-prone open-canopy environments, through the succession of environmental changes that
occurred during last millennia, namely the last-glacial-maximum, early human's colonization,
the mid-Holocene transition, and the 1-Kya expansion of agro-pastoralism.

Further prospects and conservation implications

The power of coupling genomic data to landscape genetics allowed not only identifying major
landscape components influencing effective dispersal, but also their respective effects on seed

and pollen dispersal. This surprising result warrants further investigation using higher resolution
landscape and environmental layers, not used, or not available to our study. In particular, it
would benefit from the use of forest type, soil type, land use, and climate data of better
resolution. In addition, the wind effect has been tested without considering its directionality.
Recent analytical advances allowing wind directionality integration within a landscape genetics
framework (Fernández-López & Schliep, 2018) may allow to formally test its effect on pollen
dispersal. Furthermore, while our study clearly identifies that seed and pollen are dispersed by
forest-dwelling organisms, it neither identifies these organisms nor does it clearly show that seed
and pollen do still effectively disperse among forests. These questions could be tackled (i) by
inferring pedigree data from high density population sampling, coupled with sampling of young
trees and seedlings, (ii) using field survey of potential dispersers during flowering and
fructification (e.g. camera tracking), and/or (iii) using metabarcoding approaches to assess the
interaction network within the LM forests.

While our study confirms the biological importance of the LM region, which is known
for its species richness and endemism across taxa (Goodman & Wilmé, 2006; Rakotondravony,
2006, 2009; Sgarlata *et al.*, 2019), and more specifically for the genus *Noronhia* (Hong-Wa,
2016), our results also have several implications for biodiversity conservation in the region:
- First, they underscore the conservation value of the often-overlooked intraspecific genetic
diversity, which is unexpectedly high in *N. spinifolia*.
- Second, this study highlights the importance of riparian forests of the LM region for their major
role both as corridors connecting forest fragments, which is supported by the fact that genetic
diversity in *N. spinifolia* is explained by forest cover rather than Euclidian distance, and as
vectors promoting the roles of vertebrates and insects on seed and pollen dispersal. Therefore,
actively maintaining, protecting, and reforesting riparian and corridor forests, which are likely
pivotal for the functional connectivity of *N. spinifolia* but also most native and endemic species
of the LM region (Quéméré *et al.*, 2010; Rakotoarisoa *et al.*, 2013a; Sgarlata *et al.*, 2018; Aleixo-
Pais *et al.*, 2019), remain critical conservation actions.
- Third, our study identifies the Binara forest as unique among the major forests of the LM
region and in urgent need of deeper conservation focus. Indeed, our extensive forest survey
allowed us to find and collect just a few samples in this forest, where they were found only at
unexpectedly higher altitude and wetter habitat (Fig. S1). Similarly, several other Malagasy olive

species that are mostly distributed in dry forests (e.g. *N. ankaranensis*, *N. candicans*, *N.*

christenseniana and *N. oblanceolata*; GB and JS unpublished data), were also found to occur

only at higher altitude in the mountain evergreen forests of this region (e.g. Binara and

Antsahabe). Altogether, this pattern, though unclear, echoes the peculiarities of these forests, that

likely acted as refugia for numerous taxa during drier periods (Raxworthy & Nussbaum, 1995;

Goodman & Wilmé, 2006; Rakotoarisoa *et al.*, 2013b; Sgarlata *et al.*, 2019).

Data availability

634   Raw RADseq data and RADseq mtDNA alignments have been deposited to the Short Read

635   Archive (SRA) NCBI database under the reference PRJNA632767. Organellar microsatellite

636   genotypes and mtRAD variants are available in Tables S7 and S8, respectively. All additional

637   data, scripts and materials are available to readers at 10.5281/zenodo.4764651.

Conflict of interest disclosure

639   The authors of this article declare that they have no financial conflict of interest with the content

640   of this article.

Acknowledgments

We thank the Direction Générale du Ministère de l'Environnement et des Forêts de Madagascar, Madagascar's Ad Hoc Committee for Fauna and Flora, and Organizational Committee for Environmental Research (CAFF/CORE) for permission to perform this study (permits number: [49/17] & [127/18]/MEEF/SG/DGF/DSAP/SCB.Re) and for their support. We thank the local communities of Daraina and of the LM region for their warm reception and support. We thank E. Rasolondraibe and the many local guides and cooks for sharing their incomparable expertise and help in the field, *misaotra anareo jiaby*. We thank U. Suescun, C. Verbeke and M.A. Naranjo Arcos for lab assistance, L. Chikhi for comments on an earlier version of the draft and P. Raimondeau for sharing a complete, annotated mitogenome sequence of *N. clarinerva*. This work was mostly funded through an ERA-NET BiodivERsA project: INFRAGECO (Inference, Fragmentation, Genomics, and Conservation, ANR-16-EBI3-0014). We also thank the LABEX TULIP (ANR-10-LABX-0041) and CEBA (ANR-10-LABX-25-01), and the LIA BEEG-B (Laboratoire International Associé – Bioinformatics, Ecology, Evolution, Genomics and Behaviour, CNRS). We are grateful to the Get-Plage sequencing and Genotoul bioinformatics (Bioinfo Genotoul) platforms for sequencing services and providing computing resources.

Author Contribution

JS and GB designed the experiment. JS, AER, BLP, JR, CHW and GB were pivotal to field material collection and herbarium composition. JS, SM, and GB generated the genetic data. JS conducted bioinformatics and population genetic analyses. JS and AD conducted IBR analyses. JS and GB drafted a first version of the manuscript with a significant input from CHW. All co-authors agreed with the last version of the manuscript.

Supporting Information

References

Aleixo-Pais I, Salmona J, Sgarlata GM, Rakotonanahary A, Sousa AP, Parreira B, Kun-Rodrigues C, Ralantoharijaona T, Jan F, Rasolondraibe E, *et al.* 2019. The genetic structure of a mouse lemur living in a fragmented habitat in northern Madagascar. *Conservation Genetics* 20: 229–243.

Alexander DH, Novembre J, Lange K. 2009. Fast model-based estimation of ancestry in unrelated individuals. *Genome Research* 19: 1655–1664.

Allendorf FW, Hohenlohe PA, Luikart G. 2010. Genomics and the future of conservation genetics. *Nature Reviews Genetics* 11: 697–709.

Andrews KR, Good JM, Miller MR, Luikart G, Hohenlohe PA. 2016. Harnessing the power of RADseq for ecological and evolutionary genomics. *Nature Reviews Genetics* 17: 81–92.

Andrianoelina O, Favreau B, Ramamonjisoa L, Bouvet J-M. 2009. Small effect of fragmentation on the genetic diversity of *Dalbergia monticola*, an endangered tree species of the eastern forest of Madagascar, detected by chloroplast and nuclear microsatellites. *Annals of Botany* 104: 1231–1242.

Baguette M, Van Dyck H. 2007. Landscape connectivity and animal behavior: functional grain as a key determinant for dispersal. *Landscape Ecology* 22: 1117–1129.

Baird NA, Etter PD, Atwood TS, Currey MC, Shiver AL, Lewis ZA, Selker EU, Cresko WA, Johnson EA. 2008. Rapid SNP discovery and genetic mapping using sequenced RAD markers. *PLoS One* 3: e3376.

Balkenhol N, Cushman S, Storfer A, Waits L (Eds.). 2016. *Landscape genetics: concepts, methods, applications*. Oxford: John Wiley & Sons.

Beichman AC, Huerta-Sanchez E, Lohmueller KE. 2018. Using genomic data to infer historic population dynamics of nonmodel organisms. *Annual Review of Ecology, Evolution, and Systematics* 49: 433–456.

Beier P, Majka DR, Spencer WD. 2008. Forks in the road: choices in procedures for designing wildland linkages. *Conservation Biology* 22: 836–851.

Beier P, Spencer W, Baldwin RF, McRae BH. 2011. Toward best practices for developing regional connectivity maps. *Conservation Biology* 25: 879–892.

Besnard G, Hernández P, Khadari B, Dorado G, Savolainen V. 2011. Genomic profiling of plastid DNA variation in the Mediterranean olive tree. *BMC Plant Biology* 11: 80.

Besnard G, Khadari B, Baradat P, Bervillé A. 2002. Combination of chloroplast and mitochondrial DNA polymorphisms to study cytoplasm genetic differentiation in the olive complex (*Olea europaea* L.). *Theoretical and Applied Genetics* 105: 139–144.

Besnard G, Khadari B, Navascués M, Fernández-Mazuecos M, El Bakkali A, Arrigo N, Baali-
Cherif D, Brunini-Bronzini de Caraffa V, Santoni S, Vargas P. 2013. The complex history of the
olive tree: from Late Quaternary diversification of Mediterranean lineages to primary
domestication in the northern Levant. *Proceedings of the Royal Society B: Biological Sciences*
280: 20122833.

Bianconi ME, Dunning LT, Curran EV, Hidalgo O, Powell RF, Mian S, Leitch IJ, Lundgren
MR, Manzi S, Vorontsova MS, *et al.* 2020. Contrasted histories of organelle and nuclear
genomes underlying physiological diversification in a grass species. *Proceedings of the Royal*
Society B: Biological Sciences 287: 20201960.

Bond WJ, Silander Jr JA, Ranaivonasy J, Ratsirarson J. 2008. The antiquity of Madagascar's
grasslands and the rise of C4 grassy biomes. *Journal of Biogeography* 35: 1743–1758.

Bruvo R, Michiels NK, D'Souza TG, Schulenburg H. 2004. A simple method for the calculation
of microsatellite genotype distances irrespective of ploidy level. *Molecular Ecology* 13: 2101–
2106.

Cayuela H, Boualit L, Laporte M, Prunier JG, Preiss F, Laurent A, Foletti F, Clobert J, Jacob G.
2019. Kin-dependent dispersal influences relatedness and genetic structuring in a lek system.
Oecologia 191: 97–112.

Craul M, Radespiel U, Rasolofoson DW, Rakotondratsimba G, Rakotonirainy O,
Rasoloharijaona S, Randrianambinina B, Ratsimbazafy J, Ratelolahy F, Randrianamboavaonjy
T. 2008. Large rivers do not always act as species barriers for *Lepilemur* sp. *Primates* 49: 211–
218.

Dellicour S, Prunier JG, Piry S, Eloy M-C, Bertouille S, Licoppe A, Frantz AC, Flamand M-C.
2019. Landscape genetic analyses of *Cervus elaphus* and *Sus scrofa*: comparative study and
analytical developments. *Heredity* 123: 228–241.

Dixon P. 2003. VEGAN, a package of R functions for community ecology. *Journal of*
Vegetation Science 14: 927–930.

Dodd ME, Silvertown J, Chase MW. 1999. Phylogenetic analysis of trait evolution and species
diversity variation among angiosperm families. *Evolution* 53: 732–744.

Everson DA, Boucher DH. 1998. Tree species-richness and topographic complexity along the
riparian edge of the Potomac River. *Forest Ecology and Management* 109: 305–314.

Fahrig L. 2003. Effects of habitat fragmentation on biodiversity. *Annual Review of Ecology,*
Evolution, and Systematics 34: 487–515.

Fanamby. 2010. *Plan de gestion environnemental et de sauvegarde sociale (PGESS). Etude*
d'impact environnemental et social (EIES) de la Nouvelle Aire Protégée Loky-Manambato.

Fernández-López J, Schliep K. 2018. rWind: download, edit and include wind data in ecological
and evolutionary analysis. *Ecography* 42: 804–810.

Fontdevila A. 2005. Hybrid genome evolution by transposition. *Cytogenetic and Genome Research* 110: 49–55.

Frankham R. 2010. Challenges and opportunities of genetic approaches to biological conservation. *Biological Conservation* 143: 1919–1927.

Gardiner LM, Rakotoarinivo M, Rajaovelona LR, Clubbe C. 2017. Population genetics data help to guide the conservation of palm species with small population sizes and fragmented habitats in Madagascar. *PeerJ* 5: e3248.

Gardner CJ, Waeber PO, Razafindratsima OH, Wilmé L. 2018. Decision complacency and conservation planning. *Conservation Biology* 32: 1469–1472.

Garot E, Joët T, Combes M-C, Lashermes P. 2019. Genetic diversity and population divergences of an indigenous tree (*Coffea mauritiana*) in Reunion Island: role of climatic and geographical factors. *Heredity* 122: 833–847.

Gautier L, Ranirison P, Nusbaumer L, Wohlhauser S. 2006. Aperçu des massifs forestiers de la région Loky-Manambato. *Inventaires de la faune et de la flore du nord de Madagascar dans la région Loky-Manambato, Analamerana et Andavakoera. Série Sciences Biologiques* 23: 81–99.

Godfrey LR, Crowley BE. 2016. Madagascar's ephemeral palaeo-grazer guild: who ate the ancient C4 grasses? *Proceedings of the Royal Society B: Biological Sciences* 283: 20160360.

Goodman SM, Benstead JP. 2003. *Natural history of Madagascar*. Chicago and London: University Chicago Press.

Goodman SM, Raherilalao MJ, Wohlhauser S. 2018. *The terrestrial protected areas of Madagascar: their history, description, and biota*. Chicago: University of Chicago Press.

Goodman SM, Wilmé L. 2006. Inventaires de la faune et de la flore du nord de Madagascar dans la région Loky-Manambato, Analamerana et Andavakoera. *Recherche pour le Dévelopement* 23: 1–238.

Goudie AS. 2018. *Human impact on the natural environment: Past, present and future*. Hoboken, NJ: Wiley-Blackwell.

Graves TA, Beier P, Royle JA. 2013. Current approaches using genetic distances produce poor estimates of landscape resistance to interindividual dispersal. *Molecular Ecology* 22: 3888–3903.

Hackel J, Vorontsova MS, Nanjarisoa OP, Hall RC, Razanatsoa J, Malakasi P, Besnard G. 2018. Grass diversification in Madagascar: in situ radiation of two large C3 shade clades and support for a Miocene to Pliocene origin of C4 grassy biomes. *Journal of Biogeography* 45: 750–761.

Haddad NM, Brudvig LA, Clobert J, Davies KF, Gonzalez A, Holt RD, Lovejoy TE, Sexton JO, Austin MP, Collins CD. 2015. Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances* 1: e1500052.

Hall N, Mercer L, Phillips D, Shaw J, Anderson AD. 2012. Maximum likelihood estimation of
individual inbreeding coefficients and null allele frequencies. *Genetics Research* 94: 151–161.

Hansen MC, Potapov PV, Moore R, Hancher M, Turubanova SA, Tyukavina A, Thau D,
Stehman SV, Goetz SJ, Loveland TR. 2013. High-resolution global maps of 21st-century forest
cover change. *Science* 342: 850–853.

Harper GJ, Steininger MK, Tucker CJ, Juhn D, Hawkins F. 2007. Fifty years of deforestation
and forest fragmentation in Madagascar. *Environmental Conservation* 34: 325–333.

Heller R, Nursyifa C, Garcia Erill G, Salmona J, Chikhi L, Meisner J, Korneliussen TS,
Albrechtsen A. 2021. A reference-free approach to analyze non-model RADseq data using
standard Next Generation Sequencing toolkits. *Molecular Ecology Resources* 21: 1085–1097.

Holderegger R, Buehler D, Gugerli F, Manel S. 2010. Landscape genetics of plants. *Trends in
Plant Science* 15: 675–683.

Hong-Wa C. 2016. A taxonomic revision of the genus *Noronhia* Stadtm. ex Thouars (Oleaceae)
in Madagascar and the Comoro Islands. *Boissiera* 70: 1–291.

Hong-Wa C, Besnard G. 2014. Species limits and diversification in the Madagascar olive
(*Noronhia*, Oleaceae). *Botanical Journal of the Linnean Society* 174: 141–161.

Isabel N, Holliday JA, Aitken SN. 2020. Forest genomics: Advancing climate adaptation, forest
health, productivity, and conservation. *Evolutionary Applications* 13: 3–10.

Joseph GS, Seymour CL. 2020. Madagascan highlands: originally woodland and forest
containing endemic grasses, not grazing-adapted grassland. *Proceedings of the Royal Society B*
287: 20201956.

Joseph GS, Seymour CL. 2021. The unlikely 'antiquity of Madagascar's grasslands':
Disproportionately forest☐limited endemic fauna support anthropogenic transformation from
woodland. *Journal of Biogeography* DOI: 10.1111/jbi.14132.

Kamvar ZN, Brooks JC, Grünwald NJ. 2015. Novel R tools for analysis of genome-wide
population genetic data with emphasis on clonality. *Frontiers in Genetics* 6: 208.

Keller D, Holderegger R. 2013. Damselflies use different movement strategies for short-and
long-distance dispersal. *Insect Conservation and Diversity* 6: 590–597.

Kimura M. 1983. *The neutral theory of molecular evolution*. Cambridge University Press.

Korneliussen TS, Albrechtsen A, Nielsen R. 2014. ANGSD: analysis of next generation
sequencing data. *BMC Bioinformatics* 15: 356.

Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004.
Versatile and open software for comparing large genomes. *Genome Biology* 5: R12.

Landguth EL, Cushman SA, Schwartz MK, McKelvey KS, Murphy M, Luikart G. 2010.
Quantifying the lag time to detect barriers in landscape genetics. *Molecular Ecology* 19: 4179–
4191.

Laurance WF, Nascimento HE, Laurance SG, Andrade A, Ewers RM, Harms KE, Luizao RC,
Ribeiro JE. 2007. Habitat fragmentation, variable edge effects, and the landscape-divergence
hypothesis. *PLoS One* 2: e1017.

LeBuhn G, Droege S, Connor EF, Gemmill-Herren B, Potts SG, Minckley RL, Jean RP, Kula E,
Roubik DW, Wright KW. 2015. Evidence-based conservation: reply to Tepedino et al.
Conservation Biology 29: 283–285.

Legendre P, Lapointe F-J, Casgrain P. 1994. Modeling brain evolution from behavior: a
permutational regression approach. *Evolution* 48: 1487–1499.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv preprint arXiv:1303.3997.

Lindenmayer DB, Fischer J. 2013. *Habitat fragmentation and landscape change: an ecological*
and conservation synthesis. Washington, DC: Island Press.

Ma T, Wang K, Hu Q, Xi Z, Wan D, Wang Q, Feng J, Jiang D, Ahani H, Abbott RJ. 2018.
Ancient polymorphisms and divergence hitchhiking contribute to genomic islands of divergence
within a poplar species complex. *Proceedings of the National Academy of Sciences of the United*
States of America 115: E236–E243.

Mantel N. 1967. The detection of disease clustering and a generalized regression approach.
Cancer Research 27: 209.

McRae BH. 2006. Isolation by resistance. *Evolution* 60: 1551–1561.

McRae BH, Beier P. 2007. Circuit theory predicts gene flow in plant and animal populations.
Proceedings of the National Academy of Sciences of the United States of America 104: 19885–
19890.

Meisner J, Albrechtsen A. 2018. Inferring population structure and admixture proportions in
low-depth NGS data. *Genetics* 210: 719–731.

Mona S, Ray N, Arenas M, Excoffier L. 2014. Genetic consequences of habitat fragmentation
during a range expansion. *Heredity* 112: 291–299.

Murcia C. 1995. Edge effects in fragmented forests: implications for conservation. *Trends in*
Ecology & Evolution 10: 58–62.

Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GA, Kent J. 2000. Biodiversity hotspots
for conservation priorities. *Nature* 403: 853–858.

Nater A, Greminger MP, Arora N, Schaik CP, Goossens B, Singleton I, Verschoor EJ, Warren
KS, Krützen M. 2015. Reconstructing the demographic history of orang-utans using
Approximate Bayesian Computation. *Molecular Ecology* 24: 310–327.

Nei M. 1972. Genetic distance between populations. *The American Naturalist* 106: 283–292.

Nei M. 1973. Analysis of gene diversity in subdivided populations. *Proceedings of the National*
Academy of Sciences of the United States of America 70: 3321–3323.

Nei M. 1987. *Molecular evolutionary genetics*. Columbia University Press.

Nielsen R, Korneliussen T, Albrechtsen A, Li Y, Wang J. 2012. SNP calling, genotype calling,
and sample allele frequency estimation from new-generation sequencing data. *PLoS One* 7:
e37558.

Olofsson JK, Dunning LT, Lundgren MR, Barton HJ, Thompson J, Cuff N, Ariyarathne M,
Yakandawala D, Sotelo G, Zeng K. 2019. Population-specific selection on standing variation
generated by lateral gene transfers in a grass. *Current Biology* 29: 3921–3927.

Paris JR, Stevens JR, Catchen JM. 2017. Lost in parameter space: a road map for stacks.
Methods in Ecology and Evolution 8: 1360–1373.

Payseur BA, Rieseberg LH. 2016. A genomic perspective on hybridization and speciation.
Molecular Ecology 25: 2337–2360.

Pedersen C-ET, Albrechtsen A, Etter PD, Johnson EA, Orlando L, Chikhi L, Siegismund HR,
Heller R. 2018. A southern African origin and cryptic structure in the highly mobile plains zebra.
Nature Ecology & Evolution 1: 491–498.

Pembleton LW, Cogan NO, Forster JW. 2013. St AMPP: An R package for calculation of
genetic differentiation and structure of mixed-ploidy level populations. *Molecular Ecology*
Resources 13: 946–952.

Peterman WE. 2018. ResistanceGA: An R package for the optimization of resistance surfaces
using genetic algorithms. *Methods in Ecology and Evolution* 9: 1638–1647.

Petit RJ, Hampe A. 2006. Some evolutionary consequences of being a tree. *Annual Review of*
Ecology, Evolution, and Systematics 37: 187–214.

Pillon Y, Hopkins HC, Rigault F, Jaffré T, Stacy EA. 2014. Cryptic adaptive radiation in tropical
forest trees in New Caledonia. *New Phytologist* 202: 521–530.

Poelstra J, Salmona J, Tiley GP, Schüßler D, Blanco MB, Andriambeloson JB, Bouchez O,
Campbell CR, Etter PD, Hohenlohe PA, *et al.* 2021. Cryptic patterns of speciation in cryptic
primates: microendemic mouse lemurs and the multispecies coalescent. *Systematic Biology*:
10.1093/sysbio/syaa053.

Pressey RL, Cabeza M, Watts ME, Cowling RM, Wilson KA. 2007. Conservation planning in a
changing world. *Trends in Ecology & Evolution* 22: 583–592.

Prevosti A, Ocana J, Alonso G. 1975. Distances between populations of *Drosophila subobscura*,
based on chromosome arrangement frequencies. *Theoretical and Applied Genetics* 45: 231–241.

Prunier JG, Colyn M, Legendre X, Flamand M-C. 2017. Regression commonality analyses on
hierarchical genetic distances. *Ecography* 40: 1412–1425.

Prunier JG, Colyn M, Legendre X, Nimon KF, Flamand M-C. 2015. Multicollinearity in spatial
genetics: separating the wheat from the chaff using commonality analyses. *Molecular Ecology*
24: 263–283.

Quéméré E, Amelot X, Pierson J, Crouau-Roy B, Chikhi L. 2012. Genetic data suggest a natural
prehuman origin of open habitats in northern Madagascar and question the deforestation
narrative in this region. *Proceedings of the National Academy of Sciences of the United States of*
America 109: 13028–13033.

Quéméré E, Crouau-Roy B, Rabarivola C, Louis EE, Chikhi L. 2010. Landscape genetics of an
endangered lemur (*Propithecus tattersalli*) within its entire fragmented range. *Molecular*
Ecology 19: 1606–1621.

Rakotoarisoa J-E, Raheriarisena M, Goodman SM. 2013a. A phylogeographic study of the
endemic rodent *Eliurus carletoni* (Rodentia: Nesomyinae) in an ecological transition zone of
northern Madagascar. *Journal of Heredity* 104: 23–35.

Rakotoarisoa J-E, Raheriarisena M, Goodman SM. 2013b. Late Quaternary climatic vegetational
shifts in an ecological transition zone of northern Madagascar: insights from genetic analyses of
two endemic rodent species. *Journal of Evolutionary Biology* 26: 1019–1034.

Rakotondravony HA. 2006. Patterns de la diversité des reptiles et amphibiens de la région de
Loky-Manambato. *Inventaires de la faune et de la flore du nord de Madagascar dans la région*
Loky-Manambato, Analamerana et Andavakoera. Série Sciences Biologiques 23: 101–148.

Rakotondravony HA. 2009. Aspects de la conservation des reptiles et des amphibiens dans la
région de Daraina. *Madagascar Conservation & Development* 1: 15–18.

Raxworthy CJ, Nussbaum RA. 1995. Systematics, speciation and biogeography of the dwarf
chameleons (*Brookesia*; Reptilia, Squamata, Chamaeleontidae) of northern Madagascar. *Journal*
of Zoology 235: 525–558.

Reynolds J, Weir BS, Cockerham CC. 1983. Estimation of the coancestry coefficient: basis for a
short-term genetic distance. *Genetics* 105: 767–779.

Rochette NC, Rivera-Colón AG, Catchen JM. 2019. Stacks 2: Analytical methods for paired-end
sequencing improve RADseq-based population genomics. *Molecular Ecology* 28: 4737–4754.

Salmona J, Heller R, Lascoux M, Shafer A. 2017a. Inferring demographic history using genomic
data. In: Rajora O, ed. *Population Genomics*. Cham: Springer, 511–537.

Salmona J, Heller R, Quéméré E, Chikhi L. 2017b. Climate change and human colonization
triggered habitat loss and fragmentation in Madagascar. *Molecular Ecology* 26: 5203–5222.

Salmona J, Olofsson JK, Hong-Wa C, Razanatsoa J, Rakotonasolo F, Ralimanana H,
Randriamboavonjy T, Suescun U, Vorontsova MS, Besnard G. 2020. Late Miocene origin and
recent population collapse of the Malagasy savanna olive tree (*Noronhia lowryi*). *Biological
Journal of the Linnean Society* 129: 227–243.

Schmieder R, Edwards R. 2011. Fast identification and removal of sequence contamination from
genomic and metagenomic datasets. *PLoS One* 6: e17288.

Scotti I, González-Martínez SC, Budde KB, Lalagüe H. 2016. Fifty years of genetic studies:
what to make of the large amounts of variation found within populations? *Annals of Forest
Science* 73: 69–75.

Sgarlata GM, Salmona J, Aleixo-Pais I, Rakotonanahary A, Sousa AP, Kun-Rodrigues C,
Ralantoharijaona T, Jan F, Zaranaina R, Rasolondraibe E, *et al.* 2018. Genetic differentiation and
demographic history of the northern rufous mouse Lemur (*Microcebus tavaratra*) across a
fragmented landscape in northern Madagascar. *International Journal of Primatology* 39: 65–89.

Sgarlata GM, Salmona J, Le Pors B, Rasolondraibe E, Jan F, Ralantoharijaona T,
Rakotonanahary A, Randriamaroson J, Marques AJ, Aleixo-Pais I, *et al.* 2019. Genetic and
morphological diversity of mouse lemurs (*Microcebus* spp.) in northern Madagascar: The
discovery of a putative new species? *American Journal of Primatology* 81: e23070.

Skotte L, Korneliussen TS, Albrechtsen A. 2013. Estimating individual admixture proportions
from next generation sequencing data. *Genetics* 195: 693–702.

Slatkin M. 1993. Isolation by distance in equilibrium and non-equilibrium populations. *Evolution*
47: 264–279.

Smouse PE, Long JC, Sokal RR. 1986. Multiple regression and correlation extensions of the
Mantel test of matrix correspondence. *Systematic Zoology* 35: 627–632.

Solofondranohatra CL, Vorontsova MS, Hackel J, Besnard G, Cable S, Williams J, Jeannoda V,
Lehmann CE. 2018. Grass functional traits differentiate forest and savanna in the Madagascar
central highlands. *Frontiers in Ecology and Evolution* 6: 184.

Štorchová H, Olson MS. 2004. Comparison between mitochondrial and chloroplast DNA
variation in the native range of *Silene vulgaris*. *Molecular Ecology* 13: 2909–2919.

Sutherland WJ, Pullin AS, Dolman PM, Knight TM. 2004. The need for evidence-based
conservation. *Trends in Ecology & Evolution* 19: 305–308.

Tang Q, Fung T, Rheindt FE. 2020. ResDisMapper: An r package for fine-scale mapping of
resistance to dispersal. *Molecular Ecology Resources* 20: 819–831.

Van de Paer C. 2017. *Structural diversity and contrasted evolution of cytoplasmic genomes in*
flowering plants: a phylogenomic approach in Oleaceae. PhD thesis, University of Toulouse III-
Paul Sabatier.

Van Etten J. 2012. R package gdistance: distances and routes on geographical grids (version 1.1-
4). *Journal of Statistical Software* 76: 13.

Van Strien MJ, Holderegger R, Van Heck HJ. 2015. Isolation-by-distance in landscapes:
considerations for landscape genetics. *Heredity* 114: 27–37.

Vences M. 2005. Madagascar as a model region for the study of tempo and pattern in adaptive
radiations. In: Huber BA, Sinclair BJ, Lampe KH, eds. *Molecules, Organisms, Ecosystems.*
African Biodiversity. Boston, MA: Springer, 69–84.

Vences M, Wollenberg KC, Vieites DR, Lees DC. 2009. Madagascar as a model region of
species diversification. *Trends in Ecology & Evolution* 24: 456–465.

Vieilledent G, Grinand C, Rakotomalala FA, Ranaivosoa R, Rakotoarijaona J-R, Allnutt TF,
Achard F. 2018. Combining global tree cover loss data with historical national forest cover maps
to look at six decades of deforestation and forest fragmentation in Madagascar. *Biological*
Conservation 222: 189–197.

Vieira FG, Fumagalli M, Albrechtsen A, Nielsen R. 2013. Estimating inbreeding coefficients
from NGS data: impact on genotype calling and allele frequency estimation. *Genome Research*
23: 1852–1861.

Vorontsova MS, Besnard G, Forest F, Malakasi P, Moat J, Clayton WD, Ficinski P, Savva GM,
Nanjarisoa OP, Razanatsoa J. 2016. Madagascar's grasses and grasslands: anthropogenic or
natural? *Proceedings of the Royal Society, London, Series B* 283: 20152262.

Wang Y, Lu J, Yu J, Gibbs RA, Yu F. 2013. An integrative variant analysis pipeline for accurate
genotype/haplotype inference in population NGS data. *Genome research* 23: 833–842.

Warmuth VM, Ellegren H. 2019. Genotype-free estimation of allele frequencies reduces bias and
improves demographic inference from RADSeq data. *Molecular Ecology Resources* 19: 586–
596.

Wilmé L, Goodman SM, Ganzhorn JU. 2006. Biogeographic evolution of Madagascar's
microendemic biota. *Science* 312: 1063–1065.

Wright S. 1943. Isolation by distance. *Genetics* 28: 114–138.

Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, Madden PA, Heath AC,
Martin NG, Montgomery GW. 2010. Common SNPs explain a large proportion of the
heritability for human height. *Nature Genetics* 42: 565.

Yoder AD, Campbell CR, Blanco MB, Dos Reis M, Ganzhorn JU, Goodman SM, Hunnicutt KE,
Larsen PA, Kappeler PM, Rasoloarison RM. 2016. Geogenetic patterns in mouse lemurs (genus
Microcebus) reveal the ghosts of Madagascar's forests past. *Proceedings of the National*
Academy of Sciences of the United States of America 113: 8049–8056.

1028