Efficient GNE myopathy disease modeling with mutation specific phenotypes in

human pluripotent stem cells by base editors

Running title: Mutation specific GNE modeling by BEs

Ju-Chan Park[1*], Jumee Kim[1*], Hyun-Ki Jang[2,6], Seung-Yeon Lee[1], Keun-Tae Kim[1],

Seokwoo Park[3], Hyun Sik Lee[4], Hee-Jung Choi[4], Soon-Jung Park[5], Sung-Hwan

Moon[5], Sangsu Bae[2,6] and Hyuk-Jin Cha[1#]

[1]College of Pharmacy, Seoul National University, Seoul, Republic of Korea, [2]Institute

for Convergence of Basic Sciences, Hanyang University, Seoul, Republic of Korea,

[3]Department of Biomedical Sciences, Seoul National University College of Medicine,

Seoul, South Korea, [4]Department of Biological Sciences, Seoul National University,

Seoul, South Korea, [5]Stem Cell Research Institute, T&R Biofab Co. Ltd, Siheung,

Republic of Korea, [6]Department of Chemistry, Hanyang University, Seoul, Republic

of Korea

* These authors contributed equally

To whom correspondence should be addressed to

Prof. Hyuk-Jin Cha, Ph.D.

College of Pharmacy, Seoul National University

1 Gwanak-ro, Gwanak-gu, Seoul 08826, Republic of Korea

Tel.: +82-2-880-7825; Fax: +82-2-880-9122; E-mail: hjcha93@snu.ac.kr

Keywords: Gene editing, Base editor, human pluripotent stem cells, disease modeling,

GNE myopathy, hyposialylation

Summary

Despite the great potential of disease modeling with the isogenic pairs of

human pluripotent stem cells (hPSCs), the extremely low efficiency of precise gene

editing in hPSCs remains a technical hurdle for this approach. Herein, we took

advantage of currently available base editors (BEs) to epitomize the isogenic disease

model from hPSCs. Using this method, we established 14 hPSCs that harbor point

mutations on the *GNE* gene, including four different mutations found in GNE

myopathy patients. Due to lesser activation of p53 by BEs than Cas9, a higher editing

efficiency with BEs was achieved. Four different mutations in the epimerase or kinase

domains of *GNE* revealed mutation-specific hyposialylation, which was closely

correlated to pathological clinical phenotypes. These mutation-specific

hyposialylation patterns were evident in GNE protein structure modeling.

Furthermore, treatment with a drug candidate currently under clinical trials showed a

mutation-specific drug response in GNE myopathy disease models. These data

suggest that isogenic disease models from hPSCs using BEs could serve as a useful

tool for mimicking the pathophysiology of GNE myopathy and for predicting drug

responses.

Introduction

Since the establishment of induced pluripotent stem cells (iPSCs), great

advances in both autologous stem cell therapy and disease modeling from patient-

specific iPSCs has been achieved (Rowe and Daley, 2019). The "disease-in-a-dish"

model (Tiscornia et al., 2011) enables the identification of unknown disease

mechanisms and drug screening once the disease phenotype indicator has been

determined (Shi et al., 2017). However, the phenotypes arising from the disparate

genetic backgrounds of the control and disease iPSCs are often more robust than those

of the disease itself, which hinders translationally relevant comparison (Merkle and

Eggan, 2013). The establishment of isogenic pairs from disease to control human

pluripotent stem cells (hPSCs) or vice versa using precise gene editing technology

was proposed as a way to minimize this bias (Merkle and Eggan, 2013; Musunuru,

2013).

Considering that the majority (58%) of genetic diseases result from point

mutations (Rees and Liu, 2018), precise gene editing technologies such as base

substitution based on homology-directed repair with Cas9 (hereafter referred to as

HDR) (Izmiryan et al., 2018; Jackow et al., 2019) and base editing (BE) (Osborn et al.,

2020) have been applied in human iPSCs to correct genetic mutations in patient iPSCs.

However, it is generally accepted that the gene editing efficiency of Cas9 *via* HDR in

hPSCs is extremely low compared to other cell lines (Mali et al., 2013), which

remains a technical challenge for hPSC-based disease modeling. Thus, various

approaches have been developed to improve editing efficiency (Kim et al., 2020;

Maurissen and Woltjen, 2020). One recent study demonstrated that p53-dependent

cell death, induced by a DNA double-strand break (DSB) that results from the

endonuclease activity of Cas9, is responsible for its low gene editing efficiency in

hPSCs (Ihry et al., 2018).

While HDR operates in tandem with DSB, base substitution by base editors

(BEs) is achieved through deamination of the target base by deaminase conjugated

with nickase Cas9 (nCas9), without inducing DSB (Gaudelli et al., 2017; Komor et al.,

2016; Rees and Liu, 2018). Target sequence accessibility (hereafter referred to as

"target-ability") of BEs is determined by the protospacer adjacent motif (PAM)

sequence that contains the canonical trinucleotide NGG (NGG-PAM). As the target-

ability of BEs is restricted by the PAM sequence, BEs based on the NG-PAM

sequence were recently developed to overcome this limitation (Huang et al., 2019;

Nishimasu et al., 2018).

GNE myopathy (OMIM #605820), a rare autosomal recessive degenerative

skeletal muscle disorder, is caused by bi-allelic loss of function (LOF) mutations of

the *GNE* gene. The enzymatic activity of the GNE protein depends in turn upon N-

acetylmannosamine (ManNAc) kinase activity and the epimerase activity of UDP-N-

acetylglucosamine, which is the rate-limiting step in sialic acid biosynthesis. The

reduced overall enzymatic activity of the GNE protein leads to hyposialylation, which

is thought to cause the progressive muscle weakness characteristic of GNE

myopathy(Noguchi et al., 2004). There are no FDA-approved therapies for GNE

myopathy (Nishino et al., 2015). The only drug candidate in clinical trials is N-

acetylmannosamine (ManNAc), the intermediate product in the synthesis of sialic

acid (Xu et al., 2017). Due to the unavailability of human GNE disease muscle cells,

immortalized cell lines (Pham et al., 2017) or mouse models (Chan et al., 2017) were

alternatively used to study the mechanism. However, these models are insufficient to

mimic pathophysiology in human muscle cells (Perlman, 2016). Thus, an appropriate

"GNE myopathy in a dish" model is required to characterize human GNE pathophysiology.

To establish the isogenic disease model, we operationally defined the GNE disease model as attenuated sialic acid production in GNE mutants, to serve as a pathophenotype indicator. First, we demonstrated that BEs (Huang et al., 2019) were more efficient than HDR in hPSCs due to lesser induction of p53-dependent cell death. Using this method, we produced 14 hPSCs with point mutations on the *GNE* gene. Isogenic pairs of hESCs with four different mutations and myocytes derived from these mutant hESCs revealed mutation-specific hyposialylation. Additionally, the 3D structure modeling of each GNE mutant was consistent with altered GNE enzymatic activity, correlating with different degrees of hyposialylation. The differing responses of each mutant upon drug candidate treatment suggests that the mutation status of GNE patients is an important factor to consider in clinical trials of GNE myopathy drug candidates.

Results

Target-ability of point mutations causing GNE myopathy by base editors

BEs, such as the adenine base editor (ABE) and cytosine base editor (CBE),

substitute the target base through the action of the deaminases TadA or APOBEC1,

respectively, conjugated with nCas9, which induces single-strand breaks (SSB) (Fig.

1A). Both BEs and HDR are potential strategies for the production of isogenic pairs

of disease models with hPSCs. Thus, prior to producing hPSCs for GNE myopathy

modeling, we first investigated the "target-ability" of mutations causing GNE

myopathy from the ClinVar database (https://www.ncbi.nlm.nih.gov/clinvar/)

(Landrum et al., 2016). Among the total 63 mutations found in GNE patients that are

categorized as "pathogenic" or "likely pathogenic," 34 mutations (54%) occur by a

transition point mutation (purine to purine or pyrimidine to pyrimidine) and are

targetable with ABE or CBE (Fig. 1B and Table S1). Eight mutations (12.7%) are

transversion mutations (purine to pyrimidine or vice versa), which are targetable with

either the prime editor (PE) or HDR. Among transition point mutations, for

establishing isogenic pairs for a GNE model (from mutant to wild-type or vice versa),

73.5% of wild-type (25 out of 34 mutations) (Fig. 1C) and 70.6% of mutants (24 out

of 34 mutations) (Fig. 1D) are targetable by BEs. As mentioned above, 26.5% of

wild-type (nine mutations) and 29.4% of mutants (10 mutations) cannot be edited due

to the lack of a PAM sequence. The target-ability of BEs has greatly expanded due to

advances in BE techniques using NG-PAM (Huang et al., 2019; Nishimasu et al.,

2018) (e.g., NG-ABE or NG-CBE) (Figs. 1C and D). Interestingly, the relatively

frequent spontaneous deamination of cytosine to uracil in nature may account for the

differential utility of CBE (for C to T mutation) and ABE (for T to C mutation) for

producing mutants and wild-type lines, respectively (Figs. 1C and D).

Efficiency of base editing compared to HDR in hESCs

Considering that GNE myopathy occurs by loss of function (LOF) mutations of either the epimerase or kinase domains (Celeste et al., 2014), it is necessary to examine the functional effect of mutation in these domains of the GNE protein. For this, we chose four mutations of the epimerase (R160Q and I329T) and kinase (I588T and V272M) domains that would be readily achievable by ABE, CBE, and HDR (Fig. 2A). Compared to BEs, the HDR technique is more broadly targetable to all point mutations regardless of whether they are transversion or transition mutations (Rees and Liu, 2018). Thus, we first compared the editing efficiencies of all currently available techniques, HDR (Yang et al., 2013), ABE (ABEmax, (Koblan et al., 2018)), NG-ABE (NG-ABEmax, (Huang et al., 2019)), CBE (BE4max, (Koblan et al., 2018)) and NG-CBE (NG-BE4max, (Huang et al., 2019)), to generate point mutations of target sequences in hESCs. The efficiency of base substitution for R160Q by CBE was close to 37% (Fig. 2B), while those of NG-CBE and HDR were 4.1% and 0.93%, respectively (Figs. 2B and S1A). Similarly, 32.5% base substitution was achieved by ABE for I588T, while NG-ABE yielded 11.2% (Fig. S1B) and HDR yielded only 1% (Fig. 2C) substitution. For the V727M mutation, on-target base substitution efficiencies were 55.8% and 10.1% with CBE and NG-CBE (Fig. 2D and S1C), respectively, while HDR achieved 3.0% editing efficiency (Fig. 2D). Base substitution for I329T with HDR, ABE, and NG-ABE showed similar results, wherein the efficiency of HDR was lower (4.5% for I329T) than that of ABE (8.6%) and NG-ABE (6.3%) (Figs. 2E and S1D). However, the existence of bystander sequences in the target window reduced the observed percentage of base substitution without bystander mutation in V727M (Fig. S1E). Additionally, two mutants with bystander

mutation (I329I and I329T) occurred in the I329T target due to codon degeneracy

(Fig. S1F). The GNE mutant hESCs with bystander mutations are listed in Table S2.

As low HDR efficiency compared to indel by Cas9 for GNE mutations (Fig.

S1G) demands laborious clonal selection in hPSCs (Li et al., 2016), the inefficiency

of HDR remains a serious technical limitation to be resolved for disease modeling in

hPSCs despite the advantage of broader target-ability. Furthermore, we observed that

NGG-BEs (ABE and CBE) were more efficient than NG-BEs (NG-ABE and NG-

CBE) as previously reported (Jeong et al., 2019).

P53 response by Cas9 determines editing efficiency

The extremely low gene editing efficiency with Cas9 in hPSCs (Kim et al.,

2020; Mali et al., 2013) results from p53-dependent cell death upon DSB caused by

the endonuclease activity of Cas9 (Ihry et al., 2018). hPSCs are particularly

susceptible to p53-dependent cell death once DSB occurs, which is thought to be a

protective mechanism to rigorously maintain genome integrity during development

(Weissbein et al., 2014). Thus, we hypothesized that the low editing efficiency of

HDR compared to BEs (Fig. 2) may result from different p53 responses. To explore

this hypothesis, we first established *TP53* KO hESCs by targeting exon 3 of *TP53*

(Fig. S2A). After introducing sgRNA and Cas9 in hESCs, Nutlin3, which is an

inhibitor of MDM2, was applied to stabilize p53, causing massive cell death, as

expected (Fig. S2B). After Nutlin3 treatment, the few clones that survived were *TP53*

KO hESCs (Fig. S2C). The p53-dependent response of Cas9 or BEs was assessed by

the measuring the level of *PPM1D*, a typical target gene of p53 (Fiscella et al., 1997).

When introducing Cas9 with sgRNA into hESCs as previously described (Ihry et al.,

2018), BEs only marginally affected the p53 response (Fig. 2F). Consistent with the

p53 response, significantly more hESCs remained and survived when BEs were used,

compared to the Cas9 approach, while all *TP53* KO hESCs tolerated the stress of gene

editing (Fig. 2G). The effects of gene editing techniques in WT and *TP53* KO hESCs

were monitored in real-time (Movies S1 and S2). Along with the higher rate of

survival in *TP53* KO hESCs than in wild-type hESCs (Fig. 2G), improvement of the

gene editing efficiency of Cas9 was evident in *TP53* KO hESCs (Fig. 2H).

Establishment of GNE myopathy disease models through base editors

Even with the limited target-ability of BEs (Figs. 1C and D), significant higher

gene editing efficiency of BEs than HDR may enable BEs to be currently the most

feasible approach for base substitution in hPSCs. Accordingly, we successfully

established 14 GNE mutants in total in both hESCs and hiPSCs using BEs (Table S2).

Briefly, based on H9 hESCs, two LOF mutants in the epimerase domain of GNE

generated with CBE (R160Q) and ABE (I329T) (Figs. 3A and B) and two LOF

mutants in the kinase domain using NG-ABE (I588T) and CBE (V727M) were

established (Figs. 3C and D). Other GNE mutant cell lines (I329T, R160Q, and I588T)

were established in CHA3-hESCs (Fig. S3D) and BJ-iPSCs (Figs. S3E and F) using

NG-ABE, ABE, or CBE, as listed in Table S2. Of note, since the target windows of

I329T and V727M carry bystander bases, five mutants with bystander mutations

including I329I, a silent mutation, were also established in V727M (Figs. S3B and C)

and I329T (Fig. S3D).

Mutation-specific phenotype of GNE myopathy disease models

The four established GNE mutant hESCs (GNE-hESCs) displayed typical

cellular morphology and comparable OCT4 expression (Fig. 4A). As the putative

underlying cause of GNE myopathy is hyposialylation, we first assessed levels of

sialic acid production using a sialic acid-specific *Sambucus nigra agglutinin* (SNA)

lectin binding assay in four different GNE-hESCs. As shown in Figure 4B, the levels

of lectin staining were significantly diminished in all mutant hESCs to different

degrees, relative to controls. The mean fluorescence intensity (MFI) of SNA lectin

clearly indicated that the sialylation of the protein varied depending on the mutation

(Figs. 4C and D). Of note, binding of SNA lectin indicates the level of sialylation of

protein, the final step for sialic acid biosynthesis, is a reliable indicator of GNE

enzymatic activity. In this line, MFI of SNA staining in GNE mutants relative to WT

(Fig. 4D) can be used to determine pathogenicity. Thus, we compared previously

predicted severity of each mutations by *in silico* platforms (PolyPhen2, SIFT, Align,

and Pmut) (Celeste et al., 2014) to the MFI of each GNE mutant (Fig. 4E). I329T

mutant, predicted as "severe" revealed the most obvious hyposialylation (Figs. 4D

and E). As the V727M mutation occurs in the compound heterozygous state, even

with the prevalent distribution of this variant (rs121908627), this mutation itself

would have a less serious effect on GNE myopathy according to the moderate

hyposialylation in the V727M mutant (Fig. 4C). We would expect the I588T mutant,

which is a variant of uncertain significance (VUS), to be "Pathogenic/Likely

pathogenic" since hyposialylation levels in this mutant are comparable to those

observed in the R160Q mutant, which is classified as "Pathogenic/Likely pathogenic"

(Fig. 4E). The hyposialylation of other GNE models derived from BJ-iPSC and

hCHA3 hESCs were examined. Evident hyposialylation (Fig. S4A) and recovery by

sialic acid treatment (Fig. S4B), compared to that of the silent I329I mutation

established in hCHA3 cells, ruled out the possibility of baseline effects of gene

editing itself on hyposialylation. The comparable degree of hyposialylation of R160Q

and I588T was also reproduced in GNE mutants derived from BJ-iPSCs (Figs. S4C

and D). Importantly, the levels of sialic acid in four different hPSC lines (two hESCs

and two iPSCs), differed in a cell type-specific manner depending on genetic

background (Fig. 4F) despite comparable level of mRNA expression of *GNE* (Fig.

4G). This supports the necessity of an isogenic pair for precise comparison as

highlighted previously (Merkle and Eggan, 2013).

Structural analysis predicts mutation-specific hyposialylation

The varying degrees of hyposialylation among the GNE mutants inspired us to

analyze each mutation at the molecular level. We used the GNE structure to predict

the molecular consequences of each mutation on the enzymatic activity of GNE. The

epimerization reaction of GNE requires tetramerization through the N-terminal

domain, which is stabilized upon ligand binding (Fig. 5A) (Chen et al., 2016; Ghaderi

et al., 2007). R160 stabilizes the C-terminal loop via interaction with the main chains

of D409, Q411, and E412 (Fig. 5B). This loop interacts with the residues involved in

the N-terminal dimerization interface. The R160Q mutant contains a shorter side

chain that would require loop reorganization were they to interact, which could

possibly affect epimerase oligomerization (Fig. 5B). However, the hyposialylation

effect of R160Q was less adverse than that of the I329T mutation. I329 participates in

forming an extensive network of hydrophobic interactions (Fig 5C). Importantly,

L249, located adjacent to I329, interacts directly with uridine diphosphate (UDP) (Fig.

5C). The I329T mutation, which introduces a smaller and less hydrophobic residue,

would weaken the Van der Waals interactions within the hydrophobic core, affecting

the structural integrity near the substrate-binding site. Meanwhile, the kinase domain

of GNE displays a typical bi-lobal architecture and contains a type I zinc-binding

motif characteristic of the repressor, open reading frame, kinase (ROK) family (Fig. 5A) (Martinez et al., 2012; Tong et al., 2009). Full-length GNE requires dimerization to permit ManNAc kinase activity (Martinez et al., 2012). I588 is located at the dimerization interface of the kinase C-terminal lobe (Fig. 5D); mutating this residue to less-hydrophobic threonine (Thr) affects kinase dimerization and thus its activity. Finally, V727 is buried in a hydrophobic core (Tong et al., 2009). The modeling of the V727M mutation suggests that the bulkier methionine (Met) may induce structural changes at this site to avoid steric clash (Fig. 5E). However, since this residue is positioned away from the catalytic site, the V727M mutation is expected to cause a less severe phenotype than the I329T mutation.

Structural analysis of GNE mutants shows that the I329T mutation, which can cause structural instability, is located near the active site. As a result, this mutation is expected to have the most adverse effect on GNE epimerase activity. Indeed, the I329T mutant shows the most pronounced hyposialylation (Figs. 4C and D).

Mutation-specific drug response of GNE myopathy disease models

To compensate for the hyposialylation caused by reduced GNE enzymatic activity in GNE myopathy patients, administration of ManNAc is being investigated in clinical trials for GNE patients (NCT02346461) following a safety profile study (Xu et al., 2017), despite the unsatisfactory outcome of a sialic acid clinical trial (Lochmuller et al., 2019). We therefore hypothesized that ManNAc would elicit a distinctive response in GNE patients with a mutation in the domain of ManNAc kinase because ManNAc is a direct substrate of this enzyme (Fig. 6A). To explore this possibility, we examined the recovery of hyposialylation of GNE mutants in the epimerase domain (R160Q and I329T) and kinase domain (I588T and V727M) after treatment with ManNAc. As predicted, the recovery of sialic acid production after

ManNAc treatment occurred in GNE mutants harboring mutations in the epimerase

domains (R160Q and I329T), while the effect in the kinase mutant (I588T and

V727M) was marginal (Figs. 6B and C). The dose-dependent response curve clearly

revealed the mutant-specific ManNAc response (Figs. 6D and E). We noticed that

sialic acid production was reduced in V727M, for unknown reasons (Fig. 6E). These

data suggest that the ManNAc response differs according to the mutation type of GNE.

GNE myopathy modeling in myocytes derived from GNE-hESCs

Prior to inducing the differentiation of GNE-hESCs into myocytes, we

examined whether hyposialylation would affect "pluripotency." To examine this

effect, I329T hESCs were used, as they showed the most severe hyposialylation.

Neither alkaline phosphatase activity, which is a well-characterized pluripotency

indicator, nor the mRNA expression of typical pluripotency markers were altered

(Figs. S5A and B). Additionally, teratomas from the I329T mutant developed

normally (Fig. S5C). Thus, we concluded that hyposialylation due to GNE mutation

had minimal effects on the pluripotency of hESCs. Next, to verify mutation-specific

responses in myocytes, WT and GNE-hESCs were differentiated into skeletal

myocytes (hereafter referred to as myocytes) through multiple steps (Fig. 7A). The

expression levels of typical markers for myocytes (e.g., *MYOG*, *MYH2*, and *MYH7*)

and pluripotency (e.g., *NANOG* and *SOX2*) were used to evaluate myocyte

differentiation (Fig. 7B). Differentiated myocytes showing typical elongated, rod-

shaped morphology, clearly expressed MYH3 (also known as Myosin heavy chain-

embryonic: MyHC-emb), a skeletal muscle specific contractile protein (Fig. 7C).

Notably, the mutation-specific hyposialylation observed in GNE-hESCs (Fig. 4D)

was also evident in GNE-myocytes relative to controls (Figs. 7D and E). Consistently,

exclusive effect of ManNAc on myocytes with epimerase mutation (R160Q and

I329T) was also reproduced (Figs. 7F and G).   While understanding of the

functionality of skeletal myocytes derived from hPSCs is limited, the functionality of

cardiomyocytes differentiated from hESCs (hESC-CMs) has been well-characterized

(Mummery et al., 2012). Thus, we examined whether GNE mutation affects the

functionality of cardiomyocytes, as cardiomyopathy has been occasionally reported in

GNE patients (Malicdan et al., 2014). hESC-CMs with higher Troponin T2, Cardiac

type (*TNNT2*) and lower *NANOG* expression than hESCs (Fig. S6A) showed clear

beating behavior (Movie S3). GNE mRNA expression was comparable regardless of

mutation status (Fig. 6B). Consistent mutation-specific hyposialylation and drug

responses were also observed in hESCs-CMs with different GNE mutations (Figs.

S6C). Consistently, mutation specific ManNAc response was reproduced in hESC-

CMs (Fig. S6D). The hyposialylation of hESC-CMs with the I329T mutation was

evident when compared with hESCs-CM with the I329I silent mutation (Fig. S6E).

The comparison of beating behavior of hESC-CM from WT or I329T hESCs revealed

no significant changes resulting from GNE mutation (Movies S4A and B).

Discussion

Base editors for disease modeling in hPSCs

Despite the limited target accessibility of BEs relative to HDR, the high

editing efficiency of BEs compared to HDR in hPSCs would be advantageous in

disease modeling using hPSCs. Through the development and implementation of NG-

BEs for gene editing, coverage of BEs for GNE myopathy modeling encompassed

73.5% of transition point mutations in the present study (Fig. 1). We demonstrated

that BEs were more efficient than HDR due to their lesser effect on p53 (Figs. 2F and

H). However, along with PAM-less target sequences, which account for 26.5% of

transition mutations in GNE myopathy, the production of undesirable mutations of

bystander bases with BEs, would be a significant drawback in disease modeling. Thus,

the development of "PAM-less BEs" and "BEs with narrowed editing windows"

would be advantageous for more efficient disease modeling with BE techniques.

Significance of isogenic disease model

Here, we established 14 GNE mutants from three hPSCs with different genetic

backgrounds. Hyposialylation, a typical pathological phenotype of GNE myopathy,

occurred in a mutation-specific manner regardless of hPSC cell type. However,

because the basal level of sialic acid in the wild-type of the four hPSCs (H9, hCHA3:

hESCs, BJ-iPSCs and SES8 (Lee et al., 2010): iPSCs) differed relative to the mutant

hPSCs (Fig. 4F), comparison of sialic acid between unmatched pairs could not allow

us to precisely determine the effect of mutation. Previously, various disease models

were established using iPSCs derived from patients, in which disease phenotype was

compared to genetically unmatched control iPSCs (Lee et al., 2009; Yamashita et al.,

2014). This study confirmed the previously proposed bias, introduced by the use of

cells with an unmatched genetic background to hinder precise comparisons of

pathological phenotypes [4].

Base editing technique for validating a variant of uncertain significance

Despite large numbers of single nucleotide variants (SNVs) in rare genetic

diseases, the functional consequences of mutations are often unpredictable, due to

insufficient *a priori* observations and incomplete penetrance (Findlay et al., 2018).

With isogenic disease models, one can classify these so-called "variants of uncertain

significance" (VUS) (Findlay et al., 2018) more accurately with the aid of functional

information. In the present study, for example, since I588T showed a similar level of

hyposialylation as R160Q, classified as "Pathogenic/Likely pathogenic," we inferred

that I588T may also be pathogenic. The I588T variant was originally annotated as

"uncertain significance" in the ClinVar database (Landrum et al., 2016). The efficient

induction of point mutation with BEs (or PE in the near future) would enhance further

studies seeking to validate the effect of VUS.

Consideration of mutation-specific drug response

We also demonstrated that epimerase mutants (R160Q and I329T) but not

kinase mutants (I588T and V727M) showed recovery of sialylation after ManNAc

treatment (Figs. 6D and 7G). Interestingly, while the I588T mutant showed no

recovery, the V727M mutation resulted in reduced sialylation following ManNAc

treatment, for unknown reasons. In light of these mutant-specific responses to

ManNAc, it would be advisable to use patient stratification based on mutation status

in clinical trials of ManNAc, to ensure accurate efficacy validation with ManNAc and

similar drugs.

Future prospects for drug development based on isogenic disease models

A recent study developed a high throughput (HT) phenotypic screening system to measure CFTR protein in cystic fibrosis with an isogenic pair of iPSCs (Merkert et al., 2019). Similarly, along with 3D protein modeling (Fig. 5), HT phenotypic screening with isogenic GNE myopathy disease models using the level of sialylation as a phenotypic indicator could accelerate drug development for GNE myopathy. Furthermore, by taking advantage of isogenic disease models and a drug-transcriptome database such as the Connectivity MAP (Kwon et al., 2019), candidate drugs from the FDA-approved drug library that reverse pathogenic gene signatures can easily be predicted (Kwon et al., 2020; Liu et al., 2015). The efficacy of these drug candidates can then be validated in the "GNE myopathy in a dish" model, making this strategy a strong potential option to develop drugs for GNE myopathy in the future.

Conclusion

The isogenic "GNE myopathy in a dish model," established in hPSCs by BEs with relatively high efficiency, revealed mutation-specific hyposialylation and drug responses. This work establishes important cellular tools and techniques for further mechanistic studies and development of drug screening systems.

Author contributions

HJ.C conceived the overall study design and led the experiments. JC.P and J.K mainly conducted the experiments, data analysis, and critical discussion of the results. HK.J and S.B contributed to sgRNA design and NGS analysis and provided gene editing techniques. SY.L, KT.K and S.P. produced mutant lines. HS.L and HJ.C performed 3D protein structure modeling. SJ.P and SH.M conducted the experiments with differentiated cardiomyocytes.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal

relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

This work was supported by a grant from the National Research Foundation of Korea

(2017M3A9B3061843 and 2020R1A2C2005914) and a grant from the Research of

Korea Centers for Disease Control and Prevention (2020-ER6902-00).

Materials and Methods

Live cell ratio after gene editing in WT and TP53 KO hESCs

To trace the number of live cells during gene editing, cells were documented by JuLI

stage for 36 hours after transfection. Cells were transfected with the same condition in

'Generation of *TP53* knock-out and *GNE* mutant hPSCs'. The number of cells at the

time point of 2 hours and 36 hours were counted by cell counter ImageJ plugin. To

calculate the live cell ratio, the number of cells at the time point of 36 hours were

divided by the number of cells at the time point of 2 hours.

Generation of *TP53* knock-out and *GNE* mutant hPSCs

For hPSCs transfection, cells were detached with Accutase^{TM} (561527, BD

Bioscience) and washed with DPBS for three times and resuspended to concentration

of 1 x 10^6 cell in 100 μl with Opti-MEM (31985070, Gibco). 2 μg of Cas9 or BE

vectors (Cas9, BE4max, NG-BE4max, ABEmax and NG-ABEmax cloned in pCMV

vector), 3 μg of sgRNA vector and 2 μg of ssODN (in case of HDR) was added to 100

μl of cell mixture. Electroporation was performed by NEPA-21 with 175 V of poring

pulse and 2.5 mV of transfer pulse. To enrich *TP53* KO cells, 10 μM of nutlin3 was

treated for 48 hours after 5 days from transfection.

Flow cytometry

Cells were detached with Accutase solution (561527, BD Bio-sciences) for hPSCs or

with TrypLE (12604, Gibco) for myocytes, followed by 3 times DPBS wash and then

analyzed through FACS Calibur I (BD Bioscience). For the determination of sialic

acid, cells were stained with Fluorescein SNA (FL-1301, Vector Laboratories).

CellQuest Pro soft-ware was used for FACS analysis.

Generation of skeletal muscle cells from hESCs

hESCs were differentiated to skeletal muscle cells by using commercially available

skeletal muscle differentiation medium kit (Amsbio). In brief, Cells were seeded onto

a collagen I (Sigma)-coated dish at 5,000 cells /cm2 dish and maintained for 6 days in

skeletal muscle induction medium (SKM01, Amsbio). At day 6, cells were

dissociated with 0.05% trypsin, seeded onto a collagen I-coated dish at 5,000 cells

/cm2 dish, and maintained for 6 days in myoblast cell culture medium (SKM02,

Amsbio). At day 12, when myoblasts are confluent, the medium was changed to

myotube cell culture medium (SKM03, Amsbio) and freshly changed every 3 days for

more than 6 days. For drug response test, cells were treated with ManNAc (Sigma) at

the indicated times and concentrations and the media with ManNAc was changed

every day.

Statistical analysis

The quantitative data are expressed as the mean values ± standard deviation (SD).

Student's unpaired t-tests was performed to analyze the statistical significance of each

response variable using the PRISM. p values less than 0.05 were considered

statistically significant (*, $p < 0.05$, **, $p < 0.01$, ***, $p < 0.001$ and ****, $p < 0.0001$)

Reference

Celeste, F.V., Vilboux, T., Ciccone, C., de Dios, J.K., Malicdan, M.C., Leoyklang, P., McKew, J.C., Gahl, W.A., Carrillo-Carrasco, N., and Huizing, M. (2014). Mutation update for GNE gene variants associated with GNE myopathy. Hum Mutat *35*, 915-926.

Chan, Y.M., Lee, P., Jungles, S., Morris, G., Cadaoas, J., Skrinar, A., Vellard, M., and Kakkis, E. (2017). Substantial deficiency of free sialic acid in muscles of patients with GNE myopathy and in a mouse model. PLoS One *12*, e0173261.

Chen, S.C., Huang, C.H., Lai, S.J., Yang, C.S., Hsiao, T.H., Lin, C.H., Fu, P.K., Ko, T.P., and Chen, Y. (2016). Mechanism and inhibition of human UDP-GlcNAc 2-epimerase, the key enzyme in sialic acid biosynthesis. Sci Rep *6*, 23274.

Findlay, G.M., Daza, R.M., Martin, B., Zhang, M.D., Leith, A.P., Gasperini, M., Janizek, J.D., Huang, X., Starita, L.M., and Shendure, J. (2018). Accurate classification of BRCA1 variants with saturation genome editing. Nature *562*, 217-222.

Fiscella, M., Zhang, H., Fan, S., Sakaguchi, K., Shen, S., Mercer, W.E., Vande Woude, G.F., O'Connor, P.M., and Appella, E. (1997). Wip1, a novel human protein phosphatase that is induced in response to ionizing radiation in a p53-dependent manner. Proc Natl Acad Sci U S A *94*, 6048-6053.

Gaudelli, N.M., Komor, A.C., Rees, H.A., Packer, M.S., Badran, A.H., Bryson, D.I., and Liu, D.R. (2017). Programmable base editing of A*T to G*C in genomic DNA without DNA cleavage. Nature *551*, 464-471.

Ghaderi, D., Strauss, H.M., Reinke, S., Cirak, S., Reutter, W., Lucka, L., and Hinderlich, S. (2007). Evidence for dynamic interplay of different oligomeric states of UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase by biophysical methods. J Mol Biol *369*, 746-758.

Huang, T.P., Zhao, K.T., Miller, S.M., Gaudelli, N.M., Oakes, B.L., Fellmann, C., Savage, D.F., and Liu, D.R. (2019). Circularly permuted and PAM-modified Cas9 variants broaden the targeting scope of base editors. Nat Biotechnol *37*, 626-631.

Ihry, R.J., Worringer, K.A., Salick, M.R., Frias, E., Ho, D., Theriault, K., Kommineni, S., Chen, J., Sondey, M., Ye, C., *et al.* (2018). p53 inhibits CRISPR-Cas9 engineering in human pluripotent stem cells. Nat Med *24*, 939-946.

Izmiryan, A., Ganier, C., Bovolenta, M., Schmitt, A., Mavilio, F., and Hovnanian, A. (2018). Ex Vivo COL7A1 Correction for Recessive Dystrophic Epidermolysis Bullosa Using CRISPR/Cas9 and Homology-Directed Repair. Mol Ther Nucleic Acids *12*, 554-567.

Jackow, J., Guo, Z., Hansen, C., Abaci, H.E., Doucet, Y.S., Shin, J.U., Hayashi, R., DeLorenzo, D., Kabata, Y., Shinkuma, S., *et al.* (2019). CRISPR/Cas9-based targeted genome editing for correction of recessive dystrophic epidermolysis bullosa using iPS cells. Proc Natl Acad Sci U S A.

Jeong, Y.K., Yu, J., and Bae, S. (2019). Construction of non-canonical PAM-targeting adenosine base editors by restriction enzyme-free DNA cloning using CRISPR-Cas9. Sci Rep *9*, 4939.

Kim, K.T., Park, J.C., Jang, H.K., Lee, H., Park, S., Kim, J., Kwon, O.S., Go, Y.H., Jin, Y., Kim, W., *et al.* (2020). Safe scarless cassette-free selection of genome-edited human pluripotent stem cells using temporary drug resistance. Biomaterials *262*, 120295.

Koblan, L.W., Doman, J.L., Wilson, C., Levy, J.M., Tay, T., Newby, G.A., Maianti, J.P., Raguram, A., and Liu, D.R. (2018). Improving cytidine and adenine base editors

by expression optimization and ancestral reconstruction. Nat Biotechnol *36*, 843-846.

Komor, A.C., Kim, Y.B., Packer, M.S., Zuris, J.A., and Liu, D.R. (2016). Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage. Nature *533*, 420-424.

Kwon, O.S., Kim, W., Cha, H.J., and Lee, H. (2019). In silico drug repositioning: from large-scale transcriptome data to therapeutics. Arch Pharm Res *42*, 879-889.

Kwon, O.S., Lee, H., Kong, H.J., Kwon, E.J., Park, J.E., Lee, W., Kang, S., Kim, M., Kim, W., and Cha, H.J. (2020). Connectivity map-based drug repositioning of bortezomib to reverse the metastatic effect of GALNT14 in lung cancer. Oncogene.

Landrum, M.J., Lee, J.M., Benson, M., Brown, G., Chao, C., Chitipiralla, S., Gu, B., Hart, J., Hoffman, D., Hoover, J.*, et al.* (2016). ClinVar: public archive of interpretations of clinically relevant variants. Nucleic Acids Res *44*, D862-868.

Lee, G., Papapetrou, E.P., Kim, H., Chambers, S.M., Tomishima, M.J., Fasano, C.A., Ganat, Y.M., Menon, J., Shimizu, F., Viale, A.*, et al.* (2009). Modelling pathogenesis and treatment of familial dysautonomia using patient-specific iPSCs. Nature *461*, 402-406.

Lee, T.H., Song, S.H., Kim, K.L., Yi, J.Y., Shin, G.H., Kim, J.Y., Kim, J., Han, Y.M., Lee, S.H., Lee, S.H.*, et al.* (2010). Functional recapitulation of smooth muscle cells via induced pluripotent stem cells from human aortic smooth muscle cells. Circ Res *106*, 120-128.

Li, H.L., Gee, P., Ishida, K., and Hotta, A. (2016). Efficient genomic correction methods in human iPS cells using CRISPR-Cas9 system. Methods *101*, 27-35.

Liu, J., Lee, J., Salazar Hernandez, M.A., Mazitschek, R., and Ozcan, U. (2015). Treatment of obesity with celastrol. Cell *161*, 999-1011.

Lochmuller, H., Behin, A., Caraco, Y., Lau, H., Mirabella, M., Tournev, I., Tarnopolsky, M., Pogoryelova, O., Woods, C., Lai, A.*, et al.* (2019). A phase 3 randomized study evaluating sialic acid extended-release for GNE myopathy. Neurology *92*, e2109-e2117.

Mali, P., Yang, L., Esvelt, K.M., Aach, J., Guell, M., DiCarlo, J.E., Norville, J.E., and Church, G.M. (2013). RNA-guided human genome engineering via Cas9. Science *339*, 823-826.

Malicdan, M.C., Mian, L., Leoyklang, P., Celeste, F., Despres, D., Zerfas, P., Carrillo-Carrasco, N., and Gahl, W.A. (2014). GP 52. Neuromuscular Disorders *9*, 810 %@ 0960-8966.

Martinez, J., Nguyen, L.D., Hinderlich, S., Zimmer, R., Tauberger, E., Reutter, W., Saenger, W., Fan, H., and Moniot, S. (2012). Crystal structures of N-acetylmannosamine kinase provide insights into enzyme activity and inhibition. J Biol Chem *287*, 13656-13665.

Maurissen, T.L., and Woltjen, K. (2020). Synergistic gene editing in human iPS cells via cell cycle and DNA repair modulation. Nat Commun *11*, 2876.

Merkert, S., Schubert, M., Olmer, R., Engels, L., Radetzki, S., Veltman, M., Scholte, B.J., Zollner, J., Pedemonte, N., Galietta, L.J.V.*, et al.* (2019). High-Throughput Screening for Modulators of CFTR Activity Based on Genetically Engineered Cystic Fibrosis Disease-Specific iPSCs. Stem Cell Reports *12*, 1389-1403.

Merkle, Florian T., and Eggan, K. (2013). Modeling Human Disease with Pluripotent Stem Cells: from Genome Association to Function. Cell Stem Cell *12*, 656-668.

Mummery, C.L., Zhang, J., Ng, E.S., Elliott, D.A., Elefanty, A.G., and Kamp, T.J. (2012). Differentiation of human embryonic stem cells and induced pluripotent stem cells to cardiomyocytes: a methods overview. Circ Res *111*, 344-358.

Musunuru, K. (2013). Genome editing of human pluripotent stem cells to generate human cellular disease models. Disease Models & Mechanisms *6*, 896-904.

Nishimasu, H., Shi, X., Ishiguro, S., Gao, L., Hirano, S., Okazaki, S., Noda, T., Abudayyeh, O.O., Gootenberg, J.S., Mori, H., *et al.* (2018). Engineered CRISPR-Cas9 nuclease with expanded targeting space. Science *361*, 1259-1262.

Nishino, I., Carrillo-Carrasco, N., and Argov, Z. (2015). GNE myopathy: current update and future therapy. J Neurol Neurosurg Psychiatry *86*, 385-392.

Noguchi, S., Keira, Y., Murayama, K., Ogawa, M., Fujita, M., Kawahara, G., Oya, Y., Imazawa, M., Goto, Y., Hayashi, Y.K., *et al.* (2004). Reduction of UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase activity and sialylation in distal myopathy with rimmed vacuoles. J Biol Chem *279*, 11402-11407.

Osborn, M.J., Newby, G.A., McElroy, A.N., Knipping, F., Nielsen, S.C., Riddle, M.J., Xia, L., Chen, W., Eide, C.R., Webber, B.R., *et al.* (2020). Base Editor Correction of COL7A1 in Recessive Dystrophic Epidermolysis Bullosa Patient-Derived Fibroblasts and iPSCs. J Invest Dermatol *140*, 338-347 e335.

Perlman, R.L. (2016). Mouse models of human disease: An evolutionary perspective. Evol Med Public Health *2016*, 170-176.

Pham, N.D., Pang, P.C., Krishnamurthy, S., Wands, A.M., Grassi, P., Dell, A., Haslam, S.M., and Kohler, J.J. (2017). Effects of altered sialic acid biosynthesis on N-linked glycan branching and cell surface interactions. J Biol Chem *292*, 9637-9651.

Rees, H.A., and Liu, D.R. (2018). Base editing: precision chemistry on the genome and transcriptome of living cells. Nat Rev Genet *19*, 770-788.

Rowe, R.G., and Daley, G.Q. (2019). Induced pluripotent stem cells in disease modelling and drug discovery. Nat Rev Genet *20*, 377-388.

Shi, Y., Inoue, H., Wu, J.C., and Yamanaka, S. (2017). Induced pluripotent stem cell technology: a decade of progress. Nat Rev Drug Discov *16*, 115-130.

Tiscornia, G., Vivas, E.L., and Izpisua Belmonte, J.C. (2011). Diseases in a dish: modeling human genetic disorders using induced pluripotent cells. Nat Med *17*, 1570-1576.

Tong, Y., Tempel, W., Nedyalkova, L., Mackenzie, F., and Park, H.W. (2009). Crystal structure of the N-acetylmannosamine kinase domain of GNE. PLoS One *4*, e7165.

Weissbein, U., Benvenisty, N., and Ben-David, U. (2014). Quality control: Genome maintenance in pluripotent stem cells. J Cell Biol *204*, 153-163.

Xu, X., Wang, A.Q., Latham, L.L., Celeste, F., Ciccone, C., Malicdan, M.C., Goldspiel, B., Terse, P., Cradock, J., Yang, N., *et al.* (2017). Safety, pharmacokinetics and sialic acid production after oral administration of N-acetylmannosamine (ManNAc) to subjects with GNE myopathy. Mol Genet Metab *122*, 126-134.

Yamashita, A., Morioka, M., Kishi, H., Kimura, T., Yahara, Y., Okada, M., Fujita, K., Sawai, H., Ikegawa, S., and Tsumaki, N. (2014). Statin treatment rescues FGFR3 skeletal dysplasia phenotypes. Nature *513*, 507-511.

1   Yang, L., Guell, M., Byrne, S., Yang, J.L., De Los Angeles, A., Mali, P., Aach, J., Kim-
2   Kiselak, C., Briggs, A.W., Rios, X., *et al.* (2013). Optimization of scarless human
3   stem cell genome editing. Nucleic Acids Res *41*, 9049-9061.
4

Figure legends

Figure 1. Target-ability of point mutations causing GNE myopathy by base

editors (A) Graphical description of three different gene editing technologies for

generating point mutations (APOBEC1: cytidine deaminase, UGI: Uracil-DNA

glycosylase inhibitor, TadA: adenine deaminase, TadA*: evolved TadA) (B) Pie

Chart of distribution of GNE myopathy associated mutations described as pathogenic

or likely pathogenic mutations of 63 patients in ClinVar database

(https://www.ncbi.nlm.nih.gov/clinvar/) (C and D) Pie chart of targetability of

mutations for 'wild-type (WT) to mutant' (C) or 'mutant to wild-type (WT)' (D)

modeling by base editors (left), graphical presentation of target-ability of each base

editor (right)

Figure 2. Efficiency of base editing compared to HDR in hESCs (A) Graphical

description of site of mutations in the epimerase and kinase domain of human GNE

protein (hGNE2) (left), List of mutations generated by each base editor (right) (B to E)

Target deep sequencing results showing overall base conversion rate of each sequence

by CBE (C to T) and ABE (A to G) (left), Graphical presentation of on-target editing

efficiency of indicated gene editing tools (right) for R160Q (B), I588T (C), V727M

(D) and I329T (E), Target base and PAM sequence is colored in red and green (3'-5')

respectively. (F) mRNA expression of *PPM1D* in wild-type (WT) and *TP53*

Knockout (TP53 KO) hESCs transfected with sgRNA targeting CCR5 (sgCCR5) and

indicated gene editing tools (Cas9, CBE, NG-CBE, ABE and NG-ABE) compared to

control (Cont: sgCCR5 only), *, $p < 0.05$ (G) Graphical presentation of percentage of

live cell ratio of wild-type (WT) and TP53 KO hESCs after transfection of sgCCR5

and indicated gene editing tools, ****, $p < 0.0001$ (H) Gene editing efficiency of

indicated gene editing tools in WT and TP53 KO hESCs on CCR5. Editing efficiencies are analyzed by target deep sequencing.

Figure 3. Establishment of GNE myopathy disease models through base editors
(A to D) Sanger validation of mutants of R160Q (A), I329T (B), I588T (C) and V727M (D) compared to wild-type (WT), Exons coding UDP-GlcNAc-2-epimerase and ManNAc kinase domain are colored in blue and in red respectively. Target amino acids and bases are colored in red.

Figure 4. Mutation-specific phenotype of GNE myopathy disease models (A) Microscope bright field (BF) images (top) and fluorescent images for OCT4 (bottom) of hESCs of WT and GNE mutants, DAPI for nuclear counterstaining (B) Fluorescent images of hESCs of WT and GNE mutants stained with SNA (green) and DAPI for counter nuclear staining (C) Histogram and (D) bar graph of mean fluorescence intensity (MFI) of SNA in WT and GNE mutants, ***, $p<0.001$, ****, $p<0.0001$ (E) List of predicted severity, clinical significance and mean fluorescence intensity of GNE mutants, predicted severe, medium and mild were colored in red, yellow and green respectively. (F) Mean fluorescence intensity of SNA of indicated hPSCs lines, ***, $p < 0.001$ (G) mRNA expression of *GNE* in indicated hPSCs lines

Figure 5. Structural analysis predicts mutation-specific hyposialylation (A) Structures of tetrameric epimerase and dimeric kinase domains of human GNE (PDB ID 4ZHT and 2YHW, respectively) are shown. Target mutations R160Q (B) and I329T (C) are found in the epimerase domain, while I588T (D) and V727M (E) are in the kinase domain. Mutagenesis and rotamer selection were performed using PyMOL, and mutated residues are indicated in grey. Inter-residue interactions within a distance of 4.1 Å are indicated with dashed lines.

Figure 6. Mutation-specific drug response of GNE myopathy disease models (A) Graphical description of biosynthesis of sialic acid and sialyation and roles of GNE protein as UDP-GlcNAc-2-epimerase and ManNAc kinase (B) Fluorescence microscopic images of GNE mutant hESCs stained with SNA after indicated dose of ManNAc treatment for day 2 and day 6 (C) Flow cytometry for SNA in GNE mutant hESCs 6 days after indicated dose of ManNAc treatment (D and E) Change of mean fluorescent intensity of SNA at 2 (D) and 6 (E) days after indicated concentration of ManNAc treatment

Figure 7. GNE myopathy modeling in myocytes derived from GNE-hESCs (A) Schematic diagram of myogenic differentiation from hESCs (B) mRNA expression of typical markers for skeletal myocytes (top) and pluripotency (bottom) in hESCs and myocytes derived from hESCs of WT and GNE mutants (C) Fluorescence microscope images for MYH3 in myocytes derived from WT and GNE mutant hESCs, DAPI for nuclear counterstaining (D) Fluorescent images of hESCs derived myocytes of WT and GNE mutants stained with SNA (green) and DAPI for counter nuclear staining (E) bar graph of mean fluorescence intensity (MFI) of SNA in WT and GNE mutants, **, $p<0.01$, ***, $p<0.001$, ****, $p<0.0001$ (F) Fluorescence microscopic images of GNE mutant hESCs derived myocytes stained with SNA after indicated dose of ManNAc treatment (G) Change of mean fluorescent intensity of SNA after indicated concentration of ManNAc treatment

Graphical abstract

Figure. 1

Figure. 2

Figure. 2

Figure. 3

Figure. 4

Figure. 4

E

Mutation (hGNE2)	Predicted severity					Clinical significance	Mean fluorescent intensity
	PolyPhen2	SIFT	Align	Pmut	Overall		
R160Q	Probably Damaging	Tolerated	35	Pathological	Medium	Pathogenic / Likely pathogenic	643.5
I329T	Probably Damaging	Damaging	65	Pathological	Severe	Likely pathogenic	374
I588T	Possibly Damaging	Damaging	65	Neutral	Medium	Uncertain significance	668.3
V727M	Probably Damaging	Damaging	15	Neutral	Medium	Conflicting (Pathogenic / Likely pathogenic / Likely benign / Uncertain significance)	735.3

Severe   Medium   Mild

Figure. 5

Figure. 6

Figure. 6

Figure. 7

Figure. 7

F

(µM) ManNAc

| | 0 | 250 | 1000 |

R160Q
I329T
I588T
V727M

G

R160Q · I329T
I588T · V727M

ΔMean Fluorescence

ManNAc concentration (µM)