1

A Non-Genetic, Cell Cycle Dependent Mechanism of Platinum

Resistance in Lung Adenocarcinoma

Alvaro Gonzalez-Rajal[1,2#], Kamila A Marzec[1], Rachael McCloy[3], Max Nobis[3,5], Venessa Chin[3,4,5], Jordan F. Hastings[3], Kaitao Lai[1,6], Marina Kennerson[1,6], William E Hughes[2,5,7], Vijesh Vaghjiani[8], Paul Timpson[3,5], Jason E. Cain[8,9], D. Neil Watkins[10,11], David R. Croucher[3,5,#,§] and Andrew Burgess[1,6,#,§]

[1] *ANZAC Research Institute, Concord Hospital, Concord, NSW 2139, Australia*
[2] *Garvan Institute of Medical Research, Sydney, NSW 2010, Australia.*
[3] *The Kinghorn Cancer Centre, Garvan Institute of Medical Research, Sydney, NSW 2010, Australia.*
[4] *St Vincent's Hospital Sydney, Darlinghurst, NSW 2010, Australia*
[5] *St Vincent's Hospital Clinical School, University of New South Wales, Sydney, NSW 2052, Australia.*
[6] *The University of Sydney Concord Clinical School, Faculty of Medicine and Health, Sydney, NSW 2139, Australia*
[7] *Children's Medical Research Institute, The University of Sydney, Westmead, Australia*
[8] *Hudson Institute of Medical Research, Clayton, Victoria 3168, Australia.*
[9] *Department of Molecular and Translational Medicine, School of Medicine, Nursing and Health Sciences, Monash University, Clayton, Victoria 3800, Australia*
[10] *Research Institute in Oncology and Hematology, CancerCare Manitoba, Winnipeg, MB, Canada, R3E-0V9*
[11] *Department of Internal Medicine, Rady Faculty of Health Science, University of Manitoba, Winnipeg, Canada, R3E 0W2*

[§] *These authors contributed equally to this work*

Corresponding Authors:

Dr Alvaro Gonzalez-Rajal
Email: a.rajal@garvan.org.au

A/Prof David Croucher
Email: d.croucher@garvan.org.au

A/Prof Andrew Burgess
Email: andrew.burgess@sydney.edu.au

Running Title: Cell cycle dependent platinum resistance.

Keywords: 53BP1, PARP1, BRCA1, HR, ATR, PCNA, mitosis, DSB, NHEJ

Abstract:

We previously used a pulse-based *in vitro* assay to unveil targetable signalling pathways associated with innate cisplatin resistance in lung adenocarcinoma (Hastings et al., 2020). Here we advanced this model system and identified a non-genetic mechanism of resistance that drives recovery and regrowth in a subset of cells. Using RNAseq and a suite of biosensors to track single cell fates both *in vitro* and *in vivo,* we identified that early S phase cells have a greater ability to maintain proliferative capacity, which correlated with reduced DNA damage over multiple generations. In contrast, cells in G1, late S or those treated with PARP/RAD51 inhibitors, maintained higher levels of DNA damage and underwent prolonged S/G2 phase arrest and senescence. Combined with our previous work, these data indicate that there is a non-genetic mechanism of resistance in lung adenocarcinoma that is dependent on the cell cycle stage at the time of cisplatin exposure.

Introduction:

Lung adenocarcinoma (LUAD) is the most common form of lung cancer and the leading cause of cancer-related death in Australia. Less than 15% of patients have a targetable driver mutation and therefore cannot benefit from targeted therapy (Herbst et al., 2018). Consequently, the overwhelming majority of LUAD patients receive platinum-based chemotherapy as standard of care. The anti-tumour abilities of platinum compounds were first identified over 50 years ago with the discovery of cisplatin (Kelland, 2007). Since then, cisplatin and its derivatives have become one of the most successful groups of chemotherapeutics ever developed. Platinum therapy is essentially curative in testicular cancer, with survival rates >90%, and is also a frontline treatment for small-cell lung cancer, ovarian, head and neck, bladder, and cervical cancers (Gonzalez-Rajal et al., 2020; Kelland, 2007). Unfortunately, response rates to platinum in LUAD are below 30%, due primarily to innate resistance (Herbst et al., 2018). Nearly 150 different mechanisms of platinum resistance have been identified to date (Stewart, 2007). The vast majority of these mechanisms have been derived from preclinical models that utilise continuous, high dose exposure models, well above what is physiologically achievable in patients. Unsurprisingly, the majority of these models have failed to translate into improved clinical outcomes. To overcome this, we recently demonstrated that analysis of an *in*

vitro assay that accurately models the *in vivo* drug exposure kinetics for cisplatin,
could provide therapeutically relevant insights into the signalling dynamics
associated with innate resistance (Hastings et al., 2020). Cisplatin is given to
patients as a single bolus dose, reaching a peak plasma concentration of ~14 µM (5
µg/ml), which is then rapidly cleared by the kidneys within 2-4 h (Andersson et al.,
1996; Urien and Lokiec, 2004). We therefore mimicked this *in vitro* by pulsing cells
for 2 h with the maximum plasma concentration (Hastings et al., 2020).

Once inside cells, platinum compounds can bind to DNA, RNA, and proteins
(Gonzalez-Rajal et al., 2020), however the binding to DNA, which forms platinum-
DNA adducts, are thought to be the primary mechanism for their tumour-specific
killing. Intra-strand DNA-platinum adducts are repaired by base-excision and
nucleotide excision repair during G1 (Slyskova et al., 2018). Inter-strand crosslinks
(ICL) are removed largely by the Fanconi anemia (FA) pathway (Smogorzewska,
2019), which generates single and double strand breaks that are resolved by either
the high-fidelity homologous recombination (HR) pathway during S-phase (Karanam
et al., 2012) or by the error-prone non-homologous end joining (NHEJ) pathway
during G1 and G2 phase (Enoiu et al., 2012; Slyskova et al., 2018). Consequently,
targeting DNA repair pathways has become a major focus for enhancing platinum
chemotherapies. For example, cells with defective HR repair have been shown to be
highly sensitive to combination therapy with cisplatin and PARP inhibitors in a
number of cancer types, including ovarian and breast (Tutt et al., 2018). However,
correlation between cisplatin sensitivity and impaired DNA repair has often failed to
translate clinically in LUAD (Mamdani and Jalal, 2016). In contrast, we have recently
identified TGF-β (Marini et al., 2018) and P70S6K (Hastings et al., 2020) as key
mediators of innate platinum resistance in LUAD. We now build upon these previous
results and identify in this research advance that a sub-population of cells are
capable of continued proliferation despite exposure to pulsed cisplatin. Using a
combination of cell cycle, DNA damage and replication biosensors together with real-
time single-cell fate tracking, we identified that these proliferative cells were enriched
in late G1/early S phase at the time of cisplatin exposure and were able to
sufficiently repair their DNA over multiple generations and rounds of replication.
These results increase our understanding of the complexities underlying non-genetic
resistance and recovery mechanisms in LUAD, while also highlighting mechanistic

issues with a number of current clinical trials focused on combination therapy with
cisplatin.

Results

Cells remain equally sensitive upon re-exposure to pulsed cisplatin

In our previous work (Hastings et al., 2020), we identified several targetable
signalling pathways that were associated with resistance to cisplatin in lung
adenocarcinoma cells. In this work, our goal was to analyse the innate mechanisms
that enable cell survival after the initial exposure to cisplatin. To assess this, we
analysed how cells respond to a subsequent dose of cisplatin, following recovery
from an initial exposure event. To do this, we pulsed cells with cisplatin (5 µg/ml) and
followed their response by time-lapse imaging. Cells were allowed to recover for 21-
42 days (depending on their base rate of proliferation), before being challenged
again with cisplatin, which equates to the approximate time patients normally receive
a second dose in the clinic (Figure 1A). We utilised 3 LUAD cell lines: A549 (wild-
type p53), NCI-H1573 (p53^{R248L} mutant) and NCI-H1299 (p53 null), which were all
engineered to stably express Histone H2B fused to mCherry, allowing real-time
quantitation of cell number and nuclear size. The initial pulse of cisplatin blocked the
proliferation of A549 and NCI-H1573 cells, and significantly reduced NCI-H1299 cell
numbers over a 3-day period. This was mirrored in colony formation assays, with
both A549 and NCI-H1573 showing strong suppression of colony outgrowth, while
p53-null H1299 cells were impacted to a lesser degree (Figure 1 - Supplement
Figure 1A). Interestingly, there was a less noticeable effect on cell confluence
(Figure 1B). Subsequent visual and quantitative analysis of cells by
immunofluorescence revealed a corresponding 2 to 6-fold increase in total cell and
nuclear area across all three cell lines (Figure 1C,D), accounting for the reduced
impact on confluence. Over the following 7 to 21 days of recovery (42 days for NCI-
H1573 due to slower rate of proliferation), cells eventually recovered to their pre-
pulse size and normal cell cycle distribution (Figure 1C,D and Figure 1 - Supplement
1B). In all three cell lines, subsequent cisplatin pulse of recovered cultures resulted
in a near identical response to the initial pulse, both in terms of inhibition of cell
number, reduced confluence and increased cell size (Figure 1B-D). Based on these
results, and our previous observation that all cells contained significantly increased

levels of cisplatin-DNA adducts (Hastings et al., 2020), we concluded that cells
surviving the first exposure remained equally sensitive to cisplatin and were
therefore unlikely to have acquired resistance or arisen from an intrinsically resistant
sub-clonal population within each cell line.

To assess this, we analysed the variability of cell and nuclear size after the initial
pulse of cisplatin at 3-7 days post exposure. Visual analysis identified several
colonies of cells whose size was similar to that of untreated control cells (Figure 2A-
B). We hypothesised that these cells were able to proliferate and outgrow the non-
proliferative (arrested) cells over the 21-to-42-day period. In support, a significant
increase in senescence associated Beta-galactosidase (β-gal) staining was
observed both visually and by flow cytometry (C12FDG) in A549 and to a lesser
extent H1573 and H1299 cell lines (Figure 2 - Supplement 1A-B). Increased levels of
the cyclin dependent kinase inhibitor p16, which is commonly linked with
senescence, was associated with larger cells in NCI-H1573 and H1299 cells, while in
A549 cells, which are null for p16 (Kawabe et al., 2000), p21 was similarly increased
(Figure 2 - Supplement 1C). These data suggest that in each cell line the larger cells
have a lower proliferative capacity and are likely senescent.

To test whether the surviving population arose from a subset of proliferating cells, we
utilised the LeGo RGB colour-guided clonal cell tracking system (Weber et al., 2011).
Briefly, each cell line was co-transfected with 3 different lentiviral vectors containing
either a red, green or blue fluorescent protein. Each cell randomly received a
variable amount of each plasmid resulting in a unique colour code for each cell.
Quantitative colour analysis of untreated control cells revealed that up to 64 unique
colours could be detected in A549 and NCI-H1299 cells and up to 46 colours in NCI-
H1573 cells (Figure 2 Supplement 1D). After pulsed exposure to cisplatin, single
colour colony outgrowths were clearly visible in all 3 cell lines at 3-7 days post
exposure, which was maintained at 21 for A549 and NCI-H1299 or 42-days for NCI-
H1573 cells (Figure 2C-D). Colour diversity in cisplatin recovered cells correlated
with the colony formation assay data (Figure 1, Supplement 1A), with NCI-H1299
(p53 null) cells displaying a greater variability in recovered clonal colours compared
to A549 (p53 wt) and NCI-H1573 (p53 mutant) cells (Figure 2C and Supplement 1D).
Taken together these results indicate that repopulation of the culture after the initial

pulse exposure is primarily driven by a small fraction of cells, with the total number of
clones impacted by p53 status.

To confirm these results *in vivo*, we injected A549, NCI-H1573 or NCI-H1299 cells
subcutaneously into the flanks of nude mice and allowed establishment of tumours
(150 mm^3) before administering a single treatment of carboplatin (60 mg/kg). Mice
harvested at 3 days post treatment were analysed by IHC for cell size and PCNA
positive staining, with all cell lines showing a significant increase in cell size (Figure
3A, B). Similar to the *in vitro* results, active proliferation (PCNA positivity) in A549
cells was strongly suppressed after carboplatin exposure, indicating that the majority
of cells were not proliferating. In p53 mutant NCI-1573, there was no significant
reduction in PCNA *in vivo* despite significant reduction in cell numbers *in vitro* cells.
In contrast, NCI-H1299 cells, which lack p53, did not show any significant decrease
in PCNA staining *in vivo* (Figure 3A,B). BrdU pulse labelling of *in vitro* treated cells,
showed similar results, with strong suppression of active BrdU incorporation in A549
cells, with partial and no significant inhibition seen in NCI-H1573 and NCI-H1299
cells, respectively (Figure 3 – Supplement 1). In summary, these *in vitro* and *in vivo*
data suggest that in LUAD cells, exposure to pulsed cisplatin results in a significant
proportion of cells increasing in size, with presence of functional p53 correlating with
increased senescence and reduced active replication in enlarged cells. Interestingly,
in all three lines, a sub-population of cells remain at a normal (stable) cell size,
maintained their proliferative capacity, and drove repopulation of the cell culture.
Despite this, these proliferative cells remained equally sensitive to subsequent
cisplatin treatment, indicating a non-genetic mechanism of resistance.

Differential RNAseq analysis of cisplatin treated sub-populations
To better understand the potential mechanism driving the difference between cells
that become enlarged and senescent compared to those that maintain stable size
and proliferative capacity, we performed RNAseq analysis on each unique
population. Briefly, A549 cells were pulsed with cisplatin, harvested at 72 h and
sorted into stable or enlarged cells based on size. This was determined by forward
and side scatter parameters (FSC, SSC), with gates established based on the size of
untreated control cells (Figure 4, Supplement 1A). Pre- and post-sorted cells were
then processed for RNAseq analysis (Figure 4A, B). Two-way hierarchical clustering

indicated that there were clear differences in gene expression between cisplatin treated cells that maintained a stable size compared to untreated control and enlarged cisplatin treated cells (Figure 4C). To better understand these effects, we undertook more detailed bioinformatic analysis using Ingenuity Pathway Analysis (IPA). Strong upregulation of the CDK inhibitor p21, was present in both pre-sorted cisplatin treated and post-sorted enlarged cells, matching the early flow data (Figure 2 Supplement 1C, Supplement Table 1,2). This corresponded with upregulation of p53, CHK and G2/M cell cycle checkpoint signalling, and a corresponding reduction in DNA replication and increase in senescence pathways (Figure 4D, E), correlating with the increased β-gal and reduced proliferation observed above (Figure 2 Supplement 1B and Figure 1B). Importantly, stable (size) cisplatin treated cells were significantly different from untreated controls, indicating that these cells were impacted by cisplatin exposure, similar to our previous reports where all cells contained detectible cisplatin-DNA adducts post pulsed exposure (Hastings et al., 2020). Notably, these stable cells displayed a decrease in EIF2, mTOR and p70S6K signalling, with a corresponding increase in TREM1, GP6 and IL-17F signalling pathways (Figure 4F, Supplement Table 3,4). Four-way comparative analysis further highlighted key differences between each sub-population, with stable cells all showing strong upregulation of BRCA1, ATM and DNA replication signalling pathways, which were heavily suppressed in non-proliferative cells (Figure 4G). Similarly, Gene Set Enrichment Analysis (GSEA), identified strong enrichment for cell cycle, HR directed repair, ATR and the Fanconi pathway in stable compared to enlarged cells (Figure 4 Supplement 1B,C). In summary, these data suggest that the cisplatin treated stable cells are considerably different from enlarged and untreated control cells. Specifically, stable cells do not undergo significant p53/p21 dependent cell cycle checkpoint arrest but do show prominent upregulation of DNA repair pathways involving HR/BRCA1 and ATM/ATR.

Cell cycle and p53 status at time of exposure correlates with cell fate outcomes

The above RNAseq data indicated that there were strong cell cycle dependent differences between proliferative and non-proliferative cisplatin treated cells. To better understand these differences, we utilised the FUCCI biosensor system to enable real-time cell cycle status of individual cells, as published in the previous

paper (Hastings et al., 2020). Briefly, asynchronous A549 cells stably expressing FUCCI, were pulsed with or without cisplatin and then followed by time-lapse microscopy for 72 h. Individual cells were manually tracked and scored for cell cycle status and cell fate as previously described (Caldon and Burgess, 2019; Hastings et al., 2020). The majority of control cells divided at least 2 times within the 72 h time period (Figure 5A). In contrast and as demonstrated previously (Hastings et al., 2020), cisplatin treated cells showed a range of cell cycle perturbations (Figure 5B and Figure 5 Supplement 1A), including a significant and prolonged S/G2 phase arrest (Figure 5 Supplement 1B), which correlated with a reduced number of total divisions (Figure 5 Supplement 1C). Combining this with additional scoring of cisplatin treated cells that underwent multiple (2 or more) divisions within the 72 h period, revealed an enrichment for cells that were in late G1 and early S-phase at the time of cisplatin exposure (Figure 5B). Unbiased analysis of an additional 400 cisplatin treated A549 cells, found a significant increase in the number of divisions arising from cells that were in G1/S or early S phase at the time of exposure compared to G1 phase cells (Figure 5C). Furthermore, the overwhelming majority of cells in late S or G2/M, only completed 1 division during the 72 h period. Taken together, these data suggest that there is a cell cycle dependent mechanism driving the ability of A549 cells to continue to proliferate after cisplatin exposure.

Our above data indicated that loss of p53 increased the number of clones capable of regrowth after cisplatin exposure. To assess this in greater detail we analysed asynchronous FUCCI expressing NCI-H1299 (p53 null) cells pulsed with or without (Control) cisplatin. Treatment with cisplatin significantly reduced the number of overall divisions in H1299 cells (Figure 5 Supplement 2A-C), however compared to A549 cells, this reduction was not as pronounced, in line with the cell proliferation data observed above (Figure 1B). Single cell fate tracking of NCI-H1299 cells suggested a small bias for higher rates of division in cells that were in late G1 and early S phase at the time of treatment (Figure 5, Supplement 2B-C), however this trend was not significant (Figure 5 Supplement 2D). We further assessed p53 loss by reanalysing our previously published data on siRNA knockdown of p53 in A549 cells (Hastings et al., 2020). Notably, depletion of p53 in A549 cells resulted in a corresponding increase in the number of divisions completed after pulsed exposure to cisplatin, supporting the NCI-H1299 data (Figure 5 Supplement 2E).

To validate these results *in vivo*, we implanted FUCCI expressing A549, NCI-H1573 or NCI-H1299 cells under optical windows in mice (Figure 5D). Tumours were allowed to establish before mice were given a single dose of carboplatin. Individual mice were then repeatedly imaged over 7 days post treatment. Notably, prior to cisplatin treatment, approximately 70 to 80% of cells from each line were in G1 phase (Figure 5E-F, Figure 5 Supplement 3A-B). Similar to *in vitro* results, we observed an increase in proportion of S/G2 phase cells at day 1 in all cell lines, indicating that cells were arrested in S/G2. In A549 cells, the percentage of S/G2 cells reduced gradually from Day 3-7 resulting in over 90% of A549 cells in a G1 like state (red) at 7-days post treatment (Figure 5E-F). In contrast, the percentage of S/G2 cells in both NCI-H1573 or NCI-H1299 increased until day 3, before returning to pre-treatment levels by day 7 (Figure 5 Supplement 3A-B). This mirrored our above *in vitro* data, where the presence of wild-type p53 (in A549 cells) corresponded with a higher rate of G2-exit and senescence compared to p53 null and mutant cells (Hastings et al., 2020), suggesting that the results we observed *in vitro* are recapitulated *in vivo*.

Cisplatin treatment during late G1 early S phase correlates with multiple divisions

The above data indicated that in p53 wild-type cells, those in late G1 and early S phase at the time of exposure had a greater capacity to undergo multiple division compared to cells G1 and S-G2/M phase. To assess this in greater detail, we synchronised and released FUCCI expressing A549 cells into either G1 or early S phase using either palbociclib or thymidine, respectively (Figure 6A-B). We combined these synchronisations with pulsed exposure to cisplatin at various points following release to target G1, early or late S phase populations, which were then monitored by time lapse microscopy. To target G1, cells were pulsed with cisplatin upon release from palbociclib (Cis at G1). This resulted in the majority of cells undergoing a prolonged S/G2 phase and then exiting back into a G1 like state without undergoing mitosis (G2-exit; Figure 6C), a state we described previously (Hastings et al., 2020). Notably, only 7/50 cells completed a single division, and no cells underwent multiple (2 or more) divisions during the 72 h time period (Figure 6A-D; proliferative). In contrast, targeting cells in G1/S (Cis at G1/S) using either

palbociclib or thymidine synchronisation with cisplatin, resulted in significantly more
cells (13/50 and 16/50 respectively) completing 2 or more divisions (Figure 6A-D;
proliferative). Finally, nearly all cells targeted in late S phase (Cis at S), completed
the first mitosis and then underwent prolonged S/G2 arrest and G2-exit, with only 4
out of 50 cells completing 2 divisions within the 72 h period (Figure 6B-D). Taken
together, these data indicate that cells in late G1 and early S at the time of cisplatin
exposure have a greater capacity to continue proliferating.

Disruption of DNA repair reduces ability of early S phase cells to proliferate
A major target of cisplatin is DNA, with intra and inter-strand cross-links disrupting
replication and repair, leading to stalled replication forks and the formation of double-
strand breaks (Gonzalez-Rajal et al., 2020). We therefore hypothesised that cells in
late G1/early S phase were able to repair cisplatin induced DNA damage during the
first cell cycle more efficiently than cells in early G1 or late S phase, thereby allowing
them to continue proliferation. To test this, we engineered A549 cells to stably co-
express a truncated form of 53BP1 fused to mApple (Apple-53BP1trunc), which has
previously been shown to bind double strand break sites co-marked with γH2A.X but
lacks any of the functional domains of 53BP1 (Yang et al., 2015). We combined this
with a PCNA Chromobody, where we replaced GFP with mNeonGreen, to mark sites
of active DNA replication (Figure 7A, inset). Cells were synchronised with thymidine
and pulsed with cisplatin 2 h prior to release to enrich for early S phase targeted
cells (as per Figure 6B), and then tracked by 4D-live cell imaging. Individual cells
were divided into either those in G1, early-S or mid/late-S based on the pattern of
PCNA foci (cyan), and then tracked through time (Burgess et al., 2012; Charrasse et
al., 2017). In control cells, a small number (<20) of 53BP1 positive foci (red hot) were
observed as cells underwent the first round of replication (Figure 7A,B). Daughter
and grand-daughter cells then displayed several (<5) large foci during G1 (up to 5
μm^2), which were resolved as cells entered S-phase and began replicating (Figure
7C, white arrow). Cells that were in G1 at the time of cisplatin exposure, entered S-
phase and rapidly accumulated a large number (~100) of 53BP1 positive foci, these
slowly reduced over the remainder of the time-lapse (Figure 7A-C), which
corresponded with an increase in the average size of the foci (~1 μm^2). In contrast,
cells that were in early S-phase and completed multiple (2 or more) divisions within
the 72 h timeframe, showed a rapid rise in foci number (~100), which then

decreased at the conclusion of S phase, correlating with an increase in foci size. A
small number of larger foci were present in the following G1 cell, although the size of
these foci was smaller than those observed in control daughter cells (~1 μm^2).
Interestingly, in grand-daughter and great grand-daughter cells, the size of G1 foci
increased (>2 μm^2), in-line with G1 foci observed in control cells (Figure 7A-C,
Figure 7 Supplement 1A). Finally, cells in mid-late S phase, also showed a large
number of 53BP1 foci, which increased in size as cells progressed through the first
G2 phase. Interestingly, the average number of 53BP1 foci in mid-S were higher and
were removed later, just prior to mitotic entry compared to cells from early S-phase
(Figure 7A-C, Figure 7 Supplement 1B). Furthermore, the quality of mitosis was
often reduced in cells from mid-S phase, with cells displaying chromatin bridges,
micronuclei and/or failed cytokinesis (Figure 7 Supplement 1C), correlating with the
increase in death during or after mitosis we observed previously (Figure 6C). The
subsequent daughter cells from those exposed in mid-S phase then showed a rapid
rise in the number of foci (>100) as they began replication. In contrast, the number of
53BP1 foci in early S-phase cells only increased mildly during replication and was
notably lower than the numbers observed in the first round of replication (Figure 7
Supplement 1A,B). Based on these results, we concluded that early-S phase cells
were able to either partially repair double strand breaks during the first round of DNA
replication, and/or mark damage for efficient repair in the subsequent daughter and
grand-daughter cells. In contrast, cells in G1 had much greater levels of damage and
remained arrested in the first G2 phase. Cells in mid/late S phase completed the first
division while acquiring damage but were unable to sufficiently repair the damage
before mitosis. Consequently, daughter cells with un-repaired damage had increased
rates of mitotic induced breaks as they attempted the second round of DNA
replication, leading to a strong S/G2 phase checkpoint arrest, similar to cells initially
exposed during G1.
371
Based on these results, we hypothesised that cells in early-S phase were better able
to take advantage of the high-fidelity homologous recombination (HR) pathway,
whose activity peaks in mid-S phase (Karanam et al., 2012) compared to G1 or late
S phase. To test this, we utilised the PARP inhibitor, olaparib, to trap PARP at single
strand break sites, leading to increased rates of replication fork stalling and reduced
capacity to repair DSBs by HR (Murai and Pommier, 2018). We hypothesised that

this would increase the rate of damage in all cells and reduce the ability of early-S
phase cells to repair during the first cell cycle. To test this, A549 FUCCI cells were
targeted in G1/S phase with cisplatin by synchronising with palbociclib or thymidine,
as previously described (Figure 6A,B). Cells were then treated with or without
olaparib (PARPi) for 1 h prior to pulsed cisplatin exposure and monitored by time
lapse microscopy (Figure 8A). In cells treated with cisplatin, co-treatment with PARPi
significantly reduced the total number of divisions (Figure 8B), indicating that cells
were unable to continue proliferating. This correlated with a trend toward G1 delay in
palbociclib and significant G1 delay in thymidine synchronised cells (Figure 8C).
Interestingly, although co-treatment with PARPi decreased the percentage of
proliferative (2 or more divisions) cells, there was only a small increase in death
observed (Figure 8D), indicating that PARPi alone is not sufficient to drive increased
toxicity to cisplatin in A549 cells. Importantly, inhibition of PARP did increase the rate
of 53BP1 foci formation compared to cisplatin alone in asynchronous cells (Figure
8E). Furthermore, this correlated with a significant increase in both the amount of γ-
H2AX staining and the size of cells (Figure 8F). We further validated these results by
inhibiting RAD51 with RI-1, to target HR mediated repair. RAD51 inhibition (RAD51i)
had no significant effect on cell growth compared to untreated controls (Figure 8
Supplement 1A). Single cell fate tracking of FUCCI expressing A549 cells revealed a
small but significant decrease in the number of divisions completed by early S phase
cells treated with RI-1 and cisplatin compared to cisplatin alone. This correlated with
a reduction in percentage of cells completing 2 or more divisions (proliferative) and
an increase in the percentage of cells undergoing a G2-exit phenotype (Figure 8
Supplement 1B-D). Taken together, these data suggest that targeting DNA repair
pathways during the first replication cycle results in more DSBs and pronounced
S/G2 cell cycle checkpoint arrest leading to G2-exit, likely due to a reduced ability to
repair DNA damage. This in turn reduces the ability of cells to undergo further
rounds of replication and division.
406
407

408   **Discussion:**
409   In this work, we have identified a novel, non-genetic mechanism of resistance to
410   platinum chemotherapy, which facilitates continued proliferation in a subset of LUAD
411   cells after pulsed exposure to cisplatin. These cells eventually outgrow the majority

of arrested cells over the course of 3 weeks *in vitro*. However, upon re-exposure, they remained equally sensitive, indicating that the mechanism of resistance is not hard-wired, nor did cells acquire resistance after the first exposure. Quantitative single cell fate tracking revealed that cells in late G1/early S phase at the time of exposure had a greater proliferative capacity after pulsed cisplatin exposure. This suggests that cell cycle stage at time of exposure impacts how cells respond to cisplatin. Interestingly, cisplatin treatment of head and neck (UM-SCC-38) cancer cells resulted in similar heterogenous cell cycle and cell fate responses (Luong et al., 2016), implying that the non-genetic cell cycle dependent mechanisms of resistance we observed here may translate to multiple cancer types beyond LUAD.

For cells that are in G1, which under normal physiological conditions represents the vast majority of cells (>70%) both *in vitro* and *in vivo*, the predominate response is to arrest in S/G2 during the first replication cycle (Figure 9). Notably, intracellular pH is lowest during G1, and cisplatin DNA binding is markedly increased in acidic conditions (Stewart, 2007), hence cells in G1 phase at the time of exposure may have higher levels of cisplatin-DNA adducts. In support, as G1 treated cells entered S phase, we noted increased numbers and size of 53BP1 foci, indicating wide-spread DSBs. This in turn correlated with prolonged S/G2 delays, indicating activation of DNA damage checkpoint signalling. Furthermore, we previously observed strong phosphorylation and activation of the G2 checkpoint proteins Chk1/2 across all LUAD cell lines after pulsed cisplatin exposure (Hastings et al., 2020). Similarly, RNAseq data from enlarged A549 cells showed consistent activation of G2/M checkpoint pathways. Notably, sustained S/G2 DNA damage has been shown to lead to ATR-mediated checkpoint signalling and sustained p21 expression, blocking CDK activity and preventing FOXM1 dependent G2 transcription required for mitotic entry (Saldivar et al., 2018). Prolonged p21 expression can lead to S/G2 checkpoint arrested cells undergoing a G2-exit and senescence-like state (Baus et al., 2003; Feringa et al., 2018). This provides an explanation for why cells with a functional p53/p21 pathway were more likely to undergo G2-exit, had higher rates of senescence and reduced numbers of cell divisions compared to those lacking p53.

In contrast, cells in mid-late S or G2/M phase at the time of exposure, proceeded through the first division with minimal delays, although numerous chromatin bridges

were observed between separating daughter cells, which correlated with increased
rates of mitotic and post-mitotic death (Figure 9). Similarly, loss, mutation, or as we
demonstrated previously, suppression of p53 signalling through inhibition of P70S6K
(Hastings et al., 2020), resulted in more cells entering mitosis and undergoing mitotic
and post-mitotic death. This likely explains why cells that lacked p53, were more
likely to undergo multiple aberrant divisions *in vitro* and maintain positive PCNA
staining *in vivo*, as they lacked the ability to instigate a stable S/G2 cell cycle
checkpoint arrest. Notably, chromatin bridges are often indicative of decatenation
failure and increased replication stress (Sarlós et al., 2017), and are commonly
formed after DNA damage caused by chemotherapies such as cisplatin. These
bridges are often repaired by homologous recombination (HR) during S phase (Chan
et al., 2017; Chan and West, 2018), but if repair is not completed before mitosis, they
can lead to increased rates of cytokinesis failure, chromosome instability and cell
death during or post mitotic division (Bakhoum et al., 2014; Burgess et al., 2014;
Hayashi and Karlseder, 2013). Importantly, chromosome instability, has been linked
to cancer heterogeneity, metastasis and acquisition of chemoresistance (Sansregret
et al., 2018), with single chromosome gains sufficient to reduce cisplatin mediated
cell death by delaying cells in G1 and slowing proliferation rate (Replogle et al.,
2020), highlighting the complexity and inter-dependence between innate and
acquired resistance mechanisms.

Thirdly, cells in late G1 and early S phase, showed a greater propensity to undergo
multiple divisions following exposure to cisplatin. This correlated with lower levels of
53BP1 foci during the first round of replication, which reduced further in subsequent
daughter and grand-daughter cells. A likely explanation is that early-S phase cells
would encounter cisplatin adducts during the first replication cycle, initiate repair
using high fidelity HR (Figure 9), which peaks during mid-S phase (Karanam et al.,
2012). In contrast, error-prone NHEJ, which is preferred during G1 and G2, is likely
favoured by cells exposed during G1 and late S-phase (Karanam et al., 2012). In
support, RNAseq analysis showed enrichment for HR associated BRCA1-mediated
signalling pathways in stable-sized cells compared to enlarged cisplatin treated cells.
It would be of interest to confirm this through future analysis of transcriptomes of cell
cycle phase sorted cells. Consequently, cells exposed during early S-phase likely
have more time and capacity to detect and accurately repair damage, thereby

avoiding chromatin bridges during mitosis compared to those treated later in S-
phase. In support, the 53BP1 foci resolved more quickly prior to the first mitosis in
early S phase treated cells. Post-mitotic G1 phase 53BP1 nuclear bodies increased
in size from daughter to grand-daughter cells, indicating increased efficiency in
identification and corralling of unrepaired DNA damage during the previous
replication cycle. These G1 53BP1 nuclear bodies prevent daughter cells from
encountering damaged DNA during replication (Watt et al., 2020), thereby avoiding
deleterious fork stalling. For under-replicated DNA, this provides the cell with a
second chance at repair (Spies et al., 2019). The partial repair in the first cell cycle
also likely ensures a pulsed p21 response (Hsu et al., 2019) with lower levels
preventing arrest in the following G1 phase (Barr et al., 2017), helping promote
continued proliferative capacity.
Finally, this work also indicates the potential for complications when co-administering
platinum chemotherapies with other targeted and cell cycle altering therapies such
as palbociclib. Specifically, pre- or co-administration of palbociclib with platinum may
inadvertently synchronise cells in late G1/early S phase, resulting in a higher
proportion of cells repairing the damage during the first cell cycle and maintaining
proliferative capacity. In support, current clinical trials in head and neck cancer
where both agents were given at the same time have resulted in no improvement to
cisplatin therapy and resulted in significant treatment related toxicity (Swiecicki et al.,
2020). Conversely, administering palbociclib after cisplatin (and other
chemotherapies), improves response and notably represses HR-dependent DNA
repair (Salvador-Barbero et al., 2020).
503

504 In summary, this work increases the understanding of the mechanisms driving
505 recovery from cisplatin treatment and identifies the need for novel combination
506 therapies that not only enhance cell death, but also prevent non-genetic, cell cycle
507 dependent resistance mechanisms.
508
509
510 **Materials and Methods:**

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Cell line (Homo sapiens)	A549	ATCC	CCL-185, RRID:CVCL_0023	
Cell line (Homo sapiens)	NCI-H1299	ATCC	CRL-5803, RRID:CVCL_0060	
Cell line (Homo sapiens)	NCI-H1573	ATCC	CRL-5877, RRID:CVCL_1478	
Transfected construct (human)	mVenus-hGeminin (1/110) (plasmid)	(Sakaue-Sawano et al., 2008)		
Transfected construct (human)	mCherry-hCdt1 (30/120) (plasmid)	(Sakaue-Sawano et al., 2008)		
Transfected construct (human)	Cell Cycle-Chromobody plasmid	Chromotek	ccr	Recloned into a pLVX lentiviral backbone with TagRFP replaced with mNeonGreen.
Transfected construct (human)	53BP1trunc-Apple	Addgene	69531, RRID:Addgene_69531	(Yang et al., 2015)

Transfected construct (human)	B_mCherry_IRES_neo3	Addgene	21044, RRID:Addgene_21044	(Steigeman n et al., 2009)
Transfected construct (human)	LeGO-Cer2 (Cerulean)	Addgene	27338, RRID:Addgene_27338	(Weber et al., 2008)
Transfected construct (human)	LeGO-V2 (Venus fluorescent protein)	Addgene	27340, RRID:Addgene_27340	(Weber et al., 2008)
Transfected construct (human)	LeGO-C2 (mCherry)	Addgene	27339, RRID:Addgene_27339	(Weber et al., 2008)
Antibody	Anti-p21 Waf1/Cip1 (rabbit monoclonal)	Cell Signaling Technology	2947, RRID:AB_330945	Flow 1:200
Antibody	Anti-p16 Ink4a	Abcam	AB201980, RRID: N/A https://www.abcam.com/CDKN2Ap16INK4a-antibody-1D7D2A1-ab201980.html	Flow 1:200
Antibody	Anti-gamma H2A.X (phospho S139) antibody	Cell Signaling Technology	9718, RRID:AB_2118009	Flow 1:200
Antibody	Anti-PCNA	Abcam	AB29, RRID:AB_303394	IHC 1:500 - 1:2000
Antibody	BrdU-FITC	BD Biosciences	347583, RRID:AB_400327	Flow 1:20
Fluorescent probe	Alexa Fluor 647 Phalloidin	Thermo Fisher	Thermo Fisher Scientific	1:100 of 400X stock

		Scientific		
Chemical compound, drug	Cisplatin	Hospira Australia	88S035	
Chemical compound, drug	Carboplatin	Abcam	ab120828	
Chemical compound, drug	Olaparib	Selleck Chem	S1060	
Chemical compound, drug	Palbociclib	Selleck Chem	S1116	
Chemical compound, drug	RI-1	Selleck Chem	S8077	
Chemical compound, drug	PureLink RNase A	Thermo Fisher Scientific	12091021	
Chemical compound, Substrate	C12FDG (5-Dodecanoylaminofluorescein Di-β-D-Galactopyranoside)	Thermo Fisher Scientific	D2893	
Chemical compound, stain	H33342	Sigma	B2261	1µg/ml
Chemical compou	Propidium Iodide	Thermo Fisher	P3566	

nd, stain		Scientific		
Chemical compund, drug	Thymidine	Selleck Chem	S4803	
Chemical compound, drug	BrdU (5-Bromo-2'-Deoxyuridine)	Thermo Fisher Scientific	B23151	
Commercial assay or kit	ImaGene Green™ C12FDG lacZ Gene Expression Kit	Molecular Probes	I2904	
Other	Matrigel Basement Membrane	Bio-Strategy	BDAA354230	
Software	Fiji/Image J	NIH	(Schindelin et al., 2012)	https://imagej.net/Fiji
Software	FlowJo	BD Biosciences	https://www.flowjo.com	
Software	GraphPad Prism (v9.1.0)	Graph Pad	https://www.graphpad.com	
Software	Huygens Professional	Scientific Volume Imaging (SVI)	https://svi.nl/Huygens-Professional	
Software	LAS-X	Leica	https://www.leica-microsystems.com/products/microscope-software/p/leica-las-x-ls/	

Antibodies, Plasmids, and Reagents

513 The γH2A.X (S139) (AB26350), p16 (AB201980) and PCNA (AB29) antibodies were

514 from Abcam (MA, USA), p21 antibody (2947) was purchased from Cell Signaling

515 Technology (MA, USA). BrdU-FITC antibody was purchased from BD-Biosciences

516 (BD-347583). Alexa-647 Conjugated Phalloidin antibody was purchased from

ThermoFisher Scientific (A22287). The plasmids for FUCCI live cell imaging, mVenus-hGeminin(1/110) and mCherry-hCdt1(30/120), were a kind gift from Dr Atsushi Miyawaki (Riken, Japan). The LeGO plasmids were obtained from Addgene (#27338, #27339, #27340) (Weber et al., 2008). Thymidine (S4803), olaparib (S1060), palbociclib (S1116) and RI-1 (S8077) were from Selleck Chem (MA, USA). Deoxycytidine (sc-231247) was from Santa Cruz Biotechnology (TX, USA). BrdU was purchased from ThermoFisher (B23151). Cisplatin was obtained from Hospira Australia (B23151) and Carboplatin from Abcam (ab120828).

Cell lines

All lung adenocarcinoma cell (LUAD) lines have been previously described (Hastings et al., 2020; Marini et al., 2018). The lines were cultured in Advanced RPMI (Gibco, 12633012) containing 1% FCS and 1% GlutaMAX (35050– 061, Gibco) under standard tissue culture conditions (5% CO_2, 20% O_2). All cell lines were authenticated by short tandem repeat polymorphism, single-nucleotide polymorphism, and fingerprint analyses, passaged for less than 6 months.

Stable cell lines expressing the FUCCI biosensor were generated previously (Hastings et al., 2020). H2B-mCherry cells were generated by lentiviral transfection, followed by FACS sorting of low-expressing clones. Finally, dual Chromobody and 53BP1 A549 cells, were generated by lentiviral transfection with the PCNA-chromobody, with low expressing clones isolated by cell sorting. These were then subsequently transfected (lentiviral) with a truncated form of 53BP1 fused to mApple (Apple-53BP1trunc), with cells sorted based on both mNeonGreen and mApple to isolate dual expressing clones.

Colony Formation Assay and Senescence-Associated Beta-Galactosidase assay

For colony formation assays, cells were seeded on 6-well plates, pulsed with cisplatin (or not) and one to two weeks later, colonies were stained with 0.5% crystal violet and counted using ImageJ/Fiji software. For β-Gal assays, cells were seeded on 6-well plates, pulsed with cisplatin (or not), fixed and stained at 3 days, following manufacturer's protocol (Cell Signaling Technology, #9860). Unpaired students t-tests along with bar graphs were generated using GraphPad Prism (v9.1.0).

LeGO clonal analysis

A549, NCI-H1573 and NCI-H1299 cells were transfected with LeGO lentiviral

particles (Addgene plasmids #27338, #27339, #27340) (Weber et al., 2008) following

the method described in (Weber et al., 2012). Cells were treated with/without

cisplatin and images were taken at 3 days and at 21 days (A549 and NCI-H1299) or

42 days (NCI-H1573) after cisplatin exposure. 100 images were taken per timepoint

and per condition (3 replicates) and the experiment was done twice. The total

number of clones (unique colour cues) and the number of cells within each clone

was determined. Briefly, images are opened and converted to 16bit.tif files. An image

is duplicated and converted to RGB overlay. The duplicate has background

subtracted using a rolling ball at 250 considering colours separately and using a

sliding parabaloid. The image is smoothed using a mean filter radius 5. Using the

"find maxima" function a point within individual cells is identified and then enlarged to

a circle radius of 5 pixels. These ROI are then applied as a mask to the

unprocessed, raw, image data and the average red, green and blue values within

these ROI collected and exported in .csv format. RGB values from each of the .csv

files for each of the 100 images are compiled. Data from cells where an R, G or B

value is too high or too low are removed. 512 unique colours were identified and

cells were classified and assigned to each of the 512 colours. More than 90% of all

cells were assigned to one of the 64 most represented colours, with analysis

performed using these 64 groups and positive clonal colour assigned when the

colour represented >0.1% of the population. Graphs were generated using

GraphPad Prism (v9.1.0).

Animal Experiments

Animal experiments were conducted in accordance with the Garvan/St Vincent's

Animal Ethics Committee (guidelines ARA 18_17, ARA_16_13) and in compliance

with the Australian code of practice for care and use of animals for scientific

purposes. Mice were kept in standard housing at a 12 h daylight cycle and fed *ad*

libitum. Cage enrichment refinement was undertaken with mice implanted with

mammary optical imaging windows, supplying the fully plastic IVC cages with papier-

mâché domes, feeding supplied in trays on the cage floor and soft tissues as nesting

material. For *in vivo* xenograft models A549 cells (2×10^6) were resuspended in 100

µL PBS:Matrigel (1:1) and injected subcutaneously into the flanks of BALB/c-

Fox1nuAusb mice (Australian BioResource). Tumour growth was assessed twice

weekly by calliper measurement and mice were randomised to treatment arms when

tumours reached 150 mm^3 (using the formula: width^2 x length x 0.5). Carboplatin (60

mg/kg) was delivered by a single i.p injection. Tumours were harvest at 3-7 days

post treatment and analysed by IHC for cell size and PCNA positive staining.

Implantation of Optical imaging windows

BALB/c-Foxn1nu/Ausb mice were injected with 1x10^6 A549-FUCCI subcutaneously

near the inguinal mammary fat pad. Following the development of palpable tumours,

mice were engrafted with titanium mammary imaging windows (Russell Symes &

Company) as described previously (Gligorijevic et al., 2009; Kedrin et al., 2008;

Nobis et al., 2017; Ritsma et al., 2013). Briefly, mice were treated with 5 mg/kg of the

analgesic Carprofen (Rimadyl) in pH neutral drinking water 24 h prior and up to a

minimum of 72 h post-surgery. Mice further received subcutaneous injections of

buprenorphine (0.075mg/kg, Temgesic) immediately prior and 6 h post-surgery. The

titanium window was prepared 24 h prior to surgery by gluing a 12 mm glass

coverslip (Electron Microscopy Science) using cyanoacrylate to the groove on the

outer rim of the titanium window. Following anaesthetic induction at 4% isoflurane

delivered via a vaporizer (VetFlo) supplemented with oxygen, mice were kept at a

steady 1-2% maintenance anaesthesia for the duration of the surgery on a heated

pad. The incision site was disinfected using 0.5% chlorhexidine/ 70% ethanol. A

straight incision was made into the skin above the developed subcutaneous tumour

and following blunt dissection of the skin surrounding the incision a purse string

suture (5-0 Mersilk, Ethicon) placed. The windows were then inserted and held in

place by tightening the suture, disappearing along with the skin into the groove of the

window and tied off. Mice were allowed to recover for a minimum of 72 h post-

surgery, actively foraging, feeding and grooming within minutes from being removed

from the anaesthesia respirator. A minimum of 24 h prior to imaging and treatment

mice were weaned off the Carprofen analgesic in the drinking water.

In vivo imaging

Mice were imaged under 1-2% isofluorane on a heated stage (Digital Pixel, UK) prior

to and 1 day, 2 days, 3 days and 7 days after ip injection of 60 mg/kg Carboplatinum

(Sigma) or the saline vehicle. Multi-photon imaging was performed using a Leica

DMI 6000 SP8 confocal microscope using a 25x 0.95 NA water immersion objective

on an inverted stage. For A549-FUCCI imaging the Ti:Sapphire femto-second laser
(Coherent Chameleon Ultra II, Coherent) excitation source operating at 80 MHz was
tuned to 920 nm and the RLD-HyD detectors with 460/40, 525/50 and 585/40
bandpass emission filters used to detect the second harmonic generation (SHG) of
the collagen I, mAzamiGreen and mKO2 respectively. Images were acquired at a
line rate of 400 Hz, 512x512 pixel and a line average of 8.

625
626
627

Flow Cytometry Analysis and Sorting
Samples for flow cytometry were fixed in -20 °C ethanol overnight, and then stained
with a primary antibody against p21 (Cell Signal Technology, 2947), p16 (Abcam,
ab201980) or gamma-H2A.X (S139) (Abcam, ab26350) followed by incubation with
an Alexa Fluor 647 secondary antibody (Invitrogen). Flow cytometry was performed
using a Beckman CytoFlex S. For BrdU incorporation analysis, cells were incubated
with BrdU at 10 µM for 2 h before overnight ethanol fixation at -20 °C. An antibody
against BrdU coupled with FITC (BD-347583) was used for staining and flow
cytometry was done using a Beckman CytoFlex S. For senescence assays we used
ImaGene Green™ C12FDG lacZ Gene Expression Kit (Molecular probes, I-2904).
Three days after cisplatin exposure cells were incubated for 30 min with Bafilomycin
A1 (Sigma, B1793) in RPMI medium without phenol red (Gibco) supplemented with
1% FBS before adding $C_{12}FDG$ to the media at 20 µM final concentration. Cells were
incubated for 60 minutes prior to 15 minutes fixation with 4% PFA and processed for
FACS analysis. Flow cytometry was performed using a Beckman CytoFlex S.
For cell sorting and RNAseq analysis, A549 cells were treated with or without
cisplatin (5 µg/ml) for 2 h, and then allowed to recover for 3-days. Cells were
collected by trypsinisation and then sorted using a BD FACS Aria III. The gates for
stable and enlarged cells were determined by running untreated control cells and
identifying cell size based on Forward and Side-Scatter (FSC vs SSC) area
parameters. Doublets were excluded based on area and height parameters of FSC
and SSC. Sorted cells were frozen as a pellet in dry ice and stored at -80 °C until
RNA purification.

Immunofluorescence and live cell imaging

Cells were grown on Histogrip (Life Technologies) coated glass coverslips and fixed with 3.7% formaldehyde diluted in PHEM buffer (60 mM Pipes, 25 mM hepes, 1 mM EGTA, 2 mM MgCl2) with 0.5% Triton X-100 for 10 min. All cells were washed and then blocked (3% BSA, 0,1% Tween 20 in PBS) for 30 min. Cells were incubated with primary antibodies for 2 h at room temperature in blocking solution. DNA was stained with H33342 and imaged using an EVOS FL2 Auto Imager (Thermofisher) or a Leica SP8-X confocal with white light laser using either a 20X (NA 0.75) or 63X (NA 1.40) objective. In some cases, 0.3 µm Z-sections were taken and displayed as 2D slices or maximum projections using Fiji (Image J v2.1.0/1.53c) and compiled using Adobe Photoshop CC 2020 software. Deconvolution and 3D volume renderings were performed using Huygens Professional Software (Scientific Volume Imaging, v20.04), while nuclear size analysis was performed using StarDist (Schmidt et al., 2018) plugins for Fiji/ImageJ. Live cell imaging and IncuCyte (Sartorius) proliferation assays were performed as previously described (Hastings et al., 2020; Rogers et al., 2018). Briefly, for live cell imaging, cells were seeded at 35% confluence on 6- or 12-well plates and imaged using a Leica DMI6000 with a 20X NA 0.4 objective. Images were taken every 10-20 min for up to 72 h. Individual cells were followed and scored for nuclear envelope breakdown (NEBD) and first signs of anaphase as previously described (Caldon and Burgess, 2019). Mitotic length = NEBD to anaphase, while interphase length = anaphase to next daughter cell NEBD. Only the first daughter cell to divide was followed and annotated. For IncuCyte assays, cells were seeded on 12- or 24-well plates and filmed for up to 4 days at 4 h intervals. Confluence and nuclear masks were generated and used to determine cell proliferation as previously described (McCloy et al., 2014). Statistical analysis along with box and violin plots were generated using GraphPad Prism (v9.1.0).

For 53BP1 and PCNA chromobody experiments, cells were seeded on 8-Well Ibidi Polymer Coverslip µ-Slides (#80826), synchronised with thymidine or palbociclib and pulsed with cisplatin for 2 h, before imaging on a Lecia SP8 confocal microscope fitted with a white light laser, hybrid detectors (HyD), a 63X HC PL APO CS2 (NA 1.40) objective and stage top incubator system set at 37 ^{o}C and 5% CO_2. Multiple X/Y positions, and a 10 µm z-stack (1 µm z-section) were taken every 30 min for 72 h, with 4D deconvolution and volume rendering performed with Huygens Professional (v20.04) software (Netherlands). 53BP1 and PCNA foci analysis was

686 performed on 2D-maximim intensity projections using appropriate thresholds coupled
687 with the Analyse Particles module within ImageJ/Fiji. The pattern of PCNA foci was
688 used to position cells in early, mid or late S phase, as previously described (Burgess
689 et al., 2012; Charrasse et al., 2017).
690

Immunohistochemistry

Immunohistochemistry was performed on formalin fixed paraffin embedded sections
using the Leica BOND RX (Leica, Wetzlar, Germany). Slides were first dewaxed and
rehydrated, followed by heat induced antigen retrieval performed with Epitope
Retrieval Solution 1 BOND (Leica, Wetzlar, Germany). PCNA Primary antibody was
diluted 1:500 (Abcam, ab29) in Leica antibody diluent and incubated for 60 min on
slides. Antibody staining was completed using the Bond Polymer Refine IHC protocol
and reagents (Leica, Wetzlar, Germany). Slides were counterstained on the Leica
Autostainer XL (Leica, Wetzlar, Germany). Leica CV5030 Glass Coverslipper (Leica,
Wetzlar, Germany) and brightfield images were taken on the Aperio CS2 Slide
Scanner (Leica, Wetzlar, Germany). Quantification of PCNA staining was performed
on three fields of view for each tumour section using QuPath (v0.2.3)(Bankhead et
al., 2017). Student's t-test statistical analysis along with dot plots and bar graphs
were generated using GraphPad Prism (v9.1.0).
705

RNA isolation, RNA sequencing (RNA-seq), SNV alignment and analysis

Cell pellets were obtained from the different conditions/populations. Cell pellets were
frozen in dry ice prior to storage at -80 ^{o}C. Total RNA was purified using miRNeasy
Micro Kit (Qiagen, 217084) following the manufacturer's protocol, including a DNase
treatment. RNA concentration and quality were also measured by Qubit and
Nanodrop. Samples were only used if they showed a 260/280 ratio >2.0 (Nanodrop).
RNA integrity was determined on an Agilent 2100 Bioanalyser and samples were
only used if they showed a RIN of >8. Three sets of RNA were collected per
condition. Compliant samples were sent to the Australian Genome Research Facility
(AGRF) for RNA sequencing with poly(A) selection. Briefly, 20 million 100 bp single
end RNA-seq was conducted on an Illumina NovaSeq platform. The library was
prepared using the TruSeq stranded RNA sample preparation protocol (Illumina).
The cleaned sequence reads were aligned against the *Homo sapiens* genome (Build
version hg38) and the RNA-seq aligner, "Spliced Transcripts Alignment to a

Reference (STAR)" aligner (v2.5.3a) (Dobin et al., 2013), was used to map reads to
the genomic sequence. Transcripts were assembled using the StringTie tool v1.3.3
(Pertea et al., 2015) with the read alignment (hg38) and reference annotation-based
assembly option (RABT). Raw data were deposited in the NCBI Gene Expression
Omnibus (GEO) data repository accession number GSE161800.

The raw data from each cell line was aligned to the human genome reference build
GRCh38/hg38 using STAR aligner v2.5.3a by AGRF. Single Nucleotide Variations
(SNVs) were identified using SNV caller Freebayes (v1.3.1;
https://github.com/ekg/freebayes) and annotated using Bcftools (v1.9) (Danecek and
McCarthy, 2017) with database NCBI dbSNP (v146) (Sherry et al., 2001). Heatmaps,
principal component analysis (PCA) and biological coefficient variant plots were
made using R language and software (The R Foundation) with the DESeq2 package
(Love et al., 2014). The log2 (fold change) scale was normalised and transformed by
considering library size or other normalisation factors. The transformation method
and the variance stabilising transformation (VST) (Anders and Huber, 2010) for over
dispersed counts have been applied in DESeq2. The VST is effective at stabilising
variance, because it considers the differences in size factors, such as the datasets
with large variation in sequencing depth (Love et al., 2014). Canonical Pathway
analysis of known proliferation, cell cycle, migration and cell death-related signalling
pathways were conducted using the Ingenuity Pathway Analysis software (QIAGEN),
as previously described (Johnson et al., 2020). Briefly, minimum significance cut offs
of p-value>0.05 and Z scores of >2 and <-2 were applied for pathways analysis. For
Gene Set Enrichment Analysis (GSEA) a ranked gene list was prepared from
Proliferative versus Arrest and analysed with GSEA 4.1.0 software
(https://www.gsea-msigdb.org/gsea/index.jsp) using a curated gene set of canonical
pathways (2868 gene sets) (https://www.gsea-
msigdb.org/gsea/msigdb/collections.jsp#C2) (Mootha et al., 2003; Subramanian et
al., 2005). The enrichment map was generated using Cytoscape 3.8.2 software
(https://cytoscape.org/) (Shannon et al., 2003), using p-value (<0.005) and FDR
(q<0.1) cut offs. Volcano and dot plots were generated using GraphPad PRISM
(v9.1.0) and figures compiled using Adobe Illustrator Creative Cloud (v25).

Acknowledgements

The authors would like to acknowledge Dr Atsushi Miyawaki (Riken, Japan) for

provision of the mVenus-hGeminin(1/110) and mKO-hCdt1(30/120) constructs for

FUCCI imaging. Dr Liz Caldon for her insightful and helpful comments. The Patricia

Helen Guest Fellowship for their generous support. The authors acknowledge the

ANZAC Microscopy and Flow Facility, the Sydney Informatics Hub and the use of the

University of Sydney's high performance computing cluster, Artemis.

Additional Information

Funding:

Cancer Institute NSW   2013/FRL102   David   R

Croucher

Cancer Institute NSW   15/REG/1-17   David   R

Croucher

Cancer Institute NSW   10/FRL/3-02   Andrew Burgess

National Breast Cancer Foundation   IIRS-18-103   Andrew Burgess

Tour de Cure   RSP-230-2020   Andrew Burgess

The funders had no role in study design, data collection and interpretation, or the

decision to submit the work for publication.

Author contributions

Alvaro Gonzalez Rajal, Conceptualization, Data curation, Formal analysis,

Visualization, Methodology; Kamila A Marzec, Data curation, Formal analysis,

Visualization; Rachael A McCloy, Formal analysis, Investigation, Methodology; Max

Nobis, Formal analysis, Visualization, Methodology; Venessa Chin, Resources, Data

curation, Supervision, Investigation, Methodology; Jordan F. Hastings, Formal

analysis, Visualization; Kaitao Lai, Formal analysis, Visualization, Data curation;

Marina Kennerson, Supervision, Methodology; Vijesh Vaghjiani, Data curation,

Formal analysis; William E Hughes, Data curation, Formal analysis; Paul Timpson,

Supervision, Methodology; Jason Cain, Data curation, Formal analysis, Supervision,

Methodology; D Neil Watkins, Conceptualization, Supervision, Methodology, Data
curation, Formal analysis, Visualization; David R Croucher, Conceptualization,
Formal analysis, Supervision, Funding acquisition, Investigation, Project
administration; Andrew Burgess, Conceptualization, Formal analysis, Supervision,
Validation, Investigation, Visualization, Methodology, Project administration, Funding
acquisition.
794
Animal experimentation: All experiments were carried out in compliance with the
Australian code for the care and use of animals for scientific purposes and in
compliance with Garvan Institute of Medical Research/St. Vincent's Hospital Animal
Ethics Committee guidelines (ARA_18_17, ARA_16_13).

Author ORCIDs
Alvaro Gonzalez-Rajal: https://orcid.org/0000-0002-9230-0339
Kamila A. Marzec: https://orcid.org/0000-0002-3051-2205
Rachael McCloy: https://orcid.org/0000-0001-7791-4044
Max Nobis: https://orcid.org/0000-0002-1861-1390
Venessa Chin: https://orcid.org/0000-0002-4630-4451
Marina Kennerson https://orcid.org/0000-0003-3332-5074
Kaitao Lai https://orcid.org/0000-0002-9420-9352
D. Neil Watkins: https://orcid.org/0000-0001-8218-4920
David R Croucher https://orcid.org/0000-0003-4965-8674
Andrew Burgess https://orcid.org/0000-0003-4536-9226

810
811
812

Figure Legends:
Figure 1 – Rechallenging LUAD cells with cisplatin results in similar response
profiles. (A) Schematic of rechallenging experiments. Briefly, cells stably expressing
H2B-mCherry were pulsed with 5 µg/ml cisplatin for 2 h. Cell proliferation (nuclear
number and cell size) were then tracked for up to 4 days. Cells were then allowed to
recover for 21 (A549 and NCI-H1299) or 42 days (NCI-H1573), re-culturing once
confluent, before being re-pulsed with cisplatin. **(B)** Cell confluence and cell number
were tracked for up to 4 days using IncuCyte based time-lapse imaging. Shown are
the mean +/- SD of n=3 biological repeats. **(C)** Immunofluorescence of cells at 72 h

post cisplatin treatment. Nuclei = cyan, Phalloidin-Alexa 647= red, scale bar = 10

µm. **(D)** Quantification of cell size and nuclear size from A, with a minimum of n=200

cells analysed per condition. Shown are the mean +/- SD. Statistical significance

was determined by one-way ANOVA (****$p<0.0001$, n.s = not significant).

Figure 2 – Variable cell size and clonal outgrowth in post-cisplatin treated

cells. (A) Representative images from cells treated as per Figure 1A showing

Control (Cont.) proliferative (Prolif.) and arrested (Arrest) cells. Scale bars = 100 µm.

(B) Cell and nuclear size were calculated on sub-populations of cells that appeared

to form clonal outgrowths. Quantification of cell size and nuclear size from A, with a

minimum of n=100 cells analysed per condition. Shown are the mean +/- SD.

Statistical significance was determined by one-way ANOVA (****$p<0.0001$, n.s = not

significant). **(C)** LeGo RGB colour guided clonal cell tracking system was used to

track clonal dynamics after cisplatin pulse treatment. Cells were treated as per

Figure 1A, with clonal identification and quantification **(D),** measured at 3 and 21-

days post cisplatin exposure using Image J/Fiji (42-days post cisplatin exposure for

NCI-H1573).

Figure 3 – In vivo validation of cisplatin effects on cell size and proliferation.

(A) A549 and NCI-H1299 cells were injected subcutaneously with $2x10^6$ cells into the

flanks of nude mice (n=5). Carboplatin (60 mg/kg) was delivered by a single tail-vein

injection and tumours were harvested at 3 days post treatment and analysed by IHC

for cell size and PCNA positive staining. Scale bar = 50 µm **(B)** Quantification of IHC

images from A (control n=300, carbo n=400). Shown are the mean +/- SD. Statistical

significance was determined by unpaired two-tailed Students t-test (****$p<0.0001$,

n.s = not significant).

Figure 4 – Comparative RNAseq analysis of FACS sorted cisplatin treated

cells. (A) Schematic (created with BioRender.com) describing treatment, sorting and

analysis pipeline **(B)** Representative example of pre- and post-sorted control and

cisplatin treated cells. **(C)** Hierarchical clustering of 3-independent biological repeat

experiments of all altered genes identified by RNA-seq. **(D-F)** Volcano plots

displaying significantly downregulated (blue) or upregulated (orange) genes and

subsequent IPA canonical pathway analysis. Predictions of inhibition (blue) or

activation (orange) or no change (white) states are based on the
Ingenuity®Knowledge Base, which compares the expected change with
experimental observation to all known upstream regulators. Variable stringent p-
value (>1.3) and z-score (>0.5) cut-offs were used to limit pathways to top 7–8 most
significant hits**. (G)** Hierarchical clustering of IPA comparative canonical pathway
analysis.

Figure 5 – Cell cycle and p53 status at time of exposure correlates with cell
fate outcomes
(A-B) The fate of asynchronously growing FUCCI expressing A549 individual control
(n=50) and cisplatin pulsed cells (n=100) was tracked by time lapse microscopy, with
images taken every 30 min for 72 h. Cisplatin treated cell analysis was slightly
biased for cells that underwent multiple divisions. **(C)** Violin plots of the number of
divisions from cells in (A-B), based on the cell cycle phase at time of cisplatin
exposure. Statistical significance was determined by one-way ANOVA with Tukey
correction for multiple comparisons (**p<0.01, ****p<0.0001, ns = not significant). **(D)**
Schematic (created with BioRender.com) of optical window based longitudinal *in vivo*
imaging of FUCCI A549 cells. **(E)** Representative 3D projection images from mice
imaged at day 1 and day 7 with carboplatin (Carbo) or control (Saline). **(F)**
Quantification of the proportion of red (G1), yellow (G1/S) and green (S/G2-M) cells
found in tumours (n=3) from day 0 to 7.

Figure 6 – Cell Cycle dependent regulation of cisplatin response. (A-B)
Schematic of palbociclib (Palbo) and thymidine (Thy) protocols used to synchronise
FUCCI expressing A549 cells in G1, G1/S and S phase prior to 2 h cisplatin pulse
treatment. Specifically, G1 cells were released from palbociclib and pulsed
immediately with cisplatin (Palbo + Cis at G1). G1/S cells were pulsed with cisplatin
at 6 h post release from palbociclib (Palbo + Cis at G1/S), or alternatively G1/S
phase cells targeted by treatment with cisplatin 2 h prior to release from thymidine
(Thy + Cis at G1/S). Finally, S phase cells targeted by pulsing with cisplatin at 4 h
post thymidine release (Thy + Cis at S). The fate of individual cells (n=50) was
tracked by time lapse microscopy, with images taken every 30 min for 72 h. **(C)**
Quantification of cell fate outcomes from A; including G1 arrest before mitosis (G1
ABM), G1 arrest after mitosis (G1 AAM), death before mitosis (DBM) and death after

mitosis (DAM) and Proliferative (Prolif.). **(D)** Quantification of the total number of cell divisions observed in each condition (n=50). Mean is shown, statistical significance was determined by one-way ANOVA with Tukey correction for multiple comparisons (****$p<0.0001$, *$p<0.05$).

Figure 7 – Dual DNA replication and damage biosensor analysis of cisplatin treated cells. (A) Representative maximum image projections of A549 cells co-expressing a mNeonGreen tagged PCNA chromobody (cyan) and a truncated version of 53BP1 tagged with mApple (trunc53BP1-mApple; red-hot LUT). Cells were imaged using confocal microscopy, with 10 µm thick z-stack (1 µm slice) taken every 30 min for 72 h. Scale bar 10 µm. **(B)** Quantification of the size and number of PCNA and 53BP1 foci for each cell shown in A. **(C)** 3D volume renders from cells in A for the indicated times, with cropped zoom areas (right image). White arrows indicate 53BP1 foci that reduce in size over time (min). Scale bars 5 µm and 1 µm for left and right panels respectively.

Figure 8 – Inhibition of PARP reduces ability of early S phase cells to maintain proliferative capacity. (A) Schematic of palbociclib (Palbo) and thymidine (Thy) protocols used to synchronise FUCCI expressing A549 cells in G1, early and late S phase prior to olaparib (PARPi; 1 µM) and 2 h cisplatin (5 µg/ml) pulse treatment. The fate of (n=50) individual cells was tracked by time-lapse microscopy, with images taken every 30 min for 72 h. Quantification of the total number of cell divisions **(B)** and G1 length after **(C)** first mitotic division observed in each condition. Statistical significance was determined by a one-way ANOVA test with correction for multiple comparisons (****$p<0.0001$, ***$p<0.001$, **$p<0.01$, *$p<0.05$). **(D)** Quantification of cell fate outcomes from A; including G1 arrest before mitosis (G1 ABM), G1 arrest after mitosis (G1 AAM), death before mitosis (DBM) and death after mitosis (DAM) and Proliferative (Prolif.), i.e. cells that divided 2 or more times. **(E)** Fluorescent imaging of asynchronous A549 dual biosensor cells pulsed with cisplatin for 2 h. The percentage of cells with less than 5 (<5) or more than 15 (>15) 53BP1 foci/cell after cisplatin treatment are shown. A minimum of 250 cells per timepoint and condition were counted from (n=3) biological repeats. Statistical significance was determined by two-way ANOVA (*$p<0.05$). **(F)** Thymidine synchronised cells treated as per A, were harvested, and analysed for cell size and γ-H2AX by flow

cytometry. Representative FACS plots and quantification from (n=3) biological
repeats are shown. Statistical significance was determined by one-way ANOVA with
Tukey correction for multiple comparisons (***p<0.001).

Figure 9 – Schematic outlining cell cycle dependence of cisplatin recovery.
Briefly, cells exposed in G1 undergo S/G2 phase arrest, marked by high levels of
P70S6K, p53 and p21, which results in a stable and permanent cell cycle exit from
G2 phase. Cells in late S phase likely receive platinum adducts in areas of already
duplicated DNA, with error-prone non-homologous end joining (NHEJ) favoured over
homologous recombination (HR). Combining this with an increase in cells
undergoing aberrant division results in daughter cells displaying higher rates of
damage during replication, further resulting in S/G2 arrest and senescence. In
contrast, cells in late G1/early S phase have the opportunity to detect and repair
damage by HR during the first cycle, thereby increasing chances that daughter cells
will successfully complete and repair damage, allowing continued proliferation.

Supplementary Figures and Tables
Figure 1 Supplement 1: (A) Representative images and quantification of colony
formation assays for A549, NCI-H1573 and NCI-H1299 cells treated with or without
cisplatin. Colonies were fixed, stained and counted at 10 (A549 and NCI-H1299) or
14 days (NCI-H1573) after cisplatin exposure. Scale bar = 100 μm. Shown are the
mean +/- SD of 3 biological repeats. Statistical significance was determined by
Student's t-test (****p<0.0001,***p<0.001, **p<0.01). **(B)** Long term recovery of A549
cells treated with a single pulse of 5 μg/ml was analysed by flow cytometry for cell
cycle by DNA content and cell size using FSC and SSC parameters. Shown are the
mean +/- SD of n=3 biological repeats. Statistical significance was determined by
two-way (cell cycle) or one-way (cell size) ANOVA with Tukey's multiple
comparisons test (****p<0.0001, ***p=0.0006, ns = not significant).

Figure 2 Supplement 1: (A) Representative images and quantification of A549,
NCI-H1573 and CI-H1299 cells 3 days after cisplatin exposure stained for beta-
galactosidase. Scale bar = 100 μm. Shown are the mean +/- SD of 3 biological
repeats. Statistical significance was determined by unpaired two-tailed Student's t-
test (****p<0.0001,***p<0.001,**p<0.01). **(B)** FACS analysis of beta galactosidase

activity levels (C$_{12}$FDG) versus "cell size" (FSC) and **(C)** p21, p16 and at 3-days after

cisplatin exposure. Shown are the mean +/- SD of 3 biological repeats. Statistical

significance was determined by unpaired two-tailed Student's t-test

(****p<0.0001,***p<0.001, *p<0.05). **(D)** Three biological repeat experiments of LeGo

RGB colour guided clonal cell tracking system was used to track clonal dynamics

after cisplatin pulse treatment in each cell line. Cells were treated as per Figure 1A,

with clonal identification and quantification measured at 3 and 21-days post cisplatin

exposure using Image J/Fiji (42-days post cisplatin exposure for NCI-H1573). For

summary data, only colours that represented >0.1% of total population were scored.

Figure 3 Supplement 1: Cells were pulsed cisplatin or not (0 h) for 2h and then with

BrdU for 2 h prior to harvesting at the indicated timepoints. Cells were counter-

stained with PI with DNA content and BrdU staining analysed by flow cytometry to

identify cells in G1 (BrdU negative, 2n DNA content; Active S phase = between 2n-

4n DNA and positive for BrdU, inactive S phase = 2n-4n DNA content and negative

for BrdU; G2/M = 4n DNA content. The percentage of cells +/- standard deviation for

each population from (n=3) independent repeats are shown.

Figure 4 Supplement 1: (A) Gating strategy used for sorting control and cisplatin

treated cells based on cell size and the exclusion of doublets. **(B)** Cytoscape

Enrichment map of curated gene sets of canonical pathways for Proliferative versus

Arrested cells. Each node represents a gene set and each line connecting nodes a

gene "common" to both nodes. Shown are gene sets enriched in proliferative (Red)

and Arrested (blue) cells. The size of the node represents the NES score of that

particular gene set. **(C)** GSEA Enrichment plots for Proliferative versus Arrest cells

(curated gene sets for canonical pathways). From left to right and top to bottom: Cell

Cycle, Homology Directed Repair, Fanconi Pathway and ATR Pathway. The green

curve corresponds to the ES (enrichment score) curve, which is the running sum of

the weighted enrichment score obtained from GSEA software, while the normalised

enrichment score (NES) and the corresponding FDR (false discovery rate) value are

reported within each graph.

Figure 5 Supplement 1: (A) Representative images of A549 FUCCI expressing

cells treated with (Cisplatin) or without (Control) 5 µg/ml cisplatin for 2 h and then

followed by time lapse microscopy. Images were taken every 20 min for 3 days.
Coloured arrows indicate specific mother and daughter cells followed through time.
(B) Quantification of cell cycle phase length from data generated in B and C.
Statistical significance was determined by one-way ANOVA with Tukey correction for
multiple comparisons (****$p<0.0001$, n.s = not significant). **(C)** The total number of
divisions completed by biological repeats of asynchronously growing FUCCI
expressing A549 cells pulsed with or without (control) with 5 µg/ml cisplatin for 2 h.
Data was extracted from time-lapse imaging analysis. Statistical significance was
determined by one-way ANOVA with Tukey correction for multiple comparisons
(****$p<0.0001$, ***$p<0.001$, **$p<0.01$, ns = not significant) and an unpaired two-tailed
Student's t-test for summary data (****$p<0.0001$).

1003

Figure 5 Supplement 2: (A) The total number of divisions completed by
asynchronously growing FUCCI expressing A549 (data from Figure 5 Supplement
1C) and NCI-H1299 cells treated without (control) or with 5 µg/ml cisplatin for 2 h.
Data was extracted from time-lapse imaging analysis. Statistical significance was
determined by one-way ANOVA with Tukey correction for multiple comparisons
(****$p<0.0001$, ns = not significant). **(B)** The individual fate maps of NCI-H1299 cells
(n=50 control, 100=cisplatin) from A, tracked by time lapse microscopy, with images
taken every 30 min for 72 h. **(C)** Quantification of cell fate outcomes from B;
including G1 arrest before mitosis (G1 ABM), G1 arrest after mitosis (G1 AAM),
death before mitosis (DBM) and death after mitosis (DAM) and Proliferative (Prolif.),
i.e. cells that divided 2 or more times. **(D)** Violin plots of the number of divisions
completed by cisplatin treated NCI-H1299 cells based on the cell cycle phase at time
of cisplatin exposure. Statistical significance was determined by one-way ANOVA
with Tukey correction for multiple comparisons (ns = not significant). **(E)** Additional
analysis of data from (Hastings et al., 2020) showing the total number of divisions
completed by (n=50) asynchronously growing FUCCI expressing A549 cells treated
with either scramble (siControl) or one of 2 separate siRNAs targeting p53. Statistical
significance was determined by one-way ANOVA with Tukey correction for multiple
comparisons (****$p<0.0001$, ***$p<0.001$, **$p<0.01$, ns = not significant).

1023

1024   **Figure 5 Supplement 3:** Similar to Figure 5D-F, quantification of the proportion of
1025   red (G1), yellow (G1/S) and green (S/G2-M) cells found in tumours (n=3) from day 0

to 7 of **(A)** NCI-H1573 and **(B)** NCI-H1299 placed under optical windows in mice treated with or without carboplatin.

Figure 7 Supplement 1: (A) Quantification of the size and number of PCNA and 53BP1 foci for additional cells as per Figure 7A and 7B. **(B)** Number of 53BP1 foci in cells from early S and mid/late S phase (n=4), were temporally aligned based on entry into and exit from the first mitosis. **(C)** Representative images of the first mitosis observed in cells that were in mid S-phase at the time of cisplatin exposure. White arrow heads indicate micronuclei (mn) in daughter cells, cf= cytokinesis failure. Scale bar = 10 μm.

Figure 8 Supplement 1: (A) Representative cell proliferation curves +/- SD of FUCCI expressing A549 cells treated with 5 μM of the RAD51 inhibitor IR-1 for 1 h prior to exposure for 2 h with 5 μg/ml cisplatin. Nuclear counts were determined from 5 independent fields of time lapse imaging, using the StarDist plugin for ImageJ/Fiji. **(B)** Cell fate maps for (n=100) of cells from A. **(C)** Quantification of cell fate outcomes from B; including G1 arrest before mitosis (G1 ABM), G1 arrest after mitosis (G1 AAM), death before mitosis (DBM) and death after mitosis (DAM) and Proliferative (Prolif.), i.e. cells that divided 2 or more times. **(D)** Violin plots of the number of divisions of cells from B based on the cell cycle phase at time of cisplatin exposure. Statistical significance was determined by one-way ANOVA with Tukey correction for multiple comparisons (*$p<0.05$, **$p<0.01$, ****$p<0.0001$, ns = not significant).

References:

Anders S, Huber W. 2010. Differential expression analysis for sequence count data. *Genome Biol* **11**:R106. doi:10.1186/gb-2010-11-10-r106

Andersson A, Fagerberg J, Lewensohn R, Ehrsson H. 1996. Pharmacokinetics of Cisplatin and Its Monohydrated Complex in Humans. *J Pharm Sci* **85**:824–827. doi:10.1021/js960037a

Bakhoum SF, Silkworth WT, Nardi IK, Nicholson JM, Compton DA, Cimini D. 2014. The mitotic origin of chromosomal instability. *Current biology : CB* **24**:R148 R149. doi:10.1016/j.cub.2014.01.019

1060 Bankhead P, Loughrey MB, Fernández JA, Dombrowski Y, McArt DG, Dunne PD,
1061  McQuaid S, Gray RT, Murray LJ, Coleman HG, James JA, Salto-Tellez M,
1062  Hamilton PW. 2017. QuPath: Open source software for digital pathology image
1063  analysis. *Sci Rep-uk* **7**:16878. doi:10.1038/s41598-017-17204-5

1064 Barr AR, Cooper S, Heldt FS, Butera F, Stoy H, Mansfeld J, Novák B, Bakal C. 2017.
1065  DNA damage during S-phase mediates the proliferation-quiescence decision in
1066  the subsequent G1 via p21 expression. *Nat Commun* **8**:ncomms14728.
1067  doi:10.1038/ncomms14728

1068 Baus F, Gire V, Fisher D, Piette J, Dulić V. 2003. Permanent cell cycle exit in G2
1069  phase after DNA damage in normal human fibroblasts. *The EMBO Journal*
1070  **22**:3992 4002. doi:10.1093/emboj/cdg387

1071 Burgess A, Lorca T, Castro A. 2012. Quantitative live imaging of endogenous DNA
1072  replication in mammalian cells. *Plos One* **7**:e45726.
1073  doi:10.1371/journal.pone.0045726

1074 Burgess A, Rasouli M, Rogers S. 2014. Stressing Mitosis to Death. *Frontiers Oncol*
1075  **4**:140. doi:10.3389/fonc.2014.00140

1076 Caldon CE, Burgess A. 2019. Label free, quantitative single-cell fate tracking of time-
1077  lapse movies. *Methodsx* **6**:2468–2475. doi:10.1016/j.mex.2019.10.014

1078 Chan YW, Fugger K, West SC. 2017. Unresolved recombination intermediates lead
1079  to ultra-fine anaphase bridges, chromosome breaks and aberrations. *Nat Cell Biol*
1080  **20**:92–103. doi:10.1038/s41556-017-0011-1

1081 Chan YW, West SC. 2018. A new class of ultrafine anaphase bridges generated by
1082  homologous recombination. *Cell Cycle* **17**:1–9.
1083  doi:10.1080/15384101.2018.1515555

1084 Charrasse S, Gharbi-Ayachi A, Burgess A, Vera J, Hached K, Raynaud P, Schwob
1085  E, Lorca T, Castro A. 2017. Ensa controls S-phase length by modulating Treslin
1086  levels. *Nat Commun* **8**:206. doi:10.1038/s41467-017-00339-4

1087 Danecek P, McCarthy SA. 2017. BCFtools/csq: haplotype-aware variant
1088  consequences. *Bioinformatics* **33**:btx100. doi:10.1093/bioinformatics/btx100

1089 Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson
1090  M, Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*
1091  **29**:15–21. doi:10.1093/bioinformatics/bts635

1092 Enoiu M, Jiricny J, Schärer OD. 2012. Repair of cisplatin-induced DNA interstrand
1093  crosslinks by a replication-independent pathway involving transcription-coupled
1094  repair and translesion synthesis. *Nucleic Acids Res* **40**:8953–64.
1095  doi:10.1093/nar/gks670

Feringa FM, Raaijmakers JA, Hadders MA, Vaarting C, Macurek L, Heitink L,
Krenning L, Medema RH. 2018. Persistent repair intermediates induce
senescence. *Nat Commun* **9**:3923. doi:10.1038/s41467-018-06308-9

Gligorijevic B, Kedrin D, Segall JE, Condeelis J, Rheenen J van. 2009. Dendra2
Photoswitching through the Mammary Imaging Window. *J Vis Exp Jove* 1278.
doi:10.3791/1278

Gonzalez-Rajal A, Hastings JF, Watkins DN, Croucher DR, Burgess A. 2020.
Breathing New Life into the Mechanisms of Platinum Resistance in Lung
Adenocarcinoma. *Frontiers Cell Dev Biology* **8**:305. doi:10.3389/fcell.2020.00305

Hastings JF, Rajal AG, Latham SL, Han JZ, McCloy RA, O'Donnell YE,
Phimmachanh M, Murphy AD, Nagrial A, Daneshvar D, Chin V, Watkins DN,
Burgess A, Croucher DR. 2020. Analysis of pulsed cisplatin signalling dynamics
identifies effectors of resistance in lung adenocarcinoma. *Elife* **9**:e53367.
doi:10.7554/elife.53367

Hayashi MT, Karlseder J. 2013. DNA damage associated with mitosis and
cytokinesis failure. *Oncogene* **32**:4593–4601. doi:10.1038/onc.2012.615

Herbst RS, Morgensztern D, Boshoff C. 2018. The biology and management of non-
small cell lung cancer. *Nature* **553**:446–454. doi:10.1038/nature25183

Hsu C-H, Altschuler SJ, Wu LF. 2019. Patterns of Early p21 Dynamics Determine
Proliferation-Senescence Cell Fate after Chemotherapy. *Cell* **178**:361-373.e12.
doi:10.1016/j.cell.2019.05.041

Johnson TG, Schelch K, Lai K, Marzec KA, Kennerson M, Grusch M, Reid G,
Burgess A. 2020. YB-1 Knockdown Inhibits the Proliferation of Mesothelioma
Cells through Multiple Mechanisms. *Cancers* **12**:2285.
doi:10.3390/cancers12082285

Karanam K, Kafri R, Loewer A, Lahav G. 2012. Quantitative live cell imaging reveals
a gradual shift between DNA repair mechanisms and a maximal use of HR in mid
S phase. *Mol Cell* **47**:320–9. doi:10.1016/j.molcel.2012.05.052

Kawabe S, Roth JA, Wilson DR, Meyn RE. 2000. Adenovirus-mediated p16INK4a
gene expression radiosensitizes non-small cell lung cancer cells in a p53-
dependent manner. *Oncogene* **19**:5359–5366. doi:10.1038/sj.onc.1203935

Kedrin D, Gligorijevic B, Wyckoff J, Verkhusha VV, Condeelis J, Segall JE, Rheenen
J van. 2008. Intravital imaging of metastatic behavior through a mammary
imaging window. *Nat Methods* **5**:1019–1021. doi:10.1038/nmeth.1269

Kelland L. 2007. The resurgence of platinum-based cancer chemotherapy. *Nat Rev*
Cancer **7**:573–584. doi:10.1038/nrc2167

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**:550.
doi:10.1186/s13059-014-0550-8

Luong KV, Wang L, Roberts BJ, Wahl JK, Peng A. 2016. Cell fate determination in
cisplatin resistance and chemosensitization. *Oncotarget* **7**:23383–23394.
doi:10.18632/oncotarget.8110

Mamdani H, Jalal SI. 2016. DNA repair in lung cancer: potential not yet reached.
Lung Cancer Management **5**:5–8. doi:10.2217/lmt-2016-0004

Marini KD, Croucher DR, McCloy RA, Vaghjiani V, Gonzalez-Rajal A, Hastings JF,
Chin V, Szczepny A, Kostyrko K, Marquez C, Jayasekara WSN, Alamgeer M,
Boolell V, Han JZR, Waugh T, Lee HC, Oakes SR, Kumar B, Harrison CA,
Hedger MP, Lorensuhewa N, Kita B, Barrow R, Robinson BW, Kretser DM de, Wu
J, Ganju V, Sweet-Cordero EA, Burgess A, Martelotto LG, Rossello FJ, Cain JE,
Watkins DN. 2018. Inhibition of activin signaling in lung adenocarcinoma
increases the therapeutic index of platinum chemotherapy. *Sci Transl Med*
10:eaat3504. doi:10.1126/scitranslmed.aat3504

McCloy RA, Rogers S, Caldon CE, Lorca T, Castro A, Burgess A. 2014. Partial
inhibition of Cdk1 in G 2 phase overrides the SAC and decouples mitotic events.
Cell Cycle Georget Tex **13**:1400–12. doi:10.4161/cc.28401

Mootha VK, Lindgren CM, Eriksson K-F, Subramanian A, Sihag S, Lehar J,
Puigserver P, Carlsson E, Ridderstråle M, Laurila E, Houstis N, Daly MJ,
Patterson N, Mesirov JP, Golub TR, Tamayo P, Spiegelman B, Lander ES,
Hirschhorn JN, Altshuler D, Groop LC. 2003. PGC-1α-responsive genes involved
in oxidative phosphorylation are coordinately downregulated in human diabetes.
Nat Genet **34**:267–273. doi:10.1038/ng1180

Murai J, Pommier Y. 2018. PARP Trapping Beyond Homologous Recombination and
Platinum Sensitivity in Cancers. *Annu Rev Cancer Biology* **3**:1–20.
doi:10.1146/annurev-cancerbio-030518-055914

Nobis M, Herrmann D, Warren SC, Kadir S, Leung W, Killen M, Magenau A,
Stevenson D, Lucas MC, Reischmann N, Vennin C, Conway JRW, Boulghourjian
A, Zaratzian A, Law AM, Gallego-Ortega D, Ormandy CJ, Walters SN, Grey ST,
Bailey J, Chtanova T, Quinn JMW, Baldock PA, Croucher PI, Schwarz JP,
Mrowinska A, Zhang L, Herzog H, Masedunskas A, Hardeman EC, Gunning PW,
Monte-Nieto G del, Harvey RP, Samuel MS, Pajic M, McGhee EJ, Johnsson A-
KE, Sansom OJ, Welch HCE, Morton JP, Strathdee D, Anderson KI, Timpson P.
2017. A RhoA-FRET Biosensor Mouse for Intravital Imaging in Normal Tissue
Homeostasis and Disease Contexts. *Cell Reports* **21**:274–288.
doi:10.1016/j.celrep.2017.09.022

Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. 2015.
StringTie enables improved reconstruction of a transcriptome from RNA-seq
reads. *Nat Biotechnol* **33**:290–295. doi:10.1038/nbt.3122

Replogle JM, Zhou W, Amaro AE, McFarland JM, Villalobos-Ortiz M, Ryan J, Letai A, Yilmaz O, Sheltzer J, Lippard SJ, Ben-David U, Amon A. 2020. Aneuploidy increases resistance to chemotherapeutics by antagonizing cell division. *Proc National Acad Sci* **117**:30566–30576. doi:10.1073/pnas.2009506117

Ritsma L, Steller EJA, Ellenbroek SIJ, Kranenburg O, Rinkes IHMB, Rheenen J van. 2013. Surgical implantation of an abdominal imaging window for intravital microscopy. *Nat Protoc* **8**:583–594. doi:10.1038/nprot.2013.026

Rogers S, McCloy RA, Parker BL, Gallego-Ortega D, Law AMK, Chin VT, Conway JRW, Fey D, Millar EKA, O'Toole S, Deng N, Swarbrick A, Chastain PD, Cesare AJ, Timpson P, Caldon CE, Croucher DR, James DE, Watkins DN, Burgess A. 2018. MASTL overexpression promotes chromosome instability and metastasis in breast cancer. *Oncogene* **37**:4518–4533. doi:10.1038/s41388-018-0295-z

Sakaue-Sawano A, Kurokawa H, Morimura T, Hanyu A, Hama H, Osawa H, Kashiwagi S, Fukami K, Miyata T, Miyoshi H, Imamura T, Ogawa M, Masai H, Miyawaki A. 2008. Visualizing Spatiotemporal Dynamics of Multicellular Cell-Cycle Progression. *Cell* **132**:487–498. doi:10.1016/j.cell.2007.12.033

Saldivar JC, Hamperl S, Bocek MJ, Chung M, Bass TE, Cisneros-Soberanis F, Samejima K, Xie L, Paulson JR, Earnshaw WC, Cortez D, Meyer T, Cimprich KA. 2018. An intrinsic S/G 2checkpoint enforced by ATR. *Science* **361**:806 810. doi:10.1126/science.aap9346

Salvador-Barbero B, Álvarez-Fernández M, Zapatero-Solana E, Bakkali AE, Menéndez M del C, López-Casas PP, Domenico TD, Xie T, VanArsdale T, Shields DJ, Hidalgo M, Malumbres M. 2020. CDK4/6 Inhibitors Impair Recovery from Cytotoxic Chemotherapy in Pancreatic Adenocarcinoma. *Cancer Cell* **37**:340-353.e6. doi:10.1016/j.ccell.2020.01.007

Sansregret L, Vanhaesebroeck B, Swanton C. 2018. Determinants and clinical implications of chromosomal instability in cancer. *Nature Reviews Clinical Oncology* **85**:14. doi:10.1038/nrclinonc.2017.198

Sarlós K, Biebricher A, Petermann EJG, Wuite GJL, Hickson ID. 2017. Knotty Problems during Mitosis: Mechanistic Insight into the Processing of Ultrafine DNA Bridges in Anaphase. *Cold Spring Harb Sym* **82**:187–195. doi:10.1101/sqb.2017.82.033647

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**:676. doi:10.1038/nmeth.2019

Schmidt U, Weigert M, Broaddus C, Myers G. 2018. Cell Detection with Star-convex Polygons. *Arxiv* 265–273. doi:10.1007/978-3-030-00934-2_30

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T. 2003. Cytoscape: A Software Environment for

Integrated Models of Biomolecular Interaction Networks. *Genome Res* **13**:2498–
2504. doi:10.1101/gr.1239303

Sherry ST, Ward M-H, Kholodov M, Baker J, Phan L, Smigielski EM, Sirotkin K.
2001. dbSNP: the NCBI database of genetic variation. *Nucleic Acids Res* **29**:308–
311. doi:10.1093/nar/29.1.308

Slyskova J, Sabatella M, Ribeiro-Silva C, Stok C, Theil AF, Vermeulen W, Lans H.
2018. Base and nucleotide excision repair facilitate resolution of platinum drugs-
induced transcription blockage. *Nucleic Acids Res* **46**:9537–9549.
doi:10.1093/nar/gky764

Smogorzewska A. 2019. Fanconi Anemia: A Paradigm for Understanding DNA
Repair During Replication. *Blood* **134**:SCI-32-SCI-32. doi:10.1182/blood-2019-
121229

Spies J, Lukas C, Somyajit K, Rask M-B, Lukas J, Neelsen KJ. 2019. 53BP1 nuclear
bodies enforce replication timing at under-replicated DNA to limit heritable DNA
damage. *Nat Cell Biol* **21**:1–11. doi:10.1038/s41556-019-0293-6

Steigemann P, Wurzenberger C, Schmitz MHA, Held M, Guizetti J, Maar S, Gerlich
DW. 2009. Aurora B-mediated abscission checkpoint protects against
tetraploidization. *Cell* **136**:473 484. doi:10.1016/j.cell.2008.12.020

Stewart DJ. 2007. Mechanisms of resistance to cisplatin and carboplatin. *Crit Rev
Oncol Hemat* **63**:12–31. doi:10.1016/j.critrevonc.2007.02.001

Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA,
Paulovich A, Pomeroy SL, Golub TR, Lander ES, Mesirov JP. 2005. Gene set
enrichment analysis: A knowledge-based approach for interpreting genome-wide
expression profiles. *P Natl Acad Sci Usa* **102**:15545–15550.
doi:10.1073/pnas.0506580102

Swiecicki PL, Durm G, Bellile E, Bhangale A, Brenner JC, Worden FP. 2020. A multi-
center phase II trial evaluating the efficacy of palbociclib in combination with
carboplatin for the treatment of unresectable recurrent or metastatic head and
neck squamous cell carcinoma. *Invest New Drug* **38**:1550–1558.
doi:10.1007/s10637-020-00898-2

Tutt A, Tovey H, Cheang MCU, Kernaghan S, Kilburn L, Gazinska P, Owen J,
Abraham J, Barrett S, Barrett-Lee P, Brown R, Chan S, Dowsett M, Flanagan JM,
Fox L, Grigoriadis A, Gutin A, Harper-Wynne C, Hatton MQ, Hoadley KA, Parikh
J, Parker P, Perou CM, Roylance R, Shah V, Shaw A, Smith IE, Timms KM,
Wardley AM, Wilson G, Gillett C, Lanchbury JS, Ashworth A, Rahman N, Harries
M, Ellis P, Pinder SE, Bliss JM. 2018. Carboplatin in BRCA1/2-mutated and triple-
negative breast cancer BRCAness subgroups: the TNT Trial. *Nat Med* **24**:628–
637. doi:10.1038/s41591-018-0009-7

Urien S, Lokiec F. 2004. Population pharmacokinetics of total and unbound plasma
cisplatin in adult patients. *British Journal of Clinical Pharmacology* **57**:756 763.
doi:10.1111/j.1365-2125.2004.02082.x

Watt LP, Natsum T, Saito Y, Garzón J, Dong Q, Boteva L, Gilber N, Kanemaki MT,
Hiraga S, Donaldson AD. 2020. The RIF1-Long splice variant promotes G1 phase
53BP1 nuclear bodies to protect against replication stress. *eLife*.
doi:10.7554/elife.58020

Weber K, Bartsch U, Stocking C, Fehse B. 2008. A Multicolor Panel of Novel
Lentiviral "Gene Ontology" (LeGO) Vectors for Functional Gene Analysis. *Mol*
Ther **16**:698–706. doi:10.1038/mt.2008.6

Weber K, Thomaschewski M, Benten D, Fehse B. 2012. RGB marking with lentiviral
vectors for multicolor clonal cell tracking. *Nat Protoc* **7**:839–849.
doi:10.1038/nprot.2012.026

Weber K, Thomaschewski M, Warlich M, Volz T, Cornils K, Niebuhr B, Täger M,
Lütgehetmann M, Pollok J-M, Stocking C, Dandri M, Benten D, Fehse B. 2011.
RGB marking facilitates multicolor clonal cell tracking. *Nat Med* **17**:504–509.
doi:10.1038/nm.2338

Yang KS, Kohler RH, Landon M, Giedt R, Weissleder R. 2015. Single cell resolution
in vivo imaging of DNA damage following PARP inhibition. *Sci Rep-uk* **5**:10129.
doi:10.1038/srep10129

Figure 1

Rajal et al 2021

Figure 3

Rajal et al 2021

Figure 5

Rajal et al 2021