Investigations into modifications of neuromuscular physiology by axonal transport disruptions in *Drosophila* SOD1 mutants

Tristan C. D. G. O'Harrow[1]*[†], Atsushi Ueda[1]*, Xiaomin Xing[1]*[‡], Chun-Fang Wu[1]

[1] Department of Biology, University of Iowa, Iowa City, IA 52242, USA

* Equal contribution

[†] Current address: Raymond G. Perelman Center for Cellular and Molecular Therapeutics, The Children's Hospital of Philadelphia, Philadelphia, PA 19104, USA

[‡] Current address: Department of Pharmacology, School of Medicine, University of California, Davis, CA, 95616

ABSTRACT

The reactive oxygen species (ROS)-scavenging enzyme Cu/Zn superoxide dismutase (SOD1) is an evolutionarily conserved mechanism for the maintenance of oxidative homeostasis, and missense mutations in the human SOD1 gene are associated with the motor neuron degenerative disease amyotrophic lateral sclerosis (ALS). Mutations in the *Drosophila melanogaster* SOD1 gene (*Sod*) shorten fly lifespan, attenuate motor function, and induce developmental mortality. We have previously found morphological and physiological defects at the neuromuscular junctions (NMJs) of hypomorphic Sod^{n108} and ALS model Sod^{G85R} mutant larvae. Here, we report genetic interactions causing striking modifications of *Sod* mutant phenotypes by mutations in the gene *Prickle* (*Pk*), which are linked to planar cell polarity, epilepsy, and axonal transport disruptions. *Pk* is expressed as two isoforms *prickle-spiny-legs* (*sple*) and *prickle-prickle* (*pk*), which are each suppressed by specific hypomorphic mutations (*sple1* and *pk1*). Interestingly, *Sod* phenotypes are distinctly modified depending on both the *Pk* isoform suppressed, and whether said suppression is heterozygous or homozygous. Heterozygous *sple1* and *pk1* improved the developmental survival of *Sod* mutants, whereas homozygous *sple1* and *pk1* drastically increased mortality. Further, only heterozygous *sple1* and *pk1* clearly ameliorated morphological defects at the neuromuscular junctions of *Sod* mutant larvae. Pharmacological treatment of *Sod* mutants reveals allele-specific motor neuron terminal hyperexcitability, characterized by synaptic transmissions of

extended duration and abnormal presynaptic Ca^{2+} transients. Heterozygous *sple1* mutation suppresses this *Sod* mutant terminal hyperexcitability, but *pk1* does not. We reversed this suppressive effect of *sple1* by pharmacological blockade of Ca^{2+}-activated K^+ channel *slowpoke*. Altogether, this study builds on our prior knowledge of *Sod* mutant development and physiology to show that axonal transport-linked gene mutations strikingly modify *Sod* mutant phenotypes, providing strong evidence for a role of intracellular transport in alterations of neuromuscular morphology and physiology by SOD1 mutations.

INTRODUCTION

The ROS-scavenging enzyme Cu/Zn superoxide dismutase (SOD1) is highly conserved across phyla, and is expressed in a wide variety of tissues (Uhlén et al., 2015). Consequently, mutations in the gene encoding SOD1 have pervasive consequences for physiology, developmental viability, and organismal longevity (Phillips et al., 1989). Research continues to reveal defects in motor neuron development and physiology associated with mutations in the SOD1 gene, including hypomorphic mutations and mutations linked to familial amyotrophic lateral sclerosis (FALS) (Rosen et al., 1993).

Reactive oxygen species (ROS) are known to behave as signaling molecules to play a wide variety of roles in healthy neuronal development and physiology (Thannickal & Fanburg, 2000; Milton & Sweeney, 2011). During neurodevelopment, ROS modify the biochemistry of cytoskeletal components to influence their dynamics and structure (Wilson & González-Billaut, 2015; Munnamalai & Suter, 2009). The reactive superoxide anion (O_2^-) has also been shown to regulate synaptic plasticity (Klann et al., 1998; Massaad & Klann, 2011). In *Drosophila* larval motor neurons, oxidative stress influences synaptic structure and transmission (Sanya et al., 2002), and regulates activity-dependent neuronal plasticity (Oswald et al., 2018).

Loss of SOD1 function due to genetic mutation allows for oxidative stress in excess of physiological levels. Resulting physiological disruptions include the misfolding of proteins (Berlett & Stadtman, 1997), peroxidation of lipids (Altan et al. 2003), and alteration of nucleic acids (Ali et al., 1996; Aitken & Krausz, 2001). High oxidative stress affects neurodevelopment by disrupting cytoskeletal structures (Landino et al., 2002; Munnamalai & Suter, 2009). SOD1 has been shown to support axon outgrowth and survival, and oxidative

stress induced by loss of SOD1 or treatment with a redox toxin triggers neurodegeneration.

(Fischer & Glass, 2010). Further, SOD1 knockout mice display NMJ neurotransmission

impairments and neuromuscular degeneration (Shi et al., 2014). The *n108* allele (Campbell et

al., 1985) of the *Drosophila* gene for SOD1 (*Sod*) is functionally null, and *n108* mutant larvae

and flies display hypersensitivity to oxidative stress (Phillips et al., 1989). The adults also

exhibit severely reduced lifespans (Phillips et al., 1989) and motor defects (Ruan & Wu,

2008). We have also previously shown that *n108* mutant larvae display altered synaptic

transmission at the neuromuscular junction, and supernumerary axon discharges (Ueda et al.,

manuscript in preparation), and defective neurotransmission in the adult giant fiber escape

circuit (Ueda et al., 2021).

Through mechanisms that remain to be clarified, SOD1 is linked to FALS when

specific amino acid substitutions are present. Mouse, *C. elegans*, and *Drosophila* models

expressing ALS-linked mutant SOD1 exhibit motor dysfunction or lethal paralysis (Wang et

al., 2008; Wang et al., 2009; Watson et al., 2008; Sahin et al., 2017; Agudelo et al., 2020).

Prior to detectable neurodegeneration, ALS mutant SOD1 impacts neurodevelopment and cell

metabolism. ALS mutant SOD1 localizes to actin structures, altering growth cone dynamics

and eventual neuronal morphology (Osking et al., 2019). Misfolded mutant SOD1 protein also

co-localizes with mitochondria in the axons of murine motor neuron, where mitochondrial

morphology is altered (Vande Velde et al., 2011) and mitochondrial Ca^{2+} buffering capacity is

defective (Damiano et al., 2006). The *Drosophila* ALS-linked *Sod* mutant *G85R* displays

defective larval locomotion, and consistently fails to eclose from the pupal stage (Held et al.,

2019; Sahin et al., 2017). *G85R* larvae also display proprioceptor neuron defects and

associated motor circuit dysfunction (Held et al., 2019).

In a separate report, we have characterized *Sod* mutant allele-dependent defects in

NMJ physiology and synaptic morphology (Ueda et al., Manuscript in preparation). In

screening for modifiers of synaptic transmission in *Drosophila Sod* mutants, we fortuitously

found genetic interactions between *Sod* and the planar cell polarity (PCP) and axonal

transport-linked gene *prickle* (*pk*) (Gubb & García-Bellido, 1982; Gray et al., 2011; Lin &

Gubb, 2009; Tree et al., 2002). *pk* is expressed in *Drosophila* in the form of the Pk[Pk] (Prickle

Prickle) and Pk[Sple] (Prickle Spiny-legs) isoforms. Isoform-specific loss-of-function mutants

(*pk1* and *sple1*, Gubb et al., 1999) display distinctly altered microtubule-mediated transport

and microtubule dynamics in larval motor axons, with related changes in neuronal excitability

(Ehaideb et al., 2014; Ehaideb et al., 2016). We have previously found that *pk1* and *sple1*

mutant larvae exhibit alterations in motor neuron development and synaptic transmission that

may be downstream of axon transport modifications (Ueda et al., 2022).

Here, we report the findings that *pk* isoform loss-of-function (LOF) mutations either

improve or reduce the developmental viability of *Sod* mutant *Drosophila* pupae in a genetic

dosage-dependent manner. Similarly, only heterozygous *pk* LOF suppresses NMJ morphology

defects in *Sod* mutants. We also demonstrate that *pk* LOF induces allele-specific alterations

Sod mutant NMJ transmission. Additionally, we show that selective pharmacological

blockade of ion channels reveals NMJ terminal hyperexcitability in Sod^{G85R}, and that this

phenotype is suppressed by partial LOF of the Pk^{Sple} isoform of *pk*. This work therefore

demonstrates that dysregulations of development and neuromuscular physiology induced by

Sod mutation can be altered by *pk* isoform dose manipulation, indicating potential roles of

modified intracellular transport in the generation of *Sod* mutant phenotypes.

MATERIALS & METHODS

<u>Fly stocks</u>

Flies and larvae were housed in bottles containing a cornmeal-agar medium (Frankel

& Brousseau, 1968), supplemented with 3g of yeast paste prepared as a 1:1 mixture of ddH_2O

and active dry yeast (Red Star Yeast Co., LLC). Flies were reared at a room temperature of 22

± 1°C. The *Sod* mutants examined were those of alleles *n108* (Phillips et al., 1989) (gift from

John Phillips, University of Guelph, Ontario, Canada) and *G85R* (Şahin et al. 2017) (gift from

Barry Ganetzky, University of Wisconsin-Madison, USA). The wild-type control strain used

for Figures 1, 2, 3, and 4 was Canton-S (CS). For experiments in which a motor neuron

membrane tag was needed, the stock used carried a chr. II *UAS-CD8-GFP* recombined with a

chr. II motor neuron GAL4 driver (Brand & Perrimon, 1993), either *C164-Gal4* (Torroja et

al., 1999) or *OK371-Gal4* (Mahr and Aberle, 2006). The original recombined *OK371-Gal4*

UAS-CD8-GFP (*OK371::CD8-GFP*) was obtained from the National Centre for Biological

Sciences (Bangalore, Karnataka, India). *Sod* lines used to produce the data represented in

Figure 1 were balanced over a TM6 balancer with a dominant *Tubby* (*Tb*) marker (TM6, *Tb*).

The Pk^{Pk} and Pk^{Sple} lines used in this study have been documented previously (Ehaideb et al.,

2014). The wild-type control strain used for Figure 5 was the Oregon-R (OR) used in Ehaideb

et al. (2014). All *Pk;Sod* double-mutant lines were generated in the Wu lab by conventional

genetic crosses.

<u>Developmental lethality studies</u>

129   Housing and medium conditions for the developmental lethality study (Figure 1) were

identical to those described in the main "Fly stocks" section. Flies including 20-40 females

were used to seed each bottle. Counts of deaths and eclosions were performed 16-17 d. after

seeding. The TM6, *Tb* balancer carried by *Sod* heterozygous wandering 3rd instar larvae and

pupae was used to distinguish them from *Sod* mutant homozygotes. Dead larvae appear as

darkened corpses on the wall of the housing bottle. Dead pupae were entirely black, or

contained a mixture of dark and empty patches, as confirmed by transmitted light under a

dissection scope. Juvenile flies dead during eclosion were distinguished from flies in the

process of eclosing by a second observation 15-30 min after the first. Juvenile fly corpses

were also frequently darker than their live counterparts.

<u>Larval preparations, vital staining, & immunohistochemistry</u>

140   Wandering third instar larvae were dissected, as per the "fillet preparation method", in

HL 3.1 saline (Feng et al., 2004). HL3.1 used in immunohistochemistry or vital staining

experiments contained 0.1 mM Ca^{2+}. Tetramethylrhodamine (TMRM, $C_{25}H_{25}ClN_2O_7$) (Floryk

& Houštěk, 1999) vital staining of larval preps for mitochondria was done at 100 nM in HL

3.1 for 20 minutes (in a light-impermeable chamber to prevent TMRM photobleaching), after

which the bath was replaced with TMRM-free HL 3.1 before microscopy. Before antibody

staining, larval preps were fixed in 3.7% formaldehyde in HL 3.1 for 25-30 minutes. Fixed

preps were incubated with an FITC-conjugated goat antibody to HRP at 1:50 in phosphate

buffer saline, at 4°C for 12-48 h. Anti-HRP (Jan & Jan, 1982) recognizes a neural

carbohydrate antigen in *Drosophila* (Kurosaka et al., 1991). For double-staining with anti-

HRP and anti-Dlg (monoclonal AB 4F3: Parnas et al., 2001) or anti-Futsch (monoclonal AB

22C10: Fujita et al., 1982; Zipursky et al., 1984) staining, fixed larval preps were incubated

with the primary monoclonal AB at 1:100 in phosphate buffer saline, at 4°C for 12-48 h,

rinsed, and incubated in phosphate buffer saline containing anti-HRP(conj. FITC) and a

secondary TRITC-conjugated anti-mouse IgG antibody, both at 1:50. The 4F3 and 22C10

antibodies were obtained from the Developmental Studies Hybridoma Bank (DSHB)

(University of Iowa, Iowa City, IA, USA). The FITC-conjugated anti-HRP and TRITC-

conjugated anti-mouse IgG were obtained from Jackson ImmunoResearch Laboratories (West

Grove, PA, USA). Larval NMJ images, vital and fixed, were gathered on a Leica DM IL LED

inverted microscope (Leica Microsystems Inc., Buffalo Grove, IL, USA), using the Leica

Application Suite X software.

<u>Larval NMJ morphology and mitochondrial distributions</u>

Examination of *Sod* single-mutant NMJ morphology (Figure 2) was carried out in

muscles 4 (M4) and 13 (M13), in abdominal segments A3, A4, and A5, and quantifications

from the different segments were pooled. Examination of *Pk;Sod* double-mutant NMJ

morphology (Figure 4) was carried out in muscle 4 (M4) in abdominal segments A2, A4, and

A6, and quantifications from the different segments were pooled. Counts of boutons from M4

include only type Ib boutons. Ib and Is boutons and branches in M13 were quantified

separately. Counts of specific bouton morphologies ("unsegregated" boutons and large

terminal boutons) are from M4 (Ib). For branch counting, a branch was defined as a terminal

process carrying at least two boutons (Zhong et al., 1992). A bouton was described as

incompletely segregated ("unsegregated") if the width of the narrowest point on the neck

preceding it was greater than 80% of the width of the widest point on the bouton. If multiple

unsegregated boutons appeared in a row, it could be difficult count them. In these cases,

counts were conservative. A terminal bouton was determined to be the "largest" if it was

measurably larger than any other bouton on that branch, past the last branching point.

<u>Electrophysiology</u>

All electrophysiological recordings were based on published protocols, and performed

in HL3.1 saline (Feng et al., 2004) (Composition (mM): 70 NaCl, 5 KCl, 0.1 $CaCl_2$, 4 $MgCl_2$,

10 $NaHCO_3$, 5 trehalose, 115 sucrose, 5 HEPES) (pH adjusted to 7.1). Intracellular recording

of excitatory junction potentials (EJPs) was adapted from Ueda and Wu (2006). Recording of

extracellular focal EJPs (efEJPs) was shown in Ueda and Wu (2009) and Xing and Wu

(2018a,b). For nerve-evoked action potentials, 0.1 ms stimuli were applied to the cut end of

the segmental nerve through a suction pipette. To enable analysis of presynaptic terminal

excitability, we also performed electrotonic stimulation (Wu et al., 1978; Ganetzky & Wu,

1982; 1983). Briefly: Tetrodotoxin (TTX) was applied to block Na+ channels. To achieve

direct electrotonic stimulation of the terminal, a 1 ms stimulus was applied to the cut

segmental nerve, which was sucked into the stimulation electrode, nearly up to the entry point

into the muscle, so as to allow the stimulus to passively propagate to the terminal and

depolarize the endplate membrane (Lee et al., 2014). In this manner, CaV channels

were directly triggered by local depolarization, independent from invasion of axonal Na+

action potentials. Further, a significant portion of multiple K+ channel species, including

Shaker (Kv1), were blocked by co-application of 4-aminopyridine (4-AP) and

tetraethylammonium (TEA) (Jan & Jan, 1977; Singh & Wu, 1999). Electrotonic stimulation

of the larval NMJ with TTX, 4-AP, and TEA has been shown in Lee et al. (2014). Such

manipulations allow for the examination of terminal excitability driven by Ca2+ channels

with reduced repolarization regulation by a range of K+ channels, with the exclusion of

axonal potential interplay.

Statistics

Total mortalities per genotype (Figure 1) were compared using Fisher's exact test.

Counts of NMJ boutons and branches and of enlarged terminal and unsegregated boutons

(Figures 2 and 3) were examined with Kruskal-Wallis testing between all groups within

muscle # and parameter (eg. M4, boutons), followed by rank-sum (Mann-Whitney U) post

hoc between individual groups. EJP sizes were compared with one-way ANOVA (Figure 4B).

Plateau-like EJP durations were compared by F-test (Figure 4D). Counts of NMJ

mitochondrial macropunctae (Figure 6) were compared with Kruskal-Wallis testing followed

by rank-sum (Mann-Whitney U) post hoc.

RESULTS

Lethality of *Sod* mutants during different developmental stages

Lifespan defects in adult *Drosophila* homozygous for hypomorphic *Sod* allele *n108*

(Ruan & Wu 2008) and pupal lethality in homozygotes of *n108* and the ALS-linked point-

mutant allele *G85R* (Şahin et al., 2017) have been previously characterized. We further noted

that *n108* and *G85R* mutants exhibit lethality in different phases of the pupal stage (Figure

1B). While the majority of *n108* death occurred at eclosion from the pupal stage (Figure 1A,

lower-right image), over 90% of *G85R* individuals died during pupal development (Figure

1A, middle-right image), unable to develop to the eclosion stage. Both *Sod* mutants also

displayed mild levels larval lethality (Figure 1A, upper-right image), similar between the

alleles. By comparison, *Drosophila* of control strains CS and *w1118* exhibited only very small

levels of developmental lethality that were greatly exceeded by the significant lethality in the

Sod mutants ($p < 0.001$). Heterozygous *Sod* mutants (*n108*/+ and *G85R*/+) did not differ in

lethality from the control strains (Supplemental Table 1), indicating a lack of phenotypic

dominance of *Sod* mutant lethality.

We found that the developmental lethality of *n108* and *G85R* mutants was modified by

mutations in *prickle*. Two copies of either the *pk1* or the *sple1* allele conferred full

developmental lethality to *n108* mutants, but unlike the lethality profile of the *n108* single-

mutant that generally died at eclosion, the *pk1*;*n108* and *sple1*;*n108* double-mutants died

during pupal development. *pk1*;*G85R* and *sple1*;*G85R* double-mutants remained as fully

lethal as the *G85R* single mutant, although rate of progression to eclosion was marginally

increased. Crucially, one copy of either *pk* mutant allele had the opposite effect: *Sod* mutant

lethality was suppressed. One copy of *pk1* (*pk1*/+;*n108*) lowered the overall rate of *n108*

lethality below 50%. A copy of *sple1* (*sple1*/+;*n108*) was even more effective, with lethality

reduced below 30%. Single copies of *pk1* and *sple1* (*pk1*/+;*G85R* and *sple1*/+;*G85R*)

conferred to *G85R* mutants rates of survival to adulthood of approx. 10% and 14%

respectively. These small proportions are notable when compared to the complete pre-

eclosion lethality of the *G85R* single mutant.

<u>Altered larval neuromuscular junction morphology</u>

The *Sod* mutants displayed mild alterations in NMJ outgrowth, that were readily visible in

counts of boutons per NMJ (Figure 2B). Counts of Ib branches and boutons in M4 and M13 in

G85R yielded distributions that were not significantly different from WT. While the increase

in *G85R* M13 Is branches was nonsignificant, the numbers of Is boutons were significantly

increased compared to WT ($p < 0.05$). NMJ outgrowth was increased in *n108* compared to

WT, with more more Ib boutons in M4 ($p < 0.05$) and M13 ($p < 0.001$), more Ib branches on

M13 ($p < 0.01$), and a visible but nonsignificant increase in numbers of M4 Ib branches

(Figure 2B). Where *n108* NMJ complexity was significantly greater than WT, it was also

significantly increased compared to *G85R*, indicating physiological differences between

mutants with the loss-of-function allele and those with the ALS-linked allele.

Both *Sod* mutants displayed increases in numbers of unsegregated boutons per M4 Ib branch

compared to WT (*G85R*: $p < 0.001$, *n108*: $p < 0.05$) (Figure 2C). Similarly, terminal boutons

that were enlarged compared to other boutons on the same M4 Ib branch appeared more

frequently in both *Sod* mutants than in WT (*G85R*: $p < 0.001$, *n108*: $p < 0.01$) (Figure 2C).

Unlike the *Sod* single-mutants, phenotypes of NMJ complexity (M4, Ib) in *Pk^Pk^*, *Pk^Sple^*, and

their double mutants with *G85R* were most visible in changes to the numbers of branches per

NMJ. Compared to WT, *Pk^Pk^*;*G85R* displayed significantly more boutons ($p < 0.01$), and

*Pk^Sple^*;*G85R* displayed significantly fewer ($p < 0.01$). Both double-mutants displayed fewer

branches than either corresponding *Pk* single-mutant (both $p < 0.001$), and *Pk^Sple^*;*G85R*

displayed a number of boutons similar to the *G85R* single-mutant, strikingly fewer than the

*Pk^Sple^* single-mutant. ($p < 0.001$). Both *Pk* single-mutants displayed slightly more boutons

than WT (both $p < 0.05$). None of the *Pk* single-mutants or *Pk*;*G85R* double-mutants

displayed a number of unsegregated boutons that was significantly different from WT, nor

was there a clear difference between WT and the *Pk* single-mutants in the rate at which

terminal boutons were enlarged. However, both *Pk*;*G85R* double-mutants displayed enlarged

terminal boutons at rates lower than *G85R*, but still greater than WT (both $p < 0.01$).

<u>Genetic interaction between *Sod* and *Pk* in motor neuron excitability and synaptic</u>

<u>transmission</u>

In light of the NMJ morphological defects manifested in *Sod* through its interaction with *Pk*

mutations, we examined neuromuscular transmission for related defects. When the motor

axon bundle was stimulated with 0.1 ms pulses in HL3.1 saline containing 0.2 mM Ca^{2+}, we

detected reliable EJP responses in *Sod* and *Pk* mutant larvae. In *Sod* mutants (hypomorphic

alleles *n108* and *x39*/*n108*, as well as the ALS-related *G85R* allele), EJP response upon nerve

stimulation was similar to that in WT ($p > 0.05$, Figure 4B). EJP size in the *Pk* alleles of *Pk*

(*Pk1* and *Pk30*, data combined in Figure 4) was comparable to WT, while EJP size was

significantly decreased in the mutants of the *Sple* allele ($p<0.05$, One-Way ANOVA).

However, the double-mutant *Pk^Pk1^*;*Sod^G85R^* showed unusually large EJPs (Figure 4B, $p < 0.05$,

One-Way ANOVA). Abnormal synaptic transmission can be attributed to alterations in ion

channels participating in regulation of presynaptic terminal membrane depolarization to

control Ca^{2+} influx that triggers release of synaptic vesicles from presynaptic terminals.

<u>Sod^{G85R} allele-specific motor terminal hyperexcitability</u>

Sod mutant axonal excitability having been studied (Figure 3) (Ueda et al., In

preparation), we examined the isolated electrophysiological properties of the *Sod* mutant

larval NMJ using a previously established protocol of direct electrotonic stimulation of the

axon terminal, after silencing the axon Na+ spikes via TTX (Ganetzky & Wu, 1982, 1983;

Ueda & Wu 2009; Lee et al., 2013). Varying intensities of direct electrotonic stimulation of

the presynaptic terminal revealed no clear qualitative differences between *G85R*, *n108*, and

WT in loose-patch focal recording (see methods) of efEJP (extracellular focal excitatory

junction potentials) from single synaptic boutons (Figure 4A). Exposure of the larval

preparation to the *Shaker* K_V channel blocker 4-aminopyridine (4-AP, 200 nM) induced NMJ

hyperexcitability in *G85R* larvae that was typically absent under the same condition in *n108*

and WT. The *Shaker* blockade-induced hyperexcitability of *G85R* manifested in the form of a

shoulder-like extension of the efEJP (Figure 4B). We quantified this characteristic with a

measurement of the duration at 20% amplitude beyond the baseline (Figure 6). A

presynaptically expressed Ca^{2+} indicator (GCaMP6f) also communicated the state of Ca^{2+}

currents in the recorded synaptic boutons during evoked neurotransmission, which revealed

that *G85R* also exhibited brief presynaptic Ca^{2+} currents in response to electrotonic stimuli in

the presence of 4-AP. In fact, we found that every extended efEJP was accompanied by a

spike in presynaptic GCaMP (Figure 4B, Figure 6). Given this consistent association between

augmented presynaptic Ca^{2+} currents and modified postsynaptic response, we suspect that the

prolonged efEJP is elicited by prolonged neurotransmitter release from the presynaptic

terminal.

<u>pk^{sple1} suppression of Sod^{G85R} hyperexcitability</u>

The *sple1* null allele of *prickle* exerted a striking effect on the NMJ physiology of

G85R: One copy of *sple1* was sufficient to suppress both the efEJP extension and the

simultaneous presynaptic Ca^{2+} current amplification (Figure 5B, Figure 6). That the control

sple1/+ mutant larvae exhibit mild hyperexcitability in response to 4-AP reveals that the

genetic interaction between *G85R* and *sple1* is not simply the sum of opposing phenotypes.

Further, the suppressive interaction of *sple1/+* with *G85R* physiology is *pk* allele-dependent,

and does not arise from the combination of the *pk1* allele with *G85R* (Figure 5B, Figure 6).

DISCUSSION

Our study contrasts the loss-of-function *n108* allele of *Sod* with the ALS-linked *G85R*

allele, by demonstrating clear differences in the developmental stages at which the mutants of

each allele die. The contrast is further enhanced by differences between their neuromuscular

synapse morphologies. We found potential mechanistic links between abnormal distribution

of mitochondria in Sod^{G85R} and synaptic bouton enlargement (Ueda et al., Manuscript in

preparation), consistent with the striking interaction with *Pk*, known to affect axonal transport.

Further, following pharmacological blockade of Na+ and K+ currents in our

electrophysiological experiments, the remaining Ca2+ currents appeared to initiate more

physiological actions in Sod^{G85R}, in a manner suppressed by *sple1* mutation.

<u>Developmental death in *Sod* mutants</u>

This study of *Drosophila Sod* mutant lethality extends specificity and aspects of

mutational effects on the developmental stages in which deaths occur. Specifically, we

distinguish between death of the fully developed juvenile fly during eclosion from the pupal

casing, and death mid-pupal development, which is characterized by a pupal casing that

remains closed. As a result, we established allele-dependent developmental lethality

phenotypes. The *n108* lethality occurs largely during eclosion and may be due to physical

weakness of the juvenile. The vast majority of *G85R* death occurs during pupal development,

often prior to formation of adult morphology, as seen by a lack of red adult eyes visible

through the pupal casing of late-stage pupae (Bainbridge & Bownes, 1981).

<u>Aberrant NMJ morphology and physiology</u>

Overgrowth of the Sod^{n108} NMJ has been documented previously, and another

oxidative stress-sensitive mutant (*spinster*) was shown to have similar overgrowth. Further,

the two genes show synergistic interactions (Milton et al., 2011). The *G85R* larval NMJ has

also been characterized (Held et al., 2019) although the bouton number did not seem to differ

significantly from their control. In this study, we also found no clear difference in bouton

numbers, but striking bouton morphological alterations could be demonstrated in *G85R* larvae

(Figure 2A). Additional immunostaining of proteins constituting the presynaptic machinery,

such as synaptotagmin and dynamin (Estes et al., 1996), or postsynaptic elements such as

glutamate receptors (Harris & Littleton, 2015), could elucidate the functional relevance of the

unusual bouton morphologies in *Sod* mutants.

Mutant SOD1 is the most commonly identifiable risk factor in human patients with

familial ALS (Zarei et al., 2015). By exploiting the amenability of the *Drosophila* larval

neuromuscular preparation to the study of motor synapses at an early developmental stage, we

detect clear signs of synaptic physiology and morphology alterations in *Sod* mutants.

Additionally, abnormal distributions of synaptic mitochondria in *Sod,* and severe NMJ

phenotypes in *Sod* double-mutants with transport-affecting *Pk* mutations, suggest

dysfunctional transport may contribute to early SOD1-linked pathology. Our work therefore

adds to our understanding of phenotypes and mechanisms that will facilitate further studies to

uncover critical events leading to irreversible and terminal motor neuron degeneration in

ALS.

References

Aitken RJ, Krausz C. (2001) Oxidative stress, DNA damage, and the Y chromosome.
Reproduction 122: 497-506.

Akbergenova Y, Bykhovskaia M. (2007) Synapsin maintains the reserve vesicle pool and
spatial segregation of the recycling pool in *Drosophila* presynaptic boutons. *Brain*
Research 1178: 52-64.

Ali S, Jain SK, Abdulla M, Athar M. (1996) Paraquat induced DNA damage by reactive
oxygen species. *Biochemistry and Molecular Biology International* 39(1): 63-67.

Altan O, Pabuccuoglu A, Altan A, Konyalioglu S, Bayraktar H. (2003) Effect of heat stress
on oxidative stress, lipid peroxidation, and some stress parameters in broilers. *British*
Poultry Science 44(4): 545-550.

Bainbridge SP, Bownes M. (1981) Staging the metamorphosis of *Drosophila* melanogaster.
J. Embryol. exp. Morph. 66: 57-80.

Bassuk AG, Wallace RH, Buhr A, ..., Slusarski DC, Berkovic SF, El-Shanti HI. (2008) A
homozygous mutation in human PRICKLE1 causes an autosomal-recessive
progressive myoclonus epilepsy-ataxia syndrome. *The American Journal of Human*
Genetics 83: 572-581.

Berlett BS, Stadtman ER. (1997) Protein Oxidation in Aging, Disease, and Oxidative Stress.
The Journal of Biological Chemistry 272(33): 20313-20316.

Bilsland LG, Sahai E, Kelly G, Golding M, Greensmith L, Schiavo G. (2010) Deficits in
axonal transport precede ALS symptoms in vivo. *Proc Natl Acad Sci* 107(47):
20523-20528.

Brand AH, Perrimon N. (1993) Targeted gene expression as a means of altering cell fates
and generating dominant phenotypes. *Development* 118:401-415.

Campbell SH, Hilliker AJ, Phillips JP. (1985) Cytogenetic analysis of the *cSOD* microregion in *Drosophila melanogaster*. *Genetics* 112: 205-215.

Ehaideb SN, Iyengar A, Ueda A, Iacobucci GJ, Cranston C, Bassuk AG, Gubb D, Axelrod JD, Gunawardena S, Wu CF, Manak RJ. (2014) *prickle* modulates microtubule polarity and axonal transport to ameliorate seizures in flies. *Proc. Natl. Acad. Sci. USA* 111(30): 11187-11192.

Ehaideb SN, Wignall EA, Kasuya J, Evans WH, Iyengar A, Koerselman HL, Lilienthal AJ, Bassuk AG, Kitamoto T, Manak RJ. (2016) Mutation of orthologous prickle genes causes a similar epilepsy syndrome in flies and humans. *Annals of Clinical and Translational Neurology* 3 (9): 695-707.

Estes PS, Roos J, van der Bliek A, Kelly RB, Krishnan KS, Ramaswami M. (1996) Traffic of Dynamin within individual *Drosophila* synaptic boutons relative to compartment-specific markers *J. Neurosci.* 1996 16(17): 5443-5456.

Feng Y, Ueda A, Wu CF. (2004) A modified minimal hemolymph-like solution, HL3. 1, for physiological recordings at the neuromuscular junctions of normal and mutant Drosophila larvae. *Journal of Neurogenetics*, 18(2): 377-402.

Frankel, A., & Brousseau, G. (1968). Drosophila medium that does not require dried yeast. *Drosophila Information Service* 43, 184.

Ganetzky B, Wu CF. (1982) *Drosophila* mutants with opposing effects on nerve excitability: Genetic and spatial interactions in repetitive firing. *Journal of Neurophysiology* 47(3): 501-514.

Ganetzky B, Wu CF. (1983) Neurogenetic analysis of potassium currents in *Drosophila*: Synergistic effects on neuromuscular transmission in double mutants. *Journal of Neurogenetics* 1: 17-28.

Gubb D, García-Bellido A. (1982) A genetic analysis of the determination of cuticular polarity during development in *Drosophila melanogaster*. *J. Embryol. exp. Morph.* 68: 337-57.

Gubb D, Green C, Huen D, Coulson D, Johnson G, Tree D, Collier S, Roote J. (1999) The balance between isoforms of the Prickle LIM domain protein is critical for planar polarity in *Drosophila* imaginal discs. *Genes & Development* 13: 2315-2327.

Gray RS, Roszko I, Solnica-Krezel L. (2011) Planar cell polarity: coordinating morphogenetic cell behaviors with embryonic polarity. *Developmental Cell* 21: July 19, 2011.

Harris KP, Littleton T. (2015) Transmission, development, and plasticity of synapses. *Genetics* 201: 345-375.

Held A, Major P, Sahin A, Reenan RA, Lipscombe D, Wharton KA. (2019) Circuit dysfunction in SOD1-ALS model first detected in sensory feedback prior to motor neuron degeneration is alleviated by BMP signaling. *Neurobiology of Disease* 39(12): 2347-2364.

Jan LY, Jan YN. (1976a) Properties of the larval neuromuscular junction in *Drosophila melanogaster*. *J. Physiol.* 262: 189-214.

Jan LY, Jan YN. (1976b) L-glutamate as an excitatory transmitter at the *Drosophila* larval neuromuscular junction. *J. Physiol.* 262: 215-236.

Jan YN, Jan LY, Dennis MJ. (1977) Two mutations of synaptic transmission in *Drosophila*. *Proc. R. Soc. Lond. B.* 198: 87-108.

Jan LY, Jan YN. (1982) Antibodies to horseradish peroxidase as specific neuronal markers
in *Drosophila* and in grasshopper embryos. *Proc. Natl. Acad. Sci. USA* 79: 2700-
2704.
Katz B, Miledi R. (1967a) A study of synaptic transmission in the absence of nerve
impulses. *J. Physiol.* 192: 407-436.
Katz B, Miledi R. (1967b) Tetrodotoxin and neuromuscular transmission. *Proceedings of*
the Royal Society B 167 (1006): 8-22.
Klagges BRE, Heimbeck G, Godenschwege TA, Hofbauer A, Pflugfelder GO, Reifegerste
R, Reisch D, Schaupp M, Buchner S, Buchner E. (1996) Invertebrate synapsins: A
single gene codes for several isoforms in *Drosophila*. *J. Neurosci.* 16(10): 3154-
3165.
Kurdyak P, Atwood HL, Stewart BA, Wu CF. (1994) Differential physiology and
morphology of motor axons to ventral longitudinal muscles in larval *Drosophila. The*
Journal of Comparative Neurology 350: 463-472.
Kurosaka A, Yano A, Itoh N, Kuroda Y, Nakagawa T, Kawasaki T. (1991) The structure of
a neural specific carbohydrate epitope of horseradish peroxidase antiserum. *the*
Journal of Biological Chemistry 266(7): 4168-4172.
Lee J, Ueda A, Wu CF (2014) Distinct roles of *Drosophila cacophony* and *Dmca1D* Ca2+
channels in synaptic homeostasis: Genetic interactions with *slowpoke* Ca2+-activated
BK channels in presynaptic excitability and postsynaptic response. *Dev. Neurobiol.*
141: 4548-4557.
Lee J, Wu CF. (2010) Orchestration of Stepwise Synaptic Growth by K+ and Ca2+
Channels in *Drosophila*. *J. Neurosci.* 30(47): 15921-15833.
Lin YY, Gubb D. (2009) Molecular dissection of *Drosophila* Prickle isoforms distinguishes
their essential and overlapping roles in planar cell polarity. *Developmental Biology*
325: 386-399.
Mahr A, Aberle H. (2006) The expression pattern of the *Drosophila* vesicular glutamate
transporter: A marker protein for motoneurons and glutamatergic centers in the brain.
Gene Expression Patterns 6: 299-309.
Massaad CA, Klann E. (2011) Reactive Oxygen Species in the Regulation of Synaptic
Plasticity and Memory. *Antioxidants and Redox Signaling* 14(10): 2013-2054.
Mei X, Wu S, Bassuk AG, Slusarski DC. (2013) Mechanisms of *prickle1a* function in
zebrafish epilepsy and retinal neurogenesis. *Disease Models & Mechanisms* 6: 679-
688.
Milton VJ, Jarrett HE, Gowers K, Chalak S, Briggs L, Robinson IM, Sweeney ST. (2011)
Oxidative stress induces overgrowth of the *Drosophila* neuromuscular junction. *Proc*
Natl Acad Sci 108(42): 17521-17526.
Osking Z, Ayers JI, Hildebrandt R, Skruber K, Brown H, Ryu D, ... Vitriol EA. (2019) ALS-
linked SOD1 mutants enhance neurite outgrowth and branching in adult motor
neurons. *iScience* 11: 294-304.
Paemka L, Mahajan VB, Skeie JM, ..., Carvill GL, Manak RJ, Bassuk AG. (2013)
PRICKLE1 interaction with SYNAPSIN I reveals a role in autism spectrum
disorders. *PLOS ONE* 8 (12): e80737.
Parnas D, Haghighi P, Fetter RD, Kim SW, Goodman CS. (2001) Regulation of
postsynaptic structure and protein localization by the rho-type guanine nucleotide
exchange factor dPix. *Neuron* 32: 415-424.

Phillips JP, Campbell SD, Michaud D, Charbonneau M, Hilliker, AJ. (1989) Null mutation of copper/zinc superoxide dismutase in Drosophila confers hypersensitivity to paraquat and reduced longevity. *Proc Natl Acad Sci* 86(8): 2761-2765.

Rosen DR, Siddique T, Patterson D, Figlewicz DA, Sapp P, Hentati A, ... Brown RH Jr. (1993) Mutations in Cu/Zn superoxide dismutase gene are associated with familial amyotrophic lateral sclerosis. *Nature* 362: 59-62.

Ruan, H., & Wu, C. F. (2008). Social interaction-mediated lifespan extension of *Drosophila* Cu/Zn superoxide dismutase mutants. *Proc Natl Acad Sci* 105(21): 7506-7510.

Şahin A, Held A, Bredvik K, Major P, Achilli TM, Kerson, AG, ... Reenan, R. (2017). Human SOD1 ALS mutations in a *Drosophila* knock-in model cause severe phenotypes and reveal dosage-sensitive gain- and loss-of-function components. *Genetics* 205(2): 707-723.

Singh S, Wu CF. (1999) Ionic currents in larval muscles of *Drosophila. International Review of Neurobiology* 43: 191-220.

Stewart BA, Atwood HL, Renger JJ, Wang J, Wu CF. (1994) Improved stability of *Drosophila* larval neuromuscular preparations in haemolymph-like physiological solutions. *J Comp Physiol A* 175: 179-191.

Thannickal VJ, Fanburg BL. (2000) Reactive oxygen species in cell signaling. *Am J Physiol Lung Cell Mol Physiol* 279: L1005-L1028.

Torroja L, Packard M, Gorczyca M, White K, Budnik V. (1999) The *Drosophila* β-amyloid precursor protein homolog promotes synapse differentiation at the neuromuscular junction. *The Journal of Neuroscience* 19(18): 7793-7803.

Tree DRP, Shulman JM, Rousset R, Scott MP, Gubb D, Axelrod JD. (2002) Prickle mediates feedback amplification to generate asymmetric planar cell polarity signaling. *Cell* 109: 371-381.

Ueda A, Wu CF. (2006) Distinct frequency-dependent regulation of nerve terminal excitability and synaptic transmission by I_A and I_K potassium channels revealed by *Drosophila Shaker* and *Shab* mutations. *The Journal of Neuroscience* 26(23): 6238-6248.

Ueda A, Wu CF. (2009) Role of *rut* adenylyl cyclase in the ensemble regulation of presynaptic terminal excitability: Reduced synaptic strength and precision in a *Drosophila* memory mutant. *J Neurogenet* 23: 185-199.

Vande Velde C, McDonald KK, Boukhedimi Y, ... Cleveland DW. (2011) Misfolded SOD1 Associated with Motor Neuron Mitochondria Alters Mitochondrial Shape and Distribution Prior to Clinical Onset. *PLoS ONE* 6(7): e22031.

Vasin A, Zueva L, Torrez C, Volfson A, Littleton JT, Bykhovskaia M. (2014) Synapsin regulates activity-dependent outgrowth of synaptic boutons at the *Drosophila* neuromuscular junction. *J. Neurosci.* 34(32): 10554-10563.

Wang J, Farr GW, Zeiss CJ, Rodriguez-Gil DJ, Wilson JH, ... Horwich AL. (2008) Progressive aggregation despite chaperone associations of a mutant SOD1-YFP in transgenic mice that develop ALS. *Proc Natl Acad Sci* 106(5): 1392-1897.

Wang J, Farr GW, Hall DH, Li F, Furtak K, Dreier L, ... Horwich AL. (2009) An ALS-Linked Mutant SOD1 Produces a Locomotor Defect Associated with Aggregation and Synaptic Dysfunction When Expressed in Neurons of Caenorhabditis elegans. *PLoS Genetics* 5(1): e1000350.

514 Xing X, Wu CF. (2018a) Unraveling synaptic GCaMP signals: Differential excitability and
515  clearance mechanisms underlying distinct Ca^{2+} dynamics in tonic and phasic
516  excitatory, and aminergic modulatory motor terminals in *Drosophila*. *eNeuro* 5(1):
517  e0362-17.2018 1-28
518 Xing X, Wu CF. (2018b) Inter-relationships among physical dimensions, distal–proximal
519  rank orders, and basal GCaMP fluorescence levels in Ca^{2+} imaging of functionally
520  distinct synaptic boutons at *Drosophila* neuromuscular junctions. *J Neurogenet*
521  32(3): 195-208.
522 Zarei S, Carr K, Reiley L, Diaz K, Guerra O, Fernandez Altamirano P, ... Chinea A. (2015)
523  A comprehensive review of amyotrophic lateral sclerosis. *Surgical Neurology*
524  *International* 6:171.
525 Zhong Y, Budnik V, Wu CF. (1992) Synaptic plasticity in *Drosophila* memory and
526  hyperexcitable mutants: role of cAMP cascade. *J Neurosci* 12(2): 644-651.
527 Zipursky SL, Venkatesh TR, Teplow DB, Benzer S. (1984) Neuronal development in the
528  *Drosophila* retina: Monoclonal antibodies as molecular probes. *Cell* 36: 15-26.

Figure 1. Developmental lethality in *Sod* mutants is modified by *pk* mutations. (A) Representative images of live and dead larvae and pupae, an empty pupal casing left after successful eclosion of the juvenile fly, and the carcass of a juvenile dead during eclosion. (B) *Sod* mutant survival during development is improved by one copy of either *sple1* or *pk1*. Bars show percentages of individuals dead at the 3rd instar larval stage (black), during pupal development (gray), and at eclosion from the pupal casing (white). Genotypes shown are *Sod* mutants *G85R* and *n108*, *pk* mutants *pk1* and *sple1*, *pk1/+* and *sple1/+* heterozygotes, and double-mutants of those *Sod* and *pk* alleles. Both wild-type CS and *w1118* are included as controls due to differing genetic backgrounds of *G85R* and *n108*. *n* indicated below each bar. ****P* < 0.001, *****P* < 0.0001, Fisher's exact test. Statistics are for cumulative developmental mortality.

Figure 2. Heterozygous *prickle* loss-of-function suppresses Sod^{G85R} neuromuscular junction morphology defects. (A) Representative images of type Ib NMJs from muscle 4 in wild-type (WT), Sod^{G85R} (*G85R*), heterozygous and homozygous double-mutants of *sple1* with *G85R* (*sple1/+;G85R* and *sple1;G85R*), and heterozygous and homozygous double-mutants of *pk1* with *G85R* (*pk1/+;G85R* and *pk1;G85R*). Immunostaining shows presynaptic morphology (Anti-HRP). White arrows indicate *Sod* phenotypic enlarged distal boutons. Scale bar = 10 μm. (B) Percentage of distal synaptic boutons that are enlarged in each genotype (see methods). (C) Numbers of presynaptic boutons (see methods). (D) Numbers of presynaptic branches (see methods). Bars in B, C and D indicate mean, error bars indicates SEM. *N* of NMJ = WT: 46, *G85R*: 34, *sple1/+;G85R*: 23, *sple1;G85R*: 11, *pk1/+;G85R*: 13, *pk1;G85R*: 12. **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$, one-way ANOVA followed by Tukey's test.

Figure 3. Homozygous *pk* mutations alter the size of excitatory junctional potentials (EJPs) from body wall muscles in *Sod* mutant larvae. Two different WT controls (CS and OR) are shown. EJPs were measured in HL3.1 saline containing 0.2 mM Ca^{2+}.

sod \ pk	+	pk / +	pk	sple / +	sple
+	OR 6.9 ± 1.2 (16) CS 11.9 ± 1.4 (22)	3.6 ± 0.7 (8)	14.8 ± 2.8 (12)	5.6 ± 0.9 (5)	6.4 ± 1.6 (16)
n108	15.3 ± 3.3 (7)	ND	22.0 ± 6.4 (5)	ND	8.0 ± 2.1 (6)
G85R	19.5 ± 3.4 (9)	ND	28.3 ± 4.3 (9)	24.6 ± 2.2 (6)	23.2 ± 2.5 (10)

Table 1. Muscle EJP size (mean ± SEM) in *Sod, pk,* and double mutant combinations. Wild-type CS is the control strain for *Sod* and wild-type OR is the control for *pk*. N of NMJs are indicated in parentheses. EJPs were measured in HL3.1 saline containing 0.2 mM Ca^{2+}.

Figure 4. Heterozygous *sple1* suppresses the NMJ terminal hyperexcitability of Sod^{G85R}. (A) efEJP of *sple1/+;G85R* and *pk1/+;G85R* (with TTX alone, see methods) are qualitatively similar to *G85R* and WT. Stimuli 1 ms in duration were delivered at 1 Hz, increasing in intensity from 1× to 2× threshold voltage. GCaMP (top, green) fluorescence remains at baseline during stimulation in all genotypes. Gray scale bar between the GCaMP and efEJP traces indicates time scale, time of stimulus (1/second), and stimulus intensity for the GCaMP and top efEJP trace. Expanded efEJP traces are responses at 0, 1, 3, and 5 s. (B) Presynaptic GCaMP spikes and efEJP extension are absent in *sple1/+;G85R*, indicating that one copy of the *sple1* mutant allele suppresses *Sh* blockade-induced terminal hyperexcitability of *G85R*. Retention of 4-AP-induced *G85R* hyperexcitability characteristics (GCaMP spikes and extended efEJP) in *pk1/+;G85R* demonstrates that the interaction of *sple1* with *G85R* is *prickle* allele-dependent. Arrowheads indicate efEJP "shoulder" characteristic in *G85R* and *pk1/+;G85R*.

Figure 5. Quantification of presynaptic GCaMP signal amplitude and efEJP duration in *G85R* and *pk*/+;*G85R* double-mutants. (A) Mean GCaMP signal amplitudes (peak ΔF/F) in response to 1 Hz stimuli, of intensity from 1× to 2× threshold voltage. Error bars indicate SEM. In the presence of TTX alone, GCaMP signals were always undetectable (no deviation from baseline). Thin dark green line indicates baseline. 4-AP induced large single stimulus-evoked GCaMP spikes (light green bars) in *G85R* and *pk1*/+;*G85R* larvae. WT and *sple1*/+;*G85R* larvae typically lacked these spikes. See GCaMP traces in Figure 6. (B) Mean durations (ms) of the efEJP co-occurring with the GCaMP signals quantified above. efEJP traces were measured at 20% amplitude (see methods). WT and *sple1*/+;*G85R* larvae displayed efEJP of similar duration in the absence (dark gray bars) or presence (light gray bars) of 4-AP. *G85R* and *pk1*/+;*G85R* efEJP under 4-AP were typically about twice the duration of efEJP without 4-AP. The efEJPs of extended duration were those displaying the shoulder-like characteristic indicated in Figure 6.

Figure 6. Blockade of *slowpoke* BK channels in addition to *Shaker* channels produces extreme motor terminal hyperexcitability. (A) GCaMP and efEJP traces from the TTX + 4-AP condition for WT, *G85R, sple1*/+;*G85R*, and *pk1*/+;*G85R* are shown for comparison to panel B. Expanded efEJP traces are responses at 1, 3 s. (See Figure 6 for additional and more thorough representation.) (B) Blockade of *Sh* and *slo* K^+ channels, by 4-AP and paxilline respectively, produces extreme terminal hyperexcitability that shares characteristics with the hyperexcitability observed in *G85R* and *pk1*/+;*G85R* under the influence of 4-AP alone. Paxilline-induced terminal hyperexcitability is qualitatively similar across all genotypes examined.

Figure 7. Extended NMJ transmission in *G85R* broadens whole-cell EJPs in the postsynaptic muscle, and heterozygous *sple1* ameliorates this phenotype. Double-recording of whole-cell EJPs and extracellular macropatch EJPs induced by electrotonic depolarization (●) of nerve terminals in the presence of TTX. Application of 4-AP increased the size of EJPs in all genotype. Note that sod^{G85R} exhibited broadened whole-cell EJPs associated with a shoulder of macropatch EJPs (arrow), while WT and $pk^{sple}/+;sod^{G85R}$ lacked them. Application of paxilline induced broadened whole-cell EJPs plus a shoulder of macropatch EJPs (arrow) in all genotypes. 1, 2, 3 indicates sub-threshold, near threshold, and supra-threshold (x3 ~ x5) stimulation. Macropatch EJP traces are normalized.