**Investigations into modifications of neuromuscular physiology by axonal transport disruptions in *Drosophila* SOD1 mutants**

Tristan C. D. G. O'Harrow[1]*, Atsushi Ueda[1]*, Xiaomin Xing[1]*‡, Chun-Fang Wu[1]

[1] Department of Biology, University of Iowa, Iowa City, IA 52242, USA

* Equal contribution

‡ Current address: Department of Pharmacology, School of Medicine, University of California, Davis, CA, 95616

**Abstract**

Cu/Zn superoxide dismutase (SOD1) is a cytoplasmic antioxidant enzyme, which, when mutant in humans, is linked to familial cases of the motor neurodegenerative disease amyotrophic lateral sclerosis (ALS). The *Drosophila* SOD1 gene (*Sod*) shares a highly conserved sequence with the human homolog, and this study includes examinations of the established hypomorphic *n108* allele (*Sod^{n108}*), alongside a knock-in construct of the *G85R* allele found in human ALS patients (*Sod^{G85R}*). In addition to previously documented decreased adult lifespan and attenuated motor function, we show that *Sod* mutant *Drosophila* display high mortality rates during larval and pupal development. Immunostaining of neuronal membrane at neuromuscular synapses in *Sod* mutant larvae revealed presynaptic terminals of abnormal, swollen morphology. In *Sod^{G85R}* larvae, *in vivo* mitochondrial staining demonstrated the presence of aggregated mitochondria inside the swollen synaptic terminals, and a genetically encoded GFP construct targeted to mitochondria revealed aggregates of mitochondria inside the axon bundles of *Sod^{G85R}* larvae. In whole-cell recordings of neuromuscular transmission, both *Sod^{n108}* and *Sod^{G85R}* exhibited

lower muscle input resistance and smaller miniature excitatory junction potentials (mEJPs)

compared to WT. However, evoked EJPs were similar to those of WT. Focal

electrophysiological recording showed that both $Sod^{n108}$ and $Sod^{G85R}$ NMJ

terminals displayed slightly higher release probability than WT terminals. Treatment of

$Sod^{G85R}$ with the *Shaker* channel (Kv1) blocker 4-aminopyridine (4-AP) and the broad-

spectrum $K^+$ channel blocker tetraethylammonium (TEA) induced prolonged "plateau-like"

potentials at the larval NMJ upon electrical stimulation. Altogether, this study provides a

snapshot of the alterations in mitochondrial distribution, synaptic morphology, and

neurotransmission that characterize the motor neurons of *Sod* mutants prior to

neurodegeneration and death of the organism.

**Introduction**

The generation of reactive oxygen species (ROS) is a ubiquitous feature of energy

metabolism in living cells. ROS are a class of oxygen-containing molecules that readily

oxidize other molecules upon contact, and as a result can cause a significant amount of

damage to cellular components over time (Lushchak, 2014). To mitigate this effect, a broad

variety of eukaryotic and prokaryotic organisms produce antioxidant enzymes to maintain

oxidative homeostasis (Matés et al., 1999). While ROS are produced endogenously as

byproducts of metabolic processes, including mitochondrial oxidative phosphorylation, they

can also be generated from exogenous sources like toxins, drugs, and pollutants (Chen et

al., 2003; Ali et al., 1996; Block & Calderón-Garcidueñas, 2009). The disruption of oxidative

homeostasis that occurs as a result of increased ROS load or a decrease in antioxidant

activity is termed 'oxidative stress' (Sies, 1997).

The accumulation of oxidative damage due to oxidative stress over time is considered a normal part of aging in macroorganisms, and ROS have been implicated as signaling molecules in healthy physiological processes ranging from cell differentiation to synaptic plasticity (Thannickal & Fanburg, 2000; Massaad & Klann, 2011). However, under increased oxidative stress, cellular proteins are more prone to misfolding, lipids to peroxidation, and nucleic acids to breakage and mutation (Berlett & Stadtman, 1997; Altan et al. 2003; Aitken & Krausz, 2001). Possibly as a result of these effects, increased oxidative stress has been implicated in neurodegenerative diseases, cardiovascular diseases, and cancer (Liguori et al., 2018; Uttara et al., 2009; Dhalla et al., 2000; Sosa et al.,2013).

Molecular species considered ROS include both free radicals such as the superoxide ($O_2^-$), the peroxide $O_2^{-2}$, and the hydroxyl $OH^-$, and molecules without unpaired valence electrons, such as hydrogen peroxide ($H_2O_2$) and the hydroxyl OH (Lushchak, 2014). Cells express a variety of antioxidant enzymes to neutralize these ROS and their derivatives. These enzymes include catalase, glutathione peroxidases, and superoxide dismutases (Matés et al., 1999). Evidence suggests that production of antioxidant enzymes may be generally downregulated in aged organisms, so the contributions of oxidative stress to the aging process may include not just oxidative damage accumulated over the lifespan, but also damage from increased oxidative stress later in the lifespan (Baek et al, 2016).

The antioxidant enzyme Cu/Zn superoxide dismutase (SOD1) behaves as a protein homodimer that mainly localizes to the cytosol, and specifically targets superoxide molecules ($O_2^-$) for neutralization (Perry et al., 2010; Culotta & Daly, 2013; Mondola et al., 2016). As SOD1 interacts with $O_2^-$ , the enzyme alternates between reducing $Cu^{2+}$ to oxidize $O_2^-$ to molecular oxygen ($O_2$), and oxidizing $Cu^+$ and donating protons to reduce $O_2^-$ to $H_2O_2$

(Mondola et al., 2016; Perry et al., 2010). $H_2O_2$ produced by this process is then converted by

the enzyme catalase into molecular oxygen and water ($O_2$ & $H_2O$) (Mondola et al., 2016).

The gene encoding SOD1 (*Sod*) is highly conserved across a wide range of species,

including humans, mice, the worm *Caenorhabditis elegans* (*C. elegans*), and the fruit fly

*Drosophila melanogaster*. The functionally null *n108* allele of the *Drosophila* gene for SOD1

(*Sod*) was generated by exposure of *Drosophila* to the mutagen ethyl methanesulfonate

(Campbell et al., 1985). Flies homozygous for the *n108* allele display hypersensitivity to

paraquat ($Pq^{2+}$), a toxin that induces acute oxidative stress (Phillips et al., 1989). All $Sod^{n108}$

flies in a cohort died when exposed to an aqueous solution containing 5 mM $Pq^{2+}$ for 48 hrs,

whereas over 80% of control flies survived in identical conditions (Phillips et al., 1989). Even

without exposure to toxins, $Sod^{n108}$ confers a reduced adult lifespan, with fewer than 5% of

$Sod^{n108}$ flies living longer than 20 d., compared to lifespans longer than 50 d. for over 70% of

flies in control cohorts.

In contrast, *C. elegans* that lack activity not just of their homolog of SOD1, but lack the

activity of all five SOD genes present in wild-type *C. elegans*, have a wild-type lifespan under

normal conditions, though they display increased sensitivity to pharmacologically-induced

oxidative stress (Van Raamsdonk & Hekimi, 2012). Mice null for their SOD1 gene have a

mean lifespan ~87% as long as that of their wild-type counterparts, a phenotype less striking

than the *Drosophila* lifespan phenotype (Zhang et al., 2016). The disparities between the

percent decreases in lifespan of SOD1 null mutants of different species suggest that, while

principal enzymatic functions appear to be evolutionarily conserved, the relative importance

of SOD1 in general organismal physiology or additional functions of the protein may be more

divergent. One such additional SOD1 function has been observed in yeast, where, under

elevated oxidative stress, the protein relocates to the nucleus. Association of SOD1 with

promoters in the yeast nucleus suggests that it regulates expression of 'oxidative response'

genes (Tsang et al., 2015). Some human patients with the neurodegenerative disease

amyotrophic lateral sclerosis (ALS), popularly known as Lou Gehrig's disease or motor

neuron disease, have been found to carry mutations in the gene for SOD1, and can exhibit

decreased overall SOD1 activity (Rosen et al., 1993; Saccon et al., 2013).

ALS is a fatal neurodegenerative disease in which the gradual death of upper motor neurons

in the cortex and brainstem and of lower motor neurons in the spinal cord causes the

muscles they innervate to weaken and atrophy (Rosen et al., 1993; Rowland & Shneider,

2001). As a result, patients progressively lose the ability to walk, swallow, and breathe, with

an average life expectancy of three to five years after diagnosis (Galvin et al., 2017;

Hardiman et al, 2011). Approximately 2.7 in 100,000 people are affected by ALS across

Europe and North America, with an age of onset most commonly between 50 and 65 years

(Zarei et al., 2015). Precise causes for the development of ALS are still under study, but

research-supported risk factors include specific genetic mutations, as well as cigarette

smoking (Zarei et al., 2015). More controversially proposed risk factors include physical

activity and exposure to agricultural chemicals and electromagnetic fields (Zarei et al., 2015).

Also missing from current research are clear, early signs of ALS progression. The 5-10% of

ALS cases demonstrated to have a heritable genetic component are classified as familial

ALS (FALS), and the remaining 90-95% of cases are classified as sporadic ALS (SALS)

(Zarei et al., 2015). Mutations in the gene for SOD are the most commonly identifiable risk

factor for FALS, and other mutations found in FALS patients include mutations in the genes

for the nucleic acid-binding proteins TAR DNA-binding protein 43 (TDP-43), Fused in

sarcoma (FUS), and angiogenin (Zarei et al., 2015). While ALS is believed to involve

relatively-specific degeneration of motor neurons for reasons that remain unclear, the

disease has at times been observed to appear alongside frontotemporal dementia (FTD),

which involves non-motor neuron degeneration in the frontal and temporal lobes (Mackenzie & Rademakers, 2008).

Due to its central antioxidant function, dysfunction of SOD1 in an organism as a result of mutation results in an increase in oxidative stress. Extremely high oxidative stress can cause enough damage that cells die by necrosis, a disorderly form of cell death that does not follow from distinct signaling pathways (Tang et al., 2013). However, oxidative stress has also been shown to induce cell death by apoptosis, a programmed form of cell death (Kannan & Jain, 2000). Dying motor neurons in *Drosophila* and mouse models of ALS have been shown to express apoptotic markers, making increased oxidative stress as a result of a loss of SOD1 function one possible element of the disease mechanism (Xia et al., 2012; Guegan & Prezedborski, 2003).

Mitochondria have a place in the apoptotic pathway, through the release of proteins from the intermembrane space, and human ALS patients display aberrant mitochondrial morphology and distribution (Ott et al., 2007; Smith et al., 2017). In a mouse model of SOD1-mediated ALS, mitochondria in spinal cord motor neurons displayed more rounded and swollen morphology than the mitochondria of control mice (Vande Velde et al., 2011). The mitochondrial distribution in motor neurons of control mice was relatively homogenous, but the mitochondria in the ALS-model neurons were visibly aggregated in various regions of the cell (Vande Velde et al., 2011). A mouse model has also shown that axonal transport in motor neurons is impaired, which may contribute to abnormal mitochondrial distributions in the cell (Bilsland et al., 2010; De Vos et al., 2007). In addition to their key roles in apoptosis, oxidative homeostasis, and energy metabolism, mitochondria carry out essential regulations of calcium dynamics by uptaking or releasing calcium from the cytosol (Giorgi et al., 2018).

Mice carrying an ALS-linked allele of the SOD1 gene displayed attenuated calcium buffering

capacity, possibly as a result of mitochondrial dysfunction (Beers et al., 2001). The possible

effects of a loss of mitochondrial calcium sequestering ability extend to the broad array of

physiological processes that employ calcium as an intracellular messenger. Regulation of

calcium concentrations is crucial for synaptic transmission, and human patients with ALS

display abnormal presynaptic physiology at motor synapses, with a selective loss of

excitatory amino acid transporter 2 (EAAT2) (Lin et al., 1998). A loss of EAAT2 allows for

increases in extracellular glutamate and overstimulation of glutamate receptors, which in turn

could induce excitotoxic neurodegeneration (Zarei et al., 2015). Key factors in excitotoxicity

include increases in oxidative stress and presynaptic calcium influx, both of which an

organism with mutated SOD1 and dysfunctional mitochondria would have difficulty

compensating for (Salińska et al., 2005). The finding revealing a loss of EAAT2 in ALS

humans has been replicated in a mouse model (Trotti et al., 1999).

When *Sod* carries a missense mutation at or near structural Zn-binding sites, the SOD1

protein can misfold due to structural instability, forming aggregates as a result (Forsberg et

al., 2011). The *Sod* alleles *G85R* and *H71Y*, examined in this study, are two such missense

mutations. Some mutated forms of TDP-43 and FUS have also been shown to form

aggregates, indicating that the formation of protein aggregates may be a shared mechanism

for cases of ALS caused by mutations in a variety of genes (Chen & Cohen, 2019; Patel et

al., 2015; Mackenzie & Rademakers, 2008). Mice carrying YFP-tagged transgenic ALS-

related human alleles of SOD1 displayed motor paralysis phenotypes and died before

reaching a year of age (Wang et al., 2008). Cells in sections of spinal cord from affected

animals contained fluorescent punctae. Fractionation and gel filtration of spinal cord extracts

revealed the presence of high molecular weight SOD1 multimers, heavier than either the

SOD1 monomer or dimer alone, and protein sequencing of the SOD1-containing aggregates

revealed that they contained cytoskeletal proteins, including multiple species of

neurofilament, glial fibrillary acidic protein, and vimentin (Wang et al., 2008).

The fruit fly *Drosophila melanogaster* is a model organism with well-characterized physiology

and a fully sequenced genome (Adams et al., 2000). It is also a diploid organism with just

four pairs of chromosomes, including sex-linked X and Y chromosomes, allowing for

relatively easy genetic characterization. Due to its rapid lifecycle and the high number of

progeny each female produces, it is possible to generate datasets gathered from a large

number of rapidly-aging individuals, for example in studies of ALS. As a result, *Drosophila*

studies of ALS-related mutants can provide us with significant insight into physiological

variability between individual organisms in the context of the many heritable diseases that

express variably between human patients.

As a result of approx. 62% AA homology and the conservation of essential functional sites

between human and *Drosophila* SOD1, the *Drosophila Sod* alleles *G85R* and *H71Y*,

generated by ends-out homologous recombination, are homologous to mutations found in

human familial ALS-linked *Sod* alleles (Şahin et al., 2017). *G85R* and *H71Y* were generated

on the genetic background of $w^{1118}$ (*white*), the white-eyed *Drosophila* mutant. In $Sod^{G85R}$, the

wild-type glycine residue at position 85 has been replaced with an arginine residue, and in

$Sod^{H71Y}$, the wild-type histidine residue at position 71 has been replaced with tyrosine. Both

mutation sites are in close proximity to metal ion binding sites that enable SOD1 to maintain

structural stability. Şahin et al. found that *Drosophila* homozygous for $Sod^{G85R}$ and $Sod^{H71Y}$

displayed locomotor defects as larvae, pupal eclosion defects (full pupal lethality in $Sod^{G85R}$),

and short lifespans in $Sod^{H71Y}$ adults. $Sod^{H71Y}$ adults exhibited locomotor defects, along with

a loss of efferent branches in their leg neuromuscular junctions. In spite of this, no loss of

motor neuron cell bodies in the CNS was observed, suggesting that ALS-related

neurodegeneration at the synapse could precede death of motor neurons (Şahin et al.,

2017).

*Sod* mutation-related neuromuscular defects may therefore also be observable at larval

neuromuscular junctions (NMJs). The *Drosophila* larval fillet preparation, a live dissection

method in which most internal organs are removed, and the muscle wall is laid bare,

provides easy access to several repeating segments of stereotypically arranged NMJs for

vital staining or electrophysiology-based investigations (Lee and Wu, 2010; Zhong et al.

1992; Ueda and Wu, 2006). The larval NMJ has well-documented morphology, which has

been shown to display significant plasticity under the influence of genetic perturbations,

including ion channel mutations and cAMP cascade dysregulation (Lee and Wu, 2010;

Zhong et al., 1992). A readily accessible live NMJ is also ideal for imaging mitochondria

made visible by staining, to investigate for distribution defects similar to those seen in a

mouse ALS model (Vande Velde et al., 2011).

**Materials and Methods**

<u>Fly stocks</u>

Flies and larvae were housed in bottles containing a cornmeal-agar medium (Frankel &

Brousseau, 1968). Flies were reared at a room temperature of $22 \pm 1^{\circ}$C. The *Sod* mutants

examined were those of alleles *n108* (Phillips et al., 1989), *G85R*, and *H71Y* (Şahin et al.

2017). The wild-type control strain used for experiments without genetically encoded labels

was Canton-S. For experiments in which a motor neuron membrane tag was needed, the

stock used carried a chr. II UAS-CD8-GFP recombined with a chr. II motor neuron Gal4

driver (Brand & Perrimon, 1993), either C164-Gal4 (Torroja et al., 1999) or OK371-Gal4

(Mahr and Aberle, 2006). *Sod* lines used to produce the data represented in Figure 1 were

balanced over a TM6 balancer with a dominant *Tubby* (*Tb*) marker.

<u>Developmental lethality studies</u>

Glass housing bottles used for these experiments contained cornmeal-agar medium,

supplemented with 3g of yeast paste prepared as a 1:1 mixture of $ddH_2O$ and active dry

yeast (Red Star Yeast Co., LLC). Flies including 20-40 females were used to seed each

bottle. Counts of deaths and eclosions were performed 16-17 d. after seeding.

<u>Larval preparations, vital staining, & immunohistochemistry</u>

Wandering third instar larvae were dissected, as per the "fillet preparation method", in HL 3.1

saline (Feng et al., 2004). All HL3.1 used in these experiments contained 0.1 mM $Ca^{2+}$.

Tetramethylrhodamine (TMRM, $C_{25}H_{25}ClN_2O_7$) vital staining of larval preps for mitochondria

was done at 100 nM in HL 3.1 for 20 minutes (in a light-impermeable chamber to prevent

TMRM photobleaching), after which the bath was replaced with TMRM-free HL 3.1 before

microscopy. Before antibody staining, larval preps were fixed in 3.7% formaldehyde in HL 3.1

for 25-30 minutes. Fixed preps were incubated with an FITC-conjugated goat antibody to

HRP at 1:50 in phosphate buffer saline, at 4$^o$C for 12-48 h. The antibody was obtained from

Jackson ImmunoResearch Laboratories (West Grove, PA, USA). Anti-HRP recognizes a

neural carbohydrate antigen in *Drosophila* (Kurosaka et al., 1991). Larval NMJ images, vital

and fixed, were gathered on a Leica DM IL LED inverted microscope (Leica Microsystems

Inc., Buffalo Grove, IL, USA), using the Leica Application Suite X software.

<u>Larval NMJ morphology</u>

NMJ boutons and branches were counted from muscles 4 (M4) and 13 (M13), in abdominal

segments A3, A4, and A5. Counts from the three segments were pooled after no clear

differences in the counts from different segments were found (data not shown). Counts of

boutons from M4 include only type Ib boutons, and counts from M13 include both type Ib and

type Is. For branch counting, a branch was defined as a terminal process carrying at least

two boutons (Zhong et al., 1992). An "enlarged terminal bouton" was defined as a bouton at

the very end of a branch that was measurably larger than any other bouton on that branch,

past the last branching point. This was often checked by holding a ruler up to the computer

screen, to measure the width of the terminal bouton and nearby boutons, measured

perpendicular to the directionality of the branch. "Elongated" boutons appear sausage-like,

running down the branch past the point where a normal bouton would have narrowed to the

"neck" that generally separates boutons. They had to also be of width equal to or greater

than the widths of other boutons in the branch with the more common round morphology. If

multiple elongated boutons appeared in a row, it could be difficult count them. In these

cases, counts were conservative.

<u>NMJ mitochondrial distributions</u>

Imaging was completed within 30 minutes after TMRM staining to mitigate effects of post-

dissection physiological decline on results. Boutons described as containing putative

mitochondrial aggregates are those that stained strikingly more brightly than other boutons in

the NMJ, and than the muscle staining in the background of the image. Fluorescence

intensity was not computer-quantified, and so counts of terminal boutons that were especially

brightly stained relied on the observer's judgement, but counts were conservative. Counts of

putative mitochondrial aggregates in enlarged terminal boutons were obtained from larvae in

which the NMJ membrane carried CD8-GFP driven by C164-Gal4 or OK371-Gal4, making

bouton morphology visible in the green fluorescence channel during vital staining in the red

channel.

 Electrophysiology

All electrophysiological recordings were based on published protocols. Excitatory junction

potential recording (EJP) was adapted from Ueda and Wu, 2006 .Electrotonic stimulation

with tetrodoxin, 4-AP and TEA has been shown in Lee et al. (2014) for electrotonic

stimulation. Focal recording method was shown in (Ueda and Wu, 2012) and (Xing and Wu,

2018).

Statistics

Overall distributions of deaths throughout the lifecycle (Figure 1) were compared using

Fisher's exact test. Counts of NMJ boutons and branches and of enlarged terminal and

unsegregated boutons (Figures 2 and 4) were examined with Kruskal-Wallis testing between

all groups within muscle # and parameter (eg. M4, boutons), followed by rank-sum (Mann-

Whitney U) post hoc between individual groups. Distributions of counts of NMJ mitochondrial

aggregations of different categories (No aggregation, aggr. in terminal bouton, aggr. in

terminal + neighboring bouton) (Figure 3) were compared using Fisher's exact test.

**Results**

*Sod* developmental lethality

Wild-type produces no recorded wandering 3rd instar larval death, a low rate of pupal death

at ~3%, and no death during eclosion (Figure 1). In contrast, homozygotes from all three *Sod*

mutants display clear lethality during development. *n108* and *G85R* exhibited wandering 3rd

instar larval death rates of approx. 10% and 2%, respectively. The percent death rate refers

to a percentage of the total number of individuals counted at all stages. Only 4% of

individuals of the functionally-null *n108* allele died during the pupal stage. In the case of

individuals of the ALS-linked *G85R* allele, almost all pupae died before eclosion, though 1

out of 106 individuals made it into eclosion, before dying partway through the process,

leaving the room temperature *G85R* eclosion rate at 0. 44% of *n108* individuals died during

the eclosion process. The successful eclosion rates were 42% of *n108* and 43% of *H71Y*. No

dominant or semi-dominant *Sod* phenotype in the *Sod/TM6, Tb* individuals was apparent

during the stages of life leading up to eclosion. Like wild-type, none of them displayed larval

death or death at eclosion, and none had a pupal death rate above 4% (WT rate: 3%).

<u>Larval neuromuscular junction morphology</u>

When larvae were reared at room temperature, differences in neuromuscular junction

complexity between genotypes were mild in both muscles 4 (M4) and 13 (M13) (Figure 2).

Within genotypes, trends sometimes differed slightly the two muscles. M4 counts included

only type Ib bouton/branches, and M13 counts were from a combination of type Ib and type

Is boutons/branches, so small divergences in morphological quantifications between the two

muscles could reflect contributions from different bouton types. In M4, *n108* and *H71Y* both

displayed mean bouton numbers of approx. 17, compared to the mean of 14 in WT ($p <$

0.05). *G85R* M4 NMJs had on average 12 boutons, a non-significant decrease from WT.

Branch numbers exhibited the same relationships, but mean branch numbers for all

genotypes were between 2.3 and 2.8, with no statistically significant differences. Room

temperature M13 bouton and branch numbers showed mild decreases in *n108* and *G85R*

compared to WT, 13 boutons and 2.6 branches in *n108*, and 14 boutons ($p < 0.05$) and 2.6

branches in *G85R*, compared to 16 and 2.9 in WT.

The most notable NMJ morphology results were those concerning the morphology of

individual boutons. Depending on the allele and temperature conditioning, *Sod* mutants

displayed terminal boutons that appeared enlarged relative to other boutons on the same

branch (Figure 4A), and thick, elongated boutons of morphology that could be described as

"sausage-like" (Figure 4B). NMJs in WT reared at room temperature did occasionally carry

boutons that fit the criteria (see methods) for these morphologies, but at low rates of about

0.5 enlarged terminal boutons and 0.3 elongated boutons per NMJ (Figure 4C). *n108* NMJs

carried both bouton morphology types at higher rates than WT, approx. 0.9 of each type

(compared to WT, enlarged $p < 0.05$, elongated $p < 0.001$) per NMJ. Enlarged terminal

boutons appeared most frequently in *G85R* and *H71Y*, at rates of about 1.4 and 1.5 enlarged

terminal boutons per NMJ ($p < 0.001$ in both genotypes), respectively. Elongated boutons in

occurred at a rate of about 1.1 per NMJ ($p < 0.001$) in *G85R*. *H71Y* deviated from the trend a

bit, producing elongated boutons only at a rate of ~0.6 per NMJ, which was less than *n108*

and *G85R*, but still twice as much WT ($p < 0.05$).

<u>Mitochondrial distributions at the larval NMJ</u>

TMRM staining of the larval NMJ in *Sod*$^{G85R}$ reared at RT revealed brightly stained, putative

mitochondrial aggregations (Figure 3) in over half (55%) of all terminal boutons ($p < 0.001$),

compared to just around 20% in WT (Figure 3). These apparent aggregations are in contrast

to the more diffuse staining seen in most WT NMJs. Unlike *G85R*, the *n108* allele did not

produce many mitochondrial aggregates in NMJ terminal boutons, in fact the rate was a little

lower than WT, at 17%. Very interestingly, rearing the *G85R* larvae under heat stress

actually appeared to provoke the disappearance of mitochondrial aggregations (drop to 14%,

$p < 0.001$ for comparison to *G85R* RT), while in WT, heat stress prompted a mild increase in

aggregates (~30% of terminal boutons), though this change was not statistically significant.

*G85R* reared at RT and WT reared at HT, the two conditions under which the highest rates of

terminal boutons containing aggregates appeared, also displayed the highest rates of

immediately neighboring boutons carrying aggregates. In *G85R* RT, 29% of terminal boutons

containing aggregates were accompanied by neighbors also containing aggregates, and the

rate for WT HT was 33%. This is compared to 15% in WT RT, 14% in *G85R* RT, and 12% in

*n108* RT. If aggregation in both a terminal bouton and one or more neighbors represents a

more severe phenotype than aggregation in the terminal bouton alone, then more frequent

occurrences of the mitochondrial aggregation phenotype are accompanied by increased

severity of those occurrences.

In larvae carrying expressing GFP targeted to motor neuron membrane (UAS-CD8-GFP

under the control of OK371-Gal4 or C164-Gal4), an association between bouton morphology

and the presence of mitochondrial aggregates became apparent (Figure 5C). 83% of

enlarged terminal boutons in *G85R* reared at RT contained aggregates, compared to 55% of

terminal boutons of all morphologies. The aggregation-morphology relationship was also

present and conspicuous in wild-type: although only 15% of all terminal boutons contained

mitochondrial aggregates, 63% of terminal boutons that were enlarged contained

aggregates.

Mutants of sod and pk synergistically cause defects in synaptic transmission and excitability

Seeing the striking growth and structural defects in the synaptic terminals of *Sod* and *Pk*, we

would like to see if these phenotypes indicate any severe physiological symptoms. EJP

recording from postsynaptic muscle showed that *Sod* and *Pk^{Pk}* both exhibited enhanced

transmission compared to WT and *Pk^{Sple}* (Figure 5A and B). Most strikingly, the double

mutant of *Sod* and *Pk* synergistically enhanced the EJP amplitude, suggesting the

importance of *Sod* in *Pk*-mediated transport.

Additional analysis of the presynaptic terminal excitability was performed using electrotonic

stimulation. NaV channels were all inhibited by TTX. Stimulation pulses were delivered at 1

ms duration instead of 0.1, and the stimulation electrode sucked the segmental nerve to

nearly entry point into the muscle, so as to allow the stimulation to passively propagate to the

synaptic terminals and depolarize the endplate membrane (Lee et al., 2014). In this way,

CaV channels were directly triggered to open independent from NaV effects. Last but not

least, a large portion of K+ channels were blocked by co-application of 4-AP and TEA. Such

manipulations allow for the examination of terminal excitability fully driven by Ca2+ and

potentially some less well-known K+ channels, with the axonal effects excluded. Using this

method, we again observed that *Sod* and *Pk* mutants both generated gigantic plateau EJPs

that could last more than 1 s, longer than those from WT. Although not in every case, double

mutant of sod and pk sometimes even produced plateau EJPs that lasted more than 5 or 6 s.

Similar results were obtained in the double mutant $Pk^{Pk}$/+ with $Sod^{n108}$, another allele of *Sod*

mutant by focal recording (Fig 7). The focal recording results confirmed the dominant effects

of pk allele, and indicated that the excitability defect was mainly in type Ib boutons (type Is

not shown). All these results suggest a synergistic effect between sod and pk alleles in

controlling the excitability of presynaptic terminals, which was governed by the local

expression of Ca2+ and K+ channels.

**Discussion**

<u>Developmental death in *Sod* mutants</u>

The developmental data presented in this study deviate a bit from the results produced by

Şahin et al., in that they reported a little under 30% successful eclosion for $Sod^{H71Y}$, as

opposed to our rate of over 40%. One possibility is that the food supplementation with a large

amount of yeast (see methods) induced an increase in successful eclosion. In recent years,

the Wu lab has been in the habit of adding chunks of yeast paste to the cornmeal-agar

medium our *Sod* mutants are reared on, because it seems to improve the numbers and

health of the mutant flies. Future experiments to clarify the basis and magnitude of this effect

will naturally include more control experiments without yeast added, but also studies of *Sod*

developmental health with the addition of yeast extract or heat-killed yeast in the place of our

standard active yeast. This substitution could give indications as to whether any yeast-

feeding effect is due to activity of live yeast, for example in the microbiome of *Drosophila*, in

which case live yeast would be necessary, or to a simple nutritional value provided by the

yeast, in which case yeast extract or heat-killed yeast would be sufficient to induce an effect

on developmental health.

<u>Aberrant NMJ morphology and physiology</u>

The way in which the *n108* allele operates in synergy with heat stress to produce overgrowth,

but does not produce overgrowth alone, is somewhat reminiscent of *Drosophila* mutants for

other genes. The hyperexcitable $K^+$ channels mutants *Shaker* and *ether-a-go-go* each

produce morphologically unremarkable NMJs on their own, but induce overgrowth once

expressed together in the same larvae (Zhong et al., 1992). NMJ overgrowth in the cAMP

phosphodiesterase mutant *dunce* is also magnified by the combination of *dunce* with either

*Shaker* or *ether-a-go-go* (Zhong et al., 1992). Studies of the genetic interactions (or lack

thereof) between $Sod^{n108}$ and any of these mutants will help us understand the mechanisms

that underlie overgrowth in heat-stressed *n108,* as well as the roles of oxidative stress in $K^+$

channel activity and cAMP metabolism.

The neuromuscular junction is, of course, primarily a synaptic structure, but this study stops

short of any examination of synaptic transmission in *Sod* mutants. However, current

observations from the Wu lab indicate that presynaptic failure to release neurotransmitter

upon stimulation in *n108* and *G85R* larvae reared at room temperature is not significantly

different from wild-type (Dr. Xiaomin Xing, unpublished observations). Similarly, it also

appears that *n108* reared at room temp. displays excitatory junction potentials upon nerve

stimulation that are not significantly different from those in WT (Dr. Atsushi Ueda,

unpublished observations). Further information about NMJ function in *Sod* mutants can be

extracted from the system with the use of antibody stains for synaptic machinery. In

particular, the densities and distributions of presynaptic active zones (stained by an antibody

to Bruchpilot) and postsynaptic glutamate receptors (stained with an antibody to one of the

DGluRII subunits) (Lee & Wu, 2010), can give us information on not just the functionality of

the entire NMJ, but also of individual boutons of specific morphologies, including the

enlarged terminal boutons and elongated boutons documented in this study. Staining of the

subsynaptic reticulum, using an antibody to Discs large, will give us further information on

any way in which the postsynaptic area has adapted, or failed to adapt, to boutons of

unusual morphology (Lee & Wu, 2010). Staining may also be the key understanding the

structural basis of the *Sod*-linked enlarged terminal and elongated boutons. It is unclear

whether they are built up with expanded cytoskeletal components, or simply inflated. A

practical and systematic way to approach this issue is to stain for various cytoskeletal

component, such as actin and tubulin, using fluorophore-conjugated phalloidin and antibody

stains. If the boutons are inflated beyond an earlier developmental shape and size, it is

possible that some cytoskeletal components will have been left behind in the original shape

of the boutons, and some will have moved outwards with the membrane.

Mitochondrial aggregation, between systems and ages

The mitochondrial aggregates observed at the larval NMJ and in culture could be related to dysfunctional intracellular transport, which has previously been reported in SOD1-based models of ALS (Bilsland et al., 2010; De Vos et al., 2007). In the case of the NMJ, reduced retrograde transport and/or increased anterograde transport could trap mitochondria at the end of the NMJ. This could then mean the aggregates disappear in *G85R* reared at HT because conditioning under heat stress somehow facilitates intracellular transport. But it's unclear what implications this would have for the phenotype in culture. Mitochondrial aggregates do not seem to localize to the very ends of neurites the way they localize to the ends of NMJ branches. These questions would be easier to approach with more information about the physical nature of the so-called mitochondrial aggregates. Due to the mechanics of TMRM, which is pulled into mitochondria by membrane potential (Chowdhury et al., 2016), increased fluorescence intensities due to higher membrane potential or due to mitochondrial volume may not be distinguishable. A possible solution could be to employ an indicator that is not sensitive to mitochondrial membrane potential, such as the genetically encoded mitochondrial indicator mito-roGFP (Albrecht et al., 2011). On the other hand, by design, it has the feature of becoming brighter when oxidized. This could complicate its use as a control for TMRM, but the additional information it could provide us may prove invaluable. Alternatively, dihyrorhodamine 123 (DHR123) (Ueda & Wu, 2008) is also a mitochondrial indicator relying on oxidation, though of the three it is likely the least practical: like TMRM, it relies on membrane potential to get into the mitochondria. Like mito-roGFP, DHR123 fluorescence is a reporter of oxidation, but DHR123 is not necessarily oxidized at the mitochondria, but rather in the cytoplasm, and it can be oxidized by a wide range of reactive oxygen species.

So far, the most important piece of information we have from the culture system is that there exists an early age at which an *Sod^{G85R}* mitochondrial phenotype does not strongly reveal

itself compared to wild-type. In the case of cultured giant motor neurons, that age is 3 days,

and the phenotype somehow appears in the 3 to 4 days following that. With the current data,

we can't know if the same process occurs at the larval NMJ, because larval dissections are

done at the 6-day time point, and dissections of larvae earlier than that are tremendously

difficult due to their small size. However, larvae younger than wandering 3rd instar, the stage

at which we dissect, are quite translucent, and it is possible to use a confocal microscope to

simply image fluorescence through the cuticle (Zito et al., 1999). A requirement for this

technique would be to use a genetically-encoded mitochondrial indicator, and we have one:

mito-roGFP. Concurrently, we can use the aforementioned *Drosophila* lines that express

GFP in the presynaptic membrane of NMJs to track the development of elongated boutons

and enlarged terminal boutons. Finally, confocal microscopy may enable the imaging of

smaller scale structures around or within the aggregates, be it in young or older larvae, or in

culture. And so the key to continuing this line of research appears to be the proper

application of higher-power microscopy, as well as the continued, comprehensive

exploitation of the relationship between the larval NMJ and the culture system.

**References**

Adams MD, Celniker SE, Holt RA, Evans CA, Gocayne JD, ... Venter JC. (2000) The

genome sequence of *Drosophila melanogaster*. *Science* 287: 2185-2195.

Aitken RJ, Krausz C. (2001) Oxidative stress, DNA damage, and the Y chromosome.

*Reproduction* 122: 497-506.

Albrecht SC, Barata AG, Großhans J, Teleman AA, Dick TP. (2011) In vivo mapping of

hydrogen peroxide and oxidize glutathione reveals chemical and regional specificity

of redox homeostasis. *Cell Metabolism* 14: 819-829.

Ali S, Jain SK, Abdulla M, Athar M. (1996) Paraquat induced DNA damage by reactive

oxygen species. *Biochemistry and Molecular Biology International* 39(1): 63-67.

Altan O, Pabuccuoglu A, Altan A, Konyalioglu S, Bayraktar H. (2003) Effect of heat stress on

oxidative stress, lipid peroxidation, and some stress parameters in broilers. *British*

*Poultry Science* 44(4): 545-550.

Ayala GX, Tapia R. (2003) Expression of heat shock protien 70 induced by 4-aminopyridine

through glutamate-mediated excitotoxic stress in rat hippocampus in vivo.

*Neuropharmacology* 45: 649-660.

Baek MK, Kim KO, Kwon HJ, Kim YW, Woo JH, Kim DY. (2016) Age-related changes in

antioxidative enzyme capacity in tongue of Fischer 344 rats. *Clinical and*

*Experimental Otorhinolaryngology* 9(4): 352-357.

Beers DR, Ho BK, Siklos L, Alexianu ME, Mosier DR, Mohamed AH, ... Appel SH. (2001)

Parvalbumin overexpression alters immune-mediated increases in intracellular

calcium, and delays disease onset in a transgenic model of familial amyotrophic

lateral sclerosis. *Journal of Neurochemistry* 79: 499-509.

Berlett BS, Stadtman ER. (1997) Protein Oxidation in Aging, Disease, and Oxidative Stress.

*The Journal of Biological Chemistry* 272(33): 20313-20316.

Bilsland LG, Sahai E, Kelly G, Golding M, Greensmith L, Schiavo G. (2010) Deficits in axonal

transport precede ALS symptoms in vivo. *Proc Natl Acad Sci* 107(47): 20523-20528.

Block ML, Calderón-Garcidueñas L. (2009) Air pollution: mechanisms of neuroinflammation

and CNS disease. *Trends in Neurosciences* 32(9): 506-516.

Brand AH, Perrimon N. (1993) Targeted gene expression as a means of altering cell fates

and generating dominant phenotypes. *Development* 118:401-415.

Campbell SH, Hilliker AJ, Phillips JP. (1985) Cytogenetic analysis of the *cSOD* microregion in *Drosophila melanogaster*. *Genetics* 112: 205-215.

Chen Q, Vasquez EJ, Moghaddas S, Hoppel CL, Lesnefsky EJ. (2003) Production of reactive oxygen species by mitochondria. *The Journal of Biological Chemistry* 278(38): 36027-36031.

Chen Y, Cohen TJ. (2019) Aggregation of the nucleic acid– binding protein TDP-43 occurs via distinct routes that are coordinated with stress granule formation. *J Biol Chem* 294(10): 3696-3706.

Chowdhury SR, Djordjevic J, Albensi BC, Fernyhough P. (2016) Simultaneous evaluation of substrate-dependent oxygen consumption rates and mitochondrial membrane potential by TMRM and safranin in cortical mitochondria. *Biosci Rep* 36: e00286

Culotta & Daly. (2013) Manganese Complexes: Diverse Metabolic Routes to Oxidative Stress Resistance in Prokaryotes and Yeast. *Antioxidants and Redox Signaling* 19(9): 933-944.

De Vos KJ, Chapman AL, Tennant ME, Manser C, Tudor EL, ... Grierson AJ. (2007) Familial amyotrophic lateral sclerosis-linked SOD1 mutants perturb fast axonal transport to reduce axonal mitochondria content. *Hum Mol Genet* 16(22): 2720-2728.

Dhalla NS, Temsah RM, Netticadan, T. (2000) Role of oxidative stress in cardiovascular diseases. *Journal of Hypertension*, 18(6): 655-673.

Feng, Y., Ueda, A., & Wu, C.-F. (2004). A modified minimal hemolymph-like solution, HL3. 1, for physiological recordings at the neuromuscular junctions of normal and mutant Drosophila larvae. *Journal of Neurogenetics*, 18(2): 377-402.

Forsberg K, Andersen PM, Marklund SL, Brannstrom T. (2011) Glial nuclear aggregates of superoxide dismutase-1 are regularly present in patients with amyotrophic lateral sclerosis. *Acta Neuropathol* 121: 623-634.

Frankel, A., & Brousseau, G. (1968). Drosophila medium that does not require dried yeast. *Drosophila Information Service* 43, 184.

Galvin (2017) The path to specialist multidisciplinary care in amyotrophic lateral sclerosis: A population- based study of consultations, interventions and costs. *PLoS ONE* 12(6): e0178796.

Guegan S, Prezedborski C. (2003) Programmed cell death in amyotrophic lateral sclerosis. *J Clin Invest* 111:153-161.

Giorgi C, Marchi S, Pinton P. (2018) The machineries, regulation and cellular functions of mitochondrial calcium. *Nature Reviews: Molecular Cell Biology* 19: 713-730.

Hardiman O, van den Berg LH, Kiernan MC. (2011) Clinical diagnosis and management of amyotrophic lateral sclerosis. *Nature Reviews: Neurology* 7: 639-649.

Kannan K, Jain SK. (2000) Oxidative stress and apoptosis. *Pathophysiology* 7(27) 153-163.

Kurosaka A, Yano A, Itoh N, Kuroda Y, Nakagawa T, Kawasaki T. (1991) The structure of a neural specific carbohydrate epitope of horseradish peroxidase antiserum. *the Journal of Biological Chemistry* 266(7): 4168-4172.

Landis G, Shen J, Tower J. (2012) Gene expression changes in response to again compared to heat stress, oxidative stress and ionizing radiation in *Drosophila melanogaster*. *Aging* 4(11): 768-789.

Lee J, Wu CF. (2010) Orchestration of Stepwise Synaptic Growth by K+ and Ca2+ Channels in *Drosophila*. *J. Neurosci.* 30(47): 15921-15833.

Lee J, Ueda A, Wu CF (2014) Distinct roles of *Drosophila cacophony* and *Dmca1D* Ca2+ channels in synaptic homeostasis: Genetic interactions with *slowpoke* Ca2+- activated BK channels in presynaptic excitability and postsynaptic response. *Dev. Neurobiol.* 141: 4548-4557.

Liguori I, Russo G, Curcio F, Bulli G, Aran L, Della-Morte D, Gargiulo G, ... Abete P. (2018) Oxidative stress, aging, and diseases. *Clinical interventions in Aging* 13: 757-772.

Lin CLG, Bristol LA, Jin L, Dykes-Hoberg M, Crawford T, Clawson L, Rothstein JD. (1998)

Aberrant RNA processing in a neurodegenerative disease: the cause for absent

EAAT2, a glutamate transporter, in amyotrophic lateral sclerosis. *Neuron* 20: 589-

602.

Lushchak, V. I. (2014). Free radicals, reactive oxygen species, oxidative stress and its

classification. *Chem-Biol Interact* 224: 164–175.

Mackenzie IRA, Rademakers R. (2008) Aberrant RNA processing in a neurodegenerative

disease: the cause for absent EAAT2, a glutamate transporter, in amyotrophic

lateral sclerosis. *Curr Opin Neurol* 21(6): 693-700.

Mahr A, Aberle H. (2006) The expression pattern of the *Drosophila* vesicular glutamate

transporter: A marker protein for motoneurons and glutamatergic centers in the

brain. *Gene Expression Patterns* 6: 299-309.

Massaad CA, Klann E. (2011) Reactive Oxygen Species in the Regulation of Synaptic

Plasticity and Memory. *Antioxidants and Redox Signaling* 14(10): 2013-2054.

Mates, J. M., Perez-Gomez, C., de Castro, IN. (1999). Antioxidant enzymes and human

diseases. *Clinical Biochemistry* 32(8): 595-603.

Mondola P, Damiano S, Sasso A, Santillo M. (2016) The Cu,Zn superoxide dismutase: not

only a dismutase enzyme. *Front Physiol* 7: 594.

Osking Z, Ayers JI, Hildebrandt R, Skruber K, Brown H, Ryu D, ... Vitriol EA. (2019) ALS-

linked SOD1 mutants enhance neurite outgrowth and branching in adult motor

neurons. *iScience* 11: 294-304.

Ott M, Gogvadze V, Orrenius S, Zhivotozsky B. (2007) Mitochondria, oxidative stress and

cell death. *Apoptosis* 12: 913-922.

Phillips JP, Campbell SD, Michaud D, Charbonneau M, Hilliker, AJ. (1989). Null mutation of

copper/zinc superoxide dismutase in Drosophila confers hypersensitivity to paraquat

and reduced longevity. *Proc Natl Acad Sci* 86(8): 2761-2765.

Patel A, Lee HO, Jawerth L, Maharana S, Jahnel M, Hein MY, Stoynov S, ... Alberti S. (2015)

619    A liquid-to-solid phase transition of the ALS protein FUS accelerated by disease

620    mutation. *Cell* 162, 1066-1077.

621 Peng IF, Berke BA, Zhu Y, Lee WH, Chen W, Wu CF. (2007) Temperature-dependent

622    developmental plasticity of *Drosophila* neurons: Cell-autonomous roles of

623    membrane excitability, Ca2+ influx, and cAMP signaling. *The Journal of*

624    *Neuroscience* 27(46): 12611-12622.

625 Perry JJP, Shin DS, Getzoff ED, Tainer JA. (2010) The structural biochemistry of the

626    superoxide dismutases. *Biochim Biophys Acta* 1804(2): 245-262.

627 Rosen DR, Siddique T, Patterson D, Figlewicz DA, Sapp P, Hentati A, ... Brown RH Jr.

628    (1993) Mutations in Cu/Zn superoxide dismutase gene are associated with familial

629    amyotrophic lateral sclerosis. *Nature* 362: 59-62.

630 Rowland LP, Shneider NA. (2001). Amyotrophic lateral sclerosis. *The New England Journal*

631    *of Medicine* 344(22): 1688-1700.

632 Ruan, H., & Wu, C. F. (2008). Social interaction-mediated lifespan extension of *Drosophila*

633    Cu/Zn superoxide dismutase mutants. *Proc Natl Acad Sci* 105(21): 7506-7510.

634 Saccon RA, Bunton-Stasyshyn RKA, Fisher EMC, Fratta P. (2013) Is SOD1 loss of function

635    involved in amyotrophic lateral sclerosis. *Brain* 136: 2342-2358.

636 Şahin A, Held A, Bredvik K, Major P, Achilli TM, Kerson, AG, ... Reenan, R. (2017). Human

637    SOD1 ALS mutations in a *Drosophila* knock-in model cause severe phenotypes and

638    reveal dosage-sensitive gain- and loss-of-function components. *Genetics* 205(2):

639    707-723.

640 Salińska E, Danysz W, Łazarexicz JW. (2005) The role of excitotoxicity in

641    neurodegeneration. *Folia Neuropathol* 43(4): 322-339.

Sies H. (1997) Oxidative Stress: Oxidants and Antioxidants (Physiological Society Symposium: Impaired endothelial and smooth muscle cell function in oxidative stress). *Experimental Physiology* 82: 291-295.

Smith EF, Shaw PJ, De Vos KJ. (2017) The role of mitochondria in amyotrophic lateral sclerosis. *Neurosci Lett* 710: 132933.

Sørensen JG, Nielsen MM, Kruhøffer M, Justesen J, Loeschcke V. (2005) Full genome gene expression analysis of the heat stress response in *Drosophila melanogaster*. *Cell Stress Chaperones*. 10(4): 312-328.

Sosa V, Moliné T, Somoza R, Paciucci R, Kondoh H, Lleonart ME. (2013) Oxidative stress and cancer: An overview. *Ageing Research Reviews* 12: 376-390.

Tang H, Tian E, Liu C, Wang Q, Deng H. (2013) Oxidative stress induces monocyte necrosis with enrichment of cell-bound albumin and overexpression of endoplasmic reticulum and mitochondrial chaperones. *PLoS ONE* 8(3): e59610.

Thannickal VJ, Fanburg BL. (2000) Reactive oxygen species in cell signaling. *Am J Physiol Lung Cell Mol Physiol* 279: L1005-L1028.

Torroja L, Packard M, Gorczyca M, White K, Budnik V. (1999) The *Drosophila* β-amyloid precursor protein homolog promotes synapse differentiation at the neuromuscular junction. *The Journal of Neuroscience* 19(18): 7793-7803.

Trotti D, Rolfs A, Danbolt NC, Brown RH Jr, Hediger MA. (1999) SOD1 mutants linked to amyotrophic lateral sclerosis selectively inactivate a glial glutamate transporter. *Nature Neuroscience* 2(5): 427-433.

Tsang CK, Liu Y, Thomas J, Zhang Y, Zheng XFS. (2015) Superoxide dismutase 1 acts as a nuclear transcription factor to regulate oxidative stress resistance. *Nat Commun* 5:3446.

Ueda A, Wu CF. (2006) Distinct frequency-dependent regulation of nerve terminal excitability

and synaptic transmission by $I_A$ and $I_K$ potassium channels revealed by *Drosophila*

*Shaker* and *Shab* mutations. *The Journal of Neuroscience* 26(23): 6238-6248.

Ueda A, Wu CF. (2008) Effects of *Hyperkinetic*, a beta-subunit of *Shaker* voltage- dependent

K+ channels, on the oxidation state of presynaptic nerve terminals. *J Neurogenet*

22(2): 1-13.

Uttara B, Singh AV, Zamboni P, Mahajan RT. (2009) Oxidative stress and neurodegenerative

diseases: a review of upstream and downstream antioxidant therapeutic options.

*Current Neuropharmacology* 7: 65-74.

Van Raamsdonk JM, Hekimi S. (2012) Superoxide dismutase is dispensable for normal

animal lifespan. *Proc Natl Acad Sci* 109(15): 5785-5790.

Vande Velde C, McDonald KK, Boukhedimi Y, ... Cleveland DW. (2011) Misfolded SOD1

Associated with Motor Neuron Mitochondria Alters Mitochondrial Shape and

Distribution Prior to Clinical Onset. *PLoS ONE* 6(7): e22031.

Wang J, Farr GW, Zeiss CJ, Rodriguez-Gil DJ, Wilson JH, ... Horwich AL. (2008) Progressive

aggregation despite chaperone associations of a mutant SOD1-YFP in transgenic

mice that develop ALS. *Proc Natl Acad Sci* 106(5): 1392-1897.

Wang J, Farr GW, Hall DH, Li F, Furtak K, Dreier L, ... Horwich AL. (2009) An ALS-Linked

Mutant SOD1 Produces a Locomotor Defect Associated with Aggregation and

Synaptic Dysfunction When Expressed in Neurons of Caenorhabditis elegans. *PLoS*

*Genetics* 5(1): e1000350.

Wang JW, Humphreys JM, Phillips JP, Hilliker AJ, Wu CF. (2000) A novel leg-shaking

*Drosophila* mutant defective in a voltage-gated $K^+$ current and hypersensitive to

reactive oxygen species. *Journal of Neuroscience* 20(16): 5958-5964.

Wu CF, Sakai K, Saito M, Hotta Y. (1990) Giant *Drosophila* neurons differentiated from

cytokinesis-arrested embryonic neuroblasts. *Journal of Neurobiology* 21(3): 499-

507.

Xia R, Liu Y, Yang L, Gal J, Zhu H, Jia J. (2012) Motor neuron apoptosis and neuromuscular

junction perturbation are prominent features in a *Drosophila* model of Fus-mediated

ALS. *Molecular Neurodegeneration* 7:10.

Zarei S, Carr K, Reiley L, Diaz K, Guerra O, Fernandez Altamirano P, ... Chinea A. (2015) A

comprehensive review of amyotrophic lateral sclerosis. *Surgical Neurology*

*International* 6:171.

Zhang Y, Liu Y, Walsh M, Bokov A, Ikeno Y, Jang YC, Perez VI, ... Richardson A. (2016)

Liver specific expression of Cu/ZnSOD extends the lifespan of *Sod1* null mice. *Mech*

*Ageing Dev* 154:1-8.

Zhong Y, Budnik V, Wu CF. (1992) Synaptic plasticity in *Drosophila* memory and

hyperexcitable mutants: role of cAMP cascade. *J Neurosci* 12(2): 644-651.

Zhong Y, Wu CF. (1991) Altered synaptic plasticity in Drosophila memory mutants with a

defective cyclic AMP cascade. *Science* 251(4990): 198-201.

Figure 1. Developmental lethality in *Sod* mutants.

(A) Representative images of larvae, pupae, and empty pupal casings of *Sod* mutant homozygotes (*Sod*/*Sod*) and balanced heterozygotes (*Sod*/TM6, *Tb*). *Sod* homozygotes have wild-type body morphology, and the TM6 balancer granted the heterozygotes shorter and fatter *Tubby* (*Tb*) body shape. The larvae represented are specifically of the wandering 3rd instar stage, immediately pre-pupation. The empty pupal casing are left over after successful eclosion of the juvenile fly.

(B) Images of *Sod* homozygotes dead during the wandering 3rd instar larval, pupal, or eclosion stages. Dead larvae are identifiable by their immobility and dark color and dead pupae have unevenly distributed contents and dark patches. Juvenile flies dead during eclosion were distinguished from flies in the process of eclosing by a second observation 15-30 min after the first. Juvenile fly corpses were also frequently darker than their live counterparts.

(C) Rates of lethality during development in wild-type (WT), $Sod^{G85R}$ (*G85R*/*G85R*), $Sod^{G85R}$/TM6,*Tb* (*G85R*/TM6), $Sod^{n108}$ (*n108*/ *n108*), and $Sod^{n108}$/TM6,*Tb* (*n108*/TM6). Dead individuals were counted at the 3rd instar larval stage (red), pupal stage (purple), and eclosion (blue). Rates of successful eclosions (white) are from counts of empty pupal casings. *** indicates $p < 0.001$ (Fisher's exact test). *n* of individuals are indicated at the bottom of each bar.

Figure 2. Morphological defects in larval motor synapses of *Sod* mutants.

(A) Representative NMJ of larval muscles 4 (M4) and 13 (M13) from WT, $Sod^{G85R}$, and $Sod^{n108}$, immunostained with anti-HRP (neuronal membrane, green in merged image) and anti-Dlg (subsynaptic reticulum, red in merged image). Type Ib (tonic glutamatergic) branches were examined in both muscles, and type Is (phasic glutamatergic) branches were examined in M13. Is branches are distinguishable from Ib by their smaller size , and weaker anti-HRP and anti-Dlg immunostaining. Labels on the anti-HRP images indicate branch types. The much smaller type II boutons are also labeled, but were not studied. Select portions of the M4 and M13 images are magnified to show phenotypic incomplete bouton segregation (third row, green boxes) and terminal bouton enlargement (fourth row, gold boxes) in $Sod^{G85R}$ and $Sod^{n108}$, compared to common WT variation. Note the "satellite" buds present on the magnified terminal bouton from $Sod^{n108}$. Scale bars in the upper left and lower left images are 10 μm.

(B) Per NMJ, number of branches (top) and synaptic boutons (bottom) of type Ib in M4 and M13, and type Is in M13. Branches are defined here as terminal processes composed of two or more synaptic boutons, distal of any bifurcations. Counts from larval abdominal segments 3, 4, and 5 are pooled. WT counts are plotted as black circles, $Sod^{G85R}$ (*G85R*) as maroon diamonds, $Sod^{n108}$ (*n108*) as red diamonds. Means are indicated by enlarged symbols, SEM by bars. * indicates $p < 0.05$, ** indicates $p < 0.01$, *** indicates $p < 0.001$ (Kruskal-Wallis test, rank-sum post hoc). *n* numbers (*n* of NMJ/*n* of larvae) are indicated above the genotype labels.

(C) Number of incompletely segregated (unsegregated) boutons per Ib branch per M4 NMJ (top), and percentage of Ib branches per M4 NMJ in which the terminal bouton was the largest after the last bifurcation. A bouton was described as incompletely segregated if the width of the narrowest point on the neck preceding it was greater than 80% of the width of the widest point on the bouton. The statistical parameters and their layout are identical to those of panel B.

Figure 3. Putative mitochondrial aggregates in terminal NMJ boutons of $Sod^{G85R}$ larvae, frequently present in enlarged terminal boutons.

(A) Mitochondrial live tissue staining (TMRM, first row) in muscle 4 NMJ terminal branches in wild-type (left), $Sod^{G85R}$ (center), and $Sod^{n108}$ (right) larvae. Second row is NMJ morphology, visible with a presynaptic membrane-targeted GFP (CD8-GFP), expression of which was driven by a Gal4 construct (C164 or OK371). Third row is an overlay of the first and second row. Fourth row contains magnified portions of the whole-NMJ TMRM images (first row, white boxes). Large white arrowhead indicates a brightly-stained potential aggregation of mitochondrial in the $Sod^{G85R}$ terminal bouton, smaller white arrowhead indicates bright mitochondrial staining in a non-terminal bouton. Scale bars in the upper left and lower left images are 10 μm.

(B) Percentages of branches in which putative mitochondrial aggregates appear in the terminal bouton (gray), or in the terminal bouton and one or more immediate neighbors (black), in WT, $Sod^{G85R}$, and $Sod^{n108}$. For WT vs $Sod$, *** indicates $p < 0.001$ (Fisher's exact test, comparing distributions of counts of terminal boutons without mito. aggr., with mito. aggr., and of terminal + neighboring bouton(s) with mito. aggr..).

(C) Percentages of terminal boutons with enlarged morphology (visible with CD8-GFP) in which putative mitochondrial aggregation appears, in WT and $Sod^{G85R}$ larvae. Includes counts from NMJs of muscles 4, 6, 7, 12, 13 in segments A2 through A6. $n$ of branches/NMJs/larvae: WT = 93/35/3, $G85R$ = 174/76/5, , $n108$ = 53/27/3.

Figure 4.

(A) Representative NMJ of larval muscle 4 (M4) (Ib branches) from WT, $Sod^{G85R}$, $Pk^{Pk}$, $Pk^{Sple}$, and the double mutants $Pk^{Pk};Sod^{G85R}$ and $Pk^{Sple};Sod^{G85R}$. Presynaptic neuronal membrane is immunostained with anti-HRP (green in merged image) and the microtubule-associated protein Futsch is stained with the monoclonal AB 22C10 (red in merged image). Is branches are labeled where appropriate, but were not studied here. Select portions of the full-scale images are magnified to show characteristic bouton morphologies and cytoskeletal structures (bottom three rows). Note the greater and irregular intensities of Futsch staining in $Sod^{G85R}$, $Pk^{Pk}$, and $Pk^{Pk};Sod^{G85R}$, and the strongly stained "loop" structures in $Pk^{Pk};Sod^{G85R}$. Scale bars in the full-scale and magnified HRP images are 10 μm.

(B) Per M4 NMJ, number of type Ib branches (top) and synaptic boutons (bottom). Counts from larval abdominal segments 2, 4, and 6 are pooled. WT counts are plotted as black circles, $Pk^{Pk}$ as blue inverted triangles, $Pk^{Pk};Sod^{G85R}$ ($Pk^{Pk};G85R$) as purple inverted triangles, $Sod^{G85R}$ ($G85R$) as maroon diamonds, $Pk^{Sple};Sod^{G85R}$ ($Sple;G85R$) as purple upright triangles, ($Sple$) as blue upright triangles. Means are indicated by enlarged symbols, SEM by bars. * indicates $p < 0.05$, ** indicates $p < 0.01$, *** indicates $p < 0.001$ (Kruskal-Wallis test, rank-sum post hoc). $n$ numbers ($n$ of NMJ/$n$ of larvae) are indicated above the genotype labels.

(C) Number of incompletely segregated (unsegregated) boutons per Ib branch per M4 NMJ (top), and percentage of Ib branches per M4 NMJ in which the terminal bouton was the largest after the last bifurcation. A bouton was described as incompletely segregated if the width of the narrowest point on the neck preceding it was greater than 80% of the width of the widest point on the bouton. The statistical parameters and their layout are identical to those of panel B.

**A**

5 mV
50 ms

WT    *sple*    *pk*    *sod*    *pk;sod*

**B**

ejp (mV)

| | | | | | |
|---|---|---|---|---|---|
| | a,b | a | b,c | a,b,c | c |

| | 13 | 11 | 12 | 2 | 6 |

*pk*    +    *sple*    *pk*    +    *pk*
*sod*    +    +    +    *G85R*    *G85R*

**C**

WT        *sple*        *pk*

*sod* ^G85R^        *pk;sod* ^G85R^

1 s

**D**

plateau ejp duration (s)

| | a | b,c | c | a,b | c |
|---|---|---|---|---|---|

3.0

2.0

1.0

| | 8 | 8 | 7 | 4 | 8 |

*pk*    +    *sple*    *pk*    +    *pk*
*sod*    +    +    +    *G85R*    *G85R*

Figure 5.

(A)  Increased size and variability of excitatory junctional potentials (ejps) from larval body wall muscles in *Pk* and *Pk;Sod*$^{G85R}$.  Ejps from two different muscles are shown as examples for each genotype.

(B)  a, b, and c indicates  that differences within the group is not significant ($p > 0.05$, F-test).  Error bars = SEM.  Number of NMJs are indicated.

(C)  Excitability in *Pk* and *Sod*$^{G85R}$ motor axon terminals.  Synaptic transmission was induced by direct activation of the motor axon terminals by electrotonic stimulation in the presence of a Na+ channel blocker (TTX, 3 μM) and K+ channel blockers (4-AP, 200 μM and TEA, 20 mM).  Under this condition, prolonged ejps supported by continuous transmitter release were recorded (see Text).  Amplitude is normalized to the peak.

(D)  The duration of plateau ejps were prolonged and more variable among perparations in *Pk*$^{Sple}$, *Pk*$^{Pk}$, and *Pk*$^{Pk}$;*Sod*$^{G85R}$.  a, b, and c indicates  that differences within the group is not significant ($p > 0.05$, F-test).  Error bars = SEM.  Number of NMJs are indicated.

Figure 6. Focal recording of plateau-like potentials induced at neuromuscular synapses by K$^+$ channel blockers.

(A) Sample focal recording (efEJP) traces from neuromuscular synapses of WT, Sod$^{G85R}$, Pk$^{Pk}$, Pk$^{Sple}$, Pk$^{Pk}$;Sod$^{G85R}$, and Pk$^{Sple}$;Sod$^{G85R}$. , upon electrotonic stimulation after application of TTX (3 µM), 4-AP (200 µM), and TEA (20 mM). Recordings are from the NMJ of muscles 6 and 7, in larval segments A4 and A5. Each recording site contained between 1 and 4 boutons, and could contain a mixture of type Ib and type Is boutons. Note that trace amplitude is dependent on the variable seal resistance between the recording electrode and the NMJ. All artifacts have been truncated.

(B) Durations of plateau-like efEJP. At each recording site, a series of single stimulation pulses will incrementally increasing voltage was applied. The plotted plateau efEJP duration for each site is a measurement from the first plateau observed in the stimulation series. WT counts are plotted as black circles, *Pk$^{Pk}$* as blue inverted triangles, *Pk$^{Pk}$;Sod$^{G85R}$* (*Pk$^{Pk}$;G85R*) as purple inverted triangles, *Sod$^{G85R}$* (*G85R*) as maroon diamonds, *Pk$^{Sple}$;Sod$^{G85R}$* (*Sple;G85R*) as purple upright triangles, (*Sple*) as blue upright triangles. Means are indicated by enlarged symbols, SEM by bars. *n* numbers (*n* of recording sites/*n* of NMJ/*n* of larvae) are indicated above the genotype labels.

**A** WT
 TTX&4AP&Q

6.00
6.00
6.00
6.00

100μV

50ms

**B** *sod*
 TTX&4AP&Q

6.00
6.00
6.00
6.00

**C** *pk/+*
 TTX&4AP&Q

1.08
1.90
2.70
3.50
4.40

500μV

50ms

**D** *pk/+; sod*
 TTX&4AP&Q

1.08
1.90
2.70
3.50
4.40

Figure 7. Focal recording examination of the excitability in $pk^{pk}$ and $sod^{n108}$ synaptic terminals.

Electrotonic stimuli (1 ms pulse duration) were applied to NMJ terminals at 0.1mM Ca2+ with Na$^+$ channel blockers (TTX, 3 µM) and K$^+$ channel blockers (4-AP, 200 µM and quinidine, 100 µM). Only type Ib boutons were recorded. (A) WT and  (B) $sodn108$ terminals quickly ran down after adding both 4-AP and quinidine and gave poor responses, whereas (C) $pk^{pk}$ heterozygous (over CyO-GFP) and (D) $pk^{pk}$/CyO-GFP; $sodn108$ displayed currents of striking amplitudes. Note that $pk^{pk}$/CyO-GFP; $sodn108$ has lower stimulation voltage (in V, labeled on the side) threshold and wider response current width. All artifacts have been truncated.