PP2A-Cdc55 phosphatase coordinates actomyosin ring contraction and

septum formation during cytokinesis

Yolanda Moyano-Rodríguez[1], David Vaquero[1,2], Odena Vilalta-Castany[1], Magdalena

Foltman[3,4], Alberto Sanchez-Diaz[3,4] and Ethel Queralt[1,2,*]

[1]Cell Cycle Group, Institut d'Investigació Biomèdica de Bellvitge (IDIBELL), Av. Gran

Via de L'Hospitalet 199-203, L'Hospitalet de Llobregat, Barcelona, Spain.

[2]Instituto de Biomedicina de Valencia (IBV-CSIC), C/ Jaume Roig 11, Valencia,

Spain.

[3]Instituto de Biomedicina y Biotecnología de Cantabria, Universidad de Cantabria,

CSIC, Santander, Spain

[4]Departamento de Biología Molecular, Facultad de Medicina, Universidad de

Cantabria, Santander, Spain.

*Correspondence/lead contact: equeralt@ibv.csic.es, telephone +34963391760.

ORCID: 0000-0003-0045-0039

Running head: New PP2A-Cdc55 role in cytokinesis

Keywords: Cell Cycle, cytokinesis, PP2A-Cdc55, ingression progression complexes

(IPCs), AMR contraction, septum formation

Summary

Eukaryotic cells divide and separate all their components after chromosome
segregation by a process called cytokinesis to complete cell division. Cytokinesis is
highly regulated by recruitment of the components to the division site and through
post-translational modifications such as phosphorylations. The budding yeast mitotic
kinases Cdc28-Clb2, Cdc5, and Dbf2-Mob1 phosphorylate several cytokinetic
proteins contributing to the regulation of cytokinesis. The PP2A-Cdc55 phosphatase
regulates mitosis counteracting Cdk1- and Cdc5-dependent phosphorylation. This
prompted us to propose that PP2A-Cdc55 could also be counteracting the mitotic
kinases during cytokinesis. Here, we demonstrate that primary septum formation and
actomyosin ring contraction are impaired in the absence of PP2A-Cdc55. In addition,
by *in vivo* and *in vitro* assays, we show that PP2A-Cdc55 dephosphorylates the
chitin synthase II (Chs2 in budding yeast) a component of the Ingression
Progression Complexes (IPCs) involved in cytokinesis. Interestingly, the non-
phosphorylable version of Chs2 rescues the asymmetric AMR contraction observed
in *cdc55Δ* mutant cells. Therefore, timely dephosphorylation of the Chs2 by PP2A-
Cdc55 is crucial for proper actomyosin ring contraction. These findings reveal a new
mechanism of cytokinesis regulation by the PP2A-Cdc55 phosphatase and extend
our knowledge in the involvement of multiple phosphatases during cytokinesis.

Introduction

Cytokinesis is the final event of the cell cycle and mediates the physical separation of mother and daughter cells. It is a highly ordered and regulated process that is conserved among eukaryotes. Cytokinesis must be spatially and temporally coordinated with sister chromatids segregation to avoid problems in chromosome segregation –failures can lead to aneuploidies and are associated with tumorigenesis[1]. Detailed mechanistic studies are important to understand cytokinesis relevance in normal cellular functions as well as its impact on human diseases.

In fungi and animal cells, the cytokinetic machinery comprises two major elements: septin and contractile actomyosin rings. In *Saccharomyces cerevisiae*, the process takes place at the division site, a narrow region linking mother and daughter cells known as the bud neck. At the end of G1, the Cdc42 GTPase polarizes actin cables and patches towards the new bud site, and the septin ring is assembled[2,3].The septin ring is reorganized later in S phase and becomes an hourglass-like structure[4] that will split into two rings in anaphase[5–7]. Septins serve as an anchor for various cytokinesis-related proteins including the type II myosin heavy chain, Myo1[8]. Myo1, together with actin filaments and the essential and regulatory myosin light chains Mlc1 and Mlc2, shape the actomyosin ring (AMR)[5,8–10]. The AMR leads to the furrow ingression through its contraction and constriction[11,12].

Cytokinetic regulatory proteins such as Iqg1,Hof1, Inn1 and Cyk3 are recruited at the AMR[6,13–15], which mediate the activation of the chitin synthase II, Chs2[6,16–18]. Chs2 synthesizes the primary septum (PS), concomitantly with AMR constriction[19,20], and drives the second part of its contraction [21]. These proteins are part of the Ingression Progression Complexes (IPCs)[17,22], which coordinates

primary septum formation, actomyosin ring contraction and ingression of the plasma

membrane at the division site[16,17,23–26].

Once the PS has been synthesized, the two secondary septa (SS) will be formed by

Chs3, glucan synthases (Fks1) and mannosyltransferases on both sides of the

PS[20,27,28]. Finally, cytokinesis is completed when the daughter cell synthesizes

hydrolases and chitinases to hydrolyze the remaining cell wall structures in between

SS[29].

The timing of cytokinesis must be tightly regulated to ensure it only happens when

anaphase is completed. Anaphase is regulated by two pathways: FEAR

(CdcFourteen Early Anaphase Release) and MEN (Mitosis Exit Network). Both are

coordinated to promote the activation of the Cdc14 phosphatase[30–33]. Cdc14 is

responsible for Cdk1 (Cdc28 in budding yeast) inactivation and the

dephosphorylation of Cdk1 targets during mitosis[34]. Several MEN kinases have

been found to be involved in cytokinesis[35]. Polo-like kinase, Cdc5 regulates AMR

formation and membrane ingression[36,37]. Dbf2, the downstream MEN kinase,

phosphorylates Hof1 promoting its re-localization from the septin ring to the

AMR[7,36]. Cdk1-Clb2 phosphorylates Iqg1 limiting the actin recruitment at the

division site [38,39] and Chs2 promoting its retention at the endoplasmic reticulum

(ER)[40–42]. Cdk1 also inhibits Iqg1 and Inn1 localization at the division site, and

Inn1 interaction with IPCs[38,43,44]. Later on, MEN and Cdc14 phosphatase initiate

cytokinesis by counteracting Cdk1 phosphorylation in budding yeast[39,41,44,45].

However, the inactivation of Cdk1-Clb2 activity during late anaphase is not sufficient

to trigger cytokinesis[41,46], it is also necessary the Cdc14-dependent

dephosphorylation of Iqg1, Inn1 and Chs2 for cytokinesis completion[39,41,44,45].

Chs2 translocates to the division site upon Chs2 dephosphorylation by

Cdc14[41,47].

Recent work demonstrates that multiple phosphatases shape the phospho-proteome

during mitotic exit[48,49], pointing out the contribution of other phosphatases such as

PP2A-Rts1 and PP2A-Cdc55[48]. These observations suggest thatPP2A

phosphatases could also be required for cytokinesis. Here, we propose that PP2A-

Cdc55 regulates cytokinesis through the dephosphorylation of the IPC protein, Chs2.

We have found that the PP2A-Cdc55 participates in the regulation of the

phosphorylation state of Chs2 and dephosphorylates Chs2 *in vivo* and *in vitro*. In the

absence of Cdc55, AMR contraction and the PS formation occur asymmetrically to

one side of the bud neck supporting a role for PP2A-Cdc55 in cytokinesis regulation.

Interestingly, the non-phosphorylable version of Chs2, *chs2-S133A* rescues the

asymmetric AMR contraction in *cdc55Δ* mutant cells. The phospho-mimetic *chs2-*

S133E promotes a longer Hof1 contraction time in wild-type cells, similarly to the

absence of Cdc55, suggesting that hyperphosphorylation of Chs2 alters Hof1

residence time at the division site. In conclusion, PP2A-Cdc55 coordinates AMR

contraction and PS formation via the dephosphorylation of Chs2 during cytokinesis.

Results

Defective septum formation and AMR contraction in the absence of Cdc55

It was described that PP2A-Cdc55 is localized at the division site during cytokinesis

[50]. However, neither a role for PP2A-Cdc55 during cytokinesis was demonstrated

nor the molecular mechanism by which PP2A-Cdc55 could regulate cytokinesis. This

prompted us to study the cytokinesis phenotypes of the *cdc55Δ* mutant cells.

Septins act as scaffold platforms for many proteins and impose a diffusion barrier for

regulating cell polarity, cell remodeling and cytokinesis [51]. Septins localization and

organization are highly dynamic through the cell cycle [52,53]. We wondered

whether septin structures were defective in the absence of Cdc55. Septins were

visualized *in situ* by immunofluorescence staining of Cdc11 and Shs1-HA in

asynchronous cells. The septin structures (septin rings, collar and double rings) were

indistinguishable between *cdc55Δ* and wild-type cells (Fig. S1), suggesting that

septin dynamics might not be greatly affected by the absence of Cdc55.

PP2A-Cdc55 regulates bud morphology through actin polarization and cell-wall

synthesis [54]. Actin cytoskeleton polarization at the site of bud emergence is

triggered at Start by the kinase activities of Cln1,2-Cdc28. Upon Clb2-Cdc28 kinase

inactivation at the end of mitosis, the actin cytoskeleton is directed to the neck for the

formation of the AMR [9,39]. We envisage the possibility that PP2A-Cdc55

influences also actin cytoskeleton during cytokinesis. To examine this possibility,

actin filaments were stained with rhodamine-labelled phalloidin in cells progressing

through mitosis and cytokinesis in *cdc55Δ* cells. Actin signal was found to be

depolarized in metaphase-arrested cells and localized at the bud neck in anaphase-

cytokinesis in wild-type and *cdc55Δ* cells (Fig. S2a). However, in a subpopulation of

cdc55Δ cells, actin polarization at the new bud was observed before cytokinesis had

been completed (Fig. S2b). We observed actin signals simultaneously at the division

site during cytokinesis and at the new bud site in 12.5% of *cdc55Δ* mutant cells,

whereas there was no premature actin re-polarization at the new bud site in wild-type

cells (Fig. S2b). The premature actin localization indicates that actin re-polarization

of the next cell cycle occurs before cell division in a fraction of *cdc55Δ* cells.

Therefore, PP2A-Cdc55 could act to prevent actin re-polarization until cytokinesis is

completed.

Cytokinesis in budding yeast is accomplished by the concerted action of the

actomyosin contractile ring (AMR) and the septum formation. We investigated how

these two processes occur in the absence of Cdc55. First, to investigate whether the

actomyosin contractile ring is functional, we analyzed the localization of the Myo1-

tdTomato fusion protein in cells progressing through mitosis and cytokinesis after

synchronous release from the metaphase arrest by Cdc20 depletion. It has been

described that Myo1 localizes to the division site immediately after budding.

Accordingly, in metaphase-arrested cells Myo1-tdTomato was localized at the bud

neck in control cells. Similarly, Myo1-tdTomato was also detected at the bud neck

during metaphase in *cdc55Δ*mutant cells. In anaphase, the Myo1 signal is reduced,

reflecting the contraction of the AMR, until the signal becomes a single dot and

finally disappears (Fig. 1a). The dynamics of AMR contraction were similar in the two

strains, but we observed that AMR contraction was asymmetric respect to the

centripetal axis in 86% (N=69) of *cdc55Δ* cells (Fig. 1b). In order to check whether

the asymmetric Myo1 signal is not due to an adaptive mechanism of the *cdc55Δ*

deletion mutant, we investigated the Myo1 contraction after inducing the Cdc55

degradation during metaphase using an auxin-degradation system [55]. Upon Cdc55

degradation, Myo1 contraction was also mostly asymmetric (86% of cells) (Fig. 1c).

The results indicate that the lack of PP2A-Cdc55activity promotes the asymmetric

contraction of Myo1.

We next examined whether Myo1 asymmetry could also be detected in the inactive

version of Cdc55 (*cdc55-ED*). We synchronized cells at the metaphase-anaphase

transition by Cdc20 depletion and analyzed the contraction of the Myo1-tdTomato.

We observed that 65% of *cdc55-ED* mutant cells showed an asymmetric Myo1-

tdTomato signal upon contraction (Fig. 1c). This result indicates that, similar to the

absence of Cdc55, the non-functional Cdc55 results in the alteration of AMR

contraction. Overall, we conclude that Myo1 recruitment and time of localization at

the division site are not altered in the absence of Cdc55, although AMR contraction

is defective in the absence of PP2A-Cdc55 activity. This asymmetric localization has

been reported before in some IPCs mutants [26] and suggests a dysfunctional AMR.

These results suggest that PP2A-Cdc55 is required for the correct function of AMR

and for an efficient cytokinesis.

To assess whether the asymmetric AMR contraction phenotype was not affected by

the synchronization method, we performed the assay synchronizing cells in G1 by

alpha factor in absence of Cdc55. *cdc55Δ* cells enter mitosis with a delay due to

compromised Cdk1 activity because of inhibitory Cdc28-Y19 phosphorylation [56].

To correct for this delay, we introduced the *cdc28_Y19F* allele, which is refractory to

Cdk1 inhibition. The *cdc55Δ* cells containing *cdc28_Y19F* progressed normally

through the cell cycle [57]. Again, we observed that Myo1-tdTomato constriction was

asymmetric in 80% of the *cdc55Δ cdc28_Y19F* cell population (Fig. S3). Therefore,

the asymmetric Myo1 signal in the absence of Cdc55 was observed independently of

the synchronization method used.

To study septum formation, chitin deposition at the division site was analyzed by *in*

vivo staining with calcofluor white and measured the fluorescence intensity of the

incorporated calcofluor on living cells containing Myo1-tdTomato as a control for

cytokinesis progression. We arrested cells at metaphase by Cdc20 depletion,

released them into mitosis and took images 40-50 min after the release when we

found cells at cytokinesis. Calcofluor intensity was then measured and quantified in

wild-type and *cdc55Δ* cells (Fig. 1d). There was a 45% reduction in the intensity of

the calcofluor staining in the absence of Cdc55 compared to control cells. Chitin is

incorporated in primary septum (PS) and secondary septa (SS); therefore, we cannot

distinguish at which septum the reduction occurs. For this reason, we repeated the

calcofluor staining in cells containing a deletion for*CHS3*, the chitin synthase

responsible for secondary septa formation [20]. As previously described, *chs3Δ* cells

showed wider bud necks and cytokinetic defects provoking the formation of chains of

cells [58]. A reduction of 70% in the calcofluor intensity was measured in the *chs3Δ*

background in the absence of Cdc55 (Fig. 1e). Interestingly, we noticed that 82%

(N=22) of the cells showed asymmetric Myo1 contraction in *chs3Δ* cells (Fig. 1c),

indicating defects in cytokinesis [26]. The above results indicate that primary and

secondary septa formation are reduced in cells lacking PP2A-Cdc55 activity,

suggesting that PP2A-Cdc55 has a role coordinating AMR contraction with septum

formation.

Primary septum formation and membrane invagination occur concomitantly. To

determine whether ingression of the plasma membrane could occur in *cdc55Δ* cells,

we performed time-lapse video microscopy of cells expressing the small G-protein

Ras2 fused to 3 copies of GFP to study plasma membrane dynamics. The

visualization of the plasma membrane at the site of division revealed no cytoplasm

connection between mother and daughter cells after cytokinesis, confirming that

cytoplasmic division is resolved in control and *cdc55Δ* cells (Fig. S4). In conclusion,

cdc55Δ cells have an asymmetric AMR contraction and a reduction in septa

formation, but nevertheless manage to complete cell division.

Next, to study the defects in septa formation we investigated the cytokinetic structure

by transmission electron microscopy (TEM). We synchronized cells at the

metaphase-anaphase transition by Cdc20 depletion and captured images 40-50 min

after the synchronous release. Cells synthesizing the primary septum were identified

and the structure of the septum was then examined. In wild-type cells, we observed

PS formation from the cell wall following membrane invagination (Fig. 2a), as

expected. Once the PS was finished, the two SS was formed on both sides of the PS

(Fig. 2b). Finally, the PS is degraded, and the cells physically separated (Fig. 2c).

Conversely, in *cdc55Δ* cells, the PS emerged from only one side of the division site

(Fig. 2d). From the cells performing primary septum formation, we quantified 85%

(N=25) with asymmetric PS in absence of Cdc55. Some cells showed aberrant,

thicker structures with diverse morphologies that resemble the secondary septa-like

structures, commonly referred as remedial septum (Fig. 2e). The remedial septum

was first described in IPCs mutant cells, i.e., *myo1Δ*, *chs2Δ* and *chs3Δ* cells[24].

They are chitin structures reminiscent of SS that allow cells to complete cytokinesis

when the PS is defective [24,59]. Strikingly, in *cdc55Δ* cells, the mother and daughter

cells are finally able to separate physically, probably upon formation of the remedial-

like septum. To further investigate this hypothesis, we repeated the experiments

using a *chs3Δ* background in which no secondary septa is formed (Figs. 2g-i).

Similar results were obtained, whence the *cdc55Δ chs3Δ* cells had asymmetric PS

formation and a remedial-like septum was formed (Figs. 2j-l). These results suggest

that remedial-like septum formation enables cytoplasm separation and cell division in

the absence of Cdc55.

In summary, the asymmetric PS formation detected by EM and the asymmetric AMR

contraction observed in *cdc55Δ* cells, indicates that PP2A-Cdc55 has a role

regulating the correct coordination of the AMR contraction and PS formation.

PP2A-Cdc55 regulates the dephosphorylation of Chs2

The Ingression Progression Complexes (IPCs) coordinate AMR constriction, plasma

membrane ingression and septum formation[17,18]. We envisaged that PP2A-Cdc55

may have a role during cytokinesis through the dephosphorylation of IPCs

proteins[60]. We initially investigated the genetic interactions between *cdc55Δ* and

IPCs mutants. We prepared double mutants with *cdc55Δ* and degron-conditional

mutants to induce protein degradation for the IPCs subunits[17,61]. No differences in

cell growth were observed on control plates (Fig. 3a). However, the viability of the

double mutants *cdc55Δ hof1-aid* and *cdc55Δ td-cyk3-aid* was impaired under

restrictive conditions (presence of auxin) (Fig. 3a). The synthetic sick interactions

found between *cdc55Δ* and *hof1-aid* or *td-cyk3-aid* degron mutants indicate that

Cdc55 is functionally related to the IPC subunits, suggesting that Cdc55 is involved

in the regulation of cytokinesis.

To determine whether the IPCs proteins physically interact with PP2A-Cdc55, we

performed co-immunoprecipitation experiments. We synchronized cells in

metaphase as described above, and co-purification of Chs2 during anaphase and

cytokinesis were detected in Cdc55 immunoprecipitates (Fig. 3b). Taken together,

these results suggest that Chs2 forms a complex with PP2A-Cdc55 during

progression through mitosis and cytokinesis.

To study the functional link between Cdc55 and the IPCs, we studied whether PP2A-

Cdc55 regulates the dephosphorylation of IPC proteins. We analyzed the

phosphorylation status of the IPC proteins in the absence of Cdc55 (Fig 3c and S5a-

d) and observed an increase of Chs2 phosphorylation levels during cytokinesis. First,

we synchronized cells at metaphase and released them synchronously into

anaphase to visualize the phosphorylation of Chs2 during mitosis and cytokinesis.

Mitotic spindle was stained by tubulin immunofluorescence and used as marker of

cell-cycle progression. Cdc14 release from the nucleolus was also determined as

control of mitosis progression. Cdc14 was prematurely release in metaphase in

absence of Cdc55 as previously published[57]. During the time-course, Chs2 protein

presented slow migration isoforms at metaphase and early anaphase in control cells

(Fig. 3c meta-20 min). From anaphase until early G1 (Fig. 3c; 30-50 min), two Chs2

isoforms were detected. Since Chs2 presented many different migrating bands, we

performed an alkaline phosphatase experiment to check which corresponds to

phosphorylation. In presence of alkaline phosphatase, the slower Chs2 migrating

bands collapse into the faster migrating band indicating that the upper bands are

phosphorylation events (Fig. S5e). Remarkably, in *cdc55Δ* mutant cells the slower-

migrating isoforms are maintained throughout the time-course, indicating that Chs2

was not efficiently dephosphorylated during late anaphase/cytokinesis (Fig. 3c; 40-

70min). These results suggest that Chs2 phosphorylation levels are altered in the

absence of Cdc55 during cytokinesis.

Later, to explicitly test whether Chs2 is dephosphorylated by PP2A-Cdc55, we

measured PP2A-Cdc55 phosphatase activity against Chs2 after PP2A-Cdc55

purification from metaphase-arrested cells when PP2A-Cdc55 is active[57].

Immunopurified-Chs2 was incubated with purified PP2A-Cdc55 and Chs2

dephosphorylation was visualized using an anti-phospho Ser/Thr antibody (Fig. 3d).

A 60% reduction in the Chs2 phosphorylation signal was measured after 40 min of

incubation in presence of PP2A-Cdc55. As negative control, the phosphatase assay

was done in the same conditions using separase (Esp1) as substrate.

Immunopurified-Esp1 was incubated with purified PP2A-Cdc55 and the

phosphorylation levels of Esp1 visualized with the anti-phospho Ser/Thr antibody

(Fig. S5f). Esp1 phosphorylation signal was maintained during the experiment

indicating that PP2A-Cdc55 is not able to dephosphorylate Esp1 in the same

conditions used for Chs2. This result suggests that PP2A-Cdc55 dephosphorylation

occurs specifically over Chs2.

Finally, we repeated the PP2A-Cdc55 phosphatase assay using a recombinant

fragment of Chs2 (1-629) purified from *E. coli* and previously phosphorylated by

Cdk1-Clb2. *In vitro* ^{32}P-phosphorylated Chs2 substrate was incubated with Cdc55

immunoprecipitated from control cells expressing Cdc55 and *cdc55-ED* inactive

version of PP2A-Cdc55[62]. A reduction of 84% in the Chs2 phosphorylation signal

was detected in the presence of control Cdc55. The *cdc55-ED* mutant version was

not able to dephosphorylate Chs2 significantly (Fig. 3e). These results strongly

suggest that Chs2 is dephosphorylated by PP2A-Cdc55 *in vivo* and *in vitro.*

PP2A-Cdc55 regulates IPCs symmetric localization and residence time at the

division site

IPCs proteins form a complex whose function is to coordinate AMR contraction,

plasma membrane ingression and PS formation[17,18]. For this reason, we

examined whether the absence of Cdc55 disturbs the IPCs subunits proper or timely

localization at the division site. To do this, we studied the localization of the IPCs

subunits by time-lapse microscopy in the absence of Cdc55. We arrested cells in

metaphase by Cdc20 depletion and captured images every 2 minutes after

synchronous release into anaphase by Cdc20 re-induction. Iqg1, Inn1, Hof1, Cyk3

and Chs2 were contracted asymmetrically at the division site in *cdc55Δ* cells (Fig.

4a), consistent with the asymmetry of Myo1 described above. These findings

indicate that the lack of PP2A-Cdc55 activity provokes a defective contraction of the

AMR, resulting in defective PS formation.

We then investigated the contraction and residence time of the IPCs subunits (Fig.

4b). No differences were detected in the residence time at the bud neck in Iqg1-GFP

and Inn1-GFP proteins (Fig. 4b and Fig. S6a-b). However, we observed that Hof1-

GFP and Cyk3-GFP signals took longer to complete contraction and to disappear in

cdc55Δ cells (Fig. 4b and Fig. S6c-d), further indicating that cytokinesis is affected in

absence of PP2A-Cdc55. The longer localization of Hof1 and Cyk3 at the division

site could be a result of the impaired AMR contraction.

On the contrary, the Chs2-GFP residence time in *cdc55Δ* mutant cells was reduced

compared to wild-type cells (Fig. 4b and S6e). Therefore, this result suggests that

PP2A-Cdc55 regulates Chs2 localization dynamics, probably through the

dephosphorylation of Chs2.

Timely Chs2 dephosphorylation by PP2A-Cdc55 is required for proper AMR

contraction

Our results suggest that PP2A-Cdc55 regulates Chs2-dependent processes during

cytokinesis, such as PS formation and AMR contraction, and that participates in the

dephosphorylation of Chs2. Therefore, PP2A-Cdc55 might contribute to cytokinesis

regulation by controlling the phosphorylation levels of Chs2.

To screen for proteins regulated by the PP2A-Cdc55 phosphatase we studied the

phosphoproteome in absence of Cdc55 by a quantitative analysis based on SILAC

labeling. Phosphopeptides were enriched by Tish enrichment[63] and identified by

LC-MS/MS. The screening revealed that a phosphopeptide corresponding to Chs2

protein was hyperphosphorylated in *cdc55Δ* mutant cells. The Chs2 peptide

contained one Cdk1 minimal S/TP site: S133 (Fig. 5a). The phosphosite was

detected with the highest confidence (pRS site probability 99.9% and p-value <

0.00001) and was also identified in a second phospho-proteomic study for *cdc55Δ*

mutant cells[49]. The S133 is one of the 6 Cdk1 consensus sites (S/T-P) previously

reported to be phosphorylated by Cdk1[64]. Chs2's residues S14, S60, S69 and

S100 containing the SP consensus sequence were described to be efficiently

dephosphorylated by Cdc14,and the phospho mimetic mutant at those four sites,

chs2-S4E, promotes Chs2 constitutive localization in the ER [41,47]. Cdc14 is active

as a phosphatase and prematurely released from the nucleolus in *cdc55Δ* mutant

cells[57,65]; therefore, the S133 hyperphosphorylation is a consequence of the lack

of PP2A-Cdc55 activity since Cdc14 is not defective in absence of Cdc55.

To investigate whether the S133 is a critical Chs2 phosphorylation site, we

introduced a non-phosphorylable version of Chs2, *GAL1-chs2-S133A-YFP*, in wild-

type cells and analyzed the phosphorylation levels in Chs2-YFP immunoprecipitates

using the anti-phospho Ser/Thr antibody. Chs2 phosphorylation was detected in

control cells expressing the wild-type version of Chs2-YFP (Fig. 5b). By contrast, the

signal was greatly reduced in the immunoprecipitation of the non-phosphorylable

version *chs2-S133A* (Fig. 5b). This result suggests that the S133 of Chs2 is

phosphorylated.

If the cytokinetic defects observed in *cdc55Δ* mutant cells are consequence of the

hyperphosphorylation of Chs2, the non-phosphorylable Chs2 version should rescue

the asymmetric Myo1 contraction. For this reason, we introduced the non-

phosphorylable version *GAL1-chs2-S133A-YFP* in wild-type and *cdc55Δ* mutant

strains. Cells were synchronized by Cdc20 depletion and re-addition and *GAL1-*

CHS2 versions were induced prior to metaphase release. We followed Myo1-

tdTomato and Chs2-YFP dynamics during mitosis and cytokinesis by time-lapse

microscopy. Myo1 signal was asymmetric in the presence of the control Chs2-YFP

(80% of cells; N=25), as expected in a *cdc55Δ* background (Fig. 5c). Remarkably, in

the *chs2-S133A* non-phosphorylable version Myo1 became symmetric in 90% of the

cells(N=31) (Fig. 5c). This indicates that the non-phosphorylable Chs2 mutant

rescued the asymmetric localization of Myo1 in absence of Cdc55, Next, we

wondered whether the *chs2-S133A* version also rescued the longer Hof1 contraction

time observed in *cdc55Δ* cells (see Fig. 4b) by investigating the Hof1-mCherry

dynamics. Interestingly, Hof1 contraction was similar in Chs2 and *chs2-S133A* in

absence of Cdc55 (Fig. 5d). In conclusion, in absence of Chs2 phosphorylation at

S133A, Myo1 signal is symmetric and Hof1 residence time is normal, suggesting that

the S133 is the main dephosphorylated residue regulated by PP2A-Cdc55.

Finally, to investigate whether the hyperphosphorylation of Chs2 is sufficient to alter

AMR symmetric contraction we introduced the phospho-mimetic *chs2-S133E* version

in wild-type cells (Fig. 5e). No significant increase in the asymmetric Myo1 signal in

the presence of *chs2-S133E* was observed suggesting that Chs2 phosphorylation is

not sufficient to promote asymmetric AMR contraction. Nevertheless, an increase in

the Hof1 residence time was observed in the phospho-mimetic *chs2-S133E* version

(Fig. 5f), similarly to what we observed in *cdc55Δ* mutant cells (compare to Fig. 4b).

These results suggest that PP2A-Cdc55 has a role in cytokinesis regulating Chs2

phosphorylation.

Overall, our findings demonstrate that the lack of PP2A-Cdc55 activity provokes an

asymmetric contraction of the AMR and a defective PS formation that depends on

the tight regulation of Chs2 phosphorylation. Therefore, Chs2 dephosphorylation by

PP2A-Cdc55 is essential for maintaining proper AMR contraction and septum

formation; both required for an efficient cytokinesis.

Discussion

PP2A-Cdc55 regulates Chs2 phosphorylation and its localization dynamics at the division site

The localization and function of the IPCs subunits(Iqg1, Myo1, Hof1, Cyk3, Inn1 and Chs2 proteins), which coordinate AMR contraction, plasma membrane ingression and PS formation, are tightly regulated by phosphorylation[43,44,66]. To date, only two phosphatases, Cdc14 and PP2A-Rts1, are known to have a role in cytokinesis[67], and it has been described that Cdc14 dephosphorylates several subunits of the IPCs proteins such as Chs2, Inn1 and Hof1[39,41,44,46]. A putative role for PP2A-Cdc55 in cytokinesis was also proposed because of its elongated morphology and multinucleated cells at low temperatures in *cdc55Δ* cells[68]. In addition, PP2A-Cdc55 is localized at the division site during cytokinesis[50]. However, the role of PP2A-Cdc55 during cytokinesis was not demonstrated and the molecular mechanism by which PP2A-Cdc55 regulates cytokinesis is still unknown.

Here, we describe a new function of PP2A-Cdc55 phosphatase in the regulation of cytokinesis. In summary, PP2A-Cdc55 regulates the dephosphorylation of Chs2 and it is involved in proper AMR contraction and septum formation. PP2A-Cdc55 dephosphorylates Chs2 *in vivo* and *in vitro*, indicating that it is a PP2A-Cdc55 target. The lack of dephosphorylation of Chs2 in absence of Cdc55 might be responsible of the defects seen in IPCs (Hof1) residence time (Fig. 4b) and AMR asymmetric contraction, since the *chs2-S133A* (Fig. 5) non-phosphorylable mutant rescue the asymmetric Myo1 signal in *cdc55Δ* mutant cells. In contrast, the phospho-mimetic mutant *chs2-S133E* is not sufficient to induce an asymmetric Myo1 signal in control cells, although provokes an increase in Hof1 residence time at the division site.

These results suggest that timely dephosphorylation of Chs2 by PP2A-Cdc55 is

crucial for proper AMR contraction. Since the IPCs coordinate AMR constriction,

plasma membrane ingression and septum formation, we propose that PP2A-Cdc55

is also involved in this efficient mechanism of coordination through Chs2

dephosphorylation (Fig. 6).

The role of PP2A-Cdc55 during cytokinesis

In the absence of Cdc55, Myo1 was constricted and unanchored in a timely fashion,

but the constriction was displaced from the central axis, becoming asymmetric (Fig.

1a). This phenotype is characteristic of IPCs mutants[6,26,52] and denotes a

dysfunctional AMR.

On the other hand,Hof1 asymmetric PS formation was described in *chs2*Δ cells[6];

but not *vice versa*[6,17], suggesting that Hof1 also influences PS formation.

Therefore, defective PS formation in absence of Cdc55 could also contribute to the

asymmetry observed in the other IPCs proteins.

In addition, PP2A-Cdc55 regulates septum formation. We observed that chitin

incorporation was severely reduced in *cdc55*Δ and *cdc55*Δ *chs3*Δ cells (Fig. 1d),

indicating that Chs2 activity might be impaired. Interestingly, the electron microscopy

study demonstrated that the PS appears only at one side of the division site during

its formation in absence of Cdc55 (Figs. 2d and j). Later, a remedial-like septum that

contains even cytosol fractions embedded in the septa is detected in absence of

Cdc55 (Fig. 2f and l). The remedial septum formation was previously described in

IPCs and MEN mutants with defective PS formation[13,18,69,70]. Chs3 glucan

synthases and mannosyltransferases are responsible for the SS formation and the

construction of the remedial septum[27,28,59]. However, the *chs3Δ* mutant is still

viable and capable of divide, suggesting that additional chitin synthases might be

able to incorporate chitin in the absence of Chs3. This may explain why the

remedial-like septum is still formed in the *cdc55Δ chs3Δ* double mutant (Fig. 2i and

l).

Recently, it has been shown that multiple phosphatases shape the phospho-

proteome during mitotic exit[48,49,71]. Notably, Hof1 and Chs2 phosphosites have

been identified in *cdc55* and *cdc14* mutants[48]. We demonstrated that PP2A-Cdc55

regulates cytokinesis through the Chs2 dephosphorylation and it was reported that

Cdc14-dependent dephosphorylation is required for Chs2 release from the ER[41],

supporting different and specific contribution to cytokinesis of several phosphatases.

We observed hyperphosphorylations of Hof1 and Inn1 during cytokinesis in absence

of Cdc55 (Fig. S5a-b), suggesting that PP2A-Cdc55 might also regulates other IPCs

proteins. Further investigations will be required to understand the coordinated

regulation of multiple phosphatases during the cell cycle.

Materials and methods

Yeast strains, plasmids and cell-cycle synchronization procedures

All yeast strains used in this study were derivatives of W303 and are listed in Table

S1. Epitope tagging of endogenous genes and gene deletions were performed by

gene targeting using polymerase chain reaction (PCR) products[72]. Cell

synchronization using α-factor and metaphase arrest by Cdc20 depletion and entry

into synchronous anaphase by Cdc20 re-introduction were performed as previously

described[73]. To obtain the *pGAL1-CDC55* construct, the DNA fragment containing

the *GAL1* promoter was cut with SpeI and subcloned into the *CDC55* containing

plasmid previously digested with NheI.

All the *cdc55Δ* mutant strains were freshly prepared by transformation or by crossing

strains to avoid accumulation of suppressor mutations. The introduction of the

cdc55Δ deletion was determined by PCR-genotyping and observation of the slow

growth and elongated morphology under the microscope.

Recombinant protein purification

All plasmids were freshly introduced into BL21 *E. coli*. Cells were grown in LB

medium and protein expression was induced with 0.1 mM isopropyl β-D-1-

thiogalactopyranoside (IPTG) at 25ºC overnight. Collected cells were washed with

PBS1 and frozen for at least 30 minutes.

Cells containing His_6-Cyk3, His_6-Inn1 or His_6-Hof1 plasmids[17] were resuspended in

cold lysis buffer (30 mM Tris-HCl, pH 8, 300 mM NaCl, 30 mM imidazole, 0.1%

NP40, 10 mM β-mercaptoethanol, 1 mM PMSF, complete EDTA-free tablet (Roche))

and sonicated for 6 cycles of 1 min at 25 µm amplitude. Protein extracts were

clarified and incubated with Ni-NTA magnetic beads (Thermo Fisher) at 4ºC for 1

hour. Beads were washed with 10 volumes of lysis buffer and protein eluted in PBS,

5 mM EDTA, 5 mM DTT, 0.1% NP40, 500 mM imidazole at 4ºC for 30-60 minutes.

Cells containing StreptagIII-Chs2_1-629 plasmid[17] were resuspended in cold lysis

buffer (50 mM Tris-HCl, pH 8.0, 10% glycerol, 0.1% NP-40, 10 mM $MgCl_2$, 300 mM

NaCl, 5 mM β-mercaptoethanol, 10% BugBuster® (Millipore), 1 mM PMSF, complete

EDTA-free tablet (Roche)) containing 5 U/mL of nuclease (Pierce), incubated in a

rotatory wheel at 25ºC for 20 minutes. Protein extracts were clarified and incubated

with Strep-tactin® Superflow® resin (Iba Life science) at 4ºC for 1 hour. Beads were

washed with ten volumes of lysis buffer (without BugBuster®), and protein was

eluted in 50 mM Tris-HCl, pH 8.0, 10% glycerol, 0.1% NP-40, 10 mM $MgCl_2$, 150 mM

NaCl, 5 mM β-mercaptoethanol, 2.5 mM desthiobiotin at 4ºC for 1 hour.

Co-immunoprecipitation, kinase assays and phosphatase assays

Co-immunoprecipitation assays were performed using 10^8 yeast cells, which were

resuspended in lysis buffer (50 mM HEPES-KOH pH 7.5, 70 mM KOAc, 5 mM

$Mg(OAc)_2$, 10% glycerol, 0.1% Triton X-100, 8 µg/mL of protease inhibitors

(leupeptin, pepstatin, aprotinin), 1 mM PMSF, 1.25mg/ml of benzamidin, 1 tablet of

complete protease inhibitor without EDTA (Roche), 4 mM of phosphatase inhibitors

(β-glycerophosphate, NaF and NEM) and 1 tablet of PhosStop (Roche)). Lysates

were obtained by mechanical lysis using glass beads in a Bertin disrupter (6 cycles

of 10 s at 5,000 rpm). Protein extracts were clarified by centrifugation and incubated

with α-Pk clone SV5-Pk1 (AbSerotec) orα-HA clone 12CA5 (Roche) antibodies for 1

hour. Protein extracts were then incubated for 1 hour with protein A-conjugated

Dynabeads (Life Technologies), after which the beads were washed with lysis buffer

at incremental KOAc concentrations (100 mM, 120 mM, 150 mM, and 60 mM

NaOAc). Co-immunoprecipitated proteins were eluted with loading buffer and protein

co-purification was visualized by western blot.

For the Chs2 alkaline phosphatase assay, metaphase arrested cells were collected

and native protein extracts were prepared following the immunoprecipitation

protocol. The protein extracts were incubated with alkaline phosphatase (Anthartic

phosphatase, New England Biolabs) in EBX reaction buffer (50mM HEPES/KOH pH

7.5, 100mM KCl, 2.5mM $MgCl_2$, 0.25% Triton X-100, 1M DTT with protease inhibitors

(8μg/mL (leupeptin, pepstatin, aprotinin)), 1 mM PMSF, 1.25mg/ml of benzamidin, 1

tablet of complete protease inhibitor without EDTA (Roche)) at 37ºC for 15 minutes.

To inhibit the alkaline phosphatase 2XPhosStop (Roche) was used.

To purify the PP2A-Cdc55 complex, a strain with the tandem affinity protein (TAP)

epitope, TAP-HA-Cdc55 was used. Cells from 2 L cultures were arrested in

metaphase by Cdc20 depletion, frozen in liquid nitrogen and stored at -80ºC. Cells

were resuspended in lysis buffer (0.1 M NaCl, 50 mM Tris-HCl pH 7.5, 1.5 mM $MgCl_2$

and 0.15% NP-40, 8 μg/mL of protease inhibitors (leupeptin, pepstatin, aprotinin), 1

mM PMSF, 1.25mg/ml of benzamidin, 1 tablet of complete protease inhibitor without

EDTA (Roche)) and protein extracts prepared by mechanical lysis. Protein extracts

were clarified by centrifugation at 20.000 g and incubated with 0.4 mL IgG-

Sepharose beads for 1 hour. The beads were extensively washed and incubated

with 3 μL of TEV protease at 16ºC for 2 h to remove the CBP-HA-Cdc55 from the

beads. The purified PP2A-Cdc55 was stored at -80ºC and used for the phosphatase

assays in Fig 3d and Fig. S5f.

The Chs2-HA and Esp1-HA were purified following the co-immunoprecipitation

protocol as above but using Pierce HA-magnetic beads and eluted from the beads

using 1mg/mL of HA peptide. For the phosphatase assays with purified PP2A-

Cdc55, eluted Chs2 or Esp1 were incubated with phosphatase reaction buffer (50

mM Tris-HCl pH7.4, 0.1 mM EGTA, 1 mM β-mercaptoethanol, 1mg/mL BSA) and the

purified CBP-HA-Cdc55 at 30ºC for 20-40 minutes. Reactions were terminated by

adding SDS-PAGE loading buffer. Proteins were separated by electrophoresis,

transferred to PVDF membranes and phosphorylation levels detected with the anti-

phospho Ser/Thr antibody (Cell Signaling) using the Supersignal West Atto Ultimate

Sensitivity Substrate (Pierce). Anti-HA western blots were then performed to analyze

and quantify the amount of Chs2, Esp1 and Cdc55. The phosphorylation levels were

normalized against Cdc55 and Chs2 or Esp1. Proteins were quantified using Fiji

software[74] and values of the mean and SEM calculated.

For radioactive ^{32}P kinase assays, Clb2-Pk immunoprecipitations were performed as

above and incubated with α-Pk clone SV5-Pk1 (AbSerotec) antibody. Beads were

washed with ten volumes of lysis buffer and twice with the kinase reaction buffer (50

mM Tris-HCl, pH7.4, 10 mM MgCl$_2$, 1 mM DTT). The kinase reaction (50 mM Tris-

HCl, pH7.4, 10 mM MgCl$_2$, 1 mM DTT, 5 mM β-glycerophosphate, 25 µM ATP, 10

mCi/mL ^{32}gamma-ATP and 1 µg of Streptag-Chs2) was incubated at 30ºC for 30

minutes. Kinase assays were stopped by placing the tubes on ice. The supernatant

containing the phosphorylated substrate was separated from the magnetic beads

and stored at -80ºC. An aliquot of the kinase assay was mixed with SDS-PAGE

loading buffer, proteins were separated by electrophoresis, transferred to

nitrocellulose membranes and radioactivity detected in a Typhoon FLA950 (GE

Healthcare). Immunopurified protein was quantified by western blot and the

membrane was stained with Coomassie to detect the recombinant substrate.

Proteins were quantified using Fiji software.

For *in vitro* phosphatase assays, cells containing HA$_3$-*Cdc55-ED* and HA$_3$-Cdc55

were collected from metaphase arrested cells. Immunoprecipitation was performed

as above but without phosphatase inhibitors. Beads were washed twice with

phosphatase buffer and incubated with the phosphatase reaction (50 mM Tris-HCl

pH7.4, 0.1 mM EGTA, 1 mM β-mercaptoethanol, 1mg/mL BSA and the ^{32}P-Streptag-

Chs2) at 30ºC for 40 minutes. Reactions were terminated by adding SDS-PAGE

loading buffer. Proteins were separated by electrophoresis, transferred to

nitrocellulose membranes and radioactivity was detected in a Typhoon FLA950

apparatus (GE Healthcare). Later, the membrane was used to analyze and quantify

the amount of protein immunoprecipitated by western blot. Finally, the membrane

was stained with Coomassie to detect the recombinant Chs2 substrate. Proteins

levels were quantified using Fiji software and values of the mean and SEM

calculated.

The *GAL1-YFP-Chs2* and *GAL1-YFP-Chs2-S133A* were purified following the co-

immunoprecipitation protocol as above but using GFP-Trap beads (Chromotek).

Proteins were separated by electrophoresis, transferred to PVDF membranes and

phosphorylation levels detected with the anti-phospho Ser/Thr antibody (Cell

Signaling) using the Supersignal West Atto Ultimate Sensitivity Substrate (Pierce).

Anti-GFP western blots were then performed to analyze and quantify the amount of

Chs2. The phosphorylation levels were normalized against total Chs2. Protein levels

were quantified using Fiji software and values of the mean and SEM calculated.

Microscopy techniques

Synchronized cells for time-lapse experiments were deposited in chambers (Nunc

Lab-Tek) containing concanavalin A/PBS 1 mg/mL. Images were captured every 2

minutes. Different z-stacks at 0.7-µm intervals were taken and projected onto a

single image per channel. A Zeiss-Apotome epifluorescence microscope with an

HXP 120C fluorescent lamp and a Carl Zeiss Plan-Apochromat 63x N.A 1.40 oil

immersion lens were used. The filters used were Cy3, GFP, and DAPI. For the GFP-

Ras2 experiments the Apotome was disabled. For the Cyk3-GFP time-lapse

experiments, a Carl Zeiss LSM880 confocal microscope, with a 63x N.A. objective

was used. Images were acquired using ZEN software. For the time-lapses in figure

5, a Leica SP8 confocal microscope with a 63X objective was used. Images were

acquired using LAS software and quantified and processed using Fiji software. To

determine the symmetry of the IPCs proteins, the bud neck length was measured,

the central point calculated and manually established whether the central point was

inside (symmetry) or outside (asymmetry) the dot signal.

Calcofluor staining was performed with calcofluor white MR2 (Fluorescent brightener

28, Sigma) in living cells. Synchronized cells were incubated with 50 µg/mL

calcofluor upon release from the metaphase arrest. Images were acquired,

quantified and processed as for the time-lapse experiments.

Phalloidin staining was performed on fixed cells. Cells collected during the time-

course experiments were pre-fixed with PBS containing 3.7% formaldehyde and

0.1% Triton X-100 at 25ºC for 10 minutes. Cells were then washed with PBS1 and

fixed with 3.7% formaldehyde at 25ºC for 1 hour. After fixation, cells were washed

twice with PBS1 and sedimented onto a multi-well slide previously incubated with

poly-L-lysine. Cells were stained with a PBS solution containing 50 U/mL rhodamine

phalloidin R415 (Life Technologies) for 2 hours. Cells were washed twice with PBS1.

Mounting medium containing DAPI (Vectashield) was added, cells were covered with

a coverslip and sealed with nail polish. Images were acquired, quantified and

processed as for the time-lapse experiments.

For electron microscopy, cell samples were fixed with 0.2 M phosphate buffer

(without salts) pH 7.4, containing 2.5% glutaraldehyde at 25ºC, for 1 hour. Cells were

washed three times with phosphate buffer without glutaraldehyde and rinsed with

milli-Q water and post-fixed with 1% osmium tetroxide for 2 hours. They were then

dehydrated in an acetone series (10%, 20%, 30%, 40%, 60%, 80%, and 100%) for

15-20 minutes. Ultrathin sections of 60-nm thickness were obtained using a UC6

ultramicrotome (Leica Microsystems, Austria) and stained with 2% uranyl acetate

and lead citrate. Sections were observed in a Jeol EM J1010 apparatus (Jeol,

Japan) and images were acquired at 80 kV with a 1k CCD Megaview camera.

SILAC labeling and phosphoproteomic analysis

Y859 (*MATa lys2Δ::TRP1, arg4Δ::HIS3 MET-CDC20::LEU2*) and Y858 (as Y859,

but *cdc55Δ*) cells were grown in free-methionine minimum media containing $^{13}C_6$-

lysine and -arginine (heavy, Cambridge Isotope Laboratories Inc., US) or

unmodified arginine and lysine (light), respectively. Protein extracts were prepared

by mechanical lysis using glass beads in presence of protein inhibitors (Complete

EDTA-free, Roche) and 2x phosphatase inhibitors PhosStop (Roche). Cell lysates

were mixed 1:1 and digestion with trypsin was performed. The phosphopeptides

were pre-enriched using TiO_2 chromatography (GL Sciences Inc) followed by

SIMAC purification. The mono-phosphorylated peptide fraction from the SIMAC

enrichment was further subjected to a second TiO_2 purification and pre-fractionated

by HILIC chromatography (Hydrophilic Interaction Liquid Chromatography). The

peptides were then analyzed by LC-MS/MS using a LTQ-Orbitrap Fusion Tribride

mass spectrometer (Thermo Fisher Scientific). Peptide identification was performed

using Proteome Discoverer v1.4.1.14 (Thermo Scientific) and search against Swiss

Prot/Uniprot *Saccharomyces cerevisiae* database. Peptide quantification from

SILAC labels was performed with Proteome Discoverer v1.4.

Other techniques

Protein extracts for western blots were obtained by TCA protein extraction. For the

phostag gels, 10 µM Phostag (Wako) 5-8% gels containing 10 mM $MnCl_2$ were

used. Protein gels were washed with 1 mM EDTA before transferring protein to

remove the manganese. Proteins were quantified using Fiji software, and values of

the mean and SEM calculated. To calculate the ratio of

phosphorylated/dephosphorylated Chs2, the faster migration band corresponds to

the dephosphorylated Chs2 (as shown in the alkaline phosphatase experiment Fig.

S5e). The slower migration band observed in metaphase is considered the

phosphorylated Chs2. For each triplicate, the signals were normalized against the

higher signal in the blot. To assign the cells to metaphase, anaphase and

cytokinesis we took in consideration the percentage of anaphase spindles and the

FACS profiles (not shown). Time-points with intermediate percentages of anaphase

spindles were not used in the quantification.

Antibodies used for western blots were α-HA clone 12CA5 (Roche), α-myc 9E10

(Babco), α-FLAG clone M2 (Sigma), α-Pk clone SV5-Pk1 (AbSerotec), α-Clb2 (y-

180) sc-907 (Santa Cruz Biotechnology), α-tubulin clone YOL1/34 (Serotec), α-

phosphoglycerate kinase (Life Technologies), α-HA rabbit (Sigma), α-Pgk1

(Invitrogen), and α-Chs2[17]. The secondary antibodies were: α-Mouse-HRP (GE

Healthcare), α-Rabbit-HRP (GE Healthcare), and α-Goat-HRP (GE Healthcare).

Antibodies used for immunofluorescence were: α-HA clone 12CA5 (Roche), α-Cdc14

(yE-17) sc-12045 (Santa Cruz Biotechnology), α-tubulin clone YOL1/34 (AbSerotec),

and α-Cdc11 (Santa Cruz Biotechnology). The secondary antibodies were Cy3-

labeled α-mouse (GE Healthcare), fluorescein-conjugated α-rat (Millipore), Cy3-

labeled α-goat (GE Healthcare), red TEXAS α-rabbit (Jackson Laboratories) and 488

α-mouse (Life Technologies).

Statistical analysis

All experiments were done at least three times. Statistical analyses were performed

with Prism5. Student's T-test was used to analyze the p-values of the different

assays compared. A p-value <0.05 was considered statistically significant,

p<0.0001 ****, p<0.001 ***, p < 0.01 **; p<0.05 *.

Acknowledgements

We thank the optical microscopy unit of IDIBELL, the electron microscopy units and

the proteomic services from IDIBELL and CCiT-UB for their support. The *cdc55-aid*

mutant strain was a kind gift of Adam Rudner and the *GAL-CHS2-YFP* containing

plasmids were a kind gift of Foong May Yeong. We thank all the members of our

laboratory for discussing the work and for their critical reading of the manuscript. We

thank CERCA Program/Generalitat de Catalunya for institutional support. EQ is

funded by the grants BFU2016-77975-R (co-funded by the European Regional

Development Fund, ERDF, a way to build Europe) from the Spanish Ministry of

Economy, Industry and Competitiveness (MINECO) and PID2019-109027GB-I00

from the Spanish Ministry of Science, Innovation and Universities (MCIU). ASD was

supported by the grant PID2019-106745GB-I00 from the Spanish Spanish Ministry of

Science, Innovation and Universities (co-funded by the European Regional

Development Fund) and a grant from the Consejería de Universidades,

Investigación, Medio Ambiente y Política Social del Gobierno de Cantabria.

Author contribution: YMR, DV, OVC, MF and EQ performed the experiments.
YMR, ASD and EQ designed the experiments and interpreted the data. YMR and EQ
wrote the manuscript. All authors read and discussed the manuscript.

Conflict of interest

The authors declare no competing interests.

References

1.   Fujiwara T, Bandi M, Nitta M, Ivanova E V., Bronson RT, Pellman D.

Cytokinesis failure generating tetraploids promotes tumorigenesis in p53-null

cells. Nature. 2005;437: 1043–7. doi:10.1038/nature04217

2.   Okada S, Leda M, Hanna J, Savage NS, Bi E, Goryachev AB. Daughter Cell

Identity Emerges from the Interplay of Cdc42, Septins, and Exocytosis. Dev

Cell. 2013;26: 148–61. doi:10.1016/j.devcel.2013.06.015

3. Schneider C, Grois J, Renz C, Gronemeyer T, Johnsson N. Septin rings act as

a template for myosin higher-order structures and inhibit redundant polarity

establishment. J Cell Sci. 2013;126: 3390–400. doi:10.1242/jcs.125302

4. Ong K, Wloka C, Okada S, Svitkina T, Bi E. Architecture and dynamic

remodelling of the septin cytoskeleton during the cell cycle. Nat Commun.

2014;5: 5698. doi:10.1038/ncomms6698

5. Lippincott J, Li R. Dual function of Cyk2, a cdc15/PSTPIP family protein, in

regulating actomyosin ring dynamics and septin distribution. J Cell Biol.

1998;143: 1947–1960. doi:10.1083/jcb.143.7.1947

6. Oh Y, Schreiter J, Nishihama R, Wloka C, Bi E. Targeting and functional

mechanisms of the cytokinesis-related F-BAR protein Hof1 during the cell

cycle. Mol Biol Cell. 2013;24: 1305–20. doi:10.1091/mbc.E12-11-0804

7. Meitinger F, Palani S, Hub B, Pereira G. Dual function of the NDR-kinase Dbf2

in the regulation of the F-BAR protein Hof1 during cytokinesis. Mol Biol Cell.

2013; 1–44. doi:10.1091/mbc.E12-08-0608

8. Finnigan GC, Booth EA, Duvalyan A, Liao EN, Thorner J. The carboxy-

terminal tails of septins Cdc11 and Shs1 recruit myosin-II binding factor Bni5 to

the bud neck in Saccharomyces cerevisiae. Genetics. 2015;200: 843–62.

doi:10.1534/genetics.115.176503

9. Shannon KB, Li R. The multiple roles of Cyk1p in the assembly and function of

the actomyosin ring in budding yeast. Mol Biol Cell. 1999;10: 283–96.

doi:10.1091/mbc.10.2.283

10. Fang X, Luo J, Nishihama R, Wloka C, Dravis C, Travaglia M, et al. Biphasic

targeting and cleavage furrow ingression directed by the tail of a myosin II. J

Cell Biol. 2010;191: 1333–50. doi:10.1083/jcb.201005134

11.   Mendes Pinto I, Rubinstein B, Kucharavy A, Unruh JR, Li R. Actin

Depolymerization Drives Actomyosin Ring Contraction during Budding Yeast

Cytokinesis. Dev Cell. 2012;22: 1247–60. doi:10.1016/j.devcel.2012.04.015

12.   Chew TG, Huang J, Palani S, Sommese R, Kamnev A, Hatano T, et al. Actin

turnover maintains actin filament homeostasis during cytokinetic ring

contraction. J Cell Biol. 2017;216: 2657–2667. doi:10.1083/jcb.201701104

13.   Nishihama R, Schreiter JH, Onishi M, Vallen EA, Hanna J, Moravcevic K, et al.

Role of Inn1 and its interactions with Hof1 and Cyk3 in promoting cleavage

furrow and septum formation in S. cerevisiae. J Cell Biol. 2009;185: 995–1012.

doi:10.1083/jcb.200903125

14.   Jendretzki A, Ciklic I, Rodicio R, Schmitz HP, Heinisch JJ. Cyk3 acts in

actomyosin ring independent cytokinesis by recruiting Inn1 to the yeast bud

neck. Mol Genet Genomics. 2009;282: 437–51. doi:10.1007/s00438-009-0476-

0

15.   Wloka C, Vallen EA, Thé L, Fang X, Oh Y, Bi E. Immobile myosin-II plays a

scaffolding role during cytokinesis in budding yeast. J Cell Biol. 2013;200:

271–86. doi:10.1083/jcb.201208030

16.   Devrekanli A, Foltman M, Roncero C, Sanchez-Diaz A, Labib K. Inn1 and

Cyk3 regulate chitin synthase during cytokinesis in budding yeasts. J Cell Sci.

2012;125: 5453–66. doi:10.1242/jcs.109157

17.   Foltman M, Molist I, Arcones I, Sacristan C, Filali-Mouncef Y, Roncero C, et al.

Ingression Progression Complexes Control Extracellular Matrix Remodelling

during Cytokinesis in Budding Yeast. PLoS Genet. 2016;12: e1005864.

doi:10.1371/journal.pgen.1005864

18.   Wang M, Nishihama R, Onishi M, Pringle JR. Role of the Hof1–Cyk3

770   interaction in cleavage-furrow ingression and primary-septum formation during

771   yeast cytokinesis. Mol Biol Cell. 2018;29: 597–609. doi:10.1091/mbc.E17-04-

772   0227

773   19.   Sburlati A, Cabib E. Chitin synthetase 2, a presumptive participant in septum

774   formation in Saccharomyces cerevisiae. J Biol Chem. 1986;261: 15147–52.

775   20.   Shaw JA, Mol PC, Bowers B, Silverman SJ, Valdivieso MH, Duran A, et al.

776   The function of chitin synthases 2 and 3 in the Saccharomyces cerevisiae cell

777   cycle. J Cell Biol. 1991;114: 111–23. doi:10.1083/jcb.114.1.111

778   21.   Okada H, MacTaggart B, Ohya Y, Bi E. The kinetic landscape and interplay of

779   protein networks in cytokinesis. iScience. 2021;24: 101917.

780   doi:10.1016/j.isci.2020.101917

781   22.   Foltman M, Filali-Mouncef Y, Crespo D, Sanchez-Diaz A. Cell polarity protein

782   Spa2 coordinates Chs2 incorporation at the division site in budding yeast.

783   PLoS Genet. 2018;14: e1007299. doi:10.1371/journal.pgen.1007299

784   23.   Wloka C, Bi E. Mechanisms of cytokinesis in budding yeast. Cytoskeleton.

785   2012;69: 710–726. doi:10.1002/cm.21046

786   24.   Schmidt M, Bowers B, Varma A, Roh DH, Cabib E. In budding yeast,

787   contraction of the actomyosin ring and formation of the primary septum at

788   cytokinesis depend on each other. J Cell Sci. 2002;115: 293–302.

789   25.   VerPlank L, Li R. Cell cycle-regulated trafficking of Chs2 controls actomyosin

790   ring stability during cytokinesis. Mol Biol Cell. 2005;16: 2529–43.

791   doi:10.1091/mbc.E04-12-1090

792   26.   Sanchez-Diaz A, Marchesi V, Murray S, Jones R, Pereira G, Edmondson R, et

793   al. Inn1 couples contraction of the actomyosin ring to membrane ingression

during cytokinesis in budding yeast. Nat Cell Biol. 2008;10: 395–406.

doi:10.1038/ncb1701

27. Cabib E, Sburlati A, Bowers B, Silverman SJ. Chitin synthase 1, an auxiliary

enzyme for chitin synthesis in Saccharomyces cerevisiae. J Cell Biol.

1989;108: 1665–72. doi:10.1083/jcb.108.5.1665

28. Douglas CM, Foor F, Marrinan JA, Morin N, Nielsen JB, Dahl AM, et al. The

Saccharomyces cerevisiae FKS1 (ETG1) gene encodes an integral membrane

protein which is a subunit of 1,3-β-D-glucan synthase. Proc Natl Acad Sci U S

A. 1994;91: 12907–11. doi:10.1073/pnas.91.26.12907

29. Weiss EL. Mitotic exit and separation of mother and daughter cells. Genetics.

2012;192: 1165–1202. doi:10.1534/genetics.112.145516

30. Queralt E, Uhlmann F. Cdk-counteracting phosphatases unlock mitotic exit.

Curr Opin Cell Biol. 2008/10/11. 2008;20: 661–668. doi:S0955-

0674(08)00161-0 [pii] 10.1016/j.ceb.2008.09.003

31. Baro B, Queralt E, Monje-Casas F. Regulation of mitotic exit in saccharomyces

cerevisiae. Methods in Molecular Biology. 2017. pp. 3–17. doi:10.1007/978-1-

4939-6502-1_1

32. Rock JM, Amon A. The FEAR network. Curr Biol. 2009;19: R1063-8.

doi:10.1016/j.cub.2009.10.002

33. Hotz M, Barral Y. The Mitotic Exit Network: New turns on old pathways. Trends

in Cell Biology. 2014. pp. 145–152. doi:10.1016/j.tcb.2013.09.010

34. Visintin R, Craig K, Hwang ES, Prinz S, Tyers M, Amon A. The Phosphatase

Cdc14 Triggers Mitotic Exit by Reversal of Cdk-Dependent Phosphorylation.

Mol Cell. 1998;2: 709–718. doi:10.1016/S1097-2765(00)80286-5

35. Jiménez J, Castelao BA, González-Novo A, Sánchez-Pérez M. The role of

MEN (mitosis exit network) proteins in the cytokinesis of Saccharomyces

cerevisiae. Int Microbiol. 2005;8: 33–42. doi:im2305005 [pii]

36.   Meitinger F, Boehm ME, Hofmann A, Hub B, Zentgraf H, Lehmann WD, et al.

Phosphorylation-dependent regulation of the F-BAR protein Hof1 during

cytokinesis. Genes Dev. 2011/04/19. 2011;25: 875–888.

doi:10.1101/gad.622411

37.   Yoshida S, Kono K, Lowery DM, Bartolini S, Yaffe MB, Ohya Y, et al. Polo-like

kinase Cdc5 controls the local activation of Rho1 to promote cytokinesis.

Science (80-). 2006;313: 108–111. doi:10.1126/science.1126747

38.   Naylor SG, Morgan DO. Cdk1-dependent phosphorylation of iqg1 governs

actomyosin ring assembly prior to cytokinesis. J Cell Sci. 2014;127: 1128–37.

doi:10.1242/jcs.144097

39.   Miller DP, Hall H, Chaparian R, Mara M, Mueller A, Hall MC, et al.

Dephosphorylation of Iqg1 by Cdc14 regulates cytokinesis in budding yeast.

Mol Biol Cell. 2015;26: 2913–26. doi:10.1091/mbc.E14-12-1637

40.   Zhang G, Kashimshetty R, Kwee EN, Heng BT, Foong MY. Exit from mitosis

triggers Chs2p transport from the endoplasmic reticulum to mother-daughter

neck via the secretory pathway in budding yeast. J Cell Biol. 2006;174: 207–

20. doi:10.1083/jcb.200604094

41.   Chin CF, Bennett AM, Ma WK, Hall MC, Yeong FM. Dependence of Chs2 ER

export on dephosphorylation by cytoplasmic Cdc14 ensures that septum

formation follows mitosis. Mol Biol Cell. 2012;23: 45–58. doi:10.1091/mbc.E11-

05-0434

42.   Jakobsen MK, Cheng Z, Lam SK, Roth-Johnson E, Barfield RM, Schekman R.

Phosphorylation of Chs2p regulates interaction with COPII. J Cell Sci.

2013;126: 2151–6. doi:10.1242/jcs.115915

43.   Meitinger F, Petrova B, Lombardi IM, Bertazzi DT, Hub B, Zentgraf H, et al.

Targeted localization of Inn1, Cyk3 and Chs2 by the mitotic-exit network

regulates cytokinesis in budding yeast. J Cell Sci. 2010/05/06. 2010;123:

1851–1861. doi:jcs.063891 [pii] 10.1242/jcs.063891

44.   Palani S, Meitinger F, Boehm ME, Lehmann WD, Pereira G. Cdc14-dependent

dephosphorylation of Inn1 contributes to Inn1-Cyk3 complex formation. J Cell

Sci. 2012/03/29. 2012;125: 3091–3096. doi:10.1242/jcs.106021

45.   Sanchez-Diaz A, Nkosi PJ, Murray S, Labib K, Sanchez☐Diaz A, Nkosi PJ, et

al. The Mitotic Exit Network and Cdc14 phosphatase initiate cytokinesis by

counteracting CDK phosphorylations and blocking polarised growth. EMBO J.

2012;31: 3620–3634. doi:10.1038/emboj.2012.224

46.   Kuilman T, Maiolica A, Godfrey M, Scheidel N, Aebersold R, Uhlmann F.

Identification of Cdk targets that control cytokinesis. EMBO J. 2015;34: 81–96.

doi:10.15252/embj.201488958

47.   Teh EM, Chai CC, Yeong FM. Retention of Chs2p in the ER requires N-

terminal CDK1-phosphorylation sites. Cell Cycle. 2009;8: 2964–74.

48.   Touati SA, Hofbauer L, Jones AW, Snijders AP, Kelly G, Uhlmann F. Cdc14

and PP2A Phosphatases Cooperate to Shape Phosphoproteome Dynamics

during Mitotic Exit. Cell Rep. 2019;29: 2015–2119.

doi:10.1016/j.celrep.2019.10.041

49.   Touati SA, Kataria M, Jones AW, Snijders AP, Uhlmann F. Phosphoproteome

dynamics during mitotic exit in budding yeast. EMBO J. 2018;37: e98745.

doi:10.15252/embj.201798745

50.   Gentry MS, Hallberg RL. Localization of Saccharomyces cerevisiae protein

phosphatase 2A subunits throughout mitotic cell cycle. Mol Biol Cell. 2002;13:

3477–3492. doi:10.1091/mbc.02-05-0065

51. Dobbelaere J, Barral Y. Spatial coordination of cytokinetic events by

compartmentalization of the cell cortex. Science (80-). 2004;305: 393–6.

doi:10.1126/science.1099892

52. Lippincott J, Li R. Sequential assembly of myosin II, an IQGAP-like protein,

and filamentous actin to a ring structure involved in budding yeast cytokinesis.

J Cell Biol. 1998;140: 355–66. doi:10.1083/jcb.140.2.355

53. McQuilken M, Jentzsch MS, Verma A, Mehta SB, Oldenbourg R, Gladfelter

AS. Analysis of septin reorganization at cytokinesis using polarized

fluorescence microscopy. Front Cell Dev Biol. 2017;5: 42.

doi:10.3389/fcell.2017.00042

54. Jonasson EM, Rossio V, Hatakeyama R, Abe M, Ohya Y, Yoshida S.

Zds1/Zds2–PP2A(Cdc55) complex specifies signaling output from Rho1

GTPase. J Cell Biol. 2016;212: 51–61. doi:10.1083/jcb.201508119

55. Kennedy EK, Dysart M, Lianga N, Williams EC, Pilon S, Doré C, et al.

Redundant regulation of Cdk1 tyrosine dephosphorylation in Saccharomyces

cerevisiae. Genetics. 2016;202: 903–10. doi:10.1534/genetics.115.182469

56. Minshull J, Straight A, Rudner AD, Dernburg AF, Belmont A, Murray AW.

Protein phosphatase 2A regulates MPF activity and sister chromatid cohesion

in budding yeast. Curr Biol. 1996;6: 1609–1620. doi:10.1016/S0960-

9822(02)70784-7

57. Queralt E, Lehane C, Novak B, Uhlmann F. Downregulation of PP2ACdc55

Phosphatase by Separase Initiates Mitotic Exit in Budding Yeast. Cell.

2006;125: 719–732. doi:10.1016/j.cell.2006.03.038

58.   Walker LA, Lenardon MD, Preechasuth K, Munro CA, Gow NAR. Cell wall

895   stress induces alternative fungal cytokinesis and septation strategies. J Cell

896   Sci. 2013;126: 2668–2677. doi:10.1242/jcs.118885

897   59.   Cabib E, Schmidt M. Chitin synthase III activity, but not the chitin ring, is

898   required for remedial septa formation in budding yeast. FEMS Microbiol Lett.

899   2003;224: 299–305. doi:10.1016/S0378-1097(03)00477-4

900   60.   Korinek WS, Bi E, Epp JA, Wang L, Ho J, Chant J. Cyk3, a novel SH3-domain

901   protein, affects cytokinesis in yeast. Curr Biol. 2000;10: 947–50.

902   doi:10.1016/S0960-9822(00)00626-6

903   61.   Nishimura K, Fukagawa T, Takisawa H, Kakimoto T, Kanemaki M. An auxin-

904   based degron system for the rapid depletion of proteins in nonplant cells. Nat

905   Methods. 2009;6: 917–22. doi:10.1038/nmeth.1401

906   62.   Játiva S, Calabria I, Moyano-Rodriguez Y, Garcia P, Queralt E. Cdc14

907   activation requires coordinated Cdk1-dependent phosphorylation of Net1 and

908   PP2A–Cdc55 at anaphase onset. Cell Mol Life Sci. 2019;76: 3601–3620.

909   doi:10.1007/s00018-019-03086-5

910   63.   Baro B, Játiva S, Calabria I, Vinaixa J, Bech-Serra JJ, De LaTorre C, et al.

911   SILAC-based phosphoproteomics reveals new PP2A-Cdc55-regulated

912   processes in budding yeast. Gigascience. 2018;7: giy407.

913   doi:10.1093/gigascience/giy047

914   64.   Holt LJ, Tuch BB, Villén J, Johnson AD, Gygi SP, Morgan DO, et al. Global

915   analysis of Cdk1 substrate phosphorylation sites provides insights into

916   evolution. Science. 2009;325: 1682–6. doi:10.1126/science.1172867

917   65.   Baro B, Rodriguez-Rodriguez J-A, Calabria I, Hernáez ML, Gil C, Queralt E.

918   Dual Regulation of the Mitotic Exit Network (MEN) by PP2A-Cdc55

919   Phosphatase. PLoS Genet. 2013;9: e1003966.

920   doi:10.1371/journal.pgen.1003966

921 66. Vallen EA, Caviston J, Bi E. Roles of Hof1p, Bni1p, Bnr1p, and Myo1p in

922   cytokinesis in Saccharomyces cerevisiae. Mol Biol Cell. 2000;11: 593–611.

923   doi:10.1091/mbc.11.2.593

924 67. Dobbelaere J, Gentry MS, Hallberg RL, Barral Y. Phosphorylation-dependent

925   regulation of septin dynamics during the cell cycle. Dev Cell. 2003;4: 345–57.

926   doi:10.1016/S1534-5807(03)00061-3

927 68. Healy AM, Zolnierowicz S, Stapleton AE, Goebl M, DePaoli-Roach AA, Pringle

928   JR. CDC55, a Saccharomyces cerevisiae gene involved in cellular

929   morphogenesis: identification, characterization, and homology to the B subunit

930   of mammalian type 2A protein phosphatase. Mol Cell Biol. 1991;11: 5767–

931   5780. doi:10.1128/mcb.11.11.5767

932 69. Park SY, Cable AE, Blair J, Stockstill KE, Shannnon KB. Bub2 regulation of

933   cytokinesis and septation in budding yeast. BMC Cell Biol. 2009;10: 43.

934   doi:10.1186/1471-2121-10-43

935 70. Oh Y, Chang KJ, Orlean P, Wloka C, Deshaies R, Bi E. Mitotic exit kinase

936   Dbf2 directly phosphorylates chitin synthase Chs2 to regulate cytokinesis in

937   budding yeast. Mol Biol Cell. 2012/05/11. 2012;23: 2445–2456. doi:mbc.E12-

938   01-0033 [pii] 10.1091/mbc.E12-01-0033

939 71. Godfrey M, Touati SA, Kataria M, Jones A, Snijders AP, Uhlmann F.

940   PP2A(Cdc55) Phosphatase Imposes Ordered Cell-Cycle Phosphorylation by

941   Opposing Threonine Phosphorylation. Mol Cell. 2017;65: 393-402.e3.

942   doi:10.1016/j.molcel.2016.12.018

943 72. Knop M, Siegers K, Pereira G, Zachariae W, Winsor B, Nasmyth K, et al.

Epitope tagging of yeast genes using a PCR-based strategy: More tags and

improved practical routines. Yeast. 1999;15: 963–972. doi:10.1002/(SICI)1097-

0061(199907)15:10B<963::AID-YEA399>3.0.CO;2-W

73.   Monje-Casas F, Queralt E. The Mitotic Exit Network. Fernando Monje-Casas

EQ, editor. Humana Press; 2017. doi:10.1007/978-1-4939-6502-1

74.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et

al. Fiji: An open-source platform for biological-image analysis. Nat Methods.

2012;9: 676–82. doi:10.1038/nmeth.2019

Figure legends

Figure 1. PP2A-Cdc55 regulates AMR constriction and septum formation. (a)

Absence of Cdc55 promotes the asymmetry of Myo1upon AMR contraction. Strains

Y1306 and Y1578 were arrested in metaphase and released into anaphase by

Cdc20 depletion and re-addition, and time-lapse images were captured every 2

minutes. Spc42-GFP (spindle pole body protein) was used as the control for

anaphase progression. Representative images of Myo1 in *CDC55* and *cdc55Δ* are

shown (left panel). Quantification of the contraction time of Myo1 is depicted (right

panel). (b) Quantification of the population of cells with asymmetric Myo1 constriction

in *CDC55* (N=76) and *cdc55Δ* (N=69) from (a). (c) Myo1 asymmetric localization

upon degradation or inactivation of Cdc55. Strains Y1652 (N=9), Y1653 (N=17),

Y1788 (N=35), Y1605 (N=22) and Y1596 (N=24) were arrested in metaphase and

released into anaphase by Cdc20 depletion and re-addition. *cdc55-aid* degradation

was induced by the addition of 0.5 mM auxin for 3 hours during the metaphase arrest

and auxin was maintained after Cdc20 re-addition. Time-lapse images were

captured every 2 minutes. Quantification of the cell population with asymmetric Myo1

constriction (left panel), and maximum intensity z-projection images of Myo1 (right

panel) are shown. Scale bar, 1µm. (d-e) Chitin deposition at primary and secondary

septa is regulated by PP2A-Cdc55. Strains Y1516 (N=46), Y1512 (N=49), Y1605

(N=82) and Y1596 (N=86) were arrested in metaphase by Cdc20 depletion and

released into anaphase by Cdc20 re-induction. Calcofluor was added to visualize the

septa upon the metaphase release. Student's unpaired t-test was carried out using

the Prism5 program.

Figure 2. Asymmetric primary septum formation and appearance of the

remedial-like septum in the absence of Cdc55. Strains Y1315, Y1318, Y1605,

and Y1596 were arrested in metaphase and released into anaphase by Cdc20

depletion and re-addition. Representative images from TEM are shown. Scale bar,

0.5 µm. PS, SS and RS denote primary septum, secondary septa and remedial

septum, respectively.

Figure 3. PP2A-Cdc55 dephosphorylates Chs2 during cytokinesis. (a) The

double mutants *cdc55 hof1* and *cdc55cyk3* are synthetically sick. Serial dilutions

ofW303, Y844, Y1728, Y1730, Y1747 and Y1749 strains were spotted on YPD

plates with and without 1.5 mM auxin (IAA). Cells were grown at 25ºC for 2-3 days.

(b) PP2A-Cdc55 forms a complex with Chs2. Strain Y1567 was synchronized into

anaphase progression by Cdc20 depletion and re-addition. Strain Y1318 without the

Pk-tag was used as negative control. Protein extracts were prepared at metaphase,

anaphase, and cytokinesis and Cdc55 was immunoprecipitated using the anti-Pk

antibody. Co-purification of Chs2 was analyzed by western blot. (c) PP2A-Cdc55

regulates Chs2 dephosphorylation. Strains Y1318 and Y1315 were arrested in

metaphase and released into anaphase by Cdc20 depletion and re-addition. Chs2

phosphorylation was analyzed by western blot in Phos-tag gels. Unspecific bands

were detected for Chs2 and marked as ([a]). Pgk1 levels were used as a loading

control. Mitosis progression was followed by analyzing the anaphase spindle

elongation by *in situ* immunofluorescence. At least 100 cells were scored at each

time-point. Quantification of the western blots was performed using Fiji Software and

means and SEMs are represented. Student's unpaired t-test analyses were carried

out using the Prism5 program. (d) Chs2 is dephosphorylated by PP2A-Cdc55.

Metaphase arrested cells of the strain Y695 were used to purify the PP2A-Cdc55

complex by TAP purification. Chs2-HA was purified from metaphase arrested cells of

strain Y1318. Purified Chs2-HA was incubated with the PP2A-Cdc55 complex at the

indicated times and the Chs2 phosphorylation levels were detected by western blot

using the anti-phospho Ser/Thr antibody. Representative images of one

phosphatase assay are shown. Protein levels were quantified using Fiji software.

Quantifications of the remaining Chs2 phosphorylation signal normalized to the

amount of Cdc55 and Chs2 are shown. Means and SEMs of three phosphatase

assays are represented. Student's unpaired t-test analysis was carried out using the

Prism5 program. (e) PP2A-Cdc55 dephosphorylates Chs2 *in vitro*. Strains Y824,

Y1652 and Y1653 were arrested in metaphase by Cdc20 depletion. Cdc55 and

cdc55-ED were purified by immunoprecipitation with HA antibody. Strep-tag-Chs2-1-

629 purified from *E. coli* and previously phosphorylated by Cdk1-Clb2 in a

radioactive kinase assay was used as substrate. Representative images of the

phosphatase assays are shown. Radioactive signals were detected using a multi-

purpose imaging plate in a Typhoon FLA950 apparatus (GE healthcare). Protein

levels were quantified using Fiji software. Quantifications of the remaining

phosphorylation signals normalized with respect to the amount of immunopurified

HA-Cdc55 or HA-*cdc55-ED* are shown. Means and SEMs of three phosphatase

assays are represented. Student's unpaired t-test analysis was carried out using the

Prism5 program.

1024

Figure 4. PP2A-Cdc55 is required for proper IPCs localization and contraction

at the division site. Strains Y1572 (N=27), Y1606 (N=28), Y1454 (N=14), Y1608

(N=41), Y1306 (N=32), Y1578 (N=12), Y1574 (N=33), Y1604 (N=11), Y1576 (N=10)

and Y1575 (N=8) were synchronized in mitosis by Cdc20 depletion and re-addition,

and time-lapse images were captured every 2 minutes. Spc42-GFP and Myo1-

tdTomato were used as a control for cytokinesis progression. (a) Percentages of

cells with IPCs asymmetric Myo1 contraction are shown (left panel). Representative

images of GFP-tagged proteins in *CDC5*5 and *cdc55Δ* are depicted (right panel). (b)

Quantifications of the IPCs contraction times. Student's unpaired t-tests were

performed using the Prism5 program. Scale bar, 1 µm.

1035

Figure 5. Chs2 phospho-mutants at S133 regulate AMR contraction and Hof1

residence time at the division site. (a) Hyperphosphorylation of Chs2 in S133 was

detected by mass spectrometry in absence of Cdc55. Representation of the Chs2

peptide detected by mass spectrometry containing the S133. (b) Chs2

phosphorylation is greatly reduced in the *chs2-S133A* mutant. Strains Y1901 and

Y1900 were arrested in metaphase and *GAL1-CHS2* versions were induced by

galactose addition for 3 h. Protein extracts were prepared and Chs2 and *chs2-*

S133A purified using GFP-trap columns. Chs2 phosphorylation levels were detected

by western blot using the anti-phospho Ser/Thr antibody. Representative blots of one

assay are shown (left panel). Western blot signals were quantified using Fiji

software. Quantifications of the Chs2 phosphorylation levels normalized to the

amount of Chs2 from three independent experiments are shown (right panel). Means

and SEMs are represented. Student's unpaired t-test analysis was performed using

the Prism5 program. (c) The non-phosphorylable *chs2-S133A* version rescues the

asymmetric AMR contraction in the *cdc55Δ* mutant cells. Strains Y1896 (N=25) and

Y1898 (N=31) were arrested at metaphase by Cdc20 depletion. *GAL1-CHS2*

versions were induced by galactose addition 3 hours before synchronous release

into anaphase by Cdc20 re-introduction, and time-lapse images were captured every

2 minutes to visualize Myo1 and Chs2. Representative images of Myo1 and Chs2

are shown (left panel). Percentages of cells with asymmetric Myo1 are represented

(right panel). (d) Similar Hof1 contraction times of Chs2 and *chs2-133A* in

cdc55Δ mutant cells. Strains Y1912 (N=36) and Y1913 (N=38) were arrested at

metaphase by Cdc20 depletion and *GAL1-CHS2* versions induced as in (c). Time-

lapse images were captured every 3 minutes to visualize Hof1 and Chs2.

Quantifications of the Hof1 contraction times are represented. Student's unpaired t-

tests were performed using the Prism5 program. (e) The phospho-mimetic *chs2-*

S133E is not able to induce asymmetric Myo1 signal. Strains Y1895 (N=35) and

Y1893 (N=29) were arrested at metaphase by Cdc20 depletion, *GAL1-CHS2*

versions induced as in (c), and time-lapse images were captured every 2 minutes to

visualize Myo1 and Chs2. Representative images of Myo1 and Chs2 are shown (left

panel). Percentages of cells with asymmetric Myo1 are represented (right panel). (f)

The phospho-mimetic *chs2-S133E* version promotes longer Hof1 residence time at

the division site. Strains Y1909 (N=25) and Y1911 (N=26), were arrested at

metaphase by Cdc20 depletion and *GAL1-CHS2* versions induced as in (c). Time-

lapse images were captured every 3 minutes to visualize Hof1 and Chs2.

Quantifications of the Hof1 contraction times are represented. Student's unpaired t-

tests were performed using the Prism5 program.

1073

1074 **Figure 6. Model for PP2A-Cdc55 regulation of actomyosin ring (AMR)**

1075 **contraction, primary septum (PS) and secondary septa (SS) formation**.

1076 Dephosphorylation of Chs2 by PP2A-Cdc55 coordinates AMR contraction and

1077 primary septum formation resulting in an increased Hof1 residence time at the

1078 division site.

1079

1080 **Supplemental Figure legends**

1081

1082 **Figure S1. Septin structures are not altered in the absence of Cdc55.** Cycling

1083 cells from strains Y1588 and Y1589 were fixed and immunofluorescence *in situ*

1084 performed for septins Cdc11 and Shs1-HA visualization. α-Cdc11 and α-HA (12CA5)

1085 antibodies were used. Representative images of Cdc11 and Shs1 septin' structures

1086 from *CDC55* and *cdc55Δ* cells are shown. Scale bar, 1 μm.

1087

Figure S2. Actin is polarized to the new bud site before cytokinesis completion

in *cdc55Δ* cells. Strains Y1434 and Y1435 were arrested in metaphase by Cdc20

depletion and released into anaphase by Cdc20 re-induction. Formaldehyde-fixed

cells were stained with 50 U/mL of rhodamine-phalloidin. (a) Representative images

of the actin staining and Myo1-GFP signals from *CDC55* and *cdc55Δ* at different cell

cycle stages (metaphase, anaphase, cytokinesis and the next S phase) are shown.

(b) Quantifications and representative images of the cells with re-polarized actin at

the new bud site are shown. Scale bar, 1 μm.

1096

Figure S3. Myo1 asymmetry contraction in G1-synchronized cells in absence

of Cdc55. Strains Y1631 (N=15) and Y1761 (N=12) were arrested in G1 by α-factor

addition and released into cell cycle progression by pheromone removal. Cells were

fixed with formaldehyde before taking images. Quantifications of the cells with

asymmetric Myo1 contraction are represented (left panel). Representative images of

Myo1 asymmetric contraction are shown (right panel). Scale bar, 1 μm.

1103

Figure S4. Proper cytoplasm separation during membrane abscission in the

absence of Cdc55. Strains Y1717 and Y1708 were arrested in metaphase by

Cdc20 depletion and released into anaphase by Cdc20 re-induction. Time-lapse

images were captured every 2 minutes. Membrane abscission was followed by

visualization of the GFP_3-Ras2 signal, and Myo1-tdTomato was used as a control for

cytokinesis progression. Representative images from *CDC55* (N=12) and *cdc55Δ*

(N=13) cells are shown (left panel). Images are taken from the best z-stack to

visualize the division site. Scale bar, 1 μm. Quantification of the cumulative

percentage of cells abscised is represented (right panel).

1113

Figure S5. PP2A-Cdc55 participates in the dephosphorylation of IPCs proteins

during cytokinesis. (a-d) Phosphorylation changes of Hof1, Inn1, Cyk3 and Iqg1 in

absence of Cdc55. Strains Y1314, Y1394, Y1639, Y1640, Y1437, Y1438, Y1491 and

Y1497 were arrested in metaphase and released into anaphase by Cdc20 depletion

and re-addition. Protein phosphorylations were analyzed by western blot. Pgk1 and

Myo1-Flag levels were used as a loading control. Mitosis progression was followed

by analyzing the anaphase spindle elongation by *in situ* immunofluorescence. At

least 100 cells were scored at each time-point. Quantifications of the western blots

were performed using Fiji Software and means and SEMs are represented.

Student's unpaired t-test analyses were carried out using the Prism5 program. (e)

Alkaline phosphatase assay for Chs2. Native protein extracts were prepared from

Y1318 cells arrested in metaphase and incubated with alkaline phosphatase and

alkaline phosphatase's inhibitor as indicated. PhosStop was used as the alkaline

phosphatase's inhibitor. Chs2 phosphorylation was analyzed by western blot in

Phos-tag gels. (f) Separase is not dephosphorylated by PP2A-Cdc55. Metaphase

arrested cells of the strain Y695 was used to purify the PP2A-Cdc55 complex by

TAP purification. Esp1-HA was purified from metaphase-arrested cells of strain

Y1748. Purified Esp1-HA was incubated with the PP2A-Cdc55 complex at the

indicated times and the separase phosphorylation levels were detected by western

blot using the anti-phospho Ser/Thr antibody. Representative images of one

phosphatase assay are shown. Protein levels were quantified using Fiji software.

Quantifications of the remaining separase phosphorylation signal normalized to the

amount of Cdc55 and Esp1 are shown. Means and SEMs of three phosphatase

assays are represented. Student's unpaired t-test analysis was carried out using the

Prism5 program.

1139

Figure S6. PP2A-Cdc55 is required for proper IPCs localization and contraction

at the division site. Strains Y1572, Y1606, Y1454, Y1608, Y1306, Y1578, Y1574,

Y1604, Y1576 and Y1575 were synchronized into anaphase by Cdc20 depletion and

re-addition, and time-lapse images were captured every 2 minutes. Myo1-tdTomato

was used as a control for cytokinesis progression. (a-e) Representative images of

the indicated GFP-tagged IPCs proteins and Myo1 contractions from *CDC55*

and *cdc55Δ* cells are shown. Iqg1 and Inn1 contraction at the bud neck are normal in

the absence of Cdc55. By contrast, Hof1 and Cyk3 contractions are longer in the

absence of Cdc55. Scale bar, 1 μm.

a)

CDC55

Myo1-tdTomato

-2   0   2   4   6   time (minutes)

cdc55Δ

Myo1-tdTomato

-2   0   2   4   6   time (minutes)

Contraction time

ns

● *CDC55*
■ *cdc55Δ*

time (minutes)

Myo1

b)

Asymmetric Myo1 dot

cells (%)

CDC55   *cdc55Δ*

c)

Asymmetric Myo1 dot

cells (%)

CDC55   *cdc55-AID*   *cdc55-ED*   *chs3Δ*   *cdc55Δ chs3Δ*

Myo1-tdTomato

CDC55   *cdc55-AID*   *cdc55-ED*   *chs3Δ*   *cdc55Δ chs3Δ*

d)

Calcofluor intensity

****

arbitrary units

CDC55 CHS3   *cdc55Δ CHS3*

e)

Calcofluor intensity

****

arbitrary units

CDC55 chs3Δ   *cdc55Δ chs3Δ*

a)

cdc55Δ hof1-aid
CDC55 hof1-aid
cdc55Δ HOF1
CDC55 HOF1 CYK3
cdc55Δ CYK3
CDC55 td-cyk3-aid
cdc55Δ td-cyk3-aid

YPD   1.5 mM IAA

b)

WCE   Pk IP

-Chs2-HA
-Pk-Cdc55

c)

CDC55   *cdc55Δ*

release (minutes)

ᵃ-P-Chs2-HA
-Chs2-HA

-Pgk1

18 35 63 89 19 5 3 7 11 6   21 25 33 40 50 21 17 6 4 5   anaphase spindles (% cells)

ratio phosphorylated/dephosphorylated Chs2

ns   ns   *

■CDC55
■cdc55Δ

metaphase   anaphase   cytokinesis -G1

arbitrary units

d)

PP2A-Cdc55 PPase assay

0 20 40   time (minutes)

-anti-phospho Ser/Thr Chs2
-Chs2-HA
-HA-Cdc55

Remaining Chs2 phosphorylation

**
ns

arbitrary units

0   20   40
time (minutes)

e)

PP2A-Cdc55 PPase assay

- HA-cdc55-ED HA-Cdc55

-^{32}P
-Chs2
-HA

Remaining Chs2 phosphorylation

ns

arbitrary units

cdc55-ED   Cdc55

a)

b)

a)

heavy (WT)/light(*cdc55Δ*) ratio 0,750

pRS site probability 99.9%

DNADLPVDPYHL**s**PQQQPSNNLFGSGR (S133)

b)

c)

Asymmetric Myo1 dot

d)

Hof1 Contraction

e)

Asymmetric Myo1 dot

f)

Hof1 Contraction