Diverse molecular mechanisms contribute to differential expression of human duplicated genes

Colin J. Shew[1,2], Paulina Carmona-Mora[1,3,4], Daniela C. Soto[1,2], Mira Mastoras[1], Elizabeth Roberts[1], Joseph Rosas[1,5], Dhriti Jagannathan[1], Gulhan Kaya[1], Henriette O'Geen[1], Megan Y. Dennis[1-6,†]

[1]Genome Center, [2]Integrative Genetics and Genomics Graduate Group, [3]MIND Institute, [4]Autism Research Training Program, [5]Postbaccalaureate Research Education Program, [6]Department of Biochemistry & Molecular Medicine, University of California, Davis, CA 95616

[†]Corresponding author:
Megan Y. Dennis, Ph.D.
University of California, Davis, School of Medicine
One Shields Avenue
Genome Center, 4303 GBSF
Davis, CA 95616
Email: mydennis@ucdavis.edu

ABSTRACT

Emerging evidence links genes within human-specific segmental duplications (HSDs) to traits

and diseases unique to our species. Strikingly, despite being nearly identical by sequence

(>98.5%), paralogous HSD genes are differentially expressed across human cell and tissue types,

though the underlying mechanisms have not been examined. Comparing cross-tissue mRNA

levels between humans and chimpanzees, we determined that broadly, HSD paralogs (75 genes

from 30 families) display expression patterns consistent with pseudo- or neofunctionalization. In

general, the ancestral paralog exhibited greatest expression conservation with the chimpanzee

ortholog, though exceptions suggest duplicate paralogs that may retain or supplant ancestral

functions. To understand mechanisms underlying this observed regulatory divergence, we

reanalyzed data from human lymphoblastoid cell lines (LCLs) (n=445), showing that ~75% of

derived HSD paralogs exhibit significant differential expression and a greater than two-fold

difference from their ancestral counterpart. To identify active *cis*-regulatory elements (CREs) in

HSDs, we reanalyzed ENCODE data to recover hundreds of candidate CREs in these regions.

Further, we generated ChIP-seq data for active chromatin features in an LCL using longer

Illumina reads to better distinguish peaks in paralogous regions. Some of these duplicated CREs

are sufficient to drive differential reporter activity, suggesting they may contribute to divergent

cis-regulation of paralogs. This work provides evidence that *cis*-regulatory divergence

contributes to novel expression patterns of recent gene duplicates in humans.

INTRODUCTION

Gene duplication occurs universally and is considered a major source of evolutionary novelty. For example, from a survey of representative eukaryotic genomes, the origin of over 30% of genes was attributed to duplication(Zhang 2003). Although most duplicated genes are fated to be rapidly pseudogenized, some may retain important functions and experience strong purifying selection(Lynch 2000). These genes are thought to become variously subfunctionalized (sharing the ancestral function) or neofunctionalized (gaining a novel function). Expression divergence is likely integral to this process, as spatiotemporal partitioning of function places both daughter paralogs under negative selection, helping them escape pseudogenization(Rodin and Riggs 2003; Rodin et al. 2005). It has been suggested that this may be the primary driver behind duplicate gene retention, as gene regulation can be altered relatively easily, without the need to acquire rare beneficial coding mutations(Ohno 1970). For example, mouse *Hoxa1* and *Hoxb1* genes are functionally redundant but partitioned by expression, with normal development possible from a single gene under control of regulatory elements from both paralogs(Tvrdik and Capecchi 2006). On a genome-wide scale, substantial expression divergence is observed in vertebrates following whole-genome duplications specific to teleost and salmonid fishes(Kassahn et al. 2009; Braasch et al. 2016; Lien et al. 2016; Varadharajan et al. 2018). Meta-analysis suggests that across all these species, selection on expression levels appears relaxed in one of the paralogs(Sandve et al. 2018). However, duplication of individual loci rather than whole genome events is substantially more common in vertebrates, with segmental duplications (SDs, regions defined as having >90% sequence similarity and being at least 1 kb in size(Bailey 2002)) having the potential to structurally rearrange the genome, in turn facilitating regulatory divergence and duplicate retention(Rodin et al. 2005). Although studies characterizing expression divergence of duplicated

genes in humans, mice, and yeast have identified broad patterns of dosage sharing among

daughter paralogs(Qian et al. 2010; Lan and Pritchard 2016), more recent duplications have not

been analyzed. In addition, no molecular explanations have been provided for the observed

expression changes.

Great apes have experienced a surge of SDs in the last ~10 million years, primarily interspersed

throughout the genome and potentially contributing to phenotypic differences observed between

these closely related species(Prado-Martinez et al. 2013). Human-specific SDs (HSDs), which

arose in the last ~6 million years following divergence from a common ancestor with

chimpanzee, contain genes unique to the human lineage that have compelling associations with

neurodevelopmental features(Charrier et al. 2012; Dennis et al. 2012; Florio et al. 2015; Fiddes

et al. 2018; Suzuki et al. 2018; Heide et al. 2020) and disorders(Dennis and Eichler 2016; Dennis

et al. 2017; Ishiura et al. 2019). Historically such young duplications have been poorly resolved

in genome assemblies due to their high sequence similarity. Recent sequencing efforts targeted to

HSDs have generated high-quality references for many of these loci(Steinberg et al. 2012;

Antonacci et al. 2014; O'Bleness et al. 2014; Dennis et al. 2017) resulting in the discovery of at

least 30 duplicated gene families accounting for >80 gene paralogs. Most derived HSD paralogs

encode putatively functional proteins and exhibit divergent expression patterns relative to

ancestral paralogs across numerous primary tissues, despite being nearly identical (>98%) by

sequence(Dennis et al. 2017). HSD genes also utilize novel promoters and exons exapted from

the site of insertion(Dougherty et al. 2017). Differential regulation may also be explained by

associations of species-specific active chromatin modifications at SD loci(Giannuzzi et al. 2014),

though due to historical reference errors and computational challenges in short-read mapping to

highly-similar sequences(Chung et al. 2011; Ebbert et al. 2019), regulatory features of duplicated

loci remain poorly characterized.

In this study, we characterized patterns of regulatory divergence observed for HSD genes

between human and chimpanzee cell lines and primary tissues by quantifying the cross-tissue

conservation of homologs, offering potential insight into the speed at which young duplicate

genes diverge in expression and allowing for inference of likely functional HSD paralogs. We

leveraged genomic and epigenomic data from hundreds of human lymphoblastoid cell lines

(LCLs) to identify differentially expressed (DE) ancestral-derived gene pairs and tested potential

molecular contributors to paralog-specific divergence in expression, including copy number

(CN) variation, post-transcriptional regulation, and *cis*-regulatory changes. Finally, we surveyed

the active chromatin "landscape" for HSDs by reanalyzing ENCODE chromatin

immunoprecipitation sequence (ChIP-seq) data, producing novel data with longer reads to

ameliorate data loss due to poor mapping in SDs, and functionally validating *cis*-regulatory

elements (CREs) via a reporter assay. Overall, our work demonstrates that *cis*-regulatory

divergence, among other mechanisms, drives differential expression following gene duplication

and that useful regulatory information can be rescued from existing datasets for duplicated loci.

RESULTS

Conservation of HSD gene expression following duplication

To assess the evolutionary trajectory of recent human duplicated genes, we quantified (Soneson

et al. 2015; Patro et al. 2017) expression of 75 HSD genes from 30 gene families for which high-

confidence sequences were available (Dennis et al. 2017) (Table S1). Each HSD gene family

corresponds to a single-copy chimpanzee ortholog; the human paralog syntentic with the

chimpanzee gene, if known, is considered to be the ancestral copy, while the human-specific

duplicates are derived (Figure 1A) We compared expression HSD paralogs and chimpanzee

orthologs using mRNA-sequencing (RNA-seq) data from three cell lines and four primary tissue

types (Khan et al. 2013; Pavlovic et al. 2018; Marchetto et al. 2019; Blake et al. 2020). We

generated a custom transcriptome with improved annotations of paralogous genes and quantified

expression with a lightweight mapping approach that shows greater accuracy than alignment-

based methods for paralogous genes (Soneson et al. 2015; Patro et al. 2017). To interpret the

evolutionary fate of these paralogous genes, we compared the individual or summed expression

of human paralogs within each gene family to the single chimpanzee ortholog. We found that

derived HSD paralogs tended to show lower expression than the chimpanzee ortholog, summed

family expression was mostly higher, and ancestral paralogs were less likely to be DE (9/21

expressed ancestral genes showed no differential expression across all cell/tissue types versus

6/37 of expressed derived genes) (Figure S1A, Table S2). This suggests that ancestral genes are

most likely to retain their functions as duplicate paralogs diverge or are pseudogenized. We next

considered expression correlation across the four tissue types and three cell lines as an indicator

of expression conservation between human paralogs and the chimpanzee ortholog. Our

expectation was that in the case of pseudogenization or neofunctionalization, a single paralog

would correlate best with chimpanzee ortholog, while subfunctionalized genes would show a

higher correlation of the total family expression with the chimpanzee ortholog. We identified the

most conserved paralog in each duplicate gene family (versus chimpanzee) and found that these

paralogs showed similar correlation to the overall sum expression of the gene family, while the

Figure 1. Expression patterns of HSD genes. (A) Illustration of genes residing within HSDs; the ancestral paralog (green) corresponds to the chimpanzee ortholog, while derived paralogs (orange) are human-specific. The ancestral and derived genes comprise a gene family. **(B)** Cross-tissue correlations of HSD genes or gene family summed expression versus expression of chimpanzee orthologs are shown in boxplots. For each gene family, the most conserved paralog ("highest"), all other paralogs ("lower"), and the sum of all paralogs ("sum") are plotted separately. Ancestral genes, if evolutionary status is known, are denoted as a filled circle. Asterisks denote significant differences ($p < 0.01$, Tukey post-hoc test of ANOVA) **(C)** Expression divergence of derived genes from families with at least one LCL-expressed paralog is plotted as the $\log_2$ ratio of median derived and ancestral TPM expression. Each point represents a different LCL from the Geuvadis consortium (total N=445). The gray bar indicates a two-fold expression difference. **(D)** Expression values of ancestral *DUSP22* (green) and derived *DUSP22B* (orange), stratified by CN. Points represent individual LCLs, and the number of individuals represented in each CN category is indicated over each boxplot.

remaining paralogs had significantly lower correlation with the sum ($p<0.01$, Tukey post-hoc test

of ANOVA) (Figure 1B). This pattern is consistent with maintenance of function of one paralog

and neo- or pseudofunctionalization of the others. Notably, ancestral genes were overrepresented

in the "highest" correlation category (14/22 of known status, $p<0.001$, hypergeometric test).

Nevertheless, the remaining eight paralogs showing strongest conservation of expression with

chimpanzee orthologs were derived, including *FRMPD2B*, *HIST2H2BA*, *OR2A42*, and *SERF1B*.

These genes represent interesting candidates of a derived paralog usurping the function of the

ancestral gene. Taken together, these findings suggest that HSD genes are variously

pseudogenized or neofunctionalized, with the ancestral paralog often, but not always,

maintaining its expression pattern across tissues.

Expression of human paralogs in lymphoblastoid cell lines (LCLs)

We next focused on LCLs to gain a more detailed understanding of HSD expression patterns

across hundreds of individuals with matched genomic data. We estimated transcript abundance

using RNA-seq data from 462 human LCLs (Lappalainen et al. 2013) (Table S3) and determined

that over half (43/75) of HSD paralogs are expressed (at least one transcript per million (TPM)),

with the most highly expressed genes including *ARHGAP11A, ROCK1,* the adjacent *GTF2I* and

NCF1 families, and the *DUSP22* family, whose human-specific paralog *DUSP22B* is missing

from the reference (GRCh38) (Dennis et al. 2017). Comparing expression profiles within gene

families, we found derived and ancestral paralogs globally showed divergent expression levels;

in families with at least one expressed gene, all 31 derived genes showed significant differences

from their ancestral counterpart. Twenty of these showed a difference in median TPM greater

than two-fold, and in most cases (25/31) the derived gene had lower expression (Figures 1C, S2,

and Table S4). We noted that some paralogs exhibited clusters of outlier values for

derived/ancestral expression ratios, which could not be reconciled as copy number (CN) or

population of origin differences (Figure S3). Altogether, these results indicate that HSD genes

show divergent expression patterns on LCLs.

Copy-number variation and HSD expression

While all the genes in this study were chosen for being fixed or nearly fixed in all modern human

populations (Dennis et al. 2017), SD loci are known to be subject to recurrent rearrangement and

consequently exhibit varying degrees of CN polymorphism. Anticipating that CN variation is

likely to alter gene expression levels, we sought to characterize the impact of CN on differential

expression of HSD genes. After performing paralog-specific CN genotyping (Shen and Kidd

2020) of a subset of individuals for which 1000 Genomes Illumina sequences were available

(N=445), we found gene expression was positively associated with CN in about half (28/55) of

genes in expressed families (Table S5), indicating that higher CN often but not always translates

to increased expression. Notably, derived genes tended to have higher CN (1.3-fold on average,

across all assayed genes and individuals) but lower expression overall. To account for the effect

of CN on differential expression between paralogs, we also regressed CN out of comparisons of

expressed ancestral-derived gene pair and found 23/25 derived paralogs were still DE with

respect to the ancestral (six were not tested due to paralog-specific effects of CN; Table S4). For

example, while expression of *DUSP22B* was significantly associated with CN, these effects were

insufficient to explain DE relative to *DUSP22* (Figure 1D). Thus, while CN differences alter the

mRNA levels of HSD paralogs, they do not provide an explanation for overall differential

expression of these genes.

Post-transcriptional regulation of HSD genes

In order to determine if paralogous expression differences are driven by post-transcriptional

regulation, we next considered whether HSD transcripts were being processed as nonfunctional

pseudogenes. In this scenario, paralogs might be equally transcribed but then subjected to

degradation via nonsense-mediated decay (NMD). To test this, we compared gene expression

using available RNA-seq data from human NMD-deficient LCLs (N=4) against controls (N=2)

(Nguyen et al. 2012) and found no HSD genes with significant differential expression. We also

used these RNA-seq data to compare the ratio of derived to ancestral expression between NMD-

deficient LCLs and controls and found no significant differences, though sample sizes were

likely limiting (Figure S4A). This result is largely recapitulated by paralog-specific RT-qPCR for

three DE HSD genes families (*ARHGAP11, DUSP22,* and *ROCK1*) in four LCLs treated with

NMD-inhibiting drug emetine. Ratios of *ROCK1P1/ROCK1* and *DUSP22B/DUSP22* expression

were unaltered by emetine treatment, while *ARHGAP11B*/*ARHGAP11A* expression ratio

increased closer to one, consistent with NMD affecting *ARHGAP11B*, though not completely

'rescuing' derived expression levels to equal that of the ancestral (Figure S4B). *ARHGAP11B* is

a 3′ truncation of *ARHGAP11A*, potentially explaining differences in transcript stability.

Altogether, these results suggest that while NMD may alter steady-state expression levels of

some HSD genes, it is not a primary driver of their differential expression.

We also examined HSD 3′ untranslated regions (UTRs) for recognition sites of miRNAs

expressed in LCLs (Lappalainen et al. 2013) (N=13 3′ UTRs of expressed gene families; mean

94 binding sites per UTR) using TargetScan (Agarwal et al. 2015). miRNA binding sites were

nearly identical between paralogs. The small numbers of miRNA-binding differences between

paralogs correlated with expression divergence (linear regression $p=0.01$; adjusted $R^2=0.38$;

Figure S5A), but in the opposite direction as we might predict (i.e., more differences leading to

more similar expression patterns between paralogs). When we compared paralogs with identical

and non-identical 3′ UTRs, we observed significantly lower derived expression ratios from those

that differ (Wilcoxon rank sum test p<0.01, Figure S5B). These data point to a potential role for

miRNAs in HSD transcriptional regulation, but this mechanism is unlikely to cause differential

expression within highly expressed gene families, such as the *DUSP22* and *NCF1*.

Role of *cis*-regulation in HSD differential expression

We next aimed to determine if *cis*-regulatory changes contribute to expression divergence of

HSDs. Because SDs often generate gene truncations and fusions with adjacent transcribed

sequences (Dougherty et al. 2017), we reasoned that gains or losses of promoters or UTRs would

likely cause large changes in gene expression. We compared relative expression by truncation

status (5′-, 3′-, or non-truncated) of all derived genes in expressed families to their ancestral

paralogs. Ancestral and derived genes had more similar expression levels in non-truncating

duplications, while truncated genes tended to be less expressed than their ancestral paralogs,

particularly 5' truncations compared to all other HSD genes ($p=0.06$, *t*-test; Figure S6A), in

concordance with previous findings(Dougherty et al. 2018). While we may have limited power

to detect differences with a small number of genes, these results hint that promoter activity may

contribute to differential expression patterns. Considering sequence-level changes more broadly,

however, we observed no relationship between expression divergence and pairwise nucleotide

divergence across entire duplicons or within promoters (Figure S6B–C).

Given that the vast majority of paralog-specific variants (PSVs) are unlikely to be functionally

important, we used publicly available chromatin immunoprecipitation sequencing (ChIP-seq)

datasets from the Encyclopedia of DNA Elements (ENCODE) project (Consortium and The

ENCODE Project Consortium 2012; Davis et al. 2018) to identify likely CREs (H3K4me3,

H3K4me1, H3K27ac, and RNA PolII) in a single LCL (GM12878). In each data set, we

observed a lower density of bases covered by peaks in SDs (>90% similarity) and HSDs (>98%

similarity) compared to the rest of the genome (empirical p=0.001, N=1000 replicates; Figure 2,

in yellow). We posit, as others have previously (Chung et al. 2011; McVicker et al. 2013;

Giannuzzi et al. 2014), that this discrepancy is an artifact of the high sequence similarity of SDs,

as reads originating from these regions map to multiple locations of the genome and are

accordingly discarded at the alignment stage or by mapping-quality filtering.

To recover this missing information, we implemented a pipeline that allowed reads to align to

multiple locations in the genome then, using CSEM (Chung et al. 2011), iteratively assigned

posterior probabilities to each read alignment based on the unique mapping rate of nearby reads.

Selecting the most likely alignment to allocate a read (i.e., mapping position with the highest

posterior probability), we improved peak discovery in SDs and HSDs for the aforementioned

chromatin features, erasing the depletion for all but H3K27ac, which was still substantially

improved (Figure 2, in blue). The peaks we discovered largely overlapped with the ENCODE

peaks, though RNA PolII had a large proportion of peaks unique to this analysis (Figure S7).

Figure 2. Depletion and recovery of ChIP peaks in SDs. Results are summarized for three ChIP-seq peak discovery approaches: publicly-available ENCODE peaks (yellow), peaks from multimapping and CSEM allocation of ENCODE raw data (blue), and peaks from multimapping and CSEM allocation of long ChIP data. For SDs (left) and SDs with greater than 98% sequence identity (or HSDs, right), the fraction of bases under peaks was calculated (solid vertical lines). The SD coordinates were permuted 1000 times within the human reference (GRCh38), and an expected distribution of the fraction of bases covered was generated. Empirical one-sided p-values for depletion are indicated above each plot.

Improved peak discovery using longer-read ChIP-seq

To improve our ability to align reads accurately to specific paralogs, we generated longer-read

(~500 bp insert size, 2x250 bp PE Illumina) ChIP-seq ("long ChIP") libraries (H3K4me3,

H3K27ac, H3K4me1, and RNA PolII) from the LCL GM12878. Longer reads mapped to SDs

with greater accuracy (Figure S8A), allowing for higher-confidence discovery of novel peaks in

duplicated regions using standard single-site mapping approaches (BWA mem, Figure S8B).

However, all marks except H3K4me1 were still depleted for peaks in SDs relative to the rest of

the genome. Subsequently, we analyzed the long ChIP data allowing for multiple alignments and

probabilistically assigning reads to one position (Bowtie and CSEM, Figure S8B). Long ChIP

showed increased posterior assignment probabilities with respect to the short-read ENCODE

data, and the depletion of peaks in SDs was erased for H3K4me3, H3K4me1, and PolII (Figure

2, in pink). Notably, for most libraries fewer overall peaks were identified with long ChIP versus

ENCODE data, though the peaks that do exist were largely replicated (on average, 73% of long

ChIP peaks were corresponded to an ENCODE peak (Chikina and Troyanskaya 2012); Table S6;

Figure S9). Long ChIP peaks tended to be larger, with 2.4–3.7 times as many bases per peak,

excepting H3K4me1, which had slightly smaller peaks. We verified greater enrichment of

H3K27ac at the ancestral *DUSP22* (1.1–2.9-fold difference) at three PSVs, also noting a

correlation of expression divergence and differential enrichment (Figure S10). These findings

suggest that reanalysis of ChIP-seq data can accurately identify enriched regions at HSD loci,

uncovering potentially divergent regulatory environments.

To identify putatively functional *cis*-regulatory regions within HSDs, we integrated our ChIP-seq

peaks in an 8-state chromHMM model (Ernst and Kellis 2012), providing a novel set of

candidate activating CREs, as virtually no information is available in the current ENCODE

release for SD loci. Because derived gene expression is generally lower than ancestral, we

compared putative CREs in 100-kb windows covering HSD loci (defined in Dennis, 2017). We

observed no significant differences in the fraction of bases covered by putative CREs (Wilcoxon

rank-sum test; Figure S11). We also observed no differences between ancestral and derived

regions in the individual ChIP-seq datasets (active histone marks or RNA PolII), considering

both the fraction of bases covered enrichment strength of peaks. We also failed to find a

difference in the fraction of bases covered by repressed H3K27me3 domains identified from our

multimapping analysis. Thus, explanations beyond the overall abundance of chromatin features

are needed, as important functional changes in CRE activity may not be reflected in global

differences. We did visually identify differences in presence or absence of putative *cis*-acting

elements suggesting a more nuanced approach is necessary in pinpointing mechanisms

contributing to paralog expression differences (Figure 3).

Impact of altered chromatin context due to structural variation on gene expression

HSDs are often transposed many thousands of kilobases from their ancestral paralogs, and in

some cases on entirely different chromosomes, resulting in new adjacent genomic regions that

could impact expression of genes contained within; as such, we sought to understand if *cis*-

regulatory elements outside of our duplicated regions might contribute to gene expression

divergence. To do this, we examined expression quantitative loci (eQTLs) associated with HSD

gene expression. Using our reanalyzed expression data and variant calls from the 1000 Genomes

Project (N=460) (Genomes Project, Consortium et al. 2015), we identified 42 HSD genes with

significant eQTLs (N=3,603 variants in 9,524 gene-variant pairs), representing a 1.5- to 3-fold

Figure 3. Histone H3K27ac landscape at the chromosome 7q11.23 HSD locus. The ancestral 7q11.23 HSD locus (top) and one of its derived loci (bottom) are aligned, and genes, PSVs, luciferase-tested regions (turquoise), and H3K27ac ChIP-seq signal (purple) are indicated. For each locus, publicly available ENCODE, reanalyzed ENCODE multimapping + CSEM allocation, and long ChIP multimapping + CSEM allocation data are shown.

increase from published work (Lappalainen et al. 2013; Wen et al. 2015). Sixty-four percent

(N=2,300) of associated variants were located within SDs, of which 14% (N=498) fell within

HSDs. Understanding that variants identified within SDs are likely to be unreliable

(Hartasánchez et al. 2018; Ebbert et al. 2019) we focused on eQTLs in SD-proximal regions and

found 577/1,303 eQTL variants with paralog-specific associations. For example, five SD-

proximal variants were associated with *ARHGAP11A* expression, while none were identified for

ARHGAP11B, which is located 2 Mb distal to its ancestral locus. Similarly, four eQTLs were

identified for *DUSP22* on chromosome 6, while 26 eQTLs were linked with its derived paralog

DUSP22B on chromosome 16. We also considered physical contacts generated by chromatin

looping of HSD promoters, as three-dimensional promoter-enhancer loops frequently indicate

regulatory interactions. Using loops identified in GM12878 from promoter capture Hi-C (Mifsud

et al. 2015) and H3K27ac HiChIP (Mumbach et al. 2017; Juric et al. 2019), we identified 354

and 26 regions, respectively, interacting with HSD promoters. Over 90% of these regions were

located outside of HSDs, and 169 HSD eQTLs overlapped 31 of these regions (total size ~160

kb). These findings highlight the potential for adjacent, unique sequences to drive divergent

regulation of HSDs genes.

Differential activity of *cis*-acting elements between duplicate paralogs

Using our combined datasets, we examined three HSD loci containing the gene families

expressed most highly in LCLs (*ARHGAP11, NCF1*, and *DUSP22*) to identify functional

changes in CREs that may contribute to paralogous expression divergence (Figures 3, S12–S14).

In all three cases, the ancestral paralog exhibited significantly greater expression compared to

derived paralog(s) (Figure 4A). To determine if sequence differences within CREs identified

from our chromHMM annotations are sufficient to drive differences in gene expression, we

performed luciferase-reporter assays in HeLa cells and LCLs on paralogous promoters and

enhancer candidates.

ARHGAP11A/ARHGAP11B

The promoter of *ARHGAP11B* exhibited greater activity compared to the chimpanzee ortholog

and ancestral paralog in both HeLa and LCLs (~4-fold difference in activity between HSD

paralogs, $p<5\times10^{-10}$ in both cell lines; Figures 4B, S15A, S16A). This is in contrast to mRNA

levels in LCLs, where the ancestral *ARHGAP11A* is more highly expressed. Positing that distal

elements may be driving differential expression between these paralogs, we identified putative

enhancers unique to each paralog; these comprised one downstream of the *ARHGAP11A*

duplicon and two downstream of *ARHGAP11B*. In HeLa cells, the *ARHGAP11A* element showed

weak repressive activity (0.3-fold difference, $p<2\times10^{-16}$), while the *ARHGAP11B* elements

showed modest activity over baseline (~2-fold difference, $p<2\times10^{-14}$ each), leaving the primary

driver of differential expression for these genes unknown (Figures 4C, S15B).

NCF1/NCF1B/NCF1C

Promoters of the ancestral *NCF1* and its derived paralogs *NCF1B* and *NCF1C* genes did not

exhibit significant differential activities in LCLs and modest difference in HeLa (0.8-fold

difference, $p<0.001$; Figures 4A, S15A, S16A). However, an enhancer element common to all

three paralogs showed the greatest activity for the ancestral *NCF1* paralog in both cell types.

This was concordant with differential mRNA levels (~3-fold difference over either derived in

LCL; $p<0.002$ for all comparisons) (Figures 4, S16B). Thus, this enhancer, if targeted to *NCF1*

and its paralogs, may be responsible for differences in their mRNA levels.

323
DUSP22/DUSP22B

DUSP22 (ancestral) and *DUSP22B* (derived) promoters showed differential activity concordant

with gene expression (i.e., the human ancestral paralog exhibited significantly greater activity

than both the human derived and chimpanzee ortholog) (~1.5-fold difference; $p<5\times10^{-13}$) in both

HeLa and LCLs (Figures 4B, S15A, S16A). We also tested six putative enhancers shared

between the two paralogs in HeLa cells only. Four were active elements, two showed differential

activity opposite to that of gene expression, and one element tracking with differential paralog

Figure 4. Functional characterization of putativef HSD CREs. Putative CREs (promoters and enhancers) from three HSD duplicate gene families (*ARHGAP11*, *NCF1*, and *DUSP22*) were tested in luciferase-reporter assays for function. **(A)** mRNA levels (TPM) for the three tested HSD gene families in human LCLs (N=446). **(B)** Representative luciferase reporter experiments for promoters of the paralogous HSD genes and orthologous chimpanzee sequences in HeLa cells. Significantly different activity ($p<0.05$, Tukey's post-hoc test of ANOVA) from the negative control is indicated along the top bar over each panel, and significant differences among homologous sequences are indicated between boxplots. The *p*-values for each comparison are available in Table S7. **(C)** Representative luciferase reporter experiments for candidate enhancers from the same gene families in HeLa cells, with significant activity over/under baseline indicated along the top bar, and significant differences between paralogous sequences between boxplots ($p<0.05$, Tukey's post-hoc test of ANOVA). **(D)** Cartoons indicating the relative locations of each candidate sequence within or adjacent to HSDs (thick, colored arrows), based on all experiments (Figure S15). Inactive sequences are shown with a small dot, enhancer sequences are shown with a diamond, and silencer sequences are shown with a square. Differentially active derived sequences (relative to ancestral) are marked with a plus or minus sign. Elements validated in LCLs are indicated with a check mark.

expression (Figure 4C). We subsequently validated the latter enhancer element in LCLs (~1.4-

fold difference; $p<2\times10^{-16}$) (Figure S16C). We conclude that the difference in promoter activity

is the primary driver of *DUSP22* and *DUSP22B* differential expression, though distal CREs also

likely play a role in modulating transcription.

DISCUSSION

In this work, we provide evidence that recently duplicated, human-specific genes exhibit

differential expression, at least in part due to divergent *cis*-acting regulation. Historically, these

regions have been poorly characterized genetically and epigenetically. Here, we performed an

improved analysis of gene expression and chromatin histone post-translational modification data

to maximize information retained in HSDs and gain an understanding of the regulatory

landscapes of these regions. Although previous work (Dennis et al. 2017; Dougherty et al. 2017)

has quantified HSD gene expression across many tissues, these analyses were limited to a single

canonical isoform from standard (RefSeq) annotations. This analysis is the first to consider all

annotated isoforms for each HSD gene and to improve or create annotations for derived HSD

genes. In addition, our implementation of CSEM (Chung et al. 2011) allowed ChIP-seq peaks to

be identified in HSD loci, an improvement upon widely-used ENCODE data.

We highlight three loci whose regulation is of potential relevance to human-specific traits.

Functional studies of *ARHGAP11B*, which encodes a Rho-GTPase activating protein, suggest

this gene has been neofunctionalized in relation to its ancestral paralog by its unique expression

in cortical progenitor neurons (basal radial glia) (Florio et al. 2015). Though the function(s) of

this gene in LCLs is uncertain, it causes increased cortical neurogenesis in a number of model

systems (Kalebic et al. 2018; Namba et al. 2020). Our characterization of CRE in LCLs may be

more directly related to immune phenotypes; it has been suggested that humans are more prone

to autoimmune disease than chimpanzees, particularly as a result of T and B cell response to

viral infection (Anon 2013; Anon 2017). The ancestral *NCF1*, which encodes Neutrophil

Cytosolic Factor 1, has been shown to regulate autoimmune response via T cell activation in

mice (Hultqvist et al. 2004)copy-number differences of the gene and its derived full-length

paralogs (*NCF1B* and *NCF1C*) are associated with reduced risk of systemic lupus erythematosus,

suggesting an additive functional effect (Zhao et al. 2017)(Hultqvist et al. 2004; Li et al. 2014)

Finally, *DUSP22*, which encodes a tyrosine phosphatase, is also a regulator of autoimmune

response, and *dusp22* knockout mice experience T cell proliferation, increased inflammation, and

autoimmune encephalomyelitis (Li et al. 2014). The full-length paralog *DUSP22B* is

uncharacterized, as it is missing from the human reference but located on chromosome 16p12.1

at variable CN (Figure 1D). Our characterization of promoters and distal regulatory elements of

these three gene families uncovered potential drivers of differential expression; the the ancestral

DUSP22 promoter and an enhancer element upstream of *NCF1* both show activity patterns

concordant with the mRNA levels of these gene families, for which the ancestral gene is most

highly expressed. Searching for potential *trans* effectors, we identified an ancestral *DUSP22*

deletion of four bases in a homopolymer repeat, with as many as 13 similar TFBSs found only in

the less active *DUSP22B* and chimpanzee *DUSP22* ortholog. Some of these belonged to

transcriptional repressors (ZNF394 and ZNF350). Examining predicted TFBSs within *NCF1*

promoters, which did not exhibit differential activity, we observed no gains or losses of any

TFBSs relative to chimpanzee. Puzzlingly, no predicted TFBSs were unique to the most active

NCF1 enhancer, but the paralogous *NCF1B* and *NCF1C* possessed many TFBSs that were

missing from the ancestral, at least one of which belonged to the transcriptional repressor

ZNF394. Overall, these results point to diverse mechanisms contributing to differential

expression between these three gene families and represent, to our knowledge, the first direct

comparison and verification of active CREs between recently duplicated gene paralogs in

humans.

In agreement with previous analysis of whole-genome duplications in teleost fishes (Sandve et

al. 2018), we find that the dominant scenario for HSD genes involves a single paralog under

stronger selection, resulting in maintenance of ancestral gene expression. This is largely

consistent with nonfunctionalization or neofunctionalization of the other duplicate paralog; some

gene families display patterns consistent with subfunctionalization (greater correlation of

summed expression with chimpanzee than any individual paralog), but none of these exceed a

difference in correlation of 0.05. Unlike previous work, these duplications arose much more

recently (a few versus tens to hundreds millions of years) (Braasch et al. 2016; Lien et al. 2016),

suggesting that these changes were rapidly acquired, though these analyses also rely on a much

smaller set of genes and fewer tissues. Additionally, our finding that the majority of HSD

paralogs show reduced expression is in line with a report that duplicate genes in mammals and

yeast are less expressed, probably as result of neutral evolution (Qian et al. 2010). In contrast, a

study by Lan and Pritchard (2016) found that mammalian tandem duplicated genes tend to be co-

regulated and subfunctionalized by dosage. Notably, many of the HSD genes in our study were

excluded from the previous analysis due to high sequence-identity. Further, most of our HSD

genes are interspersed many hundreds of kilobases from each other throughout the human

genome. They and others (Rodin and Riggs 2003) suggest that divergence of gene regulation is

more likely in the case of large genomic rearrangements, which can at birth place daughter

paralogs in a novel regulatory environments, including topological domains, heterochromatin, or

transcriptional hubs. We note that differential expression is still observed in even very recent (<1

million years) duplications (Dennis et al. 2017), such as the gene families *DUSP22*, *SERF1*,

SMN, *TCAF1*, and *TCAF2*. Thus, these differences were either rapid to evolve or existed at the

time of duplication.

Though this work represents an important step toward a more complete picture of the regulation

of HSD genes, there are still some technical limitations to overcome. We implemented a

mapping-free approach for transcript quantification, which circumvents multiple-mapping issues

that affect "traditional" RNA-seq analyses and is demonstrated to accurately distinguish highly

similar transcripts (Patro et al. 2017). However, reads originating from multiple transcripts are

assigned probabilistically, and, as such, some genes may appear artificially similar in expression.

For instance, *DUSP22B* expression was nonzero in individuals completely missing this paralog

(Figure 1C) due to the recent nature of this duplication (<1 million years), which has resulted in

very few PSVs differentiating the duplicates at the mRNA level. Conversely, RNA-seq

quantification of older gene families appeared entirely distinguishable between paralogs

(*HIST2H2BF*). In addition, the available contigs of complete HSD loci have only been generated

from one effectively haploid cell line (Dennis et al. 2017). As such, PSVs are not necessarily

fixed, and may also be shared between genes (Dumont 2015). Accordingly, a more advanced

understanding of genetic variation at HSD loci is required to separate similarly expressed genes

with maximum leverage. This would also allow for the identification of reliable eQTLs within

duplications. We also find, as expected, that longer read lengths allow greater bioinformatic

distinction between duplicated loci, which remains an inherent limitation of many existing data

sources, such as the relatively short reads (~30 nt) of ENCODE ChIP-seq. While longer reads are

fundamentally more informative, some discrepancies between the ENCODE and long ChIP

CSEM analyses could be a result of misallocation of reads. We were encouraged to find that a

standard single-mapping approach (bwa mem) identified novel peaks in HSD, albeit at a lower

rate than genome-wide. As such, longer ChIP read lengths can be used in future studies to further

describe the epigenetic state of duplicated loci with a broader array of epitopes and cell/tissue

types. Finally, chromatin conformation information will be necessary to physically link our

tested distal elements to their target promoters, as linear proximity within the genome is an

imperfect predictor. Like other genomic assays, these data are currently only sparsely available

in SDs.

Despite limitations, we show that HSD regions contain differentially regulated genes and have

built maps of potentially active chromatin in these regions. Using these data, we demonstrated

that recently duplicated regulatory sequences are often functional, as expected, and that small

sequence changes can significantly alter their activity. Our study did not pinpoint a universal

factor that gives rise to the observed differential expression between paralogs, and the underlying

molecular mechanisms are likely unique to each HSD gene. In the three duplicate gene families

tested, promoter activity is only sometimes concordant with overall gene expression, suggesting

that other types of regulatory elements, like enhancers and silencers, may cooperatively control

overall expression. Currently, the challenge is to determine which PSVs and non-duplicated

regions are functionally relevant. We have produced and leveraged a variety of analyses to

narrow down likely candidates by chromatin state, changes to TFBSs, expression modulation,

and physical proximity to promoters. However, the number of candidate regions is too great to

test via low-throughput methods such as luciferase reporters, and this problem is exacerbated by

the need to compare regulatory behavior across multiple cell types. We propose that massively-

parallel reporter assays could be implemented to validate and quantify CRE activity of thousands

of candidate paralogous sequences. Such data could determine to what extent HSD gene

expression is predicted by nearby CRE activity. Future studies could also integrate additional

types of data, such as targeted chromatin capture data (such as 4C or capture Hi-C), or nascent

transcription (GROseq, 5' CAGE). Finally, characterization of DNA methylation, which is

especially challenging in duplicated loci, will be vital to build a more complete picture of the

epigenetic landscape of HSD loci. This study represents a first step toward improving

quantification of gene expression and active chromatin states in recent duplications and provides

a foundation for future work characterizing regulatory and functional changes in recently

duplicated loci.

MATERIALS AND METHODS

Quantification of HSD gene expression

Due to poor annotation of many HSD paralogs, custom transcriptomes were generated to ensure

equivalent isoform models for paralogous genes, biasing against differential expression. First,

transcript sequences for ancestral genes were extracted from GENCODE and mapped to derived

human loci (contig from (Dennis et al. 2017) or GRCh38 for *SERF1B*) and a long-read

chimpanzee assembly (Kronenberg et al. 2018) using BLAT (Kent 2002). GENCODE v27

transcripts were used for human-chimpanzee comparisons, since the chimpanzee transcriptome

(Kronenberg et al. 2018) was built on this version; for human-only analyses, GENCODE v32

was used. Alignments were manually curated, and new derived transcripts were extracted from

contigs. These transcripts, in addition to HSD transcripts generated from whole-isoform

sequencing of brain tissue (Dougherty et al. 2018), were added to GENCODE (human) or the

chimpanzee (after aligning to the chimpanzee assembly) transcriptome. Expression

quantification was performed using Salmon v1.2.0 (Patro et al. 2017), the custom transcriptomes,

and reference genomes (GRCh38 or Kronenberg et al.) as decoy sequence. For paired-end data,

we used the flags "--validateMappings" and "--gcBias".. RNA-seq data were first lightly

trimmed prior to quantification using trim_galore (https://github.com/FelixKrueger/TrimGalore)

with the following flags: -q 20 --illumina --phred33 --length 20. Length-normalized TPM values

or counts per gene were obtained using the tximport package in R (Soneson et al. 2015).

Differential expression analysis

Human and chimpanzee RNA-seq data from four primary tissues (Blake et al. 2020), LCLs

(Khan et al. 2013; Blake et al. 2020), induced pluripotent stem cells (iPSCs) (Pavlovic et al.

2018), and iPSC-derived neural progenitor cells (Marchetto et al. 2019) were analyzed as

described above. Count data from chimpanzee genes were duplicated to allow for pairwise

comparison to each HSD duplicate, as well as the sum of all HSD genes in each family. Genes

expressed below the 75% percentile (corresponding to 1-2 counts per million reads) were filtered

from the analysis, leaving 16,752–18,225 genes. A linear model including species and sex was

fitted to each shared gene (N=55,461) using limma-voom (Law et al. 2014; Ritchie et al. 2015),

and differentially expressed genes were identified at a 5% false discovery rate (FDR) (Nguyen et

al. 2012).

Copy number-controlled differential expression analysis

Paralog-specific CN estimates were generated using QuicK-mer2 (Shen and Kidd 2020),whole-

genome sequence data from the 1000 Genomes Project (30X) (Fairley et al. 2020) and a custom

reference consisting of GRCh38 plus an additional contig representing the *DUSP22B* duplicon

(Dennis et al. 2017). Expression analysis was performed using RNA-seq data from LCLs

included in the Geuvadis study (Lappalainen et al. 2013) for which CN genotypes were

generated (N=445). Ancestral-derived gene pairs were compared with a linear model to identify

significant differences in $\log_2$-transformed TPM values after controlling for continuous CN

genotypes. Models were first fit with an interaction coefficient, and if no interaction was detected

($p > 0.05$), models were fit to expression and CN only. Resulting *p*-values were corrected via the

Benjamini-Hochberg procedure using the R package qvalue (http://github.com/jdstorey/qvalue)

and used to identify differential expression of ancestral-derived gene pairs at a 5% FDR. For

visualization purposes (Figure 1D), *DUSP22* CN genotypes were adjusted to known values for

GM12878 (as determined by fluorescence *in situ* hybridization in (Dennis et al. 2017)).

Cell Culture

Human LCLs were obtained from the Coriell Institute. The cells were grown in suspension in

RPMI 1640 medium (Genesee Scientific) supplemented with 15% fetal bovine serum, 100 U/mL

penicillin, and 100 µg/mL streptomycin and maintained at 37°C with 5% CO_2. To test the

impact of NMD inhibition, two million cells of each LCL (GM19204, GM18508, GM19193,

GM19238, GM12878, and S003659_Chimp1) were grown overnight and subsequently treated

with 100 µg/ml of emetine (Sigma) for seven hours (Noensie and Dietz 2001). Parallel cultures

were left untreated and grown at standard conditions. HeLa cells were grown in Dulbecco's

Modified Eagle Medium (DMEM), High Glucose, with L-Glutamine (Genesee Scientific)

supplemented with 10% fetal bovine serum (Gibco, Life Technologies), penicillin (100 U/mL)

and streptomycin (100 µg/mL) (Gibco, Life Technologies) at 37°C with 5% CO2.

RNA extraction and cDNA generation

LCLs were harvested and added to an appropriate volume of TRIzol® solution (Invitrogen™) (1

ml per 10^7 cells) and stored at -80°C for ~24 hr before extraction to ensure complete lysis of

cells. The next day, 200 µl of chloroform (Fisher Scientific) was added, and the homogenate was

shaken vigorously for 20 seconds and incubated at room temperature for 2–3 min. Samples were

spun at 10,000×g for 18 min at 4°C and the aqueous phase was transferred to a sterile RNase-

free tube. An equal volume of 100% RNAse-free ethanol was added, samples were mixed by

vortex, and then purified with an RNeasy Mini Kit (Qiagen). Samples were eluted in 30 µl

RNase-free water and stored at -80°C. Transcriptor High Fidelity cDNA Synthesis Kit (Roche)

was used for cDNA synthesis with OligodT primers. Following reverse transcription, samples

were treated with RNase A (Qiagen) at 37°C, and cDNAs were stored at -20°C.

Identification of miRNA binding sites

For ancestral paralogs of each HSD gene family, the 3′-UTR was extracted from canonical

transcript isoforms using the UCSC Genome Browser (GRCh38) and compared using blastn

(Altschul et al. 1990) against existing alignments of homologs previously generated for human,

chimpanzee, and rhesus (Dennis et al. 2017). Using TargetScan 7.0 and annotated miRNA

sequences and families (release 7.1; Sept 2016) (Agarwal et al. 2015), we identified miRNA

targets of individual human paralogs and non-human primate orthologs.

Correlation of expression skew and sequence divergence

Ancestral-derived paralog expression skew was calculated as the absolute value of

$\log_2$(derived/ancestral), using the median TPM values for each gene and a pseudocount 1×10^{-4}

(an order of magnitude below the smallest nonzero value). Sequence divergence as the pairwise

identity with the ancestral sequence was taken from (Dennis et al. 2017). Gene families were

included if at least one paralog was expressed at a level >1 TPM. For promoters, sequence

divergence was tabulated as the sum of all mismatches and alignment gaps within ±500 base

pairs of the transcription start site (Gencode v32). These quantities were correlated and the

strength of the relationship was determined with a linear regression.

ChIP assays

ChIP assays were carried out as previously described with minor modifications (O'Geen et al.

2019). GM12878 cells were cross-linked in growth media containing 1% formaldehyde (Fisher

Scientific BP531) for 10 min at room temperature and the reaction was stopped with 0.125 M

glycine. Cross-linked cells were washed twice in PBS and stored at -80°C. 2×10^6 cells were used

per ChIP assay. Cells from two biological replicates were lysed with ChIP lysis buffer (5 mM

PIPES pH8, 85 mM KCl, 1% Igepal) with a protease inhibitor (PI) cocktail (Roche). Nuclei were

collected by centrifugation at 2,000 rpm. for 5 min at 4°C and lysed in nuclei lysis buffer (50

mM Tris pH8, 10 mM EDTA, 1% SDS) supplemented with PI cocktail. Chromatin was

fragmented in microTUBEs with the E220 (Covaris) using the low cell shearing protocol (Duty

cycle 2%, PIP 105, CPB 200, 4 min) and diluted with 5 volumes of RIPA buffer (50 mM Tris pH

7.6, 150 mM NaCl, 1 mM EDTA pH8, 1% Igepal, 0.25% Deoxycholic acid). ChIP enrichment

was performed by incubation for 16 h at 4°C with the following antibodies: 2 μg H3K27ac

antibody (Active Motif #39133), 4 μg H3K4me1 antibody (Millipore 07-436), 2 μg H3K4me3

antibody (Active Motif #39915), or 2 μg RNA Polymerase II (PolII) antibody clone 8WG16

(Covance MMS-126R). RNA PolII samples were incubated for an additional hour with 2 μg

Rabbit Anti-Mouse IgG (MP Biomedical #55436). Immune complexes were bound to 20 µl

magnetic protein A/G beads (ThermoFisher) for 2 hours at 4°C. Beads were washed 2x with

RIPA, 3x with ChIP wash buffer (100 mM Tris pH8, 500 mM LiCl, 1% Deoxycholic acid) and

once with ChIP wash buffer plus 150 mM NaCl. ChIP samples were eluted in 100 µl ChIP

elution buffer (50 mM NaHCO$_3$, 1% SDS) and cross-linking reversed with addition of 0.5 M

NaCl and heating at 65°C overnight. Samples were treated with 2 μg RNaseA (Qiagen) and

DNA was purified using the QIAquick PCR Purification Kit (Qiagen). ChIP enrichments were

confirmed by qPCR with 2× SYBR FAST mastermix (KAPA Biosystems) using the CFX384

Real-Time System C1000 Touch Thermo Cycler (BioRad). ACTB primers served as positive

control and HER2 primers as negative controls (Table S8). ChIP enrichment was calculated

relative to input samples using the dC_t method ($dC_t = C_t$[HER2-ChIP]-C_t[input]). Each entire

ChIP sample was used to prepare Illumina sequencing libraries using the KAPA Hyper Prep Kit

(Roche). Adapter-ligated DNA was separated on a 2% E-Gel EX (Invitrogen) and the 500-800

bp fraction was excised and purified using the QIAquick gel extraction Kit (Qiagen). Indexed

primers were used to generate dual-indexed libraries and amplified libraries were size selected

(500–700 bp) using the PippenHT (Sage Science). Equimolar library amounts were pooled and

sequenced on the NovaSeq SP (Illumina).

Analysis of ChIP-seq data

ChIP-seq peaks obtained with the ENCODE pipeline were directly downloaded from the

ENCODE portal (Davis et al. 2018) for H3K4me3 (ENCFF228GWY), H3K27ac

(ENCFF367KIF), H3K4me1 (ENCFF453PEP), POLR2A (ENCFF455ZLJ), and H3K27me3

(ENCFF153VOQ). For "short" ChIP-seq peak calling using raw ENCODE data, GM12878

ChIP-seq reads were downloaded from the ENCODE portal for RNA Polymerase II

(ENCSR000AKA), H3K4me3 (ENCSR000BGD), H3K4me1 (ENCSR000AKF), H3K27ac

(ENCSR000AKC), and H3K27me3 (ENCFF000OBB). Illumina adapters and low quality bases

(Phred score < 20) were trimmed using Trimmomatic (Bolger et al. 2014) (parameters

SLIDINGWINDOW:4:20 MINLEN:20) and aligned to a custom reference genome (GRCh38

with an added *DUSP22B* contig) using single-end Bowtie (Langmead et al. 2009) configured to

allow multiple mappings per read (parameters a -v2 -m99). After mapping, PCR duplicates were

removed using Picard Markduplicates and secondary alignments were removed with samtools

v1.9. Multi-mapping reads were allocated to their most likely position using CSEM v2.4 (Chung

et al. 2011). CSEM was run using the --no-extending-reads option and the fragment size was

calculated with phantompeakqualtools run_SPP.R script (Landt et al. 2012). A custom script was

developed to select the alignment with the highest posterior probability as assigned by CSEM for

each multimapping read, choosing one alignment randomly in case of a tie. Peaks were called

using MACS2 callpeak (v2.2.6) on default settings using MACS2's shifting model (Zhang et al.

2008) (https://github.com/macs3-project/MACS). Broad peaks were called at a FDR of 5%,

while narrow peaks were called at a FDR of 1%. BigWig files for peak's visualization were

obtained with MACS2 bdgcmp tool and UCSC bedGraphToBigWig. For H3K27me3, which

occurs in large repressive domains, enriched regions were identified with hiddenDomains

(Starmer and Magnuson 2016). Paired-end long-ChIP reads were generated as described above

and Illumina adapters were removed using Trimmomatic (parameters SLIDINGWINDOW:4:30

MINLEN:50). Reads were mapped using both paired-end BWA mem and single-end Bowtie

allowing for multiple mappings (parameters -a -n -S -e 200 -m 99). For single-end alignments,

forward and reverse reads were concatenated into a single file and properly renamed to secure

unique reads IDs. Reads aligned with BWA were filtered by MAPQ $\geq$ 20 while reads with

multiple mappings aligned with Bowtie were allocated with CSEM and most likely alignments

were selected with the custom script. Duplicates and secondary alignments were removed as

explained above. Peaks were called using MACS2 with identical parameters used for short-reads,

adding the BAMPE option in the case of paired-end reads aligned with BWA mem.

For depletion analyses, SD coordinates were directly downloaded from UCSC Table Browser

and HSD coordinates were obtained by filtering alignments with sequence identity over 98% in

the fracMatch column, converting them to BED format and merging overlapping entries using

bedtools merge. The number of peaks and bases under peaks on each region of interest were

obtained with bedtools intersect. To obtain depletion statistics, 1000 regions of the same size as

SD and HSD were randomly sampled from the human genome GRCh38. Empirical p-values of

depletion tests were calculated as p-value = (M+1)/(N+1), where M is the number of iterations

less than the observed value and N is the number of iterations.

Additionally, mapping quality scores (MAPQ) distributions for H3K27ac following a similar

approach as explained before, but using BWA aln and BWA mem for short and long ChIP-seq

reads respectively, after PCR duplicates and secondary alignments removal. Posterior

probabilities distributions for H3K27ac were examined using the output of CSEM after selecting

the most likely alignment with the custom script. Entries in unique space were subsampled to 10

million and plots were obtained with the geom_density function in ggplot R package.

All ChIP-seq analyses are available as a TrackHub for the UCSC Genome Browser (TBD).

Further, the bioinformatic pipeline is freely available for use in Snakemake format

(Supplementary File 1, github TBD), allowing the analysis to be replicated in any cell or tissue

type of interest.

Paralog-specific validation of RNA expression and ChIP data

Following published protocols (Integrated DNA Technologies), we used the rhAMP assay in 10

µl total reaction volumes to quantify abundance of PSVs (for all assays except *ARHGAP11*

expression, the fluorophores FAM=A paralog and VIC=B paralog) as a proxy for paralog-

specific expression (RNA) and enrichment (ChIP) (Table S7). We used 10 ng total of RNA

converted to cDNA to validate gene expression for duplicated gene families *ARHGAP11*,

ROCK1, and *DUSP22*. We calculated dCt of cDNA and gDNA as Ct_{FAM}-Ct_{VIC} and ddCt as

dCt_{cDNA}-dCt_{gDNA} from the same cell line. We calculated dCt of the input and ChIP-enriched

library as Ct_{FAM}-Ct_{VIC} and ddCt as dCt_{ChIP}-dCt_{input} from the same cell line. For both expression

and ChIP analyses, the ratio of abundance of the B to the A paralog is 2^{ddCt}.

chromHMM annotations

We generated models separately for ENCODE short-read data and long ChIP, after

multimapping and allocation, using active chromatin histone modifications (H3K4me3,

H3K4me1, and H3K27ac). States corresponding to active transcription start sites and active

enhancers were identified manually. These annotations are publicly available at UCSC Genome

Browser TrackHub (TBD)).

Luciferase reporter assays

Promoters of highly and differentially expressed HSD gene families (*ARHGAP11*, *NCF1*, and

DUSP22) were chosen for screening in a reporter assay. Fragments containing the TSS and

spanning ~1 kb were amplified with KpnI and SacI restriction sites and cloned into the luciferase

reporter vector pGL3-basic (Promega). Candidate enhancers within 50 kb of genes bodies were

selected based on the presence of ChromHMM CREs in the re-analyzed data from human LCLs.

Target regions were a maximum size of 5 kb, and peaks larger than this were tiled with multiple

targets. Gateway homology arms were added to primers in accordance with the manual

(ThermoFisher), and PCR products were cloned into the entry vector pDONR221 (ThermoFisher

12536017). Expression clones for luciferase assays were generated by cloning pDONR221

inserts into the luciferase reporter pE1B (Antonellis et al. 2008) with the Gateway system.

Constructs were co-transfected (ThermoFisher Lipofectamine 3000) in equimolar amounts with

50 ng of the control plasmid pRL-TK (Renilla luciferase) into HeLa cells in 96-well plates. Cells

were at 70-90% confluence at the time of transfection. Luciferase assays were performed with

the Dual-Luciferase Reporter Assay System (Promega E1910). 48 hours post-transfection, cells

were washed with PBS, and lysed with Passive Lysis Buffer for at least 15 min sharking at 500

rpm. Lysates were stored at -80C. For LCLs, cells were split 48 and 24 hours pre-transfection to

ensure active division. Cells were counted, washed in PBS, and resuspended such that each

transfection contained 12.5×10^6 cells, 6.25 ug of test construct, and equimolar pRL-TK in RPMI.

Cells were electroporated using the Neon Transfection System in accordance with previously

published work (Tewhey et al. 2018) and recovered at a density of 3×10^6 cells/mL in pre-

warmed RPMI including 15% FBS without antibiotics. Transfection efficiencies of ~15% were

achieved. To perform luciferase assays, ~$5\times x10^5$ cells were pipetted into each well of a 96-well

plate, washed with PBS, and lysed with Passive Lysis Buffer as described for HeLa.

Luminescence measurements were performed according to the manufacturer's instructions using

a Tecan Infinite or Tecan Spark plate reader with injectors.

Transcription factors binding motifs

Alignments of cloned sequences were scanned for HOmo sapiens COmprehensive MOdel

COllection (HOCOMOCO) v11 (Kulakovskiy et al. 2018) TFBS motifs using FIMO (Grant et

al. 2011). HOCOMOCO motifs were limited to transcription factors expressed above 1 TPM in

>75% of ENCODE mRNA-seq libraries generated for GM12878 (ENCSR077AZT,

ENCLB555AQG, ENCLB555AQH, ENCLB555ANP, ENCLB555ALI, ENCLB555ANM,

ENCLB555ANN, ENCLB037ZZZ, ENCLB038ZZZ, ENCLB043ZZZ, ENCLB044ZZZ,

ENCLB041ZZZ, ENCLB042ZZZ, ENCLB045ZZZ, ENCLB046ZZZ, ENCLB700LMU,

ENCLB150CGC). Significant matches above a 5% FDR were retained for the analysis. TFBSs

were compared across homologous sequences to identify putative paralog-specific gains and

losses of binding sites.

FIGURE LEGENDS

Figure 1. Expression patterns of HSD genes. (A) Illustration of genes residing within HSDs;

the ancestral paralog (green) corresponds to the chimpanzee ortholog, while derived paralogs

(orange) are human-specific. The ancestral and derived genes comprise a gene family. **(B)** Cross-

tissue correlations of HSD genes or gene family summed expression versus expression of

chimpanzee orthologs are shown in boxplots. For each gene family, the most conserved paralog

("highest"), all other paralogs ("lower"), and the sum of all paralogs ("sum") are plotted

separately. Ancestral genes, if evolutionary status is known, are denoted as a filled circle.

Asterisks denote significant differences ($p<0.01$, Tukey post-hoc test of ANOVA) **(C)**

Expression divergence of derived genes from families with at least one LCL-expressed paralog is

plotted as the $\log_2$ ratio of median derived and ancestral TPM expression. Each point represents a

different LCL from the Geuvadis consortium (total N=445). The gray bar indicates a two-fold

expression difference. **(D)** Expression values of ancestral *DUSP22* (green) and derived

DUSP22B (orange), stratified by CN. Points represent individual LCLs, and the number of

individuals represented in each CN category is indicated over each boxplot.

Figure 2. Depletion and recovery of ChIP peaks in SDs. Results are summarized for three

ChIP-seq peak discovery approaches: publicly-available ENCODE peaks (yellow), peaks from

multimapping and CSEM allocation of ENCODE raw data (blue), and peaks from multimapping

and CSEM allocation of long ChIP data. For SDs (left) and SDs with greater than 98% sequence

identity (or HSDs, right), the fraction of bases under peaks was calculated (solid vertical lines).

The SD coordinates were permuted 1000 times within the human reference (GRCh38), and an

expected distribution of the fraction of bases covered was generated. Empirical one-sided p-

values for depletion are indicated above each plot.

Figure 3. Histone H3K27ac landscape at the chromosome 7q11.23 HSD locus. The ancestral

7q11.23 HSD locus (top) and one of its derived loci (bottom) are aligned, and genes, PSVs,

luciferase-tested regions (turquoise), and H3K27ac ChIP-seq signal (purple) are indicated. For

each locus, publicly available ENCODE, reanalyzed ENCODE multimapping + CSEM

allocation, and long ChIP multimapping + CSEM allocation data are shown.

Figure 4. Functional characterization of putativef HSD CREs. Putative CREs (promoters and

enhancers) from three HSD duplicate gene families (*ARHGAP11*, *NCF1*, and *DUSP22*) were

tested in luciferase-reporter assays for function. **(A)** mRNA levels (TPM) for the three tested

HSD gene families in human LCLs (N=446). **(B)** Representative luciferase reporter experiments

for promoters of the paralogous HSD genes and orthologous chimpanzee sequences in HeLa

cells. Significantly different activity ($p<0.05$, Tukey's post-hoc test of ANOVA) from the

negative control is indicated along the top bar over each panel, and significant differences among

homologous sequences are indicated between boxplots. The *p*-values for each comparison are

available in Table S7. **(C)** Representative luciferase reporter experiments for candidate enhancers

from the same gene families in HeLa cells, with significant activity over/under baseline indicated

along the top bar, and significant differences between paralogous sequences between boxplots

($p<0.05$, Tukey's post-hoc test of ANOVA). **(D)** Cartoons indicating the relative locations of

each candidate sequence within or adjacent to HSDs (thick, colored arrows), based on all

experiments (Figure S15). Inactive sequences are shown with a small dot, enhancer sequences

are shown with a diamond, and silencer sequences are shown with a square. Differentially active

derived sequences (relative to ancestral) are marked with a plus or minus sign. Elements

validated in LCLs are indicated with a check mark.

ACKNOWLEDGMENTS

We thank the many groups/consortia that have made their data publicly available, including Dr.

Yoav Gilad, the 1000 Genomes Project, GTEX, and ENCODE for which this research would not

be possible without its use. We also thank Dr. Colin Kern for valuable advice concerning ChIP-

seq analysis, Dr. Anthony Antonellis for sharing the pE1B enhancer reporter Gateway plasmid,

and Drs. Gerald Quon and Siobhan Brady for constructive feedback on the manuscript. This

work was supported by the National Institute of Neurological Disorders and Stroke at the

National Institutes of Health (NIH) (R00NS083627 to M.Y.D.). Statistical analysis advice was

provided by Dr. Blythe Durbin-Johnson through the MIND Institute Intellectual and

Developmental Disability Research Center, funded by the National Institute of Child Health and

Human Development (U54 HD079125). Additionally, M.Y.D. is supported as a Sloan fellow

(FG-2016-6814), P.C.M as an NIH National Institute of Mental Health T32 UC Davis Autism

Research Training Program fellow (5T32MH073124-17), D.C.S. as a Fulbright fellow, and J.R.

as an NIH National Institute of General Medical Sciences UC Davis Postbaccalaureate Research

Education Program fellow (R25GM116690).

DATA ACCESS

Long-insert ChIP-sequencing data generated for this study are available from the European

Nucleotide Archive under the accession PRJEB40356.

DISCLOSURE DECLARATION

Authors have no disclosures to declare.

REFERENCES

Agarwal V, Bell GW, Nam J-W, Bartel DP. 2015. Predicting effective microRNA target sites in
mammalian mRNAs. *eLife* [Internet] 4. Available from:
http://dx.doi.org/10.7554/elife.05005

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool.
J. Mol. Biol. 215:403–410.

Anon. 2013. Potential role of human-specific genes, human-specific microRNAs and human-
specific non-coding regulatory RNAs in the pathogenesis of Systemic Sclerosis and
Sjögren's Syndrome. *Autoimmun. Rev.* 12:1046–1051.

Anon. 2017. Are humans prone to autoimmunity? Implications from evolutionary changes in
hominin sialic acid biology. *J. Autoimmun.* 83:134–142.

Antonacci F, Dennis MY, Huddleston J, Sudmant PH, Steinberg KM, Rosenfeld JA, Miroballo
M, Graves TA, Vives L, Malig M, et al. 2014. Palindromic GOLGA8 core duplicons
promote chromosome 15q13.3 microdeletion and evolutionary instability. *Nature Genetics*
[Internet] 46:1293–1302. Available from: http://dx.doi.org/10.1038/ng.3120

Antonellis A, Huynh JL, Lee-Lin S-Q, Vinton RM, Renaud G, Loftus SK, Elliot G, Wolfsberg
TG, Green ED, McCallion AS, et al. 2008. Identification of neural crest and glial enhancers
at the mouse Sox10 locus through transgenesis in zebrafish. *PLoS Genet.* 4:e1000174.

Bailey JA. 2002. Recent Segmental Duplications in the Human Genome. *Science* [Internet]
297:1003–1007. Available from: http://dx.doi.org/10.1126/science.1072047

Blake LE, Roux J, Hernando-Herraez I, Banovich NE, Perez RG, Hsiao CJ, Eres I, Cuevas C,
Marques-Bonet T, Gilad Y. 2020. A comparison of gene expression and DNA methylation
patterns across tissues and species. *Genome Res.* 30:250–262.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence
data. *Bioinformatics* 30:2114–2120.

Braasch I, Gehrke AR, Smith JJ, Kawasaki K, Manousaki T, Pasquier J, Amores A, Desvignes
T, Batzel P, Catchen J, et al. 2016. The spotted gar genome illuminates vertebrate evolution
and facilitates human-teleost comparisons. *Nature Genetics* [Internet] 48:427–437.
Available from: http://dx.doi.org/10.1038/ng.3526

Charrier C, Joshi K, Coutinho-Budd J, Kim J-E, Lambert N, de Marchena J, Jin W-L,
Vanderhaeghen P, Ghosh A, Sassa T, et al. 2012. Inhibition of SRGAP2 Function by Its
Human-Specific Paralogs Induces Neoteny during Spine Maturation. *Cell* [Internet]
149:923–935. Available from: http://dx.doi.org/10.1016/j.cell.2012.03.034

Chikina MD, Troyanskaya OG. 2012. An effective statistical evaluation of ChIPseq dataset
similarity. *Bioinformatics* 28:607–613.

Chung D, Kuan PF, Li B, Sanalkumar R, Liang K, Bresnick EH, Dewey C, Keleş S. 2011.
Discovering transcription factor binding sites in highly repetitive regions of genomes with
multi-read analysis of ChIP-Seq data. *PLoS Comput. Biol.* 7:e1002111.

Consortium TEP, The ENCODE Project Consortium. 2012. An integrated encyclopedia of DNA
elements in the human genome. *Nature* [Internet] 489:57–74. Available from:
http://dx.doi.org/10.1038/nature11247

Davis CA, Hitz BC, Sloan CA, Chan ET, Davidson JM, Gabdank I, Hilton JA, Jain K,
Baymuradov UK, Narayanan AK, et al. 2018. The Encyclopedia of DNA elements
(ENCODE): data portal update. *Nucleic Acids Research* [Internet] 46:D794–D801.
Available from: http://dx.doi.org/10.1093/nar/gkx1081

Dennis MY, Eichler EE. 2016. Human adaptation and evolution by segmental duplication.
Current Opinion in Genetics & Development [Internet] 41:44–52. Available from:
http://dx.doi.org/10.1016/j.gde.2016.08.001

Dennis MY, Harshman L, Nelson BJ, Penn O, Cantsilieris S, Huddleston J, Antonacci F,
Penewit K, Denman L, Raja A, et al. 2017. The evolution and population diversity of
human-specific segmental duplications. *Nat Ecol Evol* 1:69.

Dennis MY, Nuttle X, Sudmant PH, Antonacci F, Graves TA, Nefedov M, Rosenfeld JA,
Sajjadian S, Malig M, Kotkiewicz H, et al. 2012. Evolution of Human-Specific Neural
SRGAP2 Genes by Incomplete Segmental Duplication. *Cell* [Internet] 149:912–922.
Available from: http://dx.doi.org/10.1016/j.cell.2012.03.033

Dougherty ML, Nuttle X, Penn O, Nelson BJ, Huddleston J, Baker C, Harshman L, Duyzend
MH, Ventura M, Antonacci F, et al. 2017. The birth of a human-specific neural gene by
incomplete duplication and gene fusion. *Genome Biol.* 18:49.

Dougherty ML, Underwood JG, Nelson BJ, Tseng E, Munson KM, Penn O, Nowakowski TJ,
Pollen AA, Eichler EE. 2018. Transcriptional fates of human-specific segmental
duplications in brain. *Genome Research* [Internet] 28:1566–1576. Available from:
http://dx.doi.org/10.1101/gr.237610.118

Dumont BL. 2015. Interlocus gene conversion explains at least 2.7 % of single nucleotide
variants in human segmental duplications. *BMC Genomics* [Internet] 16. Available from:
http://dx.doi.org/10.1186/s12864-015-1681-3

Ebbert MTW, Jensen TD, Jansen-West K, Sens JP, Reddy JS, Ridge PG, Kauwe JSK, Belzil V,
Pregent L, Carrasquillo MM, et al. 2019. Systematic analysis of dark and camouflaged
genes reveals disease-relevant genes hiding in plain sight. *Genome Biol.* 20:97.

Ernst J, Kellis M. 2012. ChromHMM: automating chromatin-state discovery and
characterization. *Nature Methods* [Internet] 9:215–216. Available from:
http://dx.doi.org/10.1038/nmeth.1906

Fairley S, Lowy-Gallego E, Perry E, Flicek P. 2020. The International Genome Sample Resource

(IGSR) collection of open human genomic variation resources. *Nucleic Acids Res.*
48:D941–D947.

Fiddes IT, Lodewijk GA, Mooring M, Bosworth CM, Ewing AD, Mantalas GL, Novak AM, van
den Bout A, Bishara A, Rosenkrantz JL, et al. 2018. Human-Specific NOTCH2NL Genes
Affect Notch Signaling and Cortical Neurogenesis. *Cell* [Internet] 173:1356–1369.e22.
Available from: http://dx.doi.org/10.1016/j.cell.2018.03.051

Florio M, Albert M, Taverna E, Namba T, Brandl H, Lewitus E, Haffner C, Sykes A, Wong FK,
Peters J, et al. 2015. Human-specific gene ARHGAP11B promotes basal progenitor
amplification and neocortex expansion. *Science* [Internet] 347:1465–1470. Available from:
http://dx.doi.org/10.1126/science.aaa1975

Genomes Project, Consortium, Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM,
Korbel JO, Marchini JL, McCarthy S, McVean GA, et al. 2015. A global reference for
human genetic variation. *Nature* 526:68–74.

Giannuzzi G, Migliavacca E, Reymond A. 2014. Novel H3K4me3 marks are enriched at human-
and chimpanzee-specific cytogenetic structures. *Genome Research* [Internet] 24:1455–
1468. Available from: http://dx.doi.org/10.1101/gr.167742.113

Grant CE, Bailey TL, Noble WS. 2011. FIMO: scanning for occurrences of a given motif.
Bioinformatics [Internet] 27:1017–1018. Available from:
http://dx.doi.org/10.1093/bioinformatics/btr064

Hartasánchez DA, Brasó-Vives M, Heredia-Genestar JM, Pybus M, Navarro A. 2018. Effect of
Collapsed Duplications on Diversity Estimates: What to Expect. *Genome Biol. Evol.*
10:2899–2905.

Heide M, Haffner C, Murayama A, Kurotaki Y, Shinohara H, Okano H, Sasaki E, Huttner WB.
2020. Human-specific ARHGAP11B increases size and folding of primate neocortex in the
fetal marmoset. *Science* [Internet]:eabb2401. Available from:
http://dx.doi.org/10.1126/science.abb2401

Hultqvist M, Olofsson P, Holmberg J, Backstrom BT, Tordsson J, Holmdahl R. 2004. Enhanced
autoimmunity, arthritis, and encephalomyelitis in mice with a reduced oxidative burst due to
a mutation in the Ncf1 gene. *Proceedings of the National Academy of Sciences* [Internet]
101:12646–12651. Available from: http://dx.doi.org/10.1073/pnas.0403831101

Ishiura H, Shibata S, Yoshimura J, Suzuki Y, Qu W, Doi K, Asem Almansour M, Kikuchi JK,
Taira M, Mitsui J, et al. 2019. Noncoding CGG repeat expansions in neuronal intranuclear
inclusion disease, oculopharyngodistal myopathy and an overlapping disease. *Nature*
Genetics [Internet] 51:1222–1232. Available from: http://dx.doi.org/10.1038/s41588-019-
0458-z

Juric I, Yu M, Abnousi A, Raviram R, Fang R, Zhao Y, Zhang Y, Qiu Y, Yang Y, Li Y, et al.
2019. MAPS: Model-based analysis of long-range chromatin interactions from PLAC-seq
and HiChIP experiments. *PLoS Comput. Biol.* 15:e1006982.

Kalebic N, Gilardi C, Albert M, Namba T, Long KR, Kostic M, Langen B, Huttner WB. 2018.
Human-specific ARHGAP11B induces hallmarks of neocortical expansion in developing
ferret neocortex. *eLife* [Internet] 7. Available from: http://dx.doi.org/10.7554/elife.41241

Kassahn KS, Dang VT, Wilkins SJ, Perkins AC, Ragan MA. 2009. Evolution of gene function
and regulatory control after whole-genome duplication: comparative analyses in vertebrates.
Genome Res. 19:1404–1418.

Kent WJ. 2002. BLAT--the BLAST-like alignment tool. *Genome Res.* 12:656–664.

Khan Z, Ford MJ, Cusanovich DA, Mitrano A, Pritchard JK, Gilad Y. 2013. Primate transcript
and protein expression levels evolve under compensatory selection pressures. *Science*
342:1100–1104.

Kronenberg ZN, Fiddes IT, Gordon D, Murali S, Cantsilieris S, Meyerson OS, Underwood JG,
Nelson BJ, Chaisson MJP, Dougherty ML, et al. 2018. High-resolution comparative
analysis of great ape genomes. *Science* [Internet] 360. Available from:
http://dx.doi.org/10.1126/science.aar6343

Kulakovskiy IV, Vorontsov IE, Yevshin IS, Sharipov RN, Fedorova AD, Rumynskiy EI,
Medvedeva YA, Magana-Mora A, Bajic VB, Papatsenko DA, et al. 2018. HOCOMOCO:
towards a complete collection of transcription factor binding models for human and mouse
via large-scale ChIP-Seq analysis. *Nucleic Acids Res.* 46:D252–D259.

Landt SG, Marinov GK, Kundaje A, Kheradpour P, Pauli F, Batzoglou S, Bernstein BE, Bickel
P, Brown JB, Cayting P, et al. 2012. ChIP-seq guidelines and practices of the ENCODE and
modENCODE consortia. *Genome Res.* 22:1813–1831.

Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient alignment
of short DNA sequences to the human genome. *Genome Biol.* 10:R25.

Lan X, Pritchard JK. 2016. Coregulation of tandem duplicate genes slows evolution of
subfunctionalization in mammals. *Science* [Internet] 352:1009–1013. Available from:
http://dx.doi.org/10.1126/science.aad8411

Lappalainen T, The Geuvadis Consortium, Sammeth M, Friedländer MR, 't PA, Monlong J,
Rivas MA, Gonzàlez-Porta M, Kurbatova N, Griebel T, et al. 2013. Transcriptome and
genome sequencing uncovers functional variation in humans. *Nature* [Internet] 501:506–
511. Available from: http://dx.doi.org/10.1038/nature12531

Law CW, Chen Y, Shi W, Smyth GK. 2014. voom: precision weights unlock linear model
analysis tools for RNA-seq read counts. *Genome Biology* [Internet] 15:R29. Available from:
http://dx.doi.org/10.1186/gb-2014-15-2-r29

Lien S, Koop BF, Sandve SR, Miller JR, Kent MP, Nome T, Hvidsten TR, Leong JS, Minkley
DR, Zimin A, et al. 2016. The Atlantic salmon genome provides insights into
rediploidization. *Nature* [Internet] 533:200–205. Available from:
http://dx.doi.org/10.1038/nature17164

Li J-P, Yang C-Y, Chuang H-C, Lan J-L, Chen D-Y, Chen Y-M, Wang X, Chen AJ, Belmont
JW, Tan T-H. 2014. The phosphatase JKAP/DUSP22 inhibits T-cell receptor signalling and
autoimmunity by inactivating Lck. *Nature Communications* [Internet] 5. Available from:
http://dx.doi.org/10.1038/ncomms4618

Lynch M. 2000. The Evolutionary Fate and Consequences of Duplicate Genes. *Science* [Internet]
290:1151–1155. Available from: http://dx.doi.org/10.1126/science.290.5494.1151

Marchetto MC, Hrvoj-Mihic B, Kerman BE, Yu DX, Vadodaria KC, Linker SB, Narvaiza I,
Santos R, Denli AM, Mendes APD, et al. 2019. Species-specific maturation profiles of
human, chimpanzee and bonobo neural cells. *eLife* [Internet] 8. Available from:
http://dx.doi.org/10.7554/elife.37527

McVicker G, van de Geijn B, Degner JF, Cain CE, Banovich NE, Raj A, Lewellen N, Myrthil
M, Gilad Y, Pritchard JK. 2013. Identification of Genetic Variants That Affect Histone
Modifications in Human Cells. *Science* [Internet] 342:747–749. Available from:
http://dx.doi.org/10.1126/science.1242429

Mifsud B, Tavares-Cadete F, Young AN, Sugar R, Schoenfelder S, Ferreira L, Wingett SW,
Andrews S, Grey W, Ewels PA, et al. 2015. Mapping long-range promoter contacts in
human cells with high-resolution capture Hi-C. *Nat. Genet.* 47:598–606.

Mumbach MR, Satpathy AT, Boyle EA, Dai C, Gowen BG, Cho SW, Nguyen ML, Rubin AJ,
Granja JM, Kazane KR, et al. 2017. Enhancer connectome in primary human cells identifies
target genes of disease-associated DNA elements. *Nat. Genet.* 49:1602–1612.

Namba T, Dóczi J, Pinson A, Xing L, Kalebic N, Wilsch-Bräuninger M, Long KR, Vaid S,
Lauer J, Bogdanova A, et al. 2020. Human-Specific ARHGAP11B Acts in Mitochondria to
Expand Neocortical Progenitors by Glutaminolysis. *Neuron* 105:867–881.e9.

Nguyen LS, Jolly L, Shoubridge C, Chan WK, Huang L, Laumonnier F, Raynaud M, Hackett A,
Field M, Rodriguez J, et al. 2012. Transcriptome profiling of UPF3B/NMD-deficient
lymphoblastoid cells from patients with various forms of intellectual disability. *Molecular
Psychiatry* [Internet] 17:1103–1115. Available from:
http://dx.doi.org/10.1038/mp.2011.163

Noensie EN, Dietz HC. 2001. A strategy for disease gene identification through nonsense-
mediated mRNA decay inhibition. *Nat. Biotechnol.* 19:434–439.

O'Bleness M, Searles VB, Dickens CM, Astling D, Albracht D, Mak AC, Lai YY, Lin C, Chu C,
Graves T, et al. 2014. Finished sequence and assembly of the DUF1220-rich 1q21 region
using a haploid human genome. *BMC Genomics* 15:387.

O'Geen H, Bates SL, Carter SS, Nisson KA, Halmai J, Fink KD, Rhie SK, Farnham PJ, Segal
DJ. 2019. Ezh2-dCas9 and KRAB-dCas9 enable engineering of epigenetic memory in a
context-dependent manner. *Epigenetics Chromatin* 12:26.

Ohno S. 1970. Evolution by Gene Duplication. Available from: http://dx.doi.org/10.1007/978-3-

642-86659-3

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. 2017. Salmon provides fast and bias-
aware quantification of transcript expression. *Nature Methods* [Internet] 14:417–419.
Available from: http://dx.doi.org/10.1038/nmeth.4197

Pavlovic BJ, Blake LE, Roux J, Chavarria C, Gilad Y. 2018. A Comparative Assessment of
Human and Chimpanzee iPSC-derived Cardiomyocytes with Primary Heart Tissues. *Sci.*
Rep. 8:15312.

Prado-Martinez J, Sudmant PH, Kidd JM, Li H, Kelley JL, Lorente-Galdos B, Veeramah KR,
Woerner AE, O'Connor TD, Santpere G, et al. 2013. Great ape genetic diversity and
population history. *Nature* 499:471–475.

Qian W, Liao B-Y, Chang AY-F, Zhang J. 2010. Maintenance of duplicate genes and their
functional redundancy by reduced expression. *Trends in Genetics* [Internet] 26:425–430.
Available from: http://dx.doi.org/10.1016/j.tig.2010.07.002

Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015. limma powers
differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids*
Res. 43:e47.

Rodin SN, Parkhomchuk DV, Riggs AD. 2005. Epigenetic changes and repositioning determine
the evolutionary fate of duplicated genes. *Biochemistry* 70:559–567.

Rodin SN, Riggs AD. 2003. Epigenetic silencing may aid evolution by gene duplication. *J. Mol.*
Evol. 56:718–729.

Sandve SR, Rohlfs RV, Hvidsten TR. 2018. Subfunctionalization versus neofunctionalization
after whole-genome duplication. *Nature Genetics* [Internet] 50:908–909. Available from:
http://dx.doi.org/10.1038/s41588-018-0162-4

Shen F, Kidd JM. 2020. Rapid, Paralog-Sensitive CNV Analysis of 2457 Human Genomes
Using QuicK-mer2. *Genes* [Internet] 11. Available from:
http://dx.doi.org/10.3390/genes11020141

Soneson C, Love MI, Robinson MD. 2015. Differential analyses for RNA-seq: transcript-level
estimates improve gene-level inferences. *F1000Res.* 4:1521.

Starmer J, Magnuson T. 2016. Detecting broad domains and narrow peaks in ChIP-seq data with
hiddenDomains. *BMC Bioinformatics* 17:144.

Steinberg KM, Antonacci F, Sudmant PH, Kidd JM, Campbell CD, Vives L, Malig M,
Scheinfeldt L, Beggs W, Ibrahim M, et al. 2012. Structural diversity and African origin of
the 17q21.31 inversion polymorphism. *Nature Genetics* [Internet] 44:872–880. Available
from: http://dx.doi.org/10.1038/ng.2335

Suzuki IK, Gacquer D, Van Heurck R, Kumar D, Wojno M, Bilheu A, Herpoel A, Lambert N,

Cheron J, Polleux F, et al. 2018. Human-Specific NOTCH2NL Genes Expand Cortical
988   Neurogenesis through Delta/Notch Regulation. *Cell* [Internet] 173:1370–1384.e16.
989   Available from: http://dx.doi.org/10.1016/j.cell.2018.03.067

990   Tewhey R, Kotliar D, Park DS, Liu B, Winnicki S, Reilly SK, Andersen KG, Mikkelsen TS,
991   Lander ES, Schaffner SF, et al. 2018. Direct Identification of Hundreds of Expression-
992   Modulating Variants using a Multiplexed Reporter Assay. *Cell* 172:1132–1134.

993   Tvrdik P, Capecchi MR. 2006. Reversal of Hox1 gene subfunctionalization in the mouse. *Dev.*
994   *Cell* 11:239–250.

995   Varadharajan S, Sandve SR, Gillard GB, Tørresen OK, Mulugeta TD, Hvidsten TR, Lien S,
996   Vøllestad LA, Jentoft S, Nederbragt AJ, et al. 2018. The Grayling Genome Reveals
997   Selection on Gene Expression Regulation after Whole-Genome Duplication. *Genome*
998   *Biology and Evolution* [Internet] 10:2785–2800. Available from:
999   http://dx.doi.org/10.1093/gbe/evy201

Wen X, Luca F, Pique-Regi R. 2015. Cross-Population Joint Analysis of eQTLs: Fine Mapping
and Functional Annotation. *PLOS Genetics* [Internet] 11:e1005176. Available from:
http://dx.doi.org/10.1371/journal.pgen.1005176

Zhang J. 2003. Evolution by gene duplication: an update. *Trends in Ecology & Evolution*
[Internet] 18:292–298. Available from: http://dx.doi.org/10.1016/s0169-5347(03)00033-8

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM,
Brown M, Li W, et al. 2008. Model-based analysis of ChIP-Seq (MACS). *Genome Biol.*
9:R137.