A fish-specific new gene evolves new functions in cichlid fishes

Langyu Gu[1,2,3,]*, Chenzheng Li[2], Xiaobing Mao[2], Zongfang Wei[2], Youkui Huang[2], Ximin He[1], Wenjun Zhou[1], Li Li[2],* Deshou Wang[2]*

[1] State Key Laboratory of Biocontrol, School of Life Sciences and School of Ecology, Sun Yat-sen University, Guangzhou, Guangdong, China.

[2] Key Laboratory of Freshwater Fish Reproduction and Development (Ministry of Education), Key Laboratory of Aquatic Science of Chongqing, School of Life Sciences, Southwest University, Chongqing, China

[3] Zoological Institute, Department of Environmental Sciences, University of Basel, Basel, Switzerland

*Correspondence:

Langyu Gu: langyugu@gmail.com

Li Li: alisir@swu.edu.cn

Deshou Wang: wdeshou@swu.edu.cn

Abstract

Genes whose dysfunction do not affect normal survival is a common phenomenon. Are they only meaningless residues during evolution? Here, we identified a fish-specific new gene, which we named *lg*. Gene knockout resulted in no obvious phenotype in zebrafish, but *lg* evolves an amino acid mutation that is under positive selection in the modern haplochromine (MH) cichlid fish lineage, the well-known adaptive radiative lineage. Moreover, the cichlid fish-specific upstream region of *lg* can drive new eGFP expression profiles in tissues related to adaptation. Besides, this homologous region from different cichlid fishes drive different eGFP expressions which is simply due to three MH-segregated SNPs mutations that are predicted to bind a hormone related transcription factor. We thus revealed an initially redundant gene evolving new functions in an adaptive radiative lineage. This further illuminates the mechanism about the emergence of new gene function with respect to evo-devo in a broad way.

Key words

lg, new gene, lineage-specific, evo-devo, fish

From the evolutionary point of view, the most adaptive organism is the
one that spends the least resources but produce the most outputs. However, it
seems that this rule does not apply to the genome organization, which is
occupied by a large proportion of genes whose dysfunction does not affect
normal survival. Does it make sense?

Nothing in biology makes sense except in the light of evolution [1]. As raw
materials for natural selection, the large portion of redundant genes provides
good opportunities for functional innovation. This is especially true for lineage-
specific genes, which can even occupy up to one-third of the genome in many
organisms [2]. Studies gradually revealed that many of them can evolve new
functions related to novelties and environmental adaptation in the particular
lineage. For example, in early brain development in human [3,4], in the creation of
novel tissues in the sea anemone [5], and in the development of a novel fan trait
on the leg in a water spider genus to explore new environment [6]. There is even a
whole lineage-specific gene family in water flea related to environmental
stimuli [7]. Lineage-specific gene thus provides a new perspective of new gene
function emergence.

It is well accepted that different organisms share deep homologous
developmental molecular bases (molecular toolkit), such as highly conserved
signalling pathways and transcription factors [8–10]. Then, how do new genes gain
new functions, and in what way are they integrated in the existing networks?
Although usually restricted due to pleiotropic effects [11], protein coding region
evolution can take place [12–14]. Compare to this, regulatory evolution is a well-
known strategy, since changes at the expression level are more tolerable and can
offer immediate phenotypic consequences [11,15]. Even a single mutation in the
upstream region can induce new expression profiles [16,17]. The relaxed selection

pressure induced by the changing expression profiles can even further prompt
the accumulation of mutations at the protein level.

In this study, we identified a fish-specific new gene, which we named *lg*.
Comparative -omics analyses and functional experiments revealed that multiple
losses of *lg* occurred in different fish lineages, and no obvious phenotype was
detected in gene knockout zebrafish, which make us assume that the ancestral
function of *lg* is not indispensable. *lg* mainly expressed in the brain and eyes in
most fishes, but evolved new expression patterns in cichlid fish lineage,
especially in the most-species richness lineage, the modern haplochromine
(MH) lineage, such as in pharyngeal cartilages and epidermal serotonin-positive
neuroepithelia cells. Further comparative genomic analyses across East African
cichlid fish phylogeny revealed that *lg* evolved a cichlid fish-specific upstream
region possessing three MH lineage-segregated SNPs and one MH lineage-
segregated amino acid mutation which is under positive selection. Transgenic
experiments confirm the new regulatory functions of this upstream region to
drive eGFP expression in zebrafish. Noticeably, the homologous sequence from
different cichlid fishes drive different eGFP expression patterns, which can be
simply due to the three MH lineage-segregated SNPs mutations that are
predicted to bind to a hormone related transcription factor. Together, our work
revealed an initially redundant fish-specific new gene evolves new regulatory
functions in an adaptive radiative cichlid fish lineage for the first time.

Results

lg **is a fish-specific gene**

To see *lg* sequence evolution, we conducted tblastx search in all the
species with available genomes in Ensembl 100. Results showed that *lg* only
appears in fishes (Fig. 1A, Table S1). To further characterize its evolution

within teleost fishes, we conducted tblastx search and gene predictions in draft genomes assembly across the whole phylogeny of teleost fishes [18]. Results showed that *lg* has been lost multiple times in different lineages (Fig. 1B), which inspired the assumption that the ancestral function of *lg* is not indispensable for fish survival.

**

lg **evolves new expression profiles in cichlid fishes, especially in the MH lineage**

To see the tissues where *lg* expressed, we investigated its expression profiles based on the combined results of available RNAseq data retrieved from NCBI and *in situ* hybridization in different fishes (Fig. 2, Table S2). *lg* mainly expressed in the brain and eyes in most fishes. Interestingly, it evolved new expression profiles in cichlid fishes in pharyngeal cartilages, as well as in the skin in the most species-richness lineage of cichlid fishes, the MH lineage [19]. These results suggest that the ancestral function of *lg* can be related to neurons, but new functions can evolve in other tissues in cichlid fishes, especially in the MH lineage.

**

No obvious phenotype was detected in *lg* **gene knockout (KO) zebrafish**

To further investigate the function of *lg*, KO was conducted using CRISPR/Cas9 in zebrafish. We successfully obtained F2 homozygous mutant (MUT) zebrafish and its wild-type siblings (WT). Sanger sequencing and Western-blot analyses confirmed the successful gene dysfunction with frameshift (5 bp deletion) and protein interruption (stop codon appeared at the 18[th] amino acid) (Fig. 3A). To see the affected gene expressions after KO, comparative transcriptomic analyses were conducted between MUT and WT at 2 dpf (days post fertilization) and 4 dpf zebrafish, respectively. The results showed that most differential expressed (DE) genes are related to immune, neurons and energy (Table S3). To see whether there are different immune

responses between MUT and WT, we examined the immune responses of larvae

zebrafish to lipopolysaccharide (LPS) by counting the numbers of lyz^+

neutrophils in the caudal hematopoietic tissue at one day after injection [20]. No

significant difference was detected (Fig. 3B). To see whether KO affect neurons

development in the brain and eyes where lg expressed in zebrafish, we

compared the phenotypic changes between MUT and WT using the brain

neuron marker, HuC [21]. Still no difference was detected (Fig. 3C). Then we paid

our attention to one top DE gene, $slc16a6$, which has been reported to be related

to energy metabolism under fasting condition [22]. With the same strategy,

however, we still could not find obvious metabolism phenotypic changes

between MUT and WT under fast condition using ORO staining (Fig. 3C). This

can be explained by the fact that $slc16a6$ is only down-regulated here instead of

entirely disrupted as in the [22], so that it can still execute its funciton. Above all,

mutation of lg resulted in no obvious phenotypes in zebrafish. These results

further support our assumption that the ancestral function of lg is not

indispensable.

lg evolves a cichlid fish-specific upstream region possessing MH-segregated

SNPs and one MH-segregated amino acid mutation that is under positive

selection

164   Noticeably, as we mentioned above, lg evolved new gene expression

profiles in cichlid fishes (Fig. 2), the classical model species for studying

adaptive radiation, which explosively radiated into hundreds to thousands of

species within a short time with very closely related genomes [23]. To characterize

its sequence evolution within cichlid fishes, we amplified the whole lg gene

including its upstream region in a total of 20 cichlid fishes representing main

lineages across East African cichlid fish phylogeny [24] (Fig. 4A). Interestingly,

both upstream region and the coding region exhibited segregation patterns with

the MH lineage, the most species-richness lineage of cichlid fishes [25] (Fig. 4A).

For example, there is one nonsynonymous amino acid mutation (A to S) located
at the 38[th] residue specific to the MH lineage which is under positive selection
(Figure 4A, Table S4). Furthermore, *lg* evolved an upstream region specific to
cichlid fishes (Fig. S1). Within this region, there are three MH lineage-
segregated SNPs mutations (Fig. 4A). Transcription factors (TFs) binding
prediction revealed that these three SNPs can set up the binding opportunities to
a hormone related TF, the glucocorticoid modulatory element binding protein 2
(GMEB2) (Fig. 4A). These results stimulate us to pay special attention to the *lg*
of cichlid fishes in the next steps.
182

183 The cichlid fish-specific upstream region indeed has regulatory functions
184 and performs different roles in different cichlid fish lineages

To see whether the cichlid fish-specific upstream region of *lg* indeed has
regulatory functions, and whether this region of MH lineage has different
regulatory roles compared to other cichlid fish lineages, we constructed two
transgenic lines with the homologous upstream region from different species as
the promoters to drive eGFP expression in zebrafish (Fig. 4A, Fig. S2). One line
is with the upstream region of a MH cichlid fish, *Pundamilia nyererei* (pT2AL-
Pn-eGFP transgenic line, Pn line). The other line is with the corresponding
homologous region of a cichlid fish belonging to a sister lineage of East African
Great lakes cichlid fishes, Nile tilapia (*Oreochromis niloticus*) [23] (pT2AL-On-
eGFP transgenic line, On line). We successfully obtained F2 generations for
both transgenic lines. Both lines can drive eGFP expressions in zebrafish,
confirming their regulatory functions, but with different patterns (Fig. 4B and
Fig. 4C). Pn line showed strong expressions in neurons in most regions of the
brain (olfactory epithelium, olfactory bulb, forebrain, optic tectum, cerebellum
and hindbrain), eyes, spinal cord, and heart valves. Besides, it also showed new
strong eGFP expressions in the hatching gland in embryos, pharyngeal arches,
chondrocytes cells in the fin and scattered epidermal serotonin-positive cells.
The positive eGFP cells were further confirmed by available antibodies (Fig.
4B). Interestingly, compared to the Pn line, eGFP fluroscence is pretty weak in

the brain and pharyngeal arches in the On line ($p<0.01$), and is hard to detect in
epidermal serotonin-positive cells and fin chondrocytes (Fig. 4C). Considering
that we obtained different F1 individuals of each line possess the same stable
eGFP expression patterns, and the eGFP expression patterns of different lines
are in consistent with the *in situ* hybridization and RNAseq results above, the
transgenic lines we constructed are stable and credible. These results indicate
that the cichlid fish-specific upstream region of *lg* indeed evolved new
regulatory functions, i.e. not only execute the ancestral expression profiles in
neurons, but also evolve new expressions in the hatching gland and pharyngeal
arches in cichlid fishes. Strong and new regulatory function in the scattered
epidermal serotonin-positive cells further evolved in the MH cichlid fish
lineage.

Different regulatory roles of the cichlid fish-specific upstream regions are
simply due to three MH lineage-segregated SNPs

To see whether the strong and new expression profiles in the Pn
transgenic line are caused by the three MH lineage segregated SNPs in the
upstream region, we constructed another transgenic line with the same upstream
region as in the Pn line but deleted these three SNPs (pT2AL-Pn-noSNPs-eGFP
line, Pn-noSNPs line) (Fig. 4C, Fig. S2). Interestingly, only with these three
SNPs deletion, Pn-noSNPs line showed clearly significant different expression
patterns with the Pn line, but similar patterns with the On line (Fig. 4C, Fig. 4D
and Table S5). These results confirmed that these SNPs are the key reason for
the different expression patterns between Pn line and On line.

Discussion

Above all, we unravelled the evolution of a fish-specific new gene, *lg*, for
the first time. No obvious phenotypic changes was detected in the *lg* KO
zebrafish. This can be due to the compensation by other genes in the
correponding gene network. This robust nature is not unexpected for immune

and brain network system [26,27], as the DE genes of the *lg* KO line are mainly
neurons and immunity related genes. Besides, lineage-specific new genes
initially usually located at the edge of the gene network [28], its disruption thus do
not affect the core gene network much. No matter what it is, mutation of *lg* does
not affect zebrafish survival. Together with its multiple losses in different fish
lineages during evolution, we propose that the ancestral function of *lg* is not
indispensable for fish survival.

Interestingly, *lg* evolved new regulatory functions in cichlid fishes,
especially in its most species-richness lineage, the MH lineage. For example, in
the hatching gland which is a key innovation to regulate the hatching period of
embryos to become free-living larvaes [29]; in pharyngeal cartilages, the well-
known tissues related to feeding and ecological niche exploration, are key to
cichlid fish adaptive radiation [30–32]; in scattered epidermal serotonin-positive
cells which are neuroepithelia cells related to environmental stimuli, such as
water flow, pheromone and hormone [33,34]; in the olfactory epithelium which is
well-known for detection of odorants, such as amino acid, hormone and
pheromone [35]. Considering that some of these tissues are derived from a key
vertebrate innovation, the neural crest cells [36], *lg* can be also causally linked to
neural crest cell development and differentiation. Together, *lg* gained new
regulatory functions in the most species-richness lineage, the MH lineage,
which can be causally linked to its adaptive radiation. Noticeably, although we
focused mainly on the regulatory function evolution of *lg* in cichlid fishes here,
the newly important function can be also evolved in their coding region.
Molecular evolutionary analyses have already shown that there is one amino
acid mutation segregated with the MH cichlid fish lineage and is under positive
selection, which might be causally linked to the adaptive radiation. Further KO
of *lg* in MH cichlid fishes will help test this hypothesis and unravel the
corresponding involved pathways. Besides, the evolution of *lg* gene function
can be also still on the way in cichlid fishes, considering that they are
evolutionarily young species [37].

Actually, the originally thought junk DNA in the genome, such as

transposable elements, non-coding regions and non-functional genes keep

giving us surprises. How can a new gene function be evovled and integrated

into the existing gene network? Here, we propose one hypothesis of the

functional evolution of *lg* that can be applied in a broad way (Fig. 5): the

relaxed selection pressure on the lineage-specific gene in the ancesral lineage

free it to accumulate mutations in both the regulatory region and the coding

region. The evolution of the regulatory region in *lg* offers immediate effects, i.e.

only with three SNPs mutations in the upstream region can provide new TFs

binding opportunities to build up new gene network connections to affect gene

expression profiles in the most species-richness MH lineage of cichlid fishes. It

will be interesting to test our predictions of the binding of a glucocorticoid

hormone related TF with these SNPs in the future. If this is true, it can connect

the organism with external environments efficiently, since glucocorticoid is a

well-known stress related hormone to regulate animal behavior to response to

the environment [38,39]. The *lg* gene we investigated here thus provides a good

example of new gene function evolution in an initially redundant lineage-

specific gene with respect to evo-devo in a broad way.

Methods

sampling

Samples of cichlid fishes for comparative genomic analyses were

provided by Prof. Walter Salzburger in University of Basel, Switzerland; cichlid

fish *Pundamila nyererei* for transgene plasmids construction was obtained from

IGV (Interessengemeinschaft Viktoriasee-Cichliden); Strains of AB (for

transgenic lines construction) and ABTü (for KO) zebrafish (*Danio rerio*) were

raised under standard condition (28.5°C, 14/10 h light/dark). Tilapia

(*Oreochromis niloticus*) were raised at 26°C, 14/10 h light/dark. Animal

experiments have been approved by the cantonal veterinary office in Basel
under permit number 2317 in University of Basel, Switzerland; the Committee
of Laboratory Animal Experimentation at Sun Yat-sen University and
Southwest University, China.
300

-omics analyses to characterize the evolution of lg across fish phylogeny

To characterize the sequence evolution of *lg*, tblastx was conducted with
-*e* 0.001 in fishes with good genome assembly in Ensembl 100 (Table S1) and
local blast v2.9.0 (tblastx) was conducted with draft genomes assembly across
the phylogeny according to the reference [18]. After getting hits, the scaffold
possessing the corresponding sequence was extracted to predict the coding
regions with Augustus 3.3.3 [40,41]. The predicted *lg* coding sequences were
further confirmed by aligning the translated amino acids with the available
sequences from Ensembl 100 using Geneious 8.0.5 [42].

.

-omics analyses to characterize the evolution of lg within cichlid fishes

To further characterize the sequence evolution of *lg* within East African
cichlid fishes, we amplified the whole gene sequence and its upstream region
(ca.10 kb in total) in different cichlid fishes. Combined with available cichlid
fish genomes, we thus studied the sequence evolution in a total of 20 cichlid
fishes representing main lineages across the phylogeny [24]. Genomic DNA was
extracted using DNeasy Blood&Tissue kit (Qiagen). PCR primers were
designed using Primer premier 5 (all the primers in this study were designed
with this software) [43]. High-Fidelity PCR Master Mix (BioLabs) and a touch-
down annealing process were used for the amplification. PGM Ion Torrent was
used to sequence long amplicons (400 bp) with CHIP 316 (Life Technologies).
Galaxy v20.09 [44] was used to trim low quality reads (window size 2, step size 1,
minimum quality score 20) and filter short reads with minimum size 40 bp. The
sequence of a cichlid fish, tilapia, retrieved from Ensembl database was used as

the reference to recheck the assembled sequences. The sequencing reads of *lg*

gene across cichlid fish phylogeny are available at NCBI (BioProject ID

PRJNA611690). The ion torrent sequencing of most species were successful

except *Limnochromis abeelei*, *Ophthalmotilapia ventralis*, and *Bathybates*

graueri, whose sequences were further obtained with Sanger sequencing

(Primers can be found in Table S6).

Genomic alignment of *lg* of cichlid fishes and other teleost fishes was

conducted with mVISTA [45,46] using the LAGAN alignment tool [47] with *P.*

nyererei as the reference. We applied the repeat masking option with fugu as the

reference. Transcription factors (TFs) binding prediction analyses were

conducted with MatInspector v8.4.1 with genomatix software suite

(https://www.genomatix.de) using default parameters with zebrafish as the reference

[48,49].

Positive selection detection of *lg* in cichlid fishes

To detect whether positive selection occurred in the evolution of *lg* in a

specific branch, a branch-site model within codeml in PAML v3.14 was

performed [50,51]. Fixed branch lengths (fix_blength = 2) retrieved under the M0

model were used. Alignment gaps and ambiguity characters were removed

(Cleandata = 1). The tree for PAML test is based on the phylogeny from the

study [24]. Rates of non-synonymous to synonymous substitutions (ω or dN/dS)

significantly larger than 1 represent that the sites in a specific target branch

(foreground branch) are under positive selection. A *chi*-square approximation

was calculated to test statistical significance. Detail statistics of branch-site

model comparisons can be found in Table S4.

CRISPR/Cas9 gene KO in zebrafish

The *lg* mutant line was generated by CRISPR/Cas9 system. The guide

RNA (gRNA) target site was selected using the CRISPOR v4.98 [52] with 20 bp

NGG PAM motif. The synthesized gRNA and Cas9 mRNA were co-injected

into the one-cell stage embryos of zebrafish at the concentration of 50 ng/ul.

The randomly picked 10 embryos at 1 dpf (days post fertilization) were lysated

and used to access functional gene knockout using PCR amplification with

primers designed at the target KO site (Table S6). Heterozygous F1 fish were

obtained by crossing F0 and wild-type (WT) zebrafish and were genotyped with

the same primers above using Sanger sequencing at the target site (Table S6).

Siblings of F1 generation harboring the same mutations were mated to generate

homozygous F2 mutants (MUT) and WT sisters (as the control).

Western blot

Since there is no available antibody for *lg*, we thus added a HA tag

(ATGTACCCATACGACGTCCCAGACTACGCT) in front of the coding

region of *lg* by PCR amplification (Table S6). The antibody of HA tag can thus

detect the fused *lg* protein if there is no disruption. To ligate the amplicons of

HA tagged coding regions of WT and MUT into pCS2 vector, the restriction

enzyme sites (*Bam*H I and *Xba* I) were added at the two ends of the amplicons

by PCR amplification (Table S6), followed by double digestion-system (NEB)

and ligation (Promega) (Table S6). The plasmids of pCS2, pCS2-lg-HAtag-WT

and pCS2-lg-HAtag-MUT (2 μg) (Fig. S2) were then transfected into HEK293T

cell line (ab255449, abcam), seperately. Protein was collected after 36 h

transfection to perform Western blot according to the standard procedure [53].

Antibodies of mouse anti-HA (901501 Biolegend, 1:2000) and mouse anti-

GAPDH (60004-1-lg Proteintech, 1:2000) were used for the detection.

Detections were performed with goat anti-mouse secondary antibody (31430

Thermo 1:5000) and enhanced chemiluminescence (ECL plus) reagent.

RNAseq analyses

To see gene expression profiles of *lg*, the available transcriptomic raw

data of different tissues in different fishes were extracted from SRA database

(Table S2). The raw reads were mapped to the corresponding transcriptome

retrieved from Ensembl using Hisat2-2.1.0 [54] with default parameters. The

reported SAM results were converted into BAM format, and finally were

recorded as count files using SAMtools v1.10 [55] followed by TPM values

calculation [56].

To see the affected genes after knockout, comparative transcriptomic

analyses were conducted between MUT and WT (Table S3). RNA extraction

was performed (each replicate pooled 30 individuals, six biological replicates

for each group) using TRIzol® reagent (Invitrogen). Six biological replicates of

WT and MUT of F2 generation were used for RNAseq analyses. Library

construction and sequencing were performed using Illumina TrueSeq RNAseq

technology at Novogene Company, China, with paired end reads length of 350

bp. The raw reads were mapped to the zebrafish transcriptome assembly using

the same methods mentioned above to get the reads count files. The differential

expressed (DE) transcripts between WT and MUT were further analyzed using

edgeR v3.20.7 (FDR<0.01). The raw data were deposited at NCBI BioProject

ID PRJNA611476. The top up- and down- 10 DE transcripts list can be found in

Table S3.

405 *in situ* **hybridization**

The gene fragments in the coding region of *lg* in zebrafish and tilapia

were amplified using PrimerSTAR (Takara) (Table S6). *lyz* probe synthesis was

conducted as described in the reference [57]. The amplified products were cloned

into pGEM®-T Easy vector (Promega). Plasmids were extracted and purified

using TIANprep Mini Plasmid Kit II (Tiangen). The insertion and direction of

probes were confirmed with Sanger sequencing using M13 PCR products
(Tsingke). RNA probes were synthesized with the DIG RNA labeling kit
(SP6/T7) (Roche). The experiment was conducted as described in the reference
[58] with biological replicates (n=30 for zebrafish, n=20 for tilapia, and n=10 for
P. nyererei). The embryos or larvae were mounted in 1% low melting point
Agarose (GBCBIO) and imaged with Zeiss Axio Imager Z2 microscopy.

Transgenic plasmids and transgenic lines construction

To see the regulatory functions of cichlid fish-specific upstream regions,
we constructed plasmids with different upstream regions as promoters to drive
eGFP expression in zebrafish (Fig. 4A and Fig. S2), i.e. 1) species in the MH
lineage (*P. nyererei*) (pT2AL-Pn-eGFP line); 2) the corresponding region of
species at the sister lineage of East African Great Lakes cichlid fishes (tilapia)
(pT2AL-On-eGFP line). 3) To see whether the three MH-segregated SNPs in
the cichlid fish-specific upstream region are important, we specifically deleted
these SNPs using site-directed mutagenesis by PCR primer extension (Table S6)
[59] to construct another transgenic line (pT2AL-Pn-noSNPs-eGFP line).
Schematic Plasmids construction can be found in Fig. S2.

DNA fragments for different plasmids construction were amplified with
PCR primers added with corresponding restriction endonuclease sites (Fig. S2,
Table S6). Fragments were then recovered (Tiangen), digested (NEB) and
ligated (Promega) into pT2AL vector following the instructions accordingly.
Successfully constructed recombinant plasmids were further confirmed by
Sanger sequencing (Tsingke). Injections were performed into one-cell stage of
zebrafish embryos. By outcrossing with WT zebrafish, we successfully obtained
F2 stable transgenic lines for all these lines (Pn line, On line, Pn-noSNPs line).
The fluroscent signals were imaged with Zeiss 880 confocal microscopy.

Immunohistochemistry (IHC)

The following primary antibodies were used: mouse anti-Pvalb7, 1:1000 provided by Prof. Masahiko Hibi, Nagoya University, Japan [60]; rabbit anti-Serotonin, 1:400 (s5545, Sigma); rabbit anti-Calretinin, 1:400 (ab702, abcam); goat anti-GFP, 1:400 (ab6673, abcam); mouse anti-HuC/HuD, 1:1000 (16A11, Invitrogen); rabbit anti-SOX10, 1:400 (ab229331, abcam); rabbit anti-PAX6 1:400 (ab209836, abcam); mouse anti-Collagen type II (II-II6B3, DSHB), rabbit anti-Cathepsin L/MEP (ab209708, abcam). Secondary antibodies are donkey to goat IgG H&L (Alexa Fluor® 488) (ab150133, abcam), donkey to mouse IgG H&L (Alexa Fluor® 647) (ab150107, abcam), donkey to rabbit IgG H&L (Alexa Fluor® 647) (ab150063, abcam) and DAPI (ab228549, abcam). Fish larvae were fixed in 4% Paraformaldehyde (Sigma) for 4 h at room temperature, followed by detail protocol as described before [61]. The fluroscent signals were imaged with Zeiss 880 confocal microscopy.

Immune response detection

To see whether there are different immune responses between MUT and WT zebrafish, we injected lipopolysaccharide (LPS), the key inflammatory components of Gram-negative bacteria into the circulation of 2 dpf zebrafish and counted the numbers of lyz^+ neutrophils in the caudal hematopoietic tissue at one day post injection. The lyz^+ whole-mount RNA *in situ* hybridization signals across eight somites of the larval CHT (caudal hematopoietic tissue) regions were manually counted. Statistics analyses can be found in Table S7.

Data Availability

The publicly available datasets that this study used are provided in the supplementary information. The RNAseq data of WT and MUT are available at NCBI (BioProject ID PRJNA611476). The ion torrent sequencing reads of *lg* gene are available at NCBI (BioProject ID PRJNA611690). The scaffolds

possess *lg* of teleost fishes across the phylogeny, the assembly of *lg* including upstream region of cichlid fishes across the phylogeny retrieved from ion torrent sequencing and sanger sequencing can be found in Dryad https://doi.org/10.5061/dryad.fxpnvx0qb. Source associated data of Fig. 3 and Fig. 4 are provided in the supplementary files. Other data are available upon reasonable request.

References

1. Dobzhansky, T. Nothing in biology makes sense except in the light of evolution. *Am. Biol. Teach.* **March,** 125–129 (1973).
2. Tautz, D. & Domazet-Lošo, T. The evolutionary origin of orphan genes. *Nat. Rev. Genet.* **12,** 692–702 (2011).
3. Bae, B.-I., Jayaraman, D. & Walsh, C. A. Genetic changes shaping the human brain. *Dev. Cell* **32,** 423–34 (2015).
4. Zhang, Y. E. & Long, M. New genes contribute to genetic and phenotypic novelties in human evolution. *Curr. Opin. Genet. Dev.* **29,** 90–6 (2014).
5. Babonis, L. S., Martindale, M. Q. & Ryan, J. F. Do novel genes drive morphological novelty? An investigation of the nematosomes in the sea anemone Nematostella vectensis. *BMC Evol. Biol.* **16,** 114 (2016).
6. Santos, M. E., Le Bouquin, A., Crumière, A. J. J. & Khila, A. Taxon-restricted genes at the origin of a novel trait allowing access to a new environment. *Science (80-.).* **358,** 386–390 (2017).
7. Khalturin, K. *et al.* A novel gene family controls species-specific morphological traits in Hydra. *PLoS Biol.* **6,** e278 (2008).
8. Garcia-Fernàndez, J. The genesis and evolution of homeobox gene clusters. *Nat. Rev. Genet.* **6,** 881–92 (2005).
9. Wu, M., Chen, G. & Li, Y.-P. TGF-β and BMP signaling in osteoblast, skeletal development, and bone formation, homeostasis and disease. *Bone Res.* **4,** 16009 (2016).
10. Wagner, G. P. & Lynch, V. J. The gene regulatory logic of transcription factor evolution. *Trends Ecol. Evol.* **23,** 377–85 (2008).
11. Wray, G. A. The evolutionary significance of cis-regulatory mutations. *Nat. Rev. Genet.* **8,** 206–216 (2007).
12. Zhang, J. Parallel adaptive origins of digestive RNases in Asian and African leaf monkeys. *Nat. Genet.* **38,** 819–23 (2006).
13. Hoekstra, H. E., Hirschmann, R. J., Bundey, R. A., Insel, P. A. & Crossland, J. P. A single amino acid mutation contributes to adaptive beach mouse color pattern. *Science* **313,** 101–4 (2006).

14.   Chen, L., DeVries, A. L. & Cheng, C.-H. C. Evolution of antifreeze
glycoprotein gene from a trypsinogen gene in Antarctic notothenioid fish.
Proc. Natl. Acad. Sci. **94,** 3811–3816 (1997).
15.   Carroll, S. B. Evolution at Two Levels: On Genes and Form. *PLoS Biol.* **3,**
e245 (2005).
16.   Hu, W., Chen, S., Zhang, R. & Lin, Y. Single nucleotide polymorphisms
in the upstream regulatory region alter the expression of myostatin. *Vitr.*
Cell. Dev. Biol. - Anim. **49,** 417–423 (2013).
17.   Marceca, C., Pfob, M., Schelling, G., Steinlein, O. K. & Eggert, M. Single
nucleotide polymorphism creating a variable upstream open reading frame
regulates glucocorticoid receptor expression. *Gene* **563,** 24–28 (2015).
18.   Malmstrøm, M. *et al.* Evolution of the immune system influences
speciation rates in teleost fishes. *Nat. Genet.* **48,** 1204–1210 (2016).
19.   Salzburger, W. The interaction of sexually and naturally selected traits in
the adaptive radiations of cichlid fishes. *Mol. Ecol.* **18,** 169–85 (2009).
20.   SH, O. & LI, Z. Hematopoiesis: An Evolving Paradigm for Stem Cell
Biology. *Cell* **132,** 631–44 (2008).
21.   Kim, C.-H. *et al.* Zebrafish elav/HuC homologue as a very early neuronal
marker. *Neurosci. Lett.* **216,** 109–112 (1996).
22.   Hugo, S. E. *et al.* A monocarboxylate transporter required for hepatocyte
secretion of ketone bodies during fasting. *Genes Dev.* **26,** 282–93 (2012).
23.   Salzburger, W. Understanding explosive diversification through cichlid
fish genomics. *Nat. Rev. Genet.* (2018). doi:10.1038/s41576-018-0043-9
24.   Meyer, B. S., Matschiner, M. & Salzburger, W. A tribal level phylogeny
of Lake Tanganyika cichlid fishes based on a genomic multi-marker
approach. *Mol. Phylogenet. Evol.* **83,** 56–71 (2015).
25.   Salzburger, W., Mack, T., Verheyen, E. & Meyer, A. Out of Tanganyika:
genesis, explosive speciation, key-innovations and phylogeography of the
haplochromine cichlid fishes. *BMC Evol. Biol.* **5,** 17 (2005).
26.   Hubel, P. *et al.* A protein-interaction network of interferon-stimulated
genes extends the innate immune system landscape. *Nat. Immunol.* **20,**
493–502 (2019).
27.   Denève, S., Alemi, A. & Bourdoukan, R. The Brain as an Efficient and
Robust Adaptive Learner. *Neuron* **94,** 969–977 (2017).
28.   Zhang, W., Landback, P., Gschwend, A. R., Shen, B. & Long, M. New
genes drive the evolution of gene interaction networks in the human and
mouse genomes. *Genome Biol.* **16,** 202 (2015).
29.   Nagasawa, T. *et al.* Evolutionary Changes in the Developmental Origin of
Hatching Gland Cells in Basal Ray-Finned Fishes. *Zoolog. Sci.* **33,** 272
(2016).
30.   Conith, A. J., Kidd, M. R., Kocher, T. D. & Albertson, R. C.
Ecomorphological divergence and habitat lability in the context of robust
patterns of modularity in the cichlid feeding apparatus. *BMC Evol. Biol.*

20, 95 (2020).

31.   Roberts, R. B., Hu, Y., Albertson, R. C. & Kocher, T. D. Craniofacial
divergence and ongoing adaptation via the hedgehog pathway. *Proc. Natl.*
Acad. Sci. U. S. A. **108,** 13194–9 (2011).

32.   Albertson, R. C., Streelman, J. T., Kocher, T. D. & Yelick, P. C.
Integration and evolution of the cichlid mandible: the molecular basis of
alternate feeding strategies. *Proc. Natl. Acad. Sci. U. S. A.* **102,** 16287–92
(2005).

33.   König, D. *et al.* Distribution and Restoration of Serotonin-
Immunoreactive Paraneuronal Cells During Caudal Fin Regeneration in
Zebrafish. *Front. Mol. Neurosci.* **12,** 227 (2019).

34.   Coccimiglio, M. L. & Jonz, M. G. Serotonergic neuroepithelial cells of the
skin in developing zebrafish: morphology, innervation and oxygen-
sensitive properties. *J. Exp. Biol.* **215,** 3881–3894 (2012).

35.   Kermen, F., Franco, L. M., Wyatt, C. & Yaksi, E. Neural circuits
mediating olfactory-driven behavior in fish. *Front. Neural Circuits* **7,** 62
(2013).

36.   Green, S. A., Simoes-Costa, M. & Bronner, M. E. Evolution of vertebrates
as viewed from the crest. *Nature* **520,** 474–82 (2015).

37.   Salzburger, W. Understanding explosive diversification through cichlid
fish genomics. *Nat. Rev. Genet.* **19,** 705–717 (2018).

38.   Nyman, C., Fischer, S., Aubin-Horth, N. & Taborsky, B. Evolutionary
conserved neural signature of early life stress affects animal social
competence. *Proceedings. Biol. Sci.* **285,** (2018).

39.   Culbert, B. M., Gilmour, K. M. & Balshine, S. Stress axis regulation
during social ascension in a group-living cichlid fish. *Horm. Behav.* **103,**
121–128 (2018).

40.   Stanke, M. & Morgenstern, B. AUGUSTUS: a web server for gene
prediction in eukaryotes that allows user-defined constraints. *Nucleic*
Acids Res. **33,** W465–7 (2005).

41.   Stanke, M., Steinkamp, R., Waack, S. & Morgenstern, B. AUGUSTUS: a
web server for gene finding in eukaryotes. *Nucleic Acids Res.* **32,** W309–
12 (2004).

42.   Kearse, M. *et al.* Geneious Basic: an integrated and extendable desktop
software platform for the organization and analysis of sequence data.
Bioinformatics **28,** 1647–9 (2012).

43.   Lalitha, S. Primer Premier 5. *Biotech Software&Internet Rep.* **1,** (2000).

44.   Afgan, E. *et al.* The Galaxy platform for accessible, reproducible and
collaborative biomedical analyses: 2018 update. *Nucleic Acids Res.* **46,**
W537–W544 (2018).

45.   Mayor, C. *et al.* VISTA : visualizing global DNA sequence alignments of
arbitrary length. *Bioinformatics* **16,** 1046–1047 (2000).

46.   Frazer, K. A., Pachter, L., Poliakov, A., Rubin, E. M. & Dubchak, I.

VISTA: computational tools for comparative genomics. *Nucleic Acids Res.* **32,** W273–9 (2004).

47. Brudno, M. *et al.* LAGAN and Multi-LAGAN: efficient tools for large-scale multiple alignment of genomic DNA. *Genome Res.* **13,** 721–31 (2003).

48. Quandt, K., Frech, K., Karas, H., Wingender, E. & Werner, T. MatInd and MatInspector: new fast and versatile tools for detection of consensus matches in nucleotide sequence data. *Nucleic Acids Res.* **23,** 4878–84 (1995).

49. Cartharius, K. *et al.* MatInspector and beyond: promoter analysis based on transcription factor binding sites. *Bioinformatics* **21,** 2933–2942 (2005).

50. Yang, Z. PAML: a program package for phylogenetic analysis by maximum likelihood. *Comput. Appl. Biosci.* **13,** 555–6 (1997).

51. Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24,** 1586–91 (2007).

52. Concordet, J.-P. & Haeussler, M. CRISPOR: intuitive guide selection for CRISPR/Cas9 genome editing experiments and screens. *Nucleic Acids Res.* **46,** W242–W245 (2018).

53. Zhao, F. *et al.* Irf8 regulates the progression of myeloproliferative neoplasm-like syndrome via Mertk signaling in zebrafish. *Leukemia* **32,** 149–158 (2018).

54. Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner with low memory requirements. *Nat. Methods* **12,** 357–360 (2015).

55. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25,** 2078–9 (2009).

56. Wagner, G. P., Kin, K. & Lynch, V. J. Measurement of mRNA abundance using RNA-seq data: RPKM measure is inconsistent among samples. *Theory Biosci.* **131,** 281–285 (2012).

57. Hou, Y. *et al.* Systemic inoculation of Escherichia coli causes emergency myelopoiesis in zebrafish larval caudal hematopoietic tissue. *Sci. Rep.* **6,** 36853 (2016).

58. Westerfield, M. *The zebrafish book. A guide for the laboratory use of zebrafish (Danio rerio). 4th edition.* (University of Oregon Press, Eugene, 2000).

59. Reikofski, J. & Tao, B. Y. Polymerase chain reaction (PCR) techniques for site-directed mutagenesis. *Biotechnol. Adv.* **10,** 535–47 (1992).

60. Nimura, T. *et al.* Role of Reelin in cell positioning in the cerebellum and the cerebellum-like structure in zebrafish. *Dev. Biol.* **455,** 393–408 (2019).

61. Barresi, M. J. F., Stickney, H. L. & Devoto, S. H. The zebrafish slow-muscle-omitted gene product is required for Hedgehog signal transduction and the development of slow muscle identity. *Development* **127,** 2189–2199 (2000).

Acknowledgements

Many thanks to Prof. Walter Salzburger, Attila Rüegg, Zhen Liu, Zhiqiang Zhou and Hesheng Xiao for the sampling and suggestions. We thank Prof. Masahiko Hibi and Prof. Kiyoshi Naruse for providing the antibody Pvalb7. We also thank Prof. Lingfei Luo, Prof. Hua Ruan, Prof. Yang Liu and Prof. Tienming Lee for the laboratory supports. Many thanks to Prof. Zuzana Musilova, Dr. Xiaogui Yi, Dr. Yandong Zhan, Dr. Fangying Zhao, Yuepeng He, Xinkai Cheng, Qing Li and Yue Gao for the supports. Thanks to Huansheng Tao and Jialing Xu for the Confocol microscopy support, Zhao Long for fluroscent microscopy support. This study was funded by grants from the National Natural Science Foundation of China (31802283), Guangdong Basic and Applied Basic Research Foundation (2020A1515010330); the Fundamental Research Funds for the Central Universities (20lgpy109); China Postdoctoral Science Foundation (2018 M633311) and Doctoral Scholarship of University of Basel, Switzerland to LG; Grants from the National Natural Science Foundation of China (31630082) to DW, and (31771623 and 31571500) to LL. Part of -omics analyses was supported by National Supercomputer Center in Guangzhou, China.

Authors contributions

LG, DW and LL conceived this work. LG carried out –omics sequencing and evolutionary analyses. LG, CL, XM and ZW conducted the *in situ* hybridization. LG carried out the IHC. XH and WZ participated the IHC experiments of serotonin-positive cells and olfactory epithelium cells. LG, CL and XM constructed plasmids of the Pn line and the On line. LG constructed plasmids of the Pn-noSNPs line. LG leads the transgenic lines and mutant lines construction with the help from CL and XM. LG took the confocal photos. YH

667 performed the western-blot experiment. ZW carried out the LPS immune
668 response experiments. LG conducted the fasting ORO staining experiments. LG
669 wrote the initial manuscript draft, and all authors commented on the manuscript
670 and approved the final version.
671
672 **Competing interests**: The authors declare no competing interests.
673
674 **Figure legends**
675
676 **Fig.1 Sequence evolution of *lg*.** a. Comparative genomic analyses showed that
677 *lg* is a fish-specific gene. b. Multiple losses occurred in different lineages within
678 teleost fishes. The reference tree is from [18].
679
Fig.2 Expression profiles of *lg* in fishes. a. Transcriptomic analyses of *lg* gene
expression patterns in adult fish tissues. *lg* mainly expressed in the brain and
eyes in most fishes, but evolved new expression patterns in the skin in the
modern haplochromine (MH) lineage of cichlid fishes. b. *in situ* hybridization in
zebrafish and cichlid fish larvae. *lg* mainly expressed in the brain, ganglion and
inner plexiform cell layers in the eyes of zebrafish. No expression was detected
in the hatching gland of 24 hpf (hours post fertilization) embryos of zebrafish.
lg highly expressed in the brain, eyes and pharyngeal arches in the
haplochromine cichlid fish, *Pundamilia nyererei*, but weakly expressed in the
cichlid fish, *Oreochromis niloticus*.
690
Fig.3 *lg* gene knockout in zebrafish and comparisons between mutant
(MUT) and wild-type (WT) zebrafish. A. *lg* is successfully disrupted using
CRISPR/Cas9. This is confirmed by both Sanger sequencing (up) and Western-
blot (down). B. Immune responses comparisons between MUT and WT using
whole mount *in situ* hybridization in the CHT (caudal hematopoietic tissue)

treated with LPS and PBS (n=10 for WT, and n=18 for MUT), respectively.

Quantification of the numbers of lyz^+ signals across eight somites were shown

in dots. n.s. not significant. Mean and SD error bar were presented. C.

phenotype comparisons between MUT and WT zebrafish. Brain and eye

phenotypes in mutants were examined with neuron marker HuC. No obvious

phenotype was found. There is no obvious mutant phenotype with ORO

staining under fasting condition, either, which detect energy metabolism [22].

Fig.4 *lg* sequence evolution and regulatory function evolution in cichlid

fishes. Aa. Both upstream region and coding region exhibited SNPs segregated

with the modern haplochromine (MH) lineage. i.e. three SNPs (GTA) in the

upstream region and one nonsynonymous amino acid change (A to S) located at

the 38[th] position which is under positive selection (Table S4). Transcription

factors binding sites prediction showed that these SNPs can set up the

opportunity to bind to glucocorticoid modulatory element binding protein 2

(GMEB2). The reference tree is from [24]. Ab. Three types of transgenic plasmids

construction. The pT2AL-On-eGFP line (On line) adopted the upstream region

of *O. niloticus* as the promoter to drive eGFP expression. The pT2AL-Pn-eGFP

line (Pn line) used the homologous upstream region of *P. nyererei* in the MH

lineage as the promoter. The pT2AL-Pn-no-SNPs-eGFP line (Pn-noSNPs line)

used the same promoter as the Pn line but deleted the three MH-segregated

SNPs (GTA). B. The fluorescence images in the Pn line. Merged figures of

double staining of eGFP and cell marker antibodies were shown if there are

available antibodies. Red is from the cell marker antibody. Green is from eGFP.

Blue is from DAPI. Pictures were taken from 4 dpf (days post fertilization)

zebrafish except the hatching gland which is from 24 hpf (hours post

fertilization) embryos. C. Different expression patterns of three transgenic lines

in zebrafish. D. Statistics of expression patterns comparisons among three

transgenic lines, focusing on cerebellum, hindbrain, and the first two pharyngeal

arches. The Pn line showed significantly stronger fluroscence than the other two
lines. With the three MH-segregated SNPs deletion, Pn-noSNPs line showed
weak fluroscent expressions, similar to the On line. Mean and SD error bars
were presented. n=6 for each measurement. Detail statistical analyses can be
found in Table S5.

Fig.5 Hypothesis of *lg* sequence and function evolution with respect to evo-
devo. *lg* is a fish-specific gene. Its ancestral function is not indispensable.
Interestingly, *lg* evolves a cichlid fish-specific upstream region, thus gained new
regulatory functions to drive gene expression in new tissues related to
adaptation, such as in pharyngeal cartilages and the hatching gland in cichlid
fishes. Especially in the most species-richness lineage, the modern
haplochromine (MH) lineage, the upstream of which not only drive stronger
eGFP expression patterns compared to the homologous region of tilapia, but
also evolved new profiles again, in epidermal serotonin-positive cells. And this
can be simply due to the three MH-segregated SNPs in the upstream region
which are predicted to bind to a hormone related transcription factor, the
glucocorticoid modulatory element binding protein 2 (GMEB2). If this is true, it
opens a new opportunity to response to the external environemnt. Besides, the
coding region of lg also possesses one MH- segregated amino acid mutation
which is under positive selection, suggesting that new function might also
evolved in the coding region which deserve further investigation. lg thus
provides a good example of new gene function emergence in a redundant
lineage-specific gene with respect to evo-devo.

 Figures

 Fig.1 Sequence evolution of *lg*.

Fig.2 Expression profiles of *lg* in fishes.

Fig.3 *lg* **gene knockout in zebrafish and comparisons between mutant (MUT) and wild-type (WT) zebrafish.**

764 **Fig.4 *lg* sequence evolution and regulatory function evolution in cichlid**

765 **fishes.**

Fig.5 Hypothesis of *lg* sequence and function evolution with respect to evo-devo.

Supplementary information

780   Table S7 The numbers of LPS immune response signals comparisons between

781   WT and MUT zebrafish

782   Supplementary Figures