1 **A new fish-specific gene evolves new functions in cichlid fishes**

2 Langyu Gu[1,2,3,]*, Chenzheng Li[2], Xiaobing Mao[2], Zongfang Wei[2], Youkui

3 Huang[2], Ximin He[1], Wenjun Zhou[1], Li Li[2,]*, Deshou Wang[2,]*

4 **1** State Key Laboratory of Biocontrol, School of Life Sciences and School of
5 Ecology, Sun Yat-sen University, Guangzhou, Guangdong, China.
6
7 **2** Key Laboratory of Freshwater Fish Reproduction and Development (Ministry
8 of Education), Key Laboratory of Aquatic Science of Chongqing, School of
9 Life Sciences, Southwest University, Chongqing, China
10
11 **3** Zoological Institute, Department of Environmental Sciences, University of
12 Basel, Basel, Switzerland
13

16 *Correspondence:

17 Langyu Gu: langyugu@gmail.com

18 Li Li: alisir@swu.edu.cn

19 Deshou Wang: wdeshou@swu.edu.cn
20
21
22

Abstract

Genes whose dysfunction does not affect normal survival are common. Are they only meaningless residues during evolution? Here, we identified a new fish-specific gene, which we named *lg*. Gene knockout resulted in no obvious phenotype in zebrafish, but *lg* evolved an amino acid mutation that was under positive selection in the modern haplochromine (MH) cichlid fish lineage, the well-known adaptive radiative lineage. Moreover, the cichlid fish-specific upstream region of *lg* drove new eGFP expression in tissues related to adaptation. Noticeably, this homologous region from different cichlid fishes drove different patterns, which is simply due to three MH-segregated SNP mutations that are predicted to bind a hormone-related transcription factor. We thus revealed an initially redundant gene evolving new functions in an adaptive radiative lineage. This further illuminates the mechanism of the emergence of new gene functions with respect to evo-devo in a broad way.

Key words
lg, new gene, lineage-specific, evo-devo, fish

From an evolutionary point of view, the most adaptive organism spends

the least resources but produces the most outputs. However, it seems that this

rule does not apply to genome organization, which is occupied by a large

proportion of genes whose dysfunction does not affect normal survival. Does

this make sense?

Nothing in biology makes sense except in the light of evolution [1]. As raw

materials for natural selection, the large portion of redundant genes provides

good opportunities for functional innovation. This is especially true for lineage-

specific genes, which can occupy up to one-third of the genome in many

organisms [2]. Studies have gradually revealed that many of these genes can

evolve new functions related to novelties and environmental adaptation in a

particular lineage. For example, in early brain development in humans [3,4], in the

creation of a novel tissue in the sea anemone [5], and in the development of a

novel fan trait on the leg in a water spider genus to explore new environments [6].

There is even a whole lineage-specific gene family in water fleas related to

environmental stimuli [7]. Lineage-specific genes thus provide a new perspective

on the emergence of new gene function.

It is well accepted that different organisms share deep homologous

developmental molecular bases (molecular toolkits), such as highly conserved

signalling pathways and transcription factors [8–10]. Then, how do new genes gain

new functions, and in what way are they integrated into the existing networks?

Although usually restricted due to pleiotropic effects [11], protein coding region

evolution can take place [12–14]. In comparison, regulatory evolution is a well-

known strategy, since changes at the expression level are more tolerable and can

offer immediate phenotypic consequences [11,15]. Even a single mutation in the

upstream region can induce new expression profiles [16,17]. The relaxed selection

pressure induced by the changing expression profiles can further prompt the
accumulation of mutations at the protein level.

In this study, we identified a new unannotated fish-specific gene, which
we named *lg* (ENSONIT00000026203). Comparative -omics analyses and
functional experiments revealed that multiple losses of *lg* occurred in different
fish lineages, and no obvious phenotype was detected in gene knockout
zebrafish, which led us to assume that the ancestral function of *lg* is not
indispensable. *lg* was mainly expressed in the brain and eyes in most fishes, but
it evolved new expression patterns, such as in pharyngeal cartilages and
epidermal serotonin-positive neuroepithelia cells, in the cichlid fish lineage,
especially in the most species-rich lineage, the modern haplochromine (MH)
lineage. Further comparative genomic analyses across the East African cichlid
fish phylogeny revealed that *lg* evolved a cichlid fish-specific upstream region
possessing three MH lineage-segregated SNPs and one MH lineage-segregated
amino acid mutation that was under positive selection. Transgenic experiments
confirm the new regulatory functions of this upstream region to drive eGFP
expression in zebrafish. Noticeably, the homologous sequence from different
cichlid fishes drove different eGFP expression patterns, which was due to the
three MH lineage-segregated SNP mutations that are predicted to bind to a
hormone-related transcription factor. Taken together, our work revealed for the
first time an initially redundant new fish-specific gene that evolved new
functions in an adaptive radiative cichlid fish lineage.

Results

***lg* is a new fish-specific gene**

To see *lg* sequence evolution, we conducted a tblastx search in all species
with available genomes in Ensembl 100. The results showed that *lg* only

appeared in fishes (Fig. 1A, Table S1). To further characterize its evolution within teleost fishes, we conducted a tblastx search and gene predictions in the draft genome assembly across the whole phylogeny of teleost fishes [18]. The results showed that *lg* was lost multiple times in different lineages (Fig. 1B), which inspired the assumption that the ancestral function of *lg* is not indispensable for fish survival.

lg evolved new expression profiles in cichlid fishes, especially in the MH lineage

To identify the tissues where *lg* was expressed, we investigated its expression profiles based on the combined results of available RNAseq data retrieved from NCBI and *in situ* hybridization in different fishes (Fig. 2, Table S2). *lg* is mainly expressed in the brain and eyes in most fishes. Interestingly, it evolved new expression profiles in cichlid fishes in pharyngeal cartilages, as well as in the skin of the most species-rich lineage of cichlid fishes—the MH lineage [19]. These results suggest that the ancestral function of *lg* can be related to neurons, but new functions can evolve in other tissues in cichlid fishes, especially in the MH lineage.

No obvious phenotype was detected in lg gene knockout zebrafish

To further investigate the function of *lg*, gene knockout was conducted using CRISPR/Cas9 in zebrafish. We successfully obtained F2 homozygous mutant (MUT) zebrafish and their wild-type siblings (WT). Sanger sequencing and Western blot analyses confirmed the successful gene dysfunction with frameshift (5 bp deletion) and protein interruption (stop codon appeared at the 18[th] amino acid) (Fig. 3A). To determine the affected gene expression after gene knockout, comparative transcriptomic analysis was conducted between MUT and WT zebrafish at 2 dpf (days post fertilization) and 4 dpf, respectively. The results showed that most differentially expressed (DE) genes were related to

immunity, neurons and energy (Table S3). To determine whether there are

different immune responses between MUT and WT, we examined the immune

responses of larval zebrafish to lipopolysaccharide (LPS) by counting the

numbers of lyz^+ neutrophils in the caudal haematopoietic tissue one day after

injection [20]. No significant difference was detected (Fig. 3B). To determine

whether gene knockout affects neuronal development in the brain and eyes

where *lg* is expressed in zebrafish, we compared the phenotypic changes

between MUT and WT using the brain neuron marker HuC [21] and still no

difference was detected (Fig. 3C). Then, we turned our attention to one top DE

gene, *slc16a6*, which has been reported to be related to energy metabolism

under fasting conditions [22]. Under the same strategy, however, we still could not

find obvious metabolic phenotypic changes between MUT and WT (Fig. 3C).

This can be explained by the fact that *slc16a6* is only downregulated here

instead of entirely disrupted, as in the reference [22], so that it can still execute its

function. Overall, mutation of *lg* resulted in no obvious phenotypes in zebrafish.

These results further support our assumption that the ancestral function of *lg* is

not indispensable.

***lg* evolved a cichlid fish-specific upstream region possessing MH-segregated**

SNPs and one MH-segregated amino acid mutation that is under positive

selection

Noticeably, as we mentioned above, *lg* evolved new gene expression

profiles in cichlid fishes (Fig. 2), the classical model species for studying

adaptive radiation, which explosively radiated into hundreds to thousands of

species within a short time with very closely related genomes [23]. To see whether

there are cichlids-specific region that might conduct the new regulatory

function, genomic alignment of *lg* among cichlid fishes and other teleost fishes

were conducted (Fig. S1). Indeed, cichlid fishes showed a large portion of

cichlid fish-specific CNS (Conserved Non-Coding Sequences) in the upstream

region. To further characterize its sequence evolution within cichlid fishes, we

amplified the whole *lg* gene, including this CNS upstream region, in a total of

20 cichlid fishes representing main lineages across the East African cichlid fish

phylogeny [24] (Fig. 4A). Interestingly, both the upstream and the coding regions

exhibited segregation patterns with the MH lineage, the most species-rich

lineage of cichlid fishes [25] (Fig. 4A). For example, within the cichlid fish

specific CNS upstream region, there are three MH lineage-segregated SNP

mutations (Fig. 4A). Transcription factors (TFs) binding prediction revealed

that these three SNPs can set up binding opportunities for a hormone-related

TF, the glucocorticoid modulatory element binding protein 2 (GMEB2) (Fig.

4A). In addition, there is one nonsynonymous amino acid mutation (A to S)

located at the 38th residue in the coding region specific to the MH lineage that is

under positive selection (Fig. 4A, Table S4). These results prompted us to pay

special attention to the *lg* of cichlid fishes in the next steps.

187   The cichlid fish-specific upstream region indeed has regulatory functions

188   and performs different roles in different cichlid fish lineages

To determine whether the cichlid fish-specific upstream region of *lg*

indeed has regulatory functions, and whether this region of the MH lineage has

different regulatory roles compared to other cichlid fish lineages, we

constructed two transgenic lines with the homologous upstream regions from

different species as the promoters to drive eGFP expression in zebrafish (Fig.

4A, Fig. S2). One line is with the upstream region of an MH cichlid fish,

Pundamilia nyererei (pT2AL-Pn-eGFP transgenic line, Pn line). The other line

is with the corresponding homologous region of a cichlid fish belonging to a

sister lineage of East African Great Lakes cichlid fishes, Nile tilapia

(*Oreochromis niloticus*) [23] (pT2AL-On-eGFP transgenic line, On line). We

successfully obtained F2 generations for both transgenic lines. Both lines can

drive eGFP expression in zebrafish, confirming their regulatory functions, but

with different patterns (Fig. 4B and Fig. 4C). The Pn line showed strong

expression in neurons in most regions of the brain (olfactory epithelium,

olfactory bulb, forebrain, optic tectum, cerebellum and hindbrain), eyes, spinal
cord, and heart valves. In addition, the Pn line also showed new strong eGFP
expression in the hatching gland in embryos, pharyngeal arches, chondrocyte
cells in the fin and scattered epidermal serotonin-positive cells in the skin. The
positive eGFP cells were further confirmed by available antibodies (Fig. 4B).
Interestingly, compared to the Pn line, eGFP fluroscence was weak in the brain
and pharyngeal arches in the On line ($p<0.01$), and was hard to detect in
epidermal serotonin-positive cells and fin chondrocytes (Fig. 4C). Considering
that we obtained different F1 individuals of each line possessing the same stable
eGFP expression patterns, and that the eGFP expression patterns of different
lines were consistent with the *in situ* hybridization and RNAseq results above,
the transgenic lines we constructed were stable and credible. These results
indicated that the cichlid fish-specific upstream region of *lg* indeed evolved new
regulatory functions, i.e., not only executing the ancestral expression profiles in
neurons but also evolving new expression patterns in the hatching gland and
pharyngeal arches in cichlid fishes. Strong and new regulatory functions further
evolved in the scattered epidermal serotonin-positive cells in the MH cichlid
fish lineage.
221

Different regulatory roles of cichlid fish-specific upstream regions are
simply due to three MH lineage-segregated SNPs

To determine whether the strong and new expression profiles in the Pn
transgenic line were caused by the three MH lineage-segregated SNPs in the
upstream region, we constructed another transgenic line with the same upstream
region as in the Pn line but deleted these three SNPs (pT2AL-Pn-noSNPs-eGFP
line, Pn-noSNPs line) (Fig. 4C, Fig. S2). Interestingly, only with these three
SNP deletions did the Pn-noSNPs line clearly show significantly different
expression patterns from the Pn line but similar patterns to the On line (Fig. 4C,
Fig. 4D and Table S5). These results confirmed that these SNPs are the key
reason for the different expression patterns between the Pn and On lines.
233

Discussion

Overall, we unravelled the evolution of a fish-specific new gene, *lg*, for the first time. No obvious phenotypic changes were detected in the *lg* gene knockout zebrafish. This can be due to compensation by other genes in the correponding gene network. This robust nature is not unexpected for immune and brain network systems [26,27], as the top DE genes of the *lg* gene knockout line are mainly neurons and immunity-related genes. In addition, lineage-specific new genes are initially usually located at the edge of the gene network [28], and their disruption thus does not greatly affect the core gene network. Regardless, mutation of *lg* does not affect zebrafish survival. Together with its multiple losses in different fish lineages during evolution, we propose that the ancestral function of *lg* is not indispensable for fish survival.

Interestingly, *lg* evolved new regulatory functions in cichlid fishes, especially in its most species-rich lineage—the MH lineage. For example, in the hatching gland, which is a key innovation in regulating the hatching period of embryos to become free-living larvae [29]; in pharyngeal cartilages, the well-known tissues related to feeding and ecological niche exploration are key to cichlid fish adaptive radiation [30–32]; in scattered epidermal serotonin-positive cells in the skin which are neuroepithelia cells related to environmental stimuli, such as by water flow, pheromones and hormones [33,34]; and in the olfactory epithelium, which is well known for the detection of odorants, such as amino acids, hormones and pheromones [35]. Given that some of these tissues are derived from a key vertebrate innovation—neural crest cells [36]—*lg* can also be causally linked to neural crest cell development and differentiation. Together, *lg* gained new regulatory functions in most species-rich lineages of cichlid fishes, the MH lineage, which can be causally linked to adaptive radiation. Noticeably, although we focused mainly on the regulatory function evolution of *lg* in cichlid fishes here, the newly important function can also evolve in their coding region. Molecular evolutionary analyses have already shown that there is one amino

acid mutation segregated with the MH cichlid fish lineage and that it is under
positive selection; this mutation might be causally linked to adaptive radiation
of cichlid fishes. Further gene knockout of *lg* in MH cichlid fishes will help test
this hypothesis and unravel the corresponding involved pathways. In addition,
the evolution of *lg* gene function may be ongoing in cichlid fishes, considering
that they are evolutionarily young species [37].

DNA in the genome that was originally thought of as junk, such as
transposable elements, noncoding regions and nonfunctional genes, remains
surprising. How can a new gene function be evolved and integrated into the
existing gene network? Here, we propose one hypothesis of the functional
evolution of *lg* that can be applied in a broad way (Fig. 5): the relaxed selection
pressure on the lineage-specific gene in the ancesral lineage frees it from
accumulating mutations in both the regulatory and the coding regions. The
evolution of the regulatory region in *lg* offers immediate effects, i.e., only three
SNP mutations in the upstream region can provide new TFs binding
opportunities to build new gene network connections to affect gene expression
profiles in the most species-rich MH lineage of cichlid fishes. It will be
interesting to test our predictions of the binding of a glucocorticoid hormone
related TF with these SNPs in the future. If this is true, these SNPs can
efficiently connect the organism with external environments, since
glucocorticoids are well-known stress-related hormones to regulate animal
behaviour in response to the environment [38,39]. The *lg* gene we investigated here
thus provides a good example of new gene function evolution in an initially
redundant lineage-specific gene with respect to evo-devo in a broad way.

Methods

sampling
Samples of cichlid fishes for comparative genomic analyses were
provided by Prof. Walter Salzburger in University of Basel, Switzerland; cichlid

fish *Pundamila nyererei* for transgene plasmids construction was obtained from

IGV (Interessengemeinschaft Viktoriasee-Cichliden); Strains of AB (for

transgenic lines construction) and ABTü (for gene knockout) zebrafish (*Danio*

rerio) were raised under standard condition (28.5°C, 14/10 h light/dark). Tilapia

(*Oreochromis niloticus*) were raised at 26°C, 14/10 h light/dark. Animal

experiments have been approved by the cantonal veterinary office in Basel

under permit number 2317 in University of Basel, Switzerland; the Committee

of Laboratory Animal Experimentation at Sun Yat-sen University and

Southwest University, China.

-omics analyses to characterize the evolution of lg across fish phylogeny

To characterize the sequence evolution of *lg*, tblastx was conducted with

-e 0.001 in fishes with good genome assembly in Ensembl 100 (Table S1) and

local blast v2.9.0 (tblastx) was conducted with draft genomes assembly across

the phylogeny according to the reference [18]. After obtaining hits, the scaffolds

possessing the corresponding sequences were extracted to predict the coding

regions with Augustus 3.3.3 [40,41]. The predicted *lg* coding sequences were

further confirmed by aligning the translated amino acids with the available

sequences from Ensembl 100 using Geneious 8.0.5 [42].

.

-omics analyses to characterize the evolution of lg within cichlid fishes

To further characterize the sequence evolution of *lg* within East African

cichlid fishes, we amplified the whole gene sequence and its upstream region

(ca.10 kb in total) in different cichlid fishes. Combined with available cichlid

fish genomes, we studied the sequence evolution in a total of 20 cichlid fishes

representing main lineages across the phylogeny [24]. Genomic DNA was

extracted using a DNeasy Blood & Tissue kit (Qiagen). PCR primers were

designed using Primer premier 5 [43]. High-fidelity PCR Master Mix (BioLabs)

and a touch-down annealing process were used for amplification. PGM Ion

Torrent was used to sequence long amplicons (400 bp) with CHIP 316 (Life

Technologies). Galaxy v20.09 [44] was used to trim low-quality reads (window

size 2, step size 1, minimum quality score 20) and filter short reads with a

minimum size of 40 bp. The sequence of a cichlid fish, tilapia, retrieved from

the Ensembl database was used as the reference to recheck the assembled

sequences. The sequencing reads of the *lg* gene across the cichlid fish

phylogeny are available at NCBI (BioProject ID PRJNA611690). The ion

torrent sequencing of most species was successful except *Limnochromis*

abeelei, *Ophthalmotilapia ventralis*, and *Bathybates graueri*, whose sequences

were further obtained from Sanger sequencing (Primers can be found in Table

S6).

Genomic alignment of *lg* of cichlid fishes and other teleost fishes was

conducted with mVISTA [45,46] using the LAGAN alignment tool [47] with *P.*

nyererei as the reference. We applied the repeat masking option with fugu as the

reference. Transcription factors (TFs) binding prediction analyses were

conducted with MatInspector v8.4.1 with the genomatix software suite

(https://www.genomatix.de) using default parameters with zebrafish as the reference

[48,49].

Positive selection detection of *lg* in cichlid fishes

To detect whether positive selection occurred in the evolution of *lg* in a

specific branch, a branch-site model within the codeml in PAML v3.14 was

performed [50,51]. Fixed branch lengths (fix_blength = 2) retrieved under the M0

model were used. Alignment gaps and ambiguous characters were removed

(Cleandata = 1). The tree for the PAML test is based on the phylogeny from the

study [24]. Rates of nonsynonymous to synonymous substitutions (ω or dN/dS)

significantly larger than 1 indicate that the sites in a specific target branch

(foreground branch) are under positive selection. A *chi*-squared approximation

was calculated to test statistical significance. Detail statistics of branch-site

model comparisons can be found in Table S4.

CRISPR/Cas9 gene knockout in zebrafish

The *lg* mutant line was generated by the CRISPR/Cas9 system. The guide

RNA (gRNA) target site was selected using the CRISPOR v4.98 [52] with a 20 bp

NGG PAM motif. The synthesized gRNA and Cas9 mRNA were coinjected

into one-cell stage zebrafish embryos at a concentration of 50 ng/µl. Ten

randomly selected embryos at 1 dpf (days post fertilization) were lysed and used

to assess functional gene knockout using PCR amplification with primers

designed at the target knockout site (Table S6). Heterozygous F1 fish were

obtained by crossing F0 and wild-type (WT) zebrafish and genotyped with the

same primers above using Sanger sequencing at the target site (Table S6).

Siblings of the F1 generation harbouring the same mutations were mated to

generate homozygous F2 mutants (MUT) and WT sisters (as the control).

Western blot

Since there was no available antibody for *lg*, we added an HA tag

(ATGTACCCATACGACGTCCCAGACTACGCT) in front of the coding

region of *lg* by PCR amplification (Table S6). The antibody of the HA tag can

thus detect the fused *lg* protein if there is no disruption. To ligate the amplicons

of HA-tagged coding regions of WT and MUT into the pCS2 vector, restriction

enzyme sites (*Bam*H I and *Xba* I) were added at the two ends of the amplicons

by PCR amplification (Table S6), followed by a double digestion system (NEB)

and ligation (Promega) (Table S6). The plasmids pCS2, pCS2-lg-HAtag-WT

and pCS2-lg-HAtag-MUT (2 µg) (Fig. S2) were then transfected into the

HEK293T cell line (ab255449, Abcam) separately. Protein was collected after

36 h of transfection to perform Western blot according to the standard

procedure [53]. Mouse anti-HA (901501 Biolegend, 1:2000) and mouse anti-

GAPDH (60004-1-lg Proteintech, 1:2000) antibodies were used for detection.

Detections were performed with goat anti-mouse secondary antibody (31430

Thermo 1:5000) and enhanced chemiluminescence (ECL plus) reagent.

RNAseq analyses

To determine the gene expression profiles of *lg*, the available raw

transcriptomic data of different tissues in different fishes were extracted from

the SRA database (Table S2). The raw reads were mapped to the corresponding

transcriptome retrieved from Ensembl using Hisat2-2.1.0 [54] with default

parameters. The reported SAM results were converted into BAM format, and

finally recorded as count files using SAMtools v1.10 [55] followed by TPM value

calculation [56].

To identify the affected genes after knockout, comparative transcriptomic

analyses were conducted between MUT and WT (Table S3). RNA extraction

was performed (each replicate contained 30 pooled individuals) using TRIzol®

reagent (Invitrogen). Six biological replicates of WT and MUT of the F2

generation were used for RNAseq analyses. Library construction and

sequencing were performed using Illumina TrueSeq RNAseq technology at the

Novogene Company, China, with a paired end read length of 350 bp. The raw

reads were mapped to the zebrafish transcriptome assembly using the same

methods mentioned above to obtain the read count files. The differentially

expressed (DE) transcripts between WT and MUT were further analysed using

the edgeR v3.20.7 package (FDR<0.01). The raw data were deposited at NCBI

BioProject ID PRJNA611476. The top 10 up- and downregulated DE transcripts

lists can be found in Table S3.

***In situ* hybridization**

The gene fragments in the coding region of *lg* in zebrafish and tilapia
were amplified using PrimerSTAR (Takara) (Table S6) and *lyz* probe synthesis
was conducted as described in the reference [57]. The amplified products were
cloned into the pGEM®-T Easy vector (Promega). Plasmids were extracted and
purified using a TIANprep Mini Plasmid Kit II (Tiangen). The insertion and
direction of probes were confirmed with Sanger sequencing using M13 PCR
products (Tsingke). RNA probes were synthesized with the DIG RNA labelling
kit (SP6/T7) (Roche). The experiment was conducted as described in the
reference [58] with biological replicates (n=30 for zebrafish, n=20 for tilapia, and
n=10 for *P. nyererei*). The embryos or larvae were mounted in 1% low melting
point agarose (GBCBIO) and imaged with a Zeiss Axio Imager Z2 microscope.
422
Transgenic plasmids and transgenic lines construction
To determine the regulatory functions of cichlid fish-specific upstream
regions, we constructed plasmids with different upstream regions as promoters
to drive eGFP expression in zebrafish (Fig. 4A and Fig. S2), i.e., 1) species in
the MH lineage (*P. nyererei*) (Pn line); and 2) the corresponding region of
species in the sister lineage of East African Great Lakes cichlid fishes (tilapia)
(On line). 3) To determine whether the three MH-segregated SNPs in the cichlid
fish-specific upstream region are important, we specifically deleted these SNPs
using site-directed mutagenesis by PCR primer extension (Table S6) [59] to
construct another transgenic line (Pn-noSNPs line). A schematic of the plasmid
construction can be found in Fig. S2.
434
DNA fragments for different plasmid constructs were amplified with
PCR primers supplemented with the corresponding restriction endonuclease
sites (Fig. S2, Table S6). Fragments were then recovered (Tiangen), digested
(NEB) and ligated (Promega) into the pT2AL vector following the instructions
accordingly. Successfully constructed recombinant plasmids were further

confirmed by Sanger sequencing (Tsingke). Injections were performed into the

one-cell stage of zebrafish embryos. By outcrossing with WT zebrafish, we

successfully obtained F2 stable transgenic lines for all these lines (Pn line, On

line and Pn-noSNPs line). The fluoscence signals were imaged with a Zeiss

880 confocal microscope.

Immunohistochemistry (IHC)

The following primary antibodies were used: mouse anti-Pvalb7, 1:1000

provided by Prof. Masahiko Hibi, Nagoya University, Japan [60]; rabbit anti-

serotonin, 1:400 (s5545, Sigma); rabbit anti-calretinin, 1:400 (ab702, Abcam);

goat anti-GFP, 1:400 (ab6673, Abcam); mouse anti-HuC/HuD, 1:1000 (16A11,

Invitrogen); rabbit anti-SOX10, 1:400 (ab229331, Abcam); rabbit anti-PAX6

1:400 (ab209836, Abcam); mouse anti-collagen type II (II-II6B3, DSHB); and

rabbit anti-cathepsin L/MEP (ab209708, Abcam). The secondary antibodies

were donkey to goat IgG H&L (Alexa Fluor® 488) (ab150133, Abcam),

donkey to mouse IgG H&L (Alexa Fluor® 647) (ab150107, Abcam), donkey to

rabbit IgG H&L (Alexa Fluor® 647) (ab150063, Abcam) and DAPI (ab228549,

Abcam). Fish larvae were fixed in 4% paraformaldehyde (Sigma) for 4 h at

room temperature, followed by a detailed protocol as described previously [61].

The fluoscence signals were imaged with a Zeiss 880 confocal microscope.

Immune response detection

To determine whether there are different immune responses between

MUT and WT zebrafish, we injected lipopolysaccharide (LPS), the key

component of Gram-negative bacteria, into the circulation of 2 dpf zebrafish

and counted the numbers of lyz^+ neutrophils in the caudal haematopoietic tissue

at one day post injection. The lyz^+ whole-mount RNA *in situ* hybridization

signals across eight somites of the larval CHT (caudal haematopoietic tissue)

regions were manually counted. Statistical analyses can be found in Table S7.

Data Availability

472   The publicly available datasets that this study used are provided in the supplementary information. The RNAseq data of WT and MUT are available at NCBI (BioProject ID PRJNA611476). The ion torrent sequencing reads of the *lg* gene are available at NCBI (BioProject ID PRJNA611690). The scaffolds possessing *lg* of teleost fishes across the phylogeny, the assembly of *lg* including the upstream region of cichlid fishes across the phylogeny retrieved from ion torrent sequencing and Sanger sequencing can be found in Dryad https://doi.org/10.5061/dryad.fxpnvx0qb. Source associated data of Fig. 3 and Fig. 4 are provided in the supplementary files. Other data are available upon reasonable request.

References

1. Dobzhansky, T. Nothing in biology makes sense except in the light of evolution. *Am. Biol. Teach.* **March,** 125–129 (1973).

2. Tautz, D. & Domazet-Lošo, T. The evolutionary origin of orphan genes. *Nat. Rev. Genet.* **12,** 692–702 (2011).

3. Bae, B.-I., Jayaraman, D. & Walsh, C. A. Genetic changes shaping the human brain. *Dev. Cell* **32,** 423–34 (2015).

4. Zhang, Y. E. & Long, M. New genes contribute to genetic and phenotypic novelties in human evolution. *Curr. Opin. Genet. Dev.* **29,** 90–6 (2014).

5. Babonis, L. S., Martindale, M. Q. & Ryan, J. F. Do novel genes drive morphological novelty? An investigation of the nematosomes in the sea anemone Nematostella vectensis. *BMC Evol. Biol.* **16,** 114 (2016).

6. Santos, M. E., Le Bouquin, A., Crumière, A. J. J. & Khila, A. Taxon-restricted genes at the origin of a novel trait allowing access to a new environment. *Science (80-.).* **358,** 386–390 (2017).

7. Khalturin, K. *et al.* A novel gene family controls species-specific morphological traits in Hydra. *PLoS Biol.* **6,** e278 (2008).

8. Garcia-Fernàndez, J. The genesis and evolution of homeobox gene clusters. *Nat. Rev. Genet.* **6,** 881–92 (2005).

9. Wu, M., Chen, G. & Li, Y.-P. TGF-β and BMP signaling in osteoblast, skeletal development, and bone formation, homeostasis and disease. *Bone*

Res. **4,** 16009 (2016).

10.   Wagner, G. P. & Lynch, V. J. The gene regulatory logic of transcription
factor evolution. *Trends Ecol. Evol.* **23,** 377–85 (2008).

11.   Wray, G. A. The evolutionary significance of cis-regulatory mutations.
Nat. Rev. Genet. **8,** 206–216 (2007).

12.   Zhang, J. Parallel adaptive origins of digestive RNases in Asian and
African leaf monkeys. *Nat. Genet.* **38,** 819–23 (2006).

13.   Hoekstra, H. E., Hirschmann, R. J., Bundey, R. A., Insel, P. A. &
Crossland, J. P. A single amino acid mutation contributes to adaptive
beach mouse color pattern. *Science* **313,** 101–4 (2006).

14.   Chen, L., DeVries, A. L. & Cheng, C.-H. C. Evolution of antifreeze
glycoprotein gene from a trypsinogen gene in Antarctic notothenioid fish.
Proc. Natl. Acad. Sci. **94,** 3811–3816 (1997).

15.   Carroll, S. B. Evolution at Two Levels: On Genes and Form. *PLoS Biol.* **3,**
e245 (2005).

16.   Hu, W., Chen, S., Zhang, R. & Lin, Y. Single nucleotide polymorphisms
in the upstream regulatory region alter the expression of myostatin. *Vitr.*
Cell. Dev. Biol. - Anim. **49,** 417–423 (2013).

17.   Marceca, C., Pfob, M., Schelling, G., Steinlein, O. K. & Eggert, M. Single
nucleotide polymorphism creating a variable upstream open reading frame
regulates glucocorticoid receptor expression. *Gene* **563,** 24–28 (2015).

18.   Malmstrøm, M. *et al.* Evolution of the immune system influences
speciation rates in teleost fishes. *Nat. Genet.* **48,** 1204–1210 (2016).

19.   Salzburger, W. The interaction of sexually and naturally selected traits in
the adaptive radiations of cichlid fishes. *Mol. Ecol.* **18,** 169–85 (2009).

20.   SH, O. & LI, Z. Hematopoiesis: An Evolving Paradigm for Stem Cell
Biology. *Cell* **132,** 631–44 (2008).

21.   Kim, C.-H. *et al.* Zebrafish elav/HuC homologue as a very early neuronal
marker. *Neurosci. Lett.* **216,** 109–112 (1996).

22.   Hugo, S. E. *et al.* A monocarboxylate transporter required for hepatocyte
secretion of ketone bodies during fasting. *Genes Dev.* **26,** 282–93 (2012).

23.   Salzburger, W. Understanding explosive diversification through cichlid
fish genomics. *Nat. Rev. Genet.* (2018). doi:10.1038/s41576-018-0043-9

24.   Meyer, B. S., Matschiner, M. & Salzburger, W. A tribal level phylogeny
of Lake Tanganyika cichlid fishes based on a genomic multi-marker
approach. *Mol. Phylogenet. Evol.* **83,** 56–71 (2015).

25.   Salzburger, W., Mack, T., Verheyen, E. & Meyer, A. Out of Tanganyika:
genesis, explosive speciation, key-innovations and phylogeography of the
haplochromine cichlid fishes. *BMC Evol. Biol.* **5,** 17 (2005).

26.   Hubel, P. *et al.* A protein-interaction network of interferon-stimulated
genes extends the innate immune system landscape. *Nat. Immunol.* **20,**
493–502 (2019).

27.   Denève, S., Alemi, A. & Bourdoukan, R. The Brain as an Efficient and

Robust Adaptive Learner. *Neuron* **94,** 969–977 (2017).

28. Zhang, W., Landback, P., Gschwend, A. R., Shen, B. & Long, M. New
genes drive the evolution of gene interaction networks in the human and
mouse genomes. *Genome Biol.* **16,** 202 (2015).

29. Nagasawa, T. *et al.* Evolutionary Changes in the Developmental Origin of
Hatching Gland Cells in Basal Ray-Finned Fishes. *Zoolog. Sci.* **33,** 272
(2016).

30. Conith, A. J., Kidd, M. R., Kocher, T. D. & Albertson, R. C.
Ecomorphological divergence and habitat lability in the context of robust
patterns of modularity in the cichlid feeding apparatus. *BMC Evol. Biol.*
20, 95 (2020).

31. Roberts, R. B., Hu, Y., Albertson, R. C. & Kocher, T. D. Craniofacial
divergence and ongoing adaptation via the hedgehog pathway. *Proc. Natl.*
Acad. Sci. U. S. A. **108,** 13194–9 (2011).

32. Albertson, R. C., Streelman, J. T., Kocher, T. D. & Yelick, P. C.
Integration and evolution of the cichlid mandible: the molecular basis of
alternate feeding strategies. *Proc. Natl. Acad. Sci. U. S. A.* **102,** 16287–92
(2005).

33. König, D. *et al.* Distribution and Restoration of Serotonin-
Immunoreactive Paraneuronal Cells During Caudal Fin Regeneration in
Zebrafish. *Front. Mol. Neurosci.* **12,** 227 (2019).

34. Coccimiglio, M. L. & Jonz, M. G. Serotonergic neuroepithelial cells of the
skin in developing zebrafish: morphology, innervation and oxygen-
sensitive properties. *J. Exp. Biol.* **215,** 3881–3894 (2012).

35. Kermen, F., Franco, L. M., Wyatt, C. & Yaksi, E. Neural circuits
mediating olfactory-driven behavior in fish. *Front. Neural Circuits* **7,** 62
(2013).

36. Green, S. A., Simoes-Costa, M. & Bronner, M. E. Evolution of vertebrates
as viewed from the crest. *Nature* **520,** 474–82 (2015).

37. Salzburger, W. Understanding explosive diversification through cichlid
fish genomics. *Nat. Rev. Genet.* **19,** 705–717 (2018).

38. Nyman, C., Fischer, S., Aubin-Horth, N. & Taborsky, B. Evolutionary
conserved neural signature of early life stress affects animal social
competence. *Proceedings. Biol. Sci.* **285,** (2018).

39. Culbert, B. M., Gilmour, K. M. & Balshine, S. Stress axis regulation
during social ascension in a group-living cichlid fish. *Horm. Behav.* **103,**
121–128 (2018).

40. Stanke, M. & Morgenstern, B. AUGUSTUS: a web server for gene
prediction in eukaryotes that allows user-defined constraints. *Nucleic*
Acids Res. **33,** W465–7 (2005).

41. Stanke, M., Steinkamp, R., Waack, S. & Morgenstern, B. AUGUSTUS: a
web server for gene finding in eukaryotes. *Nucleic Acids Res.* **32,** W309–
12 (2004).

42. Kearse, M. *et al.* Geneious Basic: an integrated and extendable desktop
software platform for the organization and analysis of sequence data.
Bioinformatics **28,** 1647–9 (2012).
43. Lalitha, S. Primer Premier 5. *Biotech Software&Internet Rep.* **1,** (2000).
44. Afgan, E. *et al.* The Galaxy platform for accessible, reproducible and
collaborative biomedical analyses: 2018 update. *Nucleic Acids Res.* **46,**
W537–W544 (2018).
45. Mayor, C. *et al.* VISTA : visualizing global DNA sequence alignments of
arbitrary length. *Bioinformatics* **16,** 1046–1047 (2000).
46. Frazer, K. A., Pachter, L., Poliakov, A., Rubin, E. M. & Dubchak, I.
VISTA: computational tools for comparative genomics. *Nucleic Acids
Res.* **32,** W273–9 (2004).
47. Brudno, M. *et al.* LAGAN and Multi-LAGAN: efficient tools for large-
scale multiple alignment of genomic DNA. *Genome Res.* **13,** 721–31
(2003).
48. Quandt, K., Frech, K., Karas, H., Wingender, E. & Werner, T. MatInd and
MatInspector: new fast and versatile tools for detection of consensus
matches in nucleotide sequence data. *Nucleic Acids Res.* **23,** 4878–84
(1995).
49. Cartharius, K. *et al.* MatInspector and beyond: promoter analysis based on
transcription factor binding sites. *Bioinformatics* **21,** 2933–2942 (2005).
50. Yang, Z. PAML: a program package for phylogenetic analysis by
maximum likelihood. *Comput. Appl. Biosci.* **13,** 555–6 (1997).
51. Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol.
Biol. Evol.* **24,** 1586–91 (2007).
52. Concordet, J.-P. & Haeussler, M. CRISPOR: intuitive guide selection for
CRISPR/Cas9 genome editing experiments and screens. *Nucleic Acids
Res.* **46,** W242–W245 (2018).
53. Zhao, F. *et al.* Irf8 regulates the progression of myeloproliferative
neoplasm-like syndrome via Mertk signaling in zebrafish. *Leukemia* **32,**
149–158 (2018).
54. Kim, D., Langmead, B. & Salzberg, S. L. HISAT: a fast spliced aligner
with low memory requirements. *Nat. Methods* **12,** 357–360 (2015).
55. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools.
Bioinformatics **25,** 2078–9 (2009).
56. Wagner, G. P., Kin, K. & Lynch, V. J. Measurement of mRNA abundance
using RNA-seq data: RPKM measure is inconsistent among samples.
Theory Biosci. **131,** 281–285 (2012).
57. Hou, Y. *et al.* Systemic inoculation of Escherichia coli causes emergency
myelopoiesis in zebrafish larval caudal hematopoietic tissue. *Sci. Rep.* **6,**
36853 (2016).
58. Westerfield, M. *The zebrafish book. A guide for the laboratory use of
zebrafish (Danio rerio). 4th edition.* (University of Oregon Press, Eugene,

2000).
59. Reikofski, J. & Tao, B. Y. Polymerase chain reaction (PCR) techniques
for site-directed mutagenesis. *Biotechnol. Adv.* **10,** 535–47 (1992).
60. Nimura, T. *et al.* Role of Reelin in cell positioning in the cerebellum and
the cerebellum-like structure in zebrafish. *Dev. Biol.* **455,** 393–408 (2019).
61. Barresi, M. J. F., Stickney, H. L. & Devoto, S. H. The zebrafish slow-
muscle-omitted gene product is required for Hedgehog signal transduction
and the development of slow muscle identity. *Development* **127,** 2189–
2199 (2000).

Acknowledgements

Many thanks to Prof. Walter Salzburger, Attila Rüegg, Zhen Liu, Zhiqiang Zhou and Hesheng Xiao for the sampling and suggestions. We thank Prof. Masahiko Hibi and Prof. Kiyoshi Naruse for providing the antibody Pvalb7. We also thank Prof. Lingfei Luo, Prof. Hua Ruan, Prof. Yang Liu and Prof. Tienming Lee for the laboratory supports. Many thanks to Prof. Zuzana Musilova, Dr. Xiaogui Yi, Dr. Yandong Zhan, Dr. Fangying Zhao, Yuepeng He, Xinkai Cheng, Qing Li and Yue Gao for the supports. Thanks to Huansheng Tao and Jialing Xu for the Confocol microscope support, Zhao Long for fluroscent microscope support. This study was funded by grants from the National Natural Science Foundation of China (31802283), Guangdong Basic and Applied Basic Research Foundation (2020A1515010330); the Fundamental Research Funds for the Central Universities (20lgpy109); China Postdoctoral Science Foundation (2018 M633311) and Doctoral Scholarship of University of Basel, Switzerland to LG; Grants from the National Natural Science Foundation of China (31630082) to DW, and (31771623 and 31571500) to LL. Part of -omics analyses was supported by National Supercomputer Center in Guangzhou, China. The English language of this article was edited by Elsevier Language Editing Services.

Authors contributions

LG, DW and LL conceived this work. LG carried out –omics sequencing

and evolutionary analyses. LG, CL, XM and ZW conducted the *in situ*

hybridization. LG carried out the IHC. XH and WZ participated the IHC

experiments of serotonin-positive cells and olfactory epithelium cells. LG, CL

and XM constructed plasmids of the Pn line and the On line. LG constructed

plasmids of the Pn-noSNPs line. LG leads the transgenic lines and mutant lines

construction with the help from CL and XM. LG took the confocal photos. YH

performed the Western blot experiment. ZW carried out the LPS immune

response experiments. LG conducted the fasting ORO staining experiments. LG

wrote the initial manuscript draft, and all authors commented on the manuscript

and approved the final version.

Competing interests: The authors declare no competing interests.

Figure legends

Fig. 1 Sequence evolution of *lg*. a. Comparative genomic analyses showed that

lg is a fish-specific gene. b. Multiple losses occurred in different lineages within

teleost fishes. The reference tree is from [18].

Fig. 2 Expression profiles of *lg* in fishes. a. Transcriptomic analyses of *lg* gene

expression patterns in adult fish tissues. *lg* was mainly expressed in the brain

and eyes in most fishes but evolved new expression patterns in the skin in the

modern haplochromine (MH) lineage of cichlid fishes. b. *In situ* hybridization

in zebrafish and cichlid fish larvae. *lg* was mainly expressed in the brain,

ganglion and inner plexiform cell layers in the eyes of zebrafish. No expression

was detected in the hatching gland of 24 hpf (hours post fertilization) zebrafish

embryos. *lg* was highly expressed in the brain, eyes and pharyngeal arches in

the MH cichlid fish, *Pundamilia nyererei*, but weakly expressed in the cichlid

fish *Oreochromis niloticus*.

Fig. 3 *lg* gene knockout in zebrafish and comparisons between mutant

(MUT) and wild-type (WT) zebrafish. A. *lg* is successfully disrupted using

CRISPR/Cas9. This was confirmed by both Sanger sequencing (top) and

Western blot (bottom). B. Immune response comparisons between MUT and

WT using whole mount *in situ* hybridization in CHT (caudal haematopoietic

tissue) treated with LPS and PBS (n=10 for WT, and n=18 for MUT),

respectively. Quantification of the numbers of lyz^+ signals across eight somites

is shown as dots. n.s. not significant. Mean and SD error bars were shown. C.

Phenotype comparisons between MUT and WT zebrafish. Brain and eye

phenotypes in mutants were examined with the neuron marker HuC. No

obvious phenotype was found. Under fasting condition, no obvious mutant

phenotype was detected with ORO staining using the same strategy as described

in the literature [22].

Fig. 4 Sequence evolution of *lg* across East African cichlid fish phylogeny

and regulatory function evolution in cichlid fishes. Aa. Both the upstream

and coding regions exhibited SNPs segregated with the modern haplochromine

(MH) lineage, i.e., three SNPs (GTA) in the upstream region and one

nonsynonymous amino acid change (A to S) located at the 38[th] position was

under positive selection (Table S4). Transcription factors binding sites

prediction showed that these SNPs can set up the opportunity to bind to

glucocorticoid modulatory element binding protein 2 (GMEB2). The reference

tree is from literature [24]. Ab. Three types of transgenic plasmids construction.

The pT2AL-On-eGFP line (On line) adopted the upstream region of

Oreochromis niloticus as the promoter to drive eGFP expression. The pT2AL-

Pn-eGFP line (Pn line) used the homologous upstream region of *Pundamilia*

nyererei in the MH lineage as the promoter. The pT2AL-Pn-no-SNPs-eGFP line

(Pn-noSNPs line) used the same promoter as the Pn line but deleted the three

MH-segregated SNPs (GTA). B. The fluorescence images in the Pn line.

Merged figures of double staining of eGFP and cell marker antibodies are

shown if there are available antibodies. Red is from the cell marker antibody.

Green is from eGFP. Blue is from DAPI. Pictures were taken from 4 dpf (days

post fertilization) zebrafish except the hatching gland which is from 24 hpf

(hours post fertilization) embryos. C. Different expression patterns of three

transgenic lines in zebrafish. D. Statistics of expression pattern comparisons

among the three transgenic lines, focusing on the cerebellum, hindbrain and the

first two pharyngeal arches. The Pn line showed significantly stronger

fluroscence than the other two lines. With the three MH-segregated SNP

deletions, the Pn-noSNPs line showed weak fluroscence expressions, similar to

the On line. Mean and SD error bars were shown. n=6 for each measurement.

Detailed statistical analyses can be found in Table S5.

Fig. 5 Hypothesis of *lg* sequence and function evolution with respect to evo-

devo. *lg* is a fish-specific gene. Its ancestral function is not indispensable. In

cichlid fishes, *lg* evolves a cichlid fish-specific regulatory region, thus gained

new regulatory functions to drive gene expression in cells related to adaptation,

such as in pharyngeal cartilages and the hatching gland. Especially in the most

species-richness lineage, the modern haplochromine (MH) lineage, this

upstream can not only drive strong eGFP expression patterns, but also further

evolved new profiles in epidermal serotonin-positive cells. This can be simply

due to the three MH-segregated SNPs in the upstream region that are predicted

to bind to a hormone related transcription factor, the glucocorticoid modulatory

element binding protein 2 (GMEB2). If this is true, it opens a new opportunity

to response to the external environemnt. In addition, the coding region of *lg* also

evolved one MH- segregated amino acid mutation that is under positive

selection, indicating that new function might also evolved in the coding region
which deserves further investigation. *lg* thus provides a good example of new
gene function emergence in a redundant lineage-specific gene with respect to
evo-devo.
757
758
759
760 **Figures**
761

763 **Fig. 1 Sequence evolution of *lg*.**

765 **Fig. 2 Expression profiles of *lg* in fishes.**

Fig. 3 *lg* **gene knockout in zebrafish and comparisons between mutant (MUT) and wild-type (WT) zebrafish.**

Fig. 4 Sequence evolution of *lg* across East African cichlid fish phylogeny and regulatory function evolution in cichlid fishes.

773 **Fig. 5 Hypothesis of *lg* sequence and function evolution with respect to evo-**

774 **devo.**

776 **Supplementary information**

787   Table S7 The numbers of LPS immune response signal comparisons between

788   WT and MUT zebrafish

789   Supplementary Figures