Disruption of orbitofrontal-hypothalamic projections in a murine ALS model and human patients

Authors: David Bayer[1,2], Stefano Antonucci[1], Hans-Peter Müller[1], Luc Dupuis[3], Tobias M. Boeckers[4,5], Albert C. Ludolph[1,5], Jan Kassubek[1], Francesco Roselli[1,5]

E-Mail: david.bayer@uni-ulm.de, stefano.antonucci@uni-ulm.de, hans-peter.mueller@uni-ulm.de, ldupuis@neuro-cnrs.unistra.fr, tobias.boeckers@uni-ulm.de, albert.ludolph@rku.de, jan.kassubek@uni-ulm.de, francesco.roselli@uni-ulm.de

Affiliation: 1. Dept. of Neurology, Ulm University, Ulm DE

2. CEMMA (Cellular and Molecular Mechanisms in Aging) research training group, Ulm DE

3. University of Strasbourg

4. Institute of Anatomy and Cell Biology, Ulm University, Ulm DE

5. German Center for Neurodegenerative Diseases-DZNE, Ulm DE

Corresponding authors: Francesco Roselli,

Center for Biomedical Research (ZBF)

Helmholtzstrasse 8-89081 Ulm (DE)

Phone: 0049-0731-500-63147

Email: francesco.roselli@uni-ulm.de

Abstract

Increased catabolism is a new clinical manifestation of Amyotrophic Lateral Sclerosis. A dysfunction of lateral hypothalamus may drive hypermetabolism in ALS; however, Its causes and anatomical substrates are unknown. We hypothesize that disruption cortico-hypothalamic circuits may impair energy homeostasis in ALS. We used rAAV2 for large-scale projection mapping and image analysis pipeline based on Wholebrain and Ilastik to quantify projections from the forebrain to the latera hypothalamus of the SOD1(G93A) ALS mouse model as well as of the Fus$^{\Delta NLS}$ ALS mouse model. Expanded projections from agranular Insula, ventrolateral orbitofrontal and secondary motor cortex to lateral hypothalamus were found in two independent cohorts of the hypermetabolic SOD1(G93A) ALS model. The non-hypermetabolic Fus$^{\Delta NLS}$ ALS mouse model display a loss of projections from motor cortex but no change in projections from insula and orbitofronal cortex. 3T DTI-MRI data on 83 ALS patients and 65 controls confirmed the disruption of the orbitofrontal-hypothalamic tract in ALS patients. Converging murine and human data demonstrate the selective disruption of hypothalamic inputs in ALS as a factor contributing to the origin of hypermetabolism.

Key words: rAAV2; Agranular Insula; Orbitofrontal Cortex; Lateral Hypothalamus; Hypermetabolism; Amyotrophic lateral sclerosis

Introduction

Amyotrophic lateral sclerosis (ALS) is traditionally conceptualized as a neurodegenerative condition primarily affecting upper motoneurons, located in primary motor cortex, and lower motoneurons, located in spinal cord, whose dysfunction and loss determines a relentless, progressive and ultimately fatal motor impairment [1]. More recently, hypermetabolism has been recognized as an additional, non-motor clinical feature of ALS [2]. Epidemiological surveys have revealed that ALS patients display increased catabolism [2, 3] and pointed out lower body-mass index constitutes a risk factor for ALS [4]. Furthermore, reduced levels of metabolic rate proxies such as plasma lipids and body fat content are predictors of survival of ALS patients [5-7], i.e. weight loss is strongly correlated with shorter survival [8]. Increased catabolism appears to be due to intrinsic hypermetabolism both in ALS patients [9-13] as well as in mutant SOD1 ALS mice [14]. Notably, metabolism and energy balance actually constitute promising targets for intervention, since increasing caloric intake has beneficial effects on survival, particularly on fast-progressing ALS patients [15].

Mechanistic insights into ALS hypermetabolism remain limited. Hypothalamic nuclei receive and integrate inputs coming from a large fraction of the brain [16-19] in order to establish the proper balance of feeding, energy storage and energy expenditure [20]. Thus, disruption of these inputs may be sufficient to drive metabolic imbalances. In fact, the overall architecture between large-scale networks appear to be disturbed in ALS [21-25] and ALS-related pathobiochemistry affects a significant proportion of cortical and subcortical structures, in a pattern evolving with disease progression [26, 27]. We have hypothesized that a similar degree of disruption and remodeling may take place in large-scale networks providing inputs to the

feeding-regulating Lateral Hypothalamic Area (LHA) and that such a disruption may coincide

with the appearance of the hypermetabolic phenotype. We have now identified the selective

disruption of projections from insular and orbitofrontal cortex to LHA in ALS mouse model as

well as in human ALS patients by a combination of retrograde rAAV-2 tracing and MRI-DTI tract

tracing.

Methods

Animals

All experimental procedures involving animals were performed in agreement with the

guidelines for the welfare of experimental animals issued by the Federal Government of

Germany; all experiments were approved by the Regierungspräsidium Tübingen under the

animal license- number 1390 and by the Ulm University Tierforschungszentrum committee. No

effort was spared to implement "3R" guidelines for animal experimentation.

We obtained from Jackson Laboratories the following strains: B6SJL73 Tg(SOD1*G93A)1Gur/J

(high-copy), (henceforth mSOD) and B6.Cg-Gt(ROSA)26Sor$^{tm6/(CAG74ZsGreen)Hze/J}$ (henceforth

ZsGreen). To generate the mSOD/ZsGreen double transgenic mice, hemizygous mSOD males

were crossed with homozygous ZsGreen$^{+/+}$ females as previously reported [21]; the progeny

included mSOD^{+}/ZsGreen$^{+/-}$ mice at the expected mendelian rate. mSOD^{+}/ZsGreen$^{+/-}$ are

henceforth going to be labelled mSOD1 and mSOD^{-}/ZsGreen$^{+/-}$will be labelled WT.

The heterozygous B6.Fus$^{\Delta NLS/+}$ (henceforth Fus) were provided by Luc Dupuis from the Faculté

de Médecine, Strasbourg, France [28]. For the generation of Fus/ZsGreen double transgenic

mice, heterozygous Fus males were crossed with homozygous ZsGreen$^{+/+}$ females. The progeny

included Fus$^-$/ZsGreen$^{+/-}$ mice at the expected mendelian rate. Fus$^-$/ZsGreen$^{+/-}$ are henceforth

going to be labelled kiFUS.

Mice were housed at 2 - 5 animals per cage, with unlimited access to food and water, a

light/dark cycle of 12/12 hr and humidity between 40% and 60%. All mice expressing mSOD

were routinely tested for motor impairment and euthanized in case of overt motor disability.

Since male and female mice have differ substantially in progression rates of clinical and

biological manifestation of motoneuron disease (e.g. [29]), the present study focused on male

mice only.

Viral vectors

Retrograde rAAV2 [30] vectors encoding for the pmSyn1-EBFP-Cre (addgene plasmid # 51507;

kindly donated by Hongkui Zeng [31]),were obtained from Addgene.

Intracerebral injection

Intracerebral injections of rAAVs were performed as previously reported [21]. Briefly, 1µl of viral

suspension with a titer of $3x10^{12}$ vg/ml and 1 µl of 1% of Fast Green were freshly mixed and 300

nl of the mixture was loaded into a pulled-glass capillary. Pulling parameters were optimized to

obtain a long and tapering capillary tip. Mice were pre-treated with buprenorphine (0.1 mg/kg)

and meloxicam (1 mg/kg) 20 min before being anesthetized with 5% sevoflurane /95% O_2 ; mice

were then positioned into a stereotactic frame (David Kopf) and kept under continuous

anesthesia with 3% sevoflurane/97% O_2. Mice were positioned on a heated pad connected to a

closed-loop system to maintain body temperature, monitored by a rectal probe, at 37°C. Upon

incision of the scalp, a burr hole was prepared using a hand-held microdrive aiming the LHA at

AP = -1.20, ML = -1.25, DV = -4.70 (according to Paxinos atlas, 2nd edition [32]) refined for the

mouse strain under consideration of the age).

Under visual control, the pulled-glass capillary (tip closed) was moved into the drilled hole and

gently pushed down to verify that meninges could be penetrated without deflection of the

capillary. Thereafter, the pulled-glass capillary was withdrawn and opened at the very tip by

gently touching with a microscissor. Before lowering the pulled-glass capillary to the final

position inside the brain, a thin layer of $DPBS^{+Ca/+Mg}$ was applied, generating a virus-free tip by

capillary force. Since brain tissue might absorb some of the liquid out of the pulled-glass

capillary by its own tissue/capillary force, during further movement, the pulled-glass capillary

was lowered not slower than 2 mm/s. The viral suspension was injected using a Picosprizer

microfluidic device; injection was performed during 5 min. The duration of one pulse was 10 ms;

since the opening diameter of each capillary could be slightly different, the injection pressure

was adapted (between 20 and 60 psi) to ensure a constant overall flow rate of 50-60 nl/min.

After injection, the capillary was kept in place for 15 min and then removed with a continuous

movement (1 mm/s). Remaining pressure in the capillary was released before withdrawing it to

prevent any residual virus suspension to be discharged into the capillary thread.

Immunohistochemistry

Mice (mSOD) were sacrificed at P (post-natal day) 40 (injection at P25) or at P110 (injection at

P95); Fus mice were injected at P255 and sacrificed at P270.

Mice were trans-cardially perfused with PBS and then 4% PFA in PBS; after 18h of post-fixation

in 4% PFA, brains were cryoprotected in 30% Sucrose/PBS, embedded in OCT and serially

sectioned using a *Leica CM1950 cryostat* into 70 µm sections from AP +2.6 mm to -3.0 mm as

previously reported [21].

The ZsGreen reporter did not require immunodetection since the native signal was very strong.

For dedicated experiments, immunostaining for SOD1 (Anti SOD1, Sigma-Aldrich, HPA001401,

1:500), misfolded SOD1 (Misfolded SOD1 (B8H10), MédiMabs, MM-0070-p, 1:250) and Fus (Anti

Fus, Sigma-Aldrich, HPA008784, 1:300) was performed.

Briefly, sections were blocked in blocking solution (3% BSA, 0.3% Triton in PBS) for 2h at room

temperature (rt). Primary antibody was diluted in blocking solution and incubated at 4°C for

72h. Secondary antibody (Invitrogen, Donkey anti-rabbit Alexa Fluor 568, # A10042; Donkey

anti-mouse Alexa Fluor 647, # A31571; anti-guinea pig 405) was applied diluted 1:500 in

blocking solution after three washing steps (PBS 20 min at rt) and incubated for 2h at rt.

Subsequently sections were washed three times again and mounted using Gold Antifade

Mountant (Invitrogen #P36930).

Image acquisition

Glass slides of serially-sectioned brains were first subject to visual assessment using a Leica

DMIL equipped with a 2.5x/0.07 objective; brains that were injected in the wrong location or

displayed obvious macroscopic artifacts were excluded at this stage and not processed further.

The remaining set of brain sections were imaged in full using a slide-scanning microscope (Leica

DMI6000B) equipped with a 5x/0.12 objective and exposure time ranging between 300 ms (405

nm) to 800ms (647 nm). All images were saved at 15304 x 28295 resolution with 16-bit depth.

Images of the cortical areas immunostained for misfolded SOD1 were acquired using a Zeiss

LSM710 confocal laser scanning microscope equipped with a 20x air objective. In total, 15

optical sections each 1μm-thick were acquired for each region of interest.

Neuron mapping

A multiple software approach was devised to analyze brain section images and properly

parcellate and quantify projections from cortex to LHA; the pipeline is summarized in Fig. 1A

and is reported here in detail.

Anatomical annotation – Firstly, a custom *Fiji* [33] macro allowed extracting single brain sections

from whole microscopic slide images while appending serial numbers. Hands-on registration of

coronal brain sections was performed by means of *WholeBrain* package in *R* [34], laying its

foundations on the Allen Brain mouse reference Atlas (ABA version 2011 [35]). In view of

accounting for tilted sections (as compared to the original ABA coronal planes) and for biological

inter-individual brain differences, a series of easily recognizable anatomical landmarks were

chosen to appropriately map the regions of interest (ROIs) along the anterior-posterior (AP)

axis, as shown in the Supplementary Table 1. The stereotactic coordinates were determined

with *openbrainmap.org*.

Neuron segmentation – Despite the flexibility of the parameters of the built-in segmentation

functions in *WholeBrain*, a systematic underestimation of neuronal counts was noticeable in

areas with high neuronal density, such as the frontal cortex, whereas a systematic

overestimation of neuronal counts took place in areas with low neuronal density, due to the

inclusion of segments of dendrites as purported neurons. In order to avoid the warping of the

neuronal counts because of these opposing biases, a separate strategy was pursued for neuron

segmentation in single brain ROIs alone. RGB images with single region outlines were generated

as explained in https://gitter.im/tractatus/Lobby and imported in *Fiji* [33]. Another *Fiji* macro

was then designed to mask and resize the RGB image in order to have a perfect match of the

original high-resolution image, to convert the former to HSB Stack, recover the outline of the

brain ROI and use it to crop the latter.

Ilastik toolkit [36] subsequently served a double purpose. The Pixel Classification workflow

enabled to discriminate neurons (foreground) from neuron-devoid brain architecture

(background); the resulting binary images (Fig. 1A, 4.3) were fed to a final *Fiji* macro to crop

highly-resolved brain ROIs (Fig.1A, 4.2) to smaller ROIs containing neurons only (Fig.1A, 4.6) and

exclude brain ROIs without LHA-projecting *somata* from further analysis. This step proved

crucial to resort to a highly trained Cell Density Counting *Ilastik* suite, since brain architecture

removal reduced the chance of generating false positives and drastically lessened the burden on

the RAM.

Ilastik training

For both pixel classification and cell density counting workflows, σ values of 1.0, 1.6, 3.5 and 5.0

were selected for the following features: Gaussian Smoothing, Laplacian of Gaussian, Gaussian

Gradient Magnitude, Difference of Gaussian, Structure Tensor Eigenvalues, Hessian of Gaussian

Eigenvalues. Twenty randomly selected whole-section images were sufficient for Pixel

Classification training; batch processing generated simple-segmentation.tiff (Figure1, A, 4.3)

images.

On the other hand, the Cell Density Counting classifier required a total of 250 randomly selected

images for training. As displayed in figure 1B, after annotating 100 images the training effect

had already reached saturation. Since accuracy of less than 90% seemed not satisfying enough,

additional 150 images were annotated to tweak out a few more percentages in accuracy.

Furthermore, we adapted the parameters of the implemented *Random Regression Forest*

algorithm (described in more detail in [37]), namely number and maximum depth of the trees

(Ntrees = 20, MaxDepth = 80), achieving a final accuracy of 94% by comparing counts to a

selection of 100 randomly selected and manually counted images. Random sampling of

probability maps exported upon batch image processing provided a further quality control

(Fig.1A, 5.2).

Determination of single brain region and injection site volumes

Volume measurements were performed with yet another custom macro in *Fiji*, measuring the

area of individual scaled cropped.tiff images (Figure1, A, 4.2) and multiplying by the thickness of

a single section (70 µm). Injection site volumes were determined by manually defining their

cross-section in every brain slice at a set image contrast and once again multiplying such areas

for the section thickness. Brains with injection sites bigger than 1 mm^3, mislocalized position or

displaying a virus backflow (visible in dorsal regions with respect to the LHA) bigger than 10% of

the whole injection site itself were excluded from further analysis.

Determination of misfSOD1 burden

A dedicated macro in Fiji was used to create a selection of misfSOD positive cells by applying the

RenyiEntropy [38] thresholding and subsequent background subtraction (rolling ball radius = 20

px) and gaussian blurring ($\sigma = 1$). Afterwards, the area was measured by restoring the selection

in the original image

MRI scanning of DTI data in ALS patients

DTI scanning study included 83 ALS patients (58 ± 14 years, 49m, ALS-FRS-R 40 ± 6, disease

duration 19 ± 16 months) and 64 controls (52 ± 11 years, 33m) according to a standardized

protocol (for details see [39]). DTI data were acquired on a 3.0 T head scanner (Allegra, Siemens

Medical, Erlangen, Germany). The standardized DTI scanning protocol was as follows: 49

gradient directions (b = 1000 s/mm²) including one b = 0 gradient directions, 52 slices, 96 x 128

voxels in-plane, slice thickness 2.2 mm, in-plane voxel size 2.2 mm x 2.2 mm, echo time 85 ms,

repetition time 7600 ms. All participating patients and controls provided written informed

consent for the study according to institutional guidelines. The study was approved by the

Ethical Committee of the University of Ulm (reference #19/12).

DTI analysis

The DTI analysis software Tensor Imaging and Fiber Tracking (TIFT [40]) was used for post

processing and statistical analysis as previously described [39]. In brief, stereotaxic

normalization to the Montreal Neurological Institute (MNI) space was performed iteratively

using study-specific templates. To map white matter microstructure, fractional anisotropy (FA)

maps were calculated from the stereotaxically normalized DTI data sets of all subjects. A

Gaussian filter of 8 mm full width at half maximum was applied for smoothing of FA maps for a

good balance between sensitivity and specificity [41] and FA maps were corrected for the

covariate age. Tractwise fractional anisotropy statistics (TFAS) was performed by statistically

comparing the FA values between the two subject groups in a given tract system [42]. The

following tract systems were focussed on: a tract from orbitofrontal regions to the

hypothalamus (orbitofrontal-hypothalamic tract), a tract from the hypothalamus to the insula

(insular-hypothalamic tract), the cingulate-hypothalamic tract, and the corticospinal tract as a

reference. Consequently, a tract-of-interest (TOI) analysis allowed for quantification of

microstructural alterations in these tract systems.

Statistical analysis

The statistical analysis was performed with the *GraphPad8* software suite. Comparison of the

neuronal counts (absolute counts or normalized) was performed by two-way ANOVA (genotype

x structure design) using Sidak´s post-hoc multiple comparisons correction; corrected p values

are provided. Normalization was obtained dividing the absolute number of neurons in each

structure for the total number of neurons counted in that brain and multiplying the ratio for

50,000. Comparison of total neuronal counts was performed by two-tailed unpaired Student´s t-

test. Comparison of misfSOD1 burden was performed by one-way ANOVA with Tuckey´s post-

hoc test.

Results

Enhanced semi-automated mouse brain segmentation with neuronal annotation for global quantification of projections with single-cell resolution.

In order to build a quantitative and anatomically accurate map of projections from cortical and subcortical areas to the lateral hypothalamus, first we set out to establish a reliable approach to identify neurons and register their position within the structure classification of the Allen Brain Anatomical reference Atlas (Fig. 1A). The image batch for software training was obtained by injecting the retrograde rAAV2 (rAAV2-retro) into the LHA of two ZsGreen reporter mice. The latter were sacrificed 15 days later and the fixed brain was serially sectioned in 70 µm-thick sections; the whole glass slide with mounted sections was scanned using a Leica epifluorescence microscope. Each atlas plate was warped onto the brain section by defining an appropriate number of corresponding anatomical landmarks with *WholeBrain* (see the Methods section), thereby matching the atlas coordinates with the actual brain sections and allowing a proper anatomical annotation (Fig. 1A, 2). Single structure outlines (cortical layers, subcortical structures, e.g. fig. 1A, 4.1) were extracted using a dedicated *R* script, size-corrected and re-overlapped onto the original image in *Fiji*; the corresponding area of the original image was then cropped (Fig. 1A, 4.2), resulting in the unpacking of the original single-section into several hundreds cropped pictures (each image was coded to be later identified), resulting in >100,000 pictures per brain. In order to identify and count the number of neurons in each fragment, *Ilastik* software suite was employed. Simple foreground/background segmentations were generated by resorting to the pixel classification workflow. This allowed to automatically clear

most of the brain autofluorescence background in *Fiji* (Fig. 1A, 4.6). This step enabled the

improvement of the discrimination of few ZsGreen+ neurons in ROI containing many more

dendrite stretches and artifacts, and also reduced the computational burden placed on *Ilastik*

cell density classifier and therefore minimizing the chance of generating artifactual counts.

Ilastik cell density classifier was trained with 250 randomly selected ROI images that were

annotated by a human operator; the accuracy of software reached 86% after 100 images and

achieved a final 94% peak (Fig. 1B) by improving the performance of the *Random Regression*

Forest algorithm with 150 additional images and tuning its parameters (trees number and

depth; see Methods). The whole procedure was then sequentially implemented so that

neuronal counts for each picture from every section were annotated and logged together with

the anatomical location within the hemisphere and the brain section as well as with the rostro-

caudal serial number of the section itself.

Figure 1: Pipeline for the quantitation of forebrain projections to LHA. A: Neurons projecting to LHA are identified by injecting rAAV-retro encoding Cre in the LHA of WT (or mSOD1) carrying a floxed ZsGreen reporter allele (panel 1). Brains are serially sectioned (2) and each section is manually registered in *WholeBrain* (3). To obtain a precise quantification of the neurons in each anatomical structure, registered sections are parcellated and background-corrected in *ImageJ* (panel 4) and the parcellated images are segmented using a purpose-trained *Ilastik* (density counting workflow) (5) to obtain the final neuronal counts per image. The *WholeBrain* dataset is also used to derive an independent anatomical annotation in the Allen Brain Atlas (backward warp transform field (atlas plate to section) 6.1, forward warp transformation field (section to atlas plate) 6.2) and 3D reconstructions (6.3). B: Training of *Ilastik* cell density counting. Accuracy improvement of the classifier according to the amount of images used for training. The increasing need of computing power is indicated by the time needed to process 60 Images. Accuracy was calculated by comparing randomly selected and manually counted images. Using a total of 250 images and adapting the parameters for the *Random Regression Forest*, a final accuracy of 94% was achieved.

Disrupted cortico-hypothalamic projections in symptomatic mSOD1 ALS mice

We exploited our enhanced registration-segmentation-annotation pipeline to establish whether

the large-scale architecture of the projections to LHA was altered in the mSOD1 mice at a

symptomatic stage (i.e., when body weight loss has appeared and clinical score is 2). To this aim,

20 mSOD1/ZsGreen+ and 20 WT/ZsGreen+ mice were injected at the age of P95 with rAAV2-

retro [30] encoding for Cre under the human synapsin promoter in the Lateral hypothalamic

area (LHA; Fig. 2A). All neurons projecting to the site of injection would take up the rAAV2-retro,

express the Cre recombinase and appear ZsGreen+. Mice were sacrificed 15 days later. Upon

serial sectioning, a series of quality control criteria were applied and brains displaying i)

improper location of the injection site ii) injection volume larger than the boundaries of the LHA

iii) large number of neurons infected along the thread of the injection capillary iv) low (<25,000)

overall number of ZsGreen+ neurons were excluded from further analysis. Six brains for each

genotype were then considered for further quantification. These brains were randomly divided

in two cohorts: cohort 1 was subjected to *WholeBrain* parcellation and cohort 2 was left for

confirmation (see below). Once cohort 1 brains were processed through the *WholeBrain*-

registration and annotation pipeline, we identified an average of about 50,000 neurons

projecting to the LHA in WT mice; when the brainstem was excluded (because anatomical

annotation of brainstem nuclei was judged unreliable based on visual inspection of the

WholeBrain-overlapped images), we identified 84 cerebral areas projecting to the LHA (full list

with absolute and normalized neuronal counts is provided in the Supplementary Table 2).

Projections were identified from a large fraction of the forebrain, with the largest component

provided by limbic areas such as Anterior Cingulate Area (ACA), Prelimbic Cortex (PL),

InfraLimbic Area (ILA), Agranular Insula (AI), as well as from subcortical structures belonging to

the limbic system such as Basolateral and Basomedial Amygdala (BLA and BMA, respectively).

Intriguingly, substantial projections were identified from motor areas (primary and secondary),

from primary sensory areas (gustatory, auditory, olfactory/piriform, visual cortex) and from

basal ganglia (Dorsal Putamen, DP), underscoring the breath of integration taking place in LHA.

The same structures appeared to project to LHA from ipsilateral and contralateral hemispheres,

although the contralateral contribution appeared to be substantially smaller (approx. 40,000

neurons from the ipsilateral hemisphere vs 10,000 from the contralateral in WT animals).

In order to obtain a pertinent comparison (i.e., excluding areas with counts <100 neurons) of

projections in WT and mSOD1 mice, we contrasted the absolute neuronal counts in 28 areas

(accounting for approx. 50% of the total projections, maintaining ipsi and contralateral dataset

distinct; PL and orbitomedial cortex as well as lateral/ventrolateral orbitofrontal cortex -ORBl/vl-

and AI were grouped together due to the uncertainty in establishing their borders and in

agreement with functional similarity. While the overall pattern of projections to LHA in

symptomatic mSOD1 mice was comparable to the WT ones, the cumulative absolute counts of

ZsGreen+ neurons for the 28 structures were significantly increasedin the mSOD cohort

considering the absolute counts (p=0.02; Supplementary Fig. 1-A). We contrasted the absolute

counts for each of the 28 areas in WT and mSOD1 mice (Supplementary Fig. 1- 1B). Two-way

ANOVA revealed a significant effect of genotype ($F_{1,112}=105.70$, p<0.0001) and post-hoc analysis

(Sidak´s multiple comparisons test) revealed significant expansion of the projections from ACA

(p<0.0001), ILA (p=0.009), ORBl/vl+AI (p<0.0001), Tenia Tecta (TT; p=0.0365), secondary motor

cortex (MOs, p<0.0001) and piriform cortex (PIR; p=0.0005). In order to compensate for the

variable total number of infected neurons in each individual brain, we then normalized the

absolute counts of neurons projecting to LHA to a nominal of 50,000 neurons, so to compute

the relative contribution of each area to the total input to LHA. Upon normalization, the 28

areas in object amounted for a comparable total number of neurons (Fig. 2B). Two-way ANOVA

on normalized data revealed once again a significant effect of genotype ($F_{1,112}$= 7.31, p=0.0079)

and post-hoc analysis revealed a significant increase in projections from ORBl/vl+AI (p=0.0012)

and from MOs (p=0.0006; Fig. 2C; also visualized in the 2D projections, Supplementary Fig. 1-C-

D as well as in the 3D Wholebrain reconstruction, Fig. 2D). The remaining 26 structures

displayed only non-significant trends toward expanded projections (PIR: p=0.21; ILA: p=0.32) or

no difference in the size of the neuronal population projection to LHA (Fig. 2C). When the

projections from the contralateral hemisphere were contrasted, we detected an effect of

genotype in the absolute counts (two-way ANOVA $F_{1,112}$= 31.49 ; p<0.0001) that was traced in

post-hoc test (Sydak´s) to a significant expansion of projections from ACA (p=0.0008), ILA and

PL+ORBm (both p<0.0001) (Supplementary Fig. 2- 2A-B). However, when the absolute counts

were normalized to account for the total number of ZsGreen$^+$ neurons in each brain, no

significant differences were identified between the genotypes (two-way ANOVA, $F_{1,112}$= 0.31,

p=0.57; Supplementary Fig. 2-C).

Since mice with neurodegenerative phenotypes may display cortical atrophy [43], we used the

WholeBrain registration to measure the volume of each cortical structure and compensate for

eventual volume loss. Indeed, we identified a significant loss of volume in mSOD1 mice (two-

way ANOVA $F_{1,112}$=28.80, p<0.0001) which was traced by post-hoc test (Sidak´s.) to the atrophy

of MOp (p=0.008) and SS (p<0.0001) with only a trend seen for MOs (p=0.33) (Supplementary

Fig. 3). Notably, ORBl/vl+AI and MOs did not display significant changes in volume.

Thus, in symptomatic mSOD1 mice (displaying the hypermetabolic phenotype), a substantial

remodeling of cortico-hypothalamic projections takes place, in particular from ORBl/vl+AI and

MOs.

Figure 2: Altered cortico-hypothalamic projection pattern in mSOD1 mice at P95. A:
Representative frontal brain sections of WT and mSOD1 mice depicting projections from
ORBl/vl+AI (significantly increased, arrow), PL+ORBm, ILA, ACA, MOs and MOp to LHA. Inset:
representative injection sites in LHA for WT and mSOD1 mice. White outlines represent LHA
boundaries. B: Sum of neurons projecting from selected 28 areas. No difference is detected in
the summarized number of neurons projecting to LHA in WT and mSOD1 from the ipsilateral
hemisphere (n=3). C: Quantification of the number of neurons (normalized for total neuronal
counts) projecting to LHA from 28 brain structures in WT and mSOD1. A significant increase in
projections from ORBl/vl+AI (p=0.0012) and from MOs (p=0.0006) is detected. D:
Representative *WholeBrain* volume reconstructions of neurons projecting to LHA in WT and
mSOD1 mice. Expansion of projections from MOs (blue) and ORBl/vl+AI (yellow) are visible
(arrows). Bars show mean ± SD. Scalebars 1 mm . *$p < 0.05$, ***$p < 0.001$, ****$p < 0.0001$.
404

Validation of cortico-hypothalamic projections abnormalities in independent operator-

registered brain cohorts

Since the identification of different cortical and subcortical areas was based on the assumption

of the overall similarity of the brain of mSOD1 and WT mice, we took into consideration the

possibility that biases in the identification of anatomical areas could have been introduced by

WholeBrain (e.g., because of atrophy), thereby mis-attributing neurons to cortical areas.

Therefore, we set out to validate our findings using the brain cohort 2 (n=3 for WT and mSOD1;

LHA injection; Supplementary Fig. 4-A). These brain images were subjected to manual

registration, i.e. each anatomical structure and cortical area was identified by a genotype-blind

operator using structural landmarks (Supplementary Table 1) and individually cropped out of

each image of each section of the brain. This approach was substantially slower than the

registration by *WholeBrain* (it required >500h of operator´s work time for the whole dataset to

be registered and cropped). The individual pictures cropped out of the brain images were then

subject to neuronal counts by *Ilastik* (since the accuracy and reproducibility of this step was

already established and quantified). In the manually-annotated dataset, the total number of

neurons projecting to LHA was found to be not significantly different in the mSOD1 compared to

the WT (Supplementary Fig. 4-B). First, we considered the absolute neuronal countings: the

contrast of the two genotypes (ipsilateral hemisphere) revealed a statistically significant

difference (two-way ANOVA $F_{1,112}$=6.28 ; p=0.0136) which was further explored in post-hoc

comparison (Sydak´s) revealing a significant expansion of projections from ORBl/vl+AI in mSOD1

mice (p=0.035) (Supplementary Fig. 4-C). Upon normalization we once again detected a

significant effect of genotype (two-way ANOVA $F_{1,112}$=23.78 ; p<0.0001) with post-hoc revealing

significant differences in ORBl/vl+AI (p<0.0001), ILA (p=0.001) and PIR (p<0.0001)

(Supplementary Fig. 4-D). When the projections from the contralateral hemisphere were

considered, no overall, genotype specific or area correlated effect was found in the absolute

counts (Supplementary Fig. 5-B); however, upon normalization, a statistically significant loss of

projections from PL+ORBm was identified (p<0.0001), together with a strong trend toward

increased projections from ORBl/vl+AI (p=0.072) (Supplementary Fig. 5-C). Thus, the manually-

registered dataset displayed substantial similarities with the *WholeBrain*-registered dataset

(expansion of the projections from ORBl/vl+AI) but also a major discrepancy: increased

projection from ILA to LHA in the manual dataset and no difference in MOs projections, whereas

the opposite was detected by *WholeBrain*. Upon closer inspection of MOs projection to LHA we

realized that the boundaries of MOs and ILA display a high degree of subjectivity when manually

drawn and it is highly likely that MOs neurons were allocated to adjacent areas in the same way

ILA was moved to neighboring areas in the dataset parcellated by the human operator.

Nevertheless, the manually-registered independent dataset confirmed the expansion of

projections from AI/ORBvl to LHA.

Unaltered projections to LHA in early presymptomatic mSOD mice

Remodelling of large-scale cortical circuits has been previously reported [21] and appears to progress over time; nevertheless, early signs of increased projections from somatosensory cortex (SS) to primary motor cortex (MOp) were detected already at early presymptomatic stages. We set out to verify if the increased projections from AI/ORBvl appeared only late, in coincidence with the appearance of the metabolic phenotype, or if they were present already at presymptomatic stage (and possibly of developmental origin). To this aim, we injected rAAV2-retro in the LHA of WT or mSOD1 at the age of P25 (Supplementary Fig. 6-A) and sacrificed the mice at P40. After quality control, we assessed the forebrain projections to LHA in the two genotypes (n=3 for each). The overall pattern of projections to LHA at P25 was comparable to the one observed at P95 and there was no difference in the absolute number of neurons projecting to LHA in the two genotypes (Supplementary Fig. 6-B). The direct contrast of the two genotypes for the absolute and relative contribution of 28 selected ipsilateral structures revealed a genotype effect in absolute (two-way ANOVA: $F_{1,112}$= 12.55 ; p=0.006) but not in the normalized dataset (two-way ANOVA: $F_{1,112}$= 1.83 ; p=0.1779). However, in both cases the post-hoc comparison did not reveal any significant difference between the two genotypes in any of the considered structures (Supplementary Fig. 6-C,D). Likewise, no statistical difference was found in the sum of projection from the contralateral hemisphere in absolute counts (Supplementary Fig. 7-A), but significant increase in projections from ACA was identified (p=0.0078) (Supplementary Fig. 7-B). Nevertheless, this difference disappeared after normalization (Supplementary Fig. 7-C) and no differences are perceptible in the 3D brain

reconstructions (Supplementary Fig. 7-D). These findings suggest that altered AI/ORBvl

connectivity to LHA arises during disease progression and it is not a developmental feature of

mSOD1 brain architecture.

ALS-related pathology in the AI/ORBvl of mSOD1 mice

We further investigated the degree of involvement of cortico-hypothalamic projections in ALS

by assessing the burden of misfolded SOD1 pathology in cortical areas projecting to LHA, in

particular those displaying altered connectivity compared to WT animals. We injected WT or

mSOD1 mice with rAAV2-retro in LHA at P95 and sacrificed them at P110. Brain sections

encompassing MOp, AI/ORBvl, PL+ORBm, (SS) were immunostained with an antibody

recognizing the misfolded conformation of SOD1 (B8H10 [44]) (Fig. 3A-B). The total burden of

misfSOD1 was different across the cortical areas (one-way ANOVA $F_{3,20}$=51.26, p<0.0001; MOp

vs. ORBl/vl+AI p<0.0001, MOp vs. PL+ORBm p<0.0001, SS vs. ORBl/vl+AI p=0.0014, ORBl/AI vs.

PL+ORBm p=0.0006), with MOp displaying the highest burden compared to SS but with

ORBl/vl+AI displaying a significantly higher burden than areas showing no increase in

projections such as SS or PL+ORBm (Fig. 3A-B). Interestingly, neurons projecting from MOs to

LHA display high levels of misfSOD1 accumulation, whereas this is not true for neurons

projecting from PL+ORBm and ORBl/vl+AI (Fig.3A).

Figure 3: Significant misfolded SOD1 burden in ORBl/vl+AI in mSOD1 mice. A: Representative images of misfSOD1+ cells in MOp, SS, ORBl/vl+AI and PL+ORBm; ZsGreen[+] neurons projecting to LHA show a strong misfSOD1 burden in MOp (arrows). Other structures show no double positive neurons (SS,ORBl/vl+AI) or weak mSOD burden (PL+ORBm; arrows). B: Burden of misfolded SOD1 (detected by the B8H10 antibody) in MOp, SS, ORBl/vl+AI and PL+ORBm. ORBl/vl+AI displays a burden significantly higher than SS or PL (p=0.0014 and p=0.0006 respectively; areas with unchanged connectivity to LHA) but significantly lower than MOp (p<0.0001). Bars show mean ± SD. Scalebars 50 µm. $**p < 0.01$, $****p < 0.0001$.

Distinct pattern of altered cortical projections to LHA in Fus mice

We further explored the alteration of projections to LHA in ALS in an independent ALS model, the kiFUS mouse. Of note, the kiFUS mouse differs from the mSOD1 mouse because it does not display an overt body-weight phenotype [45] and shows a more limited loss of spinal motoneurons and slower progression [29]. We proceeded with a similar experimental design as for the mSOD1 mice; however, since kiFUS mice display spinal motoneuron loss not before 9-10 months of age, we considered the P255 as "early symptomatic" stage. Therefore, kiFUS mice and their own WT littermates were injected at P255 into the LHA (Fig. 4A and sacrificed at P270 (n=4 for each group). Once again, there was no difference in the overall number of neurons projecting to LHA across the forebrain in absolute counts (Supplementary Fig. 8-A). When the

absolute neuronal counts of each of the 28 areas were contrasted, we detected a significant

effect of the genotype (two-way ANOVA $F_{1,166}$=8.583 ; p=0.0039) and post-hoc comparison

revealed a significant decrease in kiFUS in projections from ACA (p=0.0385), MOs (p<0.0001)

and strong trends for MOp (p=0.1543) and SS (p=0.0848) (Supplementary Fig. 8-B). Upon

normalization for 50,000 neurons, a significant difference in MOs was still detected (p<0.0001)

together with strong trends for ACA (p=0.1639) and MOp (p=0.136) (Fig. 4B-C and 2D models in

Supplementary Fig. 8-C,D). Interestingly, a significant genotype effect was also found in the

contralateral hemisphere both in absolute (Supplementary Fig. 9-A,B) and normalized data

(Supplementary Fig. 9-C) (two-way ANOVA, $F_{1,166}$= 6.57 ; p=0.0112 and $F_{1,166}$= 4.48 p=0.0356,

respectively) and differences were revealed by post-hoc analysis in loss of projections from ACA

(p=0.0121 and p=0.0137 in absolute counts and normalized, respectively) and ILA (p=0.0474) in

the absolute counts only. Of note, projections from AI/ORBvl were not affected (neither from

ipsilateral nor from contralateral). The significant loss of MOs neurons projecting to LHA was

especially affecting the most rostral part of MOs, as demonstrated by the 3D reconstruction

model of the mouse brain (Fig. 4D). Thus, distinct ALS models with divergent metabolic

phenotype display different patterns of altered projections to LHA.

Figure 4: Altered cortico-hypothalamic projection pattern in kiFUS mice at P270. A: Representative frontal brain sections of WT and kiFUS mice depicting projections from MOs (significantly decreased, arrow), PL+ORBm, ILA, ACA, and MOp to LHA. Inset: representative injection sites in LHA for WT and kiFUS mice. White outlines represent LHA boundaries. B: Sum of neurons projecting from selected 28 areas. No difference is detected in the summarized number of neurons projecting to LHA in WT and kiFUS from the ipsilateral hemisphere (n=4). C: Quantification of the number of neurons (normalized for total neuronal counts) projecting to LHA from 28 brain structures in WT and kiFUS. A significant decrease in projections from MOs (p<0.0001) and trend toward decrease projection from ACA (p=0.1639) and MOp (p=0.136) is detected (n=4). D: Representative *WholeBrain* volume reconstructions of neurons projecting to

539 LHA in WT and kiFUS mice. Loss of projections from the anterior part of MOs (blue) is visible.
540 Bars show mean ± SD. Scalebars 1 mm. *$p < 0.05$, **$p < 0.01$, ****$p < 0.001$.
541

DTI-MRI reveals altered orbitofrontal-hypothalamic tract in ALS patients.

Finally, we set out to investigate if the selective disturbance in cortico-hypothalamic projections

from AI, ORBvl and PL observed in the mSOD1 mice could also be observed in human ALS

patients. To this aim, we elected to use a 3T DTI-MRI dataset involving 83 ALS patients and 64

healthy subjects [39].Four white-matter tracts were taken into consideration: the orbitofrontal-

hypothalamic tract, the insular-hypothalamic tract, the cingulate-hypothalamic tract (Fig. 5) and

as reference the corticospinal tract which is known to be substantially altered in ALS patients

[39]; the three tracts converging on the hypothalamus were selected because they provided the

closest possible match (when accounting for the different anatomy) for the structures

investigated in the mouse model and because of their reproducible and unequivocal

identification in DTI datasets. We identified a significant decrease in Fractional Anisotropy (FA)

of the orbitofrontal-hypothalamic tract in ALS patients (ΔFA = -0.008±0.003; p<0.05), with a

magnitude approximately one-third of the FA loss in the corticospinal tract (ΔFA = -0.025±0.003;

average±SEM, p<0.001; Fig. 5). For the cingulate-hypothalamic and insular-hypothalamic tracts,

smaller FA losses were detected, which did not reach statistical significance. Thus, disturbances

of cortical projections to hypothalamus are not unique characteristics of the mSOD1 murine

model of ALS but constitute a previously unrecognized architectural phenotype shared by

human ALS patients.

Figure 5: Tract of Interest (TOI)-based analysis of DTI data from 83 ALS patients vs 64 controls. Upper panel: differences of mean fractional anisotropy (FA) values between ALS patients and controls. Lower panel: projectional views (axial, coronal, sagittal) of tract systems used for TOI analysis. TOI – tract-of-interest; *p < 0.05; **p < 0.001; error bars are the standard error of the mean (SEM).

Discussion

Here we provide converging evidence from ALS mouse models and human imaging datasets of the involvement of large-scale projections to LHA in ALS, in particular the disruption of connections from Orbitofrontal Cortex and/or Agranular Insula to LHA. This effect is independent of systematic biases introduced by software or human operators in the registration, does not appear to be an intrinsic developmental defect and it is not detected in an independent ALS model devoid of metabolic phenotypes. The findings in the murine model

display striking similarity to the disruption of the orbitofrontal-hypothalamic tract identified in a

large dataset of DTI-MRI of ALS patients.

Hypermetabolism in ALS has been repeatedly reported in human patients [9-13] as well as in

some, but not all, ALS murine models [14] and, notably, it appears to develop over time and

may predate the clinical onset of disease by several years [3, 46]. Nevertheless, the mechanisms

for this clinical manifestation of ALS are unclear.

Direct involvement of the hypothalamus has been hypothesized on the basis of the role of this

structure as the main central controller of energy homeostasis, feeding and satiety [47], has

been supported by evidence of reduced hypothalamic volume in ALS patients [48] and by the

detection of ALS-related TDP-43 pathology in a subset of ALS patients hypothalami [49].

Neurochemical abnormalities, such as reduced MCH expression, have been reported in mutant

SOD1 murine models [50]. However, these signs of intrinsic hypothalamic disturbance are not

mutually exclusive with the hypothesis that large-scale hypothalami circuits may be

dysfunctional, too. Disruption within the larger circuit regulating hypothalamic function, which

include insular, motor, orbitofrontal and limbic areas, may contribute to the hypothalamic

dysregulation by altering the equilibrium of the different inputs. Indeed, changes in neuronal

activity in hypothalamus may produce substantial modifications of the neurochemical identity

of local subpopulation, including switch in type of neurotransmitter released [51]. Here we

demonstrate that besides any pathology intrinsic to the hypothalamus, the disease

pathophysiology involves a substantial quantitative change in the inputs that hypothalamus

receives, particularly from agranular insula and ventrolateral orbitofrontal cortex. This

remodeling process coincides with the onset of body weight loss in mSOD1.

Notably, the circuit involving projections from orbitofrontal cortex and agranular insula to

lateral hypothalamus is conserved from mice and rats [52, 53] to marmosets [54] and macaques

[55] to humans, thus validating mouse as a model organism. Nevertheless, the murine

orbitofrontal cortex (including ventral and ventrolateral) and agranular insula are considered

homologous of orbitomedial cortex in primates [56], so that the study the orbitofrontal-LHA

tract in our DTI tracing study is located more medially than the ORBl/vl+AI complex in the

mouse brain.

Remarkably, altered projections from AI/ORBvl in mSOD1 mice correspond to reduced FA in the

orbitofrontal-hypothalamic (but not insular-hypothalamic) tract in humans. Although it is

noteworthy that abnormalities appear in the same brain structures in animal models and

patients with ALS, the closer examination of the findings reveal that in the murine model an

actual expansion of projections is detected whereas decreased FA in DTI is usually interpreted

as loss of integrity of axons or myelinated tracts [57]. However, it must be stressed that

SOD1$^{(G93A)}$ displays a fast progression and becomes symptomatic at a comparatively young age,

when the potential for compensatory sprouting after degeneration may be larger than in older

human patients (hence, the expansion of the projections). Thus, in human patients the DTI may

reveal a change in microstructural integrity resulting from axonal loss but would be unable to

detect any increase in the size of axonal arborization.

To the best of our knowledge, there is no demonstration that the orbitofrontal cortex and/or AI

can directly regulate metabolic rates. It must be emphasized, though, that lesion studies of

these structures have not included a rigorous, long-term testing of body-weight dynamics and

inactivation studies have been performed only in short-term settings [58-60]. On the other

hand, OFC and AI are involved in processing food characteristics, such as taste, smell and

texture [61] and display a differential response to caloric content of food depending on the

satiety of the subject (at least in humans: [62, 63]). Thus, it is conceivable that altered

connectivity between vlORB/AI and LHA may provide the latter with incorrect inputs regarding

the nutritional content of food, contributing to the dysregulation of body metabolic rates. At

least in human subjects, reduced volume of grey matter and altered microstructural integrity

(measured by apparent diffusion coefficient) in orbitofrontal cortex have been observed in

obese patients [64-66], strengthening the link between dysfunction in orbitofrontal cortex and

body weight.

Unexpectedly, our retrograde projection mapping strategy has consistently identified

projections from primary and secondary motor cortex to LHA, which appear to be affected by

the disease process both in mSOD1 and in kiFUS. This connectivity appears to be reciprocal,

since LHA projections to motor cortices have also been reported [21]. These connections cannot

be easily discounted as artifacts due to the proximity of LHA to the corticospinal tract, since they

appear to originate from both ipsi and contralateral hemispheres. It is conceivable that LHA may

receive inputs relaying the average motor activity performed or planned and may adjust energy

balance accordingly; disruption of connectivity between motor cortex and hypothalamus may

leave the latter "free running", without a proper estimate of current motor activity.

Alternatively, it is possible that any pathogenic event in motor cortex (propagating prionoids or

abnormal activity patterns) may propagate to hypothalamus and drive its dysfunction and

degeneration (as suggested for spinal cord [67]).

Conclusion

Our findings suggest that disruption of large-scale circuits providing input to LHA may contribute

to generate the metabolic phenotype observed in ALS; in particular, we have identified the

orbitofrontal-hypothalamic tract as a site of convergence of mouse projection data and human

tracing data, which may open up an independent approach to evaluate non-motor prognostic

features in ALS.

Declarations

Acknowledgements

We thank the colleagues of the German Center for Neurodegenerative Diseases (DZNE)-Ulm for

helpful comments.

Authors' contributions

DB contributed to experimental design and performed the stereotactic injections and processed

the samples for imaging and performed the image analysis using the WholeBrain registration.

SA wrote the image analysis pipeline and performed the manual parcellation study. HPM and JK

performed the analysis of the MRI-DTI dataset. AL contributed to the MRI-DTI dataset. FR and JK

designed the experiments, analyzed the data and wrote the first version of the manuscript. FR,

JK, AL, TB and LD contributed to the final version of the manuscript and to the interpretation of

the findings. All authors read and approved the final manuscript.

Conflict of interst

The authors declare no financial interest or conflict of interest.

Ethics approval and consent to participate

The animal experiments were approved by the Regierungspräsidium Tübingen under the animal

license- number 1390 and by the Ulm University Tierforschungszentrum committee.

The collection of MRI-DTI data from human subjects has been approved by the Ethical

Committee of the University of Ulm (reference #19/12)

Availability of data and materials

The datasets used and/or analysed during the current study are available from the

corresponding author on reasonable request.

Funding

FR is supported by the Thierry Latran Foundation (projects "Trials" and "Hypothals"), by the

Radala Foundation, by the Deutsche Forschungsgemeinschaft (DFG, German Research

Foundation)- Project-ID 251293561 – Collaborative Research Center (CRC) 1149 and with the

individual grant no. 431995586 (RO-5004/8-1) and no. 443642953 (RO5004/9-1), by the Cellular

690 **References:**

691 1) Hardiman O, Al-Chalabi A, Chio A, et al. Amyotrophic lateral sclerosis [published
692   correction appears in Nat Rev Dis Primers. 2017 Oct 20;3:17085]. *Nat Rev Dis Primers*.
693   2017;3:17071. Published 2017 Oct 5. doi:10.1038/nrdp.2017.71
694

695 2) Dupuis L, Pradat PF, Ludolph AC, Loeffler JP. Energy metabolism in amyotrophic lateral
696   sclerosis. *Lancet Neurol*. 2011;10(1):75-82. doi:10.1016/S1474-4422(10)70224-6
697

698 3) Peter RS, Rosenbohm A, Dupuis L, et al. Life course body mass index and risk and
699   prognosis of amyotrophic lateral sclerosis: results from the ALS registry Swabia. *Eur J
700   Epidemiol*. 2017;32(10):901-908. doi:10.1007/s10654-017-0318-z
701

702 4) Gallo V, Wark PA, Jenab M, et al. Prediagnostic body fat and risk of death from
703   amyotrophic lateral sclerosis: the EPIC cohort. *Neurology*. 2013;80(9):829-838.
704   doi:10.1212/WNL.0b013e3182840689
705

706 5) Dupuis L, Corcia P, Fergani A, et al. Dyslipidemia is a protective factor in amyotrophic
707   lateral sclerosis. *Neurology*. 2008;70(13):1004-1009.
708   doi:10.1212/01.wnl.0000285080.70324.27
709

710 6) Dorst J, Kühnlein P, Hendrich C, Kassubek J, Sperfeld AD, Ludolph AC. Patients with
711   elevated triglyceride and cholesterol serum levels have a prolonged survival in
712   amyotrophic lateral sclerosis. *J Neurol*. 2011;258(4):613-617. doi:10.1007/s00415-010-
713   5805-z
714

715 7) Lindauer E, Dupuis L, Müller HP, Neumann H, Ludolph AC, Kassubek J. Adipose Tissue
716   Distribution Predicts Survival in Amyotrophic Lateral Sclerosis. *PLoS One*.
717   2013;8(6):e67783. Published 2013 Jun 27. doi:10.1371/journal.pone.0067783
718

719 8) Marin B, Desport JC, Kajeu P, et al. Alteration of nutritional status at diagnosis is a
720   prognostic factor for survival of amyotrophic lateral sclerosis patients. *J Neurol
721   Neurosurg Psychiatry*. 2011;82(6):628-634. doi:10.1136/jnnp.2010.211474
722

723 9) Steyn FJ, Ioannides ZA, van Eijk RPA, et al. Hypermetabolism in ALS is associated with
724   greater functional decline and shorter survival. *J Neurol Neurosurg Psychiatry*.
725   2018;89(10):1016-1023. doi:10.1136/jnnp-2017-317887
726
727 10) Jésus P, Fayemendy P, Nicol M, et al. Hypermetabolism is a deleterious prognostic factor
728   in patients with amyotrophic lateral sclerosis. *Eur J Neurol*. 2018;25(1):97-104.
729   doi:10.1111/ene.13468
730
731 11) Ahmed RM, Irish M, Piguet O, et al. Amyotrophic lateral sclerosis and frontotemporal
732   dementia: distinct and overlapping changes in eating behaviour and metabolism
733   [published correction appears in Lancet Neurol. 2016 Apr;15(4):352]. *Lancet Neurol*.
734   2016;15(3):332-342. doi:10.1016/S1474-4422(15)00380-4
735
736 12) Desport JC, Torny F, Lacoste M, Preux PM, Couratier P. Hypermetabolism in ALS:
737   correlations with clinical and paraclinical parameters. *Neurodegener Dis*. 2005;2(3-
738   4):202-207. doi:10.1159/000089626
739
740 13) Bouteloup C, Desport JC, Clavelou P, et al. Hypermetabolism in ALS patients: an early and
741   persistent phenomenon. *J Neurol*. 2009;256(8):1236-1242. doi:10.1007/s00415-009-
742   5100-z
743
744 14) Dupuis L, Oudart H, René F, Gonzalez de Aguilar JL, Loeffler JP. Evidence for defective
745   energy homeostasis in amyotrophic lateral sclerosis: benefit of a high-energy diet in a
746   transgenic mouse model. *Proc Natl Acad Sci U S A*. 2004;101(30):11159-11164.
747   doi:10.1073/pnas.0402026101
748
749 15) Ludolph AC, Dorst J, Dreyhaupt J, et al. Effect of High-Caloric Nutrition on Survival in
750   Amyotrophic Lateral Sclerosis. *Ann Neurol*. 2020;87(2):206-216. doi:10.1002/ana.25661
751
752 16) González JA, Iordanidou P, Strom M, Adamantidis A, Burdakov D. Awake dynamics and
753   brain-wide direct inputs of hypothalamic MCH and orexin networks. *Nat Commun*.
754   2016;7:11395. Published 2016 Apr 22. doi:10.1038/ncomms11395
755
756 17) Barbier M, Chometton S, Pautrat A, et al. A basal ganglia-like cortical-amygdalar-
757   hypothalamic network mediates feeding behavior. *Proc Natl Acad Sci U S A*.
758   2020;117(27):15967-15976. doi:10.1073/pnas.2004914117
759
760 18) Barbier M, González JA, Houdayer C, Burdakov D, Risold PY, Croizier S. Projections from
761   the dorsomedial division of the bed nucleus of the stria terminalis to hypothalamic
762   nuclei in the mouse [published online ahead of print, 2020 Jul 17]. *J Comp Neurol*.
763   2020;10.1002/cne.24988. doi:10.1002/cne.24988
764

19) Murata K, Kinoshita T, Fukazawa Y, et al. GABAergic neurons in the olfactory cortex
projecting to the lateral hypothalamus in mice. *Sci Rep*. 2019;9(1):7132. Published 2019
May 9. doi:10.1038/s41598-019-43580-1

20) Berthoud HR. Multiple neural systems controlling food intake and body weight. *Neurosci*
Biobehav Rev. 2002;26(4):393-428. doi:10.1016/s0149-7634(02)00014-3

21) Commisso B, Ding L, Varadi K, et al. Stage-dependent remodeling of projections to motor
cortex in ALS mouse model revealed by a new variant retrograde-AAV9. *Elife*.
2018;7:e36892. Published 2018 Aug 23. doi:10.7554/eLife.36892

22) Agosta F, Canu E, Valsasina P, et al. Divergent brain network connectivity in amyotrophic
lateral sclerosis. *Neurobiol Aging*. 2013;34(2):419-427.
doi:10.1016/j.neurobiolaging.2012.04.015

23) Schulthess I, Gorges M, Müller HP, et al. Functional connectivity changes resemble
patterns of pTDP-43 pathology in amyotrophic lateral sclerosis. *Sci Rep*. 2016;6:38391.
Published 2016 Dec 8. doi:10.1038/srep38391

24) Heimrath J, Gorges M, Kassubek J, et al. Additional resources and the default mode
network: Evidence of increased connectivity and decreased white matter integrity in
amyotrophic lateral sclerosis. *Amyotroph Lateral Scler Frontotemporal Degener*.
2014;15(7-8):537-545. doi:10.3109/21678421.2014.911914

25) Dukic S, McMackin R, Buxo T, et al. Patterned functional network disruption in
amyotrophic lateral sclerosis. *Hum Brain Mapp*. 2019;40(16):4827-4842.
doi:10.1002/hbm.24740

26) Braak H, Brettschneider J, Ludolph AC, Lee VM, Trojanowski JQ, Del Tredici K.
Amyotrophic lateral sclerosis--a model of corticofugal axonal spread. *Nat Rev Neurol*.
2013;9(12):708-714. doi:10.1038/nrneurol.2013.221

27) Brettschneider J, Del Tredici K, Toledo JB, et al. Stages of pTDP-43 pathology in
amyotrophic lateral sclerosis. *Ann Neurol*. 2013;74(1):20-38. doi:10.1002/ana.23937

28) Scekic-Zahirovic J, Sendscheid O, El Oussini H, et al. Toxic gain of function from mutant
FUS protein is crucial to trigger cell autonomous motor neuron loss. *EMBO J*.
2016;35(10):1077-1097. doi:10.15252/embj.201592559

29) Ouali Alami N, Schurr C, Olde Heuvel F, et al. NF-κB activation in astrocytes drives a
stage-specific beneficial neuroimmunological response in ALS. *EMBO J*.
2018;37(16):e98697. doi:10.15252/embj.201798697

30) Tervo DG, Hwang BY, Viswanathan S, et al. A Designer AAV Variant Permits Efficient
Retrograde Access to Projection Neurons. *Neuron*. 2016;92(2):372-382.
doi:10.1016/j.neuron.2016.09.021

31) Madisen L, Garner AR, Shimaoka D, et al. Transgenic mice for intersectional targeting of
neural sensors and effectors with high specificity and performance. *Neuron*.
2015;85(5):942-958. doi:10.1016/j.neuron.2015.02.022

32) Paxinos G, Franklin KBJ. The mouse brain in stereotaxic coordinates, second edition.
(*ACADEMIC PRESS*, 2001), ISBN: 0-12-547636-1

33) Schindelin J, Arganda-Carreras I, Frise E, et al. Fiji: an open-source platform for
biological-image analysis. *Nat Methods*. 2012;9(7):676-682. Published 2012 Jun 28.
doi:10.1038/nmeth.2019

34) Fürth D, Vaissière T, Tzortzi O, et al. An interactive framework for whole-brain maps at
cellular resolution [published correction appears in Nat Neurosci. 2017 Dec 18;:]. *Nat
Neurosci*. 2018;21(1):139-149. doi:10.1038/s41593-017-0027-7

35) Lein ES, Hawrylycz MJ, Ao N, et al. Genome-wide atlas of gene expression in the adult
mouse brain. *Nature*. 2007;445(7124):168-176. doi:10.1038/nature05453

36) Berg S, Kutra D, Kroeger T, et al. ilastik: interactive machine learning for (bio)image
analysis. *Nat Methods*. 2019;16(12):1226-1232. doi:10.1038/s41592-019-0582-9

37) Fiaschi L, Koethe U, Nair R, Hamprecht FA. Learning to count with regression forest and
structured labels. *Proceedings of the 21st International Conference on Pattern
Recognition (ICPR2012)*, Tsukuba, 2012;2685-2688.

38) Sahoo P, Wilkins C, Yeager J. Threshold selection using Renyi's entropy. Pattern
Recognition. 1997;30(1):71-84. doi:10.1016/S0031-3203(96)00065-9

39) Kassubek J, Müller HP, Del Tredici K, et al. Imaging the pathoanatomy of amyotrophic
lateral sclerosis in vivo: targeting a propagation-based biological marker. *J Neurol
Neurosurg Psychiatry*. 2018;89(4):374-381. doi:10.1136/jnnp-2017-316365

40) Müller HP, Unrath A, Ludolph AC, Kassubek J. Preservation of diffusion tensor properties
during spatial normalization by use of tensor imaging and fibre tracking on a normal
brain database. *Phys Med Biol*. 2007;52(6):N99-N109. doi:10.1088/0031-9155/52/6/N01

41) Müller HP, Del Tredici K, Lulé D, et al. In vivo histopathological staging in C9orf72-
associated ALS: A tract of interest DTI study. *Neuroimage Clin*. 2020;27:102298.
doi:10.1016/j.nicl.2020.102298

42) Müller HP, Unrath A, Sperfeld AD, Ludolph AC, Riecker A, Kassubek J. Diffusion tensor
imaging and tractwise fractional anisotropy statistics: quantitative analysis in white
matter pathology. *Biomed Eng Online*. 2007;6:42. Published 2007 Nov 9.
doi:10.1186/1475-925X-6-42

43) Petrik MS, Wilson JM, Grant SC, et al. Magnetic resonance microscopy and
immunohistochemistry of the CNS of the mutant SOD murine model of ALS reveals
widespread neural deficits. *Neuromolecular Med*. 2007;9(3):216-229.
doi:10.1007/s12017-007-8002-1

44) Pickles S, Destroismaisons L, Peyrard SL, et al. Mitochondrial damage revealed by
immunoselection for ALS-linked misfolded SOD1. *Hum Mol Genet*. 2013;22(19):3947-
3959. doi:10.1093/hmg/ddt249

45) Scekic-Zahirovic J, Oussini HE, Mersmann S, et al. Motor neuron intrinsic and extrinsic
mechanisms contribute to the pathogenesis of FUS-associated amyotrophic lateral
sclerosis. *Acta Neuropathol*. 2017;133(6):887-906. doi:10.1007/s00401-017-1687-9

46) Mariosa D, Beard JD, Umbach DM, et al. Body Mass Index and Amyotrophic Lateral
Sclerosis: A Study of US Military Veterans. *Am J Epidemiol*. 2017;185(5):362-371.
doi:10.1093/aje/kww140

47) Vercruysse P, Vieau D, Blum D, Petersén Å, Dupuis L. Hypothalamic Alterations in
Neurodegenerative Diseases and Their Relation to Abnormal Energy Metabolism. *Front
Mol Neurosci*. 2018;11:2. Published 2018 Jan 19. doi:10.3389/fnmol.2018.00002

48) Gorges M, Vercruysse P, Müller HP, et al. Hypothalamic atrophy is related to body mass
index and age at onset in amyotrophic lateral sclerosis. *J Neurol Neurosurg Psychiatry*.
2017;88(12):1033-1041. doi:10.1136/jnnp-2017-315795

49) Cykowski MD, Takei H, Schulz PE, Appel SH, Powell SZ. TDP-43 pathology in the basal
forebrain and hypothalamus of patients with amyotrophic lateral sclerosis. *Acta
Neuropathol Commun*. 2014;2:171. Published 2014 Dec 24. doi:10.1186/s40478-014-
0171-1

50) Vercruysse P, Sinniger J, El Oussini H, et al. Alterations in the hypothalamic melanocortin
pathway in amyotrophic lateral sclerosis. *Brain*. 2016;139(Pt 4):1106-1122.
doi:10.1093/brain/aww004

51) Meng D, Li HQ, Deisseroth K, Leutgeb S, Spitzer NC. Neuronal activity regulates
neurotransmitter switching in the adult brain following light-induced stress. Proc Natl
Acad Sci U S A. 2018 May 15;115(20):5064-5071.

52) Floyd NS, Price JL, Ferry AT, Keay KA, Bandler R. Orbitomedial prefrontal cortical
projections to hypothalamus in the rat. *J Comp Neurol*. 2001;432(3):307-328.
doi:10.1002/cne.1105
899
900   53) Hoover WB, Vertes RP. Projections of the medial orbital and ventral orbital cortex in the
901   rat. *J Comp Neurol*. 2011;519(18):3766-3801. doi:10.1002/cne.22733
902
903   54) Roberts AC, Tomic DL, Parkinson CH, et al. Forebrain connectivity of the prefrontal cortex
904   in the marmoset monkey (Callithrix jacchus): an anterograde and retrograde tract-
905   tracing study. *J Comp Neurol*. 2007;502(1):86-112. doi:10.1002/cne.21300
906
907   55) Ongür D, An X, Price JL. Prefrontal cortical projections to the hypothalamus in macaque
908   monkeys. *J Comp Neurol*. 1998;401(4):480-505.
909
910   56) Price JL. Definition of the orbital cortex in relation to specific connections with limbic and
911   visceral structures and other cortical regions. *Ann N Y Acad Sci*. 2007;1121:54-71.
912   doi:10.1196/annals.1401.008
913
914   57) Alexander AL, Lee JE, Lazar M, Field AS. Diffusion tensor imaging of the brain.
915   *Neurotherapeutics*. 2007;4(3):316-329. doi:10.1016/j.nurt.2007.05.011
916
917   58) Kobayashi M. Macroscopic connection of rat insular cortex: anatomical bases underlying
918   its physiological functions. *Int Rev Neurobiol*. 2011;97:285-303. doi:10.1016/B978-0-12-
919   385198-7.00011-4
920
921   59) Izquierdo A. Functional Heterogeneity within Rat Orbitofrontal Cortex in Reward
922   Learning and Decision Making. *J Neurosci*. 2017;37(44):10529-10540.
923   doi:10.1523/JNEUROSCI.1678-17.2017
924
925   60) Rolls ET. The functions of the orbitofrontal cortex. *Brain Cogn*. 2004;55(1):11-29.
926   doi:10.1016/S0278-2626(03)00277-X
927
928   61) Seabrook LT, Borgland SL. The orbitofrontal cortex, food intake and obesity. *J Psychiatry*
929   *Neurosci*. 2020;45(5):304-312. doi:10.1503/jpn.190163
930
931   62) Schur EA, Kleinhans NM, Goldberg J, Buchwald D, Schwartz MW, Maravilla K. Activation
932   in brain energy regulation and reward centers by food cues varies with choice of visual
933   stimulus. *Int J Obes (Lond)*. 2009;33(6):653-661. doi:10.1038/ijo.2009.56
934
935   63) Suzuki S, Cross L, O'Doherty JP. Elucidating the underlying components of food valuation
936   in the human orbitofrontal cortex. *Nat Neurosci*. 2017;20(12):1780-1786.
937   doi:10.1038/s41593-017-0008-x
938

939   64) Raji CA, Ho AJ, Parikshak NN, et al. Brain structure and obesity. *Hum Brain Mapp*.
940   2010;31(3):353-364. doi:10.1002/hbm.20870
941

942   65) Walther K, Birdsill AC, Glisky EL, Ryan L. Structural brain differences and cognitive
943   functioning related to body mass index in older females. *Hum Brain Mapp*.
944   2010;31(7):1052-1064. doi:10.1002/hbm.20916
945

946   66) Alkan A, Sahin I, Keskin L, et al. Diffusion-weighted imaging features of brain in obesity.
947   *Magn Reson Imaging*. 2008;26(4):446-450. doi:10.1016/j.mri.2007.10.004
948

949   67) Burg T, Bichara C, Scekic-Zahirovic J, et al. Absence of Subcerebral Projection Neurons Is
950   Beneficial in a Mouse Model of Amyotrophic Lateral Sclerosis [published online ahead of
951   print, 2020 Jun 26]. *Ann Neurol*. 2020;10.1002/ana.25833. doi:10.1002/ana.25833
952

Supplementary:

Supplementary _Table 1_: Anatomical landmarks used for mapping the brain section from
anterior to posterior. The following structures provided distinctive optical characteristics for
mapping: AOB = Accessory olfactory bulb; DP = Dorsal peduncular area; fa = corpus callosum,
anterior forceps; ccg = genu of corpus callosum; aco = anterior commissure, olfactory limb; act =
anterior commissure, temporal limb; SCH = Suprachiasmatic nucleus; NLOT = Nucleus of the
lateral olfactory tract; ME = Median eminence; opt = optic tract; int = internal capsule; DG =
Dentate gyrus; CA3so = Field CA3, stratum oriens; MM = Medial mammillary nucleus.
962

Coordinate from bregma	Anatomical landmark	Section in Allen´s brain atlas
2,6	first section without AOB	27
2,2	section before beginning of DP	32
2,1	first section with DP	33
1,6	shape of fa	38
1,5	shape of fa	39
1,4	shape of fa	40
1,2	shape of fa	42
1,1	shape of fa	43
1	shape of ccg	44
0,2	shape of aco	52
0,1	shape of aco	53
0	shape of aco and act	54
-0,3	first section with SCH last section without NLOT	57
-1	shape of opt last section with NLOT	64
-1,4	first section with ME shape of otp and int	69
-2,2	first section without ME first section with ventral DG	76
-2,5	Expansion of CA3so	79
-3,1	last section with MM	85

963
964

Supplementary Table 2: List of structures of the brain projecting to LHA. For each structure, the absolute number of neurons projecting to LHA is reported as mean ± SD and independently for ipsi- and contralateral hemispheres.

	Ipsilateral raw data	Contralateral raw data	Ipsilateral normalized	Contralateral normalized
A13	85.6, ±45.9	10.4, ±6.6	58.5, ±28.2	7.1, ±3.8
AAA	297.6, ±230.0	18.8, ±7.2	225.3, ±128.1	16.3, ±7.4
ACA	2686.2, ±1373.5	897.0, ±468.9	1826.0, ±538.9	599.4, ±232.6
ACB	565.4, ±375.1	9.4, ±10.1	411.8, ±173.1	7.9, ±6.9
ADP	64.4, ±27.3	16.6, ±8.6	48.3, ±10.8	12.5, ±4.1
AHN	1174.0, ±1241.1	111.4, ±91.1	909.7, ±1146.0	84.7, ±77.7
AI	1437.8, ±1365.3	197.0, ±177.6	935.9, ±622.2	117.2, ±82.2
AON	92.2, ±90.1	6.2, ±4.4	64.6, ±41.4	4.1, ±1.6
APN	54.2, ±43.3	5.4, ±1.7	35.4, ±19.1	5.7, ±4.8
AUD	228.6, ±246.6	62.2, ±48.6	138.4, ±129.3	38.0, ±27.4
AVP	207.8, ±99.8	71.6, ±36.0	143.2, ±41.5	49.3, ±22.1
AVPV	242.0, ±43.6	71.2, ±31.5	216.0, ±99.8	70.5, ±57.7
BLA	1263.0, ±670.4	79.6, ±56.3	920.2, ±278.5	52.6, ±28.0
BMA	1405.0, ±744.5	110.4, ±67.5	1009.6, ±381.6	77.4, ±37.9
BST	2108.8, ±774.0	275.6, ±165.1	1613.9, ±236.4	188.0, ±82.2
CEA	440.4, ±315.9	14.4, ±6.8	292.6, ±129.9	12.6, ±7.0
CLA	20.0, ±11.5	4.4, ±2.2	14.8, ±4.9	3.8, ±2.1
CM	132.8, ±65.5	49.2, ±18.7	97.1, ±40.2	37.5, ±7.1
COA	730.4, ±377.3	39.2, ±31.1	539.2, ±179.2	25.4, ±17
CP	814.4, ±499.4	34.8, ±21.2	581.7, ±185.8	23.0, ±7.0
DMH	1510.0, ±729.1	233.8, ±115.7	1039.7, ±290.7	160.3, ±50
DP	618.4, ±249.1	192.2, ±128.9	535.8, ±289.4	131.5, ±54.3
ECT	255.8, ±139.4	79.0, ±50.1	172.5, ±48.9	50.5, ±26.3
ENTI	144.8, ±96.2	28.6, ±18.1	98.2, ±46.0	18.1, ±10.3
EP	252.0, ±241.5	9.4, ±8.1	175.9, ±109.7	5.8, ±4.8
FS	59.8, ±45.7	1.0, ±0.9	48.2, ±31.9	0.7, ±0.7
GP	133.8, ±83.9	9.8, ±10.7	90.2, ±27.0	6.0, ±5.3
GU	503.2, ±399.3	77.0, ±63.8	343.9, ±186.6	49.7, ±28.3
IA	127.4, ±90.9	8.0, ±5.8	92.8, ±70.4	6.5, ±5.0
ILA	2627.2, ±896.1	1192.2, ±548.6	2089.0, ±476.3	834.9, ±188.3
IMD	145.8, ±74.4	55.8, ±29.8	103.6, ±34.6	37.5, ±11.9
LH	188.4, ±99.4	32.6, ±17.2	127.9, ±41.1	21.8, ±9.0
LHA	Injection	845.8, ±509.0	Injection	573.8, ±167.2
LPO	1251.6, ±447.7	173.2, ±113.4	1000.5, ±271.0	120.5, ±51.3
LS	1453.4, ±583.2	179.6, ±78.9	1169.3, ±436.5	127.3, ±23.3
MA	139.6, ±177.3	5.8, ±4.1	97.8, ±90.1	6.0, ±6.1

MEA	712.8, ±691.0	72.4, ±103.5	499.2, ±437.3	46.6, ±66.8
MEPO	252.8, ±96.0	204.6, ±71.7	207.2, ±71.9	169.9, ±57.2
MM	126.0, ±105.5	55.2, ±49.2	86.8, ±54.7	36.4, ±29.1
MOp	1006.4, ±1321.7	71.6, ±74.0	576.5, ±705.8	40.4, ±38.2
MOs	2292.6, ±2310.5	273.6, ±224.4	1391.5, ±1146.2	168.8, ±103.4
MPN	701.2, ±246.6	177.2, ±104.2	551.6, ±124.7	121.4, ±56.4
MPO	1752.4, ±543.1	436.4, ±213.4	1384.5, ±259.7	306.7, ±83.4
MS	323.0, ±157.8	58.4, ±29.3	240.5, ±56.7	47.0, ±16.7
NDB	288.2, ±281.3	34.4, ±24.5	202.7, ±126.0	24.7, ±20.6
NLOT	74.0, ±40.3	8.0, ±5.4	55.9, ±20.7	5.5, ±3.1
OLF	125.4, ±76.6	19.0, ±11.4	97.4, ±39.3	13.5, ±4.9
ORBl	306.6, ±425.5	41.8, ±68.5	168.7, ±213.7	22.7, ±34.6
ORBm	515.4, ±261.3	142.6, ±120.6	421.2, ±183.3	104.2, ±53.7
ORBvl	184.8, ±193.9	31.4, ±42.7	110.2, ±90.7	18.6, ±20.7
OT	58.4, ±74.6	1.8, ±1.7	37.9, ±35.2	1.5, ±1.0
OV	28.2, ±9.0	17.2, ±4.2	24.9, ±12.4	16.2, ±9.8
PA	398.4, ±541.0	156.2, ±271.9	252.9, ±347.2	99.8, ±175.5
PAA	31.2, ±11.4	2.6, ±0.8	25.3, ±7.4	2.5, ±1.5
PD	17.8, ±7.0	3.6, ±3.8	14.7, ±5.9	2.6, ±2.3
PERI	235.0, ±141.4	60.8, ±34.0	156.4, ±61.6	39.9, ±15.5
PH	1304.0, ±428.6	662.0, ±209.6	1043.6, ±248.9	528.6, ±123.8
PIR	667.0, ±677.1	27.0, ±12.9	444.4, ±302.3	20.2, ±6.7
PL	1576.0, ±554.1	653.6, ±330.0	1239.8, ±254.5	466.9, ±93.6
PS	33.2, ±20.0	2.8, ±2.5	22.2, ±6.3	1.8, ±1.5
PST	37.0, ±26.9	4.0, ±4.6	41.3, ±47.5	2.9, ±2.5
PSTN	160.2, ±59.7	16.6, ±12.9	168.5, ±158.2	11.4, ±5.6
PTL	199.4, ±198.2	17.0, ±11.1	121.9, ±104.4	11.2, ±5.0
PV	239.8, ±164.1	46.2, ±27.2	171.9, ±111.7	34.1, ±14.2
PVH	449.2, ±210.4	72.4, ±46.7	339.2, ±164.4	50.9, ±33.3
PVT	960.4, ±475.5	332.4, ±231.1	686.5, ±334.9	242.4, ±193.1
RE	461.0, ±379.7	176.4, ±122.5	356.0, ±347.2	134.9, ±109.3
RSP	331.8, ±297.9	47.6, ±32.1	198.7, ±152.4	30.7, ±15.4
SF	57.8, ±32.7	13.6, ±6.0	41.9, ±21.1	9.7, ±3.6
SI	1278.0, ±723.9	71.8, ±65.0	982.8, ±364.2	46.3, ±28.1
SS	700.6, ±860.7	57.4, ±49.5	407.5, ±454	40.5, ±25.0
STN	124.4, ±78.1	20.0, ±11.8	116.4, ±96.7	13.2, ±5.1
SUM	475.2, ±235.5	211.4, ±145.1	370.8, ±110.8	144.5, ±54.0
TEa	205.6, ±122.0	78.6, ±58.0	136.7, ±45.9	49.3, ±29.6
TRS	27.8, ±18.0	20.4, ±19.1	24.1, ±16.7	14.6, ±11.3
TT	929.4, ±512.0	214.6, ±194.3	802.0, ±531.2	131.0, ±95.0
TU	531.4, ±445.1	60.8, ±36.4	353.3, ±263.6	41.7, ±18.4
VIS	68.4, ±49.9	5.0, ±4.1	42.7, ±26.3	3.6, ±3.7

VISC	138.6, ±109.1	22.0, ±21.0	86.0, ±49.0	14.3, ±9.3
VLPO	57.4, ±30.1	17.8, ±15.3	40.3, ±11.4	11.3, ±9.3
VM	821.2, ±745.3	84.8, ±34.6	523.4, ±351.8	63.8, ±20.3
VMH	1462.6, ±799.9	187.4, ±149.2	1004.7, ±629.3	137.7, ±133.4
VP	163.2, ±158.4	28.0, ±12.4	110.7, ±75.8	21.4, ±5.6
ZI	1829.0, ±912.2	263.0, ±107.7	1374.4, ±334.2	194.9, ±34.7

Supplementary Figure 1-9 are provided in a separate PDF

Supplementary Fig. 1: Altered cortico-hypothalamic projection pattern in mSOD1 mice at P95, absolute counts ipsilateral. A: Sum of neurons projecting from selected 28 areas. Significantly increased projections from ipsilateral hemisphere in mSOD mice to LHA (p=0.02; n=3). B: Quantification of the absolute number of neurons projecting to LHA from 28 brain structures in WT and mSOD1. A significant increase in projections from ACA (p<0.0001), ILA (p=0.009), ORBl/vl+AI (p<0.0001), Tenia Tecta (TT; p=0.0365), secondary Motor Cortex (MOs, p<0.0001) and Piriform cortex (PIR; p=0.0005). C: Representative *WholeBrain* ipsilateral side view reconstructions of neurons projecting to LHA in WT and mSOD1 mice. Expansion of projections from MOs are visible (arrows). D: Representative *WholeBrain* cortical top view reconstructions of neurons projecting to LHA in WT and mSOD1 mice. Expansion of projections from MOs are visible (arrows). Bars show mean ± SD. *p < 0.05, **p < 0.01, ***p < 0.001, ****p< 0.0001.

Supplementary Fig. 2: Altered cortico-hypothalamic projection pattern in mSOD1 mice at P95, contralateral hemisphere. A: Sum of neurons projecting from selected 28 areas. No difference is detected in the summarized number of neurons projecting to LHA in WT and mSOD1 from the contralateral hemisphere (n=3). B: Quantification of the absolute number of neurons projecting to LHA from 28 brain structures in WT and mSOD1. A significant increase in projections from ACA (p=0.0008), PL+ORBm and ILA (both p<0.0001) is detected. C: No difference is detected in the quantification of the normalized number of neurons projecting to LHA from 28 brain structures in WT and mSOD1. Bars show mean ± SD. *p < 0.05, **p < 0.01, ***p < 0.001, ****p< 0.0001.

Supplementary Fig. 3: Primary motor cortex and somatosensory cortex atrophy in P95 mSOD mice. Volumetric comparison of 28 areas in WT and mSOD. Significant atrophy was detected in MOp and SS. Bars show mean ± SD. *p < 0.05, **p < 0.01, ***p < 0.001, ****p< 0.0001.

Supplementary Fig. 4: Altered cortico-hypothalamic projection pattern in manual registered mSOD1 mice at P95, ipsilateral. A: Left: representative injection sites in LHA for WT and mSOD1 mice. White outlines represent LHA boundaries. Middle and right: Representative frontal brain sections of WT and mSOD1 mice depicting projections from ORBl/vl+AI (significantly increased, arrow), PL+ORBm, ILA, ACA, MOs and MOp to LHA. B: Sum of neurons projecting from selected 28 areas. No difference is detected in the summarized number of neurons projecting to LHA in WT and mSOD1 from the ipsilateral hemisphere (n=3). C: Quantification of the absolute number of neurons projecting to LHA from 28 brain structures in WT and mSOD1. A significant increase

in projections from ORBl/vl+AI (p=0.035) is detected. D: Quantification of the normalized number of neurons projecting to LHA from 28 brain structures in WT and mSOD1. A significant increase in projections from ILA (p=0.001) , ORBl/vl+AI (p<0.0001) and PIR (p<0.0001) is detected. Bars show mean ± SD. Scalebars 1 mm. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p< 0.0001$.

Supplementary Fig. 5: Altered projection pattern in manual registered mSOD1 mice at P95, contralateral. A: Sum of neurons projecting from selected 28 areas. No difference is detected in the summarized number of neurons projecting to LHA in WT and mSOD1 from the contralateral hemisphere (n=3). B: No difference is detected in the quantification of the absolute number of neurons projecting to LHA from 28 brain structures in WT and mSOD1. C: Quantification of the normalized number of neurons projecting to LHA from 28 brain structures in WT and mSOD1. A significant increase in projections from PL+ORBm (p<0.0001) is detected. Bars show mean ± SD. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p< 0.0001$.

Supplementary Fig. 6: Unaltered cortico-hypothalamic projection pattern in mSOD1 mice at P25, ipsilateral. A: Left: representative injection sites in LHA for WT and mSOD1 mice. White outlines represent LHA boundaries. Middle and right: Representative frontal brain sections of WT and mSOD1 mice depicting a similar pattern of neurons. B: Sum of neurons projecting from selected 28 areas. No difference is detected in the summarized number of neurons projecting to LHA in WT and mSOD1 from the ipsilateral hemisphere (n=3). C: No difference is detected in the quantification of the absolute number of neurons projecting to LHA from 28 brain structures in WT and mSOD1. D: No difference is detected in the quantification of the normalized number of neurons projecting to LHA from 28 brain structures in WT and mSOD1. Bars show mean ± SD. Scalebars 1 mm.

Supplementary Fig. 7: Unaltered cortico-hypothalamic projection pattern in mSOD1 mice at P25, contralateral hemisphere. A: Sum of neurons projecting from selected 28 areas. No difference is detected in the summarized number of neurons projecting to LHA in WT and mSOD1 from the contralateral hemisphere (n=3). B: Quantification of the absolute number of neurons projecting to LHA from 28 brain structures in WT and mSOD1. A significant increase in projections from ACA (p=0.0078) is detected. C: Quantification of the normalized number of neurons projecting to LHA from 28 brain structures in WT and mSOD1. No difference is detected in the quantification of the number of neurons projecting to LHA from 28 brain structures in WT and mSOD1. D: Representative *WholeBrain* volume reconstructions of neurons projecting to LHA in WT and mSOD1 mice show a similar pattern of projections in MOs (blue), ORBl/vl+AI (yellow) and PL+AI (green). Bars show mean ± SD. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p< 0.0001$.

Supplementary Fig. 8: Altered cortico-hypothalamic projection pattern in kiFUS mice at P270, absolute counts ipsilateral. A: Sum of neurons projecting from selected 28 areas. No difference is detected in the summarized number of neurons projecting to LHA in WT and kiFUS from the ipsilateral hemisphere (n=4). B: Quantification of the absolute number of neurons projecting to LHA from 28 brain structures in WT and mSOD1. A significant increase in projections from ACA

1051   (p=0.0385) and MOs (p<0.0001) is detected. C: Representative *WholeBrain* ipsilateral side view
1052   reconstructions of neurons projecting to LHA in WT and mSOD1 mice. Decrease of projections
1053   from MOs are visible (arrows). D: Representative *WholeBrain* cortical top view reconstructions
1054   of neurons projecting to LHA in WT and mSOD1 mice. Decrease of projections from MOs are
1055   visible (arrows). Bars show mean ± SD. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.
1056

Supplementary Fig. 9: Altered cortico-hypothalamic projection pattern in kiFUS mice at P270,
contralateral hemisphere. A: Sum of neurons projecting from selected 28 areas. No difference is
detected in the summarized number of neurons projecting to LHA in WT and kiFUS from the
contralateral hemisphere (n=3). B: Quantification of the absolute number of neurons projecting
to LHA from 28 brain structures in WT and mSOD1. A significant increase in projections from
ACA (p=0.0121)   and ILA (p=0.0474) is detected. C: Quantification of the normalized number of
neurons projecting to LHA from 28 brain structures in WT and kiFUS. A significant decrease in
projections from ACA (p=0.0137) is detected. Bars show mean ± SD. *$p < 0.05$, **$p < 0.01$, ***p
< 0.001, ****$p< 0.0001$.
1066

Abbreviations

ABA = Allen brain atlas, version 2011
ACA = Anterior cingulate area
aco = anterior commissure olfactory limb
act = anterior commissure temporal limb
AI = Agranular insular area
ALS = Amyotrophic lateral sclerosis
ALS-FRS-R = ALS functional rating scale revised
AOB = Accessory olfactory bulb
AP = Anteroposterior
AUD = Auditory areas
BLA = Basolateral amygdalar nucleus
BMA = Basolateral amygdalar nucleus
BSA = Bovine serum albumin
CA3so = Field CA3 stratum oriens
ccg = genu of corpus callosum
CLA = Claustrum
COA = Cortical amygdalar area
DG = Dentate gyrus
DP = Dorsal peduncular area
DPBS = Dulbecco's phosphate buffered saline
DTI = Diffusion-tensor-imaging
DV = Dorsoventral
EP = Endopiriform nucleus
fa = corpus callosum anterior forceps
FA = Fractional anisotropy

FUS = B6.FUS$^{\Delta NLS/+}$

HSB = Hue-saturation-brightness color model

GU = Gustatory areas

HY = Hypothalamus

ILA = Infralimbic area

int = internal capsule

kiFUS = FUS^{-}/ZsGreen$^{+/-}$

LA = Lateral amygdalar nucleus

LHA = Lateral hypothalamic area

ME = Median eminence

ML = Mediolateral

MM = Medial mammillary nucleus

MOp = Primary motor area

MOs = Secondary motor area

MRI = Magnetic resonance imaging

mSOD1 = mSOD^{+}/ZsGreen$^{+/-}$

mSOD = B6SJL73 Tg(SOD1*G93A)1Gur/J (high-copy)

NLOT = Nucleus of the lateral olfactory tract

OCT = Optimal cutting temperature compound

opt = optic tract

ORBl/vl = Orbital area, lateral/ventrolateral part

ORBl/vl+ AI = Group of Orbital area, lateral/ventrolateral part and Agranular insular area

ORBm = Orbital area, medial part

OT = Olfactory tubercle

P110 = Post-natal day 110

PA = Posterior amygdalar nucleus

PAA = Piriform-amygdalar area

PBS = Phosphate buffered saline

PFA = Paraformaldehyde

PIR = Piriform area

PL = Prelimbic area

PL+ ORBm = Group of Prelimbic area and Orbital area, medial part

PTL = Posterior parietal association areas

rAAV = recombinant adeno-associated virus

rAAV2-retro = retrograde recombinant adeno-associated virus 2

RAM = Random-access memory

RGB = Red-green-blue color model

Rhinal CTX = Group of Perirhinal area, Ectorhinal area and Entorhinal area, lateral part

ROI = Region of Interest

RSP = Retrosplenial area

rt = room temperature

SCH = Suprachiasmatic nucleus

SS = Somatosensory areas

Tea = Temporal association areas

TFAS = Tractwise fractional anisotropy statistics
tiff = Tagged image file format
TOI = Tract-of-interest
TT = Taenia tecta
VIS = Visual areas
VISC = Visceral area

WT vs mSOD1 P95