Stool serology: development of a non-invasive immunological method for the detection of

Enterovirus-specific antibodies in Congo gorilla feces

Youssouf Sereme[1,2], Sandra Madariaga Zarza[1,2], Hacène Medkour[1,2], Inestin Amona[1,2,3,4]

Florence Fenollar[1,5], Jean Akiana[3], Soraya Mezouar[1,2], Nicolas Orain[1,2], Joana Vitte[1,2], Bernard

Davoust[1,2], Didier Raoult[1,2] and Oleg Mediannikov[1,2*]

[1]IHU-Méditerranée Infection, Marseille, France

[2]Aix-Marseille Univ, IRD, AP-HM, MEPHI, Marseille, France

[3]Laboratoire national de santé publique, Brazzaville, Congo

[4]Faculté des Sciences et Techniques, Université Marien NGOUABI, Brazzaville, Congo

[5]Aix-Marseille Univ, IRD, AP-HM, SSA, VITROME, Marseille, France

[*]**Corresponding author**

Address: Institut Hospitalo-Universitaire Méditerranée-Infection, 19-21 Boulevard Jean Moulin,

13005 Marseille Cedex 05, France

Email: olegusss1@gmail.com

Word abstract count: 198

Word text count: 3,241

Figures*: 7*

Keywords: Gorilla, feces, immunoglobulins specifics, poliovirus, enterovirus.

Abstract

The incidence of poliovirus has been significantly reduced by 99.9% globally meanwhile vaccine-associated paralytic poliomyelitis emerges. Recently, a new recombinant virus (Enterovirus C/Poliovirus) was identified in humans working as eco-guards and gorillas in Democratic Republic of Congo, including one gorilla with polio-like sequeler. A strain of this recombinant virus (Ibou002) was also isolated from feces. In order to assess the potential role of poliovirus infection, we have developed and optimized a protocol, based on the lyophilization and solubilization of stool in a small volume, for the detection of specific antibodies. A high level of total immunoglobulins was first detected in the concentrated stool extracts. Then, specific antibodies were detected in 4/16 gorilla samples and 2/3 human samples by western blot using both poliovaccine antigen and Ibou002 antigen and by ELISA using polio vaccine antigen. The humoral responses were greater with the Ibou002 antigen both in number and intensity of bands. Thus, we suggest that this recombinant virus could cause polio-like disease in the endangered western lowland gorilla. The development of a non-invasive method for the detection of microorganism-specific immunoglobulins from fecal samples opens up new perspectives for the exploration of humoral responses of pathogens in animals and knowledge in zoonotic infectious diseases.

INTRODUCTION

Poliovirus is a virus belonging to the Picornaviridae family and the *Enterovirus* genus (Enterovirus C), which causes poliomyelitis in humans[1]. The poliovirus genome is a 7.5 kilobase single-stranded RNA surrounded by a capsid, which is composed by four proteins (VP1, VP2, VP3, and VP4)[2]. There are three poliovirus types (1, 2, and 3); the immunity to one type would not protect against the two others[3].

As a result of the global polio eradication program, initiated in 1988 with the use of live-attenuated oral poliovirus vaccine, the number of worldwide wild poliovirus cases has declined by more than 99.9%, from an estimated 350,000 cases in more than 125 endemic countries to 175 wild cases in 2019 (https://www.who.int/health-topics/poliomyelitis#tab=tab_1). Wild poliovirus types 2 and 3 were certified eradicated in 2015 and 2019, respectively[4,5]. The lack of circulation of wild poliovirus type 1 in Nigeria has been reported since 2016[14,15]. On 25 August 2020, the WHO African region was certified as free of wild poliovirus, leaving only the Eastern Mediterranean Region with endemic circulation of wild poliovirus type 1 in 2 countries (https://www.who.int/wer/2020/wer9541/en/)[6,11–13]. In parallel, another challenge arose with the emergence of vaccine-associated paralytic poliomyelitis[6]. Indeed, in the past 12 months, a total of 132 wild virus cases were reported in Afghanistan (61) and Pakistan (142), (http://polioeradication.org/polio-today/polio-now/this-week), while 605 cases of vaccine-associated paralytic poliomyelitis were detected worldwide, including in 19 African countries (406 cases), Afghanistan (101), Pakistan (80), Malaysia (1), Philippines (1), and Yemen (16). To note, among these 605 vaccine-associated cases, 424 originated from non-endemic countries and 181 from known endemic countries[6–10]. It has been also reported that 5-10% of polio deaths worldwide are due to cVDPVP[6].

The transmission of Poliovirus is fecal-oral. Humans remain the natural host, but it can also infect primates and monkeys[1,3]. There is a positive correlation between zoonotic diseases

and the emergence of infectious diseases in humans[16]. It has been shown under experimental

conditions that hominoids and monkeys can be infected with poliovirus[17,18]. A 2005 study in

Madagascar demonstrated the circulation of vaccine-derived recombinant poliovirus strains,

which are recombinant Enterovirus-C viruses, and their ability to give "polio-like" in humans[19].

A study conducted by Harvala *et al.* in Cameroon and the Democratic Republic of Congo

(DRC) screening for enteroviruses by PCR on chimpanzee and gorilla fecal samples

demonstrated both the circulation of genetically divergent variants of enteroviruses in apes and

monkeys as well as a strain (EV-A89) that they share with local human populations[20]. In

addition, *Enterovirus* species described in humans and other species as a potential source of

emerging infectious diseases in humans have been detected in monkeys[21,22].

Previously, our team reported the presence of a new Enterovirus C with a high similarity

to Coxsackievirus as well as similarity with poliovirus 1 and 2 and associated with flaccid

paralysis in chimpanzee virus EV-C99 strains[21] in the 3C, 3D, and 5'UTR regions[23]. Thus, we

hypothesized a suspicion of poliovirus infection in a gorilla male living alone in Lésio-Louna-

Léfini Nature Reserve (Republic of the Congo) based on clinical observations including facial

paralysis. Our aim was to develop and to optimize a non-invasive experimental method to

assess our hypothesis by the use of gorilla fecal samples to identify specific antiviral antibodies.

MATERIALS AND METHODS

Collection of gorilla feces samples

Fresh fecal samples were collected from western lowland gorillas (*Gorilla gorilla gorilla*) in

semi-captivity and in wild in the Lésio-Louna-Léfini Gorilla Nature Reserve, located in the

northern Pool Department, Republic of Congo. A total of 16 gorilla fecal samples were

collected. Samples were collected from Iboubikro (Lat: 03.27040; Long 15.47071) for semi-

captive gorilla youngsters and from Abio2 (Lat 03.094; Long 15.52633) for wild gorillas.

Samples were collected in individual pots (Labelians, Nemours, France) and 50 ml Falcon tubes

(Dutscher, Brumath, France). At each collection time and for each sample, information relating

to the GPS position of the place, the estimated time of decomposition of the feces as well as the

morphological and/or physical aspect was recorded. During sample collection, we observed a

male named Sid living alone on an island in front of Abio resort of Lésio-Louna-Léfini Nature

Reserve, where he was placed after being sick. We observed evident signs of facial

myodystrophy and paresis and upper limbs paresis (**Figure 1**). Three fecal samples were

collected from 3 people working on the site (eco-gardian); an anonymous identification code

was given to each sample (Ibou1, Ibou2, and Ibou3). A poliovirus serology-positive control

serum (Poliomyelitis virus kit, GenWay, San Diego, California, USA) was also used as a

control. Research and administrative authorizations for the collection of samples were granted

by the Congolese Ministries of Scientific Research and Technological Innovation

(N°003/MRSIT/DGRST/DMAST), Forest Economy and Sustainable Development represented

by the Congolese Agency for Wildlife and Protected Areas (N°0134/ACFAP-DTS) and

(N°94/MEFDD/CAB/DGACFAP-DTS) as well as the Ministry of Health and Population

(N°000220/MSP-CAB-17 and N°208/MSP/CAB.15).

Concept and optimization of the methodological approach

The principle of our approach consisted initially in developing a method of extracting

immunoglobulins which makes it possible to obtain structural and functional proteins for

humoral evaluation. The extraction buffer contained protease inhibitor cocktail to inhibit the

general degradation of proteins from stool extracts, tissues or cells by endogenous or exogenous

microorganisms' proteases and to study processes that involve blocking the activity of specific

proteases. Three gorilla and 19 human stool samples were used to optimize the concept.

Different stool treatment protocols have been tested for the detection of immunoglobulins (IgG

and IgA), mainly: (1) Filtration of stools extract from 2 g of stools plus 2 ml of buffer; (2)

Lyophilization of the filtered extract, reconstituted in 500 µl of buffer and then purification of

IgG and IgA in the reconstituted extract with protein G and pectin M; (3) Lyophilization of the

filtered extract, reconstituted in a small volume of buffer (500 µl) then concentrated by the

Amicon® Ultra-1. After the evaluation of the data, a final protocol was adopted (**Figure 2**).

Experimental protocol adopted for stool protein extraction

Two grams of each sample were solubilized in 2 µl of an extraction buffer consisting of

phosphate buffered saline solution supplemented with 4-(2-aminoethyl)-benzenesulfonyl

fluoride 4 mM, bestatin 0.26 mM, E-64 28 µM, leupeptin 2 µM and aprotinin 0.6 µM, pH 7.4,

and a protease inhibitor "SIGMAFAST™ Protease Inhibitor Tablets" (Sigma-Aldrich, Saint-

Louis, Missouri, USA). The stool-buffer mixture was incubated for 15 min at room temperature,

vortexed for 1 min, and then centrifuged at 1,500 rpm for 15 min at 4°C. The supernatant was

lyophilized for 24 h and then re-solubilized in 1 ml of distilled water. Then, 500 µl of re-

solubilized stool was concentrated using Amicon® Ultra-15 3k centrifugal filters (Merck, Saint-

Romain, France) for further studies (**Figure 2**). Concentrations of total immunoglobulins (IgA

and IgG) were evaluated by ELISA (Abcam, Paris, France) to validate the method of protein

extraction from stools (**Figure 2**).

Quantification of anti-poliovirus immunoglobulins by ELISA

Determination of specific immunoglobulins was performed (IgG and IgA) by ELISA according

the manufacturer's instructions (Poliomyelitis virus kit, GenWay, San Diego, California, USA).

The polio antigen derived from the human pathogenic poliovirus types, specifically type 1

(Brunhilde), type 2 (Lansing), and type 3 (Leon) was bound in the wells of the plate. The

diluted samples or standards were then deposited. After 1-hour incubation at room temperature,

the plate was rinsed with wash buffer. The anti-human IgG peroxidase conjugate was then

added and incubated for 30 min, followed by a wash step. Afterwards, the substrate solution

was added and incubated for 20 min. The reaction was stopped by addition of a stop solution.

The resulting dye was measured using a Thermo Fisher spectrophotometer (Uppsala, Sweden)

at a wavelength of 450 nm.

Characterization of enterovirus-specific antibodies by western blot from fecal samples

Preparation of antigen from Polio vaccine. Imovax Polio vaccine (Sanofi, Antony, France)

was used as an antigen source for the characterization of poliovirus-specific immunoglobulins

by western blot. The vaccine is a sterile suspension of polioviruses types 1 (Mahoney), 2

(MEF1), and 3 (Saukett), which are grown on Vero cells, purified, and then inactivated with

formaldehyde. Four doses of Imovax Polio were treated with the TS lysis buffer and

fractionated by sonication to release the antigen molecules. The antigen suspension was

lyophilized and re-solubilized in 100 µl of distilled water (**Figure 3**). The concentrated antigen

was measured by Bradford protein assay (Bio-Rad, Hemel Hempstead, UK), and a SDS-PAGE

was performed with 50 µg of the purified antigen and 12% of polyacrylamide.

Preparation of antigen from enterovirus strain Ibou002. The strain Ibou002[23] of recombinant

enterovirus C originally isolated from a eco-guardian of Lésio-Louna Nature Reserve was

replicated during 7 days in MRC5 cells line (ATCC® CCL-171™). The virus infected MRC5

cells were collected and centrifuged at 700 × g for 15 min (Eppendorf Centrifuge 5810R).

Supernatant was filtered by serial filtration steps to eliminate cells and cellular debris, going

throw 0.8 µm, 0.45 µm, and 0.2 µm filters. Then, virus particles were precipitated in 10% PEG-

8000 and 2.2% NaCl at 4ºC overnight, in agitation. The precipitates were centrifuged at 10,000

× g (Sorvall Evolution, SLA-3000 Recent-1 rotor, Thermo Fisher) for 30 min at 4ºC, and the

virus pellet was resuspended in HEPES saline solution (9 g NaCl, 10 mL HEPES 1M, 990 mL

distilled H_2O). The resuspended virus was purified by a 30% (wt/vol) sucrose cushion at

100,000 × g (Sorvall Discovery 90SE, Thermo Fisher) for 2 hours and 30 min at 4°C. The virus

pellet was washed in HBSS (Thermo Fisher), and centrifuged at 100,000 × g (Beckman SW 32

Ti rotor) for 30 min at 4°C. The viral particles were finally resuspended in HEPES saline and

then fractionated with TS buffer (7 M Urea, 2 M Thiourea, 4% CHAPS) to release the antigen.

The released antigen was then concentrated with the Amicon 3 kDa filter (Merck KGaA,

Darmstadt, Germany) before being analyzed by SDS gel electrophoresis, and the western blot.

The concentration of the antigen was measured. Then, 30 µg of purified were loaded onto 12%

SDS-polyacrylamide gel for analysis.

Western blot. Western blot was performed using Jess™ Simple Western system (Protein

Simple, San Jose, California, USA) according to the manufacturer's protocol[24]. This technique

is an automated capillary size separation and nano-immunoassay system for performing gel-

free, blot-free, and hands-free capillary-based immunoassay that integrates and automates the

entire protein separation and detection process with home-made antigens. To quantify the

absolute response to viral proteins from samples, we followed the manufacturer's standard

method for 12-230 kDa Jess™ separation module (SM-W004). First, poliovirus antigen (1

µg/µL) was mixed with 5X master mix (Protein Simple) to achieve a final concentration of 0.2

µg/µL in the presence of fluorescent molecular weight markers and 400 mM dithiothreitol

(DTT, Protein Simple). Then, this preparation was denatured at 95°C for 5 min. The ladder (12-

230 kDa) and the viral proteins were separated in capillaries as they migrate through a

separation matrix at 375 volts. A protein simple proprietary photoactivated capture chemistry

was used to immobilize separated viral proteins on the capillaries. Patient feces extract was

added and incubated for 30 min. After a wash step, goat HRP-conjugated anti-human

IgG/IgM/IgA antibodies (Jackson ImmunoResearch, Ely, UK) diluted 1:250 was added for 30

min. Finally, the chemiluminescent revelations of ladder and samples were established with

peroxide/luminol-S (Protein Simple). Digital image was analyzed with Compass for SW

software (v 4.1.0, Protein Simple), and quantified data of the detected proteins was reported as

the molecular weight, chemiluminescence intensity, and signal/noise ratio. Western blot

analyses were performed according this protocol using as antigens (1) the polio vaccine and (2)

the purified Ibou002 virus strain.

RESULTS

Differentiation of the feces of gorilla individuals

A previous study by our team had confirmed that each feces sample corresponded to an

individual gorilla[23]. A code was given to each of the 16 gorillas. Twelve had a G0 code

followed by a numbering from 1 to 12 and 4 had a GNLL code followed by a numbering from 1

to 4.

Selection of stool preparation protocol

Using the protocol 1 based on raw stool extracts, very low concentrations of immunoglobulins

were detected (**Supplementary figure 1**). Using the protocol 2 based on purification with

protein G and peptin M, we also noticed a loss of immunoglobulin concentrations after

purification (Data not shown). Protocol 3 based on the lyophilization of the filtered stool extract

followed by its reconstitution in a small volume allowed to obtain the best results as shown in.

In gorilla stool, the concentration mean with range of 222 (121.85-320.5) ng/ml was observed

for IgA and 123.12 (25.87-208.9) ng/ml for IgG. In human feces, the concentration mean with

range of 1600.55 (500-2978.8) ng/ml was observed for IgA and 619.7 (453.67-870.58) ng/ml

for IgG (**Figure 4**). It is therefore the latter that was selected to continue analyzes on the stool.

Quantification and detection of poliovirus-specific immunoglobulins by ELISA and

Western Blot

Quantification of poliovirus-specific immunoglobulins (IgA-IgG) by ELISA

The poliovirus-IgG/IgA evaluation of 16 gorilla and 3 human stool samples was carried

out by ELISA; IgG/IgA level was represented by the OD at 450 nm. A sample was concluded as

positive for IgG / IgA when the OD value was greater than 2 times the negative control value

for the kit used (OD positive > 0.35). Specific antibodies were detected in 4 of 16 (25%) gorilla

samples (G01, G02, G03, and G07) and 2 of 3 human fecal samples (Ibou2 and Ibou3) (**Figure**

5). Besides the recombinant Enterovirus C (strain Ibou002) was isolated from one of the two

eco-guardians (Ibou3) positive to poliovirus[22].

Detection of poliovirus- and enterovirus C-specific immunoglobulins by western blot

Protein profiles of both antigens (Imovax vaccine and strain Ibou002)

The protein profile was performed for both antigens (Imovax vaccine and strain Ibou002).

Different protein profiles were identified for each antigen by SDS-PAGE. The vaccine

displayed five bands possibly indicating five proteins with molecular weights ranging from 20-

35 kDa. We found seven bands in the antigen purified from virus, which correspond to a

molecular weight from 28 to 130 kDa (Figure 6A and B).

Western blot

The immunoglobulin evaluation by western blot showed positive samples for both

antigens source (poliovirus vaccine and Ibou002 virus strain), but with different bands profiles

between them. Bands were detected in 4 samples (G01, G02, G03, and G07) from 16 samples of

gorilla feces, and 2 samples (Ibou2 and Ibou3) from 3 human feces, which were previously

detected positive by ELISA. We also noted that all samples with bands from the vaccine

antigen, also showed bands for the virus isolated, but at different molecular weight between

antigen source (14 kDa for vaccine antigen, and 60 kDa and 116 kDa for antigen Ibou002 virus

strain).

DISCUSSION

In this study, we developed a non-invasive method for screening animal immune

responses using fresh stool samples. We also provided a proof of concept on the feasibility of

exploring immunity from feces. Indeed, the detection of the total antibody levels (IgA / IgG) in

concentrated stool extracts via the developed protocol confirms the possibility of using these

samples for the detection, quantification and characterization of specific antibodies by ELISA

and western blot. The very high concentrations of IgA in feces can be explained by the

microbial colonization of the gut that triggers the production of IgA by the gut associated

lymphoid tissue (GALT)[25]. This supports our choice to use an anti-IgA, IgG mixture in our

ELISA and western blot techniques to detect the maximum amount of specific antibodies. In

addition, we did not look for specific IgM because the concentration of total IgM in the

digestive mucosa is very low or even zero[26].

By testing a working hypothesis that circulating (probably vaccine) poliovirus[19] caused

poliomyelitis-like disease in gorillas, we attempted to identify specific anti-poliovirus

antibodies in the stools of 16 gorillas and 3 eco-guardians of the reserve. Using our original

method of antibodies extraction, we detected antibodies binding to poliovirus antigen in fecal

samples by ELISA[27], allowed us to identify positive samples among 4 of 16 gorillas and 2 of 3

eco-guardians. Such results in humans were not surprising as the majority of the population is

vaccinated against poliomyelitis in this area. However, the positive results in 4 gorillas

including Sid, the one with sequelae, were more surprising, indicating a potential circulation of

the poliovirus or a close Enterovirus in this community. Then, we performed a western blot by

Jess[TM] to characterize the specific antibodies. Western blot analysis of antibodies from gorilla

feces detected a single band of around 14 kDa using the antigens extracted from poliovirus

vaccine. These data do not allow confirmation of immunization of the gorilla with poliovirus

due to the absence of antibodies against the main proteins of the poliovirus, namely VP1, VP2,

and VP3. In addition, PCR testing for poliovirus RNA was negative[22].

We then based on the circulation of recombinant Enterovirus C[22] having at least 3 genes

very close to the poliovirus in this geographical area to hypothesize that the gorilla Sid may

suffer from the sequelae of polio-like illness caused by this virus. Indeed, these polio-like

diseases can be caused in humans by chimeric/recombinant viruses[19–21]. These diseases often

produce the typical sequelae in the form of flaccid paralysis and muscle atrophy. In order to

verify this hypothesis, we prepared the antigen from the recombinant Enterovirus C (Ibou002

strain) previously isolated from a stools sample of eco-guardian from the Lésio-Louna reserve.

We noticed that the 4 samples of gorilla feces that showed bands with the vaccine antigen were

the same samples that showed bands with the antigen from the recombinant Enterovirus C. This

concordance was also observed with the two human feces samples. Western blot analysis

allowed the detection of specific immunoglobulins to different viral proteins. Indeed, two 60

kDa and 110 kDa molecular weight bands were present in all positive samples from gorillas and

humans with the Ibou002 strain antigen, and one 14 kDa band was also present in all positive

samples with the vaccine antigen. Evaluation of the western blot results showed that the positive

specimens for the two antigen sources (polio vaccine and Ibou002 virus strain) were the same

but with different profiles. Thus, we hypothesize that the recombinant Enterovirus C that we

recently isolated from an eco-guardian may have wide circulation and it was probably

responsible for the immunization of humans and gorillas.

Given the presence of typical post-polio sequelae in the gorilla Sid, the presence of

specific antibodies binding to the Ibou002 virus strain and the recombinant nature of this virus,

we suggest that this new virus could cause polio-like disease in the endangered western lowland

gorilla. We also believe that the detection of a band in western blot with the poliovirus antigen

is the result of a non-specific cross-reaction between the antibodies induced by the recombinant

Enterovirus C strain Ibou002 and the genetically related poliovirus. Another interesting finding

is the detection of specific anti-recombinant Enterovirus C Ibou002 antibodies in the same stool

sample from the person from which the virus was isolated. This may indicate a long persistence

of the virus in the human organism and the absence of protective immunity.

CONCLUSION

Thanks to the non-invasive immunological developed method using feces, we were able

to assess the presence of specific antibodies against a recombinant Enterovirus C in gorillas and

eco-guardians from the same reserve. This method, which does not require medical or invasive

procedures for the collection of samples and allowing the detection of specific antibodies, opens

numerous prospects for application in zoonotic infectious diseases and can revolutionize the

screening of animals for emerging infections of importance, such as Ebola fever, rabies, and

coronavirus infections.

ACKNOWLEDGEMENTS

We thank Lucile Pinault and Rita Jaafar for their help and advice in the development of the experiments. This work was supported by the French Government under the "Investissements d'avenir" (Investments for the future) program managed by the "Agence Nationale de la Recherche" (reference: 10-IAHU-03). Research and administrative authorizations for the collection of samples were granted by the Congolese Ministries of Scientific Research and Technological Innovation (n°003/MRSIT/DGRST/DMAST), Forest Economy and Sustainable Development represented by the Congolese Agency for Wildlife and Protected Areas (N° 0134/ACFAP-DTS) and (N° 94/MEFDD/CAB/DGACFAP-DTS) as well as the Ministry of Health and Population (N°000220/MSP-CAB-17and N°208/MSP/CAB.15). YS, SZM, HM are supported by a "Fondation Méditerranée Infection" doctoral position. SM was supported by a post-doctoral "Fondation Méditerranée Infection" fellowship.

AUTHOR CONTRIBUTIONS

YS conducted the experiments, analyzed the data and wrote the paper; SMZ, HM, JV, NO and SM helped conduct the experiments and participated in the writing of the paper; JA, BD and IA contributed to the collection of samples; OM, DR, FF designed the project, participated in the writing of the paper and provided great support for the realization of the experiments.

ADDITIONAL INFORMATION

Competing interest statement

The authors declare no competing interests in relation to this study.

Figure legends.

Figure 1. Geographic map of the collection site and the gorilla named "Sid" with myodystrophy and paresis on the face that lives alone on an island in front of the Abio resort of the Lesio-Louna nature reserve (Republic of Congo).

Figure 2. Schematic representation of the three protocols evaluated for protein extraction from stools up to evaluation of immunoglobulins by ELISA and western blot. The protocol 3, written in black, is the one that was retained.

Figure 3. Protocol for the extraction of antigens from the polio vaccine (Imovax®) for analysis by western blot.

Figure 4. Evaluation of total IgA (A) and IgG (B) concentrations (ng/mL) in stools of 3 gorillas and 2 humans by ELISA.

Figure 5. Evaluation of poliovirus IgG/IgA by ELISA. Graph representing the specific IgG/IgA (OD) level of all samples, measured at a wavelength of 450 nm. A sample was concluded as positive for IgG / IgA when the OD value was greater than 2 times the negative control value for the kit used (OD positive > 0.35).

Figure 6. Silver stain SDS-PAGE analyses for poliovirus vaccine (A) and antigen of Ibou002 virus strain (B) proteins. In both profiles, the left lane corresponds to the molecular weight standard and the right lane to the antigen; 50 µg of poliovirus vaccine and 30 µg of antigen of Ibou004 virus strain were running in a 12% polyacrylamide gel.

Figure 7. Western blot Jess^{TM} using two different antigens, poliovirus vaccine (A) and Ibou002 virus strain (B). Anti-human IgG/IgM/IgA antibodies detection was done for 16 gorilla stools, 3 human stools, and 1 control positive serum (non-diluted and diluted 1:100).

References

1.   Fields virology. (Wolters Kluwer Health/Lippincott Williams & Wilkins, Philadelphia, 2007).

2.   Blondel, B. *et al.* [Genetic evolution of poliovirus: success and difficulties in the eradication of paralytic poliomyelitis]. *Med Trop (Mars)* **68**, 189–202 (2008).

3.   Pelletier, J. & Sonenberg, N. Internal initiation of translation of eukaryotic mRNA directed by a sequence derived from poliovirus RNA. *Nature* **334**, 320–325 (1988).

4.   Khan, F. *et al.* Progress Toward Polio Eradication - Worldwide, January 2016-March 2018. *MMWR Morb. Mortal. Wkly. Rep.* **67**, 524–528 (2018).

5.   Weekly epidemiological record: Progress towards poliomyelitis eradication. *World Health Organization (WHO)* (2019).

6.   Javelle, E. & Raoult, D. Antibiotics against poliovirus carriage: an additional tool in the polio endgame? *Clinical Microbiology and Infection* **26**, 542–544 (2020).

7.   Mbaeyi, C. *et al.* Update on Vaccine-Derived Poliovirus Outbreaks — Democratic Republic of the Congo and Horn of Africa, 2017–2018. *MMWR Morb. Mortal. Wkly. Rep.* **68**, 225–230 (2019).

8.   Saleem, A. F. *et al.* Immunogenicity of Different Routine Poliovirus Vaccination Schedules: A Randomized, Controlled Trial in Karachi, Pakistan. *The Journal of Infectious Diseases* **217**, 443–450 (2018).

9.   Chandler, J. Fighting a polio outbreak in Papua New Guinea. *The Lancet* **392**, 2155–2156 (2018).

10.   Roberts, L. Polio eradication campaign loses ground. *Science* **365**, 106–107 (2019).

11.   Akhtar, R. *et al.* Genetic Epidemiology Reveals 3 Chronic Reservoir Areas With Recurrent Population Mobility Challenging Poliovirus Eradication in Pakistan. *Clinical Infectious Diseases* **71**, e58–e67 (2020).

12.   Kalkowska, D. A., Pallansch, M. A. & Thompson, K. M. Updated modelling of the prevalence of immunodeficiency-associated long-term vaccine-derived poliovirus (iVDPV) excreters. *Epidemiol. Infect.* **147**, e295 (2019).

13.   Martinez, M. *et al.* Progress Toward Poliomyelitis Eradication — Afghanistan, January 2019–July 2020. *MMWR Morb. Mortal. Wkly. Rep.* **69**, 1464–1468 (2020).

14.   Duintjer Tebbens, R. J. & Thompson, K. M. Polio endgame risks and the possibility of restarting the use of oral poliovirus vaccine. *Expert Review of Vaccines* **17**, 739–751 (2018).

15.   Thompson, K. M. & Duintjer Tebbens, R. J. Lessons From the Polio Endgame: Overcoming the Failure to Vaccinate and the Role of Subpopulations in Maintaining Transmission. *The Journal of Infectious Diseases* **216**, S176–S182 (2017).

16.   McArthur, D. B. Emerging Infectious Diseases. *Nurs. Clin. North Am.* **54**, 297–311 (2019).

17.   Hsiung, G. D. Diagnostic virology: from animals to automation. *Yale J Biol Med* **57**, 727–733 (1984).

18. Ida-Hosonuma, M. *et al.* Host range of poliovirus is restricted to simians because of a
rapid sequence change of the poliovirus receptor gene during evolution. *Archives of Virology*
148, 29–44 (2003).

19. Rakoto-Andrianarivelo, M. *et al.* Reemergence of Recombinant Vaccine-Derived
Poliovirus Outbreak in Madagascar. *J INFECT DIS* **197**, 1427–1435 (2008).

20. Harvala, H. *et al.* Co-circulation of enteroviruses between apes and humans. *J. Gen.*
Virol. **95**, 403–407 (2014).

21. Mombo, I. M. *et al.* First Detection of an Enterovirus C99 in a Captive Chimpanzee with
Acute Flaccid Paralysis, from the Tchimpounga Chimpanzee Rehabilitation Center, Republic of
Congo. *PLoS ONE* **10**, e0136700 (2015).

22. Mombo, I. M. *et al.* African Non-Human Primates Host Diverse Enteroviruses. *PLoS*
ONE **12**, e0169067 (2017).

23. Amona, I. et al. Isolation and Molecular Characterization of Enteroviruses from Humans
and Great Apes in the Republic of Congo: Recombination within Enterovirus C Serotypes.
Microorganisms. 2020, 8(11), 1779; https://doi.org/10.3390/microorganisms8111779

24. Wu, J., Haitjema, C. H., Heger, C. D. & Boge, A. A proof-of-concept analysis of
carbohydrate-deficient transferrin by imaged capillary isoelectric focusing and in-capillary
immunodetection. *BioTechniques* **68**, 85–90 (2020).

25. Suzuki, K., Ha, S., Tsuji, M. & Fagarasan, S. Intestinal IgA synthesis: A primitive form
of adaptive immunity that regulates microbial communities in the gut. *Seminars in Immunology*
19, 127–135 (2007).

26. Fadlallah, J. *et al.* Microbial ecology perturbation in human IgA deficiency. *Sci. Transl.*
Med. **10**, eaan1217 (2018).

27. Kummitha, C. M. *et al.* A sandwich ELISA for the detection of Wnt5a. *J. Immunol.*
Methods **352**, 38–44 (2010).

Figure 1. Geographic map of the collection site and the gorilla named "Sid" with myodystrophy and paresis on the face that lives alone on an island in front of the Abio resort of the Lesio-Louna nature reserve (Republic of Congo).

Figure 2. Schematic representation of the three protocols evaluated for protein extraction from stools up to evaluation of immunoglobulins by ELISA and western blot.

Figure 3. Protocol for the extraction of antigens from the polio vaccine (Imovax®) for analysis by western blot.

Figure 4. Evaluation of total IgA (A) and IgG (B) concentrations (ng/mL) in stools of 3 gorillas and 19 humans by ELISA.

Figure 5. Evaluation of poliovirus IgG/IgA by ELISA. Graph representing the specific IgG/IgA (OD) level of all samples, measured at a wavelength of 450 nm. A sample was concluded as positive for IgG / IgA when the OD value was greater than 2 times the negative control value for the kit used (OD positive > 0.35).

Figure 6. Silver stain SDS-PAGE analyses for poliovirus vaccine (A) and antigen of Ibou004 virus strain (B) proteins. In both profiles, the left lane corresponds to the molecular weight standard and the right lane to the antigen; 50 μg of poliovirus vaccine and 30 μg of antigen of Ibou004 virus strain were running in a 12% polyacrylamide gel.

Figure 7. Western blot JessTM using two different antigens, poliovirus vaccine (A) and Ibou004 virus strain (B). Anti-human IgG/IgM/IgA antibodies detection was done for 16 gorilla stools, 3 human stools, and 1 control positive serum (non-diluted and diluted 1:100).

B.

Sample of the gorilla Sid with
polios-like sequeler

Sample from which strain
Ibou002 was isolated

Ladder G01 G02 G03 G04 G05 G06 G07 G08 G09 G010 G011 G012 GNLL01 GNLL03 GNLL04 Ibou1 Ibou2 Ibou3 Low pos. 1/100 Negative Milk-Free dil.

kDa

230-
180-

116-

66-

40-

12-

Gorillas   Eco-guardians   Controls

Supplementary figure 1. Quantification of immunoglobulins (IgG and IgA) in gorilla feces with three different fecal treatment protocols (1) Filtration of stools extract from 2 g of stools plus 2 ml of buffer; (2) Lyophilization of the filtered extract, reconstituted in 500 μl of buffer and then purification of IgG and IgA in the reconstituted extract with protein G and pectin M; (3) Lyophilization of the filtered extract, reconstituted in a small volume of buffer (500 μl) then concentrated by the Amicon® Ultra-1