Multielectrode Cortical Stimulation Selectively Induces Unidirectional Wave Propagation

Alma Halgren[1], Zarek Siegel[1,2], Ryan Golden[1,2], Maxim Bazhenov[1]

[1]Department of Medicine, University of California, San Diego, La Jolla, California, United States of America

[2]Neurosciences Graduate Program, University of California, San Diego, La Jolla, California, United States of America

ABSTRACT

Cortical stimulation is emerging as an experimental tool in basic research and a promising therapy for a range of neuropsychiatric conditions. As multielectrode arrays enter clinical practice, the possibility of using spatiotemporal patterns of electrical stimulation to induce desired physiological patterns has become theoretically possible, but in practice can only be implemented by trial-and-error because of a lack of predictive models. Experimental evidence increasingly establishes travelling waves as fundamental to cortical information-processing, but we lack understanding how to control wave properties despite rapidly improving technologies. This study uses a hybrid biophysical-anatomical and neural-computational model to predict and understand how a simple pattern of cortical surface stimulation could induce directional traveling waves via asymmetric activation of inhibitory interneurons. It reveals local circuit mechanisms to control spatiotemporal cortical dynamics and predicts interventions that can be developed to treat a broad range of cognitive disorders.

INTRODUCTION

Brain stimulation is widely used in both experimental and clinical settings. In basic research it is used to probe neural function by disrupting or hyperactivating local brain processing[1–3]. In clinical settings, direct manipulation of activity via stimulation has also been shown to be an effective in treatment of several neural and psychiatric disorders. Deep brain stimulation (DBS) has been successful in the treatment of movement disorders like Parkinson's disease[4–6], depression[7,8] and obsessive-compulsive disorder[9,10].Superficial cortical stimulation is an effective therapy for epileptic[11] and stroke patients[12]. Increasingly, electrical stimulation has also shown promise in both the restoration and enhancement of critical cognitive functions such as memory. DBS has been shown experimentally to enhance memory encoding when applied during learning[13] and closed-loop stimulation protocols have proven to be effective during periods of poor memory encoding as well as during memory recall[14–16].

While brain stimulation is sometimes conceptualized as disrupting pathological activity in order to restore normal activity, increasingly the explicit goal is to directly generate normal activity. Experimental evidence supports travelling waves as critical to normal brain activity. These propagating waves are fundamental to brain information-processing as they coordinate neural behavior across all spatial scales, from within-layer to whole-brain interactions, as well as across temporal scales, from tens to hundreds of milliseconds. By mediating communication across multiple brain areas, propagating activity putatively performs a variety of cognitive functions such as long-term memory consolidation and the processing of visual stimuli[17]. For example, sleep spindles travelling across the cortical surface at multiple scales have been hypothesized to synchronize convergent co-firing, resulting in spike-timing dependent plasticity and consequent memory consolidation[19]. Similarly, the alpha rhythm which modulates visual processing appears to be a travelling wave from association to primary areas[18]. Accordingly, the ability to predict and control traveling waves has far-reaching implications for improving and controlling cognitive function.

Currently, there is no method of reliably generating directional traveling waves with electrical stimulation. Past efforts to develop new paradigms of stimulation which reinstate particular brain activity states have largely depended upon trial and error. Recently, we described a method for modeling the effects of cortical stimulation that enables one to predict the consequences of stimulation in silico, and thereby develop stimulation protocols that achieve desired results in vivo[20]. This earlier study was limited to the effects of stimulation of a single electrode and thus did not evoke directional propagation. In this new study, we describe an initial attempt to model a multielectrode stimulation paradigm that produces unidirectional traveling waves in the cortex. With multielectrode arrays increasingly entering clinical practice[21], our model harnesses the additional complexity and control during stimulation that multielectrode protocols allow for.

The modeling approach includes two phases. In the first phase, biophysical model is used to predict spiking probability in response to a spatially-varied electric field potential in reconstructed rat somatosensory cortical neurons obtained from NeuroMorpho.org[22]. We found

that the hyperpolarization or depolarization of individual neurons varied according to cell type
and cortical depth, and also varied with respect to the polarity of the applied electric field. The
diversity in excitation response underlies the propagating wave activity that we observed in the
second phase of the model. In the second phase, we constructed a Hodgkin-Huxley model of
cortical network composed of multiple interconnected cortical columns, each containing a circuit
of inhibitory and excitatory cells across cortical layers. Approximating stimulation effects using
the activation probabilities calculated in phase one, we found that fast inhibitory activity,
coupled with excitatory cells' preference for cathodal stimulation, resulted in a unidirectionally
propagating wave of activity. These results develop the previously published model for
application to a new and potentially clinically-relevant domain, suggesting a simple multi-
electrode pattern for evoking traveling waves and providing testable predictions for experimental
confirmation and parameter optimization.

RESULTS

Building on existing *in silico* models that simulated the effects of cortical surface

stimulation from a single electrode[23], we sought to explore the spatial dynamics of stimulation by

modeling an asymmetrical three-electrode configuration (Fig 1a). The paper is organized as

follows. We first calculated electrical field created by the system of three electrodes. Next, we

estimated the activation probability for each cell type and cortical layer by computing the

activating function in biophysical reconstructions of axonal arbors. We then constructed a

cortical microcircuit model with Hodgkin-Huxley dynamics to model the network effects of

stimulation based on the previously calculated spiking probabilities.

Figure 1

Electric potential and activating function in the plane Y=0.

1a. Schematic of electrode configuration and electric potential in the plane Y=0. Each electrode
is square-shaped and measures 150 μm by 150 μm, and the spacing between each electrode is
also 150 μm. The two leftmost electrodes are negative and deliver negative 75 μA of current
each, while the rightmost electrode is positive and delivers 150 μA of current, and the
stimulation period is 200 μs. Layer depths are approximated from experimental measurements of
rat cortex[24,25,71,84] as follows: layer I is 0 to 100 μm, layer II/III is 100 to 500 μm, layer IV is 500
to 750 μm, and layer V is 750 to 900 μm. 1b. Axial potential along horizontal axonal fiber
located 100 um below the cortical surface. The axon is depolarized when the activating function
is below zero and hyperpolarized when the activating function is above zero. 1c and 1d. As a
result of 1b, horizontal fibers are activated by the cathode and hyperpolarized by the anodes (1d).
The opposite is true for vertical fibers (1c). The vast majority of direct stimulation occurs in
layers I and II/III due to the decay of electric potential with depth.

189 **Cell activation results from a combination of morphology (cell type) and depth within the**
190 **column**

191   The applied electric field potential generated by the system of three electrodes (assuming
192 homogenous tissue) is shown in Fig 1a. To estimate the probability of specific cell types being
193 activated by stimulation, we simulated the various cell types based on three-dimensional (3D)
194 morphological reconstructions of neurons derived from electron microscopy available from
195 NeuroMorpho.org[22]. The excitatory cells we considered were pyramidal cells across layers II-V,
196 while the inhibitory neurons included basket cells and Martinotti cells across layers II-V in
197 addition to layer I interneurons (Table S1). Example reconstructions as well as average axon
198 density plots per cell type/cortical layer (Fig 2) demonstrate the significant differences in axonal
199 arborization and density among the different cell types, as well as between cells of the same type
200 based in different layers.

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

Figure 2

Representative reconstructions and averaged axon density for neuronal cell types modeled.
For each neuronal cell type employed in the cortical microcircuit model, we plot both a single
representative anatomical reconstruction as well as an averaged axonal density heatmap for all
the reconstructions of that type. Cells are arranged by layer and type, where the first row depicts
layer I inhibitory neurons. The representative LI interneuron is a horizontal cell, but we average
across small, descending, and horizontal LI interneurons to obtain the axonal heatmap as well as
in our analyses. The following rows depict pyramidal, basket, and Martinotti cells across layers
II/III, IV, and V (LVa for pyramidal cells). In axonal density (AD) plots, color indicates the
averaged density computed using all available reconstructions for the given cell type. The color
scale is logarithmic for visual clarity. AD gives a sense of the general orientation and density of
axon branches for each cell type, which is key to understanding subsequently computed
activation probabilities.

The hyperpolarization or depolarization of a neuronal fiber within a constant electric field
can be modeled with one-dimensional cable theory in conjunction with the activating function.
The activating function (see Methods) computes the net transmembrane current generated by
external stimulation (while ignoring pre-existing synaptic currents). According to one-
dimensional cable theory, the activating function is the second-order spatial derivative of the
electric potential along the neuronal fiber. The case of a perfectly horizontal fiber is shown in
Figure 1b. Through this, we can draw relationships between the orientation and excitation of a
fiber in response to a given stimulation polarity. Indeed, horizontal fibers were depolarized by
cathodal stimulation and hyperpolarized by anodal stimulation, while in contrast vertical fibers
were hyperpolarized by cathodal stimulation and depolarized by anodal stimulation (Figures 1d
and 1c, respectively). While each neuron has unique axonal fibers that span three-dimensional
space, these maps of activation and suppression zones for orthogonal axonal orientations give us
insight into how each cell type will behave across the stimulated space given its average axon
density and orientation (Fig 2).

We next calculated the spiking probability in response to the applied electric field
potential for each cell type/cortical layer by averaging across the activating function results of
their respective cell reconstructions; each cell reconstruction was shuffled by rotating and
shifting along the vertical axis and multiple reconstructions were considered for each cell type
(see Methods and Komarov, et al.[20] for details). This calculation compares the overall
excitability of each reconstruction to an experimentally-derived threshold (f_{th} = 3 pA/μm^2) to
determine probability of spiking. This threshold was set twenty times higher for unmyelinated
cell types (Martinotti cells and layer I interneurons) compared to myelinated cell types
(pyramidal and basket cells) since unmyelinated fibers are relatively unexcitable and lack nodes
of Ranvier[24–26].

The results of these calculations are shown in Figure 3. The average axonal density
heatmaps in Figure 2 as well as the presence or absence of myelination explain the variation of
activation responses across cell types and cortical layers.

276   Figure 3

Probability of spiking as a function of horizontal distance from the center electrode for each cell type and cortical layer.

Average (solid line) cell spiking probability and 95% confidence intervals (shaded region) for
each reconstruction was calculated for soma locations across the entire X-Z plane of its cortical
layer by averaging spiking probability across rotations and vertical shifts of all cell
reconstructions. Activation probabilities were calculated with a cathodal stimulation current of
150 μA and an anodal stimulation current of -75 μA per electrode over a 200 μs stimulation
period. Pyramidal cells (excitatory) and Basket cells (inhibitory) are highly activated by cathodal
stimulation and are minimally activated by anodal stimulation due to their myelination and
horizontally oriented axonal arbors. Martinotti cells (inhibitory) are activated by all electrodes
with a slight preference for the anodes but lack myelination and thus show less activation overall.
LI interneurons are also unmyelinated and are minimally excited by stimulation.

Across all layers, pyramidal cells were strongly activated by the cathode and minimally
activated by the anodes but showed decreasing intensity of stimulation response with increasing
depth (Fig 3a). As shown in Figure 2, all pyramidal cells vertically span the cortical layers
regardless of soma position. However, Layer II/III and LIV pyramidal cells exhibit significant
horizontal axonal density close to the cortical surface and thus responded more strongly to
stimulation overall (and to cathodal stimulation in particular), whereas the bulk of Layer Va
pyramidal axons lie in deeper layers and lack the superficial axonal density to be adequately
stimulated above threshold. Strong overall pyramidal response to stimulation is due to their
myelinate axons.

Basket cells also exhibited a strong preference for cathodal stimulation and little
activation underneath the anodes (Fig 3b). However, their responses were significantly more
tiered according to cortical layer as compared to pyramidal cells because basket cell arborization
is localized within the same layer as the soma (Figure 2). Their preference for cathodal
stimulation is due to their largely horizontal axonal arbor that stretches out within each layer.
Basket cells were the only myelinated inhibitory cell type in our model and therefore
demonstrated a significantly stronger spiking response overall relative to Martinotti or layer I
interneurons.

Martinotti cells across all cortical layers are moderately activated by both cathodal and
anodal stimulation but showed a slight preference for the latter (Fig 3c). This is because all
Martinotti cells make universal connections with pyramidal cells via layer I (Fig 2), and
therefore the majority of their arborizations lie in vertical axonal fibers connecting the soma to
layer I, with additional density spread out horizontally across layer I. However, they exhibited a
dampened stimulation response overall due to their unmyelinated axons.

Lastly, since layer I axon fibers are unmyelinated and stay localized to layer I (resulting in
mainly horizontal arborization), layer I interneurons to show a slight preference for cathodal
stimulation but little activation overall (Fig 3d).

Cortical microcircuit model shows directional propagation when stimulated with three
electrode array

In the previous section, we estimated the activation probabilities of isolated neurons
within an applied electric field. To understand how stimulation affects the dynamics between
neurons and ultimately the overall dynamics of the cortex, we constructed and stimulated a
network model of the cortex using simplified neuron models and previously-calculated activation
probabilities. Each cortical column was modeled as a canonical microcircuit containing the same
cell types and cortical layers as the biophysical analysis above; cross-columnar connections were
formed only between excitatory cells within the same layer. A schematic of the cell types and
synaptic connections is shown in Figure 4.

Figure 4

Microcircuit diagram of a single cortical column in modeled network.

This depiction of a single cortical column details the cell types/cortical layers and the synaptic
connections between them in our microcircuit network model. Circular labels (IN – LI
interneurons; BC – basket cells; MC – Martinotti cells) indicate inhibitory neurons, while
triangular labels (PY – pyramidal cells) indicate excitatory neurons. Green connections indicate
excitatory synapses, with the circular end indicating the postsynaptic cell and the unlabeled end
indicating the presynaptic cell. Purple connections indicate inhibitory synapses, with the
perpendicular line indicating the postsynaptic end. Teal-colored connections are also excitatory
but represent connections from pyramidal cells to others in adjacent columns (cross-columnar
synapses). The only cross-column synapses present are within-layer pyramidal-pyramidal
excitatory connections to adjacent columns. Layer I includes only inhibitory interneurons, which
inhibit pyramidal cells in all three deeper layers and are reciprocally excited by the same cells.
Each of the deeper layers contains pyramidal, basket, and Martinotti cells. Within a cortical
column, pyramidal cells reciprocally excite other pyramidal cells in their same layer as well as
across cortical layers. Basket cells act as local interneurons as they only inhibit and are excited
by pyramidal cells within their own layer. In contrast, while Martinotti cells are only excited by
pyramidal cells within their own layer, they universally inhibit pyramidal cells across all layers.
In this model, inhibitory cells receive only excitatory synaptic inputs. The number of neurons in
each column is listed in Table S2.

Following a brief stimulation period in which the activation probabilities from the
biophysical calculations were applied to the model, the network was allowed to run without any
external input for 500 ms, behaving according to synaptic interactions between neurons. A raster
plot and voltage and conductance traces of one microcircuit trial are shown in Figures 5 and 6.
The trial shown is one example of the general behavior of the microcircuit in the majority of
trials (Fig 5c) in which spiking activity, particularly from LII/III pyramidal cells, propagates to
the rightward columns but not past the leftmost electrode. Given this unique spiking activity and
the biological importance of LII/III pyramidal cells in mediating communication across cortical
regions, we chose to focus our analyses on the network behavior of LII/III pyramidal cells.

381 Figure 5

Directed propagation of pyramidal activity in raster plot of microcircuit trial.

A raster plot displaying network behavior during and after stimulation in one trial of the
microcircuit simulation across all cortical columns. Each cell within the microcircuit has its own
coordinate on the y axis, and each dot is an action potential. Orange dots indicate spikes that are
directly triggered by electrical stimulation (occurs during first 5 ms), while blue dots indicate
spikes triggered via synaptic input. Figure 5a displays the first 50 ms of the simulation (the
network is silent beyond this period), and Figure 5b zooms in on the first 10 ms in the five
central columns. In Figure 5c, the number of spikes (in all cell types) was summed across the
simulated cortical columns and averaged from 150 different simulations. In the vast majority of
simulation trials, pyramidal activity in superficial layers propagates in a traveling wave to the
right but is halted underneath the anodal electrodes by primarily Martinotti cell activity.

Directionality of the stimulation-triggered wave can be explained by network inhibition

The activation probability curves in Figure 3 provide intuition into the network behavior during and immediately after the stimulation period (0-5 ms; Fig 5b). Let us first examine the column underneath the cathodal electrode (Fig 5b – column 7). While both layer II/III and layer IV pyramidal cells were predicted to be highly activated underneath the cathodal electrode (Fig. 3), only layer IV pyramidal cells were directly activated. Although all pyramidal cells were inhibited by moderate Martinotti activity, layer II/III pyramidal cells in particular were locally inhibited by strong synchronous LII/III basket cell activity because basket cell response dropped off with increasing cortical depth and because basket cells only inhibited within their own layer. Following stimulation, layer IV pyramidal cells excited layer II/III pyramidal cells and triggered a cluster of LII/III activity. There was negligible layer V excitation during stimulation and none following due to their low excitation probabilities and relatively small neuronal population.

Network behavior in response to anodal stimulation contrasted sharply with cathodal stimulation response and underpins unidirectional excitatory propagation (Fig 5b – columns 5 and 6). Although Layers II/III and IV pyramidal cells were still moderately activated by anodal stimulation (Fig 3a), very few cells were pushed above threshold due to strong inhibition. Martinotti cells showed a preference for anodal stimulation (according to Figure 3) and thus fired early in the stimulation period, inhibiting pyramidal cells across all cortical depths (since Martinotti cells universally synapsed to all pyramidal cells). This strong inhibitory force coupled with moderate superficial basket cell activity silenced almost all pyramidal activity across cortical layers.

In the column directly to the left of the electrode array (Fig 5b – column 4), there was negligible activity across all cell types and cortical layers. Not only did the electric field potential drop off significantly at this distance, but pyramidal and basket cells were already minimally activated by the anodal electrodes, and Martinotti cells were only moderately activated by the anodal electrodes due to their lack of myelination. In the absence of stimulating electric field potential or activating input from neighboring cortical columns, the first three columns exhibited no spiking activity at all (Fig 5b – columns 1-3). Hence, the excitatory pyramidal activity present underneath the electrodes did not propagate leftward past the anodal electrodes. This activity, however, did travel rightward past the electrode array, growing stronger and more synchronous as it propagated.

On the other side of the array, in the cortical column directly to the right of the cathodal electrode (Fig 5a & 5b – column 8), there was moderate direct activation of pyramidal cells and little direct activation of inhibitory cells. This follows from Figure 3, which depicts pyramidal cells continuing to be activated by the cathodal electrode. While basket cells were also moderately activated by cathodal stimulation, their joint inhibition with Martinotti cells was not enough to counter pyramidal stimulation response. This allowed for dense clusters of excitatory activity in pyramidal cells following stimulation.

In the second column to the right of the electrode array (Fig 5a – column 9), we see a
dense cluster of highly synchronized pyramidal layer II/III activity that was slightly delayed
from the activity in the column to its left. Although pyramidal cells were no longer directly
stimulated in this cortical column, this activity resulted from the rightward cross-column
propagation of excitatory signaling. Remarkably, this substantial, synchronized pyramidal
activity grew more and more synchronous as the wave propagated rightward across cortical
columns. This unidirectional propagation of excitatory signaling is an exceptional result of
asymmetrical stimulation (Fig 5a – columns 9-11).

Following the stimulation period and initial clusters of activity that die down around 10
ms post-stimulation, there were a handful of waves of activity that ping-ponged between
excitatory and inhibitory cells in columns with pyramidal excitation (Fig 5a – columns 6-11). In
these columns, pyramidal activity activated both Martinotti and basket cells, which in turn
inhibited pyramidal activity. This negative feedback loop went through a handful iterations over
the course of a few milliseconds before halting pyramidal spiking entirely.

Analysis of synaptic currents reveal mechanism of asymmetrical spiking activity

Next we analyzed voltage traces of individual neurons and synaptic dynamics to explain
the causes of pyramidal asymmetrical spiking activity (Fig 6).

Figure 6

Voltage traces for layer II/III cells and PY input conductances during simulation show how
inhibition causes directionality of traveling wave.

Each of the subplots in the grid contain data for the collection of layer II/III cells specified by
type and column. The x-axis for each subplot is time in milliseconds and is restricted to the
interval from 5 msec before stimulation to 50 msec after. The stimulation period is depicted by
the region in light green. Subplot columns, labeled at top, indicate the model columns; columns
of 4 through 10 of the total 11 are included. The first three rows depict data from pyramidal cells,
with the first row depicting the voltage trace, the second the total excitatory conductance (the
sum of all incoming pyramidal connections), and the third the total inhibitory conductance (sum
of inhibitory connections from layer I interneurons, basket cells, and Martinotti cells). The fourth
and fifth rows depict the voltage traces of basket and Martinotti cells, respectively. In each
subplot, there are light grey traces depicting each of the individual cells in the group, and their
average (within the same column, layer, and subtype) in red. Sharp increases in the light grey
trace indicate action potentials (spikes) and darker grey regions correspond to times when large
numbers of cells spiked.

Voltage traces revealed that LII/III pyramidal cells were initially depolarized underneath
the cathodal electrode (column 7), as they were highly activated by cathodal stimulation.
However, a large inhibitory conductance immediately following the stimulation period
dampened any stimulation-induced depolarization and hyperpolarized these pyramidal cells. The
inhibitory conductance then fell and the excitatory conductance rose as layer II/III pyramidal
cells gained excitatory input from neighboring layers and columns, bringing layer II/III
pyramidal cells to threshold and triggering action potentials. The peak in inhibitory conductance
midway through the pyramidal action potential was caused by pyramidal cell input into the
inhibitory cells, which initiated a negative feedback loop that quickly subsided as pyramidal cells
were silenced by inhibition. This effect occurred in all columns with substantial pyramidal
activity.

Although there were similar inhibitory and excitatory conductance dynamics in the
column underneath the central anodal electrode (column 6), the excitatory conductance was of a
smaller magnitude overall and there were fewer action potentials because pyramidal cells were
less excited by anodal stimulation. Underneath the leftmost electrode (column 5), moderate
inhibitory conductance outweighed negligible excitatory conductance, leading to minimal
pyramidal activity. There was little excitation in column 4 or any of the other leftward columns
(not shown). On the other side of the electrode array, in column 8, high initial excitatory
conductance and minimal inhibitory conductance resulted in strong initial pyramidal spiking.
Pyramidal action potentials became more and more synchronous as they travel rightward, as
evidenced by increasingly overlapped voltage traces. Together, these observations revealed the
mechanisms of activity propagation to the right but not to the left in given stimulation settings.

DISCUSSION

In this work we predict that an asymmetrical cortical stimulation protocol using a
combination of anodal and cathodal electrodes may trigger propagating excitatory activity that
shows strong directional preference. Our model has two steps: we first construct a biophysical
model to predict activation probabilities across cell types in response to an asymmetrically
applied electric field potential, and then incorporate these probabilities into a cortical
microcircuit to model the network effects of stimulation. We found that pyramidal cells and
basket cells are highly activated by the cathodal electrode and minimally activated by the anodal
electrodes due to their myelination and horizontal axonal arbors, while Martinotti cells and Layer
I interneurons are moderately activated by both electrodes but exhibit a slight preference for the
anodal stimulation as they are unmyelinated and have vertically-oriented axonal arbors. Network
model simulations revealed that this asymmetrical activation results in a traveling wave in
superficial excitatory cells that propagates away from the electrode array, past the cathodal
electrode, and into adjacent cortical columns, but does not propagate in the opposite direction
past the leftmost anodal electrodes (Fig. 7).

535 Figure 7

	COLUMN 4	COLUMN 5	COLUMN 6	COLUMN 7	COLUMN 8
PY	No direct activation + no excitatory input + small inhibitory input = little spiking	Low direct activation + low excitatory input + moderate inhibition = low spiking	Direct activation + very little inhibition = strong spiking	Direct activation + excitatory input outweighs inhibition = strong spiking	Direct activation + high excitatory input + very low inhibition = strong spiking
BC	No direct activation + no excitatory input = little spiking	Low direct activation + low excitatory input = low spiking	Direct activation + moderate excitatory input = strong spiking	Direct activation + excitatory input = strong spiking	Strong excitatory input = strong spiking in response to PYs
MC	Small activation + no excitatory input = little spiking	High direct activation = strong initial spiking	Direct activation + moderate excitatory input = moderate spiking	Direct activation + excitatory input = moderate spiking	Excitatory input = strong spiking in response to PYs

Summary of interactions resulting in unidirectional propagation.

Key interactions are depicted graphically above and summarized in text in the table below.
Columns 4 through 8 of the total 11 are included, indicated by labels at top. Cell and synaptic
labels are as in Figure 4. Orange arrows coming from the electrodes and pointing toward cells
indicate direct electrical stimulation, as opposed to synaptic inputs, depicted by the green, purple,
and teal lines. The opacity of the cell labels approximately depicts the degree of spiking activity,
with more transparent cells showing little or no spiking activity, and more opaque ones
corresponding to more actively spiking cells. The opacity of lines indicates the strength of the
input to cells, either synaptic or from the stimulating electrodes, e.g. more opaque orange arrows
indicate that the cell being pointed to was more strongly stimulated directly. The light blue
dashed arrow at top right indicates the initiation of synchronous, unidirectional propagation of
activity in the direct right of the electrodes. Key events in initiation of this unidirectional
propagation are indicated by the white numbers in black circles: ① Direct activation under (+)
[col. 7] and central (-) [col. 6] induces strong spiking in PYs, BCs, MCs. ② Under left (-) [col.
5] direct activation induces strong spiking in MCs, but little in PYs, BCs. ③ PYs in col. 5
further inhibited by MCs leads to little activity to left of electrode. ④ Strong PY activity in cols.
6 and 7 propagates rightward via cross-column PY-PY synapses, overcoming moderate
inhibition. ⑤ PY activity causes spiking in MCs and BCs as it propagates rightward, feedback
between PYs and inhibitory cells causes increasingly synchronous spiking. Layer I interneurons
have been omitted, as they did not contribute strongly to the propagation described here.

This model develops a stimulation protocol that can be experimentally verified in future
studies. We found that inhibitory cells are the cause of unidirectional propagation, which implies
that excitatory activity would exhibit unhindered propagation without inhibition. Inhibitory cells
could be deactivated optogenetically during stimulation to test model predictions. Martinotti cells
may also be deactivated separately from basket cells via genetically-targeted optogenetics to
localize the main source of inhibition.

Traveling waves in the brain

Experimental recordings have observed traveling waves during most types of cortical
processing. They relay information across a range of distances and thereby coordinate processes
such as memory, perception, language, orientation, executive functions, and more across distant
brain regions[17,27–29]. Multi-electrode recordings in human and animal subjects have demonstrated
the ubiquity of traveling waves in cognitive function[17]. A growing body of evidence therefore
places traveling waves as integral to cognition, as they facilitate fast, efficient information
transfer and local-to-global communication[30]. Multi-electrode recordings show that propagating
activity present in the human cortex is often directional, traveling from one point to another. The
ability to generate directional propagation via stimulation would allow for unique precision in
and control over induced activity. This has widespread implications for the restoration or
enhancement of high-level brain function, particularly because many neuropsychiatric disorders
are marked by abnormal or absent propagating activity.

In the model, propagating activity was confined to supragranular cortical layers. This
may not be a limitation if the goal is to reproduce natural waves, inasmuch as spontaneous
traveling waves in the human cortex have also been found to be largely confined to upper layers,
including the alpha rhythm during waking[18], and spindles and slow oscillation during sleep[31,32].

Traveling waves have long been studied with a variety of computational models, and
numerous mechanisms have been proposed to explain how activity propagates across neuronal
networks[33,34]. Cortical propagating waves that are triggered specifically by electrical stimulation
have been recorded in mammalian cortical slices[35–37], as well as in non-human mammals[38–40] and
simulated in non-mammalian computational models[41], but remain understudied in human
subjects. This previous experimental work, both *in vivo* and *in silico*, has yielded scarce evidence
of asymmetrical travelling wave propagation or reliable wave generation analogous to that
reported here. To the extent that previous work has focused on travelling wave propagation
initiated by stimulation[42], these studies examined stimulation through the skull and meninges,
and are thus not directly comparable to this model of intracranial stimulation of the cortical
surface. Many computational models exist that model the effects of stimulation on the brain,
including some that have constructed Hodgkin-Huxley microcircuits[43,44] and modeled cortical
surface stimulation[45]. However, few have modeled multi-electrode or asymmetrical stimulation
or reproduced traveling waves using surface electrodes. Many existing stimulation models have
focused on stimulation of a particular nerve[46,47] or isolated cell type[48,49], as opposed to the
functioning cortical microcircuit presented here, which can be more readily adapted to other

cortical regions. In addition, stimulation has more often been simulated in these models by a
simple application of superthreshold current or a uniform electrical field[50,51]. Thus, by combining
the two-phase biophysical model with asymmetrical, multi-polar surface stimulation, our
approach synthesizes existing achievements into a single coherent, clinically-adaptable model
that uniquely sheds light on the generation of propagating wave activity.

Clinical relevance of our findings

Brain stimulation is becoming increasingly common in clinical and experimental settings,
especially using multi-electrode arrays[52]. As such, it is pressing that we develop accurate models
of the effects that multi-electrode stimulation has on neural activity. While sometimes the
explicit goal of stimulation may be to disrupt aberrant activity to restore normal functioning,
increasingly the goal is to induce the desired brain activity directly via stimulation, as our work
demonstrates.

Changes in neural plasticity result from patterned activity, with the particular changes in
connectivity contingent on the specific timing and order of activity[53]. Stimulation protocols that
induce neural activity which continues past the stimulus duration are more likely to alter cellular
and synaptic properties in favor of the induced activity, in contrast to stimulation protocols that
briefly activate broad swaths of cells without triggering existing activity patterns. Thus, initiating
propagating waves within tailored spatiotemporal constraints is a promising way to retrain neural
networks and enhance or silence brain functions in a targeted way.

The generation of traveling waves may serve as a promising therapy for a variety of
neurological disorders. For example, it has been previously suggested that triggering propagating
activity in perilesional areas where waves are otherwise aberrant or absent may be an effective
therapy for post-stroke aphasia[54]. While the ultimate clinical applications of this technique are
uncertain, stimulation-induced traveling waves may have the potential to offset inhibition in
cortical spreading depression[55,56], reduce the risk of seizure while determining which brain tissue
to remove from epileptic patients[11], and enhance memory formation when applied during
learning or recall periods[13,14,16,57,58], as consistency in traveling wave direction is positively
correlated with working memory efficiency[59].

Limitations

In this work, we modelled a single, short stimulating pulse. However, clinical stimulation
is most often composed of longer pulses or pulse trains and is usually performed with bipolar
electrodes delivering biphasic pulses in order to prevent damaging Faradic currents[60,61]. These
stimulation paradigms modulate properties over time that are not accounted for in the current
model such as underlying dendritic and axonal dynamics as well as synaptic interactions. Thus,
our biophysical approach may be expanded in future studies to incorporate these steady-state
properties through alternative modeling approaches such as the cylinder model[62,63] or the multi-
compartmental model[64]. However, we chose to model the activation probability of the axonal
instead of the dendritic arbor in this work because experimental evidence shows that the nodes of

Ranvier, followed by the axon hillock, are the most excitable neuronal elements by far via direct stimulation[65–68] as they both have a high concentration of sodium channels[69]. In contrast, direct stimulation of the dendritic arbor generates transmembrane currents that propagate to the axon hillock, but these effects are strongly attenuated and delayed, and are negligible compared to direct stimulation of the nodes of Ranvier and axon hillock.

In the microcircuit phase of the model, the connectivity between different cell types follows a canonical microcircuit model. While this approach characterizes the main signal pathways and feedback loops present within cortical columns[70–72], finer details are not modeled such as the variability of layer II/III pyramidal cells and their projections[73,74], descending projections to inhibitory cells from excitatory cells[75], or the contribution of less common interneuron cell types. These and other characteristics such as axonal and dendritic arborization change between species and cortical areas. The cells within the cortical microcircuit model could be extended from single- to multi-compartment neurons[76] in order to distinguish tuft versus soma-targeting interneurons, which may further differentiate the inhibitory power of interneuron cell types[24]. This phase of the model may be further expanded from a two-dimensional plane to a three-dimensional circuit in the volume of cortex underneath the electrodes, which would allow for a better understanding of how activity spreads across space.

Generalization of this approach

While this work modeled a specific stimulation protocol, the approach is generalizable to a variety of clinical applications. The model can be customized in terms of the number and shape of the electrodes, the arrangement of anodal and cathodal electrodes, the intensity of the applied current, and the brain area studied (provided sufficient neuronal reconstructions are available). This model can also be modified to simulate other applied stimulating pulses as long as an activation threshold has been experimentally determined. However, further adjustments are necessary to adapt this model to human stimulation protocols such as modeling a biphasic bipolar pulse, incorporating human cell reconstructions as they become available, and adjusting model parameters to account for the thicker human pia and cortex.

Conclusions

This work models an asymmetrical stimulation paradigm that could be implemented to initiate unidirectional traveling waves in the cortex. A biophysical model is integrated with a network-computational model to predict the behavior of single neurons as well as the cortical network dynamics resulted from multielectrode stimulation. This model provides hypotheses and stimulation paradigms that can be verified experimentally and also provides an avenue for further experimental work to be guided systematically rather than in blind trial-and-error fashion. These results demonstrate how complex stimulation protocols could be harnessed to generate persistent changes in activity with the potential to restore normal brain function in neurological and psychiatric conditions.

Methods

Cell reconstruction selection:

All neuronal cell reconstructions were chosen from publicly available datasets on neuromorpho.org[22,25,77–80]. The types of cells included and their respective datasets are listed in table S1. We used multiple cell reconstructions for each cell type to account for anatomical diversity. We used experimental measurements to approximate the cutoff depths for each layer (Fig 1a)[24,25].

Calculating the electric field potential generated by the electrode array:

The electrode array modeled in this study is composed of three square electrodes (each 150 μm by 150 μm) placed linearly on the surface of the cortex. Two electrodes have negative current (-75 μA each) and one electrode has positive current (150 μA) (Fig 1a), and stimulation is applied for 200 μs. These values are in accordance with common experimental parameters[21]. Assuming that the current sources are homogenous square electrodes, we calculated the electric field potential of each electrode as follows:

$$\Phi(X,Y,Z) = \frac{\rho_e I}{4\pi A^2} \int\int_{-A/2}^{A/2} \frac{dx\,dy}{\sqrt{(X-x)^2 + (Y-y)^2 + Z^2}} \qquad (1)$$

I is net current, ρ_e is extracellular resistivity and A is the length of the square electrode edge (Fig.1a). In this study, A=150 μm and net current I is either -75 μA or 150 μA. The derivation of this formula can be found in Komarov et al.[20].

We sum all three electric field potentials at each point in space to determine the overall electric field potential.

Estimating the activating function:

Derived from one-dimensional cable theory[63,66], the transmembrane voltage dynamics of axonal segments can be modeled as follows:

$$\frac{c_m dV_i}{dt} = \left[\frac{d}{4\rho_i} \left(\frac{V_{i-1} - 2V_i + V_{i+1}}{\Delta x^2} + \frac{\Phi_{i-1} - 2\Phi_i + \Phi_{i+1}}{\Delta x^2} \right) - \sum I_n^{ion} \right] \qquad (2)$$

V_{i-1}, V_i, and V_{i+1} are transmembrane voltages of the neighboring axonal compartments (sub index indicates number of the compartment), $c_m = 1$ μF/cm^2 is a capacitance of membrane per square unit area, d is the diameter of the axon (typically between 10 μm and 1 μm), $\rho_i = 300$ Ω·cm is a resistivity of axoplasm, and Δx is a discretization parameter that defines length of the compartment. The term $\sum I_n^{ion}$ is the sum of intrinsic ionic currents such as leak currents and fast potassium and sodium currents for spike generation. From above, the activating function is the effective transmembrane current that arises due to extracellular electrical stimulation and is described as

$$f = \frac{d}{4\rho_i} \frac{\Phi_{i-1} - 2\Phi_i + \Phi_{i+1}}{\Delta x^2} \qquad (3)$$

where Φ_{i-1}, Φ_i, and Φ_{i+1} stand for extracellular potentials in the vicinity of axonal compartments.

In the limit $\Delta x \to 0$ the expression becomes $f = \frac{d}{4\rho_i}\frac{\partial^2 \Phi}{\partial x^2}$, where the x axis represents the direction

of the axonal fiber. To estimate the probability of axonal activation, we calculate the

transmembrane current generated in each small component piecewise along reconstructed axons

using the activating function. This calculation is based upon the orientation of each component in

space because the activating function is computed along the axonal direction by definition. Since

jitters may be present along the edges of reconstructions, we estimate component direction using

neighboring segments in space up to 10 microns away.

Estimating the threshold of activation

After calculating the transmembrane current generated within each axonal segment of cell

reconstructions, we used a threshold of activation to determine whether a neuronal response was

triggered. This threshold was drawn from *in vivo* experiments which define the threshold injected

current I required to induce a threshold effective current f at the axonal initial segment a distance

d from the electrode[48,81]. Komarov et al.[20] simulated one such experiment by computing the

activation current f at the axon initial segment of a layer II/III pyramidal cell while varying the

stimulation current I and distance d. This simulation used a 200 μs duration stimulating pulse,

which is typical of similar *in vivo* experiments[48,81]. The value $f = f_{th} = 3$ pA/μm^2 fully replicated

the experimentally observed current-distance relationship across varying stimulation currents and

distances. Thus, this threshold value f_{th} was used to determine whether each axonal segment was

activated by induced transmembrane current.

Computing the average activation probability per cell type

To determine whether a given neuronal reconstruction would be activated by applied current, we

used the activating function to calculate how many axonal segments had above-threshold

transmembrane current values that could initiate an axonal action potential (assumed at nodes of

Ranvier); these above-threshold axonal segments are collectively called the trigger area. The

activating function threshold was set to $f_{th} = 3$ pA/μm^2 for myelinated axons but to $f_{th} = 60$

pA/μm^2 for unmyelinated axons, since unmyelinated segments are significantly less excitable as

they have fewer sodium channels[61]. In this model, we assumed pyramidal and basket cells are

myelinated and Martinotti and Layer I interneurons are unmyelinated based upon experimental

data[25,75,82]. For each cell reconstruction at a given point in space (i.e. a given coordinate in the x-

z plane, Fig 1a), we found the probability of firing as follows:

We computed the activating function along the axonal segments of the cell reconstruction and

found the trigger area L as described above.

a. For myelinated axons, we assumed that at least one node of Ranvier must be present

746   in the trigger area in order to initiate an axonal response. To approximate the

747   probability of a node of Ranvier in the trigger area, we first discretized the trigger

748   area into segments that are the typical length of a node of Ranvier (k=1 μm). If we

749   approximate the mean internodal distance as D = 100 μm, then $p_n = k/D$ is the

750   probability that there is a node of Ranvier in a given segment. It follows that $(1 - p_n)^N$

751   is the probability that the trigger area L does not contain a node of Ranvier, where N

$= L/k$ is the number of segments of length k that can fit into a fiber of total length L.
Therefore, the overall probability p that a myelinated axonal arbor will be activated is
estimated as $p = 1 - (1 - p_n)^N = 1 - \left(\frac{D-k}{D}\right)^{L/k}$.

b. For unmyelinated axons, we assumed that any axonal segment with an activating
function above threshold ($L > 0$) resulted in an overall cell response, since the entire
membrane is exposed to extracellular space.
Given that variation in cell position and orientation within each cortical layer is present in nature,
we additionally average across neuronal rotations and vertical shifts. Thus, we computed the
average activation probability for each cell type/cortical layer as follows: for each cell
reconstruction, we first position the soma at a point along the x-axis within its cortical layer. We
then performed four rotations about the vertical axis of the cell reconstruction in three-
dimensional space, and at every rotation we calculate the likelihood of activation by computing
the activation function across the axonal arbor, as outlined above. We then averaged across these
four probabilities and set the result as the activation probability for said point. After this, we
incrementally shift the soma of the reconstruction vertically within its cortical layer and find the
mean activation probability at each point. We averaged across all vertical shifts to obtain an
approximate spiking probability for this cell reconstruction at this point along the x-axis. We
then computed the activation probability of all cell reconstructions across the x-axis in this
manner, and finally average across all reconstructions within a given cell type and cortical layer
to find the expected spiking probability in response to stimulation for each class of cell.
Computational model of the cortical circuit:
Rationale: The network model is composed of eleven interconnected cortical columns, and each
column contains layer I interneurons and pyramidal, basket, and Martinotti cells from layers II-V
(same as the biophysical analysis). The number of cells within each column is outlined in table
S2. This balance of excitatory to inhibitory cell types approximates the true cell composition of
the rat cortex, where pyramidal cells are the primary excitatory cells and basket and Martinotti
cells comprise the majority of inhibition within and across layers[24,25]. Cells were constructed to
only spike if receiving synaptic input or electrical stimulation. Each cell behaves according to
Hodgkin-Huxley dynamics, with a handful of parameters differentiating excitatory and inhibitory
cells. Basket cells were modeled as fast-spiking cells, while the other cell types were modeled as
regular-spiking cells with spike rate adaptation. Inhibitory cells were fired more quickly than
excitatory cells in response to activation because all interneurons were modeled as having a
lower leak current[56]. Some additional parameters are: a fast Na^+-K^+ spike generating mechanism
(all cells), high-threshold activated Ca^{2+} current (for pyramidal cells), and slow calcium-
dependent potassium (AHP) current (for regular spiking cells).
Equations: The membrane potential is described by following equation[83]:
$$C_m \frac{dV_i}{dt} = I_{ion}(V_i) + I_i^{syn} + I_i^{ext} + \eta \xi_i \qquad (4)$$
The ionic currents $I_{ion}(V_i)$, which are responsible for intrinsic cells dynamics, are as follows:

$$I_{ion}(V_i) = g_{Na}m_i^3h_i(V_{Na} - V_i) + g_Kn_i^4(V_K - V_i) + g_L(V_L - V_i) + g_{Ca}\frac{V_{Ca} - V_i}{1 + \exp\left(-\frac{V_i - V_{th}}{V_{shp}}\right)}$$

$$+ g_{ahp}\frac{c_i}{c_i + k_d}(V_K - V_i) \tag{5}$$

The gating variables m_i, n_i, h_i evolve according to:

$$\frac{dx_i}{dt} = \alpha_x(V_i)(1 - x_i) - \beta_x(V_i)x_i \tag{6}$$

where x_i is one of gating variables. The functions $\alpha_x(V)$ and $\beta_x(V)$ are:

$$\alpha_m(V) = \frac{0.32(54 + V)}{1 - \exp(-0.25(V + 54))} \tag{7}$$

$$\beta_m(V) = \frac{0.28(V + 27)}{\exp(0.2(V + 27)) - 1} \tag{8}$$

$$\alpha_h = 0.128\exp\left(-\frac{50 + V}{18}\right) \tag{9}$$

$$\beta_h = \frac{4}{1 + \exp(-0.2(V + 27))} \tag{10}$$

$$\alpha_n = \frac{0.032(V + 52)}{1 - \exp(-0.2(V + 52))} \tag{11}$$

$$\beta_n = 0.5\exp\left(-\frac{57 + V}{40}\right) \tag{12}$$

Calcium concentration c_i obeys the following equation:

$$\frac{dc_i}{dt} = -\alpha_{Ca}g_{Ca}\frac{V_i - V_{Ca}}{1 + \exp\left(-\frac{V_i - V_{th}}{V_{shp}}\right)} - \frac{c_i}{\tau_{Ca}} \tag{13}$$

and governs calcium-dependent hyperpolarizing potassium current

$$I_{AHP} = g_{ahp}\frac{c_i}{c_i + k_d}(V_K - V_i) \tag{14}$$

which is responsible for spike-frequency adaptation. The synaptic input was modeled according
to:

$$I_i^{syn} = \sum_j G_{ij}^{exc}s_j^{exc}(V_{syn}^{exc} - V_i) + \sum_j G_{ij}^{inh}s_j^{inh}(V_{syn}^{inh} - V_i) \tag{15}$$

where synaptic variables $s_j^{exc,inh}$ are governed by the following equation:

$$\frac{ds_j}{dt} = \alpha_s^{exc,inh}S(V_j)(1 - s_j^{exc,inh}) - \beta_s^{exc,inh}V_j \tag{16}$$

The function $S(V)$ reads:

$$S(V) = \frac{1}{1 + \exp(-100(V - 20))} \tag{17}$$

Initially, the network runs without stimulating input for 200 ms to simulate pre-existing activity.
Then the network is stimulated and runs for an additional 500 ms. To simulate electrical
stimulation, we used the binary term I_i^{ext} to inject above-threshold current into a subset of

randomly-chosen neurons within each cell type/cortical layer such that the fraction of neurons
induced to spike corresponds to the activation probabilities calculated in the biophysical phase of
the model. The term $\eta\xi_i(t)$ models spontaneous background activity as a white noise process (ξ)
with standard deviation η. All model parameters are listed in Table S2 (unless specified in the
description of simulations), and the network structure and connectivity are described in Table S3
and Table S4 respectively. Cells were synaptically coupled by excitatory (AMPA) and inhibitory
(GABA$_A$) connections. The strength and probability of connections between layers and cell types
are set according to a canonical cortical circuit[75].

Acknowledgements

This work was supported by NIH-NINDS (grant 1R01NS19553).

Competing Interests Statement

The authors declare no competing or financial interests.

References

1.   Tehovnik, E. J., Slocum, W. M. & Schiller, P. H. Differential effects of laminar
stimulation of V1 cortex on target selection by macaque monkeys. *Eur J Neurosci* **16**,
751–760 (2002).

2.   Salzman, C. D., Britten, K. H. & Newsome, W. T. Cortical microstimulation influences
perceptual judgements of motion direction. *Nature* **346**, 174–176 (1990).

3.   Halgren, E., Walter, R. D., Cherlow, D. G. & Crandall, P. H. Mental phenomena evoked
by electrical stimulation of the human hippocampal formation and amygdala. *Brain* **101**,
83–117 (1978).

4.   Blumenfeld, Z. & Brontë-Stewart, H. High Frequency Deep Brain Stimulation and Neural
Rhythms in Parkinson's Disease. *Neuropsychol. Rev.* **25**, 384–397 (2015).

5.   Papageorgiou, P., Deschner, J. & Papageorgiou, S. Effectiveness and Adverse Effects of
Deep Brain Stimulation: Umbrella Review of Meta-Analyses. *J. Neurol. Surg. Part A*
Cent. Eur. Neurosurg. **78**, 180–190 (2016).

6.   Baizabal-Carvallo, J. F. & Alonso-Juarez, M. Low-frequency deep brain stimulation for
movement disorders. *Parkinsonism Relat. Disord.* **31**, 14–22 (2016).

7.   Mayberg, H. S. *et al.* Deep Brain Stimulation for Treatment-Resistant Depression. *Neuron*
45, 651–660 (2005).

8.   Schlaepfer, T. E. *et al.* Deep Brain Stimulation to Reward Circuitry Alleviates Anhedonia
in Refractory Major Depression. *Neuropsychopharmacology* **33**, 368–377 (2008).

9.   Abelson, J. L. *et al.* Deep brain stimulation for refractory obsessive-compulsive disorder.
Biol. Psychiatry **57**, 510–516 (2005).

10.   Greenberg, B. D. *et al.* Three-Year Outcomes in Deep Brain Stimulation for Highly
Resistant Obsessive–Compulsive Disorder. *Neuropsychopharmacology* **31**, 2384–2393
(2006).

11.   Nagaraj, V. *et al.* Future of Seizure Prediction and Intervention. *J. Clin. Neurophysiol.* **32**,
194–206 (2015).

12.   Hummel, F. C. & Cohen, L. G. Non-invasive brain stimulation: a new strategy to improve
neurorehabilitation after stroke? *Lancet. Neurol.* **5**, 708–12 (2006).

13.   Suthana, N. *et al.* Memory enhancement and deep-brain stimulation of the entorhinal area.
N Engl J Med **366**, 502–510 (2012).

14.   Kucewicz, M. T. *et al.* Electrical Stimulation Modulates High γ Activity and Human
Memory Performance. *eneuro* **5**, ENEURO.0369-17.2018 (2018).

15.   Kahana, M. J., Aggarwal, E. V. & Phan, T. D. The variability puzzle in human memory. *J.*
Exp. Psychol. Learn. Mem. Cogn. **44**, 1857–1863 (2018).

16.   Ezzyat, Y. *et al.* Closed-loop stimulation of temporal cortex rescues functional networks
and improves memory. *Nat. Commun.* **9**, 365 (2018).

17.   Muller, L., Chavane, F., Reynolds, J. & Sejnowski, T. J. Cortical travelling waves:
mechanisms and computational principles. *Nat. Rev. Neurosci.* **19**, 255–268 (2018).

18.   Halgren, M. *et al.* The generation and propagation of the human alpha rhythm. *Proc. Natl.*

Acad. Sci. **116**, 23772–23782 (2019).

19.   Dickey, C. *et al.* Travelling spindles create necessary conditions for spike-timing-
dependent plasticity in humans. (2020). doi:10.1101/2020.05.21.079152

20.   Komarov, M. *et al.* Selective recruitment of cortical neurons by electrical stimulation.
PLOS Comput. Biol. **15**, e1007277 (2019).

21.   Ha, S. *et al.* Silicon-Integrated High-Density Electrocortical Interfaces. *Proc. IEEE* **105**,
11–33 (2017).

22.   Ascoli, G. A., Donohue, D. E. & Halavi, M. NeuroMorpho.Org: A Central Resource for
Neuronal Morphologies. *J. Neurosci.* **27**, 9247–9251 (2007).

23.   Komarov, M., Malerba, P., Nunez, P., Halgren, E. & Bazhenov, M. Selective Recruitment
of Cortical Neurons by Electrical Stimulation. *bioRxiv* 213017 (2017).
doi:10.1101/213017

24.   Markram, H. *et al.* Interneurons of the neocortical inhibitory system. *Nat. Rev. Neurosci.*
5, 793–807 (2004).

25.   Wang, Y. *et al.* Anatomical, physiological and molecular properties of Martinotti cells in
the somatosensory cortex of the juvenile rat. *J. Physiol.* **561**, 65–90 (2004).

26.   DeFelipe, J., Markram, H. & Rockland, K. The neocortical column. *Front. Neuroanat.* **6**,
1–2 (2013).

27.   Rubino, D., Robbins, K. A. & Hatsopoulos, N. G. Propagating waves mediate information
transfer in the motor cortex. *Nat. Neurosci.* **9**, 1549–1557 (2006).

28.   Wu, J.-Y., Xiaoying Huang & Chuan Zhang. Propagating Waves of Activity in the
Neocortex: What They Are, What They Do. *Neurosci.* **14**, 487–502 (2008).

29.   Salimpour, Y. & Anderson, W. S. Cross-Frequency Coupling Based Neuromodulation for
Treating Neurological Disorders. *Front. Neurosci.* **13**, 1–13 (2019).

30.   Turi, Z., Alekseichuk, I. & Paulus, W. On ways to overcome the magical capacity limit of
working memory. *PLOS Biol.* **16**, e2005867 (2018).

31.   Halgren, M. *et al.* Superficial Slow Rhythms Integrate Cortical Processing in Humans.
Sci. Rep. **8**, 2055 (2018).

32.   Cash, S. S. *et al.* The human K-complex represents an isolated cortical down-state.
Science (80-.). **324**, 1084–1087 (2009).

33.   Ermentrout, G. B. & Kleinfeld, D. Traveling Electrical Waves in Cortex. *Neuron* **29**, 33–
44 (2001).

34.   Breakspear, M. Dynamic models of large-scale brain activity. *Nat. Neurosci.* **20**, 340–352
(2017).

35.   Kim, U., Bal, T. & McCormick, D. A. Spindle waves are propagating synchronized
oscillations in the ferret LGNd in vitro. *J. Neurophysiol.* **74**, 1301–1323 (1995).

36.   Wu, J.-Y., Guan, L., Bai, L. & Yang, Q. Spatiotemporal Properties of an Evoked
Population Activity in Rat Sensory Cortical Slices. *J. Neurophysiol.* **86**, 2461–2474
(2001).

37.   Wester, J. C. & Contreras, D. Columnar Interactions Determine Horizontal Propagation of

975   Recurrent Network Activity in Neocortex. *J. Neurosci.* **32**, 5454–5471 (2012).

976   38.   Contreras, D., Destexhe, A., Sejnowski, T. J. & Steriade, M. Spatiotemporal patterns of
977   spindle oscillations in cortex and thalamus. *J Neurosci* **17**, 1179–1196 (1997).

978   39.   Xu, W., Huang, X., Takagaki, K. & Wu, J. Compression and Reflection of Visually
979   Evoked Cortical Waves. *Neuron* **55**, 119–129 (2007).

980   40.   Stieger, K. C., Eles, J. R., Ludwig, K. A. & Kozai, T. D. Y. In vivo microstimulation with
981   cathodic and anodic asymmetric waveforms modulates spatiotemporal calcium dynamics
982   in cortical neuropil and pyramidal neurons of male mice. *J. Neurosci. Res.* **98**, 2072–2095
983   (2020).

984   41.   Chen, E., Stiefel, K. M., Sejnowski, T. J. & Bullock, T. H. Model of traveling waves in a
985   coral nerve network. *J. Comp. Physiol. A* **194**, 195–200 (2008).

986   42.   Alekseichuk, I. *et al.* Electric field dynamics in the brain during multi-electrode
987   transcranial electric stimulation. *Nat. Commun.* **10**, 2573 (2019).

988   43.   Douglas, R. J., Koch, C., Mahowald, M., Martin, K. A. & Suarez, H. H. Recurrent
989   excitation in neocortical circuits. *Science (80-.).* **269**, 981–985 (1995).

990   44.   Haeusler, S. & Maass, W. A Statistical Analysis of Information-Processing Properties of
991   Lamina-Specific Cortical Microcircuit Models. *Cereb. Cortex* **17**, 149–162 (2006).

992   45.   Anderson, W. S., Kudela, P., Weinberg, S., Bergey, G. K. & Franaszczuk, P. J. Phase-
993   dependent stimulation effects on bursting activity in a neural network cortical simulation.
994   *Epilepsy Res.* **84**, 42–55 (2009).

995   46.   Raspopovic, S., Capogrosso, M. & Micera, S. A Computational Model for the Stimulation
996   of Rat Sciatic Nerve Using a Transverse Intrafascicular Multichannel Electrode. *IEEE*
997   *Trans. Neural Syst. Rehabil. Eng.* **19**, 333–344 (2011).

998   47.   Helmers, S. L. *et al.* Application of a computational model of vagus nerve stimulation.
999   *Acta Neurol. Scand.* **126**, 336–343 (2012).

48.   Greenberg, R. J., Velte, T. J., Humayun, M. S., Scarlatis, G. N. & De Juan, E. A
computational model of electrical stimulation of the retinal ganglion cell. *IEEE Trans.*
Biomed. Eng. **46**, 505–514 (1999).

49.   Traub, R. D., Jefferys, J. G. R., Miles, R., Whittington, M. A. & Tóth, K. A branching
dendritic model of a rodent CA3 pyramidal neurone. *J. Physiol.* **481**, 79–95 (1994).

50.   Radman, T., Ramos, R. L., Brumberg, J. C. & Bikson, M. Role of cortical cell type and
morphology in subthreshold and suprathreshold uniform electric field stimulation in vitro.
Brain Stimul. **2**, 215-228.e3 (2009).

51.   Radman, T., Su, Y., An, J. H., Parra, L. C. & Bikson, M. Spike Timing Amplifies the
Effect of Electric Fields on Neurons: Implications for Endogenous Field Effects. *J.*
Neurosci. **27**, 3030–3036 (2007).

52.   Lewis, P. M., Ackland, H. M., Lowery, A. J. & Rosenfeld, J. V. Restoration of vision in
blind individuals using bionic devices: A review with a focus on cortical visual prostheses.
Brain Res. **1595**, 51–73 (2015).

53.   Bennett, J. E. M. & Bair, W. Refinement and Pattern Formation in Neural Circuits by the

Interaction of Traveling Waves with Spike-Timing Dependent Plasticity. *PLOS Comput.*
Biol. **11**, e1004422 (2015).

54.   Beuter, A., Balossier, A., Vassal, F., Hemm, S. & Volpert, V. Cortical stimulation in
aphasia following ischemic stroke: toward model-guided electrical neuromodulation. *Biol.*
Cybern. **114**, 5–21 (2020).

55.   Liebetanz, D. *et al.* After-effects of transcranial direct current stimulation (tDCS) on
cortical spreading depression. *Neurosci. Lett.* **398**, 85–90 (2006).

56.   Santos, E. *et al.* Spreading depolarizations in a case of migraine-related stroke.
Cephalalgia **32**, 433–436 (2012).

57.   Batterink, L. J., Creery, J. D. & Paller, K. A. Phase of Spontaneous Slow Oscillations
during Sleep Influences Memory-Related Processing of Auditory Cues. *J. Neurosci.* **36**,
1401–1409 (2016).

58.   Kucewicz, M. T. *et al.* Evidence for verbal memory enhancement with electrical brain
stimulation in the lateral temporal cortex. *Brain* **141**, 971–978 (2018).

59.   Zhang, H., Watrous, A. J., Patel, A. & Jacobs, J. Theta and Alpha Oscillations Are
Traveling Waves in the Human Neocortex. *Neuron* **98**, 1269-1281.e4 (2018).

60.   Merrill, D. R., Bikson, M. & Jefferys, J. G. R. Electrical stimulation of excitable tissue:
Design of efficacious and safe protocols. *J. Neurosci. Methods* **141**, 171–198 (2005).

61.   Cogan, S. F., Ludwig, K. A., Welle, C. G. & Takmakov, P. Tissue damage thresholds
during therapeutic electrical stimulation. *J. Neural Eng.* **13**, 021001 (2016).

62.   Rall, W. Electrophysiology of a Dendritic Neuron Model. *Biophys. J.* **2**, 145–167 (1962).

63.   Tranchina, D. & Nicholson, C. A model for the polarization of neurons by extrinsically
applied electric fields. *Biophys. J.* **50**, 1139–1156 (1986).

64.   Berzhanskaya, J., Chernyy, N., Gluckman, B. J., Schiff, S. J. & Ascoli, G. A. Modulation
of hippocampal rhythms by subthreshold electric fields and network topology. *J. Comput.*
Neurosci. **34**, 369–389 (2013).

65.   Tehovnik, E. J., Tolias, A. S., Sultan, F., Slocum, W. M. & Logothetis, N. K. Direct and
Indirect Activation of Cortical Neurons by Electrical Microstimulation. *J. Neurophysiol.*
96, 512–521 (2006).

66.   Rattay, F. The basic mechanism for the electrical stimulation of the nervous system.
Neuroscience **89**, 335–346 (1999).

67.   Gustafsson, B. & Jankowska, E. Direct and indirect activation of nerve cells by electrical
pulses applied extracellularly. *J. Physiol.* **258**, 33–61 (1976).

68.   Swadlow, H. A. Monitoring the excitability of neocortical efferent neurons to direct
activation by extracellular current pulses. *J. Neurophysiol.* **68**, 605–619 (1992).

69.   Catterall, W. Localization of sodium channels in cultured neural cells. *J. Neurosci.* **1**,
777–783 (1981).

70.   Douglas, R. J., Martin, K. A. C. & Whitteridge, D. A Canonical Microcircuit for
Neocortex. *Neural Comput.* **1**, 480–488 (2008).

71.   DeFelipe, J. The neocortical column. *Front. Neuroanat.* **6**, 1–2 (2012).

72. da Costa, N. M. Whose cortical column would that be? *Front. Neuroanat.* (2010). doi:10.3389/fnana.2010.00016

73. Shepherd, G. M. G. Laminar and Columnar Organization of Ascending Excitatory Projections to Layer 2/3 Pyramidal Neurons in Rat Barrel Cortex. *J. Neurosci.* **25**, 5670–5679 (2005).

74. Staiger, J. F., Bojak, I., Miceli, S. & Schubert, D. A gradual depth-dependent change in connectivity features of supragranular pyramidal cells in rat barrel cortex. *Brain Struct. Funct.* **220**, 1317–1337 (2015).

75. Thomson, A. M., West, D. C., Wang, Y. & Bannister, A. P. Synaptic Connections and Small Circuits Involving Excitatory and Inhibitory Neurons in Layers 2-5 of Adult Rat and Cat Neocortex: Triple Intracellular Recordings and Biocytin Labelling In Vitro. *Cereb. Cortex* **12**, 936–953 (2002).

76. Bonjean, M. *et al.* Interactions between Core and Matrix Thalamocortical Projections in Human Sleep Spindle Synchronization. *J. Neurosci.* **32**, 5250–5263 (2012).

77. Staiger, J. F. *et al.* Functional diversity of layer IV spiny neurons in rat somatosensory cortex: Quantitative morphology of electrophysiologically characterized and biocytin labeled cells. *Cereb. Cortex* **14**, 690–701 (2004).

78. Schubert, D., Kötter, R., Luhmann, H. J. & Staiger, J. F. Morphology, electrophysiology and functional input connectivity of pyramidal neurons characterizes a genuine layer Va in the primary somatosensory cortex. *Cereb. Cortex* **16**, 223–236 (2006).

79. Muralidhar, S., Wang, Y. & Markram, H. Synaptic and cellular organization of layer 1 of the developing rat somatosensory cortex. *Front. Neuroanat.* **7**, 1–17 (2014).

80. Wang, Y., Gupta, A., Toledo-Rodriguez, M., Wu, C. Z. & Markram, H. Anatomical, physiological, molecular and circuit properties of nest basket cells in the developing somatosensory cortex. *Cereb. Cortex* **12**, 395–410 (2002).

81. Douglas, R. J. & Martin, K. A. A functional microcircuit for cat visual cortex. *J. Physiol.* **440**, 735–769 (1991).

82. Tomassy, G. S. *et al.* Distinct Profiles of Myelin Distribution Along Single Axons of Pyramidal Neurons in the Neocortex. *Science (80-.).* **344**, 319–324 (2014).

83. Traub, R. D., Wong, R. K., Miles, R. & Michelson, H. A model of a CA3 hippocampal pyramidal neuron incorporating voltage- clamp data on intrinsic conductances. *J Neurophysiol* **66**, 635–50. (1991).

84. White, E. L. & Keller, A. *Cortical circuits: synaptic organization of the cerebral cortex: structure, function, and theory*. (Springer, 1989).

1096 **SUPPLEMENTARY TABLES**

1097 **Table S1: Summary of datasets with reconstructed cells.**

Cell type	# of Reconstructions	Reference	Strain (Age)
Pyramidal cells (Layer II/III)	21	[74]	Wistar (P20-25)
Pyramidal cells (Layer IV)	11	[78]	Wistar (P19-21)
Pyramidal cells (Layer Va)	14	[79]	Wistar (P20-21)
Basket cells (Layer II/III)	96	[81]	Wistar (P13-15)
Basket cells (Layer IV)	82	[81]	Wistar (P13-15)
Basket cells (Layer V)	57	[81]	Wistar (P13-15)
Martinotti cells (Layer II/III)	13	[23]	Wistar (P13-16)
Martinotti cells (Layer V)	7	[23]	Wistar (P13-16)
Horizontal interneurons (LI)	59	[80]	Wistar (P13-16)
Descending interneurons (LI)	29	[80]	Wistar (P13-16)
Small interneurons (LI)	27	[80]	Wistar (P13-16)

1099 **Table S2: Model parameters for network simulations.**

Parameter	Description	Value		
C_m (μF/cm^2)	Membrane capacitance	1		
η	Standard deviation of spontaneous background activity	0.4		
V_{Na} (mV)	Na$^+$ reversal potential	50		
V_K (mV)	K$^+$ reversal potential	-90		
V_{Ca} (mV)	Ca^{2+} reversal potential	120		
V_{shp} (mV)	Slope of Ca^{2+} activation function	2.5		
V_{th} (mV)	Threshold potential	-25		
k_d	Equilibrium dissociation constant	1		
α_{Ca} (ms^{-1})	Ca^{2+} activation time constant	0.005		
τ_{Ca} (ms^{-1})	Ca^{2+} decay time constant	200		
α_s^{exc} (ms^{-1})	Excitatory synaptic activation time constant	0.6		
α_s^{inh} (ms^{-1})	Inhibitory synaptic activation time constant	0.5		
β_s^{exc} (ms^{-1})	Excitatory synaptic deactivation time constant	0.2		
β_s^{inh} (ms^{-1})	Inhibitory synaptic deactivation time constant	0.1		
V_{syn}^{exc} (mV)	Excitatory synaptic reversal potential	0		
V_{syn}^{inh} (mV)	Inhibitory synaptic reversal potential	-80		
Cell-type specific parameters		**PYs**	**BCs**	**MCs**
g_{Na} (mS/cm^2)	Na$^+$ channel conductance	30	60	60
g_K (mS/cm^2)	K$^+$ channel conductance	2.5	10	10
g_L (mS/cm^2)	Leak channel conductance	0.6	0.6	0.4
g_{Ca} (mS/cm^2)	Ca^{2+} channel conductance	0.1	0	0.1
g_{ahp} (mS/cm^2)	Slow Ca^{2+}-dependent (after-hyperpolarization) K$^+$ channel conductance	0.5	0	0.75
V_L (mV)	Leak channel reversal potential	-67	-65	-65

1101 **Table S3: Structure of the network.**

Cortical cell type	Layer	Cells/column	Total cells
IN	I	12	132
PY	II/III	100	1100
BC	II/III	100	1100
MC	II/III	24	264
PY	IV	12	132
BC	IV	12	132
MC	IV	12	132
PY	V	12	132
BC	V	12	132
MC	V	12	132
total		308	3388

1103 **Table S4: Connectivity within the network.**

Presynaptic		Postsynaptic		Cross-column	Type	Strength	Probability
Type	Layer	Type	Layer				
PY	II/III	PY	II/III	TRUE	AMPA	0.4	0.1
PY	IV	PY	IV	TRUE	AMPA	0.4	0.1
PY	V	PY	V	TRUE	AMPA	0.4	0.1
PY	II/III	PY	II/III	FALSE	AMPA	0.75	0.1
PY	II/III	BC	II/III	FALSE	AMPA	0.75	0.1
PY	II/III	MC	II/III	FALSE	AMPA	0.75	0.1
PY	II/III	PY	IV	FALSE	AMPA	0.25	0.05
PY	IV	PY	II/III	FALSE	AMPA	1.5	0.05
PY	IV	PY	IV	FALSE	AMPA	0.75	0.1
PY	IV	BC	IV	FALSE	AMPA	0.75	0.1
PY	IV	MC	IV	FALSE	AMPA	0.75	0.1
PY	IV	BC	II/III	FALSE	AMPA	1.5	0.05
PY	IV	MC	II/III	FALSE	AMPA	1.5	0.05
PY	V	PY	V	FALSE	AMPA	0.75	0.1
PY	V	PY	II/III	FALSE	AMPA	0.5	0.05
PY	V	BC	V	FALSE	AMPA	0.75	0.1
PY	V	MC	V	FALSE	AMPA	0.75	0.1
MC	II/III	PY	II/III	FALSE	GABA	0.75	0.05
MC	II/III	PY	IV	FALSE	GABA	0.75	0.05
MC	II/III	PY	V	FALSE	GABA	0.75	0.05
MC	IV	PY	II/III	FALSE	GABA	0.75	0.05
MC	IV	PY	IV	FALSE	GABA	0.75	0.05
MC	IV	PY	V	FALSE	GABA	0.75	0.05
MC	V	PY	II/III	FALSE	GABA	0.75	0.05
MC	V	PY	IV	FALSE	GABA	0.75	0.05
MC	V	PY	V	FALSE	GABA	0.75	0.05
BC	II/III	PY	II/III	FALSE	GABA	1.5	0.25
BC	IV	PY	IV	FALSE	GABA	1.5	0.25
BC	V	PY	V	FALSE	GABA	1.5	0.25
MC	V	PY	IV	FALSE	GABA	1	0.2
MC	V	PY	V	FALSE	GABA	1.5	0.25
IN	I	PY	II/III	FALSE	GABA	0.3	0.3
IN	I	PY	IV	FALSE	GABA	0.3	0.3
IN	I	PY	V	FALSE	GABA	0.3	0.3
PY	II/III	IN	I	FALSE	AMPA	0.3	0.3