1 **Brain transcriptomic changes in Japanese quail reveal roles for neurotensin and**
2 **urocortin 3 in avian parental care**
3

Patricia C. Lopes[*] and Robert de Bruijn

Schmid College of Science and Technology, Chapman University, Orange, California,
USA

* Corresponding author
Patricia C. Lopes
lopes@chapman.edu

Abstract

For many species, parental care critically affects offspring survival. But what drives animals to display parental behaviours towards young? In mammals, pregnancy-induced physiological transformations seem key in preparing the neural circuits that lead towards attraction (and reduced-aggression) to young. Beyond mammalian maternal behaviour, knowledge of the neural mechanisms that underlie parental care is severely lacking. We took advantage of a domesticated bird species, the Japanese quail, for which parental behaviour towards chicks can be induced through a sensitization procedure, a process that is not effective in all animals. We used the variation in parental responses to study neural transcriptomic changes associated with the sensitization procedure itself and with the outcome of the procedure (i.e., presence of parental behaviours). Out of the brain regions studied, we found that most differences in gene expression were located in the hypothalamus. Two genes identified are of particular interest, as no role in avian parental care was known for those genes. One is neurotensin, previously only demonstrated to be causally associated with maternal care in mammals. The other one is urocortin 3, causally demonstrated to affect young-directed neglect and aggression in mammals. Our work opens new avenues of research into understanding the neural basis of parental care in non-placental species.

Keywords: avian parental care, bed nucleus of the stria terminalis, hypothalamus, Neurotensin, nucleus taeniae, prolactin, RNA-seq, Urocortin 3

1. Introduction

Most of what is known regarding the mechanistic basis of infant-directed parental care comes from studies in mammals [1]. In mammals, pregnancy-induced physiological transformations seem key in preparing the neural circuits that lead towards attraction (and reduced-aggression) to an infant [1]. However, not all parental care is explained by pregnancy-induced changes. For example, there is also paternal care, alloparenting (parental care towards non-descendant young including adoption) and care by non-placental animals [2]. Knowledge of what neurotransmitters/neuromodulators are involved in these situations is lacking. While these forms of care are widespread in human societies [3] and in non-placental vertebrates [4], and play critical roles in offspring development [5,6], their neural underpinnings are severely understudied. Studying species that show parental care in the absence of pregnancy can therefore help elucidate the neurochemical make-up of these diverse forms of parental care.

While parental care is rare in many vertebrate taxa, it is essential for survival in the majority of avian species [7]. Although biparental care is the most frequent modality observed, birds as a group show an incredible diversity in forms of parental care, from brood parasitism, where care is lost, to cooperative breeding, where non-reproductive individuals provide care to offspring that are not their own [7]. In terms of neuroendocrine mechanisms associated with incubation and caring for young, the majority of attention has been paid to a single pituitary hormone: prolactin [8,9]. Coupled with an increase in prolactin, a decline in gonadal hormones seems to facilitate the transition between courtship and nest building into incubation behaviour [10]. The neural underpinnings of young-directed parental care in birds are, however, severely understudied.

Using a sensitization procedure developed by [11], a recent experiment by our group confirmed that it is possible to induce parental behaviours in Japanese quail through a single overnight exposure to chicks [12]. After the sensitization procedure, both sexes display chick brooding behaviour and reduced aggression towards chicks (pecking

behaviour). This procedure therefore allows us to study young-directed care at its onset, in both sexes, in the absence of pregnancy-induced effects, copulation or pharmacological hormonal priming. What is also interesting is that, while the change in behaviour after sensitization is drastic, it is not observed in all animals, i.e., some animals fail to respond to the sensitization. Conversely, a few animals not exposed to the sensitization spontaneously show some parental care when paired with chicks.

In the current study, we take advantage of this variation in parental care to understand, from a neural molecular perspective, a) what modifications occur due to the sensitization treatment, b) what genes correlate with the duration of brooding behaviour and with chick-directed aggression, and c) whether the same mechanisms are involved in care that is shown after sensitization treatment relative to care that is shown spontaneously (i.e., in the absence of sensitization treatment). We focused our transcriptomic analysis on the hypothalamus, the bed nucleus of the stria terminalis and the nucleus taeniae. We targeted these brain regions because they contain nuclei important for parental care, affiliative interactions, and agonistic interactions in birds and other vertebrates [2,4,13–16]. Our experiments highlight the importance of using intra-species variation in parental responses to elucidate neural mechanisms of parental behaviour and to discover new molecular pathways involved in avian parental care, several of which are likely conserved across other vertebrate taxa.

2. Materials and methods

(a) Parental care induction procedure and behaviour quantification

Japanese quail (*Coturnix japonica*) were raised in the lab from eggs obtained from AA Lab Eggs, Inc. (Westminster, CA) following the procedures described in [12]. Adult birds used in this experiment were kept in an 8L:16D light cycle, at a temperature range of 20-24 °C and maintained in groups of 4-5 same sex animals in cages (100 × 40 × 50 cm). We chose a short photoperiod for this experiment for two reasons: 1) male and female Japanese quail kept at short photoperiods are reproductively quiescent [17],

which reduces the influence of gonadal hormones from the study; and 2) quail

sensitization treatments are more effective at increasing brooding duration in both sexes

under short photoperiods [11,16]. The reason for more effective sensitization when

gonadal hormones are low may be linked to the natural decline in gonadal hormones

undergone by birds transitioning between courtship and nest building to a parental state

(onset of incubation) (reviewed in [10]). Two days prior to carrying out the experiments,

a replica of the wooden box (18 × 18 × 18 cm) used in the parental care sensitization

procedure was added to those cages. The day prior to the procedure, animals were

separated individually into cages identical to the group cages, which also contained a

wooden box. The wooden box was always open during these days, allowing the birds to

inspect it and go in and out. Food and water were provided *ad libitum* during the entire

experiment.

The day of the parental care sensitization procedure, each adult bird was locked inside

the wooden box in their individual cage starting at one hour before lights were off. Under

the parental induction treatment, two chicks (1-3 days old) were added to this box just

before lights off. Under the control treatment, no chicks were added to the box. Animals

were left undisturbed overnight. The morning after, the wooden box was opened, the

overnight chicks were removed, and two new chicks (1-3 days old) were added to the

cage. This swap of chicks ensured that birds in both treatments (controls and

sensitized) were being tested for their behaviours towards novel chicks (in case

familiarity influenced the outcome). The novel chicks were allowed to stay in the cage

for 20 min, during which videos were continuously recorded (Axis M1065L network

camera by Axis Communications). At the end of the 20 min, adults were euthanized by

isoflurane inhalation and decapitation. The brain was removed from the skull, flash

frozen and then stored at -80 °C until further processing. Chicks were returned to their

cages. In total, 50 adults (ages 90-110 days old) were subjected to the experimental

procedures, of which 11 females and 13 males were randomly assigned to the

sensitization treatment and 19 females and 7 males to the control treatment. Two

control females were removed from the study due to methodological failures during the

experiments, bringing the final number of control females to 17. In total, 77 chicks were

used. Behaviours towards chicks within the 20 min period of observation were scored

by observers blind to the treatment and analysed as described in [12]. The behavioural

results described here were re-plotted from [12] and were not re-analysed.

(b) Brain dissection

Brains were coronally sectioned on a Leica CM1860UV cryostat, at -18 °C. We used

separate surgical micropunches (EMS Rapid Core Instruments) to punch out the three

brain regions of interest from 100 μm slices, spaced apart by three 30 μm slices

collected onto microscope slides (Fisherbrand, item 12–550-15) for future use. The

brain regions collected were the entire hypothalamus, the nucleus taeniae (Tn) and the

bed nucleus of the stria terminalis (BnST). These regions were identified based on use

of both the quail and the chicken brain atlases [18,19]. The diameter of the

micropunchers used was 4 mm for the hypothalamus and 3 mm for the other regions.

The hypothalamus was collected from the start of the bifurcation the tractus

septomesencephalicus until the start of the posterior commissure. We started collecting

the BnST from the point when the anterior commissure was visible, using the lateral

ventricle as a guide for positioning the micropuncher, and ended after collecting 5

punches, i.e., over 1 cm coronally (as each punch is separated from the next by about

190 mm). Collection of the Tn started at the point where the occipitomesencephalic tract

splits into two and ended after 5 punches (again equivalent to 1 cm coronally). Punches

from each of these brain regions were placed in individual tubes containing 2 mm size

beads (ZR BashingBeads Lysis Tubes, Zymo Research, item S6003-50) and 1 mL of

QIAzol lysis reagent (Qiagen, item #79306). Tissue homogenization was done by

agitating the tubes for 20 s at a $7\,m\,s^{-1}$ speed (Beadbug 6 homogenizer,

Benchmark Scientific), followed by a 5 min rest period. The homogenate was then

transferred into a new tube and preserved at −80 °C until RNA isolation.

(c) RNA isolation, library preparation and sequencing

Total RNA was extracted from the aqueous layer formed after chloroform precipitation,

using the RNA Clean & Concentrator Kit-5 (Zymo Research, item # R1013) following

manufacturer's instructions and including the DNase I in-column treatment step. These

samples were then shipped on dry ice to Novogene Corporation Inc. (Chula Vista, CA,

USA), where RNA quantity, purity and integrity were assessed on an Agilent 2100

Bioanalyzer (Agilent Technologies, Santa Clara, CA), cDNA libraries were generated

using NEBNext® UltraTM RNA Library Prep Kit for Illumina® (NEB, USA) and cDNA

fragments (150 ~ 200 bp in length) were then purified using the AMPure XP System

(Beckman Coulter, Beverly, USA). Paired-end sequencing of libraries (PE150; Illumina

Novaseq 6000) was performed according to standard protocols. An average of 52

million paired-end raw reads were obtained for each sample.

(d) Mapping and differential gene expression analysis

An average of 85.3% of clean (post adapter removal and quality filtering) reads were

mapped to the Japanese quail reference genome (Coturnix_japonica_2.0, INSDC

Assembly Mar 2016, downloaded from Ensembl), representing an average of 43.7

million mapped reads per sample (Table S1 contains information on mapping statistics

per sample). Mapping was done using HISAT2 [20], and HTSeq was used to count the

number of mapped reads to each gene [21]. Differential gene expression analysis for

the effect of treatment was performed using the DESeq2 R package [22,23]. To

determine whether any genes showed significant correlations to either brooding

duration or aggressive events, we used edgeR TMM-normalized counts (edgeR

package [24]) with the limma package [25], with voom transformation [26]. While the

DESeq2 package can also accept continuous covariates in the model, the list of

significant genes detected using the limma+voom approach overlaps with the one

produced by DESeq2 but excludes genes for which the regression line does not

properly fit the points when plotted. To control for the false discovery rate due to

multiple testing, p-values were adjusted using the Benjamini-Hochberg procedure.

Genes were considered as statistically differentially expressed when adjusted p-values

were $<$ 0.05.

To examine the effect of treatment on gene expression, for each brain region, we first

tested whether the variable sex interacted with the variable treatment. In case no

differentially expressed genes (DEGs) were detected for the interaction term at $p_{adj} <$

0.05, the variable sex was excluded from the model and treatment was tested alone. To

examine covariation between gene expression levels and our two continuous variables

(brooding duration or aggression), for each brain region, we first tested for an interaction

of treatment or sex with each continuous covariate. In case no DEGs were detected for

the interaction term at $p_{adj} < 0.05$, the variable treatment was excluded from the model

and the continuous covariates were tested alone. Prior to testing, the continuous

covariates were scaled and centered, but figures show brooding duration in seconds for

simplicity of interpretation. Volcano plots for the results of treatment effects were

produced using log2 transformed DESeq2 normalized counts. A subset of

representative genes obtained from the limma-voom analysis were represented by

plotting log2 voom transformed TMM normalized counts, and a regression line with a

slope of the logFC and the coefficient for the intercept for each specific gene.

(e) Gene ontology (GO) enrichment analysis and KEGG pathway enrichment

analysis of differentially expressed genes

Functional enrichment analysis was done only for the DEGs found in the hypothalamus,

as this region contained the most DEGs both for the effect of treatment and for the

covariation with parental duration. The stringApp (v. 1.5.4) [27] within Cytoscape

software (v. 3.8.2) [28] was used to find enriched GO terms and enriched KEGG

Pathways. Up- and down-regulated DEGs were analysed separately. *Mus musculus*

was used as the background species, and the significance threshold for the false

discovery rate (FDR) was 0.05. We used the same app and settings to visualize

possible protein-protein interactions (PPI) among the protein products of these DEGs,

but this time we analysed all genes (up and down-regulated) at once. This analysis

produces a PPI enrichment p-value, which, when smaller or equal than 0.05, indicates

that the proteins being tested have more interactions among themselves than would be
expected for a random set of proteins of similar number, drawn from the genome.
220
221 **3. Results**
222
223 **(a) Effect of sensitization treatment**
224
Behavioural response. In response to spending an overnight enclosed in a box with
two chicks (sensitization treatment), the average duration of time spent brooding novel
chicks was significantly increased and not different by sex (Fig. 1a, [12]). Out of animals
in the sensitization treatment, 3 out of 11 females (27.3 %) and 3 out of 13 males (23.1
%) showed no brooding behaviour. Out of control animals, 6 out of 16 females (37.5 %
of females) and 3 out of 8 males (37.5 % of males) showed at least a few seconds of
brooding behaviour. Aggression, quantified as chick-directed pecking, was overall low
and while it tended to be lower in animals in the sensitization treatment, this difference
was not significant (Fig. 1b, [12]).
234

Figure 1. Brooding duration (a) and aggressive events (chick-directed pecking) (b) in

animals that underwent sensitization treatment or not (control). Dots represented

individual animals, coloured by sex.

Transcriptomic response. Sex did not interact with treatment in affect gene

expression in any of the brain regions. Therefore, the effect of treatment was tested in

both sexes together. Very few genes were differentially expressed in the brains of

animals that underwent the sensitization treatment relative to controls during interaction

with chicks (Fig. 2, Table S2). In the hypothalamus, only two genes were up-regulated

in the sensitization treatment relative to control: neurotensin (NTS) and urocortin 3

(UCN3). In the bed nucleus of the stria terminalis (BnST), only one gene was up-

regulated: adipocyte plasma membrane associated protein (APMAP). This same gene

was the only one that approached significance (padj = 0.07229) in the Nucleus Taeniae

(Tn). Out of the six genes down-regulated in the hypothalamus of animals that

underwent the sensitization treatment, three overlapped with down-regulated genes in

the BnST: Fos proto-oncogene, AP-1 transcription factor subunit (FOS), early growth

response 4 (EGR4) and BTG anti-proliferation factor 2 (BTG2). In addition to these

three, another five genes were down-regulated in the BnST: Kruppel like factor 2

(KLF2), neuronal PAS domain protein 4 (NPAS4), RRAD, Ras related glycolysis

inhibitor and calcium channel regulator (RRAD), salt inducible kinase 1 (SIK1) and Jun

proto-oncogene, AP-1 transcription factor subunit (JUN).

Figure 2. Volcano plots representing the $\log_2$ fold change against the corresponding –

$\log_{10}$ p_{adj}-value of all genes for each of the three brain regions studied. Differentially

expressed genes ($p_{adj} < 0.05$) are shown in red if they are up-regulated in the

sensitization treated animals relative to control and in blue if they are down-regulated.

Other genes are shown in black. The horizontal dotted line indicates the p_{adj}-value cut-

off, while the vertical dotted line marks the point for zero fold change differences. One

down-regulated gene without annotation is not show for hypothalamus (see Table S2 for

complete list).

(b) Genes and pathways associated with the duration of brooding behaviour

The only brain region where genes showed a significant correlation with the duration of

brooding behaviour was the hypothalamus. Twelve genes in this region were

significantly correlated with brooding behaviour (Table S3). With the exception of

transducer of ERBB2, 1 (TOB1), a down-regulated gene, all other genes overlap with

DEGs detected in the hypothalamus or BnST. Furthermore, DEGs that were up-

regulated in animals in the sensitization treatment relative to control increase in

expression as brooding duration increases (UCN3, Fig. 3) and down-regulated genes

decrease in expression as brooding duration increases (e.g., FOS, Fig. 3; other down-

regulated genes show the same pattern). No genes were correlated with aggressive

events in any of the brain regions studied. No interaction between brooding duration or

aggressive events and sex was found. We also found no interaction between treatment

and parental duration or treatment and aggressive events, indicating treatment did not

affect the expression of genes related to either of these traits. In other words, genes that

covary with parental duration are the same between birds in the sensitization treatment

and birds in the control treatment and, therefore, the molecular mechanisms behind

spontaneous care and care after undergoing the sensitization treatment are similar.

Figure 3. Regression lines representing the relationship between brooding duration

(sex) and FOS expression (left panel) or UCN3 expression (right panel). Statistical

details for all genes that covary with brooding duration can be found in Table S3.

To gain further insight into what pathways might be involved in brooding behaviour, we

used GO enrichment analysis and KEGG pathway enrichment analysis (summary of

results in Table I; full results in Tables S4 and S5). The GO enrichment analysis of

DEGs that negatively covary with brooding behaviour revealed enrichment in several

GO categories related with transcriptional regulation (most significant terms across all

three GO categories), response to hormones (examples of terms with non-redundant

gene combinations: response to corticosterone, response to steroid hormone, response

to progesterone, cellular response to hormone steroid stimulus, response to peptide

hormone) and with learning or memory (Table S4). In terms of genes that were either

up-regulated by the sensitization treatment (UCN3 and NTS) or that positively covaried

with brooding duration (UCN3), enriched GO terms were related to neuropeptide

307 hormone activity (MF category) and with axon terminus and extracellular region (CC
308 category).
309
310

Source	Term name	FDR	Down-regulated genes
GO:MF	transcription regulator activity	1.58E-05	FOSB, BTG2, FOS, NR4A3, TOB1, NPAS4, EGR1
GO:MF	DNA-binding transcription activator activity...	1.58E-05	FOSB, FOS, NR4A3, NPAS4, EGR1
GO:MF	DNA-binding transcription factor activity...	1.58E-05	FOSB, BTG2, FOS, NR4A3, NPAS4, EGR1
GO:MF	RNA polymerase II transcription regulatory...	1.58E-05	FOSB, FOS, NR4A3, NPAS4, EGR1, EGR4
GO:MF	RNA polymerase II core promoter...	5.30E-04	FOS, EGR1
GO:MF	histone acetyltransferase binding	5.60E-04	NR4A3, EGR1
GO:MF	SMAD binding	3.20E-03	FOS, TOB1
GO:MF	heterocyclic compound binding	6.40E-03	FOSB, FOS, NR4A3, RRAD, NPAS4, EGR1, EGR4
GO:BP	positive regulation of RNA metabolic process	1.97E-05	FOSB, BTG2, FOS, NR4A3, TOB1, NPAS4, EGR1, EGR4
GO:BP	response to corticosterone	2.65E-05	FOSB, FOS, NPAS4
GO:BP	response to steroid hormone	2.20E-04	FOSB, FOS, NR4A3, NPAS4
GO:BP	cellular nitrogen compound biosynthetic process	2.20E-04	FOSB, BTG2, FOS, ST8SIA2, NR4A3, NPAS4, EGR1, EGR4
GO:BP	skeletal muscle cell differentiation	2.20E-04	BTG2, FOS, EGR1
GO:BP	positive regulation of gene expression	2.20E-04	FOSB, FOS, NR4A3, TOB1, NPAS4, EGR1, EGR4
GO:BP	learning or memory	2.20E-04	BTG2, FOS, NPAS4, EGR1
GO:BP	behavior	2.20E-04	BTG2, FOS, NR4A3, NPAS4, EGR1
GO:BP	transcription, DNA-templated	2.20E-04	FOSB, BTG2, FOS, NR4A3, NPAS4, EGR1, EGR4
GO:BP	regulation of neuron death	2.80E-04	BTG2, FOS, NR4A3, EGR1
GO:CC	transcription regulator complex	2.53E-02	FOS, NR4A3, NPAS4
KEGG	Amphetamine addiction	1.65E-02	FOSB, FOS
KEGG	IL-17 signaling pathway	1.65E-02	FOSB, FOS
KEGG	Osteoclast differentiation	1.65E-02	FOSB, FOS
KEGG	RNA degradation	1.65E-02	BTG2, TOB1
Source	Term name	FDR	Up-regulated genes
GO:MF	neuropeptide hormone activity	2.02E-05	NTS, UCN3
GO:CC	axon terminus	2.00E-03	NTS, UCN3
GO:CC	extracellular region	2.90E-02	NTS, UCN3

311
312
Table I. Significantly enriched GO terms and KEGG pathways associated with down- or
up-regulated genes in the hypothalamus. Only top 10 terms per category are shown.
For MF and BP, filtering of terms for redundancy of enriched genes was done
(maximum Jaccard similarity of 1 allowed). Several term names are cropped for space
(see Tables S4 and S5 for full list of unfiltered terms and complete term names). MF =
Molecular function; BP = Biological Process; CC = Cellular Component.
319

In order to understand how the protein products of the DEGs may interact, if at all, we
produced a protein-protein interaction network using the StringApp plugin within
Cytoscape. The network produced showed that the majority of down-regulated genes
interacted with one another and that NTS may interact with FOS (Fig. 4). The PPI
enrichment p-value for this set of proteins is 1×10^{-16}, indicating a larger degree of
interaction among these proteins than would be expected by chance.
326
327

328
Figure 4. Protein-protein interaction network produced with STRING. Nodes with blue
colours have lower expression in animals showing increased parental behaviours.
Nodes with red colours have higher expression in animals in the sensitization treatment
(both NTS and UCN3) or in animals showing increased parental behaviours (UCN3
only). The width of the edges (arrows) corresponds to the stringdb score, which
measures how likely the interactions between two given proteins are to be true given
the available evidence. In our network, we only show edges with medium (0.4) or better
confidence.
337
338

4. Discussion

The neural mechanisms mediating parental care remain poorly understood. We took advantage of a domesticated species, the Japanese quail, for which parental behaviour towards chicks does not tend to occur spontaneously in captivity. Parental behaviour in these animals can be induced through a sensitization procedure, but the process is not 100 % effective. Using the variation in parental responses, we asked what neural molecular mechanisms: a) are altered upon sensitization treatment, b) correlate with the amount of parental response shown, and c) differ between care shown after the sensitization procedure relative to care shown in the absence of this procedure (spontaneous care). Out of the three brain regions studied, the only one where gene expression patterns correlated with the duration of brooding behaviour was the hypothalamus. Our unbiased transcriptomic approach identified two genes in this region, neurotensin and urocortin 3, that were previously only demonstrated to be causally associated with young-directed care and young-directed aggression in mammals. Transcriptomic studies focused on animals that fail to show parental care under care-promoting circumstances, or studies on those animals that show care unprompted provide great opportunities to understand the neural basis of variation in parental care.

By comparing the neural transcriptomic responses of adult quail that had undergone a sensitization procedure to those that did not undergo this process, we were able to identify genes that may be involved in activating avian parental care. One of those genes, neurotensin, was expressed at higher levels in the hypothalamus of both females and males after sensitization treatment. This finding is significant given that neurotensin levels in the hypothalamus of postpartum female mice were also observed to be elevated relative to that of virgin mice [29]. Neurotensin gene expression was also recently shown to be elevated in the preoptic area of males of one species of poison frog (*Dendrobates tinctorius*) when performing parental care relative to males not performing care [30]. Furthermore, intracerebroventricular injections of neurotensin significantly decreased maternal aggression in female mice [31], indicating the

neurotensin participates in at least one component of maternal responses in rodents.
Previously, the only reports of behavioural effects of neurotensin on birds suggested a
possible relationship between neurotensin expression in the medial preoptic nucleus
(POM) and non-vocal courtship behaviour and agonistic behaviour in male European
starlings (*Sturnus vulgaris*) [32]. Interestingly, that study also reported that male
starlings that had acquired a nest box also had higher neurotensin expression in the
POM relative to males that did not acquire boxes and the measures of non-vocal
courtship behaviour were mostly related to nest building (e.g., such as number of times
entering or landing on the nest and gathering nesting material). European starlings
show biparental care and the increase in neurotensin in a hypothalamic nucleus in that
study could have been related to nest-building activity, which, for many species, is a
component of parental behaviour.
382
Out of the hormones associated with avian parental care to date, prolactin is by far the
one that has received the most attention [9]. In precocial avian species, as is the case
with Japanese quail, prolactin tends to be elevated in females during the incubation
period and to decline when chicks hatch [8]. Exposure to prolactin maybe therefore be
needed to promote young-directed care. Prolactin release is known to be under the
control of dopamine, which can have both inhibitory and stimulatory effects depending
on the dopamine receptor subtype it acts on [9]. Neurotensin is capable of enhancing
dopamine release [33] and could, in this way, impact parental behaviour by playing a
role in controlling prolactin secretion. Neurotensin could also affect care through its
involvement in reward processing [34].
393
Galanin is a neuropeptide that has emerged as an important regulator of parental
behaviour [35]. While galanin was not detected in our study, it is interesting to note that
neurotensin neurons in the medial preoptic area of the hypothalamus overlap with
galanin neurons in this region [36]. This overlap may be suggestive of overlapping
functions for the neurotensin neurons. It is also important to highlight that whereas
neurotensin was elevated in birds exposed to the sensitization treatment, it did not
covary with the extent of parental behaviour shown. A recent study that examined

neurogenomic responses in paternal sticklebacks (*Gasterosteus aculeatus*) across

several stages of care found that galanin, for example, was elevated in early stages

(prior to hatching) but no different from controls once all eggs had hatched [37],

highlighting that the role of different neuromodulators changes with the stage of care.

Galanin may therefore be more important at a different stage of parental care in birds.

Also, further studies will therefore be needed to understand whether neurotensin

participates only in switching on parental behaviours (as is indicated by our study) or

whether it is needed for the maintenance of those behaviours.

The nonapeptides oxytocin and vasopressin have been implicated in the regulation of

social bonding and maternal care in mammals [1]. In birds, two studies (one in female

turkeys, *Meleagris gallopavo* [38] and one in female Native Thai chickens, *Gallus*

domesticus [39]) focused specifically on the transitions from egg-laying to incubation

and from incubation to brooding behaviour, showed similar patterns in terms of number

of neurons producing mesotocin (the avian homologue to oxytocin). Both studies found

that the number of mesotocin-immunoreactive neurons was higher in specific

hypothalamic nuclei of incubating than of non-incubating laying hens. While in the first

chicken study showed no major differences in mesotocin-immunoreactive neurons

between incubating and chick-brooding hens, a subsequent study on the same species

found that hens that had been incubating eggs naturally and had their eggs swapped by

chicks, had higher numbers of neurons showing mesotocin-immunoreactivity than hens

that continued incubating eggs and received no chicks [40]. Combined, these studies

suggest that mesotocin is important for transitions between parental care stages in

some bird species. As we did not detect differences in mesotocin in our study, it may be

that animals need to be actively laying or incubating eggs to express these changes in

mesotocin. It will be interesting for future studies to explore these relationships further to

understand the precise roles played by mesotocin in avian parental care behaviours.

The only other gene that was upregulated in the hypothalamus following the

sensitization procedure was urocortin 3. Urocortin 3 was also the only gene in the

hypothalamus to be positively correlated with brooding duration. In rodents, central

administration or overexpression of urocortin 3 increases stress-induced anxiety and

suppresses ingestive behaviour [41,42]. We are only aware of two reports relating

urocortin 3 to young-directed parental care. One is the poison frog study mentioned

earlier [30] where urocortin 3 was upregulated in the preotic area of frogs performing

parental care relative to non-parental ones. The other one revealed that urocortin 3

expressing neurons in the perifornical area of the hypothalamus are activated

specifically during infant-directed aggression in both male and female mice and that

optogenetic activation of these neurons triggers infant-directed aggression and neglect

[43]. While aggression levels were not significantly different between our two

treatments, they tended to be reduced after sensitization treatment. Furthermore,

urocortin 3 expression did not covary with aggressive behaviour in our study. Rather

than affecting infant-directed aggression in our system, urocortin 3 may serve different

functions that support parental care, such as altered feeding patterns [44,45].

Interestingly, recent work proposes that molecules related to feeding behaviour may

also be involved in complex social behaviours, such as parental care [46,47].

In the BnST only one gene was up-regulated in animals exposed to the sensitization

procedure: APMAP. This was also the only gene up-regulated in the Tn if we consider a

$p_{adj} < 0.1$ cut-off. The protein encoded by APMAP is important for white adipose tissue

differentiation [48] and little is known regarding it's neurobiological functions, except that

is may be involved in suppressing amyloid-beta formation in brain [49]. The connection

to parental care, if any, is unknown.

As no significant differences in gene expression was found between treatments in the

Tn and no genes in this region showed covariation with brooding behaviour, this region

is likely less critically important during the onset or production of brooding behaviour in

birds.

Most of the differentially expressed genes that were down-regulated in the

hypothalamus and BnST in animals that underwent the sensitization procedure, as well

as those genes that negatively covaried with the expression of brooding behaviour, are

considered immediate early genes [50] and are involved in regulation of transcription

(FOSB, BTG2, FOS, NR4A3, TOB1, NPAS4, EGR1, EGR4, KLF2, JUN, SIK1) and in

response to hormones (FOSB, FOS, NR4A3, NPAS4, JUN, SIK1). Very similar changes

in immediate early gene expression patterns were found in several brain regions

(including the hypothalamus; the BnST was not included in that study) in female mice,

whereby immediate early gene expression was decreased in females during pregnancy

and post-partum relative to virgin females [51]. One of the top enriched pathways

highlighted through enrichment analysis was for response to corticosterone.

Corticosterone is one of the main hormones involved in the response to stress in birds

[52]. Data from a separate study on these birds showed that there were no differences

between treatments in terms of serum corticosterone levels after the birds had been

exposed to the novel chicks [12]. One possibility is that the way in which the

hypothalamus and BnST respond to corticosterone changes in parental animals.

Prolactin, for instance, has been implicated in the modulation of the stress response

[53]. Reduced stress responsiveness may facilitate the acceptance of hatchlings in

natural settings, which may otherwise be perceived as intruders. Alternatively, it is

possible that the pathways highlighted reflect altered responsiveness to other hormones

or peptides. For example, while central injection of neurotensin can increase c-Fos

immunoreactivity in several brain nuclei (including hypothalamic nuclei) in non-parental

rats [54], two studies that compared immediate early gene responses to neurotensin

administration in maternal versus virgin female mice found a general decrease in

reactivity (decreased Egr-1 and c-Fos production) in maternal females [55,56].

A previous study in female Japanese quail found that sensitized maternal females had

increased c-Fos immunoreactivity in the BnST relative to sensitized non-maternal

females [16]. This appears to be in contrast with our finding of higher FOS expression in

the control relative to the sensitization treatment in this brain region. This difference may

be related to the time point of parental care being studied and details of each study. In

their study, the sensitization treatment consisted of exposure to chicks for 20 min

periods over 5 consecutive days. The study of immediate early gene immunoreactivity

was only done after the 5th day. Also, on the 5th day, after being allowed to interact with

the chicks for 20 min, females were separated from the test chicks for 1 h prior to brain
collection. These two major differences alone could result in different outcomes in our
studies. The [16] study may be quantifying parental behaviour after is has been
established, while our study reflects the onset of parental behaviour. Additionally, as c-
Fos protein can be detected at 20 to 90 min after the stimulus [57], the separation from
the chicks for 1 h prior to sample collection could have reflected responses related to
maternal separation rather than maternal behaviour.
501
Our study indicates that both sexes possess the neural circuitry needed to produce
brooding behaviour and that the changes in gene expression associated with this
behaviour do not differ per sex. It is, however, important to note that our study was done
under a background of low levels of gonadal hormones (short photoperiod), which may
make males and females less different. It will therefore be interesting in future studies to
assess how the photoperiod affects genomic responses associated with parental care
behaviour and whether it introduces differences between the sexes.
509
510 **5. Conclusion**
511
Recent studies have used transcriptomic approaches to compare gene expression in
specific brain areas as female birds transition from territorial defence to egg incubation
[58], or from egg-laying to incubation behaviour [59]. Still, the range of neural
mechanisms involved in post-hatching care in birds is virtually unknown. Our findings
revealed new potential modulators of avian young-directed parental care, which deserve
to be examined through further mechanistic studies and explored in other bird species
and other taxonomic groups. Particularly exciting are neurotensin and urocortin 3, which
potentially show conserved roles in parental care in mammals, amphibians, and birds,
but are extremely under-explored in the context of parental behaviour. Natural variation
in avian parental behaviour in response to young extends from abandoning the nest to
alloparenting or even adoption of non-offspring deposited by brood parasites. The
genes found to be associated with the onset of parental care and with variation in
young-directed care in our study serve as a foundation for future studies aimed at

investigating the neural underpinnings of variation in parental care within and across
species.
527
528 **Ethics**

529 Animal use and experimental design were approved by the Chapman University
530 Institutional Animal Care and Use Committee (protocol # 2019-01).
531

Data accessibility

RNA-seq mapping statistics, lists of DEGs and lists of enrichment terms have been
uploaded as part of the supplementary material. Sequencing datasets generated and
analysed during the current study are deposited in the NCBI Gene Expression Omnibus
(GEO) repository, with record GSE162378 [dataset will be made publicly available once
manuscript is accepted for publication].
538

539 **Authors' contributions**

540 PCL conceived the study, helped RDB carry out the animal experiments, collected the
541 brain samples, performed the molecular work, the data analysis and wrote the
542 manuscript. RDB carried out the animal experiments. All authors read and edited the
543 manuscript.
544

545 **Competing interests**

546 We declare we have no competing interests.

547 **Funding**

548 This work was supported by start-up funding from Chapman University to PCL; RDB
549 was supported by a Chapman University Grand Challenges Initiative fellowship.

Acknowledgements

We thank Kaelyn Bridgette for her help with animal husbandry and for NEST lab undergraduate students for support in sample processing and behavioural analysis. We thank Greg Goldsmith for providing spontaneous parental care, particularly during the pandemic.

References

1. Numan M. 2020 *The Parental Brain: Mechanisms, Development, and Evolution*. Oxford, New York: Oxford University Press.

2. Rogers FD, Bales KL. 2019 Mothers, Fathers, and Others: Neural Substrates of Parental Care. *Trends Neurosci.* **42**, 552–562. (doi:10.1016/j.tins.2019.05.008)

3. Sear R, Mace R. 2008 Who keeps children alive? A review of the effects of kin on child survival. *Evol. Hum. Behav.* **29**, 1–18. (doi:10.1016/j.evolhumbehav.2007.10.001)

4. Dulac C, O'Connell LA, Wu Z. 2014 Neural control of maternal and paternal behaviors. *Science* **345**, 765–770. (doi:10.1126/science.1253291)

5. Pluess M, Belsky J. 2009 Differential susceptibility to rearing experience: the case of childcare. *J. Child Psychol. Psychiatry* **50**, 396–404. (doi:10.1111/j.1469-7610.2008.01992.x)

6. Bales KL, Saltzman W. 2016 Fathering in Rodents: Neurobiological Substrates and Consequences for Offspring. *Horm. Behav.* **77**, 249–259. (doi:10.1016/j.yhbeh.2015.05.021)

7. Cockburn A. 2006 Prevalence of different modes of parental care in birds. *Proc. R. Soc. B* **273**, 1375–1383. (doi:10.1098/rspb.2005.3458)

8. Ohkubo T. 2017 Neuroendocrine Control of Broodiness. *Adv. Exp. Med. Biol.* **1001**, 151–171. (doi:10.1007/978-981-10-3975-1_10)

9. Smiley KO. 2019 Prolactin and avian parental care: New insights and unanswered questions. *Horm. Behav.* **111**, 114–130. (doi:10.1016/j.yhbeh.2019.02.012)

10. Ball GF, Balthazart J. 2017 2.11 - Neuroendocrine Regulation of Reproductive Behavior in Birds. In *Hormones, Brain and Behavior (Third Edition)* (eds DW Pfaff, M Joëls), pp. 217–254. Oxford: Academic Press. (doi:10.1016/B978-0-12-803592-4.00029-8)

11. Richard-Yris M-A. 1994 Comportement parental chez les gallinacés : importance du facteur émotivité sur la vitesse d'émergence des réponses parentales : apport du modèle caille japonaise. In *Comportement et adaptation des animaux domestiques aux contraintes de*

583  *l'élevage□: bases techniques du bien-être animal* (eds M Picard, R Porter, & JP Signoret),
584  pp. 61–76. Paris: INRA Editions.

585 12. de Bruijn R, Wright-Lichter JX, Khoshaba E, Holloway F, Lopes PC. 2020 Baseline
586  corticosterone is associated with parental care in virgin Japanese quail (Coturnix japonica).
587  *Horm. Behav.* **124**, 104781. (doi:10.1016/j.yhbeh.2020.104781)

588 13. Cheng M-F, Chaiken M, Zuo M, Miller H. 1999 Nucleus Taenia of the Amygdala of Birds:
589  Anatomical and Functional Studies in Ring Doves (Streptopelia risoria) and European
590  Starlings (Sturnus vulgaris). *Brain Behav. Evol.* **53**, 243–270. (doi:10.1159/000006597)

591 14. Nelson RJ, Trainor BC. 2007 Neural mechanisms of aggression. *Nat. Rev. Neurosci.* **8**, 536–
592  546. (doi:10.1038/nrn2174)

593 15. O'Connell LA, Hofmann HA. 2011 The Vertebrate mesolimbic reward system and social
594  behavior network: A comparative synthesis. *J. Comp. Neurol.* **519**, 3599–3639.
595  (doi:10.1002/cne.22735)

596 16. Ruscio MG, Adkins-Regan E. 2004 Immediate early gene expression associated with
597  induction of brooding behavior in Japanese quail. *Horm. Behav.* **46**, 19–29.
598  (doi:10.1016/j.yhbeh.2004.02.002)

599 17. Ball GF, Balthazart J. 2010 Japanese Quail as a Model System for Studying the
600  Neuroendocrine Control of Reproductive and Social Behaviors. *ILAR J* **51**, 310–325.

601 18. Baylé JD, Ramade F, Oliver J. 1974 Stereotaxic topography of the brain of the quail
602  (Coturnix coturnix japonica). *J. Physiol. (Paris)* **68**, 219–241.

603 19. Kuenzel WJ, Masson M. 1988 *A Stereotaxic Atlas of the Brain of the Chick (Gallus*
604  *domesticus).* Poultry Science Faculty Publications and Presentations. See
605  https://scholarworks.uark.edu/poscpub/1.

606 20. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019 Graph-based genome alignment
607  and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotech.* **37**, 907–915.
608  (doi:10.1038/s41587-019-0201-4)

609 21. Anders S, Pyl PT, Huber W. 2015 HTSeq—a Python framework to work with high-
610  throughput sequencing data. *Bioinformatics* **31**, 166–169.
611  (doi:10.1093/bioinformatics/btu638)

612 22. Anders S, Huber W. 2010 Differential expression analysis for sequence count data. *Genome*
613  *Biol.* **11**, R106. (doi:10.1186/gb-2010-11-10-r106)

614 23. Love MI, Huber W, Anders S. 2014 Moderated estimation of fold change and dispersion for
615  RNA-seq data with DESeq2. *Genome Biol.* **15**, 550. (doi:10.1186/s13059-014-0550-8)

24. Robinson MD, McCarthy DJ, Smyth GK. 2010 edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140.
(doi:10.1093/bioinformatics/btp616)

25. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. 2015 limma powers
differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids*
Res. **43**, e47–e47. (doi:10.1093/nar/gkv007)

26. Law CW, Chen Y, Shi W, Smyth GK. 2014 voom: precision weights unlock linear model
analysis tools for RNA-seq read counts. *Genome Biol.* **15**, R29. (doi:10.1186/gb-2014-15-2-
r29)

27. Doncheva NT, Morris JH, Gorodkin J, Jensen LJ. 2019 Cytoscape StringApp: Network
Analysis and Visualization of Proteomics Data. *J Proteome Res.* **18**, 623–632.
(doi:10.1021/acs.jproteome.8b00702)

28. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B,
Ideker T. 2003 Cytoscape: a software environment for integrated models of biomolecular
interaction networks. *Genome Res.* **13**, 2498–2504. (doi:10.1101/gr.1239303)

29. Driessen TM, Zhao C, Whittlinger A, Williams H, Gammie SC. 2014 Endogenous CNS
Expression of Neurotensin and Neurotensin Receptors Is Altered during the Postpartum
Period in Outbred Mice. *PLOS ONE* **9**, e83098. (doi:10.1371/journal.pone.0083098)

30. Fischer EK, Roland AB, Moskowitz NA, Tapia EE, Summers K, Coloma LA, O'Connell
LA. 2019 The neural basis of tadpole transport in poison frogs. *Proc. R. Soc. B* **286**.
(doi:10.1098/rspb.2019.1084)

31. Gammie SC, D'Anna KL, Gerstein H, Stevenson SA. 2009 Neurotensin inversely modulates
maternal aggression. *Neurosci.* **158**, 1215–1223. (doi:10.1016/j.neuroscience.2008.11.045)

32. Merullo DP, Cordes MA, Susan DeVries M, Stevenson SA, Riters LV. 2015 Neurotensin
neural mRNA expression correlates with vocal communication and other highly-motivated
social behaviors in male European starlings. *Physiol. Behav.* **151**, 155–161.
(doi:10.1016/j.physbeh.2015.07.019)

33. Kasckow J, Nemeroff CB. 1991 The neurobiology of neurotensin: focus on neurotensin-
dopamine interactions. *Regul. Pept.* **36**, 153–164. (doi:10.1016/0167-0115(91)90053-J)

34. Torruella-Suárez ML, McElligott ZA. 2020 Neurotensin in reward processes.
Neuropharmacol. **167**, 108005. (doi:10.1016/j.neuropharm.2020.108005)

35. Wu Z, Autry AE, Bergan JF, Watabe-Uchida M, Dulac CG. 2014 Galanin neurons in the
medial preoptic area govern parental behaviour. *Nature* **509**, 325–330.
(doi:10.1038/nature13307)

36. Kohl J *et al.* 2018 Functional circuit architecture underlying parental behaviour. *Nature* **556**,
326–331. (doi:10.1038/s41586-018-0027-0)

37. Bukhari SA, Saul MC, James N, Bensky MK, Stein LR, Trapp R, Bell AM. 2019
Neurogenomic insights into paternal care and its relation to territorial aggression. *Nat.*
Commun. **10**, 4437. (doi:10.1038/s41467-019-12212-7)

38. Thayananuphat A, Youngren OM, Kang SW, Bakken T, Kosonsiriluk S, Chaiseha Y, El
Halawani ME. 2011 Dopamine and mesotocin neurotransmission during the transition from
incubation to brooding in the turkey. *Horm. Behav.* **60**, 327–335.
(doi:10.1016/j.yhbeh.2011.06.009)

39. Chokchaloemwong D, Prakobsaeng N, Sartsoongnoen N, Kosonsiriluk S, El Halawani M,
Chaiseha Y. 2013 Mesotocin and maternal care of chicks in native Thai hens (Gallus
domesticus). *Horm. Behav.* **64**, 53–69. (doi:10.1016/j.yhbeh.2013.04.010)

40. Sinpru P, Sartsoongnoen N, Rozenboim I, Porter TE, El Halawani ME, Chaiseha Y. 2018
The effects of replacing eggs with chicks on mesotocin, dopamine, and prolactin in the
native Thai hen. *Gen. Comp. Endocrinol.* **263**, 32–42. (doi:10.1016/j.ygcen.2018.04.013)

41. Kuperman Y, Issler O, Regev L, Musseri I, Navon I, Neufeld-Cohen A, Gil S, Chen A. 2010
Perifornical Urocortin-3 mediates the link between stress-induced anxiety and energy
homeostasis. *Proc. Natl Acad. Sci. USA* **107**, 8393–8398. (doi:10.1073/pnas.1003969107)

42. Skelton KH, Owens MJ, Nemeroff CB. 2000 The neurobiology of urocortin. *Regul. Pept.* **93**,
85–92. (doi:10.1016/S0167-0115(00)00180-4)

43. Autry AE, Wu Z, Kohl J, Bambah-Mukku D, Rubinstein ND, Marin-Rodriguez B, Carta I,
Sedwick V, Dulac C. 2019 Perifornical Area Urocortin-3 Neurons Promote Infant-directed
Neglect and Aggression. *bioRxiv* , 697334. (doi:10.1101/697334)

44. Ogino M, Okumura A, Khan MSI, Cline MA, Tachibana T. 2014 Comparison of brain
urocortin-3 and corticotrophin-releasing factor for physiological responses in chicks.
Physiol. Behav. **125**, 57–61. (doi:10.1016/j.physbeh.2013.11.006)

45. Wang J, Gilbert ER, Cline MA. 2019 Hypothalamic mechanisms associated with
neuropeptide K-induced anorexia in Japanese quail (Coturnix japonica). *Comp. Biochem.*
Physiol. A Mol. Integr. Physiol. **237**, 110539. (doi:10.1016/j.cbpa.2019.110539)

46. O'Rourke CF, Renn SC. 2015 Integrating adaptive trade-offs between parental care and
feeding regulation. *Curr. Opin. Behav. Sci.* **6**, 160–167. (doi:10.1016/j.cobeha.2015.11.010)

47. Fischer EK, O'Connell LA. 2017 Modification of feeding circuits in the evolution of social
behavior. *J. Exp. Biol.* **220**, 92–102. (doi:10.1242/jeb.143859)

48. Albrektsen T, Richter HE, Clausen JT, Fleckner J. 2001 Identification of a novel integral
plasma membrane protein induced during adipocyte differentiation. *Biochem. J.* **359**, 393–
402. (doi:10.1042/0264-6021:3590393)

49. Mosser S, Alattia J-R, Dimitrov M, Matz A, Pascual J, Schneider BL, Fraering PC. 2015 The
adipocyte differentiation protein APMAP is an endogenous suppressor of Aβ production in
the brain. *Hum. Mol. Genet.* **24**, 371–382. (doi:10.1093/hmg/ddu449)

50. Bahrami S, Drabløs F. 2016 Gene regulation in the immediate-early response process. *Adv.*
Biol. Regul. **62**, 37–49. (doi:10.1016/j.jbior.2016.05.001)

51. Ray S, Tzeng R-Y, DiCarlo LM, Bundy JL, Vied C, Tyson G, Nowakowski R, Arbeitman
MN. 2015 An Examination of Dynamic Gene Expression Changes in the Mouse Brain
During Pregnancy and the Postpartum Period. *G3 (Bethesda)* **6**, 221–233.
(doi:10.1534/g3.115.020982)

52. Sapolsky RM, Romero LM, Munck AU. 2000 How do glucocorticoids influence stress
responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocr.*
Rev. **21**, 55–89. (doi:10.1210/edrv.21.1.0389)

53. Torner L. 2016 Actions of Prolactin in the Brain: From Physiological Adaptations to Stress
and Neurogenesis to Psychopathology. *Front. Endocrinol. (Lausanne)* **7**.
(doi:10.3389/fendo.2016.00025)

54. Lambert PD, Ely TD, Gross RE, Kilts CD. 1996 Neurotensin induces Fos and Zif268
expression in limbic nuclei of the rat brain. *Neuroscience* **75**, 1141–1151.
(doi:10.1016/0306-4522(96)00210-2)

55. Gammie SC, Lee G, Scotti M-AL, Stevenson SA, Gessay GM. 2012 Neurotensin induced
Egr-1 activity is altered in the postpartum period in mice. *Brain Res.* **1433**, 47–55.
(doi:10.1016/j.brainres.2011.11.017)

56. Scotti M-AL, Stevenson SA, Gammie SC. 2011 Changes in CNS response to neurotensin
accompany the postpartum period in mice. *Horm. Behav.* **60**, 177–184.
(doi:10.1016/j.yhbeh.2011.04.007)

57. Bullitt E. 1990 Expression of C-fos-like protein as a marker for neuronal activity following
noxious stimulation in the rat. *J. Comp. Neurol.* **296**, 517–530.
(doi:https://doi.org/10.1002/cne.902960402)

58. Bentz AB, Rusch DB, Buechlein A, Rosvall KA. 2019 The neurogenomic transition from
territory establishment to parenting in a territorial female songbird. *BMC Genomics* **20**, 819.
(doi:10.1186/s12864-019-6202-3)

59. Ye P, Li M, Liao W, Ge K, Jin S, Zhang C, Chen X, Geng Z. 2019 Hypothalamic
transcriptome analysis reveals the neuroendocrine mechanisms in controlling broodiness of
Muscovy duck (Cairina moschata). *PLOS ONE* **14**, e0207050.
(doi:10.1371/journal.pone.0207050)