1 **Brain transcriptomic changes in Japanese quail reveal a role for neurotensin in**
2 **avian parental care**
3

Patricia C. Lopes[*] and Robert de Bruijn

Schmid College of Science and Technology, Chapman University, Orange, California,
USA

* Corresponding author
Patricia C. Lopes
lopes@chapman.edu

Abstract

For many species, parental care critically affects offspring survival. But what drives

animals to display parental behaviours towards young? In mammals, pregnancy-

induced physiological transformations seem key in preparing the neural circuits that

lead towards attraction (and reduced-aggression) to young. Beyond mammalian

maternal behaviour, knowledge of the neural mechanisms that underlie parental care is

severely lacking. We took advantage of a domesticated bird species, the Japanese

quail, for which parental behaviour towards chicks can be induced through a

sensitization procedure, a process that is not effective in all animals. We used the

variation in parental responses to study neural transcriptomic changes associated with

the sensitization procedure itself, with the outcome of the procedure (i.e., presence of

parental behaviours) and with spontaneous parental care (i.e., in the absence of

sensitization). Out of the brain regions studied, we found that most differences in gene

expression were located in the hypothalamus. Our results highlight several molecular

pathways that may contribute to the modulation of avian parental care. We identified

one gene, neurotensin, that was previously only demonstrated to be causally associated

with parental care in mammals. Our work opens new avenues of research into

understanding the neural basis of parental care in non-mammalian species.

Keywords: avian parental care, bed nucleus of the stria terminalis, hypothalamus,

Neurotensin, nucleus taeniae, prolactin, RNA-seq, Urocortin

1. Introduction

While parental care is rare in many vertebrate taxa, it is essential for survival in the majority of avian species, with biparental care being the most frequent modality observed [1]. As a group, birds show an incredible diversity in forms of parental care, from brood parasitism, where care is lost, to cooperative breeding, where non-reproductive individuals provide care to offspring that are not their own [1]. The neural underpinnings of young-directed parental care in birds are, however, severely understudied. In terms of neuroendocrine mechanisms associated with incubation and caring for young, the majority of attention has been paid to a single pituitary hormone: prolactin [2,3]. In contrast to birds, uniparental maternal care is the most frequently observed mode of care in mammals and, therefore, the neuroendocrine basis of this behaviour is more well studied and understood than any other form of parental care across vertebrate taxa.

Using a sensitization procedure developed by [4], a recent experiment by our group confirmed that it is possible to induce parental behaviours in Japanese quail through a single overnight exposure to chicks [5]. After the sensitization procedure, both sexes display chick brooding behaviour and reduced aggression towards chicks (pecking behaviour). While the change in behaviour after sensitization is drastic, it is not observed in all animals, i.e., some animals fail to respond to the sensitization. Conversely, a few animals not exposed to the sensitization spontaneously show some parental care when paired with chicks.

In the current study, we take advantage of this variation in responses to the sensitization treatment to understand, from a neural molecular perspective, a) what modifications occur due to the sensitization treatment, b) how animals that are unable to show care after this treatment differ from parental ones, and c) how some animals are capable of spontaneously showing care during their first encounter with chicks. We focused our analysis on the hypothalamus, the bed nucleus of the stria terminalis and the nucleus taeniae. We targeted these brain regions because they contain nuclei

important for parental care, affiliative interactions, and agonistic interactions in birds and

other vertebrates [6–11]. Our experiments highlight the importance of using intra-

species variation in parental responses to elucidate neural mechanisms of parental

behaviour and to discover new molecular pathways involved in avian parental care,

several of which we believe are conserved across other vertebrate taxa.

2. Materials and methods

(a) Parental care induction procedure and behaviour quantification

Japanese quail (*Cortunix japonica*) were raised in the lab from eggs obtained from AA

Lab Eggs, Inc. (Westminster, CA) following the procedures described in [5]. Adult birds

were kept in an 8L:16D light cycle and maintained in groups of 4-5 same sex animals in

cages (100 × 40 × 50 cm). Two days prior to carrying out the experiments, a replica of

the wooden box (18 × 18 × 18 cm) used in the parental care sensitization procedure

was added to those cages. The day prior to the procedure, animals were separated

individually into cages identical to the group cages, which also contained a wooden box.

The wooden box was always open during these days, allowing the birds to inspect it

and go in and out. Food and water were provided *ad libitum* during the entire

experiment.

The day of the parental care sensitization procedure, each adult bird was locked inside

the wooden box in their individual cage starting at one hour before lights were off. Under

the parental induction treatment, two chicks (1-3 days old) were added to this box just

before lights off. Under the control treatment, no chicks were added to the box. Animals

were left undisturbed overnight. The morning after, the wooden box was opened, the

overnight chicks were removed and two new chicks (1-3 days old) were added to the

cage. This swap of chicks ensured that birds in both treatments (controls and

sensitized) were being tested for their behaviours towards novel chicks (in case

familiarity influenced the outcome). The novel chicks were allowed to stay in the cage

for 20 min, during which videos were continuously recorded (Axis M1065L network

camera by Axis Communications). At the end of the 20 min, adults were euthanized by

isoflurane inhalation and decapitation. The brain was removed from the skull, flash

frozen and then stored at -80 °C until further processing. Chicks were returned to their

cages. In total, 50 adults (ages 90-110 days old) were subjected to the experimental

procedures, of which 11 females and 13 males were randomly assigned to the

sensitization treatment and 19 females and 7 males to the control treatment. Two

control females were removed from the study due to methodological failures during the

experiments, bringing the final number of control females to 17. In total, 77 chicks were

used. Behaviours towards chicks within the 20 min period of observation were scored

by observers blind to the treatment, as described in De Bruijn et al. (2020). We used

these behavioural data to place birds in different categories for the subsequent

transcriptomics analysis. In particular, we used the duration of parental behaviour (chick

brooding, measured as time spent sitting on top of chicks) to define whether sensitized

birds were responders (>= 180 sec of parental behaviour) or non-responders (< 60 sec

of parental behaviour) to this treatment. Similarly, we classified controls birds as

showing spontaneous parental behaviour (>= 180 sec of parental behaviour) or no

parental behaviour (< 60 sec of parental behaviour). Animals with chick brooding

duration between 60 and 180 sec were removed from the data for the responsiveness

comparisons, but not for the treatment comparisons, during the transcriptomic analysis.

(b) Brain dissection

Brains were coronally sectioned on a Leica CM1860UV cryostat, at -18 °C. We used

separate surgical micropunches (EMS Rapid Core Instruments) to punch out the three

brain regions of interest from 100 µm slices, spaced apart by three 30 µm slices

collected onto microscope slides (Fisherbrand, item 12–550-15) for future use. The

brain regions collected were the entire hypothalamus, the nucleus taeniae (Tn) and the

bed nucleus of the stria terminalis (BnST). These regions were identified based on use

of both the quail and the chicken brain atlases [12,13]. Punches from each of these

brain regions were placed in individual tubes containing 2 mm size beads (ZR

BashingBeads Lysis Tubes, Zymo Research, item S6003-50) and 1 mL of QIAzol lysis

reagent (Qiagen, item #79306). Tissue homogenization was done by agitating the tubes

for 20□s at a 7□m□s$^{−□1}$ speed (Beadbug 6 homogenizer, Benchmark Scientific),

followed by a 5□min rest period. The homogenate was then transferred into a new tube

and preserved at −□80□°C until RNA isolation.

(c) RNA isolation, library preparation and sequencing

Total RNA was extracted from the aqueous layer formed after chloroform precipitation,

using the RNA Clean & Concentrator Kit-5 (Zymo Research, item # R1013) following

manufacturer's instructions and including the DNase I in-column treatment step. These

samples were then shipped on dry-ice to Novogene Corporation Inc. (Chula Vista, CA,

USA), where RNA quantity, purity and integrity were assessed on an Agilent 2100

Bioanalyzer (Agilent Technologies, Santa Clara, CA), cDNA libraries were generated

using NEBNext® UltraTM RNA Library Prep Kit for Illumina® (NEB, USA) and cDNA

fragments (150 ~ 200 bp in length) were then purified using the AMPure XP System

(Beckman Coulter, Beverly, USA). Paired-end sequencing of libraries (PE150; Illumina

Novaseq 6000) was performed according to standard protocols. An average of 52

million paired-end raw reads were obtained for each sample.

(d) Mapping and differential gene expression analysis

An average of 85.3% of clean (post adapter removal and quality filtering) reads were

mapped to the Japanese quail reference genome (Coturnix_japonica_2.0, INSDC

Assembly Mar 2016, downloaded from Ensembl), representing an average of 43.7

million mapped reads per sample (Table S1 contains information on mapping statistics

per sample). Mapping was done using HISAT2 [14], and HTSeq was used to count the

number of mapped reads to each gene [15]. Differential gene expression analysis was

performed using the DESeq2 R package [16,17].

For each sex and for each brain region, the following pairwise comparisons were

performed: birds subjected to the sensitization treatment relative to control treated birds;

birds responsive to the sensitization treatment (responders) versus non-responsive

birds (non-responders); and birds subjected to the control treatment that showed

spontaneous parental care (spontaneous) relative to birds in the same treatment that

showed no care (no-care). As only one control male showed spontaneous parental

behaviour, comparisons of this male relative to the non-parental males in the control

group were not performed. To control for the false discovery rate due to multiple

testing, *p*-values were adjusted using the Benjamini-Hochberg procedure. Genes were

considered as statistically differentially expressed when adjusted *p*-values

were $<$ 0.05. Boxplots for a subset of the differentially expressed genes were

prepared using the ggplot2 R package [18], using DESeq2 normalized counts. A

heatmap was produced for the differentially expressed genes in the hypothalamus of

control females using the pheatmap package (version 1.0.12; [19]) in R, using variance

stabilized DESeq2 counts.

(e) Gene ontology (GO) enrichment analysis and KEGG pathway enrichment

analysis of differentially expressed genes

GO enrichment analysis was done only for the differentially expressed genes (DEGs)

found in the hypothalamus of control females with spontaneous versus control females

with no parental care. The analysis was done separately for up- and down-regulated

DEGs, using ShinyGO (v0.61; [20]) with the Japanese quail as the species, and FDR

cut-off of 0.05. ShinyGO can also group genes by functional categories defined by high-

level GO terms, which are often more easily interpretable and therefore, also reported.

KEGG pathway enrichment analysis was also performed using ShinyGO with an FDR

cut-off of 0.05, but this time with mouse as the reference species, as this analysis tool is

not well developed for Japanese quail.

3. Results

(a) Effect of sensitization treatment

Behavioural response. In response to spending an overnight enclosed in a box with
two chicks (sensitization treatment), 72.7 % of females responded with parental
behaviours (responders) and all others were non-responders. Concerning the males,
69.2 % males were responders, 23.1% non-responders and one male undefined (> 60
sec of parental behaviour but < 180 sec). Out of control treated animals (no chicks
overnight), 25 % of the females showed spontaneous parental behaviour, 62.5% of
females showed no parental behaviour and two females were undefined. The duration
of parental behaviour in control animals is, however, shorter (average ± s.e.m = 396 ±
119 sec) than that of parental females in the sensitization treatment (744 ± 54 sec). Of
the control males, only 12.5 % showed spontaneous parental behaviour and all others
were non-parental.
199
Transcriptomic response. To our surprise, very few genes were differentially
expressed in the brains of animals that underwent the sensitization treatment relative to
controls during interaction with chicks (Table S2). Neurotensin (NTS) expression was
significantly higher in the hypothalamus of sensitized females and trended in that
direction in males (figure 1). In both sexes, Urocortin-3 (UCN3) was upregulated in the
hypothalamus of sensitized relative to control animals (figure 1). In females, one
additional gene was upregulated in sensitized animals: a zinc finger protein (ZFN385C).
In males, one different additional gene was upregulated in sensitized animals, calbindin
1 (CALB1), and a different gene downregulated, BTG anti-proliferation factor 2 (BTG2)
(figure 1). Patterns of BTG2 expression were similar in females (figure 1). In the bed
nucleus of the stria terminalis (BnST), two DEGs were upregulated in sensitized
females (pentatricopeptide repeat domain 1 or PTCD1, and an unidentified gene) and
two were downregulated (FOS, and salt inducible kinase 1, SIK1). In males, only one
DEG was found in this region (Kruppel like factor 2, KLF2) and it was downregulated in
sensitized relative to control animals. No genes were differentially expressed due to
treatment in either sex in the Nucleus Taeniae (Tn).

Figure 1. Expression levels of selected genes differentially expressed in the
hypothalamus of animals exposed or not exposed to the sensitization treatment. Males
and females are represented in all graphs for purposes of comparison.

(b) Differences between responders and non-responders

We compared the transcriptomic responses of the animals that responded to the
sensitization treatment by showing parental behaviours (> 180 sec of chick brooding
response; responders) to those of animals with no clear signs of parental behaviours (<
60 sec of chick brooding response; non-responders). In the hypothalamus of females,
32 genes were differentially expressed between responders and non-responders (Table
S3), several of which were either immediate early genes (EGR1, EGR2, FOS, ARC). In
males, there were only 3 hypothalamic DEGs, including COL28A1 and COL20A1, both
related to collagen formation, and anti-Mullerian hormone (AMH). The three BnST

DEGs upregulated in female responders were all unidentified. In males, there were
seven DEGs upregulated in responders in this brain region, and one downregulated. Of
the upregulated genes, one was also AMH, as found in the hypothalamus. In the Tn,
four DEGs were upregulated in female responders, two of which being unidentified and
the other two being target of EGR1 (TOE1) and t-complex-associated-testis-expressed
3 (TCT3). In males, two DEGs were upregulated in this region responders, transthyretin
(TTR) and solute carrier family 13 member 4 (SLC13A4).
239
240   **(c) Differences between females showing spontaneous parental behaviour**
241   **relative to non-parental females**
242
As four females in the control treatment showed parental behaviour when presented
with chicks (spontaneous), we compared the transcriptomic responses of these relative
to control females showing no parental behaviour (non-spontaneous). One gene, IRF5
(interferon regulatory factor 5) was upregulated in spontaneous females in all three
brain regions analysed. While in the BnST, no genes were downregulated, in the Tn
three genes were (sulfatase 1, SULF1; farnesyl diphosphate synthase, FDPS; and
tachykinin precursor 1, TAC1). Meanwhile, three thousand genes were differentially
expressed in the hypothalamus (Table S4; figure 2). Given the large number of DEGs in
this region, we used GO enrichment analysis and KEGG pathway enrichment analysis
for data reduction. The GO enrichment analysis revealed enrichment in several GO
categories related with metabolism, immune system function/immune response,
reproduction, circadian biology and behaviour (including reproductive behaviour,
locomotor behaviour, feeding behaviour, and behavioural defence response) (Table S5
and Table S6). In addition to also identifying several pathways involved in metabolism
(e.g., insulin resistance and fatty acid metabolism pathway) and immune system
responses (e.g., platelet activation and TNF signalling pathway) and one pathway
related to circadian biology (circadian rhythm), enrichment analysis of KEGG pathways
identified important hormone signalling (e.g., thyroid hormone signalling, GnRH
signalling pathway, oxytocin signalling pathway and prolactin signalling pathway) and
synaptic function pathways (e.g., cholinergic synapse, GABAergic synapse,

dopaminergic synapse, glutamatergic synapse) that are involved in reproduction and

behaviour (Figure 3; Table S7).

Figure 2. Heatmap of hypothalamic genes differentially expressed between control

females that showed spontaneous parental behaviour when placed with chicks, relative

to females that did not. Blue colours indicate lower expression and red colours higher

expression levels in parental relative to non-parental controls.

Figure 3. Number of hypothalamic downregulated DEGs (blue) or upregulated DEGs (green) in significantly enriched KEGG pathways in control females showing spontaneous parental behaviour compared to non-parental females. Pathways that were redundant, related to infectious diseases or cancer, or not brain-specific were excluded from this figure for simplicity (see Table S7 for full list).

4. Discussion

The neural mechanisms mediating parental care remain poorly understood. We took advantage of a domesticated species, the Japanese quail, for which parental behaviour towards chicks does not tend to occur spontaneously in captivity. Parental behaviour in these animals can be induced through a sensitization procedure, but the process is not 100 % effective. Using the variation in parental responses, we asked what neural molecular mechanisms are: a) altered upon sensitization, b) different between birds that show chick care behaviours in response to the sensitization relative to non-responders, and c) different between untreated animals that spontaneously show parental care relative to those that do not.

Out of the three brain regions studied, we found that most gene expression differences were located in the hypothalamus. We identified a gene, neurotensin, that was previously only demonstrated to be causally associated with parental care in mammals. Transcriptomic studies focused on animals that fail to show parental care under care-promoting circumstances, or studies on those animals that show care unprompted provide great opportunities to understand the neural basis of variation in parental care.

(a) How does sensitization modify neural responses?

By comparing the neural transcriptomic responses of adult quail that had undergone a sensitization procedure to those that did not undergo this process, we were able to identify a small number of differentially expressed genes that may be involved in avian

parental care. One of those genes, neurotensin, was expressed at higher levels in the

hypothalamus of both females and males (albeit only significantly so in females) after

sensitization. This finding is significant given that neurotensin levels in the

hypothalamus of postpartum female mice were also observed to be elevated relative to

that of virgin mice [21]. Neurotensin gene expression was also recently shown to be

elevated in the preoptic area of males of one species of poison frog (*Dendrobates*

tinctorius) when performing parental care relative to males not performing care [22].

Furthermore, intracerebroventricular injections of neurotensin significantly decreased

maternal aggression in female mice [23], indicating the neurotensin participates in at

least one component of maternal responses in rodents. Previously, the only reports of

behavioural effects of neurotensin on birds suggested a possible relationship between

neurotensin expression in the medial preoptic nucleus (POM) and non-vocal courtship

behaviour and agonistic behaviour in male European starlings (*Sturnus vulgaris*) [24].

Interestingly, that study also reported that male starlings that had acquired a nest box

also had higher neurotensin expression in the POM relative to males that did not

acquire boxes and the measures of non-vocal courtship behaviour were mostly related

to nest building (e.g., such as number of times entering or landing on the nest and

gathering nesting material). European starlings show biparental care and the increase in

neurotensin in a hypothalamic nucleus in that study could have been related to nest-

building activity, which, for many species, is a component of parental care.

Out of the hormones associated with avian parental care to date, prolactin is by far the

one that has receive the most attention [3]. In precocial avian species, as is the case

with Japanese quail, prolactin tends to be elevated in females during the incubation

period and to decline when chicks hatch [2]. Prolactin release is known to be under the

control of dopamine, which can have both inhibitory and stimulatory effects depending

on the dopamine receptor subtype it acts on [3]. Neurotensin could impact parental

behaviour by playing a role in controlling prolactin secretion, through its action in

enhancing dopamine release [25]. Neurotensin could also affect care through its

involvement in reward processing [26].

Another interesting gene that was upregulated following the sensitization procedure in
both sexes was urocortin-3. In rodents, central administration or overexpression of
urocortin-3 increases stress-induced anxiety and suppresses ingestive behaviour
[27,28]. We are only aware of two reports relating urocortin-3 to young-directed parental
care. One is the poison frog paper mentioned above [22] where urocortin-3 was
upregulated in the preotic area of frogs performing parental care relative to non-parental
one. The other one revealed that urocortin-3 expressing neurons in the perifornical area
of the hypothalamus are activated specifically during infant-directed aggression in both
male and female mice and that optogenetic activation of these neurons triggers infant-
directed aggression and neglect [29]. Given that we observed a reduction in aggression
towards chicks after the sensitization procedure [5], urocortin-3 may have an opposing
effect on infant-directed aggressive behaviour in our system or serve different functions
that support parental care, such as altered feeding patterns [30,31].
347
Calbindin 1 was only upregulated in the hypothalamus of sensitized males. Calbindin 1
has a sexually dimorphic distribution in certain brain regions, including the hypothalamic
nuclei in mice [32,33] and in regions of the song control system and related areas in
certain bird species [34–36]. In mice, calbindin 1 appears to also have sexually
dimorphic effects on behaviour, including effects on fear conditioning and on social
preferences in males [37]. Calbindin 1 has been found to limit dopamine release in
certain brain regions, which, as mentioned previously, can affect prolactin secretion
[38]. Also exclusive to sensitized males was the downregulation of BTG2. One study in
mice and zebrafish found increased neural expression of this gene in males exposed to
aggressive social encounters [39]. As we do observe aggression towards chicks in
control males [5], differences in BTG2 may be linked to aggressive interactions in our
study.
360
FOS, an immediate early gene downregulated in the BnST in sensitized females relative
to controls, is widely used as a marker of neuronal activation of specific brain regions
[40] and the difference between sensitized and control females may imply that the BnST
is activated in response to chicks only in control females. Interestingly, sensitized

parental female quail separated from their chicks show increased fos immunoreactivity
in this region [11], which may indicate an involvement of BnST in maternal separation.
As the sensitization elicited no significant gene expression changes in the Tn in neither
sex, this region is likely less critically involved during the expression of parental care in
birds.
370
371 **(b) What makes a non-responder?**
372
One important aspect of our study is that rather than excluding animals that did not
respond to the sensitization treatment, we used them to understand what neural
differences there are in animals that fail to show parental behaviours under conditions
that induce these behaviours. Female responders showed lower expression of several
immediate early genes (EGR1, EGR2, FOS, ARC) in the hypothalamus, which could be
indicative of lower hypothalamic activity after effective sensitization. Several of the
downregulated genes in responders are also associated with the modulation of synaptic
signalling and synapse reorganization (CNKSR2, NPAS4, ARC, GRIN2B, EGR2) and
with nervous system development (CPNE5, MEF2D, NR4A3, EGR2, FOS, NPAS4,
ARC, CDHR1, GRIN2B). It has been suggested that maternal experience can alter both
the functioning and the structure of the medial preoptic area (a hypothalamic nucleus) in
a way that increases neuronal responsiveness to young-related cues (reviewed in [41]).
It is therefore possible that, in females that failed to show parental behaviours, the
sensitization procedure was ineffective in triggering the necessary re-structuring of
hypothalamic nuclei. Other brain regions studied had fewer differences between female
responders and non-responders.
389
Contrary to results in females, in males, most differences between responders and non-
responders were seen in the BnST. Of particular interest are two genes that had higher
expression in responders: Anti-Müllerian Hormone (AMH) and Butyrylcholinesterase
(BCHE). Increased expression of AMH was also found in the hypothalamus of male
responders. While the most familiar role of AMH is in the regression of Müllerian ducts
in male embryos during development, AMH is also produced in the brain [42,43].

Emerging actions of AMH are being found in the control of hypothalamic function, in particular, in increasing GnRH neuron excitability and GnRH secretion [42,43]. BChE has an interesting role in inactivating ghrelin (a hormone that affects hunger, feeding and stress), and mice with overexpression of BCHE show decreased fighting [44,45]. A reduction in aggression towards young is essential for proper parental care and BChE might facilitate this in responders. The two DEGs in the Tn (Transthyretin, TTR, and the sulfate transporter SLC13A4) showed some of the greatest fold changes observed in responders relative to non-responders. TTR expression has been shown to change in mammals in ways that depend on the manipulation and brain region. For instance, TTR expression decreases in the cortex of mice exposed to chronic immobilization stress [46] but increases in the amygdala during the formation of fear memory [47]. One interpretation is, therefore, that the presence of chicks may be perceived as a stressor by non-responders. Regarding SLC13A4, a recent study in mice showed that Slc13a4 haploinsufficiency (Slc13a4$^{+/-}$) results in atypical social behaviours, including longer time to retrieve pups and reduced nest-building activity in females, and reduced social exploration and impaired social memory in both sexes [48]. While parental behaviours were only tested in females in that study (given that male mice do not display those), a Slc13a4 increase may be required for proper parental behaviours in other species.

(c) What neural changes are associated with spontaneous care?

A number of females in our experiment that had no previous experiences with chicks showed parental behaviour in the absence of sensitization. It was in the hypothalamus of these females that we found the largest number of differentially expressed genes out of all comparisons performed in our study. While these control females showed some parental behaviour, the time spent engaging in those behaviours was less than those observed in sensitized animals. The changes in gene expression in parental controls may therefore represent processes associated with the *onset* of the modifications that will be needed for a more complete parental response. For example, enrichment analysis highlighted several pathways associated with synaptic function, including

426 cholinergic, GABAergic, dopaminergic, and glutamatergic synapse, which may again
427 indicate the necessity for a hypothalamic re-structuring needed for parental care.
428
Many of the pathways detected through enrichment analysis have been linked to
parental care in mammals and other species, including thyroid hormone, GnRH,
oxytocin and prolactin signalling pathways [7,10,41,49]. The onset of parental care is
associated with reduced reproductive behaviour, reduced young-directed aggression,
and with changes in metabolism, locomotion and feeding patterns [10,49]. Pathways
associated with this array of behavioural and metabolic changes were highlighted
through enrichment analysis. Interestingly, recent work proposes that molecules related
to feeding behaviour may also be involved in complex social behaviours, such as
parental care [22,50].
438
Changes in immune function were also detected through enrichment analysis. In fact,
one DEG, IRF5, was upregulated in parental females controls across all three brain
regions analysed. IRF5 is produced in the brain by microglia (resident immune cells in
the brain) and it is known for its pro-inflammatory functions [51,52]. Extensive
neuroimmune changes occur in rodents postpartum (reviewed in [53]) and it is likely that
some of these extend to other vertebrate species (e.g., [54]).
445
446 **5. Conclusion**
447
Recent studies have used transcriptomic approaches to compare gene expression in
specific brain areas as female birds transition from territorial defence to egg incubation
[55], or from egg-laying to incubation related behaviours [56]. Still, the range of neural
mechanisms involved in post-hatching care in birds is virtually unknown. Our findings
revealed new potential modulators of avian young-directed parental care, which deserve
to be examined through further mechanistic studies and explored in other bird species
and other taxonomic groups. Particularly exciting are neurotensin and urocortin-3, which
potentially show conserved roles in parental care in mammals, amphibians, and birds,
but are extremely under-explored in the context of parental behaviour. Natural variation

in avian parental behaviour in response to young extends from abandoning the nest to

alloparenting or even adoption of non-offspring deposited by brood parasites. The

pathways and genes found to be associated with spontaneous parental care, with lack

of parental responsiveness, and with parental sensitization in our study serve as a

foundation for future studies aimed at investigating the neural underpinnings of variation

in parental care within and across species.

Ethics

Animal use and experimental design were approved by the Chapman University

Institutional Animal Care and Use Committee (protocol # 2019-01).

Data accessibility

RNA-seq mapping statistics, lists of DEGs and lists of enrichment terms have been

uploaded as part of the supplementary material. Sequencing datasets generated and

analysed during the current study are deposited in the NCBI Gene Expression Omnibus

(GEO) repository [dataset will be made publicly available once manuscript is accepted

for publication].

Authors' contributions

PCL conceived the study, helped RDB carry out the animal experiments, collected the

brain samples, performed the molecular work, the data analysis and wrote the

manuscript. RDB carried out the animal experiments. All authors read and edited the

manuscript.

Competing interests

We declare we have no competing interests.

Funding

This work was supported by start-up funding from Chapman University to PCL; RDB was supported by a Chapman University Grand Challenges Initiative fellowship.

Acknowledgements

We thank Kaelyn Bridgette for her help with animal husbandry and for NEST lab undergraduate students for support in sample processing and behavioural analysis. We thank Greg Goldsmith for providing spontaneous parental care, particularly during the pandemic.

References

1. Cockburn A. 2006 Prevalence of different modes of parental care in birds. *Proc. R. Soc. B* **273**, 1375–1383. (doi:10.1098/rspb.2005.3458)

2. Ohkubo T. 2017 Neuroendocrine Control of Broodiness. *Adv. Exp. Med. Biol.* **1001**, 151–171. (doi:10.1007/978-981-10-3975-1_10)

3. Smiley KO. 2019 Prolactin and avian parental care: New insights and unanswered questions. *Horm. Behav.* **111**, 114–130. (doi:10.1016/j.yhbeh.2019.02.012)

4. Richard-Yris M-A. 1994 Comportement parental chez les gallinacés: importance du facteur émotivité sur la vitesse d'émergence des réponses parentales: apport du modèle caille japonaise. In *Comportement et adaptation des animaux domestiques aux contraintes de l'élevage: bases techniques du bien-être animal* (eds M Picard, R Porter, & JP Signoret), pp. 61–76. Paris: INRA Editions.

5. de Bruijn R, Wright-Lichter JX, Khoshaba E, Holloway F, Lopes PC. 2020 Baseline corticosterone is associated with parental care in virgin Japanese quail (Coturnix japonica). *Horm. Behav.* **124**, 104781. (doi:10.1016/j.yhbeh.2020.104781)

6. Cheng M-F, Chaiken M, Zuo M, Miller H. 1999 Nucleus Taenia of the Amygdala of Birds: Anatomical and Functional Studies in Ring Doves (Streptopelia risoria) and European Starlings (Sturnus vulgaris). *Brain Behav. Evol.* **53**, 243–270. (doi:10.1159/000006597)

7. Dulac C, O'Connell LA, Wu Z. 2014 Neural control of maternal and paternal behaviors. *Science* **345**, 765–770. (doi:10.1126/science.1253291)

8. Nelson RJ, Trainor BC. 2007 Neural mechanisms of aggression. *Nat. Rev. Neurosci.* **8**, 536–546. (doi:10.1038/nrn2174)

9. O'Connell LA, Hofmann HA. 2011 The Vertebrate mesolimbic reward system and
social behavior network: A comparative synthesis. *J. Comp. Neurol.* **519**, 3599–
3639. (doi:10.1002/cne.22735)

10. Rogers FD, Bales KL. 2019 Mothers, Fathers, and Others: Neural Substrates of
Parental Care. *Trends Neurosci.* **42**, 552–562. (doi:10.1016/j.tins.2019.05.008)

11. Ruscio MG, Adkins-Regan E. 2004 Immediate early gene expression associated
with induction of brooding behavior in Japanese quail. *Horm. Behav.* **46**, 19–29.
(doi:10.1016/j.yhbeh.2004.02.002)

12. Baylé JD, Ramade F, Oliver J. 1974 Stereotaxic topography of the brain of the quail
(Coturnix coturnix japonica). *J. Physiol. (Paris)* **68**, 219–241.

13. Kuenzel WJ, Masson M. 1988 *A Stereotaxic Atlas of the Brain of the Chick (Gallus
domesticus).* Poultry Science Faculty Publications and Presentations. See
https://scholarworks.uark.edu/poscpub/1.

14. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019 Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotech.* **37**, 907–
915. (doi:10.1038/s41587-019-0201-4)

15. Anders S, Pyl PT, Huber W. 2015 HTSeq—a Python framework to work with high-
throughput sequencing data. *Bioinformatics* **31**, 166–169.
(doi:10.1093/bioinformatics/btu638)

16. Anders S, Huber W. 2010 Differential expression analysis for sequence count data.
Genome Biology **11**, R106. (doi:10.1186/gb-2010-11-10-r106)

17. Love MI, Huber W, Anders S. 2014 Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550.
(doi:10.1186/s13059-014-0550-8)

18. Wickham H. 2016 *ggplot2: Elegant Graphics for Data Analysis*. 2nd edn. Springer
International Publishing. See https://www.springer.com/gp/book/9783319242750.

19. Kolde R. 2019 pheatmap: Pretty Heatmaps. R package version 1.0.12.

20. Ge SX, Jung D, Yao R. 2020 ShinyGO: a graphical gene-set enrichment tool for
animals and plants. *Bioinformatics* **36**, 2628–2629.
(doi:10.1093/bioinformatics/btz931)

21. Driessen TM, Zhao C, Whittlinger A, Williams H, Gammie SC. 2014 Endogenous
CNS Expression of Neurotensin and Neurotensin Receptors Is Altered during the
Postpartum Period in Outbred Mice. *PLOS ONE* **9**, e83098.
(doi:10.1371/journal.pone.0083098)

22. Fischer EK, Roland AB, Moskowitz NA, Tapia EE, Summers K, Coloma LA,
O'Connell LA. 2019 The neural basis of tadpole transport in poison frogs. *Proc. R.*
Soc. B **286**. (doi:10.1098/rspb.2019.1084)

23. Gammie SC, D'Anna KL, Gerstein H, Stevenson SA. 2009 Neurotensin inversely
modulates maternal aggression. *Neuroscience* **158**, 1215–1223.
(doi:10.1016/j.neuroscience.2008.11.045)

24. Merullo DP, Cordes MA, Susan DeVries M, Stevenson SA, Riters LV. 2015
Neurotensin neural mRNA expression correlates with vocal communication and
other highly-motivated social behaviors in male European starlings. *Physiol. Behav.*
151, 155–161. (doi:10.1016/j.physbeh.2015.07.019)

25. Kasckow J, Nemeroff CB. 1991 The neurobiology of neurotensin: focus on
neurotensin-dopamine interactions. *Regul. Pept.* **36**, 153–164. (doi:10.1016/0167-
0115(91)90053-J)

26. Torruella-Suárez ML, McElligott ZA. 2020 Neurotensin in reward processes.
Neuropharmacol. **167**, 108005. (doi:10.1016/j.neuropharm.2020.108005)

27. Kuperman Y, Issler O, Regev L, Musseri I, Navon I, Neufeld-Cohen A, Gil S, Chen
A. 2010 Perifornical Urocortin-3 mediates the link between stress-induced anxiety
and energy homeostasis. *Proc. Natl Acad. Sci. USA* **107**, 8393–8398.
(doi:10.1073/pnas.1003969107)

28. Skelton KH, Owens MJ, Nemeroff CB. 2000 The neurobiology of urocortin. *Regul.*
Pept. **93**, 85–92. (doi:10.1016/S0167-0115(00)00180-4)

29. Autry AE, Wu Z, Kohl J, Bambah-Mukku D, Rubinstein ND, Marin-Rodriguez B,
Carta I, Sedwick V, Dulac C. 2019 Perifornical Area Urocortin-3 Neurons Promote
Infant-directed Neglect and Aggression. *bioRxiv* , 697334. (doi:10.1101/697334)

30. Ogino M, Okumura A, Khan MSI, Cline MA, Tachibana T. 2014 Comparison of brain
urocortin-3 and corticotrophin-releasing factor for physiological responses in chicks.
Physiol. Behav. **125**, 57–61. (doi:10.1016/j.physbeh.2013.11.006)

31. Wang J, Gilbert ER, Cline MA. 2019 Hypothalamic mechanisms associated with
neuropeptide K-induced anorexia in Japanese quail (Coturnix japonica). *Comp.*
Biochem. Physiol. A Mol. Integr. Physiol. **237**, 110539.
(doi:10.1016/j.cbpa.2019.110539)

32. Edelmann M, Wolfe C, Scordalakes EM, Rissman EF, Tobet S. 2007 Neuronal nitric
oxide synthase and calbindin delineate sex differences in the developing
hypothalamus and preoptic area. *Dev. Neurobiol* **67**, 1371–1381.
(doi:10.1002/dneu.20507)

33. Gilmore RF, Varnum MM, Forger NG. 2012 Effects of blocking developmental cell
death on sexually dimorphic calbindin cell groups in the preoptic area and bed
nucleus of the stria terminalis. *Biol. Sex Differ.* **3**, 5. (doi:10.1186/2042-6410-3-5)

34. Garcia-Calero E, Scharff C. 2013 Calbindin expression in developing striatum of
zebra finches and its relation to the formation of area X. *J. Comp. Neurol.* **521**, 326–
341. (doi:https://doi.org/10.1002/cne.23174)

35. Moe Y *et al.* 2016 A comparative study of sex difference in calbindin neurons among
mice, musk shrews, and Japanese quails. *Neurosci. Lett.* **631**, 63–69.
(doi:10.1016/j.neulet.2016.08.018)

36. Pinaud R, Fortes AF, Lovell P, Mello CV. 2006 Calbindin-positive neurons reveal a
sexual dimorphism within the songbird analogue of the mammalian auditory cortex.
J. Neurol. **66**, 182–195. (doi:https://doi.org/10.1002/neu.20211)

37. Harris EP, Abel JM, Tejada LD, Rissman EF. 2016 Calbindin Knockout Alters Sex-
Specific Regulation of Behavior and Gene Expression in Amygdala and Prefrontal
Cortex. *Endocrinology* **157**, 1967–1979. (doi:10.1210/en.2016-1055)

38. Brimblecombe KR, Vietti-Michelina S, Platt NJ, Kastli R, Hnieno A, Gracie CJ, Cragg
SJ. 2019 Calbindin-D28K Limits Dopamine Release in Ventral but Not Dorsal
Striatum by Regulating Ca2+ Availability and Dopamine Transporter Function. *ACS*
Chem. Neurosci. **10**, 3419–3426. (doi:10.1021/acschemneuro.9b00325)

39. Malki K *et al.* 2016 Transcriptome analysis of genes and gene networks involved in
aggressive behavior in mouse and zebrafish. *Am. J Med. Genet. B Neuropsychiatr.*
Genet. **171**, 827–838. (doi:https://doi.org/10.1002/ajmg.b.32451)

40. Sheng M, Greenberg ME. 1990 The regulation and function of c-fos and other
immediate early genes in the nervous system. *Neuron* **4**, 477–485.
(doi:10.1016/0896-6273(90)90106-P)

41. Michael Numan. 2020 *The Parental Brain: Mechanisms, Development, and*
Evolution. Oxford, New York: Oxford University Press.

42. Barbotin A-L, Peigné M, Malone SA, Giacobini P. 2019 Emerging Roles of Anti-
Müllerian Hormone in Hypothalamic-Pituitary Function. *Neuroendocrinology* **109**,
218–229. (doi:10.1159/000500689)

43. Silva MSB, Giacobini P. 2020 New insights into anti-Müllerian hormone role in the
hypothalamic–pituitary–gonadal axis and neuroendocrine development. *Cell. Mol.*
Life Sci. (doi:10.1007/s00018-020-03576-x)

44. Brimijoin S, Chen VP, Pang Y-P, Geng L, Gao Y. 2016 Physiological Roles for
Butyrylcholinesterase: A BChE-Ghrelin Axis. *Chem. Biol. Interact.* **259**, 271–275.
(doi:10.1016/j.cbi.2016.02.013)

45. Chen VP, Gao Y, Geng L, Parks RJ, Pang Y-P, Brimijoin S. 2015 Plasma
butyrylcholinesterase regulates ghrelin to control aggression. *Proc. Natl Acad. Sci.*
USA **112**, 2251–2256. (doi:10.1073/pnas.1421536112)

46. Joo Y *et al.* 2009 Chronic immobilization stress induces anxiety- and depression-like
behaviors and decreases transthyretin in the mouse cortex. *Neurosci. Lett.* **461**,
121–125. (doi:10.1016/j.neulet.2009.06.025)

47. Stork O, Stork S, Pape H-C, Obata K. 2001 Identification of Genes Expressed in the
Amygdala During the Formation of Fear Memory. *Learn. Mem.* **8**, 209–219.
(doi:10.1101/lm.39401)

48. Zhang Z, Dawson PA, Piper M, Simmons DG. 2019 Postnatal N-acetylcysteine
administration rescues impaired social behaviors and neurogenesis in Slc13a4
haploinsufficient mice. *EBioMedicine* **43**, 435–446.
(doi:10.1016/j.ebiom.2019.03.081)

49. Stolzenberg DS, Hernandez-D'Anna KL, Bosch OJ, Lonstein JS. 2019 Maternal
Behavior From a Neuroendocrine Perspective. *Oxford Research Encyclopedia of*
Neuroscience. (doi:10.1093/acrefore/9780190264086.013.237)

50. Fischer EK, O'Connell LA. 2017 Modification of feeding circuits in the evolution of
social behavior. *Journal of Experimental Biology* **220**, 92–102.
(doi:10.1242/jeb.143859)

51. Almuttaqi H, Udalova IA. 2019 Advances and challenges in targeting IRF5, a key
regulator of inflammation. *The FEBS Journal* **286**, 1624–1637.
(doi:https://doi.org/10.1111/febs.14654)

52. Mamun AA *et al.* 2020 Microglial IRF5-IRF4 regulatory axis regulates
neuroinflammation after cerebral ischemia and impacts stroke outcomes. *Proc. Natl*
Acad. Sci. USA **117**, 1742–1752. (doi:10.1073/pnas.1914742117)

53. Haim A, Julian D, Albin-Brooks C, Brothers HM, Lenz KM, Leuner B. 2017 A survey
of neuroimmune changes in pregnant and postpartum female rats. *Brain Behav.*
Immun. **59**, 67–78. (doi:10.1016/j.bbi.2016.09.026)

54. Bukhari SA, Saul MC, James N, Bensky MK, Stein LR, Trapp R, Bell AM. 2019
Neurogenomic insights into paternal care and its relation to territorial aggression.
Nat. Commun. **10**, 4437. (doi:10.1038/s41467-019-12212-7)

55. Bentz AB, Rusch DB, Buechlein A, Rosvall KA. 2019 The neurogenomic transition
from territory establishment to parenting in a territorial female songbird. *BMC*
Genomics **20**, 819. (doi:10.1186/s12864-019-6202-3)

56. Ye P, Li M, Liao W, Ge K, Jin S, Zhang C, Chen X, Geng Z. 2019 Hypothalamic
transcriptome analysis reveals the neuroendocrine mechanisms in controlling

broodiness of Muscovy duck (Cairina moschata). *PLOS ONE* **14**, e0207050.
(doi:10.1371/journal.pone.0207050)