Can citizen science analysis of camera trap data be used to study reproduction? Lessons from Snapshot Serengeti program

Thel Lucie[1], Chamaillé-Jammes Simon[2, 3, 4], Keurinck Léa[1], Catala Maxime[1], Packer Craig[5], Huebner Sarah E.[5], Bonenfant Christophe[1, 4]

[1] Laboratoire de Biométrie et Biologie Évolutive, Unité Mixte de Recherche 5558, Bâtiment 711, Université Lyon I, 43 Boulevard du 11 novembre 1918, F-69622 Villeurbanne Cedex, France.

[2] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Université Paul Valéry Montpellier 3, Montpellier, France.

[3] Mammal Research Institute, Department of Zoology & Entomology, University of Pretoria, Pretoria, South Africa.

[4] LTSER France, Zone Atelier "Hwange", Hwange National Park, Bag 62, Dete, Zimbabwe-CNRS HERD (Hwange Environmental Research Development) program.

[5] College of Biological Sciences, University of Minnesota, Saint Paul, MN 55108, U.S.A.

Corresponding author: lucie.mc.thel@gmail.com

19**Abstract**

20Ecologists increasingly rely on camera-trap data to estimate biological parameters such as

21population abundance. Because of the huge amount of data, the assistance of non-scientists is

22often sought after, but an assessment of the data quality is necessary. We tested whether

23volunteers data from one of the largest citizen science projects - Snapshot Serengeti - could

24be used to study breeding phenology. We tested whether the presence of juveniles (less than

25one or 12 months old) of species in the Serengeti: topi, kongoni, Grant's gazelle, could be

26reliably detected by the "naive" volunteers vs. trained observers. We expected a positive

27correlation between the proportion of volunteers identifying juveniles and their effective

28presence within photographs, assessed by the trained observers.

29	The agreement between the trained observers was good (Fleiss' $\kappa > 0.61$ for juveniles

30of less than one and 12 month(s) old), suggesting that morphological criteria can be used to

31determine age. The relationship between the proportion of volunteers detecting juveniles less

32than a month old and their actual presence plateaued at 0.45 for Grant's gazelle, reached 0.70

33for topi and 0.56 for kongoni. The same relationships were much stronger for juveniles

34younger than 12 months, reaching 1 for topi and kongoni. The absence of individuals < one

35month and the presence of juveniles < 12 months could be reliably assumed, respectively,

36when no volunteer and when all volunteers reported a presence of a young. In contrast, the

37presence of very young individuals and the absence of juveniles appeared more difficult to

38ascertain from volunteers' classification, given how the classification task was presented to

39them.

40	Volunteers' classification allows a moderately accurate but quick sorting of

41photograph with/without juveniles. We discuss the limitations of using citizen science

42camera-traps data to study breeding phenology, and the options to improve the detection of

43juveniles.

44**Keywords**: African ungulates, age determination, *Alcelaphus buselaphus,* camera trap,

45citizen science, *Damaliscus jimela, Nanger granti,* reproduction, Serengeti.

47**Introduction**

48Camera trapping is increasingly used for ecological monitoring due to its low cost, relative

49ease of use, and the variety of data it can supply (O'Connell et al. 2010). For instance, camera

50trap data are used to study species' occupancy and co-occurrence (Anderson et al. 2016),

51population dynamics (Karanth et al. 2006), or individual behaviour (*e.g.* vigilance behaviour:

52Chamaillé-Jammes et al. 2014, or diel activity patterns: Luo et al. 2019). A potential

53drawback of camera traps is the huge amounts of data that are generated (> 7 millions

54photographs for the Snapshot Serengeti initiative alone). Ecologists have realized that the

55benefits of continuously collecting data in the field can quickly be negated by the burden of

56database management, and visual inspection and analysis of photographs to record the

57desired data (Wearn and Glover-Kapfer 2017).

58   To process such a massive amount of information in a reasonable time, scientists have

59sought the help of non-specialists who perform diverse tasks like counting objects in

60photographs, describing picture content or identifying animal and plant species (*e.g.* McShea

61et al. 2015). Initially, part of the scientific community was sceptical about citizen science, in

62particular questioning data quality (Riesch and Potter 2014). However, volunteers have

63sometimes proven to be as efficient as experts, as for instance for the identification of large

64herbivore species in savanna ecosystems (Swanson et al. 2016). More recently, the advances

65in deep learning have led computers to become as efficient as people at identifying species,

66and, sometimes, behaviour classification problems (Norouzzadeh et al. 2018). However,

67human judgment is still valuable in particular cases where too little data are available to train

68models (*e.g.* active learning, Joshi et al. 2009), or when differences among the objects to be

69classified are subtle and classification requires some subjectivity (Miele et al. 2020, *in prep.*).

70We believe this is the case for age classification problems, for which to the best of our

71knowledge the number of precisely labelled pictures is currently too small to allow an

72efficient and reliable automatization of the process.

73   Under the assumption that the detectability of juveniles and adult females segments of

74the population is not biased by camera traps methodology, classifying individuals into age-

75classes such as juveniles and adults would allow estimates of key demographic parameters

76(*e.g.* reproductive rates) or life-history traits (*e.g.* breeding phenology), which are essential in

77estimating population growth rates (Sibly and Hone 2002). For instance, Ogutu and

78colleagues (2008) highlighted that rainfall influences the abundance of several large

79herbivore species of the Mara-Serengeti ecosystem, by acting differently on each segment of

80the populations at specific periods of the year. Furthermore, it would facilitate the study of

81the relationships between population characteristics and their environments such as between

82birth phenology, diet and food resource availability (Sinclair et al. 2000), or their potential

83evolution in the context of climate change (Visser and Both 2005). Until now, the study of

84those key demographic parameters has been mainly conducted by direct field observations

85(*e.g.* Côté and Festa-Bianchet 2001 in mountain goats, Plard et al. 2013 in roe deer).

86However, this methodology still requires an intensive and often costly field effort. Identifying

87and counting juveniles from camera traps could reduce this field effort, or allow larger-scale

88or longer-term studies, as suggested by Hofmeester et al. 2019, but could also be time-

89consuming because of tedious data processing. With the help of citizen science, data handling

90time could be substantially reduced, but the accuracy of non-specialists in detecting juveniles

91of large mammals from photographs has not yet been explored.

92   Here, we evaluate the usefulness of camera trap data annotated by citizen scientists

93online to assess the presence of juveniles of large herbivores in the photographs. We use

94photographs and citizen classifications from the Snapshot Serengeti project (Swanson et al.

952015), one of the world's largest citizen science programs, on a subset of the data. We focus

96on the detection of juveniles in three species found in the Serengeti National Park, Tanzania

97with contrasting social and neonatal behaviours: topi (*Damaliscus jimela*), kongoni

98(*Alcelaphus cokii*) and Grant's gazelle (*Nanger granti*). We first evaluate the agreement

99between trained observers from our research team, and then test the ability of the volunteers

100to detect juveniles by comparing their classification with ours. We predict a better agreement

101between trained observers for the youngest age class because determination criteria are

102clearer and easier to identify than for older juveniles (*e.g.* absence of horns). Consequently,

103the level of agreement should decrease for age classes that are based on more subjective or

104difficult-to-assess criteria (*e.g.* shape of the horns). Under the hypothesis that volunteers

105could generally identify juveniles correctly, we expect a positive relationship between the

106proportion of volunteers reporting a juvenile on a photograph and the probability of the actual

107presence of a juvenile, as determined by the trained observers. Again, we expect the

108correlation to be stronger for the youngest age class of juveniles because they are easier to

109differentiate from adults. Across species, we expect a higher agreement and correlation for

110topi and kongoni than for Grant's gazelle because the former are larger, live in smaller groups

111and have similar body growth rate between males and females (Wilson and Mittermeier

1122011), hence reducing the risk of confusion between young males and older females. Overall,

113our study details the strengths and weaknesses of camera trap data, in particular when

114classified by citizen scientists, for the study of reproductive traits such as reproductive rates

115or breeding phenology.

Materials and methods

Study site

The surveyed area within the Serengeti National Park, Tanzania, is composed of open plains and savanna woodlands. Rain mostly occurs between November and June (wet season), with mean annual rainfall increasing from 500 mm in the southeast to 1,100 mm in the northwest. This area harbours a rich community of large herbivores, composed of gregarious and migratory wildebeest (*Connochaetes mearnsi*), zebra (*Equus sp.*) and Thomson's gazelle (*Eudorcas nasalis*), but also resident populations such as Cape buffalo (*Syncerus caffer*) or warthog (*Phacochoerus africanus*) (Sinclair and Norton-Griffiths 1995). Community dynamics are driven both by herbivores, maintaining an open state of the grassland by intensive grazing (McNaughton 1985, Sinclair and Norton-Griffiths 1995) as well as large predators (*e.g.* lion (*Panthera leo*) and hyena (*Crocuta crocuta*), Sinclair and Norton-Griffiths 1995).

Camera trap data

The Snapshot Serengeti camera trap grid was deployed in 2010 in Serengeti National Park, Tanzania, to monitor lions and their prey, though the bycatch of numerous other species has proven useful as well. Running continuously since 2010, the grid spans 1,125 km² in the center of the park. We used data provided by Snapshot Serengeti camera survey recorded between July 2010 and April 2013 (Supporting information 1). The camera traps were set ~ 50 cm above ground in the centre of a 5 km² grid cell. The detection radius was approximately 45° and their field of view about 14 m (Swanson et al. 2015). Cameras took a rapid series of three pictures upon trigger of the motion and heat sensors ("capture event" in Swanson et al. 2015, hereafter called a "sequence" following Meek et al. 2014) in a few seconds interval, with a one-minute delay between sequences.

Choice of studied species and sorting steps of the dataset

Among the many large herbivore species present in the study site, we selected topi, kongoni

and Grant's gazelle due to their contrasting biology and characteristics useful to assess the

age classes of individuals. The criteria considered were (1) number of available sequences,

(2) relatively small group size, (3) presence of horns in males and females, (4) relatively large

size of the young (young of larger species are larger, and therefore criteria like horns are

easier to detect), (5) contrasting anti-predator of behaviour of the young (Supporting

information 2).

We selected the final dataset (n = 2,359 sequences) to conduct the analyses following

several sorting steps, based on the detection of the species of interest and of the presence of

juveniles from the initial complete dataset (n = 1,184,657 sequences) by the volunteers. We

then corrected this dataset thanks to the trained observers reclassifications (see details in

Table 1).

Table 1: number of sequences at each sorting step from the extraction of raw data to the selection of all the

independent sequences with at least one individual < 1 month old, for the three species of the study: topi,

kongoni and Grant's gazelle (pictures from Snapshot Serengeti program, Tanzania, between July 2010 and

April 2013). Standard deviations are calculated on the basis of the classifications of the three trained observers.

Step n°	Step name	Consists in	Classifier	Number of sequences		
				Topi	Kongoni	Grant's gazelle
1	Row data	all sequences produced during the survey	none	1,184,657		
2	Blank sorting	sequences with animals	volunteers	319,915		
3	Species sorting 1	sequences with study species	volunteers	2,299	12,431	7,723
4	Young sorting 1	sequences with at least more than 0% of volunteers identifying young	volunteers	281	1,290	1,095
5	Species	sequences with study species	trained	324	1,281	754

	sorting 2	corrected by trained observers	observers			
6	Young sorting 2	sequences with at least one young	trained observers	216 ± 1	830 ± 17	348 ± 18
7	Young sorting 3	sequences with at least one young less than one month old	trained observers	59 ± 9	137 ± 33	58 ± 3
8	Re-sight sorting 1	independent sequences (remove of resights notified by the observers)	trained observers	39 ± 7	71 ± 27	50 ± 5
8'	Re-sight sorting 2	independent sequences (remove of sequences taken less than 10 minutes after the previous one by the same camera trap)	theoretical sorting	42 ± 8	70 ± 22	40 ± 4

Assessing the presence or absence of juveniles in photographs

All Snapshot Serengeti photographs have been uploaded to the online citizen science

platform "The Zooniverse" (www.zooniverse.org) to be classified by volunteers. Each

sequence was classified by as many as 25 volunteers (*minimum* = 11, *maximum* = 57,

Swanson et al. 2016), who each identified what species was present in the sequence and

recorded whether at least one juvenile was present or not. Note that no formal definition of a

juvenile was provided to volunteers, nor were there any particular guidelines about how to

identify a juvenile. Volunteers simply searched and checked "young" in the Zooniverse

interface when they suspected the presence of non-adult individuals. Regarding age classes,

the only question volunteers had to reply to was: "Are there any young present?". For each

sequence, the volunteers' classifications were then compiled via a plurality algorithm to yield

a consensus classification, leading to a proportion of volunteers (*Pv*) who identified at least

one juvenile in each sequence (see details in Swanson et al. 2015). Here, we used all

sequences where volunteers identified topi, kongoni, and Grant's gazelle, with at least one

volunteer (*Pv* > 0%) having annotated the presence of a juvenile. We could not analyse all

175sequences for which no volunteer had reported a juvenile, as these were too many ($n = 2{,}018$,

17611,141 and 6,628 for topi, kongoni and Grant's gazelle respectively) to be reviewed

177individually. However, we checked a subset of them ($n = 1{,}000$ for each species), and the

178chance that a trained observer observed a "true" juvenile (*i.e.* of less than 12 months old, see

179age classes definition below) was < 6.5% for all three species studied. We did not correct

180observations for recaptures of the same individuals as we were only interested in the ability

181of volunteers to detect the presence of juveniles on the sequences, but not the actual number

182of juveniles.

183 Three of us (LT, LK and MC), considered here as trained observers, searched all

184sequences retained for juveniles, which were assigned to an age class when detected. We

185used previously published morphological descriptions of the studied species (*e.g.* shape and

186size of horns, size relating to the adult; Supporting information 3) to identify and age

187individuals. We distinguished between (1) juveniles < one month, (2) between one and six

188months, (3) between six and 12 months, (4) between 12 and 24 months, termed yearling

189hereafter and (5) individuals over 2 years old, termed adults hereafter. We defined age classes

190according to biological characteristics relevant to juvenile identification for each species (*e.g.*

191very young individuals for birth phenology identification, juveniles under one year for

192recruitment estimation). We recorded observers' classifications with Aardwolf software

193(Krishnappa and Turner 2014). Ultimately, we produced a dataset describing the presence or

194absence ($M_{i,s,j}$), in each sequence of individuals of any of the five age categories i, for the

195species s, by trained observer j.

197*Statistical analyses*

198We first evaluated the agreement between the three trained observers on the detection of

199individuals assigned to each age class for each species. We measured this agreement with the

200Fleiss' κ, implemented in the *"raters"* R package (Quatto and Ripamonti 2014). Fleiss' κ

201(Fleiss 1971) is the comparison of agreement between 2+ judges and the level of agreement

202expected by chance alone. It takes values between -1 and 1, values < 0 indicating an

203agreement lower to what could be expected by chance, values > 0 indicating a greater

204agreement than expected by chance, and values = 0 indicating an agreement close to random.

205We tested for significance of the difference between the Fleiss' κ using a bootstrap procedure

206following Vanbelle and Albert (2008) (Supporting information 4).

207　　　　We tested the relationship between the proportion of volunteers identifying at least

208one juvenile (*Pv*) and the probability that trained observers had identified at least one juvenile

209< 1 month (category $i = 1$). We also explored the same relationship with juveniles < 12

210months (therefore including juveniles of categories 1 to 3 above). We fitted three generalized

211estimating equation models: one linear (see equation 1 below) and two piecewise models.

212The first piecewise model was characterized by a slope on both sides of the threshold

213(equation 2), the second by a slope before and a plateau after the threshold (equation 3). We

214fitted piecewise models to search for a potential "saturation" phenomenon, whereby beyond a

215specific proportion of volunteers the probability to effectively observe a juvenile does not

216increase anymore. We also fitted the null model for comparison. All the models were fitted

217for the two age classes and for each species individually, using the *wgee* function

218implemented in the *"wgeesel"* R package (Xu et al. 2018). We selected the best model using

219the Quasi-likelihood under the Independence model Criterion QIC (thresholds selected by

220comparison of QIC of the models for each species and age class as well). It is a modification

221of the Akaike Information Criterion AICc, suitable when quasi-likelihood is used instead of

222likelihood (Pan 2001), implemented in the *"MuMIn"* R package (Barton 2019). We used a

223logit link function and a binomial distribution of errors (Agresti 2002), considering the

224proportion of volunteers identifying at least one juvenile as fixed effect, and the identity of

225the sequence as clustering variable with an exchangeable correlation structure:

$$logit(M_{s,i,j}) = \mu + \beta_{s,i} \times Pv_s \qquad \text{(eqn 1 – linear model)}$$

and

$$\mu + \beta 1_{s,i} \times Pv_s \; if \; Pv_s \leq \theta$$

$$logit(M_{s,i,j}) = \begin{cases} & \end{cases} \qquad \text{(eqn 2 – piecewise model 1)}$$

$$\mu + \beta 2_{s,i} \times Pv_s \; if \; Pv_s > \theta$$

and

$$\mu + \beta 1_{s,i} \times Pv_s \; if \; Pv_s \leq \theta$$

$$logit(M_{s,i,j}) = \begin{cases} & \end{cases} \qquad \text{(eqn 3 – piecewise model 2)}$$

$$\mu 2 \; if \; Pv_s > \theta$$

235where $M_{s,i,j}$ denotes the presence or absence of at least one juvenile of the age class i (< 1 or

23612 months) for a given sequence, for the species s and the observer j, μ is the expected mean

237probability of actual presence of at least one juvenile of the age class i when none of the

238volunteers identified the presence of a juvenile, Pv_s is the proportion of volunteers identifying

239at least one juvenile for a given sequence, for the species s and θ is the best threshold. We

240performed all analyses using the R statistical software (R Core Development Team 2019).

Results

243From the Snapshot Serengeti monitoring program, we obtained 281, 1,290 and 1,095

244sequences for topi, kongoni and Grant's gazelle respectively (Fig. 1a) sorting step "Young

245sorting 1" and Table 1). Our selection process led to a dramatic drop in the number of usable

246sequences (*i.e.* those containing at least one < 1 month old juvenile), with a tenfold reduction

247at the species level: it only remained between 7.7% and 18.1% of the complete dataset

248available for the species (*n* = 59 ± 9 SD, 137 ± 33 SD and 58 ± 3 SD for each species

249respectively, Fig. 1a) sorting step "young sorting 3" and Table 1). The largest losses of

250sequences happened during the phases of selection of sequences containing the studied

251species and then juveniles (Fig. 1a) and Table 1). Fig. 1b) suggests that we would obtain

252similar results concerning the loss of sequences after selection for sequences containing

253juveniles for any large herbivore species recorded in the Snapshot Serengeti program.

255*Figure 1: a) number of sequences available at each sorting step from the extraction of raw data to the selection*

256*of all the independent sequences with at least one individual < 1 month old, for the three species of the study:*

257*topi, kongoni, Grant's gazelle. Pictures from Snapshot Serengeti, Tanzania, between July 2010 and April 2013.*

258*Raw data: all sequences produced during the survey; blank sorting: sequences with animals; species sorting 1:*

259*sequences with study species according to the volunteers; young sorting 1: sequences with at least one*

260*volunteer identifying "young"; species sorting 2: sequences with study species corrected by trained observers;*

261*young sorting 2: sequences with at least one young according to the trained observers; young sorting 3:*

262*sequences with at least one young < 1 month old according to the trained observers; resight sorting:*

263*independent sequences once sequences taken less than 10 minutes after the previous one by the same camera*

264*trap and presenting the same species have been removed (following Palmer et al. 2018). Note the log scale for*

265*the ordinate axis, vertical bars represent the standard deviations. b) Number of sequences available for all the*

266*large herbivore species present in the study site. The proportions indicate the proportion of sequences where at*

267*least one volunteer identified juveniles for a given species.*

268   The three trained observers identified 993 ± 49 SD, 3,020 ± 180 SD and 2,128 ± 224

269SD individuals of any age class. On average, they could not assign an age class to ~11% (n =

270118 ± 68 SD), ~20% (n = 775 ± 256 SD) and 32% (n = 1,026 ± 492 SD) of the individuals for

271topi, kongoni and Grant's gazelle respectively, a significant between-species difference (χ^2 =

272262.83, *df* = 2, *p* < 0.001).

273   For all species, and as expected, the agreement between trained observers was highest

274for the youngest age class, then declined with age until the yearling class was reached (Fig.

2752). Yearlings were reasonably well classified in kongoni, but overly misclassified in topi and

276Grant's gazelle (Fig. 2). As expected, the agreement among trained observers was the highest

277for the youngest age class of juveniles, but also for the second age class and the adults, with

278Fleiss' κ values almost always > 0.61 (denoting a substantial agreement, following Landis

279and Koch 1977), except for Grant's gazelle aged between one and six months and adult topi.

280Agreement among observers was the highest for topi aged < 1 month old (Fleiss' κ = 0.78

281[0.72; 0.84]). In support of our prediction, we observed the lowest agreement for juveniles

282aged six months and older, and more obviously so for Grant's gazelle. Agreement among

283observers concerning the three first age classes pooled together (representing the juveniles)

284was very good, with Fleiss' κ values largely > 0.61 for all the species. The same holds true

285for juveniles between one and 12 months old when pooled together. Our results were globally

286consistent among the three species studied, with the highest agreement for topi, and the least

287for Grant's gazelle (Fig. 2). All estimated Fleiss' κ were significantly different from an

288agreement obtained by chance (Supporting information 4).

290Figure : Fleiss' κ denoting the level of agreement between the three trained observers (LT, LK and MC) in the

291identification of presence or absence of at least one individual belonging to any of the five age classes (< 1

292month, 1-6 months, 6-12 months, > 12 months, adults), for the three species of interest. Pictures from Snapshot

293Serengeti Program, Tanzania, between July 2010 and April 2013. The two combined age classes "1-12

294months" and "< 12 months" are also presented. Light grey dots: topi, dark grey dots: kongoni, black dots:

295Grant's gazelle. Vertical bars represent the confidence intervals. A Fleiss' κ near 1 denotes an almost perfect

296agreement, whereas a value near or < 0 means a very poor agreement between raters.

297   The model best describing our data for the age class < 1 month was the linear model

298for topi, the piecewise model with slopes on both sides of the threshold for kongoni and the

299piecewise model with slope before and plateau after the threshold for Grant's gazelle (Table

3002). The model best describing our data for the age class < 12 months was the threshold model

301with slopes on both sides of the threshold for topi, kongoni and Grant's gazelle (Table 2).

303 *: Statistics of models investigating the relationship between the proportion of volunteers identifying at*
304 *least one "young" and the probability of presence of at least one individual < 1 or 12 months, assessed by the*
305 *trained observers on a given sequence for the three species of interest: topi, kongoni and Grant's gazelle. Best*
306 *models are in bold. θ = estimated threshold (% of volunteers), QIC = Quasi-likelihood under the Independence*
307 *model Criterion, in. = intercept, estimates ± standard error and [95% confidence interval].*

Age class	Species	Model type	θ	QIC	Quasi-likelihood	Estimates
< 1 month	Topi	Null		924.6	-459.77	in. = -1.509 ± 0.133
		Linear		**630.9**	**-311.05**	**in. = -4.013 ± 0.342** **β = 0.048 ± 0.005**
		Piecewise - slope/plateau	90	635.2	-313.23	in. = -4.102 ± 0.362 $\beta1$ = 0.051 ± 0.005 μ2 = 0.626 [0.523; 0.720]
		Piecewise - slope/slope	78	634.7	-310.49	in. = -8.863 ± 1.763 $\beta1$ = 0.045 ± 0.007 $\beta2$ = 0.064 ± 0.023
	Kongoni	Null		2614.4	-1304.83	in. = -2.125 ± 0.081
		Linear		1993	-992.05	in. = -3.977 ± 0.174 β = 0.043 ± 0.003
		Piecewise - slope/plateau	90	2005.8	-998.42	in. = -4.007 ± 0.178 $\beta1$ = 0.044 ± 0.003 μ2 = 0.498 [0.434; 0.563]
		Piecewise - slope/slope	**10**	**1966.7**	**-977.58**	**in. = -10.522 ± 1.740** **$\beta1$ = 0.687 ± 0.178** **$\beta2$ = 0.039 ± 0.003**
	Grant's gazelle	Null		1236.2	-615.91	in. = -2.479 ± 0.117
		Linear		1016.6	-501.81	in. = -3.653 ± 0.181 β = 0.055 ± 0.007
		Piecewise - slope/plateau	**43**	**969.7**	**-480.75**	**in. = -4.355 ± 0.227** **$\beta1$ = 0.097 ± 0.008** **μ2 = 0.451 [0.353; 0.552]**

		Piecewise - slope/slope	4 3	974.6	-480.48	in. = -4.573 ± 0.544 $\beta1 = 0.094 \pm 0.010$ $\beta2 = 0.006 \pm 0.013$
< 12 months	Topi	Null		1241.3	-618	in. = 0.698 ± 0.111
		Linear		948	-468.64	in. = -0.866 ± 0.184 $\beta = 0.059 \pm 0.007$
		Piecewise - slope/plateau	6 0	964.6	-477.23	in. = -0.987 ± 0.191 $\beta1 = 0.069 \pm 0.007$ µ2 = 0.958 [0.926; 0.977]
		Piecewise - slope/slope	**3 0**	**946.5**	**-464.45**	**in. = -3.202 ± 0.624** $\beta1 = 0.031 \pm 0.017$ $\beta2 = 0.090 \pm 0.022$
	Kongoni	Null		4992.3	-2493.63	in. = 0.609 ± 0.054
		Linear		3726.5	-1858.95	in. = -1.139 ± 0.108 $\beta = 0.089 \pm 0.006$
		Piecewise - slope/plateau	9 0	3726.6	-1859.03	in. = -1.140 ± 0.108 $\beta1 = 0.089 \pm 0.006$ µ2 = 0.999 [0.997; 1]
		Piecewise - slope/slope	**1 2**	**3684.8**	**-1835.45**	**in. = -3.041 ± 0.271** $\beta1 = 0.213 \pm 0.029$ $\beta2 = 0.066 \pm 0.006$
	Grant's gazelle	Null		3127.4	-1561.2	in. = -0.154 ± 0.067
		Linear		2940.5	-1463.22	in. = -0.837 ± 0.140 $\beta = 0.049 \pm 0.010$
		Piecewise - slope/plateau	3 0	2866.2	-1428.27	in. = -1.285 ± 0.132 $\beta1 = 0.091 \pm 0.009$ µ2 = 0.809 [0.748; 0.859]
		Piecewise - slope/slope	**4 1**	**2866.4**	**-1425.59**	**in. = -0.009 ± 0.506** $\beta1 = 0.083 \pm 0.010$ $\beta2 = -0.029 \pm 0.013$

309   The probability of observing a juvenile when all volunteers reported one was near 1

310for juveniles < 12 months for topi and kongoni. Concerning Grant's gazelle, this probability

only reached 0.90 when 41% of the volunteers recorded the presence of young. Between 41% and 100% of volunteers identifying young in the sequences, the probability decreased (Fig. 3 d), e), f)). When investigating the presence of juveniles < 1 month, the probability that a juvenile was actually present was never greater than 0.697 (Fig. 3 a), b), c)). On the other hand, the model predicted that when no volunteer reported the presence of juveniles, the probability that there was juveniles < 1 month was under 1.8%, but there was at least 9.6% chance to observe a juvenile < 12 months.

Figure 3: relationship, as predicted from the best model (see text for details), between the proportion of

volunteers identifying the presence of "young" and the probability of presence of at least one individual < 1

(a), b), c)) and < 12 (d), e), f)) months assessed by the trained observers in a given sequence (0 < proportion of

volunteers identifying "young" ≤ 1), for the three species of interest: a) and d) topi, b) and e) kongoni and c)

and f) Grant's gazelle. Pictures from Snapshot Serengeti Program, Tanzania, between July 2010 and April

2013. Light grey dots represent the probability of presence of at least one individual < selected age class in

each sequence assessed by the three trained observers, dark grey dots represent the mean of those probabilities

for each 10% volunteers interval, vertical bars represent 95% confidence intervals. Solid line represents

predicted values from the best model. Shaded areas represent 95% confidence intervals.

Contrary to our expectations, the piecewise model characterized by two slopes was

almost always selected as the best model for the three species and both age classes. This

denotes a sudden change in the rate of detection from a certain percentage of volunteers. The

detection rate decreases for kongoni and Grant's gazelle for young < 12 months beyond 12

and 41% of volunteers voting for the presence of young respectively. The number of

volunteers who did classify a photograph was independent from the probability to detect a

young.

Discussion

Our study reveals the strength and weaknesses of using citizen-based assessment of age

classes on camera trap pictures. Clearly, citizen involvement through an online platform has

been critical for the classification of the millions of photographs collected by the Snapshot

Serengeti initiative. Also, despite leaving it to the volunteers to decide what a juvenile looks

like, volunteers' classification allows a rough, moderately accurate, but quick sorting of

sequences with/without juveniles.

Our study makes clear that, for the species studied and given the minimal guidelines

given to volunteers, the *absence* of very young (< 1 month) individuals on pictures can be

reliably assumed when no volunteer reports a presence. The sequences almost never labelled

346by volunteers as containing "young" were very unlikely to contain young under one month of

347age according to the trained observers. This suggests that volunteers rarely miss very young

348juveniles. This likely occurs because very young juveniles are distinctively smaller than

349adults, and possibly also because very young mammals share some physical characteristics

350such as relatively large eyes, long legs, short and rounded nose, all belonging to

351*Kindchenschema*, known to be very attractive stimuli for humans (Brosch et al. 2007, Golle

352et al. 2013). By contrast, the *presence* of very young individuals appears more difficult to

353ascertain from volunteers' data, and this apparently comes from the lack of guidelines given

354to volunteers. Indeed, consistent with the idea that volunteers easily identify very young

355individuals, a large consensus among volunteers around the presence of a very young juvenile

356could be a reliable indication of actual presence, but only when no older juveniles are present

357(compare Fig. 3 a), b), c) & Supporting information 4 - Fig. A). Unfortunately, the absence or

358presence of older juveniles cannot currently be known without a reassessment of the pictures

359because volunteers were not asked to differentiate between juvenile age classes. Therefore,

360the presence of very young juveniles remains difficult to ascertain. On the other hand, the

361*presence* of juveniles, irrespectively of their age, can be reliably assumed when all volunteers

362agree about a presence (Fig. 3 a), b), c)), especially for topi and kongoni: the model predicted

363that when all the volunteers reported the presence of juveniles, the probability that there was

364juveniles < 12 months was indeed at least 99.8%. This time, the *absence* of juveniles

365regardless of their age is less accurate, meaning that they are missed quite often. As they

366grow, juveniles become increasingly similar to adults, and are more likely to be mistaken for

367the latter by volunteers. Anyhow, we emphasize that detectability of juveniles is not equal

368between species (Fig. 3).

369   Among the three trained observers, we found the best agreement in the detection of

370age classes for a given sequence for topi and kongoni, suggesting that they are the easiest

371species to classify, and Grant's gazelle the hardest. The combination of small body size and

372gregariousness could explain why the determination of age was more difficult and

373heterogeneous for Grant's gazelle than for the two other species. Topi and kongoni are fairly

374large, and small body size makes the detection of some inconspicuous age criteria

375challenging (*e.g.* presence or absence of very small horn buds on the forehead of the

376individuals). On the contrary, it is certainly easier to discriminate between adults and

377juveniles in the largest herbivores, such as giraffe (*Giraffa camelopardalis*) or elephant

378(*Loxodonta africana)* because of the marked size difference between them. Discrimination

379between several juvenile age classes will certainly be difficult still as individuals are often

380only partially spotted by camera traps and subtle aging criteria are not visible. More

381generally, in species forming large herds such as wildebeest, young might be particularly

382difficult to spot and are likely missed frequently. In large groups, the body of many

383individuals overlaps, hampering our ability to see aging criteria correctly, and to age them

384accurately. This could explain why in our study for instance, Grant's gazelle frequently

385occurring in large herds reaching more than 30 individuals in some sequences, is also the one

386with the lowest identification success of juveniles. Finally, a pronounced sexual dimorphism

387in horn growth in Grant's gazelle likely led to confusions between the first two age classes as

388male horns grow faster, a young male can look like an older female when using the length of

389the horns as an aging criteria. This would also be the case for species with a sexual

390dimorphism in which only males grow horns like impala (*Aepyceros melampus*) or waterbuck

391(*Kobus defassa*). Clearly, a description of the morphological changes that occur throughout

392the development of young herbivores (*e.g.* Spinage 1976 in our species, Cunningham et al.

3932011, Dezeure et al. 2020 in other species) is of great value and substantially helps at

394reaching consistent results among different observers.

395   Another limitation of camera trapping in the context of reproductive studies is the

396level of effort required to obtain a sufficient sample size. Here, working with data from one

397of the world's largest and long-running camera trap studies, we ended up with a small

398number of sequences with juveniles < 1 month old after appropriate data selection. We

399identified four main explanations for this important reduction in exploitable sequences. First

400and foremost, our ability to determine an individual's age class depends strongly on the

401photograph quality and particularly its framing and exposure. In many cases, individuals were

402located too far from the camera or were only partly visible, or photographs were too blurry,

403dark, or overexposed to be scrutinized, leading to potentially significant loss of reproductive

404data and a high number of individuals of unknown or over-estimated age. Another potential

405source of information loss was species misidentification by volunteers. In our study, about

40630% of the sequences labelled with Grant's gazelle were misidentified because it greatly

407resembles species such as impala and Thomson's gazelle which are also present in the study

408site. Species abundance obviously directly impacts the number of sequences collected. The

409abundance of the three studied species is low compared to other ungulates in the Serengeti

410system. Sinclair reported 55,500 individuals topi, 20,700 individuals kongoni and 6,000

411individuals Grant's gazelle in the 1970s, whereas the numbers of wildebeest and zebra were

412720,000 and 240,000 individuals respectively (Sinclair and Norton-Griffiths 1995). Finally,

413the anti-predator strategy of juvenile large herbivores, known as the hider-follower gradient

414(Lent 1974, Rutberg 1987), could influence the number of sequences containing very young

415individuals. While followers become active and stick with their mother just a few hours after

416birth, hiders stay concealed in dense vegetation during their first weeks of life. The detection

417probability of hiders from camera traps should then be much lower than of followers,

418consistent with our observation of very young topi seen in a greater proportion than kongoni

419and Grant's gazelle.

Volunteer classification provides information that can reliably be used to infer the

presence of juveniles < 12 months or the *absence* of juveniles < 1 month, as these annotations

appear robust. However, because volunteers seem able to discriminate between individuals of

less than one month old and the rest of the juveniles even in the absence of any stated criteria,

more precise results could be achieved by asking them to differentiate between two age

classes of juveniles, such as "juvenile" and "newborn". The level of agreement between

trained observers in the classification of age classes according to species is also a good

indication of what kind of tasks could be successfully conducted by volunteers. When this

agreement is low (*e.g.* in the identification of topi and Grant's gazelle yearlings), one could

not expect volunteers to properly identify such an age class. We advise to limit the number of

classes the volunteers are asked to identify, and to focus on the most recognisable ones.

Another way to improve results generated via citizen science platforms could be the inclusion

of detailed information, as for species identification (Swanson et al. 2015). When a volunteer

detects a young on his/her photograph, he or she could be prompted with comprehensive keys

to age juvenile from its morphology along with a set of reference pictures or drawings.In

general, however, we would recommend to ask citizen scientists to identify newborns, *i.e.*

individuals under one month of age vs. other juveniles. To identify newborns of bovid species

with horns like in our study, volunteers would have to look for the smallest individuals (the

head does not come above the back of the female, Ogutu et al. 2008), with no evidence of

bud horns, with specific coat color (*e.g.* darker coat color in Grant's gazelle in our study, or

lighter coat color in wildebeest) or even with umbilical cord remnants. One difficulty is to

adapt the different criteria to the every species studied.

We finally suggest evaluating volunteers' classification skills by presenting them with

images of individuals of known age (captive or tagged animals for instance) and assessing

their accuracy compared to labels provided by experts. Volunteers could then be assigned

classification tasks adapted to their skills (*e.g.* species identification would belong to the

easiest tasks, whereas age classification would belong to the hardest). Snapshot Safari and
Zooniverse continue to create new modes of annotation that best leverage the public's interest
in contributing to research, and this is a logical next step for the Snapshot Safari initiative.
Overall, we find that by closely investigating the data already collected by volunteer-based
programs, data collection procedures can be adjusted to enhance the contributions of citizen
scientists to scientific research and conservation efforts.

Declarations

Acknowledgments: A PhD fellowship from the university Lyon 1 attributed to LT. This
work was performed using the computing facilities of the CC LBBE/PRABI. The authors
thank T. Michael Anderson for maintaining the camera traps in Tanzania, Zooniverse for
hosting "Snapshot Serengeti", >130,000 online volunteers for classifying the images, the
Tanzania Wildlife Research Institute and Tanzania National Parks for research permission,
and the Minnesota Supercomputing Institute (http://www.msi.umn.edu) for contributing to
data storage/processing and analysis.

Funding: A PhD fellowship from the University Lyon 1 attributed to LT.

Authors' contributions: LT, CB and SCJ conceived the ideas and designed methodology;
CP and SH organized data collection; LT, LK and MC analysed the data; LT, CB and SCJ led
the writing of the manuscript. All authors contributed critically to the drafts and gave final
approval for publication.

Data availability: Pictures available from the Labeled Image Library of Alexandria –
Biology and Conservation: http://lila.science/datasets/snapshot-serengeti and data from the
Dryad Data Repository: *to be completed*.

469**References**

470Agresti, A. 2002. Categorical data analysis, 3rd edition. – Wiley-Blackwell

471Anderson, T. M. et al. 2016. The spatial distribution of African savannah herbivores: species

472   associations and habitat occupancy in a landscape context. – Philos. Trans. R. Soc.

473   London, Ser B 371(1703): 20150314. https://doi.org/10.1098/rstb.2015.0314

474Bartoń, K. 2019. Mumin: Multi-model inference. Retrieved from https://CRAN.R-

475   project.org/package=MuMIn (R package version 1.43.10)

476Brosch, T. et al. 2007. That baby caught my eye... attention capture by infant faces. –

477   Emotion 7(3): 685–689. https://doi.org/10.1037/1528-3542.7.3.685

478Chamaillé-Jammes, S. et al. 2014. Innate threat-sensitive foraging: black-tailed deer remain

479   more fearful of wolf than of the less dangerous black bear even after 100 years of

480   wolf absence. – Oecologia 174(4): 1151–1158. https://doi.org/10.1007/s00442-013-

481   2843-0

482Côté, S. D. and Festa-Bianchet, M. 2001. Birthdate, mass and survival in mountain goat kids:

483   effects of maternal characteristics and forage quality. – Oecologia 127(2): 230–238.

484   https://doi.org/10.1007/s004420000584

485Cunningham, P. L. et al. 2011. Morphological characteristics and ageing criteria of sand

486   gazelle (gazella subgutturosa marica thomas, 1897) and their use for wildlife

487   management. – Eur. J. Wildl. Res. 57(4): 865–872. https://doi.org/10.1007/s10344-

488   011-0498-x

489Dezeure, J. et al. 2020. Developmental transitions in body color in chacma baboon infants:

490   Implications to estimate age and developmental pace. – Am. J. Phys. Anthropol.

491   e24118. https://doi.org/10.1002/ajpa.24118

492Fleiss, J. L. 1971. Measuring nominal scale agreement among many raters. – Psychol. Bull.

493   76(5): 378. https://doi.org/10.1037/h0031619

494Golle, J. et al. 2013. Sweet puppies and cute babies: Perceptual adaptation to babyfacedness

495   transfers across species. – PLoS One 8(3): e58248.

496   https://doi.org/10.1371/journal.pone.0058248

497Hofmeester, T. R. et al. 2019. Using by-catch data from wildlife surveys to quantify climatic

498   parameters and the timing of phenology for plants and animals using camera traps. –

499   Remote Sens. Ecol. Conserv. https://doi.org/10.1002/rse2.136

500Joshi, A. J. et al. 2009. Multi-class active learning for image classification. In: 2009 ieee

501   conference on computer vision and pattern recognition: pp. 2372–2379.

502   https://doi.org/10.1109/cvpr.2009.5206627

503Karanth, K. U. et al. 2006. Assessing tiger population dynamics using photographic capture–

504   recapture sampling. – Ecology 87(11): 2925–2937. https://doi.org/10.1890/0012-

505   9658(2006)87[2925:atpdup]2.0.co;2

506Krishnappa, Y. S. and Turner, W. C. 2014. Software for minimalistic data management in

507   large camera trap studies. – Ecol. Inf. 24: 11–16.

508   https://doi.org/10.1016/j.ecoinf.2014.06.004

509Landis, J.R. and Koch, G. G. 1977. The measurement of observer agreement for categorical

510   data. – Biometrics: 159–174. https://doi.org/10.2307/2529310

511Lent, P. C. 1974. Mother-infant relationships in ungulates. In: Geist V, Walther F (eds), The

512   behaviour of ungulates and its relationship to management. – IUCN Publications

513   24(1), Morges, Switzerland, pp. 14-55.

514Luo, G. et al. 2019. Habitat use and diel activity pattern of the tibetan snowcock (tetraogallus

515   tibetanus): a case study using camera traps for surveying high-elevation bird species.

516   – Avian Res. 10(1): 4. https://doi.org/10.1186/s40657-019-0144-y

517McNaughton, S. 1985. Ecology of a grazing ecosystem: the Serengeti. – Ecol. Monogr.

55(3): 259–294. https://doi.org/10.2307/1942578

McShea, W. J. et al. 2015. Volunteer-run cameras as distributed sensors for macrosystem

mammal research. – Landscape Ecol. 31(1): 55–66. https://doi.org/10.1007/s10980-

015-0262-9

Meek, P. et al. 2014. Recommended guiding principles for reporting on camera trapping

research. – Biodivers. Conserv. 23(9): 2321–2343. https://doi.org/10.1007/s10531-

014-0712-8

Miele, V. et al. 2020. Revisiting giraffe photo-identification using deep learning and network

analysis. – bioRxiv. https://doi.org/10.1101/2020.03.25.007377

Norouzzadeh, M. S. et al. 2018. Automatically identifying, counting, and describing wild

animals in camera-trap images with deep learning. – PNAS: 201719367.

https://doi.org/10.1073/pnas.1719367115

O'Connell, A. F. et al. 2010. Camera traps in animal ecology: methods and analyses. –

Springer Science & Business Media

Ogutu, J. O. et al. 2008. Rainfall influences on ungulate population abundance in the Mara-

Serengeti ecosystem. – J. Anim. Ecol. 77(4): 814-829. 10.1111/j.1365-

2656.2008.01392.x

Palmer, M. S. et al. 2018. Evaluating relative abundance indices for terrestrial herbivores

from large-scale camera trap surveys. – Afr. J. Ecol. 56(4): 791–803.

https://doi.org/10.1111/aje.12566

Pan, W. 2001. Akaike's information criterion in generalized estimating equations. –

Biometrics 57(1): 120–125. https://doi.org/10.1111/j.0006-341x.2001.00120.x

Plard, F. et al. 2013. Parturition date for a given female is highly repeatable within five roe

deer populations. – Biol. Lett. 9(1): 20120841. https://doi.org/10.1098/rsbl.2012.0841

Quatto, P. and Ripamonti, E. 2014. raters: A modification of fleiss' kappa in case of nominal

and ordinal variables. Retrieved from https://CRAN.R-project.org/package=raters (R

package version 2.0.1)

Riesch, H. and Potter, C. 2014. Citizen science as seen by scientists: Methodological,

epistemological and ethical dimensions. – Public Underst. Sci. 23(1): 107–120.

https://doi.org/10.1177/0963662513497324

Rutberg, A. T. 1987. Adaptive hypotheses of birth synchrony in ruminants: an interspecific

test. – Am. Nat. 130(5): 692–710. https://doi.org/10.1086/284739

Sibly, R. M. and Hone, J. 2002. Population growth rate and its determinants: an overview. –

Philos. Trans. R. Soc. London, Ser B 357(1425): 1153–1170.

https://doi.org/10.1098/rstb.2002.1117

Sinclair, A. et al. 2000. What determines phenology and synchrony of ungulate breeding in

serengeti? – Ecology 81(8): 2100–2111. https://doi.org/10.1890/0012-

9658(2000)081[2100:wdpaso]2.0.co;2

Sinclair, A. R. E. and Norton-Griffiths, M. 1995. Serengeti: dynamics of an ecosystem. –

University of Chicago Press

Spinage, C. 1976. Age determination of the female grant's gazelle. – Afr. J. Ecol. 14(2): 121–

134. https://doi.org/10.1111/j.1365-2028.1976.tb00157.x

Swanson, A. et al. 2016. A generalized approach for producing, quantifying, and validating

citizen science data from wildlife images. – Conserv. Biol. 30(3): 520–531.

https://doi.org/10.1111/cobi.12695

Swanson, A. et al. 2015. Snapshot Serengeti, high-frequency annotated camera trap images of

40 mammalian species in an African savanna. – Sci. Data 2: 150026.

https://doi.org/10.1038/sdata.2015.26

Vanbelle, S. and Albert, A. 2008. A bootstrap method for comparing correlated kappa

coefficients. – J. Stat. Comput. Simul. 78(11): 1009–1015.

https://doi.org/10.1080/00949650701410249

Visser, M. E. and Both, C. 2005. Shifts in phenology due to global climate change: the need

for a yardstick. – Proc. R. Soc. B 272(1581): 2561–2569.

https://doi.org/10.1098/rspb.2005.3356

Wearn, O. and Glover-Kapfer, P. 2017. Camera-trapping for conservation: a guide to best-

practices [Internet]. – WWF Conservation Technology Series 1.

Wilson, D. E. and Mittermeier, R. A. 2011. Handbook of the mammals of the world. vol 2.

Hoofed mammals. – Lynx Edicions

Xu, C. et al. 2018. wgeesel: Weighted generalized estimating equations and model selection.

Retrieved from https://CRAN.R-project.org/package=wgeesel (R package version

1.5)