1

Reproductive proteins evolve faster than non-reproductive proteins among *Solanum* species

10
11
12

Leonie C. Moyle[1*], Meng Wu[1], Matthew J. S. Gibson[1]

[1] Dept of Biology, Indiana University, Bloomington, Indiana, USA

* Correspondence:
lmoyle@indiana.edu

Keywords: female, male, molecular evolution, reproductive protein, sexual selection, *Solanum*, tomato

Abstract

Elevated rates of evolution in reproductive proteins are commonly observed in animal species, and are thought to be driven by the action of sexual selection and sexual conflict acting specifically on reproductive traits. Whether similar patterns are broadly observed in other biological groups is equivocal. Here we examine patterns of protein divergence among wild tomato species (Solanum section Lycopersicon), to understand forces shaping the evolution of reproductive genes in this diverse, rapidly evolving plant clade. By comparing rates of molecular evolution among loci expressed in reproductive and non-reproductive tissues, our aims were to test if: a) reproductive-specific loci evolve more rapidly, on average, than non-reproductive loci; b) 'male'-specific loci evolve at different rates than 'female'-specific loci; c) genes expressed exclusively in gametophytic (haploid) tissue evolve differently from genes expressed in sporophytic (diploid) tissue or in both tissue types; and d) mating system variation (a potential proxy for the expected strength of sexual selection and/or sexual conflict) affects patterns of protein evolution. We observed elevated evolutionary rates in reproductive proteins; however this pattern was most evident for female- rather than male-specific loci, both broadly and for individual loci inferred to be positively selected. These elevated rates might be facilitated by greater tissue-specificity of reproductive proteins, as faster rates were associated with more narrow expression domains. In contrast we found little evidence that evolutionary rates are consistently different in loci experiencing haploid selection (gametophytic-exclusive loci), or in lineages with quantitatively different mating systems. Overall while reproductive protein evolution is generally elevated in this diverse plant group, specific patterns of protein evolution are more complex than those reported in other (largely animal) systems, and include a more prominent role for female-specific loci among adaptively evolving genes.

1   Introduction

The rapid evolution of reproductive proteins has been observed across many different animal species, including in groups as diverse as marine invertebrates, insects, and mammals (Swanson and Vacquier 2002; Clark et al. 2006; Turner and Hoekstra 2008). This pattern is especially well established for proteins involved in male-specific functions, such as seminal fluid proteins, and sperm-egg and sperm-reproductive tract (RT) interactions. Although many factors can influence molecular evolutionary rates, two specific evolutionary forces—sexual selection and sexual conflict—have been proposed as primary drivers of this accelerated evolution, because both processes are expected to differentially affect reproductive traits and the proteins that underlie them. That is, traits mediating male-male competition and male-female mate choice experience unique selection to maximize mating and reproductive opportunities (Andersson 1994), and often appear to evolve rapidly between species (Ritchie 2007); therefore, the loci underpinning these reproductive traits might also be expected to display accelerated evolution and protein divergence between species (Swanson and Vacquier 2002, Clark et al. 2006, Vacquier and Swanson 2011). This inference is supported by observations of especially exaggerated protein evolution in male-exclusive and male-biased loci (collated in Dapper and Wade 2020), the sex that is usually subject to more intense sexual selection (Andersson 1994). Importantly, if sexual selection or conflict are the most critical factors driving rapid reproductive protein evolution, this pattern should be observable in other groups of organisms that also experience these selective forces. Nonetheless, whether reproductive proteins routinely evolve rapidly in non-animal species remains unclear.

Among these other groups, flowering plants (angiosperms) have numerous reproductive traits that can influence the operation and intensity of intrasexual competition and/or mate choice (Lloyd and Webb 1977, Willson 1979, Stephenson and Bertin 1983, Delph and Ashman 2006, Moore and Pannell 2011)—the two components of sexual selection. These include pollinator attraction traits, pollen competition traits, female reproductive tract ("pistil") traits (that influence pollen performance and fertilization success), and seed maturation traits (that can be used to exercise mate choice via selective abortion) (Willson 1994, Delph and Haven 1998, Skogsmyr and Lankinen 2002). Accordingly, angiosperms may also be expected to exhibit elevated evolutionary rates in loci underpinning these traits, similar to those inferred in animals (Clark et al. 2006). Indeed, of the few studies in angiosperms, some analyses have found evidence for faster protein evolution in reproductive loci (Szovenyi et al. 2013, Harrison et al. 2019), or greater representation of rapidly evolving genes among loci expressed in certain reproductive tissues (Gossman et al. 2016). However, others have shown more equivocal patterns (e.g. Gossman et al. 2014)—for example, sex-biased genes do not appear to evolve faster than non-sex-biased genes in several dioecious plant species (reviewed in Muyle 2019)—or even inferred greater levels of constraint on reproductive-specific proteins (e.g., Darolti et al. 2018).

These more complex findings could reflect the influence of additional factors on rates of protein evolution in plants, including the opportunity for haploid selection and the complex effects of mating system variation. In the first case, plants often express a substantial

proportion of their genome during the haploid phase of the life cycle (Hafidh et al. 2016),
thereby exposing these 'gametophytic' loci to haploid selection. In flowering plants,
gametophytic traits are largely reproductive and include 'male' functions such as pollen tube
germination, growth, and interactions with the female pistil, and 'female' functions such as
ovule signaling (Hafidh et al. 2016, Mizuta and Higashiyama 2018). Compared to diploid-
expressed loci, traits that rely on haploid gene function are expected to experience both
stronger purifying selection against deleterious alleles, and elevated fixation rates (stronger
positive selection) for advantageous alleles, because both deleterious and advantageous alleles
will be visible to haploid selection regardless of dominance (Charlesworth and Charlesworth
1992, Walsh and Charlesworth 1992, Immler and Otto 2018, Immler 2019). Moreover,
gametophytic-exclusive genes could also evolve differently from loci that are expressed in both
gametophytic and sporophytic (diploid) tissue, if the latter also experience additional
constraints, or even antagonistic effects, because of their expression during both phases of the
life cycle (Immler and Otto 2018).
104
In addition to haploid selection, plants also often exhibit substantial variation in mating system,
including among closely related species. This variation is predicted to have diverse effects on
the nature, timing, and strength of selection, acting both on reproductive genes specifically,
and across the genome more generally. In the specific case of reproductive genes, because the
mating system affects the number of reproductive partners an individual experiences, it can
directly influence the intensity of intrasexual competition, the opportunity for mate choice, and
the magnitude of conflict over reproductive decisions (Delph and Havens 1998; Skogsmyr and
Lankinen 2002; Brandvain and Haig 2005, Clark et al. 2006, Mazer et al. 2010). Selection on
traits affecting sexual competition and/or sexual conflict will be greatest in obligately
outcrossing individuals, but weak or absent in obligately self-fertilizing species where the
reproductive interests of both parents align perfectly (because they are the same individual).
Even intermediate rates of self-fertilization can diminish the strength of sexual selection and
thereby reduce, for example, the advantage of pollen competition traits (e.g. Mazer et al. 2018)
and the spread of superior pollen-expressed genes (e.g. Peters and Weis 2018). Mating
system—specifically variation in the frequency of selfing—is also proposed to have more global
effects on the efficacy of selection across the genome, regardless of whether genes have
reproductive functions. Because selfing reduces the effective population size (Ne), the overall
efficacy of selection—both against deleterious alleles, and in favor of advantageous alleles—is
expected to be reduced in selfing compared to outcrossing lineages (Charlesworth et al. 1993,
Charlesworth and Wright 2001, Mattila et al. 2018). Conversely, because selfing increases
homozygosity, it can also reduce the genetic load in a population—by increasing the exposure
of strongly deleterious recessive alleles to selection—as well as increase the early efficacy of
selection on recessive advantageous alleles by exposing them more rapidly to positive selection
(Mattila et al. 2018). Together with specific effects on reproductive genes, these complex and
sometimes antagonistic genome-wide effects might contribute to the heterogeneous
inferences that have emerged from studies of reproductive protein evolution in plants to date.
131
In this study, we examined genome-wide patterns and rates of protein evolution among four
closely-related species from the diverse plant genus Solanum. Our aims were to compare

patterns of molecular evolution in loci expressed in different classes of tissue in order to
evaluate whether: a) reproductive-specific proteins evolve more rapidly, on average, than non-
reproductive proteins; b) 'male'-specific loci evolve at different rates than 'female'-specific loci;
c) genes expressed exclusively in gametophytic tissues evolve differently from genes expressed
in sporophytic (diploid) tissue or in both tissue types; and d) mating system variation (a
potential proxy for the expected strength of sexual selection and/or sexual conflict) affects
patterns of protein evolution. In addition, to help assess underlying causes of detected
patterns, we also evaluated the influence of the magnitude and breadth of gene expression on
observed rates of protein evolution. Our goals were to understand the forces shaping the
molecular evolution of reproductive proteins in this plant group, and whether this differs from
evolutionary patterns in other protein types and/or those proposed to shape reproductive
protein evolution in animal systems.

146
147

2   Materials and Methods

2.1   Study Species

Our analyses used gene expression and sequence data from each of four species within the wild
tomato group (Solanum section Lycopersicum): *S. lycopersicum* (domesticated tomato; LA3475)
and three wild relatives—*S. pimpinellifolium* (LA1589), *S. pennellii* (LA0716), and *S. habrochaites*
(LA1777). (LA#### refers to the specific germplasm accession ID, as described in
tgrc.ucdavis.edu, from which all or most of the data were obtained; Table S1.) This group is a
clade of 12 closely-related diploid wild species along with the domesticated tomato, within the
hyperdiverse (~1300 species) plant genus *Solanum* (Sarkinen et al. 2013). All 12 wild tomato
species arose within the last ~2.5 million years, consistent with recent rapid speciation in this
clade (Pease et al. 2016a).

Mating system varies substantially across species in *Solanum* (Goldberg et al. 2010) and our
four focal species differ in their mating system, most notably in whether species are historically
self-incompatible (SI: *S. pennellii* and *S. habrochaites*) versus self-compatible (SC: *S.
lycopersicum* and *S. pimpinellifolium*). These species also have large differences in their genetic
effective population sizes (Ne)—for example, as estimated from average nucleotide
polymorphism (Pease et al. 2016a)—that are consistent with these SI vs SC mating system
differences. In *Solanum*, genetic self-incompatibility (the inferred ancestral state) is based on
molecular interactions between proteins expressed in growing pollen tubes and pistil-expressed
SRNase and other known proteins (Cruz-Garcia et al. 2003; Takayama and Isogai 2005). SI is
observed as arrest of pollen tube growth in the female reproductive tract (the pistil). Self-
compatible lineages have lost function in one or more SI-mediating proteins (Stone 2002),
thereby allowing individuals to be fertilized by their own pollen, in addition to outcrossing. As
with other angiosperms (Sicard and Lenhard 2011), transitions to SC within *Solanum* are often
followed by other morphological changes in floral size and structure, that further increase the
frequency of self-pollination and reduce the number/diversity of mating partners (Rick 1979). In
our four focal species, SI species have larger flowers, longer reproductive tracts, and more

pollen production on average, in comparison to SC species (Vosters et al. 2014), even though
the transition to SC likely occurred within the last ~0.5MY (the estimated age of the clade
containing the two SC species here; Pease et al. 2016a). Both differences in the capacity to
reject self pollen (i.e. SI versus SC), and these morphological differences, could have large
effects on the average diversity of mating partners, and therefore the opportunity for and
magnitude of intraspecific sexual selection and sexual conflict (Willson 1994, Delph and Havens
1998, Clark et al. 2006).
185
186 **2.2   Quantifying gene expression and determining tissue-specificity**
187
Our analyses used gene expression (RNAseq) data collected individually from leaf (up to five
developmental stages), root, seed, vegetative meristem, stem, flower (several stages), style,
pollen, and ovule tissues in each accession. RNAseq data were obtained from seven publically
available SRA projects (Table S1), mostly drawn from three previously published analyses
(Koenig et al. 2013, Ichihashi et al. 2014, Pease et al. 2016b), except data from ovule tissue
which has not been previously published. Note that style and pollen data were not available
specifically for *S. pimpinellifolium*. Sources of each dataset are described in detail in Table S1.
Procedures for generating the ovule RNAseq data are described in the supplementary text. Each
tissue had 1-6 replicates (generally 3) per accession (Table S1).
197
For each library within each tissue, we processed the raw reads by filtering adapter sequences
or low quality bases with TRIMMOMATIC (Bolger et al. 2014). Reads were then mapped against
tomato reference genome (ITAG 2.4) using STAR (Dobin et al. 2013). Numbers of reads mapped
onto genic regions were estimated with FEATURECOUNTS (Liao et al. 2014). We normalized the
read counts from each library by TPM (transcripts per million) and then calculated the mean
normalized read counts for each tissue of each species.
204
We first classified loci into three general classes of gene—reproductive (RP), vegetative (VG),
and general (GR) (i.e., expressed in both reproductive and non-reproductive/vegetative tissue
types). Reproductive genes were required to be expressed (TPM>2) in at least one
reproductive tissue (i.e. style, pollen, ovule), and to have no or trace expression (TPM <0.5) in
any of the remaining tissues (i.e. leaf, root, stem, seed, root, vegetative meristem), in at least
two of the three investigated species for which we had all tissues available (i.e. *S. lycopersicum*,
S. pennellii, and *S. habrochaites*). (Because *S. pimpinellifolium* lacked some of the tissue types in
our RNA-seq data, we did not include this species in the filtering/determination of tissue-
specificity.) Vegetative genes were defined similarly by requiring expression in at least one
vegetative (non-reproductive) tissue, and no/trace expression in any reproductive tissue.
Generally-expressed (GR) genes were expressed in at least one reproductive tissue and also in
at least one vegetative tissue.
217
Using similar criteria, we also identified a set of tissue-specific genes, defined as those only
expressed in one specific tissue (TPM>2) and not in any other tissue (TPM<0.5) in at least two
out of the three species for which we had all tissue types (i.e. *S. lycopersicum*, *S. pennellii*, and
S. habrochaites). For these analyses, we focused on specific tissues (i.e. leaf, root, stem, seed,

vegetative meristem, style, pollen, ovule) for which we had sufficient sampling for at least three
species (Table S1). Finally, using the same criteria and the expression domains in the three
species for which we had all tissues represented, we also classified transcripts as gametophytic-
exclusive (pollen and/or ovule expression only) versus sporophytic-exclusive versus both
gametophytic and sporophytic (pollen and/or ovule expression, and at least one other diploid
tissue), as well as ovule-specific versus expressed in both ovules and at least one sporophytic
(diploid) tissue, and pollen-specific versus expressed in both pollen and at least one sporophytic
(diploid) tissue.
230
231 **2.3 Estimating rates of protein evolution**
232
As with prior studies (Clark et al. 2006 and references therein), we used dN/dS—the ratio of the
number of non-synonymous substitutions per non-synonymous site to the number of
synonymous substitutions per synonymous site—as our estimate of protein evolution. Although
dN/dS is usually <1—because most functional loci experience some constraint on non-
synonymous changes—larger values of dN/dS are consistent with faster protein evolution. We
estimated dN/dS for each locus in our dataset, to calculate mean rates of protein evolution in
different classes of genes, and to identify the set of loci with estimated dN/dS>1—a pattern
consistent with positive selection for protein evolution at that locus. In addition, to compare
the overall dN/dS for loci on SI versus SC branches, we compared estimates from sequences
that were concatenated across groups of genes (described below).
243
To generate the sequence alignments for these analyses, we combined the RNA-seq data used
for quantifying gene expression (above) with whole-transcriptome data from the same four
accessions, previously generated in Pease et al. (2016a) or publically available (Table S2); the
latter dataset allowed us to augment our RNAseq data with additional sequences for loci whose
expression was not detected for one or more of our four species. These reads were aligned
against the tomato reference genome by STAR. The aligned data were processed using
SAMTOOLS followed by MVFTOOLS to generate the sequence alignments of orthologous genes. We
only retained transcript sequence alignments longer than 200 bps after removing gaps and
indels, leaving us with 21,216 sequence alignments (loci) for downstream analyses.
253
For each locus, we used PAML (Yang 2007) to estimate dN/dS across the branches of a given
phylogenetic topology among the four investigated species, using the input species tree
((lycopersicum, pimpinellifolium), (pennellii, habrochaites)). To do so we used the one-ratio
(M0) codeml model in PAML, that generates a single dN/dS estimate across all branches for each
locus (that is, dN/dS is fixed at one ratio for each gene). Outputs from this model were used to
compare the mean and distribution of dN/dS for genes among three broad categories
(reproductive (RP), vegetative (VE), and general (GR)), as well as between each of six tissue-
specific groups classified by their expression in only a single tissue (pollen, style, ovule, leaf,
root, or stem; vegetative meristem and seed were excluded from the latter analyses because
they had no exclusively tissue-specific genes), and between loci that had gametophytic versus
sporophytic, or gametophytic+sporophytic, domains of gene expression. We also compared loci

expressed in ovules only versus expressed in ovules and at least one sporophytic (diploid)
tissue, or in pollen only versus expressed in pollen and at least one sporophytic tissue.
267
Second, in addition to locus-by-locus dN/dS comparisons, we also evaluated overall dN/dS
estimates along SC branches (those leading to *S. lycopersicum* and *S. pimpinellifolium*) in
comparison to those along SI branches (those leading to *S. pennellii* and *S. habrochaites*).
Because the four species in our analysis are very closely related, the branch lengths for each
class of lineage (especially for SC lineages) were too short to reliably estimate lineage-specific
dN/dS for each locus individually. Therefore, for these analyses we concatenated all loci within
each class of genes to be compared, and then estimated dN/dS across the concatenated set of
loci for SI and SC branches separately. For example, for reproductive (RP) loci, we concatenated
all RP loci and estimated dN/dS across the entire concatenated set, allowing for different dN/dS
estimates (variable rates) on SI versus SC branches. Because of the small numbers of loci
concatenated for some of our individual tissues, we limited SI vs SC branch-specific estimates to
our three broad classes of loci (RP, VG, and GR) only.
280
Because the phylogeny of our species contains one internal branch, for these analyses we used
two alternative multi-rate (M2) codeml models in PAML. For the first, three-ratio, model we
allowed the internal branch to have a rate that differed from the tip branches leading either to
SC species, or to SI species. For the second, two-ratio, model, we classified the internal branch
as 'SI' (i.e. dN/dS was estimated separately for tip branches leading to SC species, versus a
background rate for the tip branches leading to SI species plus the internal branch). This
approach was used to estimate branch-specific rates (dN/dS estimates) for SI versus SC
branches for each of our three general classes of loci (RP, VG, and GR). Note we report three-
ratio results in the main text, and two-ratio results in the supplement.
290
291 **2.4   Comparing protein evolution rates**
292
293 All statistical analyses were performed in R version 3.6.3 (R Core Team 2015). For all
294 comparisons, we first removed genes showing unusually high values (dN/dS >10) from the PAML
295 output, as these were due to estimation limitations within PAML (e.g. a lack of synonymous
296 substitutions across the branch(es) used for estimation) or otherwise likely resulted from poor
297 sequence alignment.
298
To compare the mean rate of per locus protein evolution (dN/dS) between different classes of
loci, we fit generalized linear models (GLMs). Shapiro-Wilk tests of normality and quantile-
quantile plots were used to assess normality in the distribution of dN/dS; in all categories,
dN/dS was heavily skewed towards lower values and zero-inflated (reflecting the fact that most
expressed genes are under purifying selection). To accommodate this in our models, we assume
a gamma residual distribution with an identity link function, rather than the standard Gaussian
assumption, after confirming the suitability of this distribution with maximum likelihood (using
the fitdistrplus R package). Following GLMs, to compare the mean estimated dN/dS between
specific classes of loci, we made pairwise comparisons using Tukey post-hoc tests, adjusting p-
values for multiple comparisons. In addition, we compared the proportion of positively selected

genes (i.e. dN/dS>1) between different classes of loci by performing Chi-Squared tests of
independence and, for specific tests of enrichment in single categories (e.g., in RP relative to
GR), one-sided Fisher's exact tests.
312
To assess evidence for differences in dN/dS between mating systems, we aimed to compare
dN/dS estimates from concatenated data, as calculated separately for SI and SC branches. Both
biological and technical factors make these comparisons challenging to interpret. Biologically,
overall differences in Ne and therefore in the expected efficacy of positive and purifying
selection, could generate global differences in rates of protein evolution between SC and SI
lineages (see Introduction), that are unrelated to their potential differences in the historical
strength of sexual selection. Technically, in our specific dataset, the two SC species are much
more closely related than the two SI species, so there are large differences in branch lengths
(number of substitutions) over which we are estimating dN/dS values for SI versus SC branches.
Therefore, to assess whether exclusively reproductive (RP) loci had a different pattern of
molecular evolution on SI versus SC branches in comparison to other loci, we compared the
observed difference in dN/dS between SI and SC branches for RP genes to an estimate of the
'baseline' genome-wide difference in dN/dS between SI and SC branches. This baseline was
described by the SI vs SC difference in our set of generally expressed genes (GR), which is
expected to be due to factors unrelated to variation in sexual selection. Because we have only a
single estimate (from the concatenated data) for each class of gene for each branch type,
standard parametric tests can't be used. Instead we used bootstrap resampling to estimate the
variance around the mean SI vs SC difference in dN/dS in the GR dataset, and then to evaluate
whether the observed SI vs SC difference in RP genes falls outside the range of this variance (i.e.
is larger or smaller than the 95% CI of this distribution). For each bootstrap replicate (N=1000
replicates), we re-sampled 500 loci at random from the GR dataset, estimated dN/dS for SI and
SC branches and the difference between these two estimates. Our observed dN/dS difference
(SC-SI rate) was compared to the resulting distribution of 1000 replicate dN/dS differences from
GR loci.
337
338   **2.5   Assessing the effect of gene expression level on molecular evolutionary rates**
339
Numerous studies have shown that rates of protein evolution can be influenced by the
magnitude and breadth of gene expression, such that highly expressed genes and/or genes
expressed in a broader number of tissues tend to have systematically lower estimated dN/dS
(Meisel 2011). To assess the effect of the magnitude and breadth of gene expression on our
inferences, we also compared average dN/dS values between genes expressed in different
numbers of tissues (ranging from 1 (tissue-specific loci) through to 8 (expressed in all tissues
examined)), using the criteria that a gene be expressed >2 TPM in a specific tissue in all species
for which we had data on that tissue-type. Within each of the broad classes of RP, VG, and GR
loci, we also assessed the quantitative relationship between mean gene expression level
(averaged across all tissues in which a locus was expressed) and estimated dN/dS. For this
analysis we constructed a multiple regression model of dN/dS (square root transformed to
approximate normality), including both mean gene expression (TPM), tissue class (RP, VG, or
GR), and their interaction term as independent variables.

2.6   Assessing enrichment of functional categories among adaptively evolving loci

To test for functional enrichment among our loci with dN/dS > 1, we performed a Gene

Ontology (GO) term enrichment analysis using Panther version 14 (Mi et al., 2019; Ashburner et

al., 2000; The Gene Ontology Consortium, 2019). We performed separate tests for GR, VE, and

RP tissue categories, as these classes had large sample sizes appropriate for tests of

enrichment. Loci were evaluated using both the biological process and molecular function

annotation data sets, and significance was determined using Fisher's Exact tests with false

discovery rate correction.

3   Results

**3.1   Reproductive genes evolve at modestly faster rates, and have proportionally more
genes under positive selection**

Reproductive (RP) genes had an elevated mean rate of protein evolution (0.307±0.014)

compared to loci expressed in both reproductive and non-reproductive (general/GR) tissues

(0.271±0.003) (Posthoc Z=2.548, p = 0.0108) (Figure 1; Table 1; Table S3, Table S4). The mean

dN/dS for vegetative-specific (VG) loci (0.278±0.011) fell between these two other classes of

genes and did not differ from either of them (Table 1). In addition, the proportion of genes

with dN/dS>1 differed by group ($X^2(2)$= 7.9696, p=0.0186). A significantly greater proportion of

reproductive loci (28 of 670, or ~4.17%) are inferred to be evolving adaptively compared to

general loci (2.45%) (Fisher's exact test, p=0.0067; Table 1, Table S4). The proportion of

vegetative loci evolving adaptively (2.81%) was marginally lower than reproductive loci (Fisher's

exact test, p=0.093), and did not differ from general loci (Table 1; Table 4).

3.2   Female- rather than male-specific genes evolve the most rapidly

Comparisons among tissue-specific genes indicated no evidence that pollen-specific loci evolve

more rapidly than loci from female-derived tissues (style and ovule) (Figure 2; Table 2: Table

S5). Instead, style- and ovule-specific loci had the highest mean dN/dS values of all tissue

groups (0.67±0.17 and 0.34±0.08 respectively), and posthoc pairwise tests indicated that style

mean rate was significantly greater than rates in root-specific and leaf-specific loci, and ovule

mean rate was significantly greater than leaf mean rate (Figure 2; Table 2; Table S6). In

comparison, pollen loci had a mean rate (0.25±0.10) that was intermediate between these

groups of tissue-specific genes, and statistically indistinguishable from each other tissue (Table

S6). Note that many fewer loci were available for tissue-specific comparisons, especially leaf-

and stem-exclusive loci (Table 2), and two tissues—vegetative meristem and seed—had no

detected tissue-exclusive loci.

Similar to comparisons of mean rates, the proportion of genes with dN/dS>1 differed by tissue

($X^2(5)$=33.707, p=2.723 x 10^{-6}). Moreover, this proportion was significantly higher in style

tissue—where 6 of 25 loci were inferred to be evolving adaptively—compared to all other
tissues (Fisher's exact test, p= 5.046 x 10^{-5}; Table 2). For ovule-exclusive loci, the proportion
adaptive evolving loci (3 of 71) was also greater than in all other tissues (excluding style-specific
loci) (Fisher's exact test, p= 0.0376) (Table 2).
401
402 **3.3   Gametophytic-exclusive proteins do not evolve more rapidly than sporophytic-**
403 **exclusive proteins, or proteins expressed in both tissue types**
404
Although the mean rate of protein evolution in gametophytic genes (0.33±0.04) was slightly
higher than the rate of protein evolution in sporophytic-exclusive loci (0.30±0.06), this
difference was not significant (t=-0.452, P=0.588) (Table 3; Table S7). Gametophytic-exclusive
proteins also do not evolve faster than those expressed in at least one gametophytic and one
sporophytic tissue (0.27±0.03) (t=-1.498, P=0.134). Similar to mean rate comparisons, the
proportion of gametophytic loci with dN/dS>1 (3 of 86, or 3.49%) did not differ from this
proportion in sporophytic-exclusive genes (68 of 2033, or 3.34%) (Fisher's exact test, p=0.557),
and was slightly but not statistically larger than this proportion in gametophytic+sporophytic
genes (383 of 15778, or 2.43%) (Fisher's exact test, p=0.283).
414
As observed in the general gametophytic comparisons, mean rate of protein evolution in ovule-
exclusive loci (0.34±0.46) was slightly higher than the rate of protein evolution in loci expressed
in ovules and at least one sporophytic tissue (0.29±0.32), but this difference was not significant
(GLM t=1.212, p=0.225; Table S5). The proportion of ovule-exclusive loci with dN/dS>1 (3 of 71,
or 4.23%) also did not differ from the proportion of ovule+sporophytic loci with dN/dS>1 (225
of 7884, or 2.85%) (Fisher's exact test, p=0.854). Neither the mean rate of protein evolution
(GLM t=-0.037, p=0.971; Table S7) nor the proportion of loci with dN/dS>1 (Fisher's exact test,
p=0.832) differed between pollen-exclusive loci and loci expressed in pollen and at least one
sporophytic tissue (Table 3).
424
425 **3.4   Patterns of protein evolution show mating-system effects that are not specific to**
426 **reproductive loci**
427
Consistent with genome-wide effects of lower Ne in our self-compatible lineages, rates of
protein evolution were inferred to be higher overall on self-compatible branches in both GR
and RP classes of loci, although not for VG loci (Table 4); in the latter case, VG rates were
numerically higher on SI branches. Results were similar for two-ratio models (Table S8). The
observed difference in rates between SC and SI branch for RP loci (0.0338) did not fall outside
the 95% CI of bootstrapped values (0.0007-0.0797) from the broader class of GR genes; instead
33% of bootstrapped samples had a dN/dS difference smaller than observed for RP loci.
435
436 **3.5   Higher rates of protein evolution are consistently associated with greater tissue-**
437 **specificity**
438
Across loci classified according to the number of tissues in which they were expressed at >2
TPM (from 1 to 8 tissues), mean dN/dS was highest in loci expressed in a single tissue and

consistently declined across loci expressed in increasingly more tissues (Figure 3; Table 5; Table
S9, S10). Similarly, both mean expression level and tissue class (GR, RP, VG) affected the per
locus estimates of protein evolution (dN/dS), as did their interaction (Table 6), indicating highly
expressed loci generally have lower average rates of protein evolution but also that this
relationship varies depending upon the broad tissue class. The relationship between dN/dS and
mean expression level was negative for generally-expressed (GR) loci, however the 95% CI for
this slope overlapped zero for both RP and VG loci, indicating weaker relationships for these
less broadly expressed loci (Figure S1; Table S11). Classes of genes also differed in their overall
mean gene expression, which was twofold higher in RP genes (100.5±12.7) compared to GR
genes (50.1 ±1.1), and fivefold higher compared to VG loci (20.3±2.6).

These findings are generally consistent with greater evolutionary constraint on loci with
broader expression domains and, at least for broadly expressed (general) genes, greater
magnitudes of gene expression.

3.6   Adaptively evolving reproductive proteins are not enriched for specific functional
categories, but do include roles in sexual interactions

We found no evidence for enrichment of functional gene ontology categories among loci for
which we found dN/dS>1, including among reproductive loci inferred to be adaptively evolving
(data not shown). Nonetheless, among these reproductive loci, we did identify several genes
with clear functional roles in sexual interactions (Table S12).

4   Discussion

Reproductive proteins appear to evolve rapidly in many animal groups. Here we evaluated
evidence for elevated reproductive protein evolution in four closely-related flowering plant
species, along with the potential influence of three factors on observed patterns: variation in
mating system (and therefore the possible strength of sexual selection), in gametophytic gene
expression (and therefore the strength of haploid selection), and in the breadth and/or
magnitude of gene expression. We found evidence for elevated rates of reproductive protein
evolution, globally across the genome and in some specific reproductive tissues. Moreover,
these elevated rates appear to be more characteristic of female- rather than male-specific loci.
Among the three broad factors that might influence these patterns, we found that more rapid
evolution was consistently associated with more narrow domains of gene expression, but not
with expression in haploid versus diploid tissues; evolutionary rate differences between
lineages with different mating systems (self-compatible versus self-incompatible) were more
complex and not associated with specific shifts in reproductive protein evolution. Here we
discuss these findings in light of previously observed patterns of elevated reproductive protein
evolution, and the factors proposed to drive these patterns.

4.1   Female reproductive proteins tend to evolve more rapidly

Our evidence for elevated rates of reproductive protein evolution was observed both as a
modest but significant increase in mean dN/dS across all reproductive loci, and as a significantly
larger proportion of loci showing evidence of adaptive divergence between species (dN/dS>1).
Interestingly, however, this elevated evolution appears to be more clearly associated with
proteins that have female- rather than male-specific functions. Indeed, female reproductive loci
predominate among our most rapidly evolving loci, both generally and in analyses of tissue-
specific loci. Of the tissue-specific loci found to have dN/dS>1, all are expressed in a female-
specific tissue (either style or ovule).
493
This finding is intriguing. Some previous studies have detected accelerated evolution in proteins
with specific female-associated functions, for example, egg proteins involved in sperm-egg
interactions (e.g. Galindo et al. 2003). Nonetheless, most studies that include both male and
female proteins have found mixed evidence for elevated female-biased or female-specific
evolution (e.g. Wong 2010, Haerty et al. 2007; including in plants, e.g. Gossman et al. 2016),
compared to much clearer accelerated evolution in male-specific or male-biased proteins
(reviewed in Wong 2011, Wilburn and Swanson 2016). Some of this male-female difference
could simply reflect a comparative lack of knowledge about female-associated proteins, so that
these might be systematically underexamined in many comparisons. Indeed, most prior studies
in animals (reviewed, for example, in Clark et al. 2006, Dapper and Wade 2020), as well as some
in plants (Arunkumar et al. 2013, Harrison et al. 2019, and references therein), have focused
largely or entirely on reproductive proteins with male-specific functions. Here, our loci were
defined by tissue-specific patterns of gene expression, rather than from *a priori* expectations of
function, which might have helped to ameliorate any such biases.
508
Our data also indicates a role for adaptive evolution in this accelerated female protein
divergence. Elevated rates of non-synonymous change (but values of dN/dS still less than 1)
could result from relaxed purifying selection (see further below) or because positive selection
has only acted on a subset of sites within each gene, or a combination of these effects.
However, we also found a larger proportion of female (style and ovule) loci with dN/dS>1,
suggesting a concentration of adaptive female protein evolution specifically involving post-
pollination reproductive processes. One driver of this could be sexual interactions. Unlike in
many animals, where female choice can be exerted both before and after mating (the latter as
'cryptic female choice') (Andersson and Simmons 2006), in angiosperms male-female (pollen-
pistil) interactions can only act after pollen arrives at a flower. Therefore all evolutionary
dynamics involving female choice or male-female interactions must act upon loci expressed in
this relatively narrow post-mating window. Indeed, among our rapidly evolving (dN/dS>1) style
loci, we detect at least one protein with putative functions in pollen interactions (a pistil-
specific extension-like protein; Solyc02g078080), in addition to several biosynthetic enzymes
(Table S12). (In contrast, 2/3 rapid ovule-specific genes have potential functions in stress
response, possibly suggesting roles that are not directly related to sexual interactions—see
below.) Moreover, in other angiosperms, 'female' loci that show rapid adaptive divergence are
also mostly pistil-specific proteins involved in selection among different pollen genotypes (via
roles in genetic self-incompatibility) (Clark et al. 2006), again suggesting the importance of post-
pollination sexual dynamics in accelerated reproductive protein evolution.

529
Whether this potential adaptive explanation extends to explain the general elevation of female
reproductive protein evolution (Figure 2) or reproductive proteins overall (Figure 1) is not yet
clear. One prior study that compared protein evolution in male, female, and non-reproductive
tissues in *Arabidopsis thaliana* and two relatives, also found that mean dN/dS was elevated in
loci with expression biased towards female tissues (egg cell, central cell, and synergid cells:
three haploid tissues within the female gametophyte) but not in pollen-biased proteins
(Gossman et al. 2014). This pattern was attributed to relaxed selection on female-biased loci,
partly because additional site-specific tests did not consistently detect evidence of positive
selection in these loci (Gossman et al. 2014). Because our four species are very closely related,
we do not have sufficient power to assess site-specific evidence for adaptive evolution in loci
whose estimated dN/dS is less than 1. Future analyses that pair divergence-based estimates
with polymorphism data within species—that can provide alternative and often more powerful
tests of positive and relaxed selection (e.g. Gossman et al. 2016, Arunkumar et al. 2013)—will
be helpful in differentiating the relative influence of these different forms of selection on the
broader patterns of elevated dN/dS detected here.
545
Regardless, if elevated female protein evolution is at least partly driven by post-pollination
sexual interactions, it is interesting that we do not also observe elevated evolution in pollen-
acting proteins. Of prior analyses in angiosperms, several have detected broadly elevated
evolution in pollen proteins (e.g., Arunkumar et al. 2013, Gossman et al. 2016, Harrison et al.
2019). Pollen protein evolution could be more constrained than style or ovule evolution due to
factors like differences in the degree of tissue-specificity or in the operation of haploid
selection, but these explanations are not generally consistent with our data (see further below).
Alternatively, it's possible that our analysis did not capture all the male proteins relevant to
post-pollination sexual interactions. For instance, pollen loci whose expression requires direct
interactions with the style and/or other pollen tubes would not be captured in our expression
atlas (which was generated from RNA expression in mature ungerminated and *in vitro*
germinated pollen; Table S1). There is evidence in other angiosperms that pollen
transcriptomes can be actively modified during their transit through the style, resulting in
changes in the suite of proteins being expressed (e.g., Qin et al. 2009, Mizukami et al. 2019).
Accounting for and including these genes would require transcriptomes from 'mixed' post-
pollination tissue, as well as specific approaches to differentiate pollen- from stylar-expression
in these mixed samples (e.g. Pease et al. 2016b). Importantly, this limitation would also extend
to any style- or ovule-specific proteins that are only elicited during pollination, so may not be a
complete explanation for the general difference we detect between male and female loci here.
565
Other explanations for the under-accounting of pollen loci could include technical effects such
as detection limits on gene expression in pollen, and/or exclusion of pollen-expressed loci that
are especially short or rapidly evolving, due to filtering during transcript-mapping steps. In
addition, by using dN/dS comparisons only, we would not have detected protein changes that
involve duplicate, chimeric, or novel genes, and/or if different loci are the targets of selection in
different lineages. If these latter factors do contribute to our failure to detect elevated pollen
protein evolution here, it suggests that different dynamics shape pollen protein identity and/or

evolution in *Solanum* compared to other angiosperms that have shown evidence for elevated
evolution using metrics like dN/dS (e.g. Arunkumar et al. 2013, Gossman et al. 2016, Harrison
et al. 2019). Regardless, with the data that we currently have, we do not see evidence that male
functions are subject to especially intense sexual selection, particularly in comparison to female
reproductive proteins.
578
579 **4.2   Rates of protein evolution are affected by tissue-specificity but not haploid expression**
580
581 Our results can also suggest factors that shape the patterns of reproductive protein evolution
582 that we do observe, at least among three broad factors (gene expression level, haploid
583 expression, and/or mating system variation) that we addressed here.
584
Of these, we find the strongest evidence that rates of protein evolution are influenced by the
breadth and magnitude of gene expression. Many studies have shown higher rates of protein
evolution in genes with more narrow expression domains and/or lower levels of gene
expression (Meisel 2011, and references therein), a relationship that is thought to reflect lower
constraint on tissue-specific proteins and stronger purifying selection on genes with very high
expression levels or broad expression domains. We similarly found that higher rates of protein
evolution were generally associated greater tissue-limitation and lower levels of gene
expression in our species. However, we infer that tissue-limitation (a more narrow expression
domain) is likely more important for shaping reproductive protein evolution specifically, as
lower mean gene expression was not significantly associated with higher protein evolution for
reproductive genes (Figure S1, Table 4), and reproductive loci also had much higher mean gene
expression levels in comparison to generally-expressed genes and, especially, vegetative loci.
597
High tissue-specificity has been proposed as an important contributor to elevated rates of
reproductive protein evolution—and a better predictor of this than sex-biased gene expression
per se—in both animals (Meisel 2011) and some plants (Veltsos 2019). For example, in
Arabidopsis thaliana and close relatives, Harrison et al. (2019) found faster protein evolution in
both pollen-specific and tissue-specific sporophytic genes compared to loci expressed in >1
tissue, although pollen-specific loci still retain the highest evolutionary rates in this and other
analyses in these species (e.g. Szovenyi et al. 2013, Gossman et al. 2016). Arunkumar et al.
(2013) also inferred a role for tissue-limitation or specificity in elevated dN/dS in *Capsella*
grandiflora pollen proteins (although this elevation was attributed to different forms of
selection—relaxed purifying selection in pollen germ cell genes versus greater purifying and
positive selection in pollen tube loci—depending on the specific sub-tissues within pollen).
These prior inferences, and our observations, suggest that higher tissue-specificity could
facilitate broadly elevated rates of protein evolution. Nonetheless, tissue-specificity cannot
provide a complete explanation of elevated reproductive protein evolution in our dataset, as
tissue-specific non-reproductive proteins still evolve more slowly than tissue-specific (female)
reproductive proteins (Table 4).
614
In comparison, we found no rate differences between loci expressed in haploid versus diploid
tissues, and therefore little evidence for a strong effect of haploid selection in driving

differences between reproductive and non-reproductive proteins. Haploid selection has two
projected effects on haploid-expressed loci—increasing the efficacy of positive selection on
advantageous alleles, and of purifying selection on deleterious alleles (Immler and Otto 2018,
Mattila et al. 2018)—compared to diploid-expressed loci. We have limited ability to
differentiate these two effects with our particular estimate of protein evolution, but our
observations are not consistent with a strong effect of either. In terms of accelerating adaptive
protein evolution, we found no significant elevation of evolutionary rates in loci limited to
gametophytic-tissues in general, and in ovule-limited and pollen-limited loci specifically (Table
3). Similarly, we find little evidence for stronger or more widespread purifying selection on
haploid-expressed loci, which would be observed as lower mean rates of protein evolution in
gametophytic loci. Stronger purifying selection due to haploid expression has been proposed as
one reason why sex-biased loci (generally, pollen expressed loci) do not appear to evolve faster
than non-sex-biased loci in several dioecious species (Sanderson et al. 2019, Cossard et al.
2019; discussed in Muyle 2019, but see also Veltsos 2019) and, indeed, sometimes appear to
evolve more slowly (e.g., Darolti et al. 2018). Our data do not suggest stronger purifying
selection in either gametophytic-limited proteins in general, or ovule- and pollen-limited loci
specifically, in our species. However, because we have relatively few pollen-specific loci
(discussed above) we cannot exclude the possibility that intermediate rates of pollen protein
evolution result from a tension between both positive and purifying selection predicted to act
on these loci. Nonetheless, our present data indicates little evidence for a global effect of
haploid gene expression on molecular evolutionary rates.
638
639   **4.3   Mating system variation does not does not differentially influence reproductive**
640   **protein evolution**
641
Our results also did not support a strong consistent effect of mating system differences—and
therefore predicted differences in the strength of sexual selection—on patterns of reproductive
protein evolution. Interpreting these comparisons is complex because differences in genetic
effective population size are expected to have global effects on rates of molecular evolution
(Charlesworth and Wright 2001, and see Introduction) that are unrelated to sexual selection *per*
se. The broad patterns of dN/dS we observed generally reflected these expected global effects
of differences in Ne due to mating system variation: in the SC lineages, mean dN/dS was higher
in both RP and GR genes, consistent with a general relaxation of selection in more selfing
lineages (Charlesworth and Wright 2001) regardless of whether or not loci were associated with
reproductive functions (Table 4). This genome-wide effect might have overwhelmed more
subtle effects of mating system variation acting specifically on loci involved in male-male
competition and/or female choice. In addition, the transition to self-compatibility is relatively
recent (less than 1 MY) among our species, and self-compatible lineages still retain significant
potential for outcrossing (Rick 1979, Vosters et al. 2014). These factors mean there may be
relatively modest differences in the historical and current opportunity for sexual selection
among our lineages, and therefore little opportunity to observe systematic consequences of
this transition as differential changes in evolutionary rates. (Statistically, this is compounded in
our particular dataset by the two SC species branches being substantially shorter than those for
our two SI species.) This contrasts with other angiosperms, such as *A. thaliana*, where the

transition to self-compatibility is about as old but the consequent effects on outcrossing rates
has been much more severe (selfing rates can be >95%; Harrison et al. 2019, and references
therein).
664
Interestingly, several previous analyses in animal systems have similarly found equivocal
evidence for differential effects of mating system variation on the evolution of reproductive
tract proteins (reviewed in Wong 2011). For example, in studies of primates, evolutionary rates
in specific sperm and copulatory plug proteins are positively associated with common proxies
for sexual selection, such as residual testis size and estimated number of mates (e.g. Dorus et
al. 2004, Martin-Coello et al. 2009). However other studies find that phylogeny-wide signals of
positive selection in reproductive tract proteins are not associated specifically with polyandrous
(as opposed to monogamous) lineages (e.g., Ramm et al. 2008, Finn and Civetta 2010; and see
Table 1 in Wong 2011). Limitations on phylogenetic models that can assess associations
between molecular evolutionary rates and changes in phenotypic characters, and variation
among lineages in the specific targets of sexual selection (Wong 2011), might explain relatively
weak effects of mating system variation on protein evolution rates, despite the importance of
sexual selection in driving faster reproductive protein evolution. Adding more species, and
focusing on pairs of SI-SC sister species, might help in addressing some of these limitations in
future analyses within *Solanum*.
680
Alternatively, the weak effect of mating system variation—both here and in prior studies—
might be because the predominant cause of globally elevated reproductive protein evolution is
not sexual selection. For example, pathogen resistance responses are also known to be
associated with elevated rates of protein evolution—driven by antagonistic host-pathogen
evolution—and might be concentrated among reproductive proteins if mating often leads to
the exchange of pathogens (Clark et al. 2006, Wong 2011). In principle, the identity of loci
inferred to be under positive selection could be helpful in differentiating among alternative
selective agents responsible for driving this adaptive divergence. Of the genes in our dataset
that show elevated evolution, we cannot yet draw strong conclusions about whether they
better reflect roles related to sexual interactions versus responses to non-sexual environmental
factors; many of them have unknown functions or general functions that do not *a priori* provide
specific support for either inference (Table S12). We do detect at least one style-specific protein
with a clear functional connection to pollen-pistil interactions, however our rapidly-evolving
ovule-specific genes are implicated in stress responses, which might indicate a greater role for
natural rather than sexual selection in shaping adaptive responses in this class of loci.
696
697   **4.4   Consequences of rapid protein evolution for the evolution of reproductive differences**
698   **and speciation**
699
Finally, among the motivations for examining patterns of reproductive protein evolution is to
understand their possible role in lineage diversification. If reproductive proteins do indeed
evolve more rapidly than other classes of protein, then perhaps they play an outsized role in
the emergence of functional differences and reproductive isolating barriers between lineages.
For example, evidence suggests post-mating prezygotic traits frequently evolve rapidly in

animals and, when divergent between species, could contribute to reproductive isolation
(Howard 1999, Snook et al. 2009, Palumbi 2009, Castillo and Moyle 2014). Similarly, divergence
in post-pollination prezygotic reproductive molecules could be important contributors to
species barrier formation among angiosperms (Moyle et al. 2014). Here we did detect evidence
for a global elevation of protein evolution in tissues that mediate these post-pollination
prezygotic stages, especially of female-specific loci. Our observations are consistent with a role
for tissue specificity in facilitating these elevated rates, but not for haploid selection or, as yet,
for differences in selective dynamics that are associated with recent shifts in mating system.
The evolutionary factors that might drive this more rapid evolution therefore remain to be
clarified. Regardless, our results indicate that rapid reproductive divergence is especially
characteristic of proteins with female-specific functions among our species. These specific
functions can be critical for maintaining coordinated sexual signals and ensuring fertilization
post-pollination (Bernasconi et al. 2001, Swanson et al. 2004, Moyle et al. 2014). Therefore
their rapid divergence might be particularly influential in the initial stages of reproductive
isolation and, thereby, in speciation, in this highly diverse plant genus.
720
721

722 **5   Author contributions**
723
724 LCM, MW, and MJSG designed the study. MW and MJSG performed the analyses. LCM wrote
725 the manuscript with assistance from MW and MJSG.
726
727

728 **6   Funding**
729
730 This work was supported by NSF grant IOS-1127059 and NSF grant DEB-1856469 to LCM.
731
732

733 **7   Acknowledgments**
734
735 Thanks to CJ Jewell for collecting ovule tissue, Christie Bergeon for RNA extractions, and the
736 Indiana University Center for Genomics and Bioinformatics for assistance with library
737 construction and sequencing.
738

739
740
741 **Figure 1:** Per locus protein evolution (dN/dS) among loci expressed in reproductive (RP),
742 vegetative (VG), and both reproductive and non-reproductive (general/GR) tissues in Solanum
743 species. Black circles indicate means for each group. Inset: Estimated mean (circle) and 95% CI
744 of dN/dS for each group of loci.
745

746
747
748 **Figure 2:** Per locus protein evolution (dN/dS) among groups of tissue-specific loci in Solanum
749 species. Black circles indicate means for each group. Inset: Estimated mean (circle) and 95% CI
750 of dN/dS for each group of loci.
751
752

753
754
755 **Figure 3:** Per locus protein evolution (dN/dS) among groups of loci expressed (>2 TPM) in
756 different numbers of tissues (from 1 to 8) in Solanum species. Black circles indicate means for
757 each group. Inset: Estimated mean (circle) and 95% CI of dN/dS for each group of loci.
758
759

760
761
762

763
764

765 **Supplementary Figure 1:** Relationship between per locus mean level of gene expression (TPM)
766 and per locus protein evolution (dN/dS) in loci expressed in (left) general/GR, (middle)
767 reproductive/RP, and (right) vegetative/VG tissues in Solanum species. Each point indicates an
768 individual locus. Regression lines shown in black.
769

Tables

Table 1. Rates of protein evolution (dN/dS) in different broad classes of loci. dN/dS calculated for each locus using a one ratio test in PAML.

		Per locus dN/dS				Loci with dN/dS>1		
Class	N loci	mean	SE	Median	Tukey test[1]	Number	Proportion2	Fisher's exact tests[1]
General	16221	0.2709	0.0026	0.20	a	399	0.0246	a
Vegetative	923	0.2783	0.0111	0.19	ab	26	0.0282	ab
Reproductive	670	0.3071	0.0142	0.21	b	28	0.0418	b

1. see Table S3.
2. The proportion of genes with dN/dS>1 differed among classes of loci (X^2=7.9696, p=0.0186).

Table 2: Rates of protein evolution (dN/dS) in tissue-specific loci

		Per locus dN/dS				Loci with dN/dS>1	
Group	N	Mean	SE	Median	Tukey test[1]	Number	Proportion[2]
Leaf	3	0.0978	0.0657	0.00	a	0	0
Stem	2	0.1925	0.1714	0.19	abc	0	0
Root	124	0.2103	0.0693	0.19	ab	0	0
Pollen	13	0.2521	0.1046	0.16	abc	0	0
Ovule	71	0.3418	0.0809	0.25	bc	3	0.042
Style	25	0.6668	0.1685	0.17	c	6	0.240

1. see Table S6.
2. The proportion of genes with dN/dS>1 differed by tissue (X^2= 33.707, p=2.72 x 10-6). One-sided Fisher's exact tests indicated this proportion was greater in style compared to all other tissues (p=5.046 x 10-5), and greater in ovule compared to all other tissues except style (p=0.03756).

788 **Table 3: Rates of protein evolution (dN/dS) in gametophytic (GM) versus sporophytic (SP) or**
789 **gametophytic+sporophytic (GM+SP) loci.** Within each GLM, tissue groups did not differ in
790 mean dN/dS (Table S7).

Group		dN/dS			Loci with dN/dS>1		
	N	mean	SE	Median	Number	Proportion	p[1]
a) Gametophytic vs Sporophytic							
Gametophytic only	86	0.3275	0.0576	0.24	3	0.0349	0.5574
Sporophytic only	2033	0.2958	0.0586	0.21	68	0.0334	
b) Gametophytic vs GM+SP							
Gametophytic only	86	0.3275	0.0393	0.24	3	0.0349	0.2833
Gametophytic + sporophytic	15778	0.2685	0.0394	0.20	383	0.0243	
c) Ovule loci							
Ovule only	71	0.3418	0.0036	0.25	3	0.0423	0.8542
Ovule + sporophytic	7884	0.2874	0.0448	0.21	225	0.0285	
d) Pollen loci							
Pollen only	13	0.2521	0.0679	0.16	0	0.0000	0.8316
Pollen + sporophytic	277	0.2496	0.0695	0.18	4	0.0144	

792
793 1. Fisher's exact tests
794
795
796 **Table 4: Estimates of dN/dS for SC and SI branches separately, for loci concatenated within**
797 **each broad class of genes.** Results are from 3 ratio model in PAML, with input species tree:
798 ((Slyc #1, Spim #1), (Spen #2, Shab #2)).
799

Class	# genes	# codons	Null model	Variable rate model	2*LnL	p value	SI-branch dN/dS	SC-branch dN/dS	Internal Branch	SC-S diff.
GR	16221	6619178	-28576265.18	-28575900.38	729.61282	3.686 10x-159	0.2224	0.2617	0.1901	0.039
RP	904	110171	-478937.9344	-478932.4222	11.024336	0.00404	0.2285	0.2623	0.1999	0.03
VE	1300	78355	-344551.0442	-344546.389	9.310318	0.00951	0.2267	0.2042	0.1850	-0.02

801 **Table 5: Rates of protein evolution (dN/dS) in loci according to breadth of tissue expression.**

# Tissues expressed in	N loci	Mean sqrt(dN/dS)	SE	Median	Posthoc Tukey test[1]
1	1892	0.3429	0.0088	0.24	d
2	1528	0.3080	0.0125	0.22	cd
3	901	0.3183	0.0148	0.23	cd
4	930	0.3053	0.0142	0.23	cd
5	1083	0.3022	0.0135	0.24	c
6	1507	0.2919	0.0122	0.21	c
7	7206	0.2498	0.0094	0.19	b
8	3173	0.2213	0.0098	0.16	a

804 1. see Table S10

807 **Table 6: Relationship between mean gene expression level and estimated dN/dS, across loci**
808 **in each broad class (GR, RP, VG) of loci (Linear model)**

	DF	SS	MS	F	Pr > F
Expression level	1	15.44	15.4355	243.85	2.2 x 10-16
Class	2	1	0.4987	7.88	0.00038
Expression level x Class	2	3.34	1.6689	26.36	3.72 x 10-12
	15086	954.94	0.0633		

810
811

8   References

ANDERSSON M, SIMMONS LW. 2006 Sexual selection and mate choice. Trends in Ecology & Evolution. 21(6):296-302.

ANDERSSON MB. 1994 *Sexual Selection*. Princeton University Press, Princeton, NJ.

ARUNKUMAR R, JOSEPHS EB, WILLIAMSON RJ, WRIGHT SI. 2013 Pollen-specific, but not sperm-specific, genes show stronger purifying selection and higher rates of positive selection than sporophytic genes in *Capsella grandiflora*. Mol Biol Evol. 30(11):2475-2486.

ASHBURNER M, BALL CA, BLAKE JA, BOTSTEIN D, BUTLER H, CHERRY JM, DAVIS AP, DOLINSKI K, DWIGHT SS, EPPIG JT, HARRIS MA, HILL DP, ISSEL-TARVER L, KASARSKIS A, LEWIS S, MATESE JC, RICHARDSON JE, RINGWALD M, RUBIN GM, SHERLOCK G, GENE ONTOLOGY C. 2000 Gene Ontology: tool for the unification of biology. Nature Genetics. 25(1):25-29.

BERNASCONI G, ASHMAN TL, BIRKHEAD TR, BISHOP JDD, GROSSNIKLAUS U, KUBLI E, MARSHALL DL, SCHMID B, SKOGSMYR I, SNOOK RR, TAYLOR D, TILL-BOTTRAUD I, WARD PI, ZEH DW, HELLRIEGEL B. 2004 Evolutionary ecology of the prezygotic stage. Science. 303(5660):971-975.

BOLGER AM, LOHSE M, USADEL B. 2014 Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics. 30(15):2114-2120.

BRANDVAIN Y, HAIG D. 2005 Divergent mating systems and parental conflict as a barrier to hybridization in flowering plants. American Naturalist. 166(3):330-338.

CASTILLO DM, MOYLE LC. 2014 Intraspecific sperm competition genes enforce post-mating species barriers in Drosophila. Proceedings of the Royal Society B-Biological Sciences. 281(1797).

CHARLESWORTH B, MORGAN MT, CHARLESWORTH D. 1993 The effect of deleterious mutations on neutral molecular variation Genetics. 134(4):1289-1303.

CHARLESWORTH D, CHARLESWORTH B. 1992 The effects of selection in the gametophytic stage on mutational load Evolution. 46(3):703-720.

CHARLESWORTH D, WRIGHT SI. 2001 Breeding systems and genome evolution. Current Opinion in Genetics & Development. 11(6):685-690.

CLARK NL, AAGAARD JE, SWANSON WJ. 2006 Evolution of reproductive proteins from animals and plants. Reproduction. 131(1):11-22.

CONSORTIUM TGO. 2019 The Gene Ontology Resource: 20 years and still GOing strong. Nucleic Acids Research. 47(D1):D330-D338.

COSSARD GG, TOUPS MA, PANNELL JR. 2019 Sexual dimorphism and rapid turnover in gene expression in pre-reproductive seedlings of a dioecious herb. Annals of Botany. 123(7):1119-1131.

CRUZ-GARCIA F, HANCOCK CN, MCCLURE B. 2003 S-RNase complexes and pollen rejection. Journal of Experimental Botany. 54(380):123-130.

DAPPER AL, WADE MJ. 2020 Relaxed Selection and the Rapid Evolution of Reproductive Genes. Trends in Genetics. 36(9):640-649.

DAROLTI I, WRIGHT AE, PUCHOLT P, BERLIN S, MANK JE. 2018 Slow evolution of sex-biased genes in the
reproductive tissue of the dioecious plant Salix viminalis. Molecular Ecology. 27(3):694-708.

DELPH LF, ASHMAN TL. 2006 Trait selection in flowering plants: how does sexual selection contribute?
Integrative and Comparative Biology. 46(4):465-472.

DELPH LF, HAVENS K. 1998 Pollen competition in flowering plants. In: *Sperm competition and sexual*
selection. (TR B, AP M, eds), pp. 149-173. Academic Press, San Diego, CA.

DOBIN A, DAVIS CA, SCHLESINGER F, DRENKOW J, ZALESKI C, JHA S, BATUT P, CHAISSON M,
GINGERAS TR. 2013 STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 29(1):15-21.

DORUS S, EVANS PD, WYCKOFF GJ, CHOI SS, LAHN BT. 2004 Rate of molecular evolution of the seminal
protein gene SEMG2 correlates with levels of female promiscuity. Nature Genetics. 36(12):1326-1329.

FINN S, CIVETTA A. 2010 Sexual Selection and the Molecular Evolution of ADAM Proteins. Journal of
Molecular Evolution. 71(3):231-240.

GALINDO BE, VACQUIER VD, SWANSON WJ. 2003 Positive selection in the egg receptor for abalone
sperm lysin. Proceedings of the National Academy of Sciences of the United States of America.
100(8):4639-4643.

GOLDBERG EE, KOHN JR, LANDE R, ROBERTSON KA, SMITH SA, IGIC B. 2010 Species selection maintains
self-incompatibility. Science. 330(6003):493-495.

GOSSMANN TI, SALEH D, SCHMID MW, SPENCE MA, SCHMID KJ. 2016 Transcriptomes of Plant
Gametophytes Have a Higher Proportion of Rapidly Evolving and Young Genes than Sporophytes.
Molecular Biology and Evolution. 33(7):1669-1678.

GOSSMANN TI, SCHMID MW, GROSSNIKLAUS U, SCHMID KJ. 2014 Selection-Driven Evolution of Sex-
Biased Genes Is Consistent with Sexual Selection in Arabidopsis thaliana. Molecular Biology and
Evolution. 31(3):574-583.

HAERTY W, JAGADEESHAN S, KULATHINAL RJ, WONG A, RAM KR, SIROT LK, LEVESQUE L, ARTIERI CG,
WOLFNER MF, CIVETTA A, SINGH RS. 2007 Evolution in the fast lane: Rapidly evolving sex-related genes
in drosophila. Genetics. 177(3):1321-1335.

HAFIDH S, FILA J, HONYS D. 2016 Male gametophyte development and function in angiosperms: a
general concept. Plant Reproduction. 29(1-2):31-51.

HARRISON MC, MALLON EB, TWELL D, HAMMOND RL. 2019 Deleterious Mutation Accumulation in
Arabidopsis thaliana Pollen Genes: A Role for a Recent Relaxation of Selection. Genome Biology and
Evolution. 11(7):1939-1951.

HOWARD DJ. 1999 Conspecific sperm and pollen precedence and speciation. Annual Review of
Ecology and Systematics. 30:109-132.

IMMLER S. 2019 Haploid Selection in "Diploid" Organisms. In: *Annual Review of Ecology, Evolution, and*
Systematics, Vol 50, Vol. 50: *Annual Review of Ecology Evolution and Systematics* (Futuyma DJ, ed), pp.
219-236.

IMMLER S, OTTO SP. 2018 The evolutionary consequences of selection at the haploid gametic stage.
American Naturalist. 192(2):241-249.

ICHIHASHI Y, AGUILAR-MARTINEZ JA, FARHI M, CHITWOOD DH, KUMAR R, MILLON LV, PENG J,
MALOOF JN, SINHA NR. 2014 Evolutionary developmental transcriptomics reveals a gene
network module regulating interspecific diversity in plant leaf shape. Proceedings of the
National Academy of Sciences of the United States of America. 111(25):E2616-E2621.

KOENIG D, JIMENEZ-GOMEZ JM, KIMURA S, FULOP D, CHITWOOD DH, HEADLAND LR, KUMAR R,
COVINGTON MF, DEVISETTY UK, TAT AV, TOHGE T, BOLGER A, SCHNEEBERGER K, OSSOWSKI S,
LANZ C, XIONG GY, TAYLOR-TEEPLES M, BRADY SM, PAULY M, WEIGEL D, USADEL B, FERNIE AR,
PENG J, SINHA NR, MALOOF JN. 2013 Comparative transcriptomics reveals patterns of selection
in domesticated and wild tomato. Proceedings of the National Academy of Sciences of the
United States of America. 110(28):E2655-E2662.

LANKINEN A, HYDBOM S, STRANDH M. 2017 Sexually antagonistic evolution caused by male-male
competition in the pistil. Evolution. 71(10):2359-2369.

LIAO Y, SMYTH GK, SHI W. 2014 featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics. 30(7):923-930.

LLOYD DG, WEBB CJ. 1977 Secondary sexual characters in plants Botanical Review. 43(2):177-216.

MARTIN-COELLO J, DOPAZO H, ARBIZA L, AUSIO J, ROLDAN ERS, GOMENDIO M. 2009 Sexual selection
drives weak positive selection in protamine genes and high promoter divergence, enhancing sperm
competitiveness. Proceedings of the Royal Society B-Biological Sciences. 276(1666):2427-2436.

MATTILA TM, LAENEN B, SLOTTE T. 2018 Population genomics of transitions to selfing in Brassicaceae
model systems. Arxiv. http://arxiv.org/abs/1809.04872.

MAZER SJ, HENDRICKSON BT, CHELLEW JP, KIM LJ, LIU JW, SHU J, SHARMA MV. 2018 Divergence in
pollen performance between Clarkia sister species with contrasting mating systems supports
predictions of sexual selection. Evolution. 72(3):453-472.

MAZER SJ, HOVE AA, MILLER BS, BARBET-MASSIN M. 2010 The joint evolution of mating system and
pollen performance: Predictions regarding male gametophytic evolution in selfers vs. outcrossers.
Perspectives in Plant Ecology Evolution and Systematics. 12(1):31-41.

MCCLURE B, CRUZ-GARCIA F, ROMERO C. 2011 Compatibility and incompatibility in S-RNase-based
systems. Annals of Botany. 108(4):647-658.

MEISEL RP. 2011 Towards a More Nuanced Understanding of the Relationship between Sex-Biased
Gene Expression and Rates of Protein-Coding Sequence Evolution. Molecular Biology and Evolution.
28(6):1893-1900.

MI H, HUANG X, MURUGANUJAN A, TANG H, MILLS C, KANG D, THOMAS PD. 2019 PANTHER version
14: more genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools. . Nucleic
Acids Research. 47(D1):D419-D426.

MIZUKAMI AG, INATSUGI R, JIAO J, KOTAKE T, KUWATA K, OOTANI K, OKUDA S, SANKARANARAYANAN
S, SATO Y, MARUYAMA D, IWAI H, GARENAUX E, SATO C, KITAJIMA K, TSUMURAYA Y, MORI H,
YAMAGUCHI J, ITAMI K, SASAKI N, HIGASHIYAMA T. 2016 The AMOR Arabinogalactan Sugar Chain
Induces Pollen-Tube Competency to Respond to Ovular Guidance. Current Biology. 26(8):1091-1097.

MIZUTA Y, HIGASHIYAMA T. 2018 Chemical signaling for pollen tube guidance at a glance. Journal of
Cell Science. 131(2).

MOORE JC, PANNELL JR. 2011 Sexual selection in plants. Current Biology. 21(5):R176-R182.

MOYLE LC, JEWELL CP, KOSTYUN JL. 2014 Fertile approaches to dissecting mechanisms of premating
and postmating prezygotic reproductive isolation. Current Opinion in Plant Biology. 18:16-23.

MUYLE A. 2019 How different is the evolution of sex-biased gene expression between plants and
animals? A commentary on: 'Sexual dimorphism and rapid turnover in gene expression in
prereproductive seedlings of a dioecious herb'. Annals of Botany. 123(7):IV-V.

PEASE JB, HAAK DC, HAHN MW, MOYLE LC. 2016A Phylogenomics reveals three sources of adaptive
variation during a rapid radiation. PLoS Biology. 14(2):e1002379.

PEASE JB, GUERRERO RF, SHERMAN NA, HAHN MW, MOYLE LC. 2016B Molecular mechanisms of
postmating prezygotic reproductive isolation uncovered by transcriptome analysis. Molecular Ecology.
25(11):2592-2608.

PETERS MAE, WEIS AE. 2018 Selection for pollen competitive ability in mixed-mating systems.
Evolution. 72(11):2513-2536.

QIN Y, LEYDON AR, MANZIELLO A, PANDEY R, MOUNT D, DENIC S, VASIC B, JOHNSON MA, PALANIVELU
R. 2009 Penetration of the Stigma and Style Elicits a Novel Transcriptome in Pollen Tubes, Pointing to
Genes Critical for Growth in a Pistil. Plos Genetics. 5(8).

R CORE TEAM. 2015. R: A language and environment for statistical computing. Vienna, Austria:
Sponsored by the R Foundation for Statistical Computing.

RAMM SA, OLIVER PL, PONTING CP, STOCKLEY P, EMES RD. 2008 Sexual selection and the adaptive
evolution of mammalian ejaculate proteins. Molecular Biology and Evolution. 25(1):207-219.

RICK CM. 1979 Biosystematic studies in Lycopersicon and closely related species of Solanum. In: *The*
biology and taxonomy of the Solanaceae. (Hawkes JG, Lester RN, Skelding AD, eds), pp. 667-679.
Academic Press Inc., New York.

RITCHIE MG. 2007 Sexual selection and speciation. Annual Review of Ecology, Evolution, and
Systematics. 38:79-102.

SANDERSON BJ, WANG L, TIFFIN P, WU ZQ, OLSON MS. 2019 Sex-biased gene expression in flowers,
but not leaves, reveals secondary sexual dimorphism in Populus balsamifera. New Phytologist.
221(1):527-539.

SARKINEN T, BOHS L, OLMSTEAD RG, KNAPP S. 2013 A phylogenetic framework for evolutionary study
of the nightshades (Solanaceae): a dated 1000-tip tree. Bmc Evolutionary Biology. 13.

SICARD A, LENHARD M. 2011 The selfing syndrome: a model for studying the genetic and evolutionary
basis of morphological adaptation in plants. Annals of Botany. 107(9):1433-1443.

SKOGSMYR I, LANKINEN A. 2002 Sexual selection: an evolutionary force in plants. Biological Reviews.
77(4):537-562.

SLOTTE T, BATAILLON T, HANSEN TT, ST ONGE KR, WRIGHT SI, SCHIERUP MH. 2011 Genomic

Determinants of Protein Evolution and Polymorphism in Arabidopsis. Genome Biology and Evolution.
3:1210-1219.

SNOOK RR, CHAPMAN T, MOORE PJ, WEDELL N, CRUDGINGTON HS. 2009 Interactions between
the sexes: new perspectives on sexual selection and reproductive isolation. Evolutionary
Ecology. 23(1):71-91.

STEPHENSON AG, BERTIN RI. 1983 Male competition, female choice, and sexual selection in plants. In:
Pollination Biology. (L R, ed), pp. 109-151. Academic Press, New York, NY.

STONE JL. 2002 Molecular mechanisms underlying the breakdown of gametophytic self-incompatibility.
Quarterly Review of Biology. 77(1):17-32.

SWANSON R, EDLUND AF, PREUSS D. 2004 Species specificity in pollen-pistil interactions.
Annual Review of Genetics. 38:793-818.

SWANSON WJ, VACQUIER VD. 2002 The rapid evolution of reproductive proteins. Nature Reviews
Genetics. 3(2):137-144.

SWANSON WJ, VACQUIER VD. 2002 Reproductive protein evolution. Annual Review of Ecology and
Systematics. 33:161-179.

SZOVENYI P, RICCA M, HOCK Z, SHAW JA, SHIMIZU KK, WAGNER A. 2013 Selection Is No More Efficient
in Haploid than in Diploid Life Stages of an Angiosperm and a Moss. Molecular Biology and Evolution.
30(8):1929-1939.

TAKAYAMA S, ISOGAI A. 2005 Self-incompatibility in plants. Annual Review of Plant Biology. 56:467-
489.

THE GENE ONTOLOGY CONSORTIUM. 2019 The Gene Ontology Resource: 20 years and still GOing
strong. Nucleic Acids Research. 47(D1):D330-D338.

TURNER LM, HOEKSTRA HE. 2008 Causes and consequences of the evolution of reproductive proteins.
International Journal of Developmental Biology. 52(5-6):769-780.

VACQUIER VD, SWANSON WJ. 2011 Selection in the Rapid Evolution of Gamete Recognition Proteins in
Marine Invertebrates. Cold Spring Harbor Perspectives in Biology. 3(11).

VELTSOS P. 2019 Not all sex-biased genes are the same. New Phytologist. 221(1):10-11.

VOSTERS SL, JEWELL CP, SHERMAN NA, EINTERZ F, BLACKMAN BK, MOYLE LC. 2014 The timing of
molecular and morphological changes underlying reproductive transitions in wild tomatoes (Solanum
sect. Lycopersicon). Molecular Ecology. 23(8):1965-1978.

WALSH NE, CHARLESWORTH D. 1992 Evolutionary interpretations of differences in pollen-tube growth
rates Quarterly Review of Biology. 67(1):19-37.

WILBURN DB, SWANSON WJ. 2016 From molecules to mating: Rapid evolution and biochemical studies
of reproductive proteins. Journal of Proteomics. 135:12-25.

WILLSON MF. 1979 Sexual selection in plants American Naturalist. 113(6):777-790.

WILLSON MF. 1994 Sexual selection in plants - perspective and overview American Naturalist. 144:S13-
S39.

WONG A. 2010 Testing the effects of mating system variation on rates of molecular evolution in
primates. Evolution. 64(9):2779-2785.

WONG A. 2011 The molecular evolution of animal reproductive tract proteins: what have we learned
from mating-system comparisons? International Journal of Evolutionary Biology. 2011(908735):1-9.

YANG Z. 2007 PAML 4: Phylogenetic Analysis by Maximum Likelihood. Molecular Biology and Evolution.
24(8):1586-1591.

1005
1006