CellKb Immune: a manually curated database of hematopoietic marker gene sets from 7 species for rapid cell type identification

Ajay Patil[1], Ashwini Patil[1*]

[1]Combinatics, Ichikawa-shi, Chiba, Japan.

* Correspondence:
Ashwini Patil
ashwini@combinatics.com

Keywords: single-cell, cell types, marker genes, immune cells, annotation, database, cell type
prediction

Abstract

Using single-cell RNA sequencing, it is now possible to get the detailed gene expression pattern in
each cell for millions of cells in a single experiment under varying conditions. Annotating cells by
assigning a type or state to them is important for correct interpretation of results. In spite of current
advances in technology, assigning cell types in single-cell datasets remains a bottleneck due to the
lack of a comprehensive reference database and an accurate search method in a single tool. CellKb
Immune is a knowledgebase of manually collected, curated and annotated marker gene sets from cell
types in the immune response in human, mouse and 5 other model organisms. It allows users to find
matching cell types in literature given a list of genes within a species and across multiple species. We
evaluated the contents and search methodology of CellKb Immune using a leave-one-out approach.
We further used CellKb Immune to annotate previously defined marker gene sets from a bulk RNA-
seq dataset to confirm its accuracy and coverage. CellKb Immune provides an easy-to-use online tool
with a reliable method to find matching cell types and annotate cells in single-cell experiments. It is
available at https://www.cellkb.com/immune.

1   Introduction

Single-cell RNA-seq is widely used to study transcriptional patterns of genes in individual cells.
Various computational tools have been developed for the annotation of cell types from single-cell
experiments, given a reference database and a gene list (1,2). Several single cell reference datasets or
atlases have also been prepared (3–5), or compiled into aggregate reference databases of cell types
and marker gene sets (6–9). There are currently two primary methods used for annotating cells from
single-cell experiments. 1) Manually searching through scientific literature to find marker genes
associated with cell types being studied. This requires prior knowledge of the cell types potentially
present in the sample and fails to identify rare cell types without established markers. 2) Selecting a
single cell reference database/ dataset from literature, downloading and processing the data,
identifying and installing a computational method to assign cell types, and finally, writing a program
in R or Python to perform the cell type assignment. This method usually limits a researcher to using a
single, or a few, datasets as a reference in a single species. Both these methods require significant
time and effort but fail to use the vast amount of cell type marker gene sets previously published.
Reliable annotations require a resource that contains a comprehensive reference database of cell types

and marker gene sets from literature along with a search method that allows users to look for
matching cell types.

CellKb Immune addresses these issues by collecting and aggregating a large amount of immune-
related cell type signatures across 7 species from literature and providing an online interface to find
matching annotated cell types easily, quickly and accurately.

2   Materials and Methods

2.1   Data collection and curation

The CellKb Immune database consists of marker gene sets with associated cell type annotations
collected from literature. The marker genes and their cell type annotations are taken primarily from
published single-cell RNA-seq experiments. All publications related to single-cell experiments are
taken from PubMed and manually screened to select those identifying cell type specific gene
expression patterns.

Cell type specific marker genes are defined as those identified by authors based on their significant
change in expression within a group of cells as compared to all other cells in the experiment. The
marker gene sets are extracted from tables, figures, supplementary materials and raw, or processed,
data of publications describing these experiments. If calculated based on the raw/processed
expression data, cell type specific markers are identified using the methods described by the authors
in the publication, along with cell type annotations given for the single cells.

Marker gene sets are selected from published experimental studies if they satisfy the following
criteria: 1) Raw data related to the experiment is deposited in public databases, 2) The marker gene
sets are publicly available for download, 3) The experimental methods and conditions are adequately
described in the manuscript to allow detailed annotation, 4) The number of cells and samples studied
is clearly specified, 5) The computational methods used to normalize, filter and cluster cell types,
along with identification of cluster-specific genes is clearly described, 6) Associated values such as
average expression, fold change, statistical significance, raw UMI or read counts, are available for
marker genes (marker gene sets specified only as ranked gene lists without associated values are
included in some cases), 7) A majority of the gene identifiers in the marker gene set are valid and can
be mapped to the latest version of the Ensembl database (14). Single-cell experiments without cell
type annotations are not used to calculate cell type specific marker gene sets.

Detailed sample and treatment conditions, including sample counts and cell counts are also stored in
CellKb Immune. Information about disease state of the samples and the cell types enriched in disease
samples are also noted. All data in CellKb Immune is referenced and linked to the PubMed identifier
of the source publication.

CellKb Immune also contains marker genes extracted from select bulk RNA-seq and microarray
experiments from public databases. These include selected gene signatures from Human Protein
Altas (10), MSig-db (11) and previously published bulk RNA-seq datasets of immune cell types
(1,12). Cell-type specific interaction networks are created using marker gene sets of all cell types and
protein-protein interactions with reliability scores from the HitPredict database (13).

2.2 Data harmonization and annotation

Data harmonization is essential for rapid search and retrieval of matching cell types, within and
across species. Given that there is no standard format currently in place for sharing cell type specific
marker gene sets, CellKb Immune performs a number of operations to harmonize the data collected.
Marker gene sets are extensively curated to identify valid genes and cell types. Genes given in the
published marker gene sets are mapped to valid Ensembl identifiers using all possible names,
synonyms and identifiers. Assigning a common gene identifier to the marker genes is an important
part of data harmonization in CellKb Immune. It also helps identify orthologous genes when using a
gene list from one species to search for cell types in another.

Cell types, tissue names and disease conditions are assigned standardized ontology terms using Cell
Ontology (15), Uberon Ontology (16) and Disease Ontology (17). Associated values given by
authors with each signature are also stored in CellKb Immune. These include, but are not limited to,
rank, score, average expression, log fold change and corrected/uncorrected p-values. Marker gene
sets that are published in the form of ranked gene lists without associated values are also included in
the database if they have been accurately annotated. Unranked gene lists without associated values
are not included in CellKb Immune.

Each cell type is assigned a reliability score that gives an indication of how similar it is to other cell
types of the same ontology. The reliability score is calculated as the ratio of the average Spearman's
correlation coefficient of a marker gene set with others of the same cell type to that of marker gene
sets of different cell types. The higher the reliability score of a marker gene set, the closer it is to
other marker gene sets of the same cell type and, hence, the more likely it is to be accurate. This
helps identify marker gene sets that are incorrect, calculated from erroneous cluster assignments by
authors, or that have been calculated from mixtures of cell types.

Finally, all marker genes and their annotations are stored in a uniform format with indexed identifiers
enabling rapid search and retrieval of the data.

2.3 Data searching

In CellKb Immune, users can search for cell type specific marker gene sets by giving a list of ranked
query genes. Query genes given by the user are compared with every marker gene set in the database
and the best match is identified using a match score. The match score is based on the number of
common genes between the query and cell type, their ranks, their rank differences and the total
number of significant genes in the cell type. This results in cell types sharing highly ranked genes
with the query being assigned higher match scores. This method also accounts for differences in the
sizes of gene lists between the query and various cell types, such that cell types with fewer marker
genes are not disregarded. To search cell types in a species different from that of the query gene list,
CellKb Immune identifies the orthologous genes in the species of interest using Ensembl Compara
(18) before searching for matching cell types.

3   Results

3.1   Database contents

CellKb Immune contains author-defined immune cell type marker gene sets manually collected from
publications describing mainly single-cell, and selected bulk RNA-seq or microarray experiments
(1,10,11) (Figure 1). Cell type marker gene sets are assigned a Cell Ontology term (15), where
available. Tissues, organs or body parts of the experimental sample are assigned Uberon ontology
terms (16). Disease ontology terms are provided where the marker gene sets are obtained from a
diseased sample (17). All marker gene sets reference their source publication with a link to its
PubMed identifier.

CellKb Immune contains:

• 6,604 marker gene sets
• 560 unique annotated hematopoietic cell types (Cell Ontology CL:0000988 and all its descendants).
• 439 publications selected from approximately 10,000 studies published between 2013 and 2021.
• 852,305 protein-protein interactions from HitPredict (13)
• 7 species - *H. sapiens*, *M. musculus*, *D. rerio*, *R. norvegicus*, *M. fascicularis*, *M. mulatta*, *D.*
melanogaster

Figure 2 shows the number of marker gene sets for each species in CellKb Immune.

3.2   Functionality

The CellKb Immune web interface allows users to search matching cell types and their marker gene
sets within and across species. It can be used to annotate cells from single-cell RNA-seq data, given a
list of statistically significant cluster-specific genes ranked by decreasing log fold change. Associated
values such as log fold changes and p-values are also accepted as input by CellKb Immune. Genes in
the user query are compared with every marker gene set in the database using a rank-based search
method and a match score is computed. To identify matching cell types in another species, CellKb
Immune identifies the orthologous genes between the query and the target species before searching.

CellKb Immune is limited to marker gene sets of hematopoietic cell types. However, the user is
notified if the query gene list matches non-hematopoietic cell types in 5 out of top 10 cell type
matches.

3.3   Use case: Assigning cell types in a single-cell RNA-seq experiment

To demonstrate the functionality of CellKb Immune, we used it to assign cell types to a single-cell
RNA-seq dataset of human peripheral blood mononuclear cells from the Single Cell Portal (19).

1) To query CellKb Immune for matching cell types, we used the cell type specific genes, along with
their associated log fold changes and p-values calculated in the study for primary lineage clusters
defined by the authors. Cell-type specific genes for each cluster were limited to those that were
upregulated at least two-fold and had an adjusted p-value less than or equal to 0.05. The filtered
query genes were ranked in descending order of their log fold change before submission to CellKb
Immune (Table S1).

2) Users have an option to search for matching cell types taken single-cell and/or bulk experiments,
and filter the search based on cell types, organs/tissues, diseases and other conditions. No filters were
set for this use case.

3) The CellKb Immune results display a heatmap with the rank-based score of the top 100 matching
cell types (Figure 3).

4) Details of the experiment containing the matching cell type, such as PubMed ID, condition,
number of cells, cell type ontology, etc. are shown in tabular format for the top 10 hits. Other
statistics such as the statistical significance by Fisher's exact test, Pearson's correlation coefficient
and Jaccard index are also calculated for each cell type match. These results are available for
download.

5) Users can also view the top 10 overlapping genes between the query and the marker gene set of the
matching cell type.

6) Cell type specific protein-protein interaction network containing the user query genes and the top
10 overlapping genes are also displayed.

CellKb Immune was able to correctly identify 7 of the 8 cell types assigned by the authors. One cell
type which was assigned as "Ambiguous/Potential Doublets", was matched to Plasmablasts by
CellKb Immune. Table S1 gives the top 10 matching cell types identified by CellKb Immune for
each of the 8 lineage clusters of cells.

The cells in this dataset were also classified into 16 clusters using k-means clustering. We used the
cluster specific genes for these 16 clusters to find cell type annotations with CellKb Immune (Table
S1). Cell types predicted by CellKb Immune are identical to the author-defined annotations. The top
10 hits for each gene list given to CellKb Immune is shown in Table S1.

3.4   Evaluation

To check the database quality and cell type matching performance of the CellKb Immune search
method, we used the leave-one-out approach. We matched each of the 2,223 mouse marker gene sets
with cell ontologies in CellKb Immune with all others in the database. 1,568 (70.5%) of gene sets
were correctly matched to another gene set with the exact same ontology or its parent, while for 427
(19.2%) gene sets, the ontology of the first hit was a sibling i.e. a descendant of the same parent
ontology (Table 1). Thus, 89.7% of marker gene sets were matched to another cell type with the same
ontology, or its parent or sibling (Table S2).

We further evaluated the performance of the CellKb Immune by checking the number of correct cell
types assigned to gene lists obtained from a bulk RNA-seq dataset of mouse immune cell types (12)
from Immgen (21). Using the most significantly upregulated genes for each cell type in the query
dataset, we searched the CellKb Immune database for matching cell types. We then compared the cell
ontology of the top 10 hits with that of the ontology assigned to the cell types in the query dataset. Of
the 98 cell type signatures representing 40 distinct cell types, CellKb Immune was able to find the
exact matching ontology or its parent as the first hit in 61 instances (62.2%). In another 18 cell types
(18.4%), the ontology of the first hit was a sibling i.e. a descendant of the same parent ontology as
that in the bulk RNA-seq dataset (Table 1). Thus, a total of 79.5% of cell types in the query dataset
had their first hit with the same ontology, its parent or sibling (Table S3).

These results show that 1) the CellKb Immune database includes high quality cell type signatures
from single-cell experiments, and 2) the search methodology used by CellKb Immune to find
matching cell types is able to recapitulate the original cell types from within its own database and
other unseen bulk and single-cell datasets.

4   Discussion

CellKb Immune has been designed as a resource to assign cell types to cells in single-cell RNA-seq
experiments. CellKb Immune addresses several issues in existing single-cell reference databases and
tools used for this purpose.

CellKb Immune consists of user-defined marker gene sets from literature. These carry significant
knowledge since the cell types in the form of cell clusters are often selected by authors based on
biological information. CellKb Immune is able to capture and aggregate this biological information.
On the other hand, several reference databases reanalyze the raw expression patterns using a common
pipeline while ignoring the methodology and cluster definitions used by the authors in the original
study, thus losing valuable biological information.

CellKb Immune provides deep annotations, in the form of extensive cell type information and
description, curated and validated marker genes ranked by significance, along with fold change and
significance values associated with gene expression. Additionally, CellKb Immune provides a web-
based interface to find matching cell types in published data sets given a user gene list, independent
of the experimental platform, analysis methods and insufficient marker gene sets. Thus, users do not
need to spend time integrating data and search methods programmatically.

Unlike other methods which require the presence of associated expression or fold change values and
the same number of genes in all target cell types, the rank-based search method used by CellKb
Immune enables searching through marker gene sets of differing sizes, in the absence of expression
fold changes, and independent of experimental platform and pre-processing methods. Searching of
cell types happens by direct comparison of the query gene list with the marker genes of every cell
type in CellKb Immune using a fast and reliable method without any type of score aggregation or
machine learning. As a result, CellKb Immune provides information about the matching cell type in
the form of the publication, experimental details and the overlapping genes between the query and
matching marker gene set. This allows users to further explore the study to which their cell type
matches.

Since CellKb Immune contains marker gene sets from multiple publications for a single a cell type, it
can also be used to evaluate the validity of known marker genes for the cell type of interest. The
reliability score assigned to each cell type marker gene set can also be used to identify those that are
incorrectly annotated in publications. Additionally, comparing the query gene list with marker gene
sets from multiple publications helps assess the accuracy of the matching cell types found by CellKb
Immune.

Thus, CellKb Immune provides an easy-to-use reference database with a fast and reliable method to
find matching cell types and marker gene sets to help annotate cells in single-cell experiments in a
single tool.

5 Conflict of Interest

Ashwini is CEO at Combinatics Inc., a privately-held company. Ajay and Ashwini are stock-holders in Combinatics Inc.

6 Author Contributions

Ashwini conceived of the project and collected the data. Ajay and Ashwini designed and developed the database and web application, and prepared the manuscript.

7 References

1. Aran D, Looney AP, Liu L, Wu E, Fong V, Hsu A, Chak S, Naikawadi RP, Wolters PJ, Abate AR, et al. Reference-based analysis of lung single-cell sequencing reveals a transitional profibrotic macrophage. *Nat Immunol* (2019) doi:10.1038/s41590-018-0276-y

2. Pliner HA, Shendure J, Trapnell C. Supervised classification enables rapid annotation of cell atlases. *Nat Methods* (2019) doi:10.1038/s41592-019-0535-3

3. Rozenblatt-Rosen O, Stubbington MJT, Regev A, Teichmann SA. The Human Cell Atlas: From vision to reality. *Nature* (2017) doi:10.1038/550451a

4. Han X, Wang R, Zhou Y, Fei L, Sun H, Lai S, Saadatpour A, Zhou Z, Chen H, Ye F, et al. Mapping the Mouse Cell Atlas by Microwell-Seq. *Cell* (2018) doi:10.1016/j.cell.2018.02.001

5. Schaum N, Karkanias J, Neff NF, May AP, Quake SR, Wyss-Coray T, Darmanis S, Batson J, Botvinnik O, Chen MB, et al. Single-cell transcriptomics of 20 mouse organs creates a Tabula Muris. *Nature* (2018) doi:10.1038/s41586-018-0590-4

6. Ner-Gaon H, Melchior A, Golan N, Ben-Haim Y, Shay T. JingleBells: A Repository of Immune-Related Single-Cell RNA–Sequencing Datasets. *J Immunol* (2017) doi:10.4049/jimmunol.1700272

7. Zhang X, Lan Y, Xu J, Quan F, Zhao E, Deng C, Luo T, Xu L, Liao G, Yan M, et al. CellMarker: A manually curated resource of cell markers in human and mouse. *Nucleic Acids Res* (2019) doi:10.1093/nar/gky900

8. Franzén O, Gan LM, Björkegren JLM. PanglaoDB: A web server for exploration of mouse and human single-cell RNA sequencing data. *Database* (2019) doi:10.1093/database/baz046

9. Papatheodorou I, Moreno P, Manning J, Fuentes AMP, George N, Fexova S, Fonseca NA, Füllgrabe A, Green M, Huang N, et al. Expression Atlas update: From tissues to single cells. *Nucleic Acids Res* (2020) doi:10.1093/nar/gkz947

10. Uhlen M, Zhang C, Lee S, Sjöstedt E, Fagerberg L, Bidkhori G, Benfeitas R, Arif M, Liu Z,

Edfors F, et al. A pathology atlas of the human cancer transcriptome. *Science (80-)* (2017)
doi:10.1126/science.aan2507

11. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A,
Pomeroy SL, Golub TR, Lander ES, et al. Gene set enrichment analysis: A knowledge-based
approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* (2005)
doi:10.1073/pnas.0506580102

12. H Y, CA L, RN R, SA R, B M, A W, F D, A C, A M, C D, et al. The cis-Regulatory Atlas of
the Mouse Immune System. *Cell* (2019) **176**:897-912.e20. doi:10.1016/J.CELL.2018.12.036

13. López Y, Nakai K, Patil A. HitPredict version 4: Comprehensive reliability scoring of physical
protein-protein interactions from more than 100 species. *Database* (2015) **2015**:
doi:10.1093/database/bav117

14. Howe KL, Achuthan P, Allen J, Allen J, Alvarez-Jarreta J, Amode MR, Armean IM, Azov
AG, Bennett R, Bhai J, et al. Ensembl 2021. *Nucleic Acids Res* (2021) **49**:D884–D891.
doi:10.1093/NAR/GKAA942

15. Bard J, Rhee SY, Ashburner M. An ontology for cell types. *Genome Biol* (2005)
doi:10.1186/gb-2005-6-2-r21

16. Mungall CJ, Torniai C, Gkoutos G V., Lewis SE, Haendel MA. Uberon, an integrative multi-
species anatomy ontology. *Genome Biol* (2012) doi:10.1186/gb-2012-13-1-r5

17. Schriml LM, Arze C, Nadendla S, Chang YWW, Mazaitis M, Felix V, Feng G, Kibbe WA.
Disease ontology: A backbone for disease semantic integration. *Nucleic Acids Res* (2012)
doi:10.1093/nar/gkr972

18. Herrero J, Muffato M, Beal K, Fitzgerald S, Gordon L, Pignatelli M, Vilella AJ, Searle SMJ,
Amode R, Brent S, et al. Ensembl comparative genomics resources. *Database* (2016) **2016**:96.
doi:10.1093/DATABASE/BAV096

19. Single Cell Portal. Available at:
https://singlecell.broadinstitute.org/single_cell/study/SCP345/ica-blood-mononuclear-cells-2-
donors-2-sites?scpbr=immune-cell-atlas [Accessed October 25, 2021]

20. Hurskainen M, Mižíková I, Cook DP, Andersson N, Cyr-Depauw C, Lesage F, Helle E,
Renesme L, Jankov RP, Heikinheimo M, et al. Single cell transcriptomic analysis of murine
lung development on hyperoxia-induced damage. *Nat Commun* (2021) **12**:1565.
doi:10.1038/s41467-021-21865-2

21. Aguilar SV, Aguilar O, Allan R, Amir EAD, Angeli V, Artyomov MN, Asinovski N, Astarita
J, Austen KF, Bajpai G, et al. ImmGen at 15. *Nat Immunol* (2020) doi:10.1038/s41590-020-
0687-4

8 Figures

Figure 1. CellKb Immune workflow – data collection, harmonization and search interface.

Figure 2. Number of marker gene sets for each species in CellKb Immune.

Figure 3. CellKb Immune user interface used to search matching cell types given a gene list.

9   Tables

Table 1. Performance evaluation of CellKb Immune

Evaluation database	Marker gene sets	CellKb Immune performance		Total ontologies matched
		Correct ontology match	Parent ontology match	
CellKb Immune Mouse (leave-one-out)	2,223	1,568 (70.5%)	427 (19.2%)	**1,995 (89.7%)**
Yoshida et al. (12) **Mouse (all-against-all)**	98	61 (62.2%)	18 (18.4%)	**79 (80.6%)**

10   Supplementary Material

Table S1. List of cluster-specific genes for 8 and 16 pre-defined clusters in human peripheral blood
mononuclear cells used as input to CellKb Immune. Top 10 results of matching cell types obtained
from CellKb Immune.

Table S2. Results of the leave-one-out evaluation in CellKb Immune shows the first hit of each
mouse cell type marker gene set compared to all other cell types.

Table S3. First hit of each cell type marker gene set in a bulk RNA-seq dataset of immune cell types
compared to all cell types in CellKb Immune mouse.

11   Data Availability Statement

CellKb Immune is available at https://www.cellkb.com/immune. All data used in CellKb Immune is
publicly available and appropriately referenced in the web application.

Search cell types using ranked gene list

View mapping of gene names to Ensembl IDs

View heatmap of matching cell types with gene scores

View cell type specific protein interaction network

View matching marker genes in hit cell type

View table of matching cell types with source details