Species-resolved sequencing of low-biomass microbiomes by 2bRAD-M

Zheng Sun[1,4,8,§], Shi Huang[1,4,8,§,#], Pengfei Zhu[1,4,8], Lam Tzehau[3], Helen Zhao[3], Jia Lv[2,4], Rongchao

Zhang[5], Lisha Zhou[5], Qianya Niu[5], Xiuping Wang[5], Meng Zhang[6], Gongchao Jing[1,4,8], Zhenmin

Bao[2,4,7], Jiquan Liu[3], Shi Wang[2,4,7,#], Jian Xu[1,4,8,#]

[1]Single-Cell Center, CAS Key Laboratory of Biofuels and Shandong Key Laboratory of Energy

Genetics, Qingdao Institute of BioEnergy and Bioprocess Technology, Chinese Academy of Sciences,

Qingdao, Shandong, China

[2]MOE Key Laboratory of Marine Genetics and Breeding and Sars－Fang Centre, Ocean University

of China, Qingdao, Shandong, China

[3]Procter & Gamble Singapore Innovation Center, Singapore

[4]Qingdao National Laboratory of Marine Science and Technology, Qingdao, Shandong, China

[5]Qingdao OE Biotechnology Company Limited, Qingdao, Shandong, China

[6]Key Laboratory of Dairy Biotechnology and Engineering, Ministry of Education, Inner Mongolia

Agricultural University, Hohhot, China

[7]Laboratory of Tropical Marine Germplasm Resources and Breeding Engineering, Sanya

Oceanographic Institution, Ocean University of China, Sanya, China

[8]University of Chinese Academy of Sciences, Beijing, China

[§]Equal contribution

[#]Corresponding to: xujian@qibebt.ac.cn (Jian Xu), Shi Wang (swang@ouc.edu.cn) and Shi Huang

(huangshi@qibebt.ac.cn)

Fax: (86) 532-80662654, Phone: (86) 532-80662651

Running title: Low-biomass microbiome sequencing

Keywords: IIB restriction enzyme, reduced metagenome sequencing, low-biomass samples,

microbiome

Abstract

Microbiome samples with low microbial biomass or severe DNA degradation remain challenging for amplicon-based (e.g., 16S/18S-rRNA) or whole-metagenome sequencing (WMS) approaches. Here, we introduce 2bRAD-M, a "reduction" metagenome-sequencing strategy that simultaneously produces species-level bacterial, archaeal, and fungal profiles for low-biomass samples, yet is cost-competitive with 16S rRNA gene analysis. For mock communities, 2bRAD-M can accurately generate species-level taxonomic profiles for otherwise hard-to-sequence samples with (*i*) low biomass of merely 1 pg of total DNA, (*ii*) high host DNA contamination (99%), and (*iii*) severely fragmented DNA (50-bp) from degraded samples. Tests of 2bRAD-M on stool, skin and environment-surface samples deliver successful reconstruction of comprehensive, high-resolution microbial profiles with agreement across 16S-rRNA, WMS and existing literature. In addition, it enables microbial profiling in formalin-fixed paraffin-embedded (FFPE) cervical tissue samples which were recalcitrant to conventional approaches due to the low amount and heavy degradation of microbial DNA, and discriminated healthy tissue, pre-invasive cancer and invasive cancer via species-level microbial profiles with 91.1% accuracy. Therefore, 2bRAD-M greatly expands the reach of microbiome sequencing.

Introduction

Metagenome sequencing, widely used to derive the taxonomic profile of microbiome, typically adopts two strategies that target (*i*) amplicons of phylogenetic "marker genes" (e.g., 16S rRNA for bacteria and archaea, and 18S rRNA or internal transcribed spacer (ITS) for fungi) or (*ii*) the whole genomes (whole metagenome shotgun; WMS). Although less costly, marker gene analyses can be limited in taxonomic resolution (i.e. at the genus level) and susceptible to PCR bias in composition and abundance estimates[1]; moreover they are usually unable to capture a landscape-like view that includes bacteria, archaea, fungi and virus due to the lack of universal primers. In contrast, by sequencing the total DNA, WMS can resolve species- or strain-level taxonomy, and offers a landscape-like view that includes all domains of organisms [2, 3]. However, WMS requires a rather high amount of DNA as the starting material (≥50 ng preferred, 20 ng at a minimum), and are usually unable to tackle DNA samples that are low in biomass, heavily degraded or dominated by host DNA [1, 4]. Moreover, WMS is typically much more costly, due to the much higher sequencing volume required for covering the whole genome (instead of just the marker gene). Therefore, new methods should be developed that cost-efficiently produce accurate, species-resolution, landscape-like taxonomic profiles for challenging samples like low-biomass, high-host-contaminated and degraded microbiomes.

Restriction site-Associated DNA sequencing (RADseq) that utilizes restriction enzymes to digest genomic DNA from a board range of organisms and sequences only the digested fragments has been applied to genotype variable genetic markers as well as for microbial species detection[5, 6]. Although RADseq has demonstrated the ability to produce species-resolution microbiome profiles with lower costs [7-10], the large size variation of DNA fragments after enzyme digestion results in a

high bias in PCR-based amplification and thus the low fidelity of reconstructed taxonomic profile[11,]

[12]. Moreover, RADseq has not been thoroughly and systematically benchmarked against marker

gene-based or WMS approaches [7-10]. Therefore, it remains unclear whether RADseq can provide

accurate, landscape-like taxonomic profiles for low-biomass and severely degraded microbiome

samples.

To address these challenges, here we proposed "2bRAD sequencing for Microbiome"

(2bRAD-M) method that utilizes Type IIB restriction enzymes to produce exclusively iso-length

DNA fragments of 21-33 bp (depending on the selection of enzymes) for sequencing. This approach

reduces the bias in PCR-based fragment amplification, and thus ensures the high fidelity of

taxonomic profile. This offers significant benefits especially for low-biomass, heavily contaminated

or degraded microbial DNA which requires more PCR cycles. Tests on mock and actual

microbiome samples illustrate that of 2bRAD-M, by sequencing just 1% of genomes, accurately

generates species-level taxonomic profiles for challenging samples: (*i*) of merely 1 pg total DNA, (*ii*)

of 99% host DNA contamination or (*iii*) consisting of highly degraded fragments just 50-bp long.

For real stool, skin and environment-surface samples, it accurately reconstructs a comprehensive,

species-resolution profile of bacteria, archaea and fungi. Furthermore, microbiome in the

formalin-fixed paraffin-embedded (FFPE) tissue samples which were otherwise recalcitrant to

sequencing can now be analyzed, and a species-resolved classifier discriminated the healthy tissue,

pre-invasive cancer and invasive cancer with 91.1% accuracy. The ability to profile low-biomass

microbiomes at the species level is pivotal to expanding the boundary of known microbial world.

Results

The principle and workflow of 2bRAD-M

The principle of 2bRAD-M is (**Fig. 1**): (*i*) reliable enzyme-digested sequence tags can be

derived that are specific to high-resolution taxa (e.g., species or strain) yet universally applicable for

a broad range or all of bacterial, archaeal and fungal genomes; (*ii*) these taxa-specific, iso-length

sequence tags can be evenly amplified and sequenced, and (*iii*) the tag sequences can be mapped to

reference genomes to reconstruct faithfully the taxonomic composition.

Specifically, the experimental workflow has two steps (**Methods**): (*i*) BcgI (a commercially

available Type IIB restriction enzymes) is used, as an example, to digest total genomic DNA

extracted from microbiome samples. BcgI recognizes the sequence of $CGA-N_6-TGC$ in the genomic

DNA and cleaves on both upstream (12-10 bp) and downstream (10-12 bp) of this signature [13],

producing short and iso-length DNA (32bp without sticky ends) across all loci [14, 15]. (*ii*) These

so-called "2bRAD fragments" are ligated to adaptors, amplified and then sequenced.

In the computational workflow (**Methods**), the foundation is a unique 2bRAD tag database

("2b-Tag-DB"), which contains taxa-specific 2bRAD tags identified from all the sequenced bacteria,

fungi and archaea genomes. Mapping the 2bRAD reads against 2b-Tag-DB thus identifies the

presence of species in a sample. Subsequently, to estimate relative abundance of the identified taxa,

the mean read coverage of all 2bRAD tags specific to each taxon is derived. To improve utilization

rate of reads and classification accuracy, a secondary, sample-specific 2b-Tag-DB was dynamically

derived from only those candidate taxa identified in a particular sample, which produces more

species-specific 2bRAD tags than the original 2b-Tag-DB and results in more accurate modeling of

relative abundance of taxa (**Methods**).

The feasibility of 2bRAD-M for microbiome analysis via *in silico* simulation

To identify the taxa-specific tags, we first downloaded 173,165 microbial genomes (171,927

bacteria, 293 fungi and 945 archaea; representing 26,163 species) from NCBI RefSeq (Oct, 2019) to

create a 2b-Tag-DB via *in-silico* restriction digestion of these genomes using BcgI restriction enzyme.

This yields an average of 3,010 iso-length (32bp) 2bRAD tags per genome, and a total of 114,132,487

BcgI-digested unique species-specific 2bRAD tags that are of single-copy within a genome (average

1,949 per species genome; **Supplementary Methods**). Besides BcgI, species-specific 2bRAD tags

were also identified for each of the other 15 Type IIB enzymes (AlfI, AloI, BaeI, BplI, BsaXI, BslFI,

Bsp24I, CjeI, CjePI, CspCI, FalI, HaeIV, Hin4I, PpiI and PsrI; **Table S1**) to establish the usability of

all Type IIB enzymes for 2bRAD-M. Overall, the number of 2bRAD-M tags within a genome and

their GC content are highly consistent with the length and the GC% of genomes (r>0.98) (**Fig. S1**) [15]

for all the 16 Type IIB enzymes, suggesting unbiased, broadly applicable representation of the

microbial genomes by these tags.

To test whether these species-specific tags enable detection and abundance profiling of all

known species in a community (**Methods**), a simulated 50-species microbiome was generated (one

genome per species; randomly selected from RefSeq; **Table S2**; **Methods**) and profiled using the

default 2b-Tag-DB and sample-specific 2b-Tag-DB for each of the 16 Type IIB restriction enzymes.

Performance of 2bRAD-M in species detection was assessed via precision and recall (with relative

abundance threshold of 0.0001), while that of species abundance evaluated via L2 similarity score (a

metric of similarity adapted from L2 distance [16]), by comparing to the ground truth (**Fig. 2**).

Precision, recall and L2 similarity of the taxonomic profiling are all remarkably high (average for the

16 enzymes - precision=98.0%, recall=98.0%, L2 similarity=96.9%), and this is achieved with an

average genome coverage of 1.50% among the 50 selected genomes (**Fig. 2**).

High reproducibility and sensitivity of 2bRAD-M under challenging conditions

To assess 2bRAD-M ability to handle low biomass, highly degraded or heavily contaminated samples, we first constructed a mock community (Mock-CAS) consisting of five prevalent oral or gut bacterial species in equal proportion (**Methods**). With this mock community stock, three series of samples were prepared : (*i*) "LoA" (low amount) – samples with total DNA amounts from 50 ng, 20 ng, 10 ng, 1 ng, 100 pg, 10 pg to only 1 pg; (*ii*) "HiD" (high degradation):– 100ng degraded DNA samples where the DNA mixture was randomly sheared by DNAse I into fragments of mostly 150-bp and 50-bp in length; (*iii*) "HoC" (host contamination): – 100ng samples spiked with human DNA to simulate 90% or 99% contamination by host genome. Three technical replicates were included for each of the LoA (n=7*3), HiD (n=2*3) and HoC (n=2*3) groups. Each of the 33 samples from the three series was then sequenced by 2bRAD-M. In addition, a sample of 100ng DNA was then prepared from the mock community stock and sequenced by WMS to be referenced as the positive control (WMS of 50ng or lower DNA failed).

To compare the results, we employed Centrifuge[17] to map WMS (150-bp) and 2bRAD-M (32-bp) generated reads to the reference microbial genomes for taxonomic abundance estimation. For 2bRAD-M, tags from the actual data of each sample covered avg. 97.1% *in silico* predicted tags from the five genomes, indicating high consistency between observed and expected 2bRAD tags (which is key to accurate and reliable taxonomic profiling). Moreover, for each sample, the 2bRAD-M results are highly consistent among the three biological replicates (avg. L2 similarity: 95.4%; **Fig. 3a**).

Notably, for the LoA, HiD and HoC groups, average L2 similarity between 2bRAD-M profiles and the reference positive control is 88.9%, 84.6% and 89.6% respectively, with global average

being 88.2% (ranging from 82.1% to 90.4%; **Fig. 3a**). These observations are consistent across the

technical replicates, indicating high reproducibility of 2bRAD-M. Specifically, in the LoA group,

the L2 similarity of the 1 pg sample can still realize a respectable 83.5% as compared to 90.1% for

the 50 ng sample (**Fig. 3a**). This suggests that 2bRAD-M offers high sensitivity and stable

performance in low biomass samples over a broad DNA-amount range (from 50 ng to 1 pg). In the

HiD group, the L2 similarity is 87.2% and 82.1% for the 150bp- and 50bp-samples respectively,

indicating DNA degradation did not have a large negative effect and the 2b-RAD-M can effectively

accommodate severe DNA degradation while providing reliable results. The L2 similarity for the

HoC group is 89.4% in the 90% host-contaminated samples and 89.7% in 99%-host-contaminated

samples, suggesting 2bRAD-M's ability to provide reliable microbial profiling in the presence of

host-DNA contamination.

To further probe 2bRAD-M proficiency in profiling highly complex microbial community at

low biomass setting, we used ATCC MSA1002, a standard, commercially available mock

community consisting of 20 bacterial species (from 18 genera) with at equal DNA abundance

among species[18] (**Methods**). Samples with total DNA amount range from 10 ng, 1 ng, 100 pg, 10 pg

and 1 pg were prepared from this mock for 2bRAD-M profiling. As a reference control, a 100 ng

sample was profiled with 16S-rRNA and WMS approach (the 1pg ~10ng samples failed). The

precision for 2bRAD-M is at 90.1%, 76.0%, 46.3%, 40.0% and 14.0% for the 10 ng, 1 ng, 100 pg,

10 pg and 1pg samples respectively; this is in contrast to the 29.8% of WMS-100ng and the 17.3%

of 16S-rRNA-100ng (**Fig 3b**). This demonstrates that 2bRAD-M offers much lower false positives

in detecting the species than 16S-rRNA and WMS. For recall, 2bRAD-M exhibits 100.0%, 95.0%,

95.0%, 80.0%, 75.0% for 10 ng, 1 ng, 100 pg, 10 pg and 1 pg respectively, as compared to the 100.0%

of WMS-100ng and the 90.0% of 16S-rRNA-100ng. Therefore, 2bRAD-M offers comparable level

of sensitivity as WMS and 16S-rRNA with down to 100 pg of DNA amount. The L2 similarity for

2bRAD-M is 90.1%, 89.1%, 88.5%, 84.8% and 48.5% for 10 ng, 1 ng, 0.1 ng, 10 pg and 1pg

respectively, as compared to 86.3% for WMS-100ng and 70.0% for 16S-rRNA-100ng. Taken as a

whole, these results evidently demonstrated 2bRAD-M's ability to profile complex microbiota with

a high level of sensitively and specificity.

Validation of 2bRAD-M in real samples (fecal)

To assess the performance of 2bRAD-M on real samples, we performed and compared

2bRAD-M and 16S rRNA sequencing on human fecal samples (n=3). In addition, each sample was

also subjected to ultra-deep WMS sequencing (mean of 437 million reads or 239.73 Gb per sample)

with the resultant taxonomic profiles used as evaluation reference[19].

Taxonomic profiles from 2bRAD-M are compared with those from 16S rRNA sequencing (**Fig.**

4a-c) and WMS (**Fig. 4d-f**; **Supplementary Methods**). Specifically, at the genus-level, results of

2bRAD-M and 16S rRNA are highly consistent (mean Pearson correlation R=0.997 and mean L2

similarity L2=92.0%), and avg. 95.27% of genus identified by 16S were also detected by 2bRAD-M

(**Table S3**). Importantly, at the species level, 2bRAD-M and the WMS are also concordant, as

evidenced by a high Pearson correlation (R=0.974) and high L2 similarity (up to 88.6%). Moreover,

only 3.35% of the taxa in WMS were not identified in 2bRAD-M. Thus 2bRAD-M can produce

highly complete and accurate species-level profiles that are equivalent to WMS.

One determinant of quality (and cost) of taxonomic profiling is sequencing depth [20]. To probe

how it influences 2bRAD-M performance, the fecal sequencing datasets were subsampled to various

depths for 2bRAD-M (average 14 million reads per sample, subsampled at 0.1, 1, 2, 3, 5 to 14 million

sequences per sample) or WMS (average 437 million reads per sample, subsampled at 0.5, 10, 20, 30,

40, 50, 100, and 200 million sequences per sample) respectively. Then we asked how many 2bRAD

sequences are required to accurately quantify key ecological metrics such as alpha diversity (Shannon

index), beta diversity (Bray-Curtis), and species compositions (species richness). In each of the fecal

samples, with 20-40 million reads per sample, WMS identified averagely 170-182 species, but the

Shannon diversity and species abundance estimates are far from saturation (i.e., reaching the values

derived by sequencing ~400 million reads). In contrast, with just 3-4 million reads per sample,

2bRAD-M identified 173-188 species in each fecal samples and yield Shannon diversity and species

abundance estimates comparable with those from the deepest (~14 million) read depth (**Fig. S2**).

To evaluate the optimal sequencing depth of 2bRAD-M for species-level taxonomic profiling,

we evaluated the similarity of taxonomic profiles derived at each reduced sequencing depth to its

original taxonomic profile using L2 similarity score. For 2bRAD-M, a sequencing depth of over 0.5

million reads per samples can achieve a L2 similarity score of 98.9% (1 million for 99.1%), which

however requires many more (at least about 200 million) reads per sample in WMS (**Fig. 4g**).

Overall, with ~3 million reads (i.e., 60 Mb of sequencing data) per sample, 2bRAD-M can generate

consistent, accurate and stable alpha diversity estimates and taxonomic profile at the species level.

2bRAD-M enables species-resolved analysis of low-biomass skin, home and car samples

To assess 2bRAD-M performance on actual low-biomass samples, we collected samples from

human skin surface (underarm, n=20; **Table S4**) for 16S-rRNA and 2bRAD-M analysis (WMS was

aborted due to too-low DNA amounts for library construction; **Methods**). To gauge the potential

impact of contaminating DNA (introduced from regents, in workflow, etc.) in the low-biomass

samples [21], MSA1002 was included as a control. MSA1002 profiling revealed the expected microbial

community profile while unforeseen microbes were absent which indicates minimal contamination.

Between 16S rRNA and 2bRAD-M, a high degree of consistency in taxonomic profiles at the genus

level was observed, with avg. L2 similarity of 81.1% (**Table 1**; **Fig. S3**). In both methods,

Staphylococcus (2bRAD-M: avg. 47.17%; 16S rRNA: avg. 44.43%) and *Corynebacterium*

(2bRAD-M: avg. 10.38%; 16S rRNA: avg. 15.06%) are recognized as dominant microbes,

consistent with existing literature [22, 23]. Interestingly, unsupervised clustering at genus-level

uncovered two distinct clusters among the 20 underarm samples, with one featuring higher

abundance of *Staphylococcus* observed in both 16S-rRNA and 2bRAD-M (2.41-fold, p=4.6E-06,

Wilcoxon rank-sum test, for 16S rRNA; 3.20 fold, p=1.08E-05, Wilcoxon rank-sum test, for

2bRAD-M; **Fig. 5a**). Notably, the species-level profile from 2bRAD-M readily revealed that the

observed difference in *Staphylococcus* was defined by higher relative abundance of *S. epidermidis*

(4.63-fold, p=4.3E-04), with no significant change in other *Staphylococcus* spp. between the clusters

(**Fig. 5b**), which suggests interesting biological hypothesis on *S. epidermidis* functionality (notably,

S. epidermidis can protect against pathogens such as *S. aureus* by producing antimicrobial peptides

that modulate skin microbiome dysbiosis in atopic dermatitis [24]). These observations would have

been missed with 16S rRNA, thus 2bRAD's ability to resolve such species-level heterogeneity (e.g.

Staphylococcus spp.) in low-biomass samples can potentially enable novel biological insights.

We further explored the application of 2bRAD-M, by testing on in-door environmental

low-biomass samples, which are of significant health implications due to their contact with humans

[25]. Surfaces of floor mat (n=5) and cushion (n=3) in car, and interior surfaces of a house (n=4) were

sampled and subjected to both 16S rRNA and 2bRAD-M (similarly, WMS was aborted due to the

low DNA amounts; **Table S4**). A high L2 similarity (average 85.6% among samples; **Fig. S3**)

between 2bRAD-M and 16S rRNA was observed, suggesting stable performance of 2bRAD-M for

such in-door samples. At the species level (**Table S5**), we identified 4929, 2335 and 1626 taxa for

the three niches respectively, whose dominant bacterial species are highly distinct: (*i*) floor mats are

dominated by *Kocuria rosea* (23.23%), *Psychrobacter 1501_2011* (11.44%) and *Acinetobacter*

johnsonii (10.43%); (*ii*) cushions are dominanted by *Cutibacterium acnes* (42.19%), *Lactobacillus*

delbrueckii (10.62%) and *Ralstonia pickettii* (10.14%); (*iii*) home surfaces are mainly colonized by

Streptococcus mitis (49.64%), *Prevotella copri* (39.14%) and *Megamonas funiformis* (18.36%).

Rarefaction of sequence depth (e.g. sequenced 2b-tags) via alpha diversity, beta diversity and

species-level compositions supports high robustness of 2bRAD-M for such low-biomass in-door

samples (**Fig. S4**).

Similar to WMS, 2bRAD-M enabled species-level profiles for fungi along with bacteria. In

general, the relative abundance of fungi in the underarm and indoor environment samples is

extremely low (0.83%) as compared to bacteria (99.16%) (**Table S6**). Possible reasons for this

observation include: (*i*) micro-environments of these samples are not friendly fungal growth[26]; (*ii*) the

relatively small number of available fungi genomes curated in our reference database limits the

discovery of the fungi population. Nonetheless, among all the sites, the samples taken from car

cushion are found to harbor the highest amount of fungi (2.33%) with the surfaces from home being

the lowest (0.06%). In addition, the 2bRAD-M species-level profile unveils distinctive patterns of

fungal composition among the various sites (**Fig 5c**). *Malassezia restricta* and *Malassezia globose*,

known commensals on human skin surface[27], were found as the top abundance fungi at underarm and

most of the indoor environmental samples. It was possible that the *Malassesia* sp. was transferred

from human, given the likely high contact frequency between the skin and these indoor

environmental surfaces. Conversely, *Alternaria alteranta* was mostly found in the indoor

environments but absence in the underarm which is in agreement with existing reports[28, 29]. Thus the

fungi profiles derived from 2bRAD-M are consistent with literature, demonstrating the ability to

reliably profile fungi (simultaneously with bacteria) in low-biomass samples.

Taking advantage of 2bRAD-M's capability in profiling bacterial and fungal species

simultaneously, occurrence-network analysis revealed negative correlation between the human skin

commensal yeast of *Malassezia globosa* (which is associated with Seborrheic Dermatitis [30]) and the

aforementioned *S. epidermidis* (recently proposed as a gatekeeper of healthy skin [31]; Spearman

coefficient of -0.569; **Fig. 5d**). Such landscape-like, species-level correlations for low-biomass

microbiomes can potentially reveal novel bacteria-fungi interactions.

2bRAD-M enables tumor microbiome profiling from FFPE tissue samples

Microbiota in human tumor or blood tissues were recently associated with the types,

developmental stages or chemotherapeutic efficiency of cancer [32-36]. Formalin-fixed,

paraffin-embedded (FFPE) tissue, the gold standard of preserving tumor biopsy specimens [37, 38],

represents a vast, irreplaceable historical clinical resource of enormous value for cancer microbiome

studies[39], however profiling microbiome from FFPE tissues has been challenging due to the low

microbial biomass, high human DNA contamination, severe DNA damage and cross-links by

chemical modification [40]. Therefore 2bRAD-M is an ideal approach to elucidate the microbial

community in FFPE samples. Here, we collected FFPE cervical tissue samples from 15 healthy

controls (H), 15 pre-invasive cancerous (PreC; benign) and 15 invasive cancerous patients (InvaC;

malignant), and subjected these samples to 2bRAD-M sequencing. DNA from the FFPE tissue was

extracted from an area of $3cm^2$ with 4μm thickness. On average 25 ng/ul of DNA in each sample was

detected as smears of size under 500 bp in agarose gels (**Fig. S5**; **Methods**), suggesting extremely

low concentration and highly fragmented nature of DNA (both human and microbes [40]) in FFPE

samples.

The microbiome of the FFPE samples is mostly dominated by bacteria species (n=243) with

minimal fungal species (n=2; **Table S5**). The alpha diversity (Shannon and Simpson index) in

healthy controls (H) is significantly lower than PreC and InvaC (p=0.044, Kruskal test; **Fig. 6a**).

Among the identified bacterial species (**Fig. S5**), samples in the PreC and InvaC groups are

significantly enriched with *Methyloversatilis discipulorum* (p-value=1.2e-5), *Mycobacterium*

tuberculosis (p-value=0.004), *Methyloversatilis universalis* (p-value=1.8e-5), *Ferrovibrio K5*

(p-value=0.001) and *Pseudomonas aeruginosa* (p-value=6.8e-5) (each increase in relative

abundance from H to PreC and to InvaC; **Fig. 6b**). Conversely, *Lactobacillus* spp. (*L. paracasei*, *L.*

vaccinostercus,, L. pentosus and *L. plantarum*) are greatly enriched in H (average abundance of

63.1%, in contrast to 33.9% and 32.9% in PreC and InvaC respectively, **Fig. 6b**); this is consistent

with a previous study[36] showing depletion of the *Lactobacillales* genera in "fresh", non-FFPE

cervical cancerous tissues. Notably, *Lactobacillus paracasei* shows anticancer potential against

cervix cancer cells (HeLa) *in vitro*[41]. Furthermore, the enrichment of *Mycobacterium tuberculosis*,

Pseudomonas aeruginosa and *Staphylococcus aureus* in PreC- and InvC-phase PPFE samples is

consistent with "fresh" tissue based studies that reported related genus-level (via 16S [32]) or

species-level (via WMS [36, 42]) taxa. Taken as a whole, 2bRAD-M has successfully captured the

microbiome structure in FFPE samples, and is able to reveal previously unknown discriminative

microbial features between healthy and cancerous tissues. These features may serve as potential

indicative novel markers for tumor onset and progression diagnosis. To evaluate this potential, we

applied Random Forest on the FFPE taxonomic profiles at the species-level, and the model

distinguishes H, PreC and InvaC samples with 91.1% accuracy (ten-fold cross validation; **Fig. 6c**).

Notably, we can achieve maximized discriminative performance (AUC: 0.96) with the fewer

features of the nine most important species in the RF model (**Fig. 6d**; **Table S7**). Thus, 2bRAD-M

offers a viable option for microbiome profiling on the vast archive of historical FFPE sample with

potential application in early diagnosis and treatment of cancer.

Discussion

Using mock microbiomes plus actual samples from stool, skin, environment-surface and frozen

FFPE tissues, we demonstrated the ability of 2bRAD-M to profile microbiome samples that are

challenging by using marker-gene or WMS sequencing approaches. Three key features are

highlighted from this research. *Firstly*, it can analyze samples with low biomass (down to 1 pg),

severe degradation or high contamination. This advantage is built on the ability to (*i*) provide an

unbiased (i.e., without the large size variation of DNA fragments) and reduced representation of

metagenomes for sequencing, (*ii*) evenly cover all restriction sites, and (*iii*) generate short, iso-length

reads and thus reliable quantification of tag abundance. With such advantage, 2bRAD-M is valuable

to profiling microbiome samples from sparsely populated niches (e.g., indoor surfaces, blood and

skin), precious clinical specimen or heavily degraded or contaminated tissues (e.g., FFPE sections

and archaeological samples). Notably cervical cancer can be effectively cured in the preinvasive

period [43], yet detection of this asymptomatic stage remains challenging which results in unnecessary

delays in diagnosis and treatment [44]; therefore, 2bRAD-M sequencing of FFPE cervical cancer

microbiomes, which is challenging for WMS or 16S rRNA approaches, promises to turn the

enormous clinical repositories of FFPE tissue specimens into a treasure trove of microbiome-driven

discoveries [37].

Secondly, it can provide species-level taxonomic information for bacteria, archaea and fungi

simultaneously. The multiple species-specific 2bRAD-M tags can uniquely identify the species

from the microbial community, as resolution at the "species" level instead of the "genus" level can

be crucial [45]. In this case, the ability to extract the species-level cancer-stage-specific markers from

FFPE tissues, as well as to distinguish *S. epidermidis*, *S. hominis* and *S. aureus* from underarm skin

samples, are crucial to diagnosis and mechanistic understanding of inflammation [24]. Interestingly,

the overall ratio of fungal and bacterial abundance, a potential indicator of ecological balance[46], is

0.31% for underarm, 0.07% for home and 1.27% for car (significantly different), suggesting

characteristic ecological features among the sites.

Finally, using 2bRAD-M, the species-level taxonomic profiling can be achieved with a much

lower sequencing cost than WMS. For example, only 1~5% of the sequencing data of WMS are

required by 2bRAD-M to produce a taxonomic profile of equivalent accuracy, resulting in a cost

reduction of 20-100 fold. Moreover, multi-isoRAD allows sequencing of five concatenated 2bRAD

tags altogether via a single, 100–150 bp read from Illumina paired-end (PE) library [14], which further

reduces the sequencing cost. The cost advantage is significant when high sequencing depth is

required to address questions like unveiling the true microbial diversity or recovering rare but

functionally important species in complex microbiota [20, 47]. Thus 2bRAD-M appears especially

suitable for such samples or circumstances.

Notably, the performance of 2bRAD-M is limited by the availability and the potential bias of

related reference genomes (e.g., 171927 bacterial and 945 archaeal genomes, yet only 293 fungal

genomes in RefSeq), although the situation is rapidly improving with the steady increase of

sequenced genomes for both cultured and uncultured microbes (e.g. via single-cell sequencing [48]).

With the key advantages versus marker genes analysis and WMS, 2bRAD-M is particularly suitable

for cost-effectively deriving a species-resolution taxonomic profiling for samples with low biomass

(down to 1 pg), severe degradation or high contamination, thus greatly expands the opportunities in

microbiome research.

Materials and Methods

Sample preparation and sequencing

Two mock microbial communities were used to validate the stability, sensitivity and precision

of 2bRAD-M. The first consists of five evenly mixed bacterial strains including *Streptococcus*

mutans UA159, *Streptococcus agalactiae* ATCC13813, *Staphylococcus aureus* ATCC29213,

Escherichia coli DH5α and *Lactobacillus fermentum* ATCC9338. Then three circumstances

that simulated "challenging" microbiome samples were produced (with three replicates for each

sample): (1) LoA: samples with low amount DNA; A concentration gradient from 10ng to 1pg was

designed, with one tenth of the concentration retained each time. (2) HiD: samples with highly

degraded DNA; Two samples with DNA length about 150bp and 50bp were included. (3) HoC:

samples with host DNA contamination, which consist of either 90% or 99% human DNA.

The second mock used in this paper is the 20-Strain Even Mix Genomic Material (named as Mock

MSA1002, 3.87 ng/uL; frozen 50 μL in Tris-HCl pH 8.5) which is purchased from ATCC. This mock

comprises genomic DNA prepared from fully sequenced, characterized, and authenticated ATCC

Genuine Cultures that were selected based on relevant phenotypic and genotypic attributes, such as

Gram stain, GC content, genome size and spore formation.

Fecal and FFPE tissue samples The study protocol complies with the ethical guidelines of the

1975 Declaration of Helsinki and was approved by the DOST ethical committee and conducted

according to ICH guidelines for Good Clinical Practice. Written informed consent and photography

consent were obtained from each subject before enrollment. Three healthy adult individuals were

enrolled as volunteers and fecal samples were collected for deep WMS, 16S rRNA gene amplicon

and 2bRAD-M sequencing for comparison.

Totally 45 cervical cancer-related sections in FFPE blocks (with a thickness of 4μm and an area

of 3cm^2) underwent microbiome profiling via 2bRAD-M. The original tissue samples include 15

healthy, 15 cancerous and 15 severer cancerous ones (determined by morphological evidence of

polymorphonuclear infiltration). Each of the 45 samples was from a distinct individual, i.e., a

cross-sectional design. To prepare the FFPE blocks, fresh tissue samples were fixed in phosphate

buffered formalin for 24–48 h, followed by tissue processing for 10 h, and paraffin embedding for

20 min [49]. Post-fixation processing of the tissues was completed in a histopathological laboratory

using consistent processor protocols over years. Prior to microbiome profiling, the FFPE blocks were

already stored at 17 °C–22 °C and 20%–60% humidity levels for 1-2 years. For these FFPE tissue

sections, attempts to construct 16S amplicon or WMS libraries for microbiome sequencing both

failed, due to the low quality of initial DNA (**Fig. S6**).

Underarm, home and car sampling collection protocol & processing For the underarm

sampling protocol, individuals were subjected to a washout period where the use of anti-bacterial

products were not allowed for 4-5 days. After the washout period, 22mm D-squame tape strip

(Cuderm) was applied onto the lower underarm skin surface (without hair) using a pressure

applicator. The tape strips were then pre-treated with 200 mg of 0.1 mm Zirconia Silica beads and

bead beat using Qiagen TissueLyser II (Valencia, CA) at 30 Hz for 3 minutes to lyse (via

mechanical force) and dislodge the biomass from the tape strip. After dislodging the biomass from

tape strip, complete cells lysis is achieved via further enzymatic reaction by 2% w/v lysozyme, 0.05%

w/v lysostaphin and 1.2% Triton-X in TE buffer, followed by DNA extraction using Qiagen

DNeasy Blood and Tissue Kit (Cat no. 69504, Valencia, CA). For the home and car sampling, swab

was used for wiping the surface of cushion and carpet in car, and toy and toilet seat in home.

Specifically, wipe the swab for 20 times on an area of $4 \times 4cm^2$. After collection, transfer the

disposable swab to the tube containing sample storage liquid immediately, and break the swab

along the crease. Then close the lid of the tube and make sure that there is no leakage. Finally, put

the sample preservation solution tube in the biosafety bag for DNA extraction.

DNA extraction, 16S rRNA and shotgun metagenomic sequencing Genomic DNA was

extracted from each fecal containing tube using the Tissue and Blood DNA Isolation kit (Qiagen,

Valencia, CA) following the manufacturer's instructions with slight modifications. PCR

amplification of the V1-V3 hypervariable regions of 16S rRNA genes was performed using the

primer set (27F/534R) and followed the protocol developed by the Human Microbiome Project. PCR

amplification reactions in triplicate for each sample were pooled at approximately equal amounts and

sequenced, via the Illumina MiSeq 250 platform. All sequences were pre-processed following the

standard QIIME (v.1.9) pipeline. Downstream bioinformatics analysis was performed using

Parallel-Meta 3, a software package for comprehensive taxonomic and functional comparison of

microbial communities. Clustering of OTUs was conducted at the 97% similarity level using a

pre-clustered version of the Refseq database.

Paired-end metagenomic sequencing was performed for the intestinal microbiota from six

individuals via the Illumina HiSeq 2500 platform, yielding 239.73 ± 12.34 GB per sample

(paired-end reads, with average fragment insert size of 350 bp and average read length of 150 bp).

The reads were quality controlled by Trimmomatic (Sliding window 4:20; Minlength:100;

MinPhred:25; Percentage of MinPhred: 80), and finally 429,016,382 ± 6,570,335 clean reads per

sample were generated, and then profiled by mOTUs2 using default parameters.

2bRAD-M sequencing The 2bRAD-M library preparation basically followed the original

protocol developed by Wang et al. (2012) [15] with minor modifications. Library preparation began

with the digestion of 1pg–200 ng genomic DNA in a 15-μl reaction using 4 U BcgI (NEB) at 37 ℃

for 3 h. 5ul of digested product was run on a 1% agarose gel to verify digestion. Next, ligation

reaction was performed at 4 ℃ for 16 h in a 20 μl volume containing 10 μl of digested product, 0.2

μM each of library-specific adaptors (Ada1 and Ada2), 1 mM ATP (NEB), 1×T4 DNA Ligase

Buffer and 800 U T4 DNA ligase (NEB). Then heat inactivation was performed for BcgI at 65 ℃

for 20 min.

Ligation products were amplified in 40-μl PCRs, each composed of 7 μl ligated DNA, 0.1 μM

each primer (Primer1 and Primer2 for Illumina), 0.3 mM dNTP, 1× Phusion HF buffer and 0.4 U

Phusion high fidelity DNA polymerase (NEB). PCR was conducted in a DNA Engine Tetrad 2

thermal cycler (Bio-Rad) with 16-28 cycles of 98 ℃ for 5 s, 60 ℃ for 20 s and 72 ℃ for 10 s and

then a final extension of 10 min at 72 ℃. The target band (Illumina: ~100 bp) was excised from 8%

(wt/vol) polyacrylamide gel, and the DNA diffused from the gel into nuclease-free water for 12 h at

4 ℃. Finally, barcodes were introduced by means of PCR with platform-specific barcode-bearing

primers. 40-μl PCR reaction contained 50 ng of gel-extracted PCR product, 0.2 μM of each primer

(Primer1 and Primer3 for Illumina), 0.6 mM dNTP, 1× Phusion HF buffer and 0.8 U Phusion

high-fidelity DNA polymerase; seven cycles of the PCR profile listed above were performed. PCR

products were purified by QIAquick PCR purification kit (Qiagen, Valencia, CA) and subjected to

Illumina HiSeq platform sequencing. All primer and adaptor sequences are provided at **Table S8**.

Data analysis

Identification of species-specific 2bRAD-M markers from the most comprehensive genome

database Firstly, totally 173,165 microbial genomes (including bacteria, fungi and archaea) were

downloaded from NCBI RefSeq database. Then, built-in Perl scripts (GitHub:

https://github.com/shihuang047/2bRAD-M) were used to sample restriction fragments from

microbial genomes by each of 16 type 2B restriction enzymes, which formed a huge 2bRAD

microbial genome database. The set of 2bRAD tags sampled from each genome was assigned under

the GCF number, as well as GCF's taxonomic information corresponding to the whole genome.

Finally, all 2bRAD tags from each GCF that occur once within the genome were compared with those

to all the others. Those 2bRAD tags are specific to a species-level taxon (having no overlap with other

species') were developed as species-specific 2bRAD markers, collectively forming a 2bRAD marker

database.

Simulation of 2bRAD-M sequencing data We designed a complex microbial composition

containing 50 bacterial species from various niches such as oral, intestinal and environmental. Given

a specified species composition, their genomes were merged into one meta-genome file accordingly

with one complete genome as the smallest operational unit. A built-in read simulator was then used to

produce about 10 million 32-bp reads for each sample from this merged genome using a SNP rate of

0.1% and an indel rate of 0.1%.

Calculation of relative abundance Firstly, to identify microbial species within each sample, all

sequenced 2bRAD tags after quality control were mapped (using a built-in Perl script) against the

2bRAD marker database which contains all 2bRAD tags theoretically unique to each of 26,163

microbial species in RefSeq database (**Fig. 1**). To control the false-positive in the species

identification, G score was derived for each species identified within a sample as below, which is a

harmonious mean of read coverage of 2bRAD markers belongs to a species and number of all

possible 2bRAD markers of this species. The threshold of G score for a false positive discovery of

microbial species was set 5.

$$\text{G score}_{species\ i} = \sqrt{S_i \times t_i}$$

S: the number of reads assigned to all 2bRAD markers belonging to species *i* within a sample.

t: number of all 2bRAD markers of species *i* that have been sequenced within a sample

We estimated the relative abundance of each microbial species in a sample using the formula as

below. We first calculated the average read coverage of all 2bRAD markers for each species, which

represent the number of individuals belonging to a species present in a sample at a given sequencing

depth. The relative abundance of a given species is then calculated as the ratio of the number of

microbial individuals belonging to a species against the total number of individuals from known

species that can be detected within a sample, the default G score is 5 .

$$\text{Relative abundance}_{species\ i} = \frac{S_i/T_i}{\sum_{i=1}^{n} S_i/T_i}$$

S: the number of reads assigned to all 2bRAD markers of species *i* within a sample

T: the number of all theoretical 2bRAD markers of species i

Calculation of Precision, Recall and L2 similarity For overall performance assessment, we

applied precision and recall to evaluate the accuracy of microbial identification, while L2 distance[16]

was employed to evaluate the accuracy of abundance estimation within a sample. Precision is the

proportion of true positive species against the total number of species identified by a method, whereas

recall is defined as the proportion of true positive species against the number of all species actually

existed in a sample. To evaluate the accuracy of abundance profiles, we calculate the L2 distance

between ground-truth abundance profile to each of taxonomic profiles at a given taxonomic level

(e.g., species or genus) produced by metagenomic sequencing methods. To visualize the

benchmarking results more intuitively, we further employed L2 similarity calculated as 1-L2 distance

for performance comparisons.

Diagnosis model of cancer samples Random forest models were trained to identify cancer status

using the taxonomy profiles on the species level. Default parameters of the R implementation of

algorithm were applied (R package 'randomForest', ntree=5,000, using default mtry of p/3 where p is

the number of input taxa). The performance of RF models based on microbiota was evaluated with a

ten-fold cross-validation approach.

Additional details are provided in **Supplemental Methods**.

Acknowledgments

We thank Dr. Rob Knight and Dr. Yoshiki Vázquez-Baeza for the helpful discussion on

technical issues. This work was funded by Grant 31800088 from National Natural Science

Foundation and 2019M652501 from China Postdoctoral Science Foundation. S. W. acknowledges

the support by Taishan Scholar Fund of Shandong Province of China.

Code availability

The computational pipeline can be downloaded at Github:

https://github.com/shihuang047/2bRAD-M.

Competing interest

Rongchao Zhang, Lisha Zhou, Qianya Niu, Xiuping Wang are affiliated to Qingdao OE

Biotechnology Company. Lam Tzehau, Helen Zhao and Jiquan Liu are affiliated with Procter &

Gamble Company.

509
510

References

1.   Knight, R. et al. Best practices for analysing microbiomes. *Na Rev Microbiol* **16**, 410-422
(2018).
2.   Cho, I. & Blaser, M.J. The human microbiome: at the interface of health and disease. *Nat*
Rev Genet **13**, 260-270 (2012).
3.   Franzosa, E.A. et al. Sequencing and beyond: integrating molecular 'omics' for microbial
community profiling. *Nat Rev Microbiol* **13**, 360-372 (2015).
4.   Quince, C., Walker, A.W., Simpson, J.T., Loman, N.J. & Segata, N. Shotgun metagenomics,
from sampling to analysis (vol 35, pg 833, 2017). *Nat Biotechnol* **35**, 1211-1211 (2017).
5.   Davey, J.W. et al. Genome-wide genetic marker discovery and genotyping using
next-generation sequencing. *Nat Rev Genet* **12**, 499-510 (2011).
6.   Andrews, K.R., Good, J.M., Miller, M.R., Luikart, G. & Hohenlohe, P.A. Harnessing the
power of RADseq for ecological and evolutionary genomics. *Nat Rev Genet* **17**, 81-92
(2016).
7.   Grewe, F., Huang, J.-P., Leavitt, S.D. & Lumbsch, H.T. Reference-based RADseq resolves
robust relationships among closely related species of lichen-forming fungi using
metagenomic DNA. *Sci Rep* **7** (2017).
8.   Liu, M. et al. Evaluation of ddRAD seq for reduced representation metagenome sequencing.
PeerJ **5** (2017).
9.   Holmes, I. & Rabosky, A.R.D. Natural history bycatch: a pipeline for identifying
metagenomic sequences in RADseq data. *PeerJ* **6** (2018).
10.   Hess, M.K. et al. A restriction enzyme reduced representation sequencing approach for
low-cost, high-throughput metagenome profiling. *PloS One* **15**, e0219882 (2020).
11.   Davey, J.W. et al. Special features of RAD Sequencing data: implications for genotyping.
Mol Ecol **22**, 3151-3164 (2013).
12.   Aird, D. et al. Analyzing and minimizing PCR amplification bias in Illumina sequencing
libraries. *Genome Biol* **12**, R18 (2011).
13.   Wang, S. et al. MethylRAD: a simple and scalable method for genome-wide DNA
methylation profiling using methylation-dependent restriction enzymes. *Open Biol* **5** (2015).
14.   Wang, S. et al. Serial sequencing of isolength RAD tags for cost-efficient genome-wide
profiling of genetic and epigenetic variations. *Nat Protoc* **11**, 2189-2200 (2016).
15.   Wang, S., Meyer, E., McKay, J.K. & Matz, M.V. 2b-RAD: a simple and flexible method for
genome-wide genotyping. *Nat Methods* **9**, 808-810 (2012).
16.   Ye, S.H., Siddle, K.J., Park, D.J. & Sabeti, P.C. Benchmarking Metagenomics Tools for
Taxonomic Classification. *Cell* **178**, 779-794 (2019).
17.   Kim, D., Song, L., Breitwieser, F.P. & Salzberg, S.L. Centrifuge: rapid and sensitive
classification of metagenomic sequences. *Genome Res* **26**, 1721-1729 (2016).
18.   Amos, G.C.A. et al. Developing standards for the microbiome field. *Microbiome* **8**, 98
(2020).
19.   Hillmann, B. et al. Evaluating the Information Content of Shallow Shotgun Metagenomics.
mSystems **3**, e00069-e00018 (2018).

20. Zaheer, R. et al. Impact of sequencing depth on the characterization of the microbiome and
resistome. *Sci Rep* **8**, 5890 (2018).
21. de Goffau, M.C. et al. Recognizing the reagent microbiome. *Nat Microbiol* **3**, 851-853
(2018).
22. Lam, T.H. et al. Understanding the microbial basis of body odor in pre-pubescent children
and teenagers. *Microbiome* **6**, 213 (2018).
23. James, A.G., Austin, C.J., Cox, D.S., Taylor, D. & Calvert, R. Microbiological and
biochemical origins of human axillary odour. *FEMS Microbiol Ecol* **83**, 527-540 (2013).
24. Nakatsuji, T. et al. Antimicrobials from human skin commensal bacteria protect against
Staphylococcus aureus and are deficient in atopic dermatitis. *Sci Transl Med* **9**, eaah4680
(2017).
25. Lax, S. et al. Longitudinal analysis of microbial interaction between humans and the indoor
environment. *Science* **345**, 1048-1052 (2014).
26. Adams, R.I., Miletto, M., Taylor, J.W. & Bruns, T.D. The Diversity and Distribution of
Fungi on Residential Surfaces. *PloS One* **8**, e78866 (2013).
27. Saunders, C.W., Scheynius, A. & Heitman, J. Malassezia Fungi Are Specialized to Live on
Skin and Associated with Dandruff, Eczema, and Other Skin Diseases. *PLOS Pathog* **8**,
e1002701 (2012).
28. Rocchi, S. et al. Indoor Microbiome: Quantification of Exposure and Association with
Geographical Location, Meteorological Factors, and Land Use in France. *Microorganisms* **8**,
341 (2020).
29. Sharpe, R.A., Thornton, C.R., Tyrrell, J., Nikolaou, V. & Osborne, N.J. Variable risk of
atopic disease due to indoor fungal exposure in NHANES 2005–2006. *Clin Exp Allergy* **45**,
1566-1578 (2015).
30. Greaves, M.W. & Camp, R.D. Prostaglandins, leukotrienes, phospholipase, platelet
activating factor, and cytokines: an integrated approach to inflammation of human skin.
Arch Dermatol Res **280 Suppl**, S33-41 (1988).
31. Gallo, R.L. S. epidermidis influence on host immunity: more than skin deep. *Cell Host*
Microbe **17**, 143-144 (2015).
32. Leinwand, J.C. & Miller, G. Microbes as biomarkers and targets in pancreatic cancer. *Nat*
Rev Clin Oncol **16**, 665-666 (2019).
33. Geller, L.T. et al. Potential role of intratumor bacteria in mediating tumor resistance to the
chemotherapeutic drug gemcitabine. *Science* **357**, 1156-+ (2017).
34. Guglielmi, G. GUT MICROBES JOIN THE FIGHT AGAINST CANCER. *Nature* **557**,
482-484 (2018).
35. Nejman, D. et al. The human tumor microbiome is composed of tumor type-specific
intracellular bacteria. *Science* **368**, 973-980 (2020).
36. Klein, C. et al. Relationship between the Cervical Microbiome, HIV Status, and
Precancerous Lesions. *mbio* **10** (2019).
37. Poore, G.D. et al. Microbiome analyses of blood and tissues suggest cancer diagnostic
approach. *Nature* **579**, 567-574 (2020).
38. Walker, S.P., Tangney, M. & Claesson, M.J. Sequence-Based Characterization of
Intratumoral Bacteria-A Guide to Best Practice. *Front Oncol* **10** (2020).

39. Pinto-Ribeiro, I. et al. Evaluation of the Use of Formalin-Fixed and Paraffin-Embedded
Archive Gastric Tissues for Microbiota Characterization Using Next-Generation Sequencing.
Int J Mol Sci **21**, 1096 (2020).

40. Do, H. & Dobrovic, A. Sequence Artifacts in DNA from Formalin-Fixed Tissues: Causes
and Strategies for Minimization. *Clin Chem* **61**, 64-71 (2015).

41. Riaz Rajoka, M.S. et al. Anticancer potential against cervix cancer (HeLa) cell line of
probiotic Lactobacillus casei and Lactobacillus paracasei strains isolated from human breast
milk. *Food Funct* **9**, 2705-2715 (2018).

42. Johansen, I.S., Thomsen, V.O., Forsgren, A., Hansen, B.F. & Lundgren, B. Detection of
Mycobacterium tuberculosis complex in formalin-fixed, paraffin-embedded tissue
specimens with necrotizing granulomatous inflammation by strand displacement
amplification. *J Mol Diagn* **6**, 231-236 (2004).

43. Patel, A. Benign vs Malignant Tumors. *JAMA Oncol 6, 1488* (2020).

44. Fontaine, P.L., Saslow, D. & King, V.J. ACS/ASCCP/ASCP guidelines for the early
detection of cervical cancer. *Am Fam Physician* **86**, 501, 506-507 (2012).

45. Reese, A.T. & Dunn, R.R. Drivers of Microbiome Biodiversity: A Review of General Rules,
Feces, and Ignorance. *mBio* **9**, e01294-18 (2018).

46. Wagg, C., Schlaeppi, K., Banerjee, S., Kuramae, E.E. & van der Heijden, M.G.A.
Fungal-bacterial diversity and microbiome complexity predict ecosystem functioning. *Nat*
Commun **10**, 4841 (2019).

47. Pereira-Marques, J. et al. Impact of Host DNA and Sequencing Depth on the Taxonomic
Resolution of Whole Metagenome Sequencing for Microbiome Analysis. *Front Microbiol*
10, 1277 (2019).

48. Lasken, R.S. & McLean, J.S. Recent advances in genomic DNA sequencing of microbial
species from single cells. *Nat Rev Genet* **15**, 577-584 (2014).

49. Rahmadwati, Naghdy, G., Ros, M. & Todd, C. in Proceedings of the 2011 2nd International
Congress on Computer Applications and Computational Science. (eds. F.L. Gaol & Q.V.
Nguyen) 235-243 (Springer Berlin Heidelberg, Berlin, Heidelberg; 2012).

Tables

Table 1. High concordance of dominant microbial taxa in 16S rRNA and 2bRAD-M profiling

results of 20 underarm samples.

Top genus in 16S	Relative abundance	Rank	Corresponding species in 2bRAD-M	Relative abundance	Rank
Staphylococcus	44.43%	1	*Staphylococcus epidermidis*	27.56%	1
			Staphylococcus hominis	12.57%	2
			Staphylococcus capitis	4.78%	5
			Staphylococcus haemolyticus	1.78%	11
			The other 4 species	0.48%	
			SUM	47.17%	
Corynebacterium	15.06%	2	*Corynebacterium ureicelerivorans*	2.58%	8
			Corynebacterium jeddahense	2.41%	9
			Corynebacterium aurimucosum	1.80%	10
			Corynebacterium pseudogenitalium	1.29%	15
			Corynebacterium kefirresidentii	1.23%	17
			Corynebacterium tuberculostearicum	1.08%	19
			The other 47 species	2.08%	
			SUM	10.38%	
Uruburuella	5.67%	3	*Uruburuella suis*	5.80%	4
Enterococcus	4.49%	4	*Enterococcus devriesei*	1.68%	12
			Enterococcus dispar	1.48%	13
			The other 3 species	0.45%	
			SUM	3.61%	
Moraxella	02.34%	5	*Moraxella osloensis*	2.77%	7
			The other 1 species	0.04%	
			SUM	2.81%	
SUM	**72.00%**		**SUM**	**63.97%**	

Figures

Figure 1. Scheme of the 2bRAD-M workflow. In the library preparation module of 2bRAD-M

pipeline, DNA samples were first digested using a type IIB restriction enzyme. The resulting 2bRAD

fragments were enriched and amplified for DNA sequencing. In the computational module of

2bRAD-M pipeline, we employed a prebuilt 2bRAD marker database to perform taxonomic profiling

on 2bRAD data. The resulting 2bRAD-M data were used for microbial identification and relative

abundance calculation. Firstly, all reads were mapped against the default unique 2bRAD marker

database to identify all candidate species in a 2bRAD-M sample. Next, to accurately estimate the

abundance of identified species, we increased the number of taxa-specific 2bRAD markers for each

candidate species by reconstructing a reduced 2bRAD marker database. A reduced 2bRAD marker

database contains more 2bRAD markers specific to each candidate species than those in the default

database. All the 2bRAD sequences were then remapped to this reduced marker database for

abundance estimation of candidate species. In principal, the relative abundance of a given species was

calculated as the read coverage of all species-specific 2bRAD markers. For more information, please

refer to **Methods**.

Figure 2. **Benchmark measurements of 2bRAD-derived taxonomic profile.** Simulated microbial

community data consisting of 50 microbes were profiled by each of the 16 Type IIB restriction

enzymes. The scatter plots indicate the correlation of the taxonomic abundance estimated from

2bRAD-M with the expected abundance for each enzyme. Percentage number indicated in each plot

represent the average genome coverage (50 microbes) by 2bRAD-M reads.

Figure 3. Comparison of taxonomic profiling results of 2bRAD-M, shotgun WGS and 16S rRNA sequencing methods in mock microbial communities. (a) 2bRAD-M performance in samples with low amount (LoA), high DNA degradation (HiD) and high host DNA contamination (HoC) based on mock community of five bacteria species (Mock-CAS). LoA includes samples with mixture of human DNA (90% and 99%) and bacterial DNA; HiD includes samples with fragmented bacterial DNA (50bp or 150bp). HoC includes samples with different DNA concentrations (50 ng, 20 ng, 10 ng, 1 ng, 0.1 ng, 0.01 ng, and 0.001 ng). Three technical replicates were included for each group. The L2 score that measures the microbial composition similarity between 2bRAD-M and WMS is shown on the head of each stacked bar plot. (b) Benchmark of WMS, 16S rRNA and 2bRAD-M approaches based on ATCC MSA1002 mock community. Each stacked bar plot in the left panel shows the resultant taxonomic profile of a given metagenomic method prepared at a specified

665   DNA amount. The left panel indicates the corresponding L2 similarity, precision and recall score as

666   compared to the ground truth.

Figure 4. Comparison of taxonomic profiles and desired sequencing depth of 2bRAD-M, 16S and WMS in fecal samples. (**a-c**) Comparison of taxonomic profiles at the genus level between 16S rRNA and 2bRAD-M. (**d-f**) Comparison of taxonomic profiles at the species level between WMS and 2bRAD-M. (**g**) The rarefaction analysis of 2bRAD-M and WMS samples. The species-level compositions in the subsampled data at each given sequencing depth were compared to the pre-rarefaction result for each method.

Figure 5. 2bRAD-M analysis of low biomass samples collected from underarm, home and car

surfaces. (a) Relative abundance of bacteria at genus-level (>1%) identified in underarm by 16S

rRNA sequencing (left panel) and 2bRAD-M (right panel) with the samples stratified by cluster

group. (**b**) Relative abundance of bacteria at species-level taxa (>1%) identified in the underarm by

2bRAD-M with the samples stratified by cluster group. (**c**) Heat map shows identified fungal species

682 via 2bRAD-M in the underarm, home and car samples. (**d**) Co-occurrence network of bacterial and

683 fungal species based on 20 underarm samples profiled by 2bRAD-M. Each circle represents a

684 species and its size refers to the degree centrality score. Blue/red edges indicate positive/negative

685 spearman coefficients.

Figure 6. Microbiome-based diagnosis of FFPE thin section from cervical cancer samples as

enabled by 2bRAD-M. (**a**) Shannon and Simpson index among the groups. (**b**) Comparison of

differential species and *Lactobacillus spp.* among the groups. (**c**) The Random Forest classifier for

discriminating cancer and healthy samples. In the ternary plot, each dot represents a FFPE sample.

The axes indicate the microbiome-based probability of severer cervical cancer, cervical cancer and

health on a FFPE sample. The closer one sample is to an apex, the more likely it is predicted as to be

corresponding disease states (i.e., severer cervical cancer, cervical cancer and health). (**d**) Feature

selection by rebuilding Random Forest classifiers using a series of reduced sets of features. The

scatter plot shows that nine variables (species) in a reduced RF model (i.e., the AUC plot on the right)

can maximize model performance.