A multi-omics dataset for the analysis of Frontotemporal Dementia genetic subtypes

Authors

Kevin Menden[1,*] (KM); Margherita Francescatto[1] (MF); Tenzin Nyima[1] (TN); Cornelis Blauwendraat[1,2] (CB); Ashutosh Dhingra[1] (AD); Melissa Castillo-Lizardo[1] (MC); Noémia Fernandes[1] (NF); Lalit Kaurani[4] (LK); Deborah Kronenberg-Versteeg[1,3] (DKV); Burcu Atasu[1,3] (BA); Eldem Sadikoglou[1] (ES); Barbara Borroni[6] (BB); Salvador Rodriguez-Nieto[1] (SRN); Javier Simon-Sanchez[1,3] (JSS); Andre Fischer[4] (AF); David Wesley Craig[7] (DWC); Manuela Neumann[1] (MN); Stefan Bonn[1,5] (SB); Patrizia Rizzu[1] (PR); Peter Heutink[1,*] (PH).

Affiliations

German Center for Neurodegenerative Diseases, Tübingen, Germany
National Institute on Aging, National Institutes of Health, Bethesda, MD, USA
Hertie Institute for Clinical Brain Research, Tübingen, Germany
German Center for Neurodegenerative Diseases, Göttingen, Germany
Institute of Medical Systems Biology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany
Neurology Unit, Department of Clinical and Experimental Sciences, University of Brescia, Italy
University of Southern California, California, USA
corresponding author(s): Peter Heutink (peter.heutink@dzne.de)

Abstract

Understanding the molecular mechanisms underlying frontotemporal dementia (FTD) is essential for the development of successful therapies. Systematic studies on human post-mortem brain tissue of patients with genetic subtypes of FTD are currently lacking. The Risk and Modyfing Factors of Frontotemporal Dementia (RiMod-FTD) consortium therefore has generated a multi-omics dataset for genetic subtypes of FTD to identify common and distinct molecular mechanisms disturbed in disease. Here, we present multi-omics datasets generated from the frontal lobe of post-mortem human brain tissue from patients with mutations in MAPT, GRN and C9orf72 and healthy controls. This data resource consists of four datasets generated with different technologies to capture the transcriptome by RNA-seq, and small RNA-seq, and supplemented this date with CAGE-seq, and methylation profiling. We show concrete examples on how to use the resulting data and confirm current knowledge about FTD and identify new processes for further investigation. This extensive multi-omics dataset holds great value to reveal new research avenues for this devastating disease.

Background & Summary

Frontotemporal Dementia (FTD) is a devastating pre-senile dementia characterized by progressive deterioration of the frontal and anterior temporal lobes[1]. The most common symptoms include severe changes in social and personal behaviour as well as a general blunting of emotions. Clinically, genetically, and pathologically there is considerable overlap with other neurodegenerative diseases including Amyotrophic Lateral Sclerosis (ALS), Progressive Supranuclear Palsy (PSP) and Cortical Basal Degeneration (CBD)[2]. Research into FTD has made major advances over the past decades. Up to 40% of cases[3] have a positive family history and up to 60% of familial cases can be explained by mutations in the genes Microtubule Associated Protein Tau (MAPT), Granulin (GRN) and C9orf72[4] which has been key to the progress in our understanding of its molecular basis. Several other disease-causing

genes have been identified that account for a much smaller fraction of cases[5]. Mutations in MAPT lead to accumulation of the Tau protein in neurofibrillary tangles in the brain of patients while mutations in GRN and C9orf72 lead to the accumulation of TDP-43[6], as well as dipeptide repeat proteins (DPRs) and RNA foci in the case of C9orf72[7].

As of today, no therapy exists that halts or slows the neurodegenerative process of FTD and to develop successful therapies there is an urgent need to determine whether a common target and therapy can be identified that can be exploited for all patients, or whether the distinct genetic, clinical, and pathological subgroups need tailored treatments. Therefore, the development of remedies relies heavily on a better understanding of the molecular and cellular pathways that drive FTD pathogenesis in all FTD subtypes.

Although our knowledge of FTD pathogenesis using molecular and cellular biology approaches has significantly advanced during recent years, a deep mechanistic understanding of the pathological pathways requires simultaneous profiling of multiple regulatory mechanisms.
Post-mortem human brain tissue is an important source for studying the disturbance of molecular processes in patients with FTD. However, systematic studies on human post-mortem brain tissue with genetic subtypes of FTD are currently lacking. Therefore, the Risk and modifying factors in Frontotemporal Dementia (RiMod-FTD) consortium has generated a multi-omics data resource with the focus on mutations in the three most common causal genes for FTD: MAPT, GRN and C9orf72.

Here, we report four datasets from the frontal lobe of post-mortem human brain from RiMod-FTD samples. We present RNA-seq, CAGE-seq, smRNA-seq, and methylation datasets from matched samples, which enable precise profiling of transcriptional dysregulations in genetic FTD subtypes. The RNA-seq dataset can be used to identify general transcriptional differences in the FTD subtypes as well as potential alternative splicing events. The regulation of the transcriptome can be studied using the other three datasets. CAGE-seq enables the detection of active or inactive promoters and thereby helps to pinpoint potentially disease-relevant transcription factors. Methylation and smRNA-seq datasets allow researchers to identify regulatory mechanisms that lead to down-regulation of certain genes. These different aspects of the transcriptome can be studied in detail due to matched samples for all these datasets.

We show that known differences and commonalities of the investigated FTD subtypes can be recapitulated with initial analyses of the datasets, such as for instance neuronal loss due to neurodegeneration. More importantly, interesting new aspects about these FTD subtypes can be easily obtained from this dataset. We therefore believe that a thorough analysis of these data from the RiMod-FTD project can reveal helpful new insights about FTD and believe that researchers will find these datasets helpful for their own studies. To make the data easily accessible, we provide a gene browser which allows to compare expression and methylation information between disease groups for specific genomic locations at www.rimod-ftd.org.

96
Figure 1: Schematic overview of the RiMod-FTD project idea. a) Samples from post-mortem
human brain tissue of patients with FTD caused by mutations in MAPT, GRN and C9orf72 and
from healthy controls are collected. Multi-omics profiling is performed on all samples to gain
detailed insights into disease mechanisms. Integrative data analysis is performed to gain new
insights and the data is made available for further studies. The goal of the RiMod-FTD project
is to further extend this data resource with fitting datasets.
103

Assay	Control	FTD-MAPT	FTD-GRN	FTD-C9orf72
RNA-seq	16	11	7	13
smRNA-seq	12	10	8	12
CAGE-seq	16	11	8	13
Methylation	14	13	7	13

104 **Table 1**: Sample numbers of the different disease groups for the available assays.
105

Methods

107
Donor samples employed in this study
Tissues were obtained under a Material Transfer Agreement from the Netherlands Brain
Bank, and additional samples were provided by the Queen Square Brain Bank of Neurological
Disorders and MRC, King College London.
GFM and GTM tissue from each subject was divided into three pieces for transcriptomic,
proteomic, and epigenetic experiments in a dry-ice bath using precooled scalpels and
plasticware.
115
Genetic analysis
Genomic DNA was isolated from 50 mg of GFM frozen brain tissue by using the Qiamp DNA
mini kit (Qiagen) following the manufacturer protocol. DNA concentration and purity were
assessed by nanodrop measurement. DNA integrity was evaluated by loading 100 nanogram
per sample on a 0,8% agarose gel and comparing size distribution to a size standard.
Presence of C9orf hexanucleotide repeat expansion (HRF) in post-mortem brain tissues was
confirmed by primed repeat PCR according to established protocols. Reported mutations for
MAPT and GRN were verified by sanger sequencing.
124
125 **Transcriptomic procedures**
126
RNA isolation from human brain tissue

Total RNA for CAGE-seq and RNAseq was isolated from ±100mg of frozen brain tissue with
TRIzol reagent (Thermo Fischer Scientific) according to the manufacturer recommendation,
followed by purification with the RNeasy mini columns (Qiagen) after DNAse treatment.
Total RNA for smallRNA-seq was isolated from frozen tissue using the TRIzol reagent
(ThermoFischer Scientific). After isopropanol precipitation and 80% ethanol rinsing RNA
pellet was resuspended in RNAse free water and up to 10 micrograms of RNA was incubated
with 2U of Ambion DNAse I (ThermoFischer) at 37°C for 20 minutes. DNA-free RNA samples
were then further purified by phenol-chloroform-isoamyl-alchol extraction followed by
ethanol precipitation.
137
138 **RNA QC**
139 For each RNA sample, RNA concentration (A260) and purity (A260/280 and A260/230) were
140 determined by Nanodrop measurement and RNA integrity (RIN) was assessed on a
141 Bioanalyser 2100 system and/or Tape station 41200 (Agilent Technologies Inc.)
142
143 **RNAseq libraries**
144 Total RNAseq libraries were prepared from 1 microgram of total RNA from frozen brain
145 tissue using the TruSeq Stranded Total RNA with Ribo-Zero Gold kit (Illumina) according to
146 the protocol specifications. RNAseq libraries were sequenced on a Hiseq2500 and HISeq4000
147 on a 2x100 bp paired end (PE) flow cell (Illumina) at an average of 100M PE/sample.
148
149 **CAGE-seq libraries**
150 CAGE-seq libraries were prepared from 5 micrograms of RNA from frozen brain tissues
151 according to a published protocol55. Libraries were sequenced on a HiSeq 2000 and/or
152 HiSeq2500 on a 1x50 bp single read (SR) flow cell (Illumina) at an average of 20M
153 reads/sample. On average, 21,212,923 reads were generated per sample.
154
smallRNAseq libraries
Small RNA-seq libraries were prepared from 1 microgram of total RNA from NPC-derived
neurons and 300 nanograms of microglia after miRNA mimics and inhibitors transfection,
using the mRNA TrueSeq Stranded kit (Illumina). mRNAseq libraries were sequenced on a
NextGen550 on a 75 cycles flow cell (Illumina). Small RNAseq libraries from frozen tissue
were prepared starting from 2 micrograms of total RNA using the Nextflex Small RNA-seq kit
v3 (Bioo Scientific) and the NEBNext Small RNA library prep set for Illumina (New England
Biolabs). Libraries were sequenced on a NextSeq550 on a 75 cycles flow cell.
163
Methylation assay
To assess the methylation status of over 850000 CpG sites in promoter, gene body and
enhancer regions we have used the MethylationEPIC bead chip arrays (Illumina).
Bisulfite conversion of genomic DNA, genome amplification, hybridization to the beadchips,
washing, staining, and scanning procedure was performed by Atlas Biolabs (Atlas Biolabs,
Berlin, Germany). Cases and controls DNAs were distributed randomly across each array.
170
RNA-seq processing and analysis
Raw FastQ files were processed using the RNA-seq pipeline from nf-core (nf-core/rnaseq
v1.3)[8], with trimming enabled. Gene quantification was subsequently done using Salmon
(v0.14.1)[9] on the trimmed FastQ files. Alignment and mapping were performed against the
human genome hg38. On average, 82,165,192 reads could be uniquely mapped, which
relates to on average 88.6% uniquely mapped reads per sample. In total, 59,270 transcripts
could be identified. DESeq2 (v.1.26.0)[10] was used to perform differential expression analysis.
We corrected for the covariates gender and PH-value.
179

Cell type deconvolution

We performed cell type deconvolution on the RNA-seq data using Scaden[11]. For training we used the human brain training dataset used in the Scaden publication. Each ensembl model was trained for 5000 steps. Cell type deconvolution was then performed with the trained Scaden model on the RNA-seq count data. Relative changes in cell type composition were quantified by first calculating the average fractions of a cell type for all groups and then calculating the percentual change of cell fractions compared to the average control fractions. This allows to detect relative changes in cell type compositions.

CAGE-seq processing and analysis

Sequencing adapters and barcodes in CAGE-seq FastQ files were trimmed using Skewer (v.0.1.126)[12]. Sequencing artefacts were removed using TagDust (v1.0)[13]. Processed reads were then aligned against the human genome hg38 using STAR (v.2.4.1)[14]. On average, 16,306,077 could be uniquely mapped per sample (76% uniquely mapped on average reads per sample). CAGE detected TSS (CTSS) files were created using CAGEr (v1.10.0)[15]. With CAGEr, we removed the first G nucleotide if it was a mismatch. CTSS were clustered using the 'distclu' method with a maximum distance of 20 bp. For exact commands used we refer to the reader to the scripts used in this pipeline: https://github.com/dznetubingen/cageseq-pipeline-mf. In total, we could identify 47,298 different peaks.

smRNA-seq processing and analysis

After removing sequencing adapters, all FastQ files were uploaded to OASIS2[16] for analysis. On average, 3,430,613 (+/- 1,365,407) reads could be uniquely mapped per sample. Subsequent differential expression analysis was performed on the counts yielded from OASIS2, using DESeq2 and correcting for gender and PH-value, as was done for the RNA-seq data. Additionally, we added a batch variable to the design matrix to correct for the two different batches of this dataset. In total, 2904 human smRNA genes could be detected.

Methylation data processing and analysis

The Infinium MethylationEPIC BeadChip, which consists of 866,091 CpG locations, data was analyzed using the minfi R package[17]. We removed all sites with a detection P-value above 0.01, on sex chromosomes and with single nucleotide polymorphisms (SNPs), leaving 810,290 loci for analysis. Data normalization was done using stratified quantile normalization. Sites with a standard deviation below 0.1 were considered uninformative and filtered out, leaving 170,595 sites that were used to perform a principal component analysis.

Data Records

All datasets have been submitted to the European Phenome-Genome Archive (EGA) under the study ID EGAS00001004895.

Technical Validation

RNA Integrity

We only included samples with a RNA integrity value above 5. The mean RIN value is 6.9 with a standard deviation of 1.0.

Methylation Data Quality Control and Normalization

The detection P-value has been calculated for all positions and samples. This value is calculated by comparing the total DNA signal to the background level, which is estimated using negative control positions. All samples had very low detection P-values, with a maximum value of 0.0005 (Fig. 2a). We used stratified quantile normalization to normalize

231   the methylation signal for every position. From Fig. 2 a) and b) it is visible that this
232   normalization can be used to even out the signal between samples in this dataset.
233

234
235   **Figure 2:** Methylation data QC statistics. **a)** Per-sample Detection P-values calculated with
236   the minfi R-package. **b)** and **c)** Density plots of methylation beta values before and after
237   quantile normalization, respectively.
238
239
Principal Component Analysis of Datasets
We performed a principal component analysis (PCA) on the counts for all sequencing
datasets and the M-values of the Methylation dataset (Fig. 3 a-d). As can be expected from
post-mortem human brain data, the datasets are heterogeneous. However, for all assays are
difference of the control samples to the different FTD subgroups can be seen. This shows
that there is a clear transcriptomic difference between brains from healthy patients and
from those with FTD. The PCA therefore confirms that the data is suitable for the
investigation of the disease.
248

249
250 **Figure 3**: Principal component analyses of the different datasets. Shown are the first two
251 principal components on the x and y axis. The samplesa are coloured according to the
252 disease group. **a)** RNA-seq dataset **b)** smRNA-seq dataset **c)** CAGE-seq dataset **d)** Methylation
253 dataset
254
Cell type composition of samples
To evaluate how the predicted cell type composition of the RiMod-FTD samples reflects the
disease, we performed cell type deconvolution with the RNA-seq data using the Scaden
algorithm (see Methods). As expected, neuronal cells make up the largest fractions for all
samples, although neuronal fractions are smaller in samples with FTD (Figure 4 b).
Comparison of percentage changes of average cell type fractions compared to controls
reveals that microglial fractions are particularly large in samples with FTD-GRN. Furthermore,
all disease subtypes show a strong increase in endothelial cell fractions compared to
controls. The decrease in neuronal cells is largely caused by a smaller number of excitatory
neurons in all disease subtypes, while inhibitory neuron fractions are not decreasing. These
results reflect, to a large extent, what is known about the biology of FTD.
266

268 **Figure 4**: Predicted cell composition of samples and disease groups. **a)** Difference in
269 percentage of cell compositions per group and cell type, compared to the healthy controls.
270 **b)** Stacked bar plot showing the sample-wise predicted cell composition.
271

Usage Notes

Inspection of Assay Results in RiMod-FTD Browser
The data from the different assays of the RiMod-FTD project can be easily inspected using
the RiMod-FTD Browser at https://www.rimod-ftd.org. The website allows to query the RNA-
seq expression levels of single genes. It is easily visible, for instance, that GRN RNA levels are
lower in brains from patients with GRN mutations, compared to the other groups (Fig. 5a).
When selecting a gene, the surrounding genomic location is additionally shown with RNA-
seq expression levels for genes in the region, CAGE-seq peaks and methylation levels. All
data can be grouped by disease, pathology, sex, mutated gene and specific mutation (Fig. 5b-
c). The browser does allow for quick inspection of FTD-related genes.
282

Figure 5: Exemplary view of RiMod-FTD genome browser. a) Normalized RNA-seq expression
levels of GRN for the different disesae groups in the RiMod-FTD project. b) Browser view a
selected gene (GRN). c) CAGE-seq peaks and methylation levels for the currently selected
region in the browser. d) Gene expression levels for genes in the selected region of the
genome browser.

Code Availability

The code for all analyses and figures of this study has been deposited at
https://github.com/dznetubingen/rimod-ftd-dataset.

Acknowledgements

Post-mortem brain tissue was obtained from the Dutch Brain Bank, Netherlands Institute for
Neuroscience, Amsterdam, and from the London Neurodegenerative Disease Brain Bank,
King's College London, London, UK. The London Neurodegenerative Disease Brain Bank is
part of the Brains for Dementia Research Initiative.

Author contributions

PH initiated and designed the project, planned, and interpreted the experiments and wrote
the manuscript.
KM was involved in all bioinformatic data analyses and wrote the manuscript.
AF, LK, PR and NF performed the small RNA-seq experiments.
CB, PR, NF and MC performed RNA-seq and CAGE-seq experiments.
MF and TN analysed the CAGE-seq data.

BA analysed the RNA-seq data.
PR and SB planned and interpreted analyses and wrote the manuscript.

Competing interests
The authors declare no competing interests.

References

1. Bang, J., Spina, S. & Miller, B. L. Frontotemporal dementia. *The Lancet* vol. 386 1672–1682 (2015).
2. Panza, F. *et al.* Development of disease-modifying drugs for frontotemporal dementia spectrum disorders. *Nat. Rev. Neurol.* **16**, 213–228 (2020).
3. Seelaar, H., Rohrer, J. D., Pijnenburg, Y. A. L., Fox, N. C. & van Swieten, J. C. Clinical, genetic and pathological heterogeneity of frontotemporal dementia: a review. *J. Neurol. Neurosurg. Psychiatry* **82**, 476–486 (2011).
4. Olszewska, D. A., Lonergan, R., Fallon, E. M. & Lynch, T. Genetics of Frontotemporal Dementia. *Current Neurology and Neuroscience Reports* vol. 16 (2016).
5. Sirkis, D. W., Geier, E. G., Bonham, L. W., Karch, C. M. & Yokoyama, J. S. Recent Advances in the Genetics of Frontotemporal Dementia. *Curr. Genet. Med. Rep.* **7**, 41–52 (2019).
6. Sieben, A. *et al.* The genetics and neuropathology of frontotemporal lobar degeneration. *Acta Neuropathologica* vol. 124 353–372 (2012).
7. Balendra, R. & Isaacs, A. M. C9orf72-mediated ALS and FTD: multiple pathways to disease. *Nat. Rev. Neurol.* 1 (2018) doi:10.1038/s41582-018-0047-2.
8. Ewels, P. A. *et al.* nf-core: Community curated bioinformatics pipelines. *bioRxiv* 610741 (2019) doi:10.1101/610741.
9. Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast and bias-aware quantification of transcript expression. *Nat. Publ. Gr.* **14**, (2017).
10. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, (2014).
11. Menden, K. *et al.* Deep learning–based cell composition analysis from tissue expression profiles. *Sci. Adv.* **6**, eaba2619 (2020).
12. Jiang, H., Lei, R., Ding, S. W. & Zhu, S. Skewer: A fast and accurate adapter trimmer for next-generation sequencing paired-end reads. *BMC Bioinformatics* **15**, 182 (2014).
13. Lassmann, T. TagDust2: A generic method to extract reads from sequencing data. *BMC Bioinformatics* **16**, 24 (2015).
14. Dobin, A. & Gingeras, T. R. Mapping RNA-seq Reads with STAR. in *Current Protocols in Bioinformatics* (2015). doi:10.1002/0471250953.bi1114s51.
15. Haberle, V., Forrest, A. R. R., Hayashizaki, Y., Carninci, P. & Lenhard, B. CAGEr: Precise TSS data retrieval and high-resolution promoterome mining for integrative analyses. *Nucleic Acids Res.* **43**, e51–e51 (2015).
16. Rahman, R. U. *et al.* Oasis 2: Improved online analysis of small RNA-seq data. *BMC Bioinformatics* **19**, 54 (2018).
17. Aryee, M. J. *et al.* Minfi: A flexible and comprehensive Bioconductor package for the analysis of Infinium DNA methylation microarrays. *Bioinformatics* **30**, 1363–1369 (2014).
18. Leek, J. T., Johnson, W. E., Parker, H. S., Jaffe, A. E. & Storey, J. D. The sva package for removing batch effects and other unwanted variation in high-throughput experiments. *Bioinformatics* **28**, 882–3 (2012).