Integrated multi-omics analysis reveals common and distinct dysregulated pathways for genetic subtypes of Frontotemporal Dementia

Kevin Menden[1,*] (KM); Margherita Francescatto[1] (MF); Tenzin Nyima[1] (TN); Cornelis Blauwendraat[1,2] (CB); Ashutosh Dhingra[1] (AD); Melissa Castillo-Lizardo[1] (MC); Noémia Fernandes[1] (NF); Lalit Kaurani[4] (LK); Deborah Kronenberg-Versteeg[1,3] (DKV); Burcu Atarsu[1,3] (BA); Eldem Sadikoglou[1] (ES); Barbara Borroni[6] (BB); Salvador Rodriguez-Nieto[1] (SRN); Javier Simon-Sanchez[1,3] (JSS); Andre Fischer[4] (AF); David Wesley Craig[7] (DWC); Manuela Neumann[1] (MN); Stefan Bonn[1,5] (SB); Patrizia Rizzu[1] (PR); Peter Heutink[1,*] (PH).

Affiliations

1 German Center for Neurodegenerative Diseases, Tübingen, Germany

2 National Institute on Aging, National Institutes of Health, Bethesda, MD, USA

3 Hertie Institute for Clinical Brain Research, Tübingen, Germany

4 German Center for Neurodegenerative Diseases, Göttingen, Germany

5 Institute of Medical Systems Biology, University Medical Center Hamburg-Eppendorf,

Hamburg, Germany

6 Neurology Unit, Department of Clinical and Experimental Sciences, University of Brescia,

Italy

7 University of Southern California, California, USA

* corresponding authors

Peter Heutink: peter.heutink@dzne.de

Kevin Menden: kevin.menden@qbic.uni-tuebingen.de

Abstract

Understanding the molecular mechanisms underlying frontotemporal dementia (FTD) is

essential for the development of successful therapies. Here we integrated transcriptomic and

epigenomic analyses of postmortem human brains of FTD patients with mutations in MAPT,

GRN and C9orf72 and detected common and distinct dysregulated cellular pathways

between patient groups. Our results highlight that excitatory neurons are the most vulnerable

neuronal cell type and that vascular aberrations are a common hallmark in FTD. Via

integration of multi-omics data, we detected several transcription factors and pathways

which regulate the strong neuroinflammation observed in FTD-GRN. Small RNA-seq data

and verification experiments in cellular models identified up-regulated miRNAs that inhibit

cellular trafficking pathways in FTD and lead to microglial activation. These findings shed

light on novel mechanistic and pathophysiological hallmarks of FTD. The data represent the

1[st] phase of a multi-omics, multi-model data resource for FTD research which allows in-

depth molecular research into disease mechanisms that will further mechanistic FTD

research.

Introduction

Frontotemporal Dementia (FTD) is a devastating pre-senile dementia characterized by

progressive deterioration of the frontal and anterior temporal lobes [1]. The most common

symptoms include severe changes in social and personal behaviour as well as a general

blunting of emotions. Clinically, genetically, and pathologically there is considerable overlap

with other neurodegenerative diseases including Amyotrophic Lateral Sclerosis (ALS),

Progressive Supranuclear Palsy (PSP) and Cortical Basal Degeneration (CBD) [2]. Research

into FTD has made major advances over the past decades. Up to 40% of cases [3] have a

positive family history and up to 60% of familial cases can be explained by mutations in the

genes Microtubule Associated Protein Tau (MAPT), Granulin (GRN) and C9orf72 [4] which

has been key to the progress in our understanding of its molecular basis. Several other

disease-causing genes have been identified that account for a much smaller fraction of

cases [5]. Mutations in MAPT lead to accumulation of the Tau protein in neurofibrillary tangles

in the brain of patients while mutations in GRN and C9orf72 lead to the accumulation of

TDP-43 [6], as well as dipeptide repeat proteins (DPRs) and RNA foci in the case of C9orf72 [7].

As of today, no therapy exists that halts or slows the neurodegenerative process of FTD and

in order to develop successful therapies there is an urgent need to determine whether a

common target and therapy can be identified that can be exploited for all patients, or

whether the distinct genetic, clinical and pathological subgroups need tailored treatments.

Therefore, the development of remedies relies heavily on a better understanding of the

molecular and cellular pathways that drive FTD pathogenesis in all FTD subtypes.

Although our knowledge of FTD pathogenesis using molecular and cellular biology

approaches has significantly advanced during recent years, a deep mechanistic

understanding of the pathological pathways requires simultaneous profiling of multiple

regulatory mechanisms. As neurodegenerative diseases develop over time, it is furthermore

important to examine temporal changes. While post-mortem human brain tissue represents

the end-stage of disease, well-defined rodent models can be used to address the temporal

component. Lastly, experimental validation of derived hypotheses can be achieved in cellular

systems, such as neurons derived from induced pluripotent stem cells (iPSCs) as well as in

rodent models.

The Risk and modifying factors in Frontotemporal Dementia (RiMod-FTD) consortium[8]

generates a multi-model and multi-omics data resource with the focus on mutations in the

three most common causal genes: MAPT, GRN and C9orf72. The data resource will consist

of multi-omics datasets from multiple post-mortem human brain regions, and matching iPSC

derived neurons and brain tissue of transgenic mouse models at different time points.

Here, we report on data derived from the post-mortem human brain RNA-seq, CAGE-seq,

smRNA-seq and methylation datasets from RiMod-FTD. We identified dysregulation of

overlapping pathways in all disease groups that indicates converging disease mechanisms

manifesting during disease progression. Using deconvolution analysis, we have identified

changes in cellular composition that are either common or distinct to genetic subgroups.

Through integration of smRNA-seq and CAGE-seq data, we could furthermore highlight

potential regulatory molecules that might play important roles in FTD pathogenesis, within

the identified functional gene modules and pathways.

89 Results

90 Multi-omics Data Resource for Frontotemporal Dementia

We have analysed data from brain tissue from diseased patients carrying mutations in the

MAPT (n=17), GRN (n=11) or C9orf72 (n=17) genes and non-demented controls (n=16).

The average age of FTD groups was lower than that of healthy controls (Table S1). We

obtained tissue from up to 7 regions for each brain. The temporal and frontal lobes are the

most affected areas in FTD, but we also obtained material from the occipital lobe,

hippocampus, caudate, putamen and cerebellum for verification experiments. We performed

CAGE-seq, smRNA-seq and quantitative proteomics (Miedema et al., manuscript in

preparation) on tissue from frontal and temporal lobes, and generated methylation and RNA-

seq data for frontal lobe tissue. Additional data types such as ATAC-seq, transgenic mouse

models and iPS derived data are planned for future releases. The resulting, comprehensive

multi-omics data resource enables the study of disease mechanisms in FTD subtypes to

greater detail than single genomics experiments (Fig. 1).

Figure 1: Graphical overview of the RiMod-FTD project. a In phase 1, Human Post-

mortem brain tissue samples from multiple regions of patients with mutations in GRN, MAPT

and C9orf72 have been collected and used for multi-omics data generation. Grey shading

indicates datasets that will be completed in future phases. The datasets have been

integrated and analysed and published to be accessible as FTD resource. In subsequent

phases additional datatypes will be added (i.e. ATAC-seq) and the resource will be extended

with data from matching mouse models and iPSC derived celltypes. **b** The multi-omics

approach allows to profile multiple regulatory features of gene expression, including

enhancer- and promoter-based regulation, epigenetic regulation, alternative splicing, post-

transcriptional regulation (miRNAs) and regulation of translation (proteomics).

116   In the current study, we have integrated RNA-seq, CAGE-seq, smRNA-seq and methylation

117   data from the Gyrus Frontalis Medialis (GFM) (Table S2), as this brain region is strongly

118   affected in FTD.

Differential gene expression analysis and cellular deconvolution

of the GFM in FTD.

To identify general gene expression patterns in the GFM of patients with FTD, we performed

differential gene expression (DGE) and principal component analysis (PCA) using RNA-seq

data. The PCA indicates considerable heterogeneity between samples, as can be expected

from post-mortem human brain tissue (Fig. 2a). However, a difference between FTD cases

and control samples is clearly visible. Differentially expressed genes (DEGs) were calculated

for all disease groups (FTD-MAPT, FTD-GRN, FTD-C9orf72) compared to controls while

controlling for gender and pH-value (see Methods). We observed the largest number of

DEGs (adj. P-value < 0.05) for FTD-GRN, followed by FTD-MAPT and FTD-C9orf72 (Fig.

2b). DGE of smRNA-seq data yielded 78, 21 and 39 differentially expressed miRNAs in

FTD-MAPT, FTD-GRN and FTD-C9orf72, respectively (Fig. 2c).

Due to the neurodegenerative nature of FTD, it is likely that there exists a systematic

difference in cell composition between cases and controls which can affect DGE analysis

due to differences in gene expression between cell types - a problem which has often been

overlooked in tissue expression studies. Here, we account for this problem by applying a

conservative filtering approach and removing DEGs that are associated with changing

cellular composition (see Methods). All further analyses were based on the filtered set of

DEGs, unless otherwise specified. Note that this method could only be applied to the total

RNA-seq dataset because similar cell type specificity data (here, single-cell RNA-seq data)

was not available for other data types.

Activation of extracellular matrix (ECM) associated pathways and circulatory system development.

We next performed pathway enrichment analysis with DEGs from the RNA-seq data using

go:Profiler[9] to identify the most affected cellular pathways. Down-regulated genes are

strongly enriched for mitochondrial and oxidative phosphorylation pathways in both FTD-

GRN and FTD-MAPT (Fig. 2d, Fig. S1), indicating a dysfunctional energy metabolism - a

well-known hallmark of many neurodegenerative diseases[10]. Neuronal system pathways are

enriched among down-regulated genes for both groups as well. This might be explained by

dysfunctional neurons that have not yet undergone apoptosis or by a general impairment of

neuronal function caused by the disease. Other significantly down-regulated pathways

include ubiquitin-dependent protein metabolism and vesicle-mediated transport (FTD-GRN).

In all three groups, up-regulated genes are enriched for extracellular matrix (ECM)

associated pathways and circulatory system development (Fig. S1). Genes involved in

Hippo-signalling are enriched in FTD-GRN and FTD-MAPT (Fig. 2d), and immune system

related genes are enriched in FTD-GRN. ECM dysregulation, in particular, has been

implicated with several neurodegenerative diseases. For instance, studies in mouse models

showed that tau pathology can lead to ECM reorganization and that reducing ECM proteins

could reverse memory deficits in an AD model[11,12]. While the role of the ECM in FTD

remains unknown, our results suggest a prominent involvement in end-stage FTD.

We also specifically examined the DEGs with the largest fold-changes in the RNA-seq data

because large expression fold-changes often signify strong dysregulation. These results

support the importance of ECM in FTD as for all disease groups, multiple matrix

metalloproteinase enzymes (MMPs) are among the DEGs with the largest LFCs (Fig. S2,

Fig. S3A). Elevated RNA levels of MMP genes have been reported for many

neurodegenerative diseases, and MMPs target a wide range of ECM [13]indicating their

importance in neurodegenerative mechanisms[14]. Protein interaction networks of up-

regulated genes in FTD-MAPT and FTD-GRN show the central importance of MMPs in

these networks (Fig. S3 B & C).

Figure 2: **Gene- and Pathway-level transcriptional changes in FTD. a** Principal

component analysis of variance stabilized RNA-seq expression values, coloured by group. **b**

Overlap between RNA-seq DEGs from different disease groups. **c** Overlap between

smRNA-seq DEGs from different disease groups. **d** Enriched Reactome pathways in RNA-

seq up- and down-regulated DEGs. Shown are the ten most significant pathways per group;

the x-axis signifies the negative log10 P-value. Colour corresponds to adjusted P-value and

node size corresponds to the number of genes in a pathway. **e,f** Best candidates for active

and inactive TFs in FTD-GRN and FTD-MAPT, respectively. The x-axis signifies the

negative log10 P-value. **g** Overlap of DMPs in different disease groups.

Regulatory mechanisms associated with differential expression

To better understand relevant regulatory mechanisms leading to these gene expression

changes, we generated a set of candidate driver transcription factors (TFs) using the GFM

CAGE-seq data. CAGE-seq cluster counts, when assigned to the closest gene, correlate

well with RNA-seq expression data (average sample-wise correlation coefficient: 0.6, Fig.

S4). We used the CAGE-seq data to predict candidate driver TFs for up- and down-

regulated genes (see Methods for details). TEAD2, a TF central to the Hippo signalling

pathway, is the only predicted active TF common to FTD-GRN and FTD-MAPT (Fig. 2e),

while there is greater overlap among inactive TFs (here: inactive TF = has down-regulated

targets, Fig. 2f). Moreover, we performed miRNA-target gene mapping to evaluate potential

regulatory roles of miRNAs. Expression values of miRNAs were correlated with their

predicted targets using matching samples from the RNA-seq data. Only miRNA-target pairs

with considerable negative correlation were retained (see Methods).

DNA methylation is another important regulatory mechanism that can affect gene

expression. We generated Illumina Infinium EPIC methylation data from the GFM and

considered the most variable CpG sites (28,173) corrected for possible confounding effects

using surrogate variable analysis (SVA) to perform differential methylation analysis (see

Methods). We detected 18,126, 9,478 and 5,282 significantly differentially methylated

positions (DMPs) for FTD-MAPT, FTD-GRN, and FTD-C9orf72, respectively (Fig. 2g). The

C9orf72 repeat expansion is known to be associated with hypermethylation[15] and we

confirmed in our data that a CpG site located at the 5'-end of the C9orf72 gene, only 14 bp

away from the repeat expansion, is hypermethylated (log fold-change: 0.6, Fig. S4A).

Pathway enrichment analysis of genes in proximity to DMPs yielded enrichment of genes

involved in nervous system development for hypermethylated CpG sites. Genes close to

hypomethylated sites were enriched for system development and vasculature development

(Fig. S5C). As hypermethylation of CpG sites at promoter regions is associated with

decreased expression, this indicates epigenetically controlled expression inhibition of genes

important for neuronal function, or remnants of cell composition effects that could not be

entirely alleviated by SVA (see Methods). Performing biological age prediction using the

methylation data resulted in underestimated age predictions for all groups, albeit to a lesser

extent for FTD groups, which indicates accelerated aging in FTD (Fig. S5B).

217 Vulnerability of excitatory neurons and enrichment of

218 endothelial cells

To identify vulnerable cell types and disease-related cell composition changes, we inspected

the results from the RNA-seq deconvolution analysis (Methods) with respect to genetic FTD

subtypes. As expected, fractions of neuronal cells are systematically lower in all FTD groups

compared to controls (Fig. S6). Consequently, virtually all other cell types show increasing

percentages. We therefore calculated the percentage-wise change for each cell type and

assessed statistical significance (see Methods). Strongest neuronal loss was observed in

FTD-GRN, followed by FTD-MAPT and FTD-C9orf72 (Fig. 3a, Table S3), which agrees with

studies that have shown that the frontal lobe is most strongly affected in FTD-GRN[16–18].

Moreover, neuronal loss can be primarily attributed to loss of excitatory neurons, while

fractions of inhibitory are not significantly different to controls (Table S3). Our results confirm

findings from recent studies that found excitatory neurons to be especially vulnerable to tau

pathology[19] and we specifically detected an important role of glutamatergic

neurotransmission in FTD [20,21]. Closer examination of the KEGG pathway 'glutamatergic

synapse' suggests that AMPA receptors are mainly affected, while we could not see signs of

dysregulation for NMDA receptors (Fig. S7 A-C). Analysis of candidate regulator TFs

highlighted the TF Early Growth Response 3 (EGR3), targets of which are enriched for

glutamatergic synapse genes (Fig. S7D), indicating involvement in excitatory neuronal

function.

To validate our computational deconvolution, we considered the fractions of excitatory

neurons as a proxy of neurodegeneration and correlated them with manually determined

degeneration scores by a neuropathologist (Fig. 3b, Methods). Indeed, excitatory neuron

fractions show strong negative correlation with pathology scores (Pearson's correlation

coefficient = -0.78, P-value = 2.8e-07), thereby providing experimental confirmation of our

computational predictions.

The strongest growth in percentage compared to the baseline is observed for endothelial

cells in FTD-MAPT and FTD-C9orf72 disease groups, but not FTD-GRN, where microglial

cells show the strongest increase. Circulatory system development is among the most

significantly up-regulated biological processes in all three disease groups (Fig. 3 c & d). The

role of the circulatory system in FTD is relatively unexplored. However, Bennet et al. recently

found increased vasculature growth in mouse models of FTD-MAPT with a P301L

mutation[22]. Interestingly, endothelial enrichment in FTD-MAPT is particularly strong in

patients with a P301L mutation (Fig. 3a). Another recent study observed a particular

microvascular structure with increased frequency in brains of patients with frontotemporal

lobar degeneration (FTLD)[23] and Park et al. have shown that soluble tau can interfere with

nitric oxide production and thus lead to reduced vasodilation of blood vessels, ultimately

leading to insufficient blood supply [24].

Figure 3: Cell composition changes in FTD. a Percentage change of averaged fractions

per group compared to the average of the control group. Genetic subtypes are indicated with

different colours. **b** Regression of excitatory neuron fractions (y-axis) against neuropathology

scores (x-axis). **c**, **d** PPI networks of genes up-regulated in FTD-MAPT and FTD-GRN (log-

fold-change > 1), respectively. Genes involved in the biological process "circulatory system

development" are labelled. **e** Heatmap of EWCE analysis results for HumanBase modules of

FTD-GRN and FTD-MAPT. Different modules are lined up on the x-axis, different cell types

on the y-axis. Tile colour signifies the EWCE P-value. Tiles with P-values above 0.1 are

marked grey.

To better understand transcriptional changes and regulatory mechanisms, we performed

tissue-specific functional module detection with HumanBase[25] and assessed cell type

specificity of modules using EWCE[26]. Both for FTD-MAPT and FTD-GRN, most modules

show specificity for a few cell types (Fig. 3e). Up-regulated modules in both groups are

significantly enriched for endothelial genes (P-value < 0.1). Genes within these modules

have been associated with blood vessel development (FTD-MAPT M6-up) and endothelial

cell growth (FTD-GRN M4-up) by HumanBase (Fig. 3 c & d), further supporting a distinct

involvement of endothelial genes in these FTD subtypes.

279   Increased Inflammatory response in FTD-GRN

In patients with FTD-GRN, microglial fractions show an even larger relative increase than

endothelial cells, indicating increased microglial activity. The protein encoded by GRN is well

known for its importance to lysosomal function, is highly expressed in microglia and well-

known for having important functions in the immune system [27,28]. Microglia are also slightly

enriched in FTD-MAPT (P-value = 0.037) but not in FTD-C9orf72 (P-value = 0.475).

Here, we have observed a prominent increase in microglial cell fractions and up-regulation of

immune system pathways in FTD-GRN, a feature of GRN deficiency that has been

frequently shown in mouse models[28–30]. We therefore wanted to further characterize

potential underlying regulatory mechanisms. First, we examined FTD-GRN modules for

enrichment of immune system-related terms. Indeed, several up-regulated modules are

enriched for genes related to the immune system, while we could not find enrichment among

down-regulated modules. The module FTD-GRN M1-up contains genes important for

neutrophil migration and response to interleukins (Fig. 4 a & d). Both modules M3-up and

M4-up contain genes relevant to NF-kappa-B (NFkB) signalling, as well as genes involved in

tumour necrosis factor (TNF) production (Fig. 4 b & c, respectively). Finally, the module M6-

up is enriched for genes involved in T cell activation. Modules M3-up, M4-up and M6-up are

furthermore enriched for microglial-specific genes (Fig. 3e). Interestingly, several

necroptosis-related genes are up-regulated (M1-up: TLR3, TLR8, RIPK3; M4-up: RIPK2),

suggesting this pathway as a potential driver of neuronal death. While we did not detect

prominent signals for neuroinflammation in FTD-MAPT, the FTD-MAPT module M3-up

contains several genes involved in T cell and TNF signalling (EZR, RAB29, CARD8, HIPK1).

However, neuroinflammation is much less prominent in FTD-MAPT and FTD-C9orf72

compared to FTD-GRN.

Figure 4: **Neuroinflammation in FTD-GRN. a** Up-regulated HumanBase modules in FTD-GRN with most significant terms. **b** Protein-protein interaction (PPI) network (made with String-DB) of FTD-GRN M1 up-module. Genes involved in necroptosis, interleukin response and neutrophil migration are indicated **c** PPI network of FTD-GRN M4-up module. Genes involved in NFkB signalling are indicated. **d** PPI network of FTD-GRN M3-up module. Genes involved in NFkB signalling and CEPBD are indicated. **e** and **f** KEGG pathway enrichment of predicted targets of TFs NFKB2 and RELA, respectively. **g** and **h** GO:BP pathway enrichment of predicted targets of TFs KLF3 and SP1, respectively.

Inspection of our candidate regulator TFs indicated the TFs Nuclear Factor Kappa B Subunit

2 (NFKB2) and RELA which together form the NFkB signalling complex, as potential drivers

in FTD-GRN (Fig. 2e). Enrichment analysis of predicted NFKB2 and RELA targets in FTD-

GRN indeed revealed TNF signalling and NFkB signalling as the most significantly enriched

KEGG pathways (Fig. 4 e & f). Furthermore, enrichment analysis indicated targets of the TFs

SP1 and KLF3 as highly enriched among genes in the FTD-GRN M3-up module. Predicted

KLF3 targets are enriched for immune system genes (Fig. 4g). SP1 target genes do not

show a strong enrichment but have roles in actin cytoskeleton organization and endothelial

cell differentiation, among others (Fig. 4h). We also investigated predicted targets of down-

regulated miRNAs and genes proximal to hypomethylated CpG sites for involvement in the

immune system in FTD-GRN but could not detect any significant immune system-relevant

enrichment suggesting they do not play a major role in regulating the immune response.

To closer examine which parts of the NFkB and TNF signalling pathways are affected in

FTD-GRN and in FTD in general, we inspected fold-changes of genes from the

corresponding KEGG pathways. Interestingly, the pro-inflammatory cytokine Interleukin 1

Beta (IL1B) is down-regulated in all disease groups, although only significantly in FTD-MAPT

(Fig. S9A). Similarly, the inflammatory cytokine Interleukin 6 (IL6) has negative fold-changes

in all disease groups. Downstream effector genes with positive fold-changes include multiple

chemokines, Interleukin 18 Receptor 1 (IL18R1) and several metalloproteinases.

GRN deficiency in mouse models leads to overactivation of microglia[28] and GRN deficiency

leads to NFkB overactivation in microglia in a GRN loss-of-function mouse model[29].

Hyperactivation of TNF by NFkB signalling has been linked to obsessive-compulsive

behaviour (OCB) and inhibition of NFkB was sufficient to rescue the behavioural symptoms

linking these TFs to the OCB observed in patients.

Dysfunctional energy metabolism and cellular trafficking in FTD

Among the most significantly down-regulated pathways in FTD-GRN and FTD-MAPT are

several pathways involved in energy metabolism and oxidative phosphorylation (Fig. 2d, Fig.

S1). Inspection revealed the modules FTD-GRN M1-down and FTD-MAPT M1-down as

being most significantly associated with the term NADH dehydrogenase complex assembly

(Fig. 5 d & e, Tables S4 and S5). Further inspection of the FTD-MAPT and FTD-GRN M1-up

modules revealed that they contain several NADH:Ubiquinone Oxidoreductase Subunit

genes (Fig. 5 b & c), which are necessary for functional oxidative phosphorylation and hence

energy production. The FTD-GRN module is moreover enriched for genes involved in

intracellular transport and autophagy. The FTD-GRN M1-down module contains several

genes associated with FTD or ALS: Superoxide Dismutase 1 (SOD1), Dynactin Subunit 1

(DCTN1), PTEN Induced Kinase 1 (PINK1), Huntingtin (HTT), and CHCHD10. All these

genes show lower expression values in every genetic subgroup, although they do not reach

significant levels in all groups (Fig. 5a).

-

Figure 5: Impaired oxidative phosphorylation and cellular trafficking in FTD. a

Expression levels (variance stabilized with DESeq2) of the genes CHCHD10, PINK1, SOD1,

DCTN1 and HTT in different groups. **b** STRING-DB PPI of FTD-MAPT M1-down module.

Genes involved in oxidative phosphorylation are labelled. **c** PPI of FTD-GRN M1-down

module. Genes involved in NADH dehydrogenase complex assembly and mitophagy are

labelled, as well as CHCHD10. **d, e** Most significant results from pathway enrichment

analysis with g:Profiler (GO biological process) for the FTD-GRN M1-down module and the

FTD-MAPT M1-down module, respectively. Node colour corresponds to adjusted P-value

and node size to term size. **f** Most significant results from enrichment analysis (Reactome) of

targets of up-regulated miRNAs in all disease groups. **g, h** Heatmaps of intersection-over-

union scores between predicted miRNA targets and down-regulated modules in FTD-GRN

and FTD-MAPT, respectively. **i** Top enrichment results of g:Profiler (GO biological process)

for the FTD-GRN M3-down module. Node colour corresponds to adjusted P-value and node

size to term size. **j** PPI network (String-DB) of FTD-GRN M3-down module. Predicted targets

of up-regulated miRNAs are labelled. **k** PPI network of predicted targets of up-regulated

miRNAs in FTD-MAPT.

Cellular transport is thought to play a key role in FTD pathogenesis as impaired trafficking

can affect protein and mitochondria homeostasis. Here, we show that mitochondrial function

is strongly impaired in end-stage FTD and that transport pathways are tightly connected to

this pathology. We thus looked for potential regulatory mechanisms driving the pathological

changes. Enrichment analysis of targets of up-regulated miRNAs in all disease groups

revealed cellular localization as the most significantly enriched biological process (GO:BP)

and membrane trafficking as the most significant Reactome pathway (Fig. 5f). Up-regulated

miRNAs in FTD therefore seem to primarily target cellular transport pathways and might play

important roles in dysfunctional transportation.

To detect modules and genes predominantly targeted by up-regulated miRNAs, we

calculated the intersection-over-union (IoU) of up-regulated miRNA targets with down-

regulated modules for FTD-GRN and FTD-MAPT (Fig. 5 g & h). The FTD-GRN M3-down

module is most strongly targeted by miRNAs and contains genes involved in metabolic

processes and cellular localization (Fig. 5i, Table S4). Five miRNAs have putative target

genes in this module: hsa-miR-150-5p, hsa-miR-142-3p, hsa-miR-193a-3p, hsa-miR-148a-

3p and hsa-miR-363-3p, which are all significantly up-regulated in FTD-MAPT as well,

except hsa-miR-363-3p. We generated networks of the above-mentioned candidate miRNAs

combined with a PPI network of the FTD-GRN M3-down module (Fig. 5j) and a PPI network

of all predicted targets in FTD-MAPT (Fig. 5k), as we could not detect a similar module in

FTD-MAPT. In total, we observed 31 common putative miRNA targets in both networks.

Next, we selected hsa-miR-193a-3p, hsa-miR-150-5p and hsa-miR-19b-3p for further

characterization in iPSC models (Fig. 6a). The first two miRNAs are DE in all three disease

groups and have many targets among module genes (Fig. 5 j & k). The miRNA hsa-miR-

19b-3p is up-regulated in all disease groups, although it does not reach significance after

(FTD-MAPT and FTD-C9orf72) or before (FTD-GRN) multiple testing correction.

Nevertheless, down-regulated genes were predicted to be enriched for targets of hsa-miR-

19b-3p by g:Profiler, the miRNA is known to inhibit autophagy [31] and it is highly expressed in

neurons. We performed RNA-seq on iPSC-derived neurons and microglia (Methods) that

were transfected with miRNA mimics and inhibitors for the three selected miRNAs. Here, we

focused on the mimic experiments, as the mimics should in theory reproduce the effects of

miRNA overexpression. Inhibition and mimicking of miR-150-5p in neurons had only minor

effects in neurons, while in microglia, the miR-150-5p mimic had strong effects, leading to

237 down-regulated and 236 up-regulated DEGs, enriched for cellular transport and immune

system pathways of the latter and nervous system development of the former (Fig. 6b).

Inhibition of miR-150-5p had even stronger effects (3221 DEGs), indicating an important

function of this miRNA in microglia. Transfection of miR-193a-3p mimic and inhibitor only

resulted in significant expression changes in microglia, where the mimic had strong effects

with 1756 down-regulated and 1474 up-regulated genes. Up-regulated genes were enriched

of mitochondrial functions like oxidative phosphorylation, while down-regulated genes were

enriched for localization and vesicle-mediated transport pathways (Fig. 6c).

In neurons, the miR-19b-3p mimic resulted in 89 down- and 137 up-regulated DEGs

(inhibitor: 8 down-regulated, 31 up-regulated). Genes down-regulated by the mimic and up-

regulated in the inhibitor experiment are involved in neuronal system pathways, enriched for

miR-19b-3p targets and share 17 common genes, thus providing evidence for these genes

to be regulated by miR-19b-3p. In microglia, stronger effects of the miR-19b-3p mimic

compared to the inhibitor were observed (1518 compared to 608 DEGs). Genes down-

regulated by the miR-19b-3p mimic were enriched for catabolic processes, autophagy and

vesicle-mediated transport, up-regulated genes were enriched for cell cycle and immune

system related genes (Fig. 6d). These results provide strong evidence that hsa-miR-19b-3p

and hsa-miR-193a-3p indeed regulate cellular trafficking pathways. Furthermore, hsa-miR-

150-5p is important for microglia function and up-regulation could lead to immune system

activation.

Figure 6: Effects of miRNA mimic and inhibitor experiments in iPSC-derived microglia.

a Boxplots of normalized expression values for the selected miRNAs. **b**, **c**, **d** The top ten most significantly enriched biological processes of up- and down-regulated genes after transfection with mimics for miR-150-5p, miR-193a-3p and miR-19b-3p, respectively. Node size corresponds to the number of genes in the biological process term and node colour corresponds to the P-value adjusted for multiple testing.

Discussion

Here, we present the data from the 1st phase of the RiMod-FTD project, a multi-omics, multi-model data resource for FTD research that aims to understand the role of distinct genetic

risk factors in the disease. Generated by the RiMod-FTD consortium over several years, the

resource depicts a valuable tool for FTD researchers that will help to accelerate scientific

progress towards a better understanding of relevant disease mechanisms in FTD. Additional

multi-omics data from iPSC derived cell types, transgenic mouse models and other brain

regions will be added over time.

By studying post-mortem tissue of the GFM we detected the largest transcriptional

dysregulation and greatest neuronal loss in FTD caused by mutations in GRN, agreeing with

previous findings of a more pronounced frontal lobe atrophy in FTD-GRN compared to other

subtypes[32]. Our deconvolution analysis indicates that excitatory neurons are the most

affected cell type in all genetic subtypes of FTD, which was confirmed in the RiMod-FTD

proteomics data (Mediema et al., manuscript in preparation) as well. Recently, evidence

from multiple studies has accumulated pointing toward a strong involvement of glutamatergic

synapses in FTD[33]. While it has been previously reported that densities of both ionotropic

glutamate receptors, AMPA and NMDA receptors, are reduced in post-mortem brain tissue

of FTD patients[21], we see evidence that AMPA receptors are particularly affected.

Intriguingly, a recent single-nucleus RNA-seq study in a GRN mouse model could show that

hyperactivation of microglia leads to selective loss of excitatory neurons[34], which confirms

our hypothesis that excitatory neurons are especially vulnerable in FTD.

Strong neuroinflammation is a distinct feature of FTD-GRN, which is confirmed in our data

and by increased microglial cell numbers in this FTD sub-type, in line with recent findings

showing increased microglial burden in FTD-GRN[35,36]. Using an integrative approach, we

identified TFs: NFKB2, RELA, KLF3 and SP1 as key inflammatory drivers, leading to

activation of the NFkB- and TNF-signalling pathways.

We found indicators of activated necroptosis, suggesting this pathway as potential cause for

cell death. The necroptosis cell death pathway is deregulated in several neurodegenerative

disorders[37], and a recent study has shown that TBK1, a genetic cause of ALS and FTD (here

down-regulated in FTD-GRN), is an endogenous inhibitor of RIPK1, an upstream regulator of

RIPK3[38]. The authors showed that embryonic lethality of TBK1-knockout mice is dependent

on RIPK1 activity, suggesting that the necroptosis pathway is indeed an important player in

FTD pathogenesis. In a recent review, Molnár and colleagues have discussed several

available drugs that could potentially regulate necroptosis[39], highlighting the potential of this

pathway as a drug target for developing therapies for FTD.

Our pathway enrichment and deconvolution analyses pointed toward increased blood vessel

abundance and growth in FTD brains compared to controls, which is consistent with the

results from the RiMod-FTD proteomics data (Mediema et al., manuscript in preparation). It

is generally not known how and if the vasculature system is involved in FTD pathogenesis,

although recent studies have observed abnormalities in a mouse model of tau pathology and

post-mortem human brains[22,23]. To our knowledge, angiogenesis as a pathological feature in

several genetic FTD subtypes has not been reported before and therefore depicts an

important subject for FTD research.

In all three disease groups, we have observed prominent up-regulation of ECM pathways

and MMP enzymes, suggesting MMPs as important regulators in FTD pathogenesis. While it

has been increasingly recognized that MMPs are important regulators in many

neurodegenerative diseases [40,41], the role of MMPs in FTD pathogenesis has not been

investigated in depth. In mouse models of ALS, inhibition of the MMPs MMP2 and MMP9

could indeed prolong survival and reduce symptoms[42,43]. Moreover, TIMP3, which is up-

regulated in our data, was found to be partly responsible for neuronal apoptosis in an ALS

model [44], which points towards TIMP3 as a potential apoptosis mechanism in FTD. MMPs

are furthermore tightly involved in the inflammatory response, and can activate the tumour

necrosis factor (TNF) gene[45]. Inflammatory cytokines, hypoxia and reactive oxygen species

can lead to the activation of MMPs[40,46] and MMPs can digest the ECM, stimulate increased

production of growth factors and thereby promote the growth of blood vessels, providing a

potential causal link to the prominent enrichment of endothelial cells[46]. Given their important

biological functions and their involvement in all genetic FTD subgroups, it will be important to

further investigate how MMPs contribute to FTD and whether they can be exploited as drug

targets, as MMP inhibition in model system has shown promising results[47,48].

Impaired cellular trafficking mechanisms is very likely a key feature of FTD pathogenesis and

it has been shown multiple times that FTD-causal mutations lead to trafficking deficits [49–51].

However, it is not always clear which mechanisms continue to dysfunctional transport

mechanisms. Here, using multi-omics data and validation experiments, we show that

elevated expression of several miRNAs contributes to the inhibition of genes important for

cellular transport. Additional studies are necessary to further validate this hypothesis, which

directly suggests several miRNAs as putative drug targets.

While our study is reasonably powered, increasing the sample size for individual groups

such as FTD-C9orf72 would further increase the power of our analysis and help to better

define which pathways are truly distinct to certain subtypes. It will therefore be an objective

for future iterations of the RiMod-FTD resource to include larger numbers of samples.

To conclude, we present here an integrated multi-omics analysis on data from Phase 1 of

the RiMod-FTD project and developed new hypotheses on FTD disease mechanisms. The

data presented here highlights several regulator molecules important for FTD pathogenesis

and their consequences such as vascular abnormalities and thereby we show the value of

an integrated multi-omics data analysis for hypothesis generation and testing. The RiMod-

FTD data will be freely accessible to the scientific community through the European

Genome-phenome Archive (**EGA**) and a dedicated RiMod-FTD web application

(https://www.rimod-ftd.org) thus enabling scientists to derive new mechanisms and

hypotheses from the data.

Methods

Donor samples employed in this study

Post mortem human brains

Tissues were obtained under a Material Transfer Agreement from the Netherlands Brain

Bank, and additional samples were provided by the Queen Square Brain Bank of

Neurological Disorders and MRC, King College London. Demographic details about human

brain samples are summarized in Table S1.

GFM and GTM tissue from each subject was divided into three pieces for transcriptomic,

proteomic and epigenetic experiments in a dry-ice bath using precooled scalpels and

plasticware.

hIPS-derived NGN2 neurons and miRNA mimics and inhibitors transfection

smNPC were derived from hiPSc cells (Cell line id: GM23280 obtained from the Coriell

Institute) using the protocol described by Reinhardt et al[52]. The differentiation protocol from

smNPC to neurons involves over-expression of Neurogenin-2 (NGN2) using a modified

version of the NGN2 lentiviral inducible vector system (single vector

pLV_TRET_hNgn2_UBC_BSD_T2A_rtTA3). The detailed description about protocol,

reagents and media composition is available in Dhingra et al.[53].

Briefly, stable NGN2 smNPC are grown for six days in expansion medium N2B27

supplemented with CHIR99021 (CHIR) 3 µM, Purmorphamine (PMA) 0.5 µM and L-ascorbic

acid 2-phosphate magnesium (AA) 64 mg/l. For differentiation, cells are plated (80,000

cells/cm2) onto Poly L-orithine and laminin coated plates in N2B27 medium supplemented

with doxycycline (dox) at 2.5 µg/mL, and 2 µM DAPT. On day 4 of differentiation,

transfection was performed in n=3 replicate plates using lipofectamine RNAiMax

(ThermoFisher Scientific) with a final concentration of miRNA mimic and inhibitors (miR-19b-

3p and miR-1505p mimics and inhibitors from Qiagen and miR-193a-3p mimic and inhibitor

from ThermoFisher Scientific) in the range of 5 to 10 nM as per the manufactures' guidelines

along with their corresponding controls. Next day (day 5 of differentiation), the complete

media was changed with N2B27 media supplemented with dox, 10 ng/mL brain-derived

neurotrophic factor (BDNF), 10 ng/mL glial cell-derived neurotrophic factor (GDNF), 10

ng/mL neurotrophic factor 3 (NT-3), 1 µg/mL Laminin, and 10 µM DAPT. Thereafter, half

media was changed on day 8 of differentiation. On day 11, cells were gently washed with

PBS and processed for RNA isolation.

hIPS-derived microglia and miRNA mimics and inhibitors transfection

hiPSCs were differentiated as previously described (van Wilgenburg et al[54]). In brief, 3 x

10^6 iPSCs were seeded into an Aggrewell 800 well (STEMCELL Technologies) to form

embryoid bodies (EBs), in mTeSR1 and fed daily with medium plus 50ng/ml BMP4 (Miltenyi

Biotec), 50ng/ml VEGF (Miltenyi Biotec), and 20ng/ml SCF (R&D Systems). Four-day EBs

were then differentiated in 6-well plates (15 EBs/well) in X-VIVO15 (Lonza) supplemented

with 100ng/ml M-CSF (Miltenyi Biotec), 25ng/ml IL-3 (Miltenyi Biotec), 2mM Glutamax

(Invitrogen Life Technologies), and 0.055mM beta-mercaptoethanol (Thermo Fisher

Scientific), with fresh medium added weekly. Microglial precursors emerging in the

supernatant after approximately 1 month were collected and isolated through a 40um cell

strainer and plated in N2B27 media supplemented with 100 ng/ml M-CSF, 25 ng/ml

interleukin 34 (IL-34) for differentiation. Thereafter, the media is reshred every 2 days

supplemented with 100 ng/ml M-CSF, and 25 ng/ml IL-34. The cells were cultured for

additional 6 days with media refresh every 2 days. On day 7 of maturation, transfection was

performed in n=3 replicate plates using lipofectamine RNAiMax with a final concentration of

miRNA mimics and inhibitors in the range of 5 to 10 nM as per the manufactures' guidelines

along with their corresponding controls (miR-19b-3p and miR-1505p mimics and inhibitors

from Qiagen and miR-193a-3p mimic and inhibitor from ThermoFisher Scientific). Next day

complete media was refreshed. On day 11, cells were gently washed with PBS and

processed for RNA isolation.

Genetic analysis

Genomic DNA was isolated from 50 mg of GFM frozen brain tissue by using the Qiamp DNA

mini kit (Qiagen) following the manufacturer protocol. DNA concentration and purity were

assessed by nanodrop measurement. DNA integrity was evaluated by loading 100

nanogram per sample on a 0,8% agarose gel and comparing size distribution to a size

standard.

Presence of C9orf72-HRE in postmortem brain tissues and hIPS cells was confirmed by

primed repeat PCR according to established protocols. Reported mutations for MAPT and

GRN were verified by sanger sequencing.

Transcriptomic procedures

RNA isolation from human brain tissue

Total RNA for CAGE-seq and RNAseq was isolated from ±100mg of frozen brain tissue with

TRIzol reagent (Thermo Fischer Scientific) according to the manufacturer recommendation,

followed by purification with the RNeasy mini columns (Qiagen) after DNAse treatment.

Total RNA for smallRNA-seq was isolated from frozen tissue using the TRIzol reagent

(ThermoFischer Scientific). After isopropanol precipitation and 80% ethanol rinsing RNA

pellet was resuspended in RNAse free water and up to 10 micrograms of RNA was

incubated with 2U of Ambion DNAse I (ThermoFischer) at 37°C for 20 minutes. DNA-free

RNA samples were then further purified by phenol-chloroform-isoamyl-alchol extraction

followed by ethanol precipitation.

RNA isolation from smNPC-derived neurons and microglia

Total RNA was isolated from NGN2 driven neurons and microglia cells after transfection with

miRNA mimics and inhibitors. Briefly at day 11 of transfection cells were carefully rinsed with

PBS and lysed in Qiazol buffer (Qiagen). Further DNAse treatment and purification were

carried out with the miRNeasy micro kit (Qiagen) according to the manufacturer protocol.

RNA QC

For each RNA sample, RNA concentration (A_{260}) and purity ($A_{260/280}$ and $A2_{60/230}$) were

determined by Nanodrop measurement and RNA integrity (RIN) was assessed on a

Bioanalyser 2100 system and/or Tape station 41200 (Agilent Technologies Inc.)

CAGE-seq libraries

CAGE-seq libraries were prepared from 5 micrograms of RNA from frozen brain tissues

according to a published protocol[55]. Libraries were sequenced on a HiSeq 2000 and/or

HiSeq2500 on a 1x50 bp single read (SR) flow cell (Illumina) at an average of 20M

reads/sample.

RNAseq libraries

Total RNAseq libraries were prepared from 1 microgram of total RNA from frozen brain

tissue using the TruSeq Stranded Total RNA with Ribo-Zero Gold kit (Illumina) according to

the protocol specifications. RNAseq libraries were sequenced on a Hiseq2500 and

HISeq4000 on a 2x100 bp paired end (PE) flow cell (Illumina) at an average of 100M

PE/sample.

smallRNAseq libraries

Small RNA-seq libraries were prepared from 1 microgram of total RNA from NPC-derived

neurons and 300 nanograms of microglia after miRNA mimics and inhibitors transfection,

using the mRNA TrueSeq Stranded kit (Illumina). mRNAseq libraries were sequenced on a

NextGen550 on a 75 cycles flow cell (Illumina). Small RNAseq libraries from frozen tissue

were prepared starting from 2 micrograms of total RNA using the Nextflex Small RNA-seq kit

v3 (Bioo Scientific) and the NEBNext Small RNA library prep set for Illumina (New England

Biolabs). Libraries were sequenced on a NextSeq550 on a 75 cycles flow cell.

Methylation assay

To assess the methylation status of over 850000 CpG sites in promoter, gene body and

enhancer regions we have used the MethylationEPIC bead chip arrays (Illumina).

Bisulfite conversion of genomic DNA, genome amplification, hybridization to the beadchips,

washing, staining and scanning procedure was performed by Atlas Biolabs (Atlas Biolabs,

Berlin, Germany). Cases and controls DNAs were distributed randomly across each array.

HumanBase Module Analysis

Functional gene modules were generated using the HumanBase tool at:

https://hb.flatironinstitute.org/. We divided DEGs into up- and down-regulated genes as we

were looking for active and repressed modules in FTD. Modules were downloaded for further

analysis. Cell type enrichment analysis was performed for genes of each modules using

EWCE[56] as described further down.

RNA-seq processing and analysis

Raw FastQ files were processed using the RNA-seq pipeline from nf-core (nf-core/rnaseq

v1.3) [57], with trimming enabled. Gene quantification was subsequently done using Salmon

(v0.14.1)[58] on the trimmed FastQ files. Alignment and mapping were performed against the

human genome hg38. DESeq2 (v.1.26.0)[59] was used to perform differential expression

analysis. We corrected for the covariates gender and PH-value. Genes were considered

differentially expressed when having a Benjamini-Hochberg corrected P-value below 0.05.

Cell type deconvolution and filtering

We performed cell type deconvolution on the RNA-seq data using Scaden[60]. For training we

used the human brain training dataset used in the Scaden publication. Each ensembl model

was trained for 5000 steps. To filter differentially expressed genes for false positives caused

by cell composition bias, we first calculated the correlation of gene expression with cell type

fraction. Then, we calculated a cell type specificity score as defined in Skene et al. [56] for

each gene available in the scRNA-seq dataset from Darmanis et al. [61]. We filtered out all

genes that had a specificity score of at least 0.5 and a positive correlation of at least 0.4 with

the cell type fractions of the most specific cell type. False positive DEGs that are caused by

systematic increase or decrease of a specific cell type will show high correlation with the cell

type fractions[62] and can thus be identified and removed from the analysis. A specificity score

of 0.5 means that half of the total gene expression for a certain gene can be attributed to a

single cell type, assuming a uniform cell type composition. The cut-offs for specificity score

and correlation were chosen based on an informed decision. Relative changes in cell type

composition were quantified by first calculating the average fractions of a cell type for all

groups and then calculating the percentual change of cell fractions compared to the average

control fractions. This allows to detect relative changes in cell type compositions. Statistical

significance between cell type fractions of groups was assessed using a t-test in the R

language.

Cell type enrichment analysis

We performed cell type enrichment analysis of genesets using the EWCE R package[56]. Cell

type specificity of genes was calculated from the single-cell RNA-seq cortex dataset of

Darmanis and colleagues[61]. EWCE analysis was done following instructions from

https://github.com/NathanSkene/EWCE.

CAGE-seq processing and analysis

Sequencing adapters and barcodes in CAGE-seq FastQ files were trimmed using Skewer

(v.0.1.126)[62]. Sequencing artefacts were removed using TagDust (v1.0)[63]. Processed reads

were then aligned against the human genome hg38 using STAR (v.2.4.1)[64]. CAGE detected

TSS (CTSS) files were created using CAGEr (v1.10.0)[65]. With CAGEr, we removed the first

G nucleotide if it was a mismatch. CTSS were clustered using the 'distclu' method with a

maximum distance of 20 bp. For exact commands used we refer to the reader to the scripts

used in this pipeline: https://github.com/dznetubingen/cageseq-pipeline-mf.

Transcription factor activity analysis

To identify candidate regulatory transcription factors, we first performed differential

expression analysis with all CAGE-seq clusters (see RNA-seq analysis). Then, we extracted

the sequence 600 bp up-stream and 300 bp downstream around all detected clusters. We

used Homer[66] to look for significant TFBS enrichment in the regions around up- and down-

regulated clusters (similar to [67]). TFBS motifs were downloaded from the JASPAR

database[68]. When calculating enrichment, we considered all extracted regions that are not

part of the set of interest as background. The complete pipeline can be found at

https://github.com/KevinMenden/tf-activity. We selected all TFs with significant enrichment

(p-value <= 0.001) for either up-regulated or down-regulated CAGE clusters as candidate

regulators. We considered genes as potential targets of a TF if a TFBS could be found in

their promoter region. As an additional filter, we selected only TFs with evidence for

differential expression in the RNA-seq data (adj. P-value < 0.05, not filtered for cell

composition).

smRNA-seq processing and analysis

After removing sequencing adapters, all FastQ files were uploaded to OASIS2 [69] for

analysis. Subsequent differential expression analysis was performed on the counts yielded

from OASIS2, using DESeq2 and correcting for gender and PH-value, as was done for the

RNA-seq data. Additionally, we added a batch variable to the design matrix to correct for the

two different batches of this dataset. For the target prediction analysis, we first downloaded

all targets from mirBase[70]. Then, we correlated the expression of miRNAs with their

predicted targets using matching samples from the RNA-seq data. We removed all predicted

targets with a correlation above -0.4, thus only considering miRNA-target pairings with high

negative correlation.

Methylation data processing and analysis

The Infinium MethylationEPIC BeadChip data was analyzed using the minfi R package[71]. We

removed all sites with a detection P-value above 0.01, on sex chromosomes and with single

nucleotide polymorphisms (SNPs). Data normalization was done using stratified quantile

normalization. Sites with a standard deviation below 0.1 were considered uninformative and

filtered out, to increase detection power. Surrogate variable analysis[72] was performed to

determine confounding factors. Differential methylation analysis was done using the limma

package[73] and controlling for the detected surrogate variables. Sites with a Benjamini-

Hochberg [74] adjusted P-value below 0.05 were considered differentially methylated.

Age prediction

We predicted the biological age of donors using the methylation data and the Wenda

algorithm[75]. Training data was kindly provided by the authors of Wenda. We subsetted the

data for CpG sites found in our data (11,729) sites and performed the prediction as

described at https://github.com/PfeiferLabTue/wenda.

Analysis of mRNA-seq data from cellular models

This section describes the analysis of mRNA-seq data generated for the miRNA mimic and

inhibitor experiments. FastQ files were mapped and gene counts quantified using Salmon

and differential expression analysis was performed with DESeq2 (see post-mortem brain

RNA-seq analysis). DEGs were examined for pathway enrichment using go:Profiler.

Assessment of degeneration

For assessment of neurodegeneration, H&E stained paraffin sections of the frontal and

temporal cortex were graded as absent (0), mild (1), moderate (2) and severe (3) based on

the presence of spongiosis, neuronal loss and gliosis.

Data Availability

All data used in this study and published as phase 1 of the RiMod-FTD resource have been

deposited at the European Phenome-genome Archive (EGA) under accession number

EGAS00001004895.

Code Availability

The code used for generating the analysis results is made freely available in the GitHub

repository https://github.com/dznetubingen/rimod-ftd-paper.

References

1. Bang, J., Spina, S. & Miller, B. L. Frontotemporal dementia. *Lancet* **386**, 1672–1682 (2015).

2. Panza, F. *et al.* Development of disease-modifying drugs for frontotemporal dementia spectrum disorders. *Nat. Rev. Neurol.* **16**, 213–228 (2020).

3. Seelaar, H., Rohrer, J. D., Pijnenburg, Y. A. L., Fox, N. C. & van Swieten, J. C. Clinical, genetic and pathological heterogeneity of frontotemporal dementia: a review. *J. Neurol. Neurosurg. Psychiatry* **82**, 476–486 (2011).

4. Olszewska, D. A., Lonergan, R., Fallon, E. M. & Lynch, T. Genetics of Frontotemporal Dementia. *Current Neurology and Neuroscience Reports* vol. 16 (2016).

5. Sirkis, D. W., Geier, E. G., Bonham, L. W., Karch, C. M. & Yokoyama, J. S. Recent Advances in the Genetics of Frontotemporal Dementia. *Curr. Genet. Med. Rep.* **7**, 41–52 (2019).

6. Sieben, A. *et al.* The genetics and neuropathology of frontotemporal lobar degeneration. *Acta Neuropathologica* vol. 124 353–372 (2012).

7. Balendra, R. & Isaacs, A. M. C9orf72-mediated ALS and FTD: multiple pathways to disease. *Nat. Rev. Neurol.* 1 (2018) doi:10.1038/s41582-018-0047-2.

8. RiMod-FTD | JPND. https://www.neurodegenerationresearch.eu/initiatives/annual-calls-for-proposals/closed-calls/risk-factors-2012/risk-factor-call-results/rimod-ftd/.

9. Raudvere, U. *et al.* g:Profiler: a web server for functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic Acids Res.* **47**, W191–W198 (2019).

10. Golpich, M. *et al.* Mitochondrial Dysfunction and Biogenesis in Neurodegenerative diseases: Pathogenesis and Treatment. *CNS Neurosci. Ther.* **23**, 5–22 (2017).

11. Li, Y., Li, Z.-X., Jin, T., Wang, Z.-Y. & Zhao, P. Tau Pathology Promotes the Reorganization of the Extracellular Matrix and Inhibits the Formation of Perineuronal Nets by Regulating the Expression and the Distribution of Hyaluronic Acid Synthases.

J. Alzheimer's Dis. **57**, 395–409 (2017).

12. Végh, M. J. *et al.* Reducing hippocampal extracellular matrix reverses early memory
deficits in a mouse model of Alzheimer's disease. *Acta Neuropathol. Commun.* **2**,
(2014).

13. Lu, P., Takai, K., Weaver, V. M. & Werb, Z. Extracellular Matrix degradation and
remodeling in development and disease. *Cold Spring Harb. Perspect. Biol.* **3**, (2011).

14. Duits, F. H. *et al.* Matrix Metalloproteinases in Alzheimer's Disease and Concurrent
Cerebral Microbleeds. *J. Alzheimer's Dis.* **48**, 711–720 (2015).

15. Xi, Z. *et al.* Hypermethylation of the CpG Island Near the G 4 C 2 Repeat in ALS with
a C9orf72 Expansion. *Am J Hum Genet* 981–989 (2013)
doi:10.1016/j.ajhg.2013.04.017.

16. Biswas, M. H. U. *et al.* MMP-9 and MMP-2 Contribute to Neuronal Cell Death in iPSC
Models of Frontotemporal Dementia with MAPT Mutations. *Stem Cell Reports* **7**, 316–
324 (2016).

17. Young, J. J., Lavakumar, M., Tampi, D., Balachandran, S. & Tampi, R. R.
Frontotemporal dementia: latest evidence and clinical implications. *Ther. Adv.*
Psychopharmacol. **8**, 33–48 (2018).

18. Cash, D. M. *et al.* Patterns of gray matter atrophy in genetic frontotemporal dementia:
results from the GENFI study. *Neurobiol. Aging* **62**, 191–196 (2018).

19. Fu, H., Hardy, J. & Duff, K. E. Selective vulnerability in neurodegenerative diseases.
Nat. Neurosci. **21**, 1350–1358 (2018).

20. Palese, F. *et al.* Anti-GluA3 antibodies in frontotemporal dementia: effects on
glutamatergic neurotransmission and synaptic failure. *Neurobiol. Aging* **86**, 143–155
(2020).

21. Murley, A. G. & Rowe, J. B. Neurotransmitter deficits from fronto temporal lobar
degeneration. *Brain* **141**, 1263–1285 (2018).

22. Bennett, R. E. *et al.* Tau induces blood vessel abnormalities and angiogenesis-related
gene expression in P301L transgenic mice and human Alzheimer's disease. *Proc.*
Natl. Acad. Sci. U. S. A. **115**, E1289–E1298 (2018).

23. Ek Olofsson, H. & Englund, E. A cortical microvascular structure in vascular dementia,
Alzheimer's disease, frontotemporal lobar degeneration and nondemented controls: a
sign of angiogenesis due to brain ischaemia? *Neuropathol. Appl. Neurobiol.* **45**, 557–
569 (2019).

24. Park, L. *et al.* Tau induces PSD95–neuronal NOS uncoupling and neurovascular
dysfunction independent of neurodegeneration. *Nat. Neurosci.* **23**, 1079–1089 (2020).

25. Greene, C. S. *et al.* Understanding multicellular function and disease with human
tissue-specific networks. *Nat. Genet.* **47**, 569–576 (2015).

26. Skene, N. G. & Grant, S. G. N. Identification of Vulnerable Cell Types in Major Brain
Disorders Using Single Cell Transcriptomes and Expression Weighted Cell Type
Enrichment. *Front. Neurosci.* **10**, 16 (2016).

27. Paushter, D. H., Du, H., Feng, T. & Hu, F. The lysosomal function of progranulin, a
guardian against neurodegeneration. *Acta Neuropathol.* **136**, 1 (2018).

28. Lui, H. *et al.* Progranulin Deficiency Promotes Circuit-Specific Synaptic Pruning by
Microglia via Complement Activation. *Cell* **165**, 921–935 (2016).

29. Krabbe, G. *et al.* Microglial NFκB-TNFα hyperactivation induces obsessive-
compulsive behavior in mouse models of progranulin-deficient Frontotemporal
dementia. *Proc. Natl. Acad. Sci. U. S. A.* **114**, 5029–5034 (2017).

30. Martens, L. H. *et al.* Progranulin deficiency promotes neuroinflammation and neuron
loss following toxin-induced injury. *J. Clin. Invest.* **122**, 3955–3959 (2012).

31. Zou, M. *et al.* Autophagy inhibition of hsa-miR-19a-3p/19b-3p by targeting TGF-β R II
during TGF-β1-induced fibrogenesis in human cardiac fibroblasts. *Sci. Rep.* **6**, 1–15
(2016).

32. Whitwell, J. L. *et al.* Neuroimaging signatures of frontotemporal dementia genetics:
C9ORF72, tau, progranulin and sporadics. *Brain* **135**, 794–806 (2012).

33. Benussi, A. *et al.* Toward a glutamate hypothesis of frontotemporal dementia. *Front.*
Neurosci. **13**, (2019).

34. Zhang, J. *et al.* Neurotoxic microglia promote TDP-43 proteinopathy in progranulin
deficiency. *Nature* (2020) doi:10.1038/s41586-020-2709-7.

35. Woollacott, I. O. C. & Rohrer, J. D. The clinical spectrum of sporadic and familial
forms of frontotemporal dementia. *J. Neurochem.* **138**, 6–31 (2016).

36. Sakae, N. *et al.* Microglia in frontotemporal lobar degeneration with progranulin or
C9ORF72 mutations. *Ann. Clin. Transl. Neurol.* **6**, 1782–1796 (2019).

37. Yuan, J., Amin, P. & Ofengeim, D. Necroptosis and RIPK1-mediated
neuroinflammation in CNS diseases. *Nat. Rev. Neurosci.* **20**, 19–33 (2019).

38. Xu, D. *et al.* TBK1 Suppresses RIPK1-Driven Apoptosis and Inflammation during
Development and in Aging. *Cell* **174**, 1477-1491.e19 (2018).

39. Molnár, T. *et al.* Current translational potential and underlying molecular mechanisms
of necroptosis. *Cell Death Dis.* **10**, 1–21 (2019).

40. Brkic, M., Balusu, S., Libert, C. & Vandenbroucke, R. E. Friends or Foes: Matrix
Metalloproteinases and Their Multifaceted Roles in Neurodegenerative Diseases.
Mediators Inflamm. **2015**, (2015).

41. Rivera, S., García-González, L., Khrestchatisky, M. & Baranger, K. Metalloproteinases
and their tissue inhibitors in Alzheimer's disease and other neurodegenerative
disorders. *Cell. Mol. Life Sci.* **76**, 3167–3191 (2019).

42. Kiaei, M. *et al.* Matrix metalloproteinase-9 regulates TNF-α and FasL expression in
neuronal, glial cells and its absence extends life in a transgenic mouse model of
amyotrophic lateral sclerosis. *Exp. Neurol.* **205**, 74–81 (2007).

43. Lorenzl, S. *et al.* The matrix metalloproteinases inhibitor Ro 26-2853 extends survival
in transgenic ALS mice. *Exp. Neurol.* **200**, 166–171 (2006).

44. Lee, J. K. *et al.* Tissue inhibitor of metalloproteinases-3 (TIMP-3) expression is
increased during serum deprivation-induced neuronal apoptosis in vitro and in the
G93A mouse model of amyotrophic lateral sclerosis: A potential modulator of Fas-
mediated apoptosis. *Neurobiol. Dis.* **30**, 174–185 (2008).

45. Vandenbroucke, R. E. *et al.* Matrix metalloproteinase 13 modulates intestinal
epithelial barrier integrity in inflammatory diseases by activating TNF. *EMBO Mol.*
Med. **5**, 1000–1016 (2013).

46. Rempe, R. G., Hartz, A. M. S. & Bauer, B. Matrix metalloproteinases in the brain and
blood-brain barrier: Versatile breakers and makers. *J. Cereb. Blood Flow Metab.* **36**,

1481–1507 (2016).

47.   Woo, M. S., Park, J. S., Choi, I. Y., Kimf, W. K. & Kim, H. S. Inhibition of MMP-3 or -9
868   suppresses lipopolysaccharide-induced expression of proinflammatory cytokines and
869   iNOS in microglia. *J. Neurochem.* **106**, 770–780 (2008).

870   48.   Garcia-Alloza, M. *et al.* Matrix metalloproteinase inhibition reduces oxidative stress
871   associated with cerebral amyloid angiopathy in vivo in transgenic mice. *J. Neurochem.*
872   **109**, 1636–1647 (2009).

873   49.   Aoki, Y. *et al.* C9orf72 and RAB7L1 regulate vesicle trafficking in amyotrophic lateral
874   sclerosis and frontotemporal dementia. *Brain* 887–897 (2017)
875   doi:10.1093/brain/awx024.

876   50.   Yarwood, R., Hellicar, J., Woodman, P. G. & Lowe, M. Membrane trafficking in health
877   and disease. *DMM Dis. Model. Mech.* **13**, (2020).

878   51.   Wren, M. C. *et al.* Frontotemporal dementia-associated N279K tau mutant disrupts
879   subcellular vesicle trafficking and induces cellular stress in iPSC-derived neural stem
880   cells. *Mol. Neurodegener.* **10**, 46 (2015).

881   52.   Reinhardt, P., Glatza, M., Hemmer, K., Tsytsyura, Y. & Thiel, C. S. Derivation and
882   Expansion Using Only Small Molecules of Human Neural Progenitors for
883   Neurodegenerative Disease Modeling. *PLoS One* **8**, 59252 (2013).

884   53.   Dhingra, A. *et al.* Automated production of human induced pluripotent stem cell-
885   derived cortical and dopaminergic neurons with integrated live-cell monitoring. *J. Vis.*
886   *Exp.* **2020**, 1–29 (2020).

887   54.   van Wilgenburg, B., Browne, C., Vowles, J. & Cowley, S. A. Efficient, Long Term
888   Production of Monocyte-Derived Macrophages from Human Pluripotent Stem Cells
889   under Partly-Defined and Fully-Defined Conditions. *PLoS One* **8**, (2013).

890   55.   Takahashi, H., Lassmann, T., Murata, M. & Carninci, P. 5′ end–centered expression
891   profiling using cap-analysis gene expression and next-generation sequencing. *Nat.*
892   *Protoc.* **7**, 542–561 (2012).

893   56.   Skene, N. G. & Grant, S. G. N. Identification of Vulnerable Cell Types in Major Brain
894   Disorders Using Single Cell Transcriptomes and Expression Weighted Cell Type
895   Enrichment. *Front. Neurosci.* **10**, 16 (2016).

896   57.   Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: Summarize analysis
897   results for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048
898   (2016).

899   58.   Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast
900   and bias-aware quantification of transcript expression. *Nat. Publ. Gr.* **14**, (2017).

901   59.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
902   dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, (2014).

903   60.   Menden, K. *et al.* Deep learning–based cell composition analysis from tissue
904   expression profiles. *Sci. Adv.* **6**, eaba2619 (2020).

905   61.   Darmanis, S. *et al.* A survey of human brain transcriptome diversity at the single cell
906   level. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 7285–7290 (2015).

907   62.   Jiang, H., Lei, R., Ding, S. W. & Zhu, S. Skewer: A fast and accurate adapter trimmer
908   for next-generation sequencing paired-end reads. *BMC Bioinformatics* **15**, 182 (2014).

909   63.   Lassmann, T. TagDust2: A generic method to extract reads from sequencing data.

910   *BMC Bioinformatics* **16**, 24 (2015).

911   64.   Dobin, A. & Gingeras, T. R. Mapping RNA-seq Reads with STAR. in *Current Protocols*
912   *in Bioinformatics* (2015). doi:10.1002/0471250953.bi1114s51.

913   65.   Haberle, V., Forrest, A. R. R., Hayashizaki, Y., Carninci, P. & Lenhard, B. CAGEr:
914   Precise TSS data retrieval and high-resolution promoterome mining for integrative
915   analyses. *Nucleic Acids Res.* **43**, e51–e51 (2015).

916   66.   Heinz, S. *et al.* Simple Combinations of Lineage-Determining Transcription Factors
917   Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. *Mol.*
918   *Cell* **38**, 576–589 (2010).

919   67.   Arenillas, D. J. *et al.* CAGEd-oPOSSUM: motif enrichment analysis from CAGE-
920   derived TSSs. *Bioinformatics* **32**, 2858–60 (2016).

921   68.   Khan, A. *et al.* JASPAR 2018: update of the open-access database of transcription
922   factor binding profiles and its web framework. *Nucleic Acids Res.* **46**, D260–D266
923   (2018).

924   69.   Rahman, R. U. *et al.* Oasis 2: Improved online analysis of small RNA-seq data. *BMC*
925   *Bioinformatics* **19**, 54 (2018).

926   70.   Kozomara, A., Birgaoanu, M. & Griffiths-Jones, S. MiRBase: From microRNA
927   sequences to function. *Nucleic Acids Res.* **47**, D155–D162 (2019).

928   71.   Aryee, M. J. *et al.* Minfi: A flexible and comprehensive Bioconductor package for the
929   analysis of Infinium DNA methylation microarrays. *Bioinformatics* **30**, 1363–1369
930   (2014).

931   72.   Leek, J. T. & Storey, J. D. Capturing Heterogeneity in Gene Expression Studies by
932   Surrogate Variable Analysis. *PLoS Genet.* **3**, e161 (2007).

933   73.   Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-
934   sequencing and microarray studies. *Nucleic Acids Res.* (2015)
935   doi:10.1093/nar/gkv007.

936   74.   Benjamini, Y. & Hochberg, Y. Controling the False Discovery Rate: A Practical and
937   Powerful Approach to Multiple Testing. *J. R. Stat. Soc.* (1995).

938   75.   Handl, L., Jalali, A., Scherer, M., Eggeling, R. & Pfeifer, N. Weighted elastic net for
939   unsupervised domain adaptation with application to age prediction from DNA
940   methylation data. in *Bioinformatics* vol. 35 i154–i163 (Oxford University Press, 2019).

Acknowledgements

Post-mortem brain tissue was obtained from the Dutch Brain Bank, Netherlands Institute for Neuroscience, Amsterdam, and from the London Neurodegenerative Disease Brain Bank, King's College London, London, UK. The London Neurodegenerative Disease Brain Bank is part of the Brains for Dementia Research Initiative.

Author contributions

The project was initiated and designed by PH. Small RNA-seq experiments were performed by AF, LK, PR and NF. RNA-seq and CAGE-seq experiments were performed by CB, PR, NF, MC. iPSC-derived neuron and microglia experiments were performed by AD and DKV.

Analysis of CAGE-seq data was done by MF, TN and KM. JSS, BA and KM analysed the RNA-seq data. KM analysed the smRNA-seq and methylation data. PH, SB, PR planned and interpreted all performed analyses. KM, PH, PR and SB wrote the manuscript.

Funding

This work was funded in part by the EU Joint Programme - Neurodegenerative Disease Research (JPND) project: Risk and Modifying factors for FTD (RiMod-FTD), NOMIS Foundation and BMBF IDSN. DKV is in receipt of an Alexander von Humboldt Research Fellowship.

RiMod-FTD Partners

Prof. Dr. P. Heutink: Kevin Menden; Margherita Francescatto; Tenzin Nyima; Cornelis Blauwendraat; Burcu Atarsu; Javier Simon Sanchez; Stefan Bonn; Patrizia Rizzu;

German Center for Neurodegenerative Diseases (DZNE)-Tübingen

Otfried Müller Strasse 23

967 72076 Tübingen

968 Germany

969 Tel.: +49 (0)7071 9254 050

970 E-Mail: peter.heutink@dzne.de

972 Prof. Dr. J. Hardy:

973 Reta Lila Weston Research Laboratories, Dept. of

974 Molecular Neuroscience, UCL Institute of

975 Neurology, London WC1N 3BG, England.

976 Tel: +44 (0) 207 679 4297.

977 E-Mail: j.hardy@UCL.AC.UK

979 Prof. Dr. A. Brice

980 Research Centre of the Brain and Spine Institute

981 (CRicm UMR_S975), Pitié-Salpêtrière Hospital, 47

982 boulevard de l'Hôpital, 75013 Paris, France

983 Tel: +33 1 572746 82

984 E-Mail: alexis.brice@upmc.fr

986 Prof. Dr. J.C. van Swieten; Fenne Riemslagh

987 Prof. Dr. A.B. Smit: Pim van Nierop; Suzanne Miedema

Dept. Neurology, ErasmusMC, 's-Gravendijkwal 230

3015 CE Rotterdam, Netherlands.

Tel: +31 107033274

E-Mail: j.c.vanswieten@erasmusmc.nl

Prof. Dr. P. Longone/

Prof. Dr. E. Buratti/

Prof. Dr. A. Alberici

Molecular Neurobiology Unit

Santa Lucia Foundation

Via del Fosso di Fiorano 64

00179 Roma Italia

Tel. +39-06501703151

E-Mail: p.longone@hsnatalucia.it

1002

Prof. Dr. P. van Damme

Laboratory for Neurobiology (Vesalius Research

Center) O&N IV Herestraat 49 - bus 912.

3000 Leuven, Belgium

Tel. +32 16 3 73181

E-Mail: philip.vandamme@vib-kuleuven.be

1009

1010   Dr. A. Sandelin

1011   Department of Biology, Copenhagen University,

1012   Ole Maaoles vej 5,

1013   DK2200 Copenhagen, Denmark.

1014   Tel +45 35321285/ Email: albin@binf.ku.dk

1015

Prof. Dr. A. Fischer

Dept. of Psychiatry and Psychotherapy, University

of Göttingen, German Center for Neurodegenerative

Diseases (DZNE) Göttingen, Grisebach Str. 5,

37077 Göttingen, Germany.

Tel: +49 551 30178.

E-Mail: afische2@gwdg.de

1023

Prof. Dr. S. Lichtenthaler

Technical University Munich (TUM) and German

Center for Neurodegenerative Diseases (DZNE),

Max-Lebsche Platz 30, 81377 Munich, Germany.

Tel: +49 89 7095 8405

E-Mail: stefanlichtethaler@dzne.de

1030

Prof. Dr. E. Mandelkow + Prof. Dr. E. Mandelkow + Marta Anglada

German Center for Neurodegenerative Diseases

(DZNE), Ludwig-Erhard-Allee 2 53175 Bonn,

Germany.

Tel: +49 (0) 228 / 43302-630

E-Mail: eckhard.mandelkow@dzne.de

1037

1038 Competing Interests

1039   The authors declare no competing interests.

1040