1   ***Streptococcus agalactiae* MprF synthesizes a novel cationic glycolipid and**

2   **contributes to brain entry and meningitis**

4   Luke R. Joyce[a], Haider S. Manzer[b], Jéssica da C. Mendonça[b], Ricardo Villarreal[b], Kelly
5   S. Doran[b#,] Kelli L. Palmer[a#], and Ziqiang Guan[c#]

7   [a]Department of Biological Sciences, The University of Texas at Dallas, Richardson, TX,

8   75080

9   [b]Department of Immunology and Microbiology, University of Colorado School of

10   Medicine, Aurora, CO, 80045

[c]Department of Biochemistry, Duke University Medical Center, Durham, NC, 27710

Running title: GBS MprF glycolipid synthesis and role in meningitis

#Corresponding authors

Kelly S. Doran: Kelly.Doran@CUAnschutz.edu
Kelli L. Palmer: Kelli.Palmer@UTDallas.edu
Ziqiang Guan: Ziqiang.Guan@Duke.edu
18

19   **Keywords:** Group B *Streptococcus*, Lipidomics, Glycolipid, Meningitis, MprF
20
21   Word counts:
22   Abstract: 187
23   Importance: 114
24   Body (incl to end of acknowledgements): 1166.

Abstract

Group B *Streptococcus* (GBS) is a major human pathogen, causing meningitis and severe infection in newborns, yet little is known about its lipid membrane. Here, we investigated the GBS lipid membrane and identify a novel cationic glycolipid, lysyl-glucosyl-diacylglycerol (Lys-Glc-DAG). Multiple peptide resistance factor (MprF) is highly conserved in many bacterial pathogens and plays a critical role in resistance against cationic antimicrobial peptides, cationic bacteriocins, and antibiotics. The MprF protein has been shown to catalyze the amino-acylation of the anionic phospholipid phosphatidylglycerol (PG). Most notably, MprF uses L-lysyl-tRNA to esterify PG with a positively charged lysine to produce lysyl-phosphatidylglycerol (Lys-PG). We demonstrate through heterologous host expression and gene deletion that the GBS MprF has an expanded substrate repertoire and is the biosynthetic enzyme responsible for both Lys-Glc-DAG and Lys-PG biosynthesis in GBS. Furthermore, we demonstrate that MprF contributes specifically to meningitis pathogenesis at the blood-brain barrier both *in vitro* and in an *in vivo* hematogenous murine infection model but does not contribute to bloodstream survival. These results greatly expand our knowledge of MprF functions and reveal insights into the survival mechanisms and pathogenesis of meningitis caused by GBS.

Importance

The lipidomes of many important Gram-positive human pathogens remain largely uncharacterized. By investigating the lipid membrane of *Streptococcus agalactiae* (Group B *Streptococcus*; GBS), an etiological agent responsible for meningitis and severe

diseases in newborns, we uncovered a novel glycolipid and biochemical activity of

Multiple peptide resistance factor (MprF) in GBS. MprF is known to add lysine to

phosphatidylglycerol (PG), forming Lys-PG. We show in GBS that it also adds lysine to

the glycolipid, glucosyl-diacylglycerol (Glc-DAG), to form Lys-Glc-DAG. We demonstrate

the GBS MprF contributes to brain entry and meningitis pathogenesis in mice. These

results expand our knowledge of MprF functions and reveal insights into the survival

mechanisms and pathogenesis of meningitis caused by GBS.

Observation

Streptococcus agalactiae (Group B *Streptococcus*; GBS) is a Gram-positive bacterium that colonizes the lower genital and gastrointestinal tracts of ~30% of healthy women (1, 2). GBS causes sepsis and pneumonia in neonates and is a leading cause of neonatal meningitis, resulting in long-lasting neurological effects in survivors (3-5). New therapeutic and preventative approaches and a more complete understanding of GBS pathogenesis are needed to mitigate the devastating impact of GBS on neonates. Despite the critical role of the bacterial membrane in host-pathogen interactions, little is known about this for GBS. Here, we investigated GBS membrane lipids using normal phase liquid chromatography (NPLC) coupled with electrospray ionization (ESI) high-resolution tandem mass spectrometry (HRMS/MS).

GBS lipidomic profile

The membrane lipids of three GBS clinical isolates of representative serotypes were characterized: COH1 (6), A909 (7), and CNCTC 10/84 (serotypes III, 1a, and V, respectively) (8). Common Gram-positive bacterial lipids were identified by normal phase LC coupled with negative ion ESI/MS/MS, including diacylglyercol (DAG), monohexosyldiacylglycerol (MHDAG), dihexosyldiacylglycerol (DHDAG), phosphatidylglycerol (PG), and lysyl-phosphatidylglycerol (Lys-PG), as shown by the negative total ion chromatogram (TIC) (Fig. 1A).

Surprisingly, the positive TIC (Fig. 1B, Supplemental Figure S1) shows highly abundant peaks of unknown identity at the retention time ~25-29 min. The mass spectra (Fig. 1C) and LC retention times of this lipid do not match with any other bacterial lipids we have

analyzed or exact masses in lipidomic databases. Tandem MS (MS/MS) in the positive

ion mode (Fig. 1D), negative ion mode (Fig. 1E), and high-resolution mass measurement

(Fig. 1C) allowed us to propose lysyl-glucosyl-diacylglyerol (Lys-Glc-DAG) (Fig. 1F) as

the structure of this unknown lipid. Observed and exact masses of Lys-Glc-DAG are

shown in Table S1. The assignment of glucose was based on the observation that

glucosyl-diacylglycerol (Glc-DAG) is a major membrane component of GBS and other

streptococci (9), and results from an isotopic labeling experiment using ^{13}C-labeled

glucose (Glucose-^{13}C$_6$). The assignment of lysine modification was supported by an

isotopic labeling experiment with deuterated lysine (lysine-*d4*). The expected mass shifts

(+4 Da) were observed in both molecular ions and MS/MS product ions (Supplemental

Figure S2). Comparison of both MS/MS spectra of labeled (Glucose-^{13}C$_6$) and unlabeled

Lys-Glc-DAG indicates the lysine residue is linked to the 6-position of glucose

(Supplemental Figure S2). Lys-Glc-DAG consists of several molecular species with

different fatty acyl compositions resulting in different retention times and multiple,

unresolved TIC peaks (~25-29 min).

GBS MprF synthesizes Lys-PG and Lys-Glc-DAG

The enzyme MprF (multiple peptide resistance factor) catalyzes the aminoacylation of PG

with lysine in some Gram-positive pathogens (10, 11). GBS MprF is responsible and

sufficient for synthesizing both Lys-Glc-DAG and Lys-PG. Deletion of *mprF* from COH1

abolishes both Lys-Glc-DAG and Lys-PG synthesis, which are restored by

complementation (Fig. 1G). Deletion of GBS *mprF* does not confer a growth defect in

Todd-Hewitt broth or tissue culture medium. The oral colonizer *Streptococcus mitis* does

not encode *mprF* or synthesize Lys-PG, but synthesizes Glc-DAG and PG (12, 13).

Heterologous expression of GBS *mprF* in *S. mitis* results in Lys-Glc-DAG and Lys-PG

production (Fig. 1H), while expression of *Enterococcus faecium mprF* results in only Lys-

PG production (Fig. 1H), as expected (14). Biosynthetic pathways involving MprF are

shown in Fig. 1I.

MprF contributes to GBS pathogenesis

We investigated whether MprF contributes to GBS invasion into brain endothelial cells

and development of meningitis. To mimic the human blood-brain barrier (BBB), we utilized

the human cerebral microvascular endothelial cell line hCMEC/D3. *In vitro* assays for

adhesion and invasion were performed as described previously (9, 15, 16). There was no

significant difference in the ability of Δ*mprF* compared to WT and complement cells to

attach to hCMEC/D3 cells (Fig. 2A). However, we observed a significant decrease in the

amount of Δ*mprF* recovered from the intracellular compartment of hCMEC/D3 cells (Fig.

2A). The reduced invasion phenotype was confirmed in the hypervirulent serotype V

strain, CJB111 (17, 18) (Supplemental Figure S3). Intracellular survival requires GBS to

survive low pH conditions in lysosomes (pH 4.5 – 5.5) (19), and Δ*mprF* is unable to survive

low pH conditions (Fig. 2B). This suggests that MprF promotes GBS invasion into the

BBB, and intracellular survival.

We hypothesized that these *in vitro* phenotypes of Δ*mprF* would translate into a

diminished ability to penetrate the BBB and produce meningitis *in vivo*. Using our

standard model of GBS hematogenous meningitis (9, 15) mice were challenged with

either WT GBS or Δ*mprF*. Mice were sacrificed at 72 h to determine bacterial loads in

blood and brain tissue. We recovered significantly less CFU in the brains of Δ*mprF*-

infected mice compared to the WT-infected mice (Fig. 2C). However, there was no

significant difference in CFU recovered from the bloodstream (Fig. 2D), demonstrating

that Δ*mprF* does not have a general *in vivo* growth defect. Furthermore, mice challenged

with WT GBS had significantly more leukocyte infiltration, meningeal thickening and

neutrophil chemokine, KC, in brain homogenates compared to Δ*mprF* mutant-infected

animals (Fig. 2E-G). Taken together, *mprF* contributes to GBS penetration into the brain

and to the pathogenesis of meningitis *in vivo*.

136 Conclusion

Bacteria use enzymes such as MprF to fine-tune cell envelope properties for stress

response and virulence. We discovered that GBS MprF uniquely synthesizes a novel and

highly abundant cationic glycolipid, Lys-Glc-DAG, as well as Lys-PG. This establishes

that GBS capitalizes on MprF to modulate charges of both glycolipids and phospholipids

at the membrane. Deletion of *mprF* impacts GBS virulence. Interestingly, this effect is

observed for meningitis, and not for bacteremia, demonstrating that MprF and/or MprF-

synthesized lipids play specific roles in BBB penetration but not *in vivo* survival in general.

In future studies, generating GBS MprF variants that synthesize only Lys-Glc-DAG or

Lys-PG at wild-type levels will allow for identification of the specific roles each lipid plays

in meningitis.

Our identification of the novel Lys-Glc-DAG glycolipid rationalizes further study of the

lipidomes of human pathogens. The decreased *in vivo* pathogenicity of Δ*mprF* identifies

GBS MprF as a candidate for targeting by antimicrobial strategies. Moreover, that Lys-

Glc-DAG is a major GBS membrane component holds promise that Lys-Glc-DAG could

be utilized as a specific molecular biomarker for GBS diagnostics.

Materials and methods

Methods were performed using established techniques; see Supplemental Text S1.

Strains listed in Table S2 and primers Table S3. Illumina sequence reads are deposited

in the Sequence Read Archive, accession PRJNA675025.

Conflict of interest

The authors have declared that no conflict of interest exists.

Acknowledgements

We thank Kathryn Patras at the University of California San Diego for the CNCTC 10/84

strain and Moutusee Islam in Kelli Palmer's lab at The University of Texas at Dallas for

E. faecium 1,231,410 DNA.

The work was supported in part by T32 5T32AI052066-18 for H.S.M, the Coordenação

de Aperfeiçoamento de Pessoal de Nível Superior, Brazil (CAPES; finance code 001 to

J.D.C.M), by grant R01NS116716 from the National Institutes of Health (NIH) to K.S.D

and associated supplement from NINDS to R.V, NIH grant R21AI130666 and the Cecil

H. and Ida Green Chair in Systems Biology Science to K.P, NIH grant R56AI139105 to

K.P and Z.G, and NIH grant U54GM069338 to Z.G.

References

1.   Wilkinson HW. 1978. Group B Streptococcal Infection in Humans. Annual Review
of Microbiology 32:41-57.
2.   Doran KS, Nizet V. 2004. Molecular pathogenesis of neonatal Group B
Streptococcal infection: No longer in its infancy. Molecular Microbiology 54:23-
31.
3.   Hall J, Adams NH, Bartlett L, Seale AC, Lamagni T, Bianchi-Jassir F, Lawn JE,
Baker CJ, Cutland C, Heath PT, Ip M, Le Doare K, Madhi SA, Rubens CE, Saha
SK, Schrag S, Sobanjo-Ter Meulen A, Vekemans J, Gravett MG. 2017. Maternal
Disease With Group B *Streptococcus* and Serotype Distribution Worldwide:
Systematic Review and Meta-analyses. Clinical Infectious Siseases 65:S112-
S124.
4.   Schuchat A. 1998. Epidemiology of Group B Streptococcal disease in the United
States: shifting paradigms. Clinical Microbiology Eeviews 11:497-513.
5.   Edwards MS, Rench MA, Haffar AA, Murphy MA, Desmond MM, Baker CJ. 1985.
Long-term sequelae of Group B Streptococcal meningitis in infants. The Journal
of Pediatrics 106:717-22.
6.   Kuypers JM, Heggen LM, Rubens CE. 1989. Molecular analysis of a region of
the Group B *Streptococcus* chromosome involved in type III capsule expression.
Infection and Immunity 57:3058-65.
7.   Lancefield RC, McCarty M, Everly WN. 1975. Multiple mouse-protective
antibodies directed against Group B Streptococci. Special reference to antibodies
effective against protein antigens. The Journal of Experimental Medicine
142:165-79.
8.   Wilkinson HW. 1977. Nontypable Group B Streptococci isolated from human
sources. Journal of Clinical Microbiology 6:183-4.
9.   Doran KS, Engelson EJ, Khosravi A, Maisey HC, Fedtke I, Equils O, Michelsen
KS, Arditi M, Peschel A, Nizet V. 2005. Blood-brain barrier invasion by Group B
Streptococcus depends upon proper cell-surface anchoring of lipoteichoic acid.
The Journal of Clinical Investigation 115:2499-507.
10.   Peschel A, Jack RW, Otto M, Collins LV, Staubitz P, Nicholson G, Kalbacher H,
Nieuwenhuizen WF, Jung G, Tarkowski A, Van Kessel KPM, Van Strijp JAG.
2001. *Staphylococcus aureus* resistance to human defensins and evasion of
neutrophil killing via the novel virulence factor MprF is based on modification of
membrane lipids with L-lysine. Journal of Experimental Medicine 193:1067-1076.
11.   Roy H, Ibba M. 2008. RNA-dependent lipid remodeling by bacterial multiple
peptide resistance factors. Proceedings of the National Academy of Sciences of
the United States of America 105:4667-4672.

12.   Joyce LR, Guan Z, Palmer KL. 2019. Phosphatidylcholine Biosynthesis in Mitis
Group Streptococci via Host Metabolite Scavenging. Journal of Bacteriology
201:e00495-19.

13.   Adams HM, Joyce LR, Guan Z, Akins RL, Palmer KL. 2017. *Streptococcus mitis*
and *S. oralis* Lack a Requirement for CdsA, the Enzyme Required for Synthesis
of Major Membrane Phospholipids in Bacteria. Antimicrobial Agents and
Chemotherapy 61:e02552-16.

14.   Roy H. 2009. Tuning the properties of the bacterial membrane with
aminoacylated phosphatidylglycerol. IUBMB Life 61:940-953.

15.   Deng L, Spencer BL, Holmes JA, Mu R, Rego S, Weston TA, Hu Y, Sanches GF,
Yoon S, Park N, Nagao PE, Jenkinson HF, Thornton JA, Seo KS, Nobbs AH,
Doran KS. 2019. The Group B Streptococcal surface antigen I/II protein, BspC,
interacts with host vimentin to promote adherence to brain endothelium and
inflammation during the pathogenesis of meningitis. PLoS Pathogens
15:e1007848.

16.   Spencer BL, Deng L, Patras KA, Burcham ZM, Sanches GF, Nagao PE, Doran
KS. 2019. Cas9 contributes to Group B Streptococcal colonization and disease.
Frontiers in Microbiology 10:1-15.

17.   Faralla C, Metruccio MM, De Chiara M, Mu R, Patras KA, Muzzi A, Grandi G,
Margarit I, Doran KS, Janulczyk R. 2014. Analysis of two-component systems in
group B *Streptococcus* shows that RgfAC and the novel FspSR modulate
virulence and bacterial fitness. mBio 5:e00870-14.

18.   Spencer BL, Chatterjee A, Duerkop BA, Baker CJ, Doran KS. 2021. Complete
Genome Sequence of Neonatal Clinical Group B Streptococcal Isolate CJB111.
Microbiology Resource Announcements 10.

19.   Mu R, Cutting AS, Del Rosario Y, Villarino N, Stewart L, Weston TA, Patras KA,
Doran KS. 2016. Identification of CiaR Regulated Genes That Promote Group B
Streptococcal Virulence and Interaction with Brain Endothelial Cells. PLoS One
11:e0153891.

Figure legends

Fig 1. Total ion chromatogram (TIC) of LC/MS in A) negative ion mode, B) positive ion mode shows a major unknown lipid eluting at ~25-29 min. C) Positive ESI/MS showing the [M+H]$^+$ ions of the unknown lipid. D) Positive ion MS/MS spectrum of [M+H]$^+$ at m/z 885.6 and E) negative ion MS/MS spectrum of [M-H]$^-$ at m/z 883.6 of the unknown lipid. F) Lys-Glc-DAG (16:0/18:1) is proposed as the structure of the unknown lipid. G) TIC showing loss of Lys-Glc-DAG and Lys-PG in COH1$\Delta mprF$ which is present when $mprF$ is complemented *in trans*. H) Lys-Glc-DAG and Lys-PG is only present in *S. mitis* when expressing GBS $mprF$ compared to Lys-PG only when expressing *E. faecium mprF*. "*" denotes methylcarbamate of Lys-Glc-DAG, an extraction artifact due to the use of chloroform. I) Biosynthetic pathways involving MprF.

Fig 2. A) *In vitro* assays for adherence and invasion of hCMEC cells indicates *mprF* contributes to invasion but not adherence to brain endothelium (mean of 3 replicate experiments with 4 technical replicates, mean and SEM). B) pH-adjusted medium growth indicates $\Delta mprF$ cannot survive in low pH conditions, mean and SD. Groups of CD-1 mice were injected intravenously with COH1 WT or COH1$\Delta mprF$ strains and bacterial counts were assessed in the C) brain and D) blood after 72h. Representative data from 2 independent experiments are shown (WT, $n = 20$; $\Delta mprF$, $n = 19$). E) Hematoxylin-eosin-stained brain sections from representative mice infected with WT (top) or $\Delta mprF$ mutant (bottom); arrows indicate meningeal thickening and leukocyte infiltration. F) Quantification of meningeal thickening using ImageJ. G) KC chemokine production measured by ELISA. Panels C,D,F, and G) median indicated. Statistical analyses performed using GraphPad Prism: A) One-way ANOVA with Tukey's multiple comparisons test; C,D,F) unpaired two-tailed t-test; G) Mann-Whitney U test; p-values indicated; ns, no significance (p-value > 0.05).

Supplemental files

Supplemental Text S1: Materials and Methods

Supplemental Figure S1. Detection of Lys-PG and Lys-Glc-DAG in *S. agalactiae* A909 and *S. agalactiae* CNCTC 10/84.

Supplemental Figure S2. Isotopic incorporation of deuterated lysine and ^{13}C-labeled glucose into Lys-Glc-DAG and Lys-PG.

Supplemental Figure S3. *In vitro* hCMEC adhesion and invasion of CJB111 strains.

Supplemental Table S1. Observed and calculated exact masses of the [M+H]$^+$ ions of Lys-Glc-DAG molecular species in *S. agalactiae* COH1.

278   Supplemental Table S2. Strains and plasmids used in this study.

279   Supplemental Table S3. Primers used in this study.

Fig 1. Total ion chromatogram (TIC) of LC/MS in A) negative ion mode, B) positive ion mode shows a major unknown lipid eluting at ~25-29 min. C) Positive ESI/MS showing the [M+H]$^+$ ions of the unknown lipid. D) Positive ion MS/MS spectrum of [M+H]$^+$ at m/z 885.6 and E) negative ion MS/MS spectrum of [M-H]$^-$ at m/z 883.6 of the unknown lipid. F) Lys-Glc-DAG (16:0/18:1) is proposed as the structure of the unknown lipid. G) TIC showing loss of Lys-Glc-DAG and Lys-PG in COH1Δ*mprF* which is present when *mprF* is complemented *in trans*. H) Lys-Glc-DAG and Lys-PG is only present in *S. mitis* when expressing GBS *mprF* compared to Lys-PG only when expressing *E. faecium mprF*. "*" denotes methylcarbamate of Lys-Glc-DAG, an extraction artifact due to the use of chloroform. I) Biosynthetic pathways involving MprF.

Fig 2. A) *In vitro* assays for adherence and invasion of hCMEC cells indicates *mprF* contributes to invasion but not adherence to brain endothelium (mean of 3 replicate experiments with 4 technical replicates, mean and SEM). B) pH-adjusted medium growth indicates Δ*mprF* cannot survive in low pH conditions, mean and SD. Groups of CD-1 mice were injected intravenously with COH1 WT or COH1Δ*mprF* strains and bacterial counts were assessed in the C) brain and D) blood after 72h. Representative data from 2 independent experiments are shown (WT, *n* = 20; Δ*mprF, n* = 19). E) Hematoxylin-eosin-stained brain sections from representative mice infected with WT (top) or Δ*mprF* mutant (bottom); arrows indicate meningeal thickening and leukocyte infiltration. F) Quantification of meningeal thickening using ImageJ. G) KC chemokine production measured by ELISA. Panels C,D,F, and G) median indicated. Statistical analyses performed using GraphPad Prism: A) One-way ANOVA with Tukey's multiple comparisons test; C,D,F) unpaired two-tailed t-test; G) Mann-Whitney U test; p-values indicated; ns, no significance (p-value > 0.05).

Supplemental Text, Figures, and Tables

Streptococcus agalactiae **MprF synthesizes a novel cationic glycolipid and contributes to**

brain entry and meningitis

Luke R. Joyce[a], Haider S. Manzer[b], Jéssica da C. Mendonça[b], Ricardo Villarreal[b], Kelly S.
Doran[b#,] Kelli L. Palmer[a#], and Ziqiang Guan[C#]

[a]Department of Biological Sciences, The University of Texas at Dallas, Richardson, TX, 75080

[b]Department of Immunology and Microbiology, University of Colorado School of Medicine,

Aurora, CO, 80045

[c]Department of Biochemistry, Duke University Medical Center, Durham, NC, 27710

[#]Corresponding authors

Supplemental Text S1: Materials and Methods

Bacterial strains, media, and growth conditions

GBS strains were grown statically at 37°C in Todd-Hewitt Broth (THB) and *S. mitis* strains were

grown statically at 37°C and 5% CO_2, unless otherwise stated. Streptococcal chemically defined

medium (20) was diluted from stock as described (21) with 1% w/v glucose (referred to as DM),

slightly modified from (22), unless otherwise stated. *Escherichia coli* strains were grown in

Lysogeny Broth (LB) at 37°C with rotation at 225 rpm. Kanamycin and erythromycin (Sigma-

Aldrich) were supplemented to media at 50 µg/mL and 300 µg/mL for *E. coli,* respectively, or 300

µg/mL and 5 µg/mL, respectively, for streptococcal strains.

Routine molecular biology techniques

All PCR reactions utilized Phusion polymerase (Thermo Fisher). PCR products and restriction

digest products were purified using GeneJET PCR purification kit (Thermo Fisher) per

manufacturer protocols. See Table S3 for primers. Plasmids were extracted using GeneJET

plasmid miniprep kit (Thermo Fisher) per manufacturer protocols. Restriction enzyme digests

utilized XbaI, XhoI, and PstI (New England Biolabs) for 3 h at 37°C in a water bath. Ligations

utilized T4 DNA ligase (New England Biolabs) at 16°C overnight or Gibson Assembly Master Mix

(New England Biolabs) per manufacturer protocols where stated. All plasmid constructs were

sequence confirmed by Sanger sequencing (Massachusetts General Hospital DNA Core or CU

Anschutz Molecular Biology Core).

Deuterated lysine and $^{13}C_6$-D-glucose isotope tracking

A GBS COH1 colony was inoculated into 15 mL of DM containing 450 μM lysine-*d4* (Cambridge Isotopes Laboratories) or a single COH1 colony was inoculated into 10 mL DM supplemented with 0.5% w/v $^{13}C_6$D-glucose (U-13C6, Cambridge Isotopes Laboratories) for overnight growth for lipidomic analysis described below.

Construction of MprF expression plasmids

Genomic DNA was isolated using the Qiagen DNeasy Blood and Tissue kit per the manufacturer's protocol with the exception that cells were pre-treated with 180 μL 50 mg/mL lysozyme, 25 μL 2500 U/mL mutanolysin, and 15 μL 20 mg/mL pre-boiled RNase A and incubated at 37°C for 2 h. The *mprF* genes from GBS COH1, (GBSCOH1_1931), GBS CJB111 (ID870_10050), and *E. faecium* 1,231,410 (EFTG_00601) were amplified and either Gibson ligated into pABG5Δ*phoZ* (23) or ligated into pDCErm (24). Plasmid constructs were transformed into chemical competent *E. coli*. Briefly, chemically competent cells were incubated for 10 min on ice with 5 μL of Gibson reaction before heat shock at 42°C for 70 sec, then placed on ice for 2 min before 900 μL of cold SOC medium was added. Outgrowth was performed at 37°C, with shaking at 225 rpm, for 1 h. Cultures were plated on LB agar plates containing 50 μg/mL kanamycin. Colonies were screened by PCR for presence of the *mprF* insert.

Expression of *mprF* in *S. mitis*

Natural transformation was performed as previously described (12). Briefly, precultures were thawed at room temperature, diluted in 900 μL of THB, further diluted 1:50 in prewarmed 5 mL THB, and incubated for 45 min at 37°C. 500 μL of culture was then aliquoted with 1 μL of 1 mg/ml

competence-stimulating peptide (EIRQTHNIFFNFFKRR) and 1 μg/mL plasmid. Transformation

reaction mixtures were cultured for 2 h at 37°C in microcentrifuge tubes before being plated on

THB agar supplemented with 300 μg/mL kanamycin. Single transformant colonies were cultured

in 15 mL THB overnight. PCR was used to confirm the presence of the *mprF* insert on the plasmid.

Plasmids were extracted and sequence confirmed as described above. Lipidomics was performed

as described below in biological triplicate.

Construction of *mprF* deletion plasmid

Regions ~2 kb upstream and downstream of the GBS COH1 *mprF* (GBSCOH1_1931) or CJB111

(ID870_10050) were amplified using PCR. Plasmid, pMBSacB (25), and the PCR products were

digested using appropriate restriction enzymes and ligated overnight. 7 μL of the ligation reaction

was transformed into chemically competent *E. coli* DH5α as described above, except that

outgrowth was performed at 28°C with shaking at 225 rpm for 90 min prior to plating on LB agar

supplemented with 300 μg/mL erythromycin. Plates were incubated at 28°C for 72 h. Colonies

were screened by PCR for correct plasmid construction. Positive colonies were inoculated into 50

mL LB media containing antibiotic and incubated at 28°C with rotation at 225 rpm for 72 h.

Cultures were pelleted using a Sorvall RC6+ centrifuge at 4,280 x *g* for 6 min at room temperature.

Plasmid was extracted as described above except the cell pellet was split into 5 columns to

prevent overloading and serial eluted into 50 μL. Plasmid construction was confirmed via

restriction digest using XhoI and XbaI, and the insert was PCR amplified and sequence-verified.

Generation of electrocompetent GBS cells for *mprF* knockout

Electrocompetent cells were generated as described (25) with minor modifications. Briefly, a GBS COH1 or CJB111 colony was inoculated in 5 mL M17 medium (BD Bacto) with 0.5% glucose and grown overnight at 37°C. The 5 mL was used to inoculate a second overnight culture of 50 mL pre-warmed filter-sterilized M17 medium containing 0.5% glucose, 0.6% glycine, and 25% PEG 8000. The second overnight was added to 130 mL of the same medium and grown for 1 h at 37°C. Cells were pelleted at 3,200 x *g* in a Sorvall RC6+ at 4°C for 10 min. Cells were washed twice with 25 mL cold filter-sterilized GBS wash buffer containing 25% PEG 8000 and 10% glycerol in water, and pelleted as above. Cell pellets were re-suspended in 1 mL GBS wash buffer and either used immediately for transformation or stored in 100 µL aliquots at -80°C until use.

Deletion of GBS COH1 and CJB111 *mprF*

Electrocompetent cells were generated as described (25) with minor modifications. The double crossover homologous recombination knockout strategy was performed as described previously (16, 25, 26) with minor modifications. 1 µg of plasmid was added to electrocompetent GBS cells and transferred to a cold 1 mm cuvette (Fisher or BioRad). Electroporation was carried out at 2.5 kV on an Eppendorf eporator. 1 mL of THB containing 12.5% PEG 8000, 20 mM $MgCl_2$, and 2 mM $CaCl_2$ was immediately added and then the entire reaction was transferred to a glass culture tube. Outgrowth was at 28°C for 2 h followed by plating on THB agar supplemented with 5 µg/mL erythromycin. Plates were incubated for 48 h at 28°C. A single colony was cultured overnight in 5 mL THB with 5 µg/mL erythromycin at 28°C. The culture was screened via PCR for the plasmid insert with the initial denaturing step extended to 10 min. The overnight culture was diluted 1:1000 THB containing 5 µg/mL erythromycin and incubated overnight at 37°C to promote single cross over events. The culture was then serial diluted and plated on THB agar plates with antibiotic and

incubated at 37°C overnight. Colonies were screened for single crossover events by PCR. Single

crossover colonies were inoculated in 5 mL THB at 28°C to promote double crossover events.

Overnight cultures were diluted 1:1000 into 5 mL THB containing sterile 0.75 M sucrose and

incubated at 37°C. Overnight cultures were serial diluted and plated on THB agar and incubated

at 37°C overnight. Colonies were patched onto THB agar with and without 5 μg/mL erythromycin

to confirm loss of plasmid. Colonies were also screened by PCR for the loss of *mprF*. Colonies

positive for the loss of *mprF* were inoculated into 5 mL THB at 37°C. Cultures were stocked and

gDNA extracted as described above, with minor modifications. Sequence confirmation of the

mprF knockout was done via Sanger sequencing (Massachusetts General Hospital DNA Core or

CU Anschutz Molecular Biology Core). The mutant was grown overnight in 15 mL THB at 37°C

and pelleted at 6,150 x *g* for 5 min in a Sorvall RC6+ centrifuge at room temperature for lipid

extraction as described. Genomic DNA of COH1Δ*mprF* was isolated as described above and

whole genome sequencing was performed in paired-end reads (2 by 150 bp) on the Illumina

NextSeq 550 platform at the Microbial Genome Sequencing Center (Pittsburgh, PA).

Complementation of *mprF* in COH1Δ*mprF* and CJB111Δ*mprF*

Electrocompetent COH1Δ*mprF* were generated as previously described (27). Briefly, GBSΔ*mprF*

was inoculated into 5 mL THB with 0.6% glycine and grown overnight. The culture was expanded

to 50 mL in pre-warmed THB with 0.6% glycine and grown to an OD_{600} nm of 0.3 and pelleted for

10 min at 3200 x *g* at 4°C in a Sorvall RC6+ floor centrifuge. The pellet was kept on ice through

the remainder of the protocol. The pellet was washed twice with 25 mL and once with 10 mL of

cold 0.625 M sucrose and pelleted as above. The cell pellet was resuspended in 400 μL of cold

20% glycerol, aliquoted in 50 μL aliquots, and used immediately or stored at -80°C until use.

Electroporation was performed as described above, with recovery in THB supplemented with 0.25

M sucrose, and plated on THB agar with kanamycin at 300 μg/mL.

Acidic Bligh-Dyer extractions

Centrifugation was performed using a Sorvall RC6+ centrifuge. Cultures were pelleted at 4,280 x

g for 5 min at room temperature unless otherwise stated. The supernatants were removed, and

cell pellets were stored at -80°C until acidic Bligh-Dyer lipid extractions were performed as

described (12). Briefly, cell pellets were resuspended in 1X PBS (Sigma-Aldrich) and transferred

to Coring Pyrex glass tubes with PTFR-lined caps (VWR), followed by 1:2 vol:vol

chloroform:methanol addition. Single phase extractions were vortexed periodically and incubated

at room temperature for 15 minutes before 500 x *g* centrifugation for 10 min. A two-phase Bligh-

Dyer was achieved by addition of 100 µL 37% HCl, 1 mL CHCl$_3$, and 900 µl of 1X PBS, which

was then vortexed and centrifuged for 5 min at 500 x *g.* The lower phase was removed to a new

tube and dried under nitrogen before being stored at -80°C prior to lipidomic analysis.

Liquid Chromatography/Electrospray Ionization Mass Spectrometry

Normal phase LC was performed on an Agilent 1200 quaternary LC system equipped with an

Ascentis silica HPLC column (5 µm; 25 cm by 2.1 mm; Sigma-Aldrich) as described previously

(28, 29).   Briefly, mobile phase A consisted of chloroform-methanol-aqueous ammonium

hydroxide (800:195:5, vol/vol), mobile phase B consisted of chloroform-methanol-water-aqueous

ammonium hydroxide (600:340:50:5, vol/vol), and mobile phase C consisted of chloroform-

methanol-water-aqueous ammonium hydroxide (450:450:95:5, vol/vol). The elution program

consisted of the following: 100% mobile phase A was held isocratically for 2 min, then linearly

increased to 100% mobile phase B over 14 min, and held at 100% mobile phase B for 11 min.

The LC gradient was then changed to 100% mobile phase C over 3 min, held at 100% mobile

phase C for 3 min, and, finally, returned to 100% mobile phase A over 0.5 min and held at 100%

mobile phase A for 5 min. The LC eluent (with a total flow rate of 300 µl/min) was introduced into

the ESI source of a high-resolution TripleTOF5600 mass spectrometer (Sciex, Framingham, MA).

Instrumental settings for negative-ion ESI and MS/MS analysis of lipid species were: IS = -4,500

V, CUR = 20 psi, GSI = 20 psi, DP = -55 V, and FP = -150V. Settings for positive-ion ESI and

MS/MS analysis were: IS = +5,000 V, CUR = 20 psi, GSI = 20 psi, DP = +55 V, and FP = +50V.

The MS/MS analysis used nitrogen as the collision gas. Data analysis was performed using

Analyst TF1.5 software (Sciex, Framingham, MA).

pH-adjusted THB growth

Approximately 30 mL of fresh THB were adjusted to different pH values, measured using a Mettler

Toledo FiveEasy pH/MV meter, and sterile-filtered using 0.22 µM syringe filters. A final volume of

200 µL culture medium was aliquoted per well in a flat-bottom 96 well plate (Falcon); culture media

were not supplemented with antibiotics. Overnight cultures of GBS strains were used to inoculate

the wells to a starting OD_{600nm} 0.02 per well. Plates were incubated for 24 h at 37°C before OD_{600nm}

was read using a BioTek MX Synergy 2 plate reader. This experiment was performed in biological

triplicate.

hCMEC cell adherence and invasion assays

Human Cerebral Microvascular Endothelial cells hCMEC/D3 (obtained from Millipore) were grown

in EndoGRO-MV complete media (Millipore, SCME004) supplemented with 5% fetal bovine

serum (FBS) and 1 ng/ml fibroblast growth factor-2 (FGF-2; Millipore). Cells were grown in tissue

culture treated 24 well plates and 5% CO_2 at 37°C.

Assays to determine the total number of bacteria adhered to host cells or intracellular bacteria

were performed as described previously (15, 16). Briefly, bacteria were grown to mid log phase

(OD_{600nm} 0.4-0.5) and normalized to 1×10^8 to infect cell monolayers at a multiplicity of infection

(MOI) of 1 (1×10^5 CFU per well). The total cell-associated GBS were recovered after 30 min

incubation. Cells were washed slowly five times with 500 µL 1X PBS (Sigma) and detached by

addition of 100 µL of 0.25% trypsin-EDTA solution (Gibco) and incubation for 5 min before lysing

the eukaryotic cells with the addition of 400 µL of 0.025% Triton X-100 (Sigma) and vigorous

pipetting. The lysates were then serially diluted and plated on THB agar and incubated overnight

to determine CFU. Bacterial invasion assays were performed as described above except infection

plates were incubated for 2 h before incubation with 100 µg gentamicin (Sigma) and 5 µg penicillin

(Sigma) supplemented media for an additional 2 h to kill all extracellular bacteria, prior to being

trypsinized, lysed, and plated as described. Experiments were performed in biological triplicate

with four technical replicates per experiment.

Murine model of GBS hematogenous meningitis

All animal experiments were conducted under the approval of the Institutional Animal Care and

Use Committee (#00316) at the University of Colorado Anschutz Medical Campus and

performed using accepted veterinary standards. The murine meningitis model was performed as

previously described (16, 30, 31). Briefly, 7-week-old male CD1 (Charles River) mice were

challenged intravenously with 1×10^9 CFU of WT COH1 or the isogenic Δ*mprF* mutant. At 72 h

post-infection, mice were euthanized and blood and brain tissue were harvested, homogenized,

and serially diluted on THB agar plates to determine bacterial CFU.

Histology and ELISA

Mouse brain tissue was frozen in OCT compound (Sakura) and sectioned using a CM1950

cryostat (Leica). Sections were stained using hematoxylin and eosin (Sigma) and images were

taken using a BZ-X710 microscope (Keyence). Images were analyzed using ImageJ software.

Meningeal thickening was quantified from sections taken from three different mice per group, and

six images per slide. Meningeal thickening was quantified across two points per image. KC protein

from mouse brain homogenates was detected by enzyme-linked immunosorbent assay according

to the manufacturer's instructions (R&D systems).

Supplemental Figure S1. Detection of Lys-PG and Lys-Glc-DAG in *S. agalactiae* A909 and
S. agalactiae CNCTC 10/84. Positive TICs (left panels) showing the presence of Lys-PG and
Lys-Glc-DAG in S. *agalactiae* A909 and S. *agalactiae* CNCTC 10/84. Mass spectra (right panels)
show the [M+H]⁺ ions of Lys-Glc-DAG.

Supplemental Figure S2. Isotopic incorporation of deuterated lysine and ¹³C-labeled glucose into Lys-Glc-DAG and Lys-PG. The lipid extracts of *S. agalactiae* COH1 cultured in DM, DM supplemented with 450 µM L-lysine-*d4* (4,4,5,5-D4), or in DM containing 0.5% w/v D-Glucose (U-¹³C₆) were analyzed by LC-ESI/MS in the positive ion mode. A) Negative ESI/MS of [M-H]⁻ ions of major Lys-Glc-DAG species in *S. agalactiae* COH1 when cultured in DM supplemented with lysine-*d4*. The incorporation of lysine-*d4* into Lys-Glc-DAG is evidenced by an upward *m/z* shift of 4 Da of the [M-H]⁻ ion (from *m/z* 883 to *m/z* 887). B) MS/MS of [M-H]⁻ at *m/z* 883.6 produces a deprotonated lysine residue at *m/z* 145. C) MS/MS of [M-H]⁻ at *m/z* 887.6 produces a deprotonated lysine-*d4* residue at *m/z* 149. D) [M+H]⁺ ions of major Lys-PG species in *S. agalactiae* COH1 cultured in DM supplemented with lysine-*d4.* The incorporation of lysine-*d4* in Lys-PG is evidenced by an upward *m/z* shift of 4 Da from unlabeled Lys-PG (blue dot) to labeled Lys-PG (red dot). E) MS/MS of 885.6. A major product ion at *m/z* 291.1 is derived from glucose-lysine residue. F) MS/MS of *m/z* 900.7 (containing fifteen ¹³C atoms). The presence of *m/z* 297.1 (with 6 Da shift) is consistent with glucose in Lys-Glc-DAG is replaced with D-Glucose (U-¹³C₆). The other nine ¹³C atoms are incorporated into the DAG portion of Lys-Glc-DAG. Furthermore, MS/MS data indicate that lysine is linked to the C6 position of glucose by the fragmentation schemes for forming *m/z* 189 ion from the unlabeled Lys-Glc-DAG and *m/z* 191 ion from the ¹³C-labeled Lys-Glc-DAG.

Supplemental Figure S3. *In vitro* hCMEC adhesion and invasion of CJB111 strains. *In vitro*
assays for adherence and invasion of hCMEC cells indicates *mprF* contributes to invasion but not
adherence to brain endothelium. Data indicates the percentage of the initial inoculum that was
recovered. Experiments were performed three times with each condition in quadruplicate. Data
from one representative experiment is shown, mean and standard deviation indicated. One-Way
ANOVA with Tukey's multiple comparisons statistical test was used. P-values indicated; ns, not
significant.

554 **Supplemental Table S1. Observed and calculated exact masses of the [M+H]$^+$ ions of Lys-**
555 **Glc-DAG molecular species in *S. agalactiae* COH1.**

Lys-Glc-DAG[1]	[M+H]$^+$	
	Observed mass	Exact mass
C28:1	801.575	801.583
C28:0	803.595	803.599
C30:1	829.610	829.615
C30:0	831.624	831.630
C32:2	855.623	855.630
C32:1	857.644	857.646
C34:2	883.657	883.622
C34:1	885.672	885.677
C36:2	911.686	911.693
C36:1	913.703	913.709

556 [1]The numbers before and after colons indicate the total acyl chain carbon atoms and double
557 bonds, respectively.

 Supplemental Table S2. Strains and plasmids used in this study.

Organism	Strain	Description	Ref
S. agalactiae	ATCC BAA-1176 (COH1)	Wild-type *S. agalactiae* strain, serotype III	(6)
	COH1Δ*mprF*	*mprF* (GBSCOH1_1931) deletion strain	This work
	COH1Δ*mprF(*pABG5)	Empty vector control strain	This work
	COH1Δ*mprF*(pGBSMprf)	Expresses GBS *mprF* from P$_{prtF}$ in pABG5Δ*phoZ*	This work
	COH1(pABG5)	Empty vector control	This work
	CJB111	Wilt-type *S. agalactiae* strain, serotype V	(17, 18)
	CJB111Δ*mprF*	*mprF* (ID870_10050) deletion strain	This work
	CJB111Δ*mprF*(pDCErm)	Empty vector control strain	This work
	CJB111Δ*mprF*(pJMprF)	Expresses GBS *mprF* in pDCErm	This work
	ATCC BAA-1138 (A909)	Wild-type *S. agalactiae* strain, serotype Ia	(7)
	CNCTC 10/84	Wild-type *S. agalactiae* strain, serotype V. Obtained from Dr. K Patras, UCSD	(8, 32)
S. mitis	ATCC 49456	Wild-type *S. mitis* type strain, also known as NCTC 12261	(33)
	ATCC 49456(pABG5)	Empty vector control	This work
	ATCC 49456(pGBSMprF)	Expresses GBS *mprF* from P$_{prtF}$ in pABG5Δ*phoZ*	This work
	ATCC 49456(pEfmMprF1)	Expresses *E. faecium mprF1* from P$_{prtF}$ in pABG5Δ*phoZ*	This work
	ATCC 49456(pEfmMprF2)	Expresses *E. faecium mprF1* from P$_{prtF}$ in pABG5Δ*phoZ*	This work
E. coli	DH5α	Plasmid cloning host; F$^-$, φ80*lacZ*ΔM15, *rec*A1, *end*A1, *hsd*R17, *pho*A, *s up*E44, λ$^-$ *thi*-1, *gyr*A96, *rel*A1	(34)
	DH5α(pABG5)	Empty vector control	This work
	MC1061	Plasmid cloning host; F$^-$, araD139, Δ(araABC-leu)7696, Δ(lac)X74, galU, galK, hsdR2, (r$_K^-$ m$_K^+$), mcrB1, rpsL, (Str^r)	(35)
	MC1061(pDCErm)	Empty vector control	This work

DH5α(pGBSMprF)	Expresses COH1 *mprF* (GBSCOH1_1931) from P_{prtF} in pABG5Δ*phoZ*	This work
DH5α(pEfmMprF1)	Expresses *E. faecium mprF1* from P_{prtF} in pABG5Δ*phoZ*	This work
DH5α(pEfmMprF2)	Expresses *E. faecium mprF1* from P_{prtF} in pABG5Δ*phoZ*	This work
DH5α(pMBMprFKO)	Allelic exchange plasmid containing ~2 kb sequence flanking GBSCOH1_1931	This work
MC1061(pJMprFKO)	Allelic exchange plasmid containing ~2 kb sequence flanking ID870_10050	This work
MC1061(pJMprF)	Expresses CJB11 *mprF* from $P_{tetM/erm}$ in pDCErm	This work
E. faecium	1,231,410	Wild type *E. faecium* strain	(36)

Plasmid	Description	Ref
pABG5Δ*phoZ*	Constitutive expression vector for streptococci with the P_{prtF} promoter. Confers kanamycin resistance. Referred to as pABG5 throughout the text	(23)
pGBSMprF	pABG5Δ*phoZ* expressing COH1 *mprF* (GBSCOH1_1931) from P_{prtF}	This work
pEfmMprF1	pABG5Δ*phoZ* expressing *E. faecium* 1,231,410 *mprF1* (EFTG_00601) from P_{prtF}	This work
pMBSacB	Allelic exchange plasmid for *S. agalactiae*. Confers erythromycin resistance and sucrose sensitivity	(25)
pMBMprFKO	Knockout plasmid containing ~2 kb sequence flanking GBSCOH1_1931	This work
pJMprFKO	Knockout plasmid containing ~2 kb sequence flanking ID870_10050	This work
pDCErm	Consitituitive expression vector for streptococcus from $P_{tetM/erm}$	(24)
pJMprF	pDCErm expressing CJB111 *mprF* (ID870_10050)	This work

 Supplemental Table S3. Primers used in this study.

Primer	5' – 3' sequence	Use
GBS_MprF_F	GAGAGGTCCCTTTCCTTGAAAAAGCTAATTGAAAAAGTC	Amplify GBSCOH1_1931 for Gibson assembly
GBS_MprF_R	ACCAATACCTTTATCTTATTTAACAATCTTAATTTTACTATC	Amplify GBSCOH1_1931 for Gibson assembly
Faec_MprF1_F	GAGAGGTCCCTTTCCTTGTTAAAAAATACCATACAATG	Amplify EFTG_00601 for Gibson assembly
Faec_MprF1_R	ACCAATACCTTTATCTTAATACTTTCTTCGTATCC	Amplify EFTG_00601 for Gibson assembly
MpF_SacII	ACGTCACCGCGGTTGAAAAAGCTAATTGAAAAAGTC	Amplify CJB111 *mprF* ID870_10050 for ligation
MpR_BamHI	ACGTCAGGATCCTTATTTAACAATCTTAATTTTACTATC	Amplify CJB111 *mprF* ID870_10050 for ligation
pABG5-5'	GGAAAGGGACCTCTCTTCCTAAAC	Linearize pABG5Δ*phoZ* for Gibson assembly
pABG5-3'	GATAAAGGTATTGGTAAATAACAAA	Linearize pABG5Δ*phoZ* for Gibson assembly
Expression plasmid sequencing		
GBS_S1	GAATGGAATAATATAGTAGGCT	For sequencing pGBSMprF/pJMprF, amplifies with pABG5_Fup2/ pF
GBS_S2	GATTGTATCCCTTATTCC	For sequencing pGBSMprF/pJMprF, amplifies with GBS_S3
GBS_S3	CGATTCAATAGCTTCAC	For sequencing pGBSMprF/pJMprF, amplifies with GBS_S2
GBS_S4	GATAAAGGCTCTACTGG	For sequencing pGBSMprF/pJMprF, amplifies with pABG5_FDwn/pR
pABG5_FDwn	CCAATAATAATGACTAGAGAAG	For pABG5 plasmid insert sequencing
pABG5_Fup2	CAAAAGGTTTCGACTTTTCACC	For pABG5 plasmid insert sequencing
EF1_S1	GAATAACGCTGATCAAAGT	For sequencing pEfmMprF1, amplifies with pABG5_Fup2
EF1_S2	TGCCAAGAGAAATAGTC	For sequencing pEfmMprF1, amplifies with EF1_S3
EF1_S3	ACAATCTCTTCGCTTG	For sequencing pEfmMprF1, amplifies with EF1_S2
EF1_S4	CCAACTGTTCTTCTCCAA	For sequencing pEfmMprF1, amplifies with pABG5_FDwn
pF	AGCGCTAGGAGGAAAC	For pDCErm plasmid insert sequencing
pR	CCCATGCCATCTCCAATC	For pDCErm plasmid insert sequencing
GBSCOH1_1931 knockout plasmid construction, sequencing, and integration screening		
Mp1F_PstI	ACGTCACTGCAGTTCAATTAGCTTTTTCAACAATTTC	Amplifies upstream fragment from within GBSCOH1_1931/ID870_10050 leaving 6 codons, with Mp1R_XhoI
Mp1R_XhoI	ACGTCACTCGAGGCTGTTTATGGTGCTTTG	5' most primer of upstream fragment, amplifies with Mp1F_PstI
Mp2F_XbaI	ACGTCATCTAGAGAAAAGGCTAGATTACGAAC	3' most primer of downstream fragment, amplifies with Mp2R_PstI
Mp2R_PstI	ACGTCACTGCAGGTTAAATAAGCTTTATTTGGCA	Amplifies downstream fragment leaving 2 codons and stop codon of GBSCOH1_1931/ID870_10050, with Mp2F_XbaI
T7 promoter	TAATACGACTCACTATAGGG	Amplifies with MpS5F below to sequence plasmid, amplifies with T3 promoter for insert screening and plasmid presence
T3 promoter	AATTAACCCTCACTAAAGGG	Amplifies with MpS3R below, amplifies with T7 promoter for insert screening and plasmid presence
Int_F	GCTAATTGAACTGCAGGTTAAATAAG	Anneals at *mprF* knockout site, amplifies with Out_R for single integration screening

Out_R	GCTATTATATTTAGTGGTTTAATTGG	Anneals outside recombination arms, amplifies with Int_F, for single integration screening

Genomic knockout region sequencing

MpS3F	CATTAGCTAGTCTTATCGGAG	Anneals outside integration arms, amplifies with MpS3R
MpS3R	ACAGCTACTTGGTAGTTCA	Amplifies with MpS3F
MpS4F	GCTACTAAGGCAAGATACG	Amplifies with MpS4R, knockout screening and plasmid sequencing primer
MpS4R	ATGGTCAGCGATGGTG	Amplifies with MpS4F, knockout screening and plasmid sequencing primer
MpS5F	CATAAGCGAAATAACTTGAG	Amplifies with MpS5R
MpS5R	GTATACAACGGCTTGATTGG	Anneals outside integration arms, amplifies with MpS5F

References

1. Wilkinson HW. 1978. Group B Streptococcal Infection in Humans. Annual Review of Microbiology 32:41-57.
2. Doran KS, Nizet V. 2004. Molecular pathogenesis of neonatal Group B Streptococcal infection: No longer in its infancy. Molecular Microbiology 54:23-31.
3. Hall J, Adams NH, Bartlett L, Seale AC, Lamagni T, Bianchi-Jassir F, Lawn JE, Baker CJ, Cutland C, Heath PT, Ip M, Le Doare K, Madhi SA, Rubens CE, Saha SK, Schrag S, Sobanjo-Ter Meulen A, Vekemans J, Gravett MG. 2017. Maternal Disease With Group B *Streptococcus* and Serotype Distribution Worldwide: Systematic Review and Meta-analyses. Clinical infectious diseases : an official publication of the Infectious Diseases Society of America 65:S112-S124.
4. Schuchat A. 1998. Epidemiology of Group B Streptococcal disease in the United States: shifting paradigms. Clinical Microbiology Reviews 11:497-513.
5. Edwards MS, Rench MA, Haffar AA, Murphy MA, Desmond MM, Baker CJ. 1985. Long-term sequelae of Group B Streptococcal meningitis in infants. The Journal of Pediatrics 106:717-22.
6. Kuypers JM, Heggen LM, Rubens CE. 1989. Molecular analysis of a region of the Group B *Streptococcus* chromosome involved in type III capsule expression. Infection and Immunity 57:3058-65.
7. Lancefield RC, McCarty M, Everly WN. 1975. Multiple mouse-protective antibodies directed against Group B Streptococci. Special reference to antibodies effective against protein antigens. The Journal of Experimental Medicine 142:165-79.
8. Wilkinson HW. 1977. Nontypable Group B Streptococci isolated from human sources. Journal of Clinical Microbiology 6:183-4.
9. Doran KS, Engelson EJ, Khosravi A, Maisey HC, Fedtke I, Equils O, Michelsen KS, Arditi M, Peschel A, Nizet V. 2005. Blood-brain barrier invasion by Group B *Streptococcus* depends upon proper cell-surface anchoring of lipoteichoic acid. The Journal of Clinical Investigation 115:2499-507.
10. Peschel A, Jack RW, Otto M, Collins LV, Staubitz P, Nicholson G, Kalbacher H, Nieuwenhuizen WF, Jung G, Tarkowski A, Van Kessel KPM, Van Strijp JAG. 2001. *Staphylococcus aureus* resistance to human defensins and evasion of neutrophil killing via the novel virulence factor MprF is based on modification of membrane lipids with L-lysine. Journal of Experimental Medicine 193:1067-1076.
11. Roy H, Ibba M. 2008. RNA-dependent lipid remodeling by bacterial multiple peptide resistance factors. Proceedings of the National Academy of Sciences of the United States of America 105:4667-4672.
12. Joyce LR, Guan Z, Palmer KL. 2019. Phosphatidylcholine Biosynthesis in Mitis Group Streptococci via Host Metabolite Scavenging. Journal of Bacteriology 201:e00495-19.
13. Adams HM, Joyce LR, Guan Z, Akins RL, Palmer KL. 2017. *Streptococcus mitis* and *S. oralis* Lack a Requirement for CdsA, the Enzyme Required for Synthesis of Major Membrane Phospholipids in Bacteria. Antimicrobial Agents and Chemotherapy 61:e02552-16.

14. Roy H. 2009. Tuning the properties of the bacterial membrane with aminoacylated phosphatidylglycerol. IUBMB Life 61:940-953.

15. Deng L, Spencer BL, Holmes JA, Mu R, Rego S, Weston TA, Hu Y, Sanches GF, Yoon S, Park N, Nagao PE, Jenkinson HF, Thornton JA, Seo KS, Nobbs AH, Doran KS. 2019. The Group B Streptococcal surface antigen I/II protein, BspC, interacts with host vimentin to promote adherence to brain endothelium and inflammation during the pathogenesis of meningitis. PLoS pathogens 15:e1007848.

16. Spencer BL, Deng L, Patras KA, Burcham ZM, Sanches GF, Nagao PE, Doran KS. 2019. Cas9 contributes to Group B Streptococcal colonization and disease. Frontiers in Microbiology 10:1-15.

17. Faralla C, Metruccio MM, De Chiara M, Mu R, Patras KA, Muzzi A, Grandi G, Margarit I, Doran KS, Janulczyk R. 2014. Analysis of two-component systems in group B Streptococcus shows that RgfAC and the novel FspSR modulate virulence and bacterial fitness. mBio 5:e00870-14.

18. Spencer BL, Chatterjee A, Duerkop BA, Baker CJ, Doran KS. 2021. Complete Genome Sequence of Neonatal Clinical Group B Streptococcal Isolate CJB111. Microbiology resource announcements 10.

19. Mu R, Cutting AS, Del Rosario Y, Villarino N, Stewart L, Weston TA, Patras KA, Doran KS. 2016. Identification of CiaR Regulated Genes That Promote Group B Streptococcal Virulence and Interaction with Brain Endothelial Cells. PLoS One 11:e0153891.

20. Van De Rijn I, Kessler RE. 1980. Growth characteristics of Group A Streptococci in a new chemically defined medium. Infection and Immunity 27:444-448.

21. Chang JC, LaSarre B, Jimenez JC, Aggarwal C, Federle MJ. 2011. Two Group A Streptococcal peptide pheromones act through opposing rgg regulators to control biofilm development. PLoS Pathogens 7:e1002190.

22. Gupta R, Shah P, Swiatlo E. 2009. Differential gene expression in *Streptococcus pneumoniae* in response to various iron sources. Microbial Pathogenesis 47:101-109.

23. Granok AB, Parsonage D, Ross RP, Caparon MG. 2000. The RofA binding site in *Streptococcus pyogenes* is utilized in multiple transcriptional pathways. Journal of bacteriology 182:1529-40.

24. Jeng A, Sakota V, Li Z, Datta V, Beall B, Nizet V. 2003. Molecular genetic analysis of a group A *Streptococcus* operon encoding serum opacity factor and a novel fibronectin-binding protein, SfbX. J Bacteriol 185:1208-17.

25. Hooven TA, Bonakdar M, Chamby AB, Ratner AJ. 2019. A Counterselectable Sucrose Sensitivity Marker Permits Efficient and Flexible Mutagenesis in *Streptococcus agalactiae*. Applied and Environmental Microbiology 85:1-13.

26. Holo H, Nes IF. 1989. High-frequency transformation, by electroporation, of *Lactococcus lactis* subsp. *cremoris* grown with glycine in osmotically stabilized media. Applied and Environmental Microbiology 55:3119-3123.

27. Framson PE, Nittayajarn A, Merry J, Youngman P, Rubens CE. 1997. New genetic techniques for Group B Streptococci: High-efficiency transformation, maintenance of temperature-sensitive pWV01 plasmids, and mutagenesis with Tn917. Applied and Environmental Microbiology 63:3539-3547.

28. Tan BK, Bogdanov M, Zhao J, Dowhan W, Raetz CRH, Guan Z. 2012. Discovery
of a cardiolipin synthase utilizing phosphatidylethanolamine and
phosphatidylglycerol as substrates. Proceedings of the National Academy of
Sciences 109:16504-16509.
29. Li C, Tan BK, Zhao J, Guan Z. 2016. In vivo and in vitro synthesis of
phosphatidylglycerol by an *Escherichia coli* cardiolipin synthase. Journal of
Biological Chemistry 291:25144-25153.
30. Kim BJ, Hancock BM, Bermudez A, Del Cid N, Reyes E, van Sorge NM, Lauth X,
Smurthwaite CA, Hilton BJ, Stotland A, Banerjee A, Buchanan J, Wolkowicz R,
Traver D, Doran KS. 2015. Bacterial induction of Snail1 contributes to blood-
brain barrier disruption. The Journal of Clinical Investigation 125:2473-83.
31. Banerjee A, Kim BJ, Carmona EM, Cutting AS, Gurney MA, Carlos C, Feuer R,
Prasadarao NV, Doran KS. 2011. Bacterial Pili exploit integrin machinery to
promote immune activation and efficient blood-brain barrier penetration. Nature
Communications 2:462.
32. Hooven TA, Randis TM, Daugherty SC, Narechania A, Planet PJ, Tettelin H,
Ratner AJ. 2014. Complete Genome Sequence of *Streptococcus agalactiae*
CNCTC 10/84, a Hypervirulent Sequence Type 26 Strain. Genome
Announcements 2:e01338-14.
33. Kilian M, Mikkelsen L, Henrichsen J. 1989. Taxonomic Study of Viridans
Streptococci: Description of *Streptococcus gordonii* sp. nov. and Emended
Descriptions of *Streptococcus sanguis* (White and Niven 1946), *Streptococcus*
oralis (Bridge and Sneath 1982), and *Streptococcus mitis* (Andrewes and Horder
1906). International Journal of Systematic Bacteriology 39:471-484.
34. Taylor RG, Walker DC, McInnes RR. 1993. *E. coli* host strains significantly affect
the quality of small scale plasmid DNA preparations used for sequencing. Nucleic
Acids Research 21:1677-1678.
35. Casadaban MJ, Cohen SN. 1980. Analysis of gene control signals by DNA fusion
and cloning in Escherichia coli. J Mol Biol 138:179-207.
36. Palmer KL, Carniol K, Manson JM, Heiman D, Shea T, Young S, Zeng Q, Gevers
D, Feldgarden M, Birren B, Gilmore MS. 2010. High-quality draft genome
sequences of 28 *Enterococcus* sp. isolates. Journal of Bacteriology 192:2469-70.