Temperature effects on individual-level parasitism translate into predictable effects on parasitism in populations

Devin Kirk[1,2], Mary I. O'Connor[2], Erin A. Mordecai[1]

AFFILIATIONS:

1. Department of Biology, Stanford University, Stanford, USA.

2. Department of Zoology, University of British Columbia, Vancouver, Canada.

ABSTRACT

Parasitism – the interaction between a parasite and its host – is expected to change in a warmer future, but the direction and magnitude of this change is uncertain. One challenge is understanding whether warming effects will be similar on individual hosts (e.g., parasite burden) compared to on population-level parasitism (e.g., prevalence, R_0). Examining thirteen empirical systems, we found a strong positive relationship between the thermal optima of individual- and population-level parasitism. We also found that parasitism thermal optima were close to host performance thermal optima in mosquito – parasite systems but not in non-mosquito – parasite systems. A simple mechanistic model showed how population-level parasitism thermal optima can be similar to individual-level parasitism thermal optima even under conditions where parasite transmission has a considerably higher thermal optimum. These results provide a key step towards finding general rules for how warming temperatures should affect parasitism in individuals, populations, and ecosystems.

INTRODUCTION

Organisms increasingly face temperatures warmer than their optima for physiological performance due to increasing and more variable temperatures caused by anthropogenic climate change. Changes in temperature can alter species interactions (Thomas & Blanford 2003; Kordas *et al.* 2011), and there is growing evidence that climate change has impacted host – parasite interactions and infectious disease dynamics around the world (Koelle *et al.* 2005; Rohr *et al.* 2011; Lafferty & Mordecai 2016; Claar & Wood 2020). Indeed, a series of major disease outbreaks in marine species have been linked to climate warming (e.g., seastar wasting disease, Harvell *et al.* 2019; coral die-offs, Bruno *et al.* 2007; and black abalone die-offs, Ben-Horin *et al.* 2013). These and other outbreaks have made predicting effects of temperature and changing climate on infectious disease dynamics an urgent priority (Altizer *et al.* 2013). Importantly, temperature affects biological systems across levels of organization, from individual physiology and infection burden to host population and disease dynamics (Fig. 1). This means that understanding the effects of temperature on both individuals and populations is critical for disease mitigation efforts as the world warms. Yet this cross-scale investigation of thermal disease ecology remains an open research gap.

Host–parasite systems involve multiple biological processes at more than one level of biological organization: individual host performance, individual parasite performance, host population dynamics, parasite population dynamics within a host, and parasite population dynamics among hosts within a population. Each of these biological processes and the traits that underlie them may be temperature dependent. One challenge for anticipating severe consequences of parasitism is to understand how temperature effects emerge at the host population level even when there can be a range of thermal responses among other biological

processes influencing the host and parasite (Fig. 1). For example, multiple host and parasite traits

and their respective temperature dependencies (typically described using *thermal performance*

curves, or TPCs) influence the effects of temperature on individual-level parasitism (Kirk *et al.*

2018). Moreover, individual-level parasitism outcomes, such as parasite burden (McCallum *et al.*

2017) and parasite-induced host mortality (Day 2001), can affect parasite transmission which

subsequently affects population-level parasitism, meaning that the thermal responses of

individual- and population-level parasitism are not independent.

In a simple case with temperature-independent transmission rates, we may expect the

thermal response of population-level parasitism to match the response of individual-level

parasitism. However, the temperature dependence of population-level parasitism is unlikely to be

so straightforward. Several potentially temperature-dependent traits can affect parasite

transmission, including demographic traits that influence host population dynamics and

behavioral traits of hosts, and these traits can therefore alter how individual-level parasitism

scales to the population level (Fig. 1). For example, activity rate of individuals can vary with

temperature (Casey 1976), thereby likely modifying the contact rate between susceptible and

infected individuals and the overall transmission rate. Additionally, the density of susceptible

hosts, often a key factor in parasite transmission, depends on demographic processes such as

birth and death rates, which can be strongly influenced by temperature (Savage *et al.* 2004;

Amarasekare & Savage 2012; Bernhardt *et al.* 2018). If the thermal responses of contact rates

and the density of susceptible hosts are distinct from the thermal response of individual-level

parasitism, we may also expect the thermal response of population-level disease to be distinct

from that at the individual level. Understanding the relationship between parasitism at these

levels is critical for understanding how disease dynamics, individual fitness, population

dynamics, and evolution will change with climate-driven shifts in parasitism regimes.

One framework for understanding temperature-driven changes in disease centers on the

thermal mismatch hypothesis (Cohen *et al.* 2017). Complementary to the approach we outline

above and in Figure 1, which emphasizes parasitism at the host individual and host population

levels, the thermal mismatch hypothesis relates the thermal response of parasitism to that of the

host. The thermal mismatch hypothesis predicts that parasitism is maximized at temperatures

away from the host's optimum—i.e., at cool temperatures for warm-adapted species and at warm

temperatures for cold-adapted species—at both the host individual and host population levels

(i.e., individual- and population-level parasitism TPCs should peak at temperatures offset from

the host optimum in what is called a thermal mismatch; Cohen *et al.* 2017, 2019a, b). Thermal

mismatches were first documented in amphibian – chytrid fungus (*Batracochytrium*

dendrobatidis, *Bd*) systems at both the individual and population levels (Cohen *et al.* 2017).

More recently, Cohen and colleagues (2020) analyzed >2000 host–parasite combinations at the

population level and found that hosts from cool and warm climates had increased disease

prevalence at warm and cool temperatures, respectively (i.e., there was evidence for population-

level thermal mismatches). However, outside of amphibian – *Bd* systems, it is unclear if thermal

mismatches at the population-level will generally correspond with thermal mismatches at the

individual level. How the optimal temperature for both individual- and population-level

parasitism compares to the optimal temperature for the host may mediate how climate change

will affect host – parasite interactions: decreased host performance at warmer temperatures could

either be compounded by increasing parasitism or mitigated by decreased parasitism.

Here, we re-analyze data from thirteen host – parasite systems to first test the hypothesis

that the thermal optimum of population-level parasitism (e.g., prevalence, basic reproduction

number R_0) is positively related to the thermal optimum of individual-level parasitism (e.g.,

parasite burden) across host – parasite systems. Next, we compare thermal optima of individual-

and population-level parasitism to the thermal optimum of host performance in the thirteen

systems to test the hypothesis that thermal mismatches at the individual level correspond with

thermal mismatches at the population level. Finally, we use a mechanistic model to illustrate how

even under conditions where parasite transmission has a thermal optimum much higher than

individual-level parasitism, the thermal optimum of population-level parasitism can still be

closer to the individual level than that of transmission. Our findings provide a first step towards

finding general rules for how warming temperatures should affect parasitism across biological

scale.

108
Figure 1. Illustration of a simplified host-parasite system that highlights temperature-
dependent processes occur and interact across levels of biological organization. Temperature
can influence parasite growth and host defenses against infection, which can ultimately lead to a
thermal response for individual-level parasitism (e.g., parasite burden; a). Similarly, temperature
can affect contact rates between hosts or hosts and parasites, the probability of infection after
contact, and host density, all of which can lead to variation in parasite transmission rate across
temperature (b). In some cases, the thermal response of individual-level parasitism (a) may also
partially influence the thermal response of parasite transmission (b), as higher parasite burden
can confer a higher probability of infection after contact or can alter contact rates between hosts.
Together, the thermal responses of individual-level parasitism (a) and parasite transmission (b)
will help determine the thermal response of population-level parasitism (e.g., parasite prevalence
or the basic reproduction number R_0; c).

133	**METHODS**

134	*Empirical analyses*

We tested the hypothesis that the thermal optimum of population-level parasitism is

positively related to the thermal optimum of individual-level parasitism across host – parasite

systems. Because it can be difficult to summarize an entire thermal performance curve (TPC)

with a single metric, we chose to focus on and compare the temperatures at which the curves are

maximized—the thermal optimum (T_{opt})—which is also more widely available for many host –

parasite systems than the full TPC. Though it does not represent the complete thermal response,

T_{opt} is a useful metric for comparison because it determines the range of temperatures at which

warming has a positive effect on parasitism (for $T < T_{opt}$) versus a negative effect (for $T > T_{opt}$).

Other metrics, such as thermal breadth and the critical thermal minimum and maximum, would

provide additional information about how climate change could affect host – parasite interactions

at the range margins. After testing for a relationship in the thermal optima of parasitism across

levels, we compared T_{opt} for individual- and population-level parasitism to T_{opt} of host

performance to determine if parasitism generally peaked at temperatures away from host thermal

optima as predicted by the thermal mismatch hypothesis (Cohen *et al.* 2017, 2019a, b), and if so,

whether these thermal mismatches at the individual level corresponded with thermal mismatches

at the population level.

We sought out systems for which measures of host performance (e.g., lifetime

reproduction), individual-level parasitism (e.g., parasite burden, parasite reproduction within the

host, time spent infected), and population-level parasitism (e.g., R_0, prevalence) were

documented across temperatures, allowing us to identify T_{opt} for each of these measures. While

not an exhaustive list, we identified thirteen systems that matched these requirements (Table 1):

four mosquito – virus systems (Mordecai *et al.* 2013; Shocket *et al.* 2018a; Tesla *et al.* 2018;

Mordecai *et al.* 2019; Shocket *et al.* 2020), four mosquito – malaria parasite systems (Villena *et*

al. 2020), two *Daphnia* – parasite systems (*D. magna* – *O. colligata*, Kirk *et al.* 2018, 2020; *D.*

dentifera – *M. bicuspidata*, Shocket *et al.* 2018), two amphibian – *B. dendrobatidis* (*Bd*, the

causative agent of chytridiomycosis) systems (Cohen *et al.* 2017), and one crab – rhizocephalan

barnacle parasite system (*E. depressus* – *L. panopaei*; Gehman *et al.* 2018). While the mosquito

systems are mainly studied with respect to human disease, here we leverage the rich data on their

thermal dependence to explore thermal scaling of the mosquito – parasite interaction, which in

turn affects transmission to humans. Using these systems, we tested the null hypothesis that there

is no relationship between the thermal optima of parasitism across scales by regressing

population-level parasitism T_{opt} against individual-level parasitism T_{opt} using the *lm* function in

the R package *stats* (R Core Team 2020).

168   Ideally, the metrics used for host performance, individual-level parasitism, and

population-level parasitism would be the same across systems; however, that is not the case

because different studies report different metrics. For example, the four mosquito–virus systems

use R_0 as the metric for population-level parasitism, while the four mosquito–malaria systems

use a slightly different metric, transmission suitability, which ignores the temperature-

independent human density and human recovery rate parameters. Table 1 lists the metrics used

for each system, with further details listed in the Supporting Information. We report a single host

T_{opt} value for eleven of the systems, but two host T_{opt} values (individual and population) for the

amphibian systems. This is because individual-level parasitism for cold-adapted and warm-

adapted amphibians was measured in the lab across one and two host species, respectively

(Cohen *et al.* 2017), but the thermal response of population-level parasitism was reported as *Bd*

prevalence in the field as a function of environmental temperature for 235 surveyed species,

where cold- and warm-adapted amphibians were categorized as those at locations where 50-year

mean temperature was <15°C or >20°C, respectively (Cohen *et al.* 2017). We therefore used

separate T_{opt} values for our individual-level host performance (measured as thermal preference in

the lab) and population-level host performance (measured as the mean climatic temperature

experienced in the field across surveyed species).

While we were generally constrained to use whichever metric of parasitism or host

performance was reported for a study, we had access to TPCs of several different metrics in the

mosquito – parasite systems. We chose to use infected days as our measure of individual-level

parasitism in these eight systems because it is a composite metric of thermal performance data on

parasite development rate within the mosquito, vector competence (the mosquito's ability to

acquire and transmit the parasite), and mosquito survival. However, we were interested in how

this choice affected our findings of how closely T_{opt} of individual-level parasitism matched T_{opt}

of population-level parasitism and T_{opt} of host performance. Generally, using vector competence

as a metric of individual-level parasitism gave similar results to our main metric of infected days,

but parasite development rate tended to exhibit higher T_{opt} values than the other two metrics (Fig.

S1). The Supporting Information contains further details on both our model and the methods

used to analyze the empirical systems.

202   TABLE 1. Thermal optima of parasitism and host performance for thirteen host – parasite systems.

Host – parasite system	Measure of host performance	Measure of individual-level parasitism	Measure of population-level parasitism	Host performance T_{opt} (°C)	Individual-level parasitism T_{opt} (°C)	Population-level parasitism T_{opt} (°C)	Reference
Culex pipiens – West Nile virus	Adult reproduction weighted by lifespan	Infected days	R_0	15.9	23.5	24.5	Shocket et al. 2020
Aedes aegypti - dengue virus	Adult reproduction weighted by lifespan	Infected days	R_0	27.6	29.3	29.1	Mordecai et al. 2017
Aedes aegypti - Zika virus	Adult reproduction weighted by lifespan	Infected days	R_0	27.7	28.9	28.9	Tesla et al. 2018
Culex annulirostris – Ross River virus	Adult reproduction weighted by lifespan	Infected days	R_0	25.6	24.0	26.4	Shocket et al. 2018a
Anopholes gambiae - *Plasmodium vivax*	Adult reproduction weighted by lifespan	Infected days	Transmission suitability	24.4	23.6	25.0	Villena et al. 2020
Anopholes gambiae - *Plasmodium falciparum*	Adult reproduction weighted by lifespan	Infected days	Transmission suitability	24.4	23.4	25.0	Villena et al. 2020
Anopholes stephensi - *Plasmodium vivax*	Adult reproduction weighted by lifespan	Infected days	Transmission suitability	24.0	23.0	24.6	Villena et al. 2020
Anopholes stephensi - *Plasmodium falciparum*	Adult reproduction weighted by lifespan	Infected days	Transmission suitability	24.0	22.9	24.8	Villena et al. 2020
Cold-adapted amphibians - *Batrachochytrium dendrobatidis*	Temperature preference for individuals; mean climatic temperature for populations †	Parasite growth rate on host	Prevalence	17.9 (10.5) †	26.0	20.5	Cohen et al. 2017
Warm-adapted amphibians - *Batrachochytrium dendrobatidis*	Temperature preference for individuals; mean climatic temperature for populations †	Parasite growth rate on host	Prevalence	23.5 (23.9) †	12.0	15.9	Cohen et al. 2017
Eurypanopeus depressus - *Loxothylacus panopaei*	Uninfected host survival	Parasite lifetime reproduction	R_0	18.3	15.9	14.7	Gehman et al. 2018
Daphnia dentifera - *Metschnikowia bicuspidata*	Uninfected intrinsic growth rate	Spore load	R_0	26.0*	20.0*	26.0	Shocket et al. 2018b
Daphnia magna – *Ordospora colligata*	Lifetime reproduction	Spore load	R_0	16.2*	11.8*	18.8	Kirk et al. 2018, 2020

* T_{opt} for these values was determined as the temperature at which the measure was maximized along a range of discrete
experimental temperatures, rather than the temperature maximizing the measure along a fitted continuous curve, as is the case for
each other value in the table. For discrete temperature range used, see Supporting Information.
† Since amphibian measures of individual-level and population-level parasitism were undertaken in different conditions and
across a range of populations and species (laboratory experiments and field observations, respectively; Cohen *et al.* 2017), we
used separate T_{opt} values for our individual-level host performance (measured as thermal preference in the lab) and population-
level host performance (measured as the mean climatic temperature experienced in the field across surveyed species). See
methods for further details.

Model

After testing for relationships between parasitism T_{opt} across levels of organization in thirteen empirical systems, we sought to determine whether differences in the thermal responses of transmission rate and individual-level parasitism could lead to large differences in thermal optima between individual-level and population-level parasitism. Based on our findings in the empirical systems, we hypothesized that even if the rates and quantities underlying transmission (contact rate and host density, respectively) had thermal optima that were much higher than individual-level parasitism, the resulting population-level parasitism T_{opt} would still tend closer to individual-level parasitism T_{opt} than to transmission T_{opt}. We explored this using a heuristic model that relates a measure of population-level parasitism, the basic reproduction number R_0, to host and parasite parameters, some of which are measured at the level of the individual host. We modeled different scenarios for temperature dependence of these parameters. We first described a single theoretical thermal performance curve for individual-level parasitism using a Brière function that is maximized at an intermediate temperature (i.e., T_{opt} = 20.6°C). The Brière function (Eq. 1) is an asymmetrical unimodal function that is often used to describe thermal performance in biological systems (Briere *et al.* 1999), where c is a positive rate constant, T_{min} and T_{max} are the minimum and maximum temperatures, respectively, T is temperature, and $B(T)$ is set to 0 when $T > T_{max}$ or $T < T_{min}$. For this example, we define host individual-level parasitism as parasite burden, or the number of parasites within or on the host:

$$B(T) = cT(T - T_{min})(T_{max} - T)^{1/2} \qquad \text{Eq. 1}$$

We next developed a simple trait-based model of population-level parasitism by modeling the basic reproduction number of the parasite (R_0; Eq. 2).

$$R_0 = \frac{\chi(T) \cdot \sigma(B(T)) \cdot S(T)}{\mu(T) + \alpha(B(T)) + \gamma}$$ Eq. 2

Here, parameters that are functions of individual-level parasitism ($B(T)$) are parasite-induced host mortality (α) and the probability of infection after contact (σ). Background mortality rate of hosts (μ), a function of temperature, and host recovery rate (γ) are both individual-level rates but are assumed here to not be affected by individual-level parasitism. Contact rate between susceptible and infected hosts (χ) and host density (S), both of which are temperature-sensitive but neither of which are assumed to vary with individual-level parasitism, affect parasite transmission (the numerator in Eq. 2). We made this distinction explicit because we wanted to highlight that certain traits can vary with the number of parasites within a host, thereby connecting the individual and population levels, while other traits will affect population-level parasitism independent from individual-level burden.

We assume linear relationships between individual-level parasite burden and both the probability of infection after contact (σ) (though non-linear relationships are also possible; McCallum *et al.* 2017), and parasite-induced mortality (α) (Anderson & May 1978), respectively. Because σ and α are proportional to individual-level parasite burden, both parameters are maximized at the same temperature as burden ($T_{opt} = 20.6°C$). We also set background mortality rate (μ) to share the same thermal optimum, meaning that μ is minimized at 20.6°C and then increases at colder and warmer temperatures as described by a quadratic function, used rather than an asymmetrical Brière because mortality often shows relatively symmetrical responses across temperature (van der Have 2002; Angilletta Jr. 2009). We assumed that recovery rate was temperature-independent, though that need not be the case in all systems.

To test whether different thermal responses of contact rate between infected and susceptible hosts (χ) and susceptible host density (S) would shift the thermal optima of

population-level parasitism away from the thermal optima of individual-level parasitism, we allowed χ and S to each be described by one of three functions: a Brière function that has identical temperature-dependence as individual-level parasitism (i.e., $T_{opt} = 20.6°C$), a Brière function that is shifted along the x-axis to peak at higher temperatures ($T_{opt} = 30°C$), and a temperature-independent function in which the parameter takes the same value (the average value of the Brière functions) at each temperature. We calculated R_0 for each of the nine combinations, resulting in six unique R_0 responses as some combinations are redundant (e.g., the scenario with contact rate peaking at 20.6°C and density peaking at 30°C gives the same R_0 response as the scenario with contact rate peaking at 30°C and density peaking at 20.6°C).

RESULTS

Empirical analyses

The thirteen empirical systems we analyzed demonstrated a range of thermal optima for individual-level parasitism (11.8°C – 29.3°C) and population-level parasitism (14.7°C – 29.1°C). Population-level parasitism T_{opt} was significantly positively associated with individual-level parasitism T_{opt} (Fig. 2; $p = 0.0005$, adjusted $R^2 = 0.65$, n = 13). Of the thirteen systems, nine exhibited thermal optima at individual and population levels that were within 2.5°C of each other. The largest difference across scales appeared in the *D. magna – O. colligata* system in which individual-level parasitism peaked at a temperature that is 7°C cooler than the peak for population-level parasitism. Nine of the thirteen systems exhibited higher thermal optima for population-level parasitism than individual-level parasitism, though in most cases this difference was small (Fig. 2; points above the dashed 1:1 line).

Figure 2. Empirical estimates of the optimal temperature (T_{opt}) for population-level parasitism versus T_{opt} of individual-level parasitism for thirteen host – parasite systems. We found a significant positive relationship between T_{opt} of population-level parasitism and T_{opt} of individual-level parasitism (p = 0.0005, adjusted R^2 = 0.65; solid line), and found that nine of the thirteen systems exhibited higher thermal optima for population-level parasitism than for individual-level parasitism (i.e., fall above the dashed 1:1 line). Circles represent mosquito – parasite systems and diamonds indicate non-mosquito – parasite systems.

Next, we compared how the thermal optima of parasitism at both levels compared to T_{opt} for host performance (Fig. 3). If a system was situated at the origin in Fig. 3, it would have individual-level parasitism, population-level parasitism, and host performance maximized at the same temperature (i.e., no thermal mismatches exist). Displacement from the origin represents parasitism peaking at temperatures away from where host performance peaks (i.e., thermal mismatches at one or both levels). We found that over half (7/13) of the systems were situated close to the origin (within Euclidian distance of 2.5°C, represented by the circle in Fig. 3).

However, all seven of these are mosquito – parasite systems, with the only mosquito system exhibiting strong thermal mismatches being *Cx. pipiens* – West Nile virus, in which both infected days and R_0 peak at temperatures nearly 10ºC higher than host performance (Fig. 3). The cold adapted amphibian – *B. dendrobatidis* system was also situated in the far upper-right quadrant (Fig. 3), showing strong thermal mismatches at both levels of parasitism, as shown in Cohen et al. 2017. Two systems showed cold-temperature mismatches at both levels: the crab *E. depressus* – *L. panopaei* and warm-adapted amphibian – *B. dendrobatidis* host – parasite systems. The zooplankton *D. dentifera* – *M. bicuspidata* system exhibited a thermal mismatch at cold temperatures at the individual level but no mismatch at the population level. Finally, the zooplankton *D. magna* – *O. colligata* system showed a distinct pattern of differing thermal mismatches across scales (Fig. 3): individual-level parasitism was maximized at a lower temperature than host performance while R_0 peaked at a higher temperature than host performance. None of the thirteen systems showed positive individual-level mismatches but negative population-level mismatches; therefore, the lower-right quadrant of Fig. 3 is empty. The six systems that experienced thermal mismatches generally supported our hypothesis that mismatches would occur at both scales and in the same direction, as more of these systems were situated in the upper-right or lower-left quadrants of Fig. 3 compared to the upper-left or lower-right quadrants.

Figure 3. Thermal matches and mismatches tended to be correlated across scales. The difference between T_{opt} of population-level parasitism and host performance (y-axis) is plotted against the difference between T_{opt} of individual-level parasitism and host performance (x-axis) for thirteen host – parasite systems. Systems situated at the origin had population-level parasitism, individual-level parasitism, and host performance all maximized at the same temperature (i.e., no thermal mismatches exist), while displacement from the origin represents parasitism peaking at temperatures away from where host performance peaks (i.e., thermal mismatches at individual or population levels). All seven of the system situated close to the origin (within Euclidian distance of 2.5°C, represented by the black circle) were mosquito – parasite systems (circles; in contrast to systems with non-mosquito hosts: diamonds).

Model

We explored different scenarios for how thermal responses of traits related to

transmission could shift the thermal optimum of population-level parasitism away from that of

individual-level parasitism. As expected, we found that when contact rate and host density share

the same thermal optimum as individual-level parasitism or are temperature-independent,

population-level parasitism is also maximized at this temperature (20.6°C; Fig. 4). Alternatively,

if either one of these traits has a thermal optimum that occurs at a higher temperature than

individual-level parasitism, this difference can pull the population-level parasitism thermal

optimum away from that of the individual level and towards the thermal optimum of the trait

(30.0°C; Fig. 4). However, even in the most extreme scenario we explored in which both contact

and host density peaked at 30.0°C (nearly 10°C warmer than individual-level parasitism T_{opt}), we

found that R_0 T_{opt} occurred at a temperature only 3.3°C warmer than individual-level parasitism

T_{opt} (Fig. 4). Intuitively, this is because population-level R_0 is constrained by individual-level

parasite burden, which is equal to 0 at and beyond its critical thermal maximum temperature of

25°C, therefore leading to $R_0 = 0$ at and above 25°C in all nine scenarios (Fig. 4c) regardless of

the thermal response of contact rate and host density.

In addition to small changes in the thermal optima, we found that the thermal response of

contact rate and host density affected the magnitude of population-level parasitism. For example,

when both contact rate and host density followed identical thermal responses to individual-level

parasitism, it led to a maximum R_0 value that was nearly double the second highest maximum R_0

value (Fig. 4c). The three scenarios in which both contact rate and host density did not share a

thermal optimum with individual-level parasitism and were either temperature-independent or

peaked at 30.0°C resulted in the three lowest maximum R_0 values (Fig. 4c).

In sum, the model confirms the intuition that if rates or quantities that are related to parasite transmission rate, such as contact rate and host density, are either temperature-independent or have a thermal response that peaks at the same temperature as individual-level parasitism, we should expect T_{opt} of population-level parasitism to match that of individual-level parasitism. By contrast, if these key rates and quantities show thermal responses distinct from individual-level parasitism, it can lead to differences in the thermal responses of parasitism across biological scale, though we may still expect the thermal response of individual-level parasitism to constrain the population-level response from exhibiting a T_{opt} that is considerably different from the T_{opt} of individual-level parasitism.

Figure 4. Individual parasite burden and transmission-related traits combine to determine the temperature-dependence of population-level R_0. Individual-level parasitism (a) interacts with traits and quantities related to transmission such as contact rate and host density (b) to determine population-level parasitism (c). We explored how one specific thermal response of individual-level parasite burden (a) can interact with different combinations of responses for contact rate and host density that were either temperature-independent (yellow line, b), temperature-dependent and peaking at the identical temperature as burden (blue line, b), or temperature-dependent and peaking at a higher temperature than burden (red line, b). These combinations generated six unique thermal responses for R_0 (c). The scenario with the warmest T_{opt} for R_0 (red line, c) exhibited a thermal optimum that was only 3.3°C warmer than individual-level parasitism T_{opt} (a).

DISCUSSION

Temperature impacts parasitism at the individual host level by influencing parasite growth and host defenses against infection, ultimately leading to differential levels of parasite burden across temperature. This thermal response of individual-level parasitism is intrinsically tied to the thermal response of parasitism at the population level (Fig. 1, 4). In our examination of thirteen empirical systems, we found a significant positive relationship between T_{opt} for parasitism at the two levels (Fig. 2). Additionally, while individual- and population-level parasitism both peaked at temperatures away from the host optimum in some systems, supporting the thermal mismatch hypothesis, this was not the case in seven of the eight mosquito – parasite systems (Fig. 3). Generally, our results suggest that information on the temperature-dependence, and specifically the thermal optima, of a host – parasite system at either the individual- or population-level should provide a useful—though not quantitatively exact—baseline for predicting temperature-dependence at the other level in a variety of host – parasite systems.

We found a significant positive relationship between T_{opt} of population-level parasitism and T_{opt} of individual-level parasitism and found that in the majority of systems, population-level parasitism tended to peak at warmer temperatures than individual-level parasitism (Fig. 2). The greatest differences in T_{opt} were observed in the zooplankton *D. magna – O. colligata* system, cold-adapted amphibians – *B. dendrobatidis* system, and the zooplankton *D. dentifera – M. bicuspidata* system (Fig. 2, Table 1). It is difficult to definitively parse out the causes of T_{opt} differences in the two amphibian systems since individual-level *Bd* parasitism was measured on either one or two frog species in the lab while population-level prevalence by the same parasite was synthesized across many field studies encompassing 235 host species (Cohen *et al.* 2017). However, the *D. magna – O. colligata* system can provide an illuminating example. Contact rate

in this host – microsporidian parasite system is maximized at 30.1°C (Kirk *et al.* 2019). This is

nearly 20°C warmer than T_{opt} for individual-level parasitism (11.8°C) and is thus one of the key

factors that pulls the thermal optimum of R_0 away from individual-level parasitism T_{opt} to a

warmer temperature (18.8°C). Moreover, in the mosquito – parasite systems, which tended to

have high population-level parasitism T_{opt} (Fig. 2), biting rate—a driver of host contact—usually

has one of the highest optimal temperatures of all measured traits (Mordecai *et al.* 2013, 2017,

2019; Shocket *et al.* 2018a, 2020; Villena *et al.* 2020). These empirical systems are in line with

our model results that show that transmission-related traits with high thermal optima can

somewhat shift thermal optima of R_0 away from that of individual-level parasitism, though not to

such an extent that population-level parasitism will peak at the same temperature as the

transmission-related traits (Fig. 4). However, these few systems do not exhaustively represent all

host – parasite systems. More empirical work measuring thermal performance of traits related to

parasite transmission in other types of host – parasite systems, such as those with helminth or

bacterial parasites, can help to broaden our perspective on scaling of parasitism from individuals

to populations.

The thermal mismatch hypothesis (Cohen *et al.* 2017, 2019a, b, 2020) has been proposed

as an approach for understanding the effects of climate change on host – parasite systems,

predicting that T_{opt} for parasitism should occur at temperatures away from host T_{opt} (i.e., a

thermal mismatch) due to larger organisms (hosts) generally having narrower thermal breadth

than smaller organisms (parasites; Rohr *et al.* 2018). While thermal mismatches have been

shown to occur widely at the population level across systems (Cohen *et al.* 2020), we tested the

hypothesis that thermal mismatches at the individual level will correspond with similar thermal

mismatches at the population level. Of the five non-mosquito systems we explored—including

the two amphibian – *Bd* systems investigated previously (Cohen et al. 2017)—parasitism tended to peak at temperatures away from the host's optimum in each, meaning these systems experience thermal mismatches at one or both levels. If our hypothesis was to be supported, we would expect most systems experiencing thermal mismatches to fall in the upper-right or lower-left quadrants of Fig. 3, rather than the upper-left or lower-right. We found that this was generally the case, with two systems experiencing thermal mismatches at warmer temperatures at both levels (Fig. 3; upper-right) and two systems experiencing thermal mismatches at cooler temperatures at both levels (Fig. 3; lower-left). The major exception was the *D. magna – O. colligata* system, in which host T_{opt} occurs at a temperature intermediate to the relatively cool T_{opt} for individual-level parasitism and relatively warm T_{opt} population-level parasitism (Fig. 3; upper-left). The other zooplankton host system we explored, *D. dentifera – M. bicuspidata,* exhibited a thermal mismatch at cooler temperatures at the individual level but no mismatch at the population level.

Evidence for thermal mismatches occurring at either level was weak in the mosquito – parasite systems. Seven of eight mosquito – parasite systems were situated close to the origin (Fig. 3), suggesting that parasitism and host performance are maximized at close to the same temperatures and that no thermal mismatches are occurring at either level. The exception, *Cx. pipiens* – West Nile virus, in which parasitism at both levels peaks at temperatures much higher than host performance because *Cx. pipiens* adult lifespan is much longer at low temperatures (Shocket *et al.* 2020), is a system with a predominantly temperate distribution in North America, compared to more the sub-tropical and tropical distributions of malaria, dengue, Zika, Ross River virus, and their respective mosquito vectors. Notably, when testing for thermal mismatches at the population level, Cohen *et al.* (2020) also found that thermal mismatch effects were strongest in

systems without vectors (or intermediate hosts). Why we observed this pattern in mosquito systems will require further investigation in different vector-borne systems across biological scale, including those systems with non-mosquito vectors such as ticks or sandflies.

We propose that situating host – parasite systems along both individual- and population-level axes as in Fig. 3 can provide a diagnostic framework for looking for thermal mismatches across biological scales. While our results do not conclusively address how general thermal mismatches at both levels may be, our finding that individual- and population-level T_{opt} are strongly related (Fig. 2) provides a general rule of thumb that, if a host – parasite system does experience thermal mismatches, they are likely to occur at both scales and in the same direction (though note the exception of *D. magna – O. colligata*; Fig. 3). This has major implications for our understanding of the effects of climate change on host-parasite systems, because if T_{opt} for both individual- and population-level parasitism is at a higher temperature than T_{opt} for host performance, warming may not only decrease host performance itself, but also lead hosts to experience increasing effects of parasitism (Fig. 3; upper-right quadrant). Alternatively, if T_{opt} for parasitism is at temperatures cooler than host performance optima, decreased host performance at warmer temperatures may be partially offset by decreased effects of parasitism (Fig. 3; lower-left quadrant). This is in line with recent projections that climate change may lead to hosts from cooler and warmer climates experiencing increased and decreased parasitism, respectively (Cohen *et al.* 2020).

To better understand how temperature affects parasitism and to make better predictions for how climate change will affect host – parasite systems, we highlight two important research directions. First, when possible, studies that use experiments to investigate effects of temperature at the individual level can also be used to parameterize simple models of population-level

parasitism (a 'bottom-up' approach). Second, field studies investigating population-level parasitism under different climate or weather conditions, often undertaken by measuring the proportion of individuals infected, should also aim to record measures of individual-level parasitism (a 'top-down' approach). The best direct measure would be to record the parasite burden or an analogous metric measuring the parasite on or within the host, but in cases where this is not possible, indirect measures such as host condition may still be informative.

Overall, we found a strong relationship between thermal optima of parasitism across scales in the thirteen systems we analyzed, and our results suggest that in many host – parasite systems, individual-level outcomes may beget similar outcomes across temperature at the population level. Many studies have investigated the effects of temperature on parasitism (reviewed, for example, in Marcogliese 2008; Lafferty 2009; Rohr *et al.* 2011; Altizer *et al.* 2013; Lafferty & Mordecai 2016; Claar & Wood 2020), though often at only one of either the individual or population levels. Our findings build upon these studies by emphasizing that how temperature affects individual-level parasitism can scale up to affect parasite transmission, prevalence, and the potential for epidemics in host populations. While this is an important first step toward predicting the effects of climate change on systems by allowing researchers to primarily focus on parasitism at one level, future empirical work that investigates the thermal dependence of a wider range of host – parasite systems across scales, including in systems with different types of hosts (e.g., plants, non-amphibian vertebrates) and other types of parasites (e.g., helminths, bacterial parasites), will be necessary for investigating the generality of this result. More generally, tying together more theory and empirical results to provide general predictions for how climate change will affect parasitism from individuals to ecosystems in a

diverse range of hosts and parasites remains an urgent priority as accelerating climate change

makes likely catastrophic temperature increases of > 2°C increasingly likely by 2050.

ACKNOWLEDGEMENTS

We thank members of the Mordecai and O'Connor labs for valuable feedback on the manuscript.

EAM and DK were funded by NIH (R35GM133439); EAM was funded by NSF (DEB-1518681

and DEB-2011147, also supported by the Fogarty International Center), a King Center for

Global Development seed grant, and a Terman Award. Figure 1 was created using

BioRender.com.

REFERENCES

Altizer, S., Ostfeld, R.S., Johnson, P.T.J., Kutz, S. & Harvell, C.D. (2013). Climate Change and Infectious Diseases: From Evidence to a Predictive Framework. *Science*, 341, 514–519.

Amarasekare, P. & Savage, V. (2012). A Framework for Elucidating the Temperature Dependence of Fitness. *The American Naturalist*, 179, 178–191.

Anderson, R.M. & May, R.M. (1978). Regulation and Stability of Host-Parasite Population Interactions: I. Regulatory Processes. *The Journal of Animal Ecology*, 47, 219.

Angilletta Jr., M.J. (2009). Thermal Adaptation: A Theoretical and Empirical Synthesis. In: *Thermal adaptation: a theoretical and empirical synthesis*. Oxford University Press, pp. 88–125.

Ben-Horin, T., Lenihan, H.S. & Lafferty, K.D. (2013). Variable intertidal temperature explains why disease endangers black abalone. *Ecology*, 94, 161–168.

Bernhardt, J.R., Sunday, J.M. & O'Connor, M.I. (2018). Metabolic Theory and the Temperature-Size Rule Explain the Temperature Dependence of Population Carrying Capacity. *The American Naturalist*, 192, 687–697.

Briere, J.-F., Pracros, P., Le Roux, A.-Y. & Pierre, J.-S. (1999). A Novel Rate Model of Temperature-Dependent Development for Arthropods. *Environ Entomol*, 28, 22–29.

Bruno, J.F., Selig, E.R., Casey, K.S., Page, C.A., Willis, B.L., Harvell, C.D., *et al.* (2007). Thermal Stress and Coral Cover as Drivers of Coral Disease Outbreaks. *PLoS Biol*, 5, e124.

Casey, T.M. (1976). Activity Patterns, Body Temperature and Thermal Ecology in Two Desert Caterpillars (Lepidoptera: Sphingidae). *Ecology*, 57, 485–497.

Claar, D.C. & Wood, C.L. (2020). Pulse Heat Stress and Parasitism in a Warming World. *Trends in Ecology & Evolution*, 35, 704–715.

Cohen, J.M., Civitello, D.J., Venesky, M.D., McMahon, T.A. & Rohr, J.R. (2019a). An interaction between climate change and infectious disease drove widespread amphibian declines. *Glob Change Biol*, 25, 927–937.

Cohen, J.M., McMahon, T.A., Ramsay, C., Roznik, E.A., Sauer, E.L., Bessler, S., *et al.* (2019b). Impacts of thermal mismatches on chytrid fungus *Batrachochytrium dendrobatidis* prevalence are moderated by life stage, body size, elevation and latitude. *Ecol Lett*, 22, 817–825.

Cohen, J.M., Sauer, E.L., Santiago, O., Spencer, S. & Rohr, J.R. (2020). Divergent impacts of warming weather on wildlife disease risk across climates. *Science*, 370, eabb1702.

Cohen, J.M., Venesky, M.D., Sauer, E.L., Civitello, D.J., McMahon, T.A., Roznik, E.A., *et al.* (2017). The thermal mismatch hypothesis explains host susceptibility to an emerging infectious disease. *Ecol Lett*, 20, 184–193.

Day, T. (2001). Parasite transmission modes and the evolution of virulence. *Evolution*, 55, 2389–2400.

Gehman, A.-L.M., Hall, R.J. & Byers, J.E. (2018). Host and parasite thermal ecology jointly determine the effect of climate warming on epidemic dynamics. *Proc Natl Acad Sci USA*, 115, 744–749.

van der Have, T.M. (2002). A proximate model for thermal tolerance in ectotherms. *Oikos*, 98, 141–155.

Kirk, D., Jones, N., Peacock, S., Phillips, J., Molnár, P.K., Krkošek, M., *et al.* (2018). Empirical evidence that metabolic theory describes the temperature dependency of within-host parasite dynamics. *PLoS Biol*, 16, e2004608.

Kirk, D., Luijckx, P., Jones, N., Krichel, L., Pencer, C., Molnár, P., *et al.* (2020). Experimental evidence of warming-induced disease emergence and its prediction by a trait-based mechanistic model. *Proc. R. Soc. B.*, 287, 20201526.

Kirk, D., Luijckx, P., Stanic, A. & Krkošek, M. (2019). Predicting the Thermal and Allometric Dependencies of Disease Transmission via the Metabolic Theory of Ecology. *The American Naturalist*, 193, 661–676.

Koelle, K., Pascual, M. & Yunus, M. (2005). Pathogen adaptation to seasonal forcing and climate change. *Proc. R. Soc. B*, 272, 971–977.

Kordas, R.L., Harley, C.D.G. & O'Connor, M.I. (2011). Community ecology in a warming world: The influence of temperature on interspecific interactions in marine systems. *Journal of Experimental Marine Biology and Ecology*, 400, 218–226.

Lafferty, K.D. (2009). The ecology of climate change and infectious diseases. *Ecology*, 90, 888–900.

Lafferty, K.D. & Mordecai, E.A. (2016). The rise and fall of infectious disease in a warmer world. *F1000Res*, 5, 2040.

Marcogliese, D.J. (2008). The impact of climate change on the parasites and infectious diseases of aquatic animals. *Rev. scii. tech. Off. init. Epiz.*, 27, 467–484.

McCallum, H., Fenton, A., Hudson, P.J., Lee, B., Levick, B., Norman, R., *et al.* (2017). Breaking beta: deconstructing the parasite transmission function. *Phil. Trans. R. Soc. B*, 372, 20160084.

Mordecai, E.A., Caldwell, J.M., Grossman, M.K., Lippi, C.A., Johnson, L.R., Neira, M., *et al.* (2019). Thermal biology of mosquito-borne disease. *Ecol Lett*, 22, 1690–1708.

Mordecai, E.A., Cohen, J.M., Evans, M.V., Gudapati, P., Johnson, L.R., Lippi, C.A., *et al.* (2017). Detecting the impact of temperature on transmission of Zika, dengue, and chikungunya using mechanistic models. *PLoS Negl Trop Dis*, 11, e0005568.

Mordecai, E.A., Paaijmans, K.P., Johnson, L.R., Balzer, C., Ben-Horin, T., de Moor, E., *et al.* (2013). Optimal temperature for malaria transmission is dramatically lower than previously predicted. *Ecol Lett*, 16, 22–30.

R Core Team. 2020. R: a language and environment for statistical computing. https://www.R-project.org.

Rohr, J.R., Civitello, D.J., Cohen, J.M., Roznik, E.A., Sinervo, B. & Dell, A.I. (2018). The complex drivers of thermal acclimation and breadth in ectotherms. *Ecol Lett*, 21, 1425–1439.

Rohr, J.R., Dobson, A.P., Johnson, P.T.J., Kilpatrick, A.M., Paull, S.H., Raffel, T.R., *et al.* (2011). Frontiers in climate change–disease research. *Trends in Ecology & Evolution*, 26, 270–277.

Savage, V.M., Gillooly, J.F., Brown, J.H., West, G.B. & Charnov, E.L. (2004). Effects of Body Size and Temperature on Population Growth. *The American Naturalist*, 163, 429–441.

Shocket, M.S., Ryan, S.J. & Mordecai, E.A. (2018a). Temperature explains broad patterns of Ross River virus transmission. *eLife*, 7, e37762.

Shocket, M.S., Strauss, A.T., Hite, J.L., Šljivar, M., Civitello, D.J., Duffy, M.A., *et al.* (2018b). Temperature Drives Epidemics in a Zooplankton-Fungus Disease System: A Trait-Driven Approach Points to Transmission via Host Foraging. *The American Naturalist*, 191, 435–451.

Shocket, M.S., Verwillow, A.B., Numazu, M.G., Slamani, H., Cohen, J.M., El Moustaid, F., *et al.* (2020). Transmission of West Nile and five other temperate mosquito-borne viruses peaks at temperatures between 23°C and 26°C. *eLife*, 9, e58511.

Tesla, B., Demakovsky, L.R., Mordecai, E.A., Ryan, S.J., Bonds, M.H., Ngonghala, C.N., *et al.* (2018). Temperature drives Zika virus transmission: evidence from empirical and mathematical models. *Proc. R. Soc. B*, 285, 20180795.

Thomas, M.B. & Blanford, S. (2003). Thermal biology in insect-parasite interactions. *Trends in Ecology & Evolution*, 18, 344–350.

Villena, O.C., Ryan, S.J., Murdock, C.C. & Johnson, L.R. (2020). Temperature impacts the transmission of malaria parasites by Anopheles gambiae and Anopheles stephensi mosquitoes. *bioRxiv*, doi: https://doi.org/10.1101/2020.07.08.194472.