1 **TITLE:** Temperature effects on individual-level parasitism translate into predictable effects on
2 parasitism in populations
3
4
5
ABSTRACT

Parasitism is expected to change in a warmer future, but the direction and magnitude of this

change is uncertain. One challenge is understanding whether warming effects will be similar on

parasitism within or on individual hosts (e.g., parasite load) compared to on population-level

parasitism (e.g., prevalence, R_0). We adapted a simple temperature-dependent model and

simulated several scenarios for individual- and population-level parasitism. Our model found that

small differences in the underlying biology of host–parasite systems can substantially alter the

expected relationship between the thermal optima of parasitism across levels of organization. In

thirteen empirical host–parasite systems, we found a strong positive correlation between the

thermal optima of individual- and population-level parasitism, suggesting that the effects of

warming on parasitism may often be in the same direction across levels. We also found that

parasitism thermal optima were close to host performance thermal optima in mosquito–parasite

systems but not in non-mosquito–parasite systems. Generally, our results suggest that

information on the temperature-dependence, and specifically the thermal optima, of a host–

parasite system at either the individual- or population-level should provide a useful—though not

quantitatively exact—baseline for predicting temperature-dependence at the other level in a

variety of host–parasite systems.

INTRODUCTION

Climate change is causing organisms to increasingly face temperatures warmer than their

optima for physiological performance. These changes in temperature can alter species

interactions (Thomas & Blanford 2003; Kordas *et al.* 2011), and there is now growing evidence

that climate change has impacted interactions between individual hosts and their parasites as well

as parasite dynamics and outbreaks in host populations (Koelle *et al.* 2005; Bruno *et al.* 2007;

Rohr *et al.* 2011; Ben-Horin *et al.* 2013; Lafferty & Mordecai 2016; Harvell *et al.* 2019; Claar &

Wood 2020). Predicting the effects of temperature and changing climate on infectious disease is

an urgent priority (Altizer *et al.* 2013), yet research tends to focus on the level of either an

individual host's biology or of host populations, but rarely both.

Understanding how individual-level *per capita* interactions in host–parasite systems scale

to the population level is key for mitigating disease (Fenton 2008). Indeed, the many biological

processes that occur in host–parasite systems occur across more than one level of biological

organization, and bridging these levels is an important research area (Handel & Rohani 2015).

These processes include those that affect individual host performance, individual parasite

performance, host population dynamics, parasite population dynamics within a host, and parasite

population dynamics among hosts within a population. Crucially, each of these processes and the

traits that underlie them may be temperature dependent. However, whether we should expect the

effects of temperature on individual-level parasitism (e.g., on parasite load or on within-host

parasite population growth rate) to match the effects of temperature on population-level

parasitism (e.g., on parasite prevalence or on the basic reproduction number R_0) remains an open

research gap, as does whether small differences in the underlying host–parasite interaction or

thermal biology should affect how thermal effects scale between levels. Understanding the

effects of temperature on both individuals and populations is critical for disease mitigation

efforts as the world warms because if the effects are similar across levels of organization, we can

leverage observations of the thermal dependence of parasitism at one level to inform predictions

in more complex systems. However, if effects differ across levels, using observations on the

thermal dependence at one level to extrapolate to the other could provide erroneous predictions

for how warming will affect disease in a system.

Obtaining, summarizing, and comparing entire thermal responses can be difficult and

sometimes unfeasible. Fortunately, most thermal response curves have a characteristic non-linear

shape with a single thermal optimum at which performance is maximized (T_{opt}) (Huey &

Stevenson 1979; Huey & Kingsolver 1989; Angilletta 2006). In this study, we therefore focus on

and compare thermal optima, as this data is more widely available for many host–parasite

systems than full thermal response curves. Though it does not represent the complete thermal

response, T_{opt} is a useful metric for comparison both across levels of organization and among

taxa because it determines the range of temperatures at which warming has a positive effect on

parasitism (for $T < T_{opt}$) versus a negative effect (for $T > T_{opt}$). Other metrics, such as thermal

breadth and the critical thermal minimum and maximum, would provide additional information

about how climate change could affect host–parasite interactions at the margins of thermal

tolerance.

The thermal optima of parasitism could either match or differ across biological levels of

organization. First, we may expect the optimal temperature for parasitism at the individual level

to reflect the optimal temperature of parasitism in populations because processes across the two

levels can be linked (Ewald 1983; Frank 1996; Mideo *et al.* 2011; Handel & Rohani 2015). For

example, if the rate-limiting or rate-determining process for population-level parasitism is

directly related to individual-level parasitism, the effects of temperature on individual-level

parasitism may propagate directly to the population level. From this perspective, the thermal

optima of parasitism at the two levels are unlikely to be independent in many systems. This

could occur if the number of parasites within a host (i.e., individual-level parasitism) is the rate-

determining process affecting the rate of parasite transmission (Ben-Ami *et al.* 2008; McCallum

et al. 2017) or the rate at which hosts are killed by the parasite (Day 2001), both of which

subsequently affect population-level parasitism metrics such as R_0 or prevalence (Anderson &

May 1979).

On the other hand, temperature may not have the same effects across levels even in cases

where population-level parasitism is dependent on the dynamics of individual-level parasitism.

Indeed, when considering the effects of anti-parasite treatments instead of temperature, models

and field experiments have shown that the effects of treatments on individual hosts do not

necessarily lead to equivalent effects in host populations (Fenton 2013; Pedersen & Antonovics

2013). The process of parasite transmission among hosts can potentially decouple population

dynamics of hosts and parasites from dynamics within host individuals, particularly if the traits

that shape the transmission process are temperature dependent. For example, individual activity

rate can vary with temperature (Casey 1976), thereby modifying the contact rate between

susceptible and infected individuals and the overall transmission rate. If the thermal optima of

traits that affect transmission are much warmer or much cooler than that of individual-level

parasitism, we may expect the thermal optimum of population-level parasitism to be pulled away

from the individual-level optimum towards that of these traits.

Beyond exploring correlations between the thermal optima of parasitism across levels,

considering how the optima for parasitism are related to the optima for their uninfected potential

hosts can provide a more holistic view of how warming should affect a host–parasite system.

One potential framework for doing this is the thermal mismatch hypothesis (Cohen *et al.* 2017).

This hypothesis predicts that parasitism is maximized at temperatures away from the host's

optimum—i.e., at cool temperatures for warm-adapted species and at warm temperatures for

cold-adapted species—at both the host individual and host population levels (i.e., individual- and

population-level parasitism thermal responses should peak at temperatures offset from the host

optimum in what is called a thermal mismatch)(Cohen *et al.* 2017, 2019a, b). Thermal

mismatches were first documented in amphibian–chytrid fungus (*Batracochytrium*

dendrobatidis, *Bd*) systems at both the individual and population levels (Cohen *et al.* 2017).

More recently, Cohen et al. (2020) analyzed >2000 host–parasite combinations at the population

level and found evidence for population-level thermal mismatches: hosts from cool climates had

increased disease prevalence at warm temperatures and vice versa. However, outside of

amphibian–*Bd* systems, it is unclear if systems with thermal mismatches at the population level

will also exhibit thermal mismatches at the individual level. How the optimal temperatures for

individual- and population-level parasitism compare to the optimal temperature for the host may

mediate how climate change will affect host–parasite interactions: decreased host performance at

warmer temperatures could either be compounded by increasing parasitism or mitigated by

decreased parasitism.

Here, we first develop simple temperature-dependent models for individual-level

parasitism and population-level parasitism to examine the temperature dependence of parasitism

across levels. We then simulate several different scenarios—in which thermal responses of

population-level processes are or are not related to the thermal responses of individual-level

parasitism—to investigate how changes in the underlying processes and thermal biology of a

host–parasite system can alter the expected relationship between parasitism T_{opt} across levels.

Next, after using our model to explore possible relationships between parasitism T_{opt} across

levels, we sought to describe the observed relationship in nature. We identify thirteen systems

that match our data requirements and re-analyze the data to investigate if the thermal optimum of

population-level parasitism is positively related to the thermal optimum of individual-level

parasitism across empirical host–parasite systems. Finally, we compare thermal optima of

individual- and population-level parasitism to the thermal optimum of host performance in the

thirteen systems to test if the thermal mismatch hypothesis holds across levels in these systems.

Our findings provide a first step towards finding general rules for how warming temperatures

will affect parasitism across biological levels of organization.

128
129
130
131   **METHODS**

132   *Model*

To formalize our intuitive arguments for why we may or may not expect the thermal

optimum of parasitism in individual hosts to occur at the same temperature as the thermal

optimum for parasitism in host populations, we adapted a simple temperature-dependent model

for disease spread in a population that can be related to the thermal response for parasitism in an

individual host. Here, we define host individual-level parasitism as parasite load, or the number

of parasites within or on the host.

We first described the thermal response of individual-level parasitism using a Brière

function. The Brière function (Eq. 1) is an asymmetrical unimodal function that is often used to

describe thermal performance in biological systems (Briere *et al.* 1999), where *c* is a positive rate

constant, T_{min} and T_{max} are the minimum and maximum temperatures, respectively, T is

temperature, and $B(T)$ is set to 0 when $T > T_{max}$ or $T < T_{min}$.

$$B(T) = cT(T - T_{min})(T_{max} - T)^{1/2} \qquad \text{Eq. 1}$$

Next, we developed a simple trait-based model of population-level parasitism by

modeling the basic reproduction number of the parasite (R_0; Eq. 2):

$$R_0 = \frac{\chi(T) \cdot \sigma(T) \cdot S(T)}{\mu(T) + \alpha(T) + \gamma(T)} \qquad \text{Eq. 2}$$

Here, χ is the contact rate between susceptible and infected hosts, σ is the probability of

infection after contact, S is the density of susceptible hosts, μ is the background mortality rate, α

is the parasite-induced mortality rate, and γ is the host recovery rate. We modeled all six

parameters as temperature-dependent, and for all parameters except $\mu(T)$ we assumed thermal

responses described by the Brière function (the same functional form used to describe the

thermal response of individual-level parasitism; Eq. 1). We modeled the thermal response of

background mortality rate using a concave-up quadratic function where mortality is minimized at

the thermal optimum and increases at cooler and warmer temperatures, as mortality often shows

relatively symmetrical responses across temperature (van der Have 2002; Angilletta 2009).

We simulated the model by randomly drawing different thermal response function

parameters {c, T_{min}, T_{max}} for each temperature-dependent trait (individual-level parasitism and

parameters in Eq. 2). For each of 1000 simulations, T_{min} was drawn from a uniform distribution

between 0-15, T_{max} was equal to T_{min} plus a value drawn from a uniform distribution between 10-

20, and the rate constant c was drawn from a uniform distribution between 0.5-1.3. Similarly, for

the concave-up quadratic function that describes background mortality rate, in each of the 1000

simulations we drew parameter a from a uniform distribution between 0.0008-0.0009, b from a

uniform distribution between 0.02-0.03, and c from a uniform distribution between 0.1-1.

Thermal responses were scaled to realistic magnitudes for each temperature-dependent trait (e.g.,

all thermal responses for the contact rate parameter were scaled by a factor of 0.001), and we set

background mortality rate to a minimum of 0.005 (see Supporting Information for more detail).

To explore if changes in the underlying biology of the host–parasite systems can alter the

expected relationship between parasitism T_{opt} across levels, we modeled four main scenarios: (1)

all temperature-dependent traits were independent; (2) parasite-induced mortality rate was

proportional to parasite load (Anderson & May 1978); (3) per-contact infection probability was

proportional to parasite load (McCallum *et al.* 2017); and (4) both parasite-induced mortality and

per-contact infection probability were proportional to parasite load. We also explored seven

additional model scenarios to determine the sensitivity of our results to our assumptions

regarding both thermal response shape and the structure of the mechanistic model for population

disease spread. These scenarios included changes to how thermal response function parameters

were drawn, the breadth of the thermal response functions, the minimum parasite load across the

temperature range, and the model used to describe population-level disease spread (see

Supporting Information for more details). While the model scenarios explored in the main text

and the Supporting Information represent only a small subset of the possible relationships

between the thermal dependence of parasitism across levels, our aim was to investigate whether

small changes to the underlying biology can result in qualitatively different patterns in the

relationship of thermal optima in parasitism across levels, rather than to exhaustively explore

potential outcomes.

188	***Empirical systems***
189	After using our model to investigate potential relationships for the thermal optima of
190	parasitism across levels, we sought to describe the observed relationship in empirical systems.
191	We searched for systems for which measures of host performance (e.g., lifetime reproduction),
192	individual-level parasitism (e.g., parasite load, parasite reproduction within the host, time spent
193	infected), and population-level parasitism (e.g., R_0, prevalence) were documented across
194	temperatures, allowing us to identify T_{opt} for each of the measures. While not an exhaustive list,
195	we identified thirteen systems that matched these requirements (Table 1): four mosquito–virus
196	systems (Mordecai *et al.* 2013; Shocket *et al.* 2018a; Tesla *et al.* 2018; Mordecai *et al.* 2019;
197	Shocket *et al.* 2020), four mosquito–malaria parasite systems (Villena *et al.* 2020), two
198	*Daphnia*–parasite systems (*D. magna–O. colligata*, Kirk *et al.* 2018, 2020; *D. dentifera–M.*
199	*bicuspidata*, Shocket *et al.* 2018b), two amphibian–*B. dendrobatidis* (*Bd*, the causative agent of
200	chytridiomycosis) systems (Cohen *et al.* 2017), and one crab–rhizocephalan barnacle parasite
201	system (*E. depressus–L. Panopaei*, Gehman *et al.* 2018). While the mosquito systems are mainly
202	studied with respect to human disease, here we leverage the rich data on their thermal
203	dependence to explore thermal scaling of the mosquito–parasite interaction, which in turn affects
204	transmission to humans.
205	Host performance, individual-level parasitism, and population-level parasitism were
206	measured using different metrics in different systems (Table 1; Supporting Information). We
207	report a single host T_{opt} value for eleven of the systems, but two host T_{opt} values (individual and
208	population) for the amphibian systems. This is because individual-level parasitism for cold-
209	adapted and warm-adapted amphibians was measured in the lab across one and two host species,
210	respectively (Cohen *et al.* 2017), but the thermal response of population-level parasitism was

reported as *Bd* prevalence in the field as a function of environmental temperature for 235

surveyed species, where cold- and warm-adapted amphibians were categorized as those at

locations where 50-year mean temperature was <15°C or >20°C, respectively (Cohen *et al.*

2017). We therefore used separate T_{opt} values for our individual-level host performance

(measured as thermal preference in the lab) and population-level host performance (measured as

the mean climatic temperature experienced in the field across surveyed species).

While we were generally constrained to use whichever metric of parasitism or host

performance was reported for a study, we had access to thermal response data for several

different metrics in the mosquito–parasite systems. We chose to use infected days as our measure

of individual-level parasitism in these eight systems because it is a composite metric using

thermal performance data on parasite development rate within the mosquito, vector competence

(the mosquito's ability to acquire and transmit the parasite), and mosquito survival. We were also

interested in how this choice affected our findings of how closely T_{opt} of individual-level

parasitism matched T_{opt} of population-level parasitism and T_{opt} of host performance. Generally,

using vector competence as a metric of individual-level parasitism gave similar results to our

main metric of infected days, but parasite development rate tended to exhibit higher T_{opt} values

than the other two metrics (Fig. S10). The Supporting Information contains further details on

both our model and the methods used to analyze the empirical systems.

Using these thirteen systems, we tested for a significant correlation by comparing

population-level parasitism T_{opt} to individual-level parasitism T_{opt} using the *cor.test* function in

the R package *stats* (R Core Team 2020; method = *pearson*). In some of these systems, metrics

used to calculate individual-level parasitism T_{opt} are also a subset of the metrics used to calculate

population-level parasitism T_{opt}. For example, the thermal response of mosquito mortality rate is

one of several components used to calculate individual-level parasitism in mosquito–parasite

systems, and also one of many components used to calculate population-level parasitism in the

same systems. As a result, observed thermal optima for population-level parasitism are not

independent from observed thermal optima for individual-level parasitism. While this can affect

the interpretation of the observed correlation across systems and whether or not it should be

considered statistically significant, we argue that the partial dependence of population-level

parasitism on what occurs at the individual-level is biologically realistic for many systems.

Finally, using the same systems, we compared T_{opt} for individual- and population-level

parasitism to T_{opt} of host performance to determine if parasitism generally peaked at

temperatures away from host thermal optima as predicted by the thermal mismatch hypothesis

(Cohen *et al.* 2017, 2019a, b), and if so, whether these thermal mismatches at the individual level

corresponded with thermal mismatches at the population level.

246
247

248   TABLE 1. Thermal optima of parasitism and host performance for thirteen host – parasite
249   systems.
250

Host – parasite system	Measure of host performance	Measure of individual-level parasitism	Measure of population-level parasitism	Host performance T_{opt} (°C)	Individual-level parasitism T_{opt} (°C)	Population-level parasitism T_{opt} (°C)	Reference
Culex pipiens – West Nile virus	Adult reproduction weighted by lifespan	Infected days	R_0	15.9	23.5	24.5	Shocket et al. 2020
Aedes aegypti - dengue virus	Adult reproduction weighted by lifespan	Infected days	R_0	27.6	29.3	29.1	Mordecai et al. 2017
Aedes aegypti - Zika virus	Adult reproduction weighted by lifespan	Infected days	R_0	27.7	28.9	28.9	Tesla et al. 2018
Culex annulirostris – Ross River virus	Adult reproduction weighted by lifespan	Infected days	R_0	25.6	24.0	26.4	Shocket et al. 2018a
Anopholes gambiae - Plasmodium vivax	Adult reproduction weighted by lifespan	Infected days	Transmission suitability	24.4	23.6	25.0	Villena et al. 2020
Anopholes gambiae - Plasmodium falciparum	Adult reproduction weighted by lifespan	Infected days	Transmission suitability	24.4	23.4	25.0	Villena et al. 2020
Anopholes stephensi - Plasmodium vivax	Adult reproduction weighted by lifespan	Infected days	Transmission suitability	24.0	23.0	24.6	Villena et al. 2020
Anopholes stephensi - Plasmodium falciparum	Adult reproduction weighted by lifespan	Infected days	Transmission suitability	24.0	22.9	24.8	Villena et al. 2020
Cold-adapted amphibians - *Batrachochytrium dendrobatidis*	Temperature preference for individuals; mean climatic temperature for populations †	Parasite growth rate on host	Prevalence	17.9 (10.5) †	26.0	20.5	Cohen et al. 2017
Warm-adapted amphibians - *Batrachochytrium dendrobatidis*	Temperature preference for individuals; mean climatic temperature for populations †	Parasite growth rate on host	Prevalence	23.5 (23.9) †	12.0	15.9	Cohen et al. 2017
Eurypanopeus depressus - Loxothylacus panopaei	Uninfected host survival	Parasite lifetime reproduction	R_0	18.3	15.9	14.7	Gehman et al. 2018
Daphnia dentifera - Metschnikowia bicuspidata	Uninfected intrinsic growth rate	Spore load	R_0	26.0*	20.0*	26.0	Shocket et al. 2018b
Daphnia magna – Ordospora colligata	Lifetime reproduction	Spore load	R_0	16.2*	11.8*	18.8	Kirk et al. 2018, 2020

* T_{opt} for these values was determined as the temperature at which the measure was maximized
along a range of discrete experimental temperatures, rather than the temperature maximizing the
measure along a fitted continuous curve, as is the case for each other value in the table. For
discrete temperature range used, see Supporting Information.
† The first number listed is individual-level host T_{opt}. The number in parentheses is population-
level T_{opt}.
257

RESULTS

Differences in host–parasite biology and the expected relationship between parasitism T_{opt}

across levels

As hypothesized, the model showed that T_{opt} for individual- and population-level

parasitism may be correlated or uncorrelated, depending on the relationships among the trait

thermal responses. When the thermal responses of all parameters in the population-level

parasitism model were independent from each other and from the thermal response of individual-

level parasitism, we did not observe any relationship between the thermal optima of parasitism

across levels (Fig. 1a). Similarly, there was no relationship between thermal optima in the

scenario in which the thermal response of parasite-induced mortality was proportional to the

thermal response of parasite load (Fig. 1b). This means that in this scenario, even though a

parameter that partially determined the thermal response of population-level parasitism was

directly dependent on the thermal response of individual-level parasitism, we still did not

observe a relationship between the thermal optima. Conversely, when the probability of infection

after contact was proportional to parasite load across the thermal range, we observed a strong,

positive relationship between the thermal optima for parasitism across levels (Fig. 1c). Though

the relationship was significant and positive, it does not fall upon the 1:1 line (95% confidence

interval; Fig. 1c). Additionally, while many simulated host–parasite systems are observed below

the 1:1 line (i.e., individual-level parasitism T_{opt} was greater than population-level parasitism

T_{opt}), there were no systems in which population-level parasitism T_{opt} greatly exceeded

individual-level parasitism T_{opt} under these model conditions. Finally, when both probability of

infection after contact and parasite-induced mortality were proportional to parasite load across

the thermal range, the correlation between individual-level parasitism T_{opt} and population-level

282 parasitism T_{opt} was strongly positive (Fig. 1d), closely resembling the scenario in which only

283 probability of infection after contact was proportional to parasite load (Fig. 1c).

Figure 1. Thermal optima of individual parasite burden may be either uncorrelated (top panels) or positively correlated (bottom panels) with thermal optima of R_0 at the population level. The thermal optima of parasitism in 1000 simulated host–parasite systems at both the population level (R_0) and individual level (parasite load) for four different modeled scenarios: a) no relationship between any parameters in the R_0 model (Eq. 2) and parasite load (Eq. 1); b) parasite-induced mortality in the R_0 model is proportional to parasite load; c) probability of infection after contact in the R_0 model is proportional to parasite load; d) both parasite-induced mortality and probability of infection after contact in the R_0 model are proportional to parasite load. Dashed black lines represent the 1:1 lines. Solid black lines and shaded bands represent the best fit linear regressions and 95% confidence intervals, respectively.

We found qualitatively similar results for several of our additional model scenarios.

Specifically, we found similar results when thermal response parameters were drawn from

normal distributions (Fig. S2), when parameter thermal breadth was larger (Fig. S3), when peak

contact rates occurred at warmer temperatures (Fig. S4), and when the population-level

parasitism model (Eq. 1) was altered to allow for chronic infections (Fig. S7) or for

environmentally-transmitted parasites (Fig. S8). However, we found that adding a baseline level

of individual-level parasitism across the temperature range, such that individual-level parasitism

(and probability of infection when it is modeled as proportional to individual-level parasitism) is

low but not zero when $T < T_{min}$ and $T > T_{max}$, led to qualitatively different results when

probability of infection was proportional to individual-level parasitism. In this case, a number of

systems exhibited population-level parasitism T_{opt} much warmer than individual-level parasitism

T_{opt} (Fig. S5c-d), a pattern we did not observe in Figs. 1c-d. Allowing for individual-level

parasitism to have a baseline level and for peak contact rates to occur at warmer temperatures led

to even more systems with population-level parasitism $T_{opt} \gg$ individual-level parasitism T_{opt}

(Fig. S6c-d).

314   ***Relationship between parasitism T_{opt} across levels in empirical systems***

The thirteen empirical systems we analyzed demonstrated a range of thermal optima for

individual-level parasitism (11.8°C – 29.3°C) and population-level parasitism (14.7°C – 29.1°C).

We found that population-level parasitism T_{opt} was significantly correlated with individual-level

parasitism (Fig. 2; Pearson correlation = 0.83, p = 0.0005, n = 13). Of the thirteen systems, nine

exhibited thermal optima at individual and population levels that were within 2.5°C of each

other. The largest difference across levels appeared in the *D. magna–O. colligata* system in

which individual-level parasitism peaked at a temperature that is 7°C cooler than the peak for

population-level parasitism. Nine of the thirteen systems exhibited higher thermal optima for

population-level parasitism than individual-level parasitism, though in most cases this difference

was small (Fig. 2; points above the dashed 1:1 line).

Figure 2. The thermal optima of population-level parasitism are positively correlated with

the thermal optima of individual-level parasitism in empirical systems. Points show

estimates of the optimal temperature (T_{opt}) for population-level parasitism versus T_{opt} of

individual-level parasitism for thirteen host–parasite systems. We found a significant positive

correlation between T_{opt} of population-level parasitism and T_{opt} of individual-level parasitism

(Pearson correlation = 0.83, p = 0.0005, n = 13), and found that nine of the thirteen systems

exhibited higher thermal optima for population-level parasitism than for individual-level

parasitism (i.e., fall above the dashed 1:1 line). Circles represent mosquito – parasite systems and

diamonds indicate non-mosquito – parasite systems.

Relationship between parasitism T_{opt} and host performance T_{opt} in empirical systems

Next, we compared the thermal optima of parasitism at both levels to T_{opt} for host

performance (Fig. 3). If a system was situated at the origin in Fig. 3, it would have individual-

level parasitism, population-level parasitism, and host performance maximized at the same

temperature (i.e., no thermal mismatches exist). Displacement from the origin represents

parasitism peaking at temperatures away from where host performance peaks (i.e., thermal

mismatches at one or both levels). We found that over half (7/13) of the systems were situated

close to the origin (arbitrarily defined here as within Euclidian distance of 2.5°C, represented by

the circle in Fig. 3). However, all seven of these are mosquito–parasite systems, with the only

mosquito system exhibiting strong thermal mismatches being *Cx. pipiens*–West Nile virus, in

which both infected days and R_0 peak at temperatures nearly 10°C higher than host performance

(Fig. 3). The cold adapted amphibian–*B. dendrobatidis* system was also situated in the far upper-

right quadrant (Fig. 3), showing strong thermal mismatches at both levels of parasitism, as

shown in Cohen *et al.* (2017). Two systems showed cold-temperature mismatches at both levels:

the crab *E. depressus*–*L. panopaei* and warm-adapted amphibian–*B. dendrobatidis* host–parasite

systems. The zooplankton *D. dentifera*–*M. bicuspidata* system exhibited a thermal mismatch at

cold temperatures at the individual level but no mismatch at the population level. Finally, the

zooplankton *D. magna*–*O. colligata* system showed a distinct pattern of differing thermal

mismatches across levels (Fig. 3): individual-level parasitism was maximized at a lower

temperature than host performance while R_0 peaked at a higher temperature than host

performance. None of the thirteen systems showed positive individual-level mismatches but

negative population-level mismatches; therefore, the lower-right quadrant of Fig. 3 is empty. The

six systems that experienced thermal mismatches generally supported our hypothesis that if

mismatches occured they would be at both levels and in the same direction, as more of these

systems were situated in the upper-right or lower-left quadrants of Fig. 3 compared to the upper-

left or lower-right quadrants.

364
Figure 3. Thermal matches and mismatches tended to be correlated across levels of
biological organization. The difference between T_{opt} of population-level parasitism and host
performance (y-axis) is plotted against the difference between T_{opt} of individual-level parasitism
and host performance (x-axis) for thirteen host–parasite systems. Systems situated at the origin
had population-level parasitism, individual-level parasitism, and host performance all maximized
at the same temperature (i.e., no thermal mismatches exist), while displacement from the origin
represents parasitism peaking at temperatures away from where host performance peaks (i.e.,
thermal mismatches at individual or population levels). All seven of the systems situated close to
the origin (within Euclidian distance of 2.5°C, represented by the black circle) were mosquito –
parasite systems (circles; in contrast to systems with non-mosquito hosts: diamonds).
375

DISCUSSION

If we are to best predict how warming temperatures will affect host–parasite systems, we

need to understand if effects on parasitism at one level of organization correspond with similar

effects at another level. Moreover, understanding how these effects on parasitism compare to

thermal effects on uninfected hosts can offer a more complete lens into how climate change will

affect hosts. Our model found that small differences in the underlying biology of host–parasite

systems can substantially alter the expected relationship between the thermal optima of

parasitism in individual hosts and host populations, ranging from strong positive correlations to

no correlation of T_{opt} across levels (Fig. 1). In our examination of thirteen empirical systems, we

found a significant positive relationship between T_{opt} for parasitism at the two levels (Fig. 2).

Additionally, while individual- and population-level parasitism both peaked at temperatures

away from the host optimum in some systems, supporting the thermal mismatch hypothesis, this

was not the case in seven of the eight mosquito–parasite systems (Fig. 3), suggesting thermal

mismatches may be more common in certain types of host–parasite systems. Generally, our

results show that information on the temperature-dependence, and specifically the thermal

optimum, of a host–parasite system at either the individual- or population-level should provide a

useful—though not quantitatively exact—baseline for predicting temperature dependence at the

other level in a variety of host–parasite systems.

We found a significant positive correlation between T_{opt} of population-level parasitism

and T_{opt} of individual-level parasitism, suggesting that the effects of warming on parasitism may

often be in the same direction across levels. However, we also found that population-level

parasitism tended to peak at slightly warmer temperatures than individual-level parasitism in the

majority of systems (Fig. 2), meaning that these systems may experience small temperature

ranges in which warming leads to increases in population-level parasitism but decreases in

individual-level parasitism. The greatest differences in T_{opt} were observed in the zooplankton *D.*

magna–O. colligata system, the cold-adapted amphibians–*B. dendrobatidis* system, and the

zooplankton *D. dentifera–M. bicuspidata* system (Fig. 2, Table 1). It is difficult to definitively

parse out the causes of T_{opt} differences in the two amphibian systems since individual-level *Bd*

parasitism was measured on either one or two frog species in the lab while population-level

prevalence by the same parasite was synthesized across many field studies encompassing 235

host species (Cohen *et al.* 2017). However, the *D. magna–O. colligata* system can provide an

illuminating example. Contact rate in this host–microsporidian parasite system is maximized at

30.1°C (Kirk *et al.* 2019). This is nearly 20°C warmer than T_{opt} for individual-level parasitism

(11.8°C) and is thus one of the key factors that pulls the thermal optimum of R_0 away from

individual-level parasitism T_{opt} to a warmer temperature (18.8°C). Moreover, in the mosquito–

parasite systems, which tended to have high population-level parasitism T_{opt} (Fig. 2), biting

rate—a driver of host contact—usually has one of the highest optimal temperatures of all

measured traits (Mordecai *et al.* 2013, 2017, 2019; Shocket *et al.* 2018a; Villena *et al.* 2020).

These few systems represent only a small slice of the diversity of host–parasite systems that

exist; however, they may represent a relatively greater portion of the diversity in thermal trait

biology and temperature-dependent metabolic rates because these rates are often conserved

across systems (Brown *et al.* 2004; Dell *et al.* 2011; Molnár *et al.* 2017)**.** More empirical work

measuring thermal responses of traits related to parasite transmission in other types of host–

parasite systems, such as those with helminth or bacterial parasites—both of which have been

shown to be affected by climate (Ben-Haim *et al.* 2003; Ben-Horin *et al.* 2013; Mignatti *et al.*

2016)—can help broaden our perspective on the scaling of parasitism from individuals to

populations in light of climate change.

The positive relationship observed for T_{opt} across levels in the thirteen empirical systems

qualitatively matched the positive relationship observed in the two modeled scenarios in which

probability of infection was proportional to parasite load (Fig. 1c-d). However, this should not be

taken as evidence that these empirical systems follow the same described dynamics or meet the

same assumptions of these modeled scenarios, as the underlying biology differs greatly between

many of these systems, and the models that have been used to describe their dynamics differ both

between the systems and from the simple model developed here (e.g., Briggs *et al.* 2010;

Mordecai *et al.* 2017; Gehman *et al.* 2018; Shocket *et al.* 2018b; Kirk *et al.* 2020). Taken

together, this means that the model results (Figs. 1, S2-S8) are useful for our understanding of

how different T_{opt} relationships are possible under different plausible scenarios, but should not be

used as statistical null hypotheses for our thirteen empirical systems. The contribution of this

approach—pairing mechanistic models and thermal responses across biological levels under

different assumptions—is that we can observe how different systems or thermal biology can lead

to different patterns in T_{opt} across levels. Indeed, while our additional model scenarios generally

supported our main finding that individual- and population-level parasitism can be correlated or

uncorrelated and showed that this is qualitatively robust across different model structures and

assumptions regarding thermal response shape, they also illustrated how changes to thermal

biology can alter these patterns. For example, if peak contact rates tend to occur at warmer

temperatures than other traits, we may observe more systems that have population-level

parasitism $T_{opt} >$ individual-level parasitism T_{opt} (Fig. S4). Also, despite not observing systems

in the far upper-left space (where population-level parasitism $T_{opt} \gg$ individual-level parasitism

T_{opt}) for our model simulations when probability of infection scaled with individual-level

parasitism (Fig. 1c-d), additional scenarios showed that this is possible if individual-level

parasitism does not decrease completely to zero when $T < T_{min}$ and $T > T_{max}$ (Figs. S5-S6).

The thermal mismatch hypothesis (Cohen *et al.* 2017, 2019a, b, 2020) has been proposed

as an approach for understanding the effects of climate change on host–parasite systems,

predicting that T_{opt} for parasitism should occur at temperatures away from host T_{opt} (i.e., a

thermal mismatch) due to larger organisms (hosts) generally having narrower thermal breadth

than smaller organisms (parasites; Rohr *et al.* 2018). While empirical evidence shows that

thermal mismatches do occur at the population level across many systems (Cohen *et al.* 2020),

the novel hypothesis we tested here is that thermal mismatches at the individual level will

correspond with similar thermal mismatches at the population level. Of the five non-mosquito

systems we explored—including the two amphibian–*Bd* systems investigated previously—

parasitism tended to peak at temperatures away from the host's optimum in each, meaning that

these systems experience thermal mismatches at one or both levels. If our hypothesis was to be

supported, we would expect most systems experiencing thermal mismatches to fall in the upper-

right or lower-left quadrants of Fig. 3, rather than the upper-left or lower-right. We found that

this was generally the case, with two systems experiencing thermal mismatches at warmer

temperatures at both levels (Fig. 3; upper-right) and two systems experiencing thermal

mismatches at cooler temperatures at both levels (Fig. 3; lower-left). The major exception was

the *D. magna–O. colligata* system, in which host T_{opt} occurs at a temperature intermediate to the

relatively cool T_{opt} for individual-level parasitism and relatively warm T_{opt} population-level

parasitism (Fig. 3; upper-left). The other zooplankton host system we explored, *D. dentifera–M.*

bicuspidata, exhibited a thermal mismatch at cooler temperatures at the individual level but no

mismatch at the population level.

Evidence for thermal mismatches occurring at either level was weak in the mosquito–

parasite systems. Seven of eight mosquito–parasite systems were situated close to the origin (Fig.

3), suggesting that parasitism and host performance are maximized at close to the same

temperatures and that no thermal mismatches are occurring at either level. Notably, when testing

for thermal mismatches at the population level, (Cohen *et al.* 2020) also found that thermal

mismatch effects were strongest in systems without vectors (or intermediate hosts). The

exception in our study, *Cx. pipiens*–West Nile virus, in which parasitism at both levels peaks at

temperatures much higher than host performance because *Cx. pipiens* adult lifespan is much

longer at low temperatures (Shocket *et al.* 2020), is a system with a predominantly temperate

distribution in North America, compared to the more sub-tropical and tropical distributions of

malaria, dengue, Zika, Ross River virus, and their respective mosquito vectors. Why we

observed this pattern in mosquito systems will require further investigation in different vector-

borne systems across biological levels, including those systems with non-mosquito vectors such

as ticks or sandflies.

We propose that situating host–parasite systems along both individual- and population-

level axes as in Fig. 3 can provide a diagnostic framework for looking for thermal mismatches

across biological levels. While our results do not conclusively address how general thermal

mismatches at both levels may be, our finding that individual- and population-level T_{opt} are

strongly related (Fig. 2) provides a general rule of thumb that, if a host–parasite system does

experience thermal mismatches, they are likely to occur at both levels and in the same direction

(though note the exception of *D. magna–O. colligata*; Fig. 3). This has major implications for

our understanding of the effects of climate change on host-parasite systems, because if T_{opt} for

both individual- and population-level parasitism is at a higher temperature than T_{opt} for host

performance, warming may not only decrease host performance itself, but also lead hosts to

experience increasing effects of parasitism (Fig. 3; upper-right quadrant). Alternatively, if T_{opt}

for parasitism is at temperatures cooler than host performance optima, decreased host

performance at warmer temperatures may be partially offset by decreased effects of parasitism

(Fig. 3; lower-left quadrant). This is in line with recent projections that climate change may lead

to hosts from cooler climates to experience increased parasitism, and hosts from warmer climates

to experience decreased parasitism (Cohen *et al.* 2020).

To better understand how temperature affects parasitism and to make better predictions

for how climate change will affect host–parasite systems across biological levels, we highlight

two important research directions. First, when possible, studies that use experiments to

investigate effects of temperature at the individual level can also be used to parameterize simple

models of population-level parasitism (a 'bottom-up' approach). Second, field studies

investigating population-level parasitism under different climate or weather conditions, often

undertaken by measuring the proportion of individuals infected, should also aim to record

measures of individual-level parasitism (a 'top-down' approach). The best direct measure would

be to record the parasite load or an analogous metric measuring the parasite on or within the host,

but in cases where this is not possible, indirect measures such as host condition may still be

informative.

Overall, after our model illustrated that either positive or flat relationships between T_{opt}

across levels were plausible, we found a strong positive relationship in the thirteen systems we

analyzed, suggesting that the effects of warming on individual-level outcomes may beget similar

outcomes at the population level in various host–parasite systems. Many studies have

investigated the effects of temperature on parasitism (reviewed, for example, in Marcogliese

2008; Lafferty 2009; Rohr *et al.* 2011; Altizer *et al.* 2013; Lafferty & Mordecai 2016; Claar &

Wood 2020), though often at only one of either the individual or population levels. Our findings

build upon these studies by emphasizing that how temperature affects individual-level parasitism

can scale up to affect parasite transmission, prevalence, and the potential for epidemics in host

populations, though not always in a 1:1 manner. While this is an important first step toward

predicting the effects of climate change on systems by allowing researchers to primarily focus on

parasitism at one level when necessary, future empirical work that investigates the thermal

dependence of a wider range of host–parasite systems across levels, including in systems with

different types of hosts (e.g., plants, non-amphibian vertebrates) and other types of parasites

(e.g., helminths, bacterial parasites), will be necessary for investigating the generality of this

result. More broadly, tying together more theory and empirical results to provide general

predictions for how climate change will affect parasitism from individuals to ecosystems in a

diverse range of hosts and parasites remains an urgent priority as accelerating climate change

makes potentially catastrophic temperature increases of $> 2°C$ increasingly likely by 2050.

LITERATURE CITED

Altizer, S., Ostfeld, R.S., Johnson, P.T.J., Kutz, S. & Harvell, C.D. (2013). Climate Change and Infectious Diseases: From Evidence to a Predictive Framework. *Science*, 341, 514–519.

Anderson, R.M. & May, R.M. (1978). Regulation and Stability of Host-Parasite Population Interactions: I. Regulatory Processes. *The Journal of Animal Ecology*, 47, 219.

Anderson, R.M. & May, R.M. (1979). Population biology of infectious diseases: part 1. *Nature*, 280, 361–367.

Angilletta, M.J. (2006). Estimating and comparing thermal performance curves. *Journal of Thermal Biology*, 31, 541–545.

Angilletta, M.J. (2009). *Thermal adaptation: a theoretical and empirical synthesis*. Oxford University Press.

Ben-Ami, F., Regoes, R.R. & Ebert, D. (2008). A quantitative test of the relationship between parasite dose and infection probability across different host–parasite combinations. *Proc. R. Soc. B*, 275, 853–859.

Ben-Haim, Y., Zicherman-Keren, M. & Rosenberg, E. (2003). Temperature-regulated bleaching and lysis of the coral Pocillopora damicornis by the novel pathogen Vibrio coralliilyticus. *AEM*, 69, 4236–4242.

Ben-Horin, T., Lenihan, H.S. & Lafferty, K.D. (2013). Variable intertidal temperature explains why disease endangers black abalone. *Ecology*, 94, 161–168.

Briere, J.-F., Pracros, P., Le Roux, A.-Y. & Pierre, J.-S. (1999). A Novel Rate Model of Temperature-Dependent Development for Arthropods. *Environ Entomol*, 28, 22–29.

Briggs, C.J., Knapp, R.A. & Vredenburg, V.T. (2010). Enzootic and epizootic dynamics of the chytrid fungal pathogen of amphibians. *Proceedings of the National Academy of Sciences*, 107, 9695–9700.

Brown, J.H., Gillooly, J.F., Allen, A.P., Savage, V.M. & West, G.B. (2004). TOWARD A METABOLIC THEORY OF ECOLOGY. *Ecology*, 85, 1771–1789.

Bruno, J.F., Selig, E.R., Casey, K.S., Page, C.A., Willis, B.L., Harvell, C.D., *et al.* (2007). Thermal Stress and Coral Cover as Drivers of Coral Disease Outbreaks. *PLoS Biol*, 5, e124.

Casey, T.M. (1976). Activity Patterns, Body Temperature and Thermal Ecology in Two Desert Caterpillars (Lepidoptera: Sphingidae). *Ecology*, 57, 485–497.

Claar, D.C. & Wood, C.L. (2020). Pulse Heat Stress and Parasitism in a Warming World. *Trends in Ecology & Evolution*, 35, 704–715.

Cohen, J.M., Civitello, D.J., Venesky, M.D., McMahon, T.A. & Rohr, J.R. (2019a). An interaction between climate change and infectious disease drove widespread amphibian declines. *Glob Change Biol*, 25, 927–937.

Cohen, J.M., McMahon, T.A., Ramsay, C., Roznik, E.A., Sauer, E.L., Bessler, S., *et al.* (2019b). Impacts of thermal mismatches on chytrid fungus Batrachochytrium dendrobatidis prevalence are moderated by life stage, body size, elevation and latitude. *Ecol Lett*, 22, 817–825.

Cohen, J.M., Sauer, E.L., Santiago, O., Spencer, S. & Rohr, J.R. (2020). Divergent impacts of warming weather on wildlife disease risk across climates. *Science*, 370, eabb1702.

Cohen, J.M., Venesky, M.D., Sauer, E.L., Civitello, D.J., McMahon, T.A., Roznik, E.A., *et al.* (2017). The thermal mismatch hypothesis explains host susceptibility to an emerging infectious disease. *Ecol Lett*, 20, 184–193.

Day, T. (2001). Parasite transmission modes and the evolution of virulence. *Evolution*, 55, 2389–2400.

Dell, A.I., Pawar, S. & Savage, V.M. (2011). Systematic variation in the temperature dependence of physiological and ecological traits. *Proceedings of the National Academy*

579    *of Sciences*, 108, 10591–10596.

580 Ewald, P.W. (1983). Host-Parasite Relations, Vectors, and the Evolution of Disease Severity.
581    *Annu. Rev. Ecol. Syst.*, 14, 465–485.

582 Fenton, A. (2008). Worms and germs: the population dynamic consequences of microparasite-
583    macroparasite co-infection. *Parasitology*, 135, 1545–1560.

584 Fenton, A. (2013). Dances with worms: the ecological and evolutionary impacts of deworming
585    on coinfecting pathogens. *Parasitology*, 140, 1119–1132.

586 Frank, S.A. (1996). Models of Parasite Virulence. *The Quarterly Review of Biology*, 71, 37–78.

587 Gehman, A.-L.M., Hall, R.J. & Byers, J.E. (2018). Host and parasite thermal ecology jointly
588    determine the effect of climate warming on epidemic dynamics. *Proc Natl Acad Sci USA*,
589    115, 744–749.

590 Handel, A. & Rohani, P. (2015). Crossing the scale from within-host infection dynamics to
591    between-host transmission fitness: a discussion of current assumptions and knowledge.
592    *Phil. Trans. R. Soc. B*, 370, 20140302.

593 Harvell, C.D., Montecino-Latorre, D., Caldwell, J.M., Burt, J.M., Bosley, K., Keller, A., *et al.*
594    (2019). Disease epidemic and a marine heat wave are associated with the continental-
595    scale collapse of a pivotal predator (Pycnopodia helianthoides). *Sci. Adv.*, 5, eaau7042.

596 van der Have, T.M. (2002). A proximate model for thermal tolerance in ectotherms. *Oikos*, 98,
597    141–155.

598 Huey, R.B. & Kingsolver, J.G. (1989). Evolution of thermal sensitivity of ectotherm performance.
599    *Trends in Ecology & Evolution*, 4, 131–135.

600 Huey, R.B. & Stevenson, R.D. (1979). Integrating Thermal Physiology and Ecology of
601    Ectotherms: A Discussion of Approaches. *American Zoologist*, 19, 357–366.

602 Kirk, D., Jones, N., Peacock, S., Phillips, J., Molnár, P.K., Krkošek, M., *et al.* (2018). Empirical
603    evidence that metabolic theory describes the temperature dependency of within-host
604    parasite dynamics. *PLoS Biol*, 16, e2004608.

605 Kirk, D., Luijckx, P., Jones, N., Krichel, L., Pencer, C., Molnár, P., *et al.* (2020). Experimental
606    evidence of warming-induced disease emergence and its prediction by a trait-based
607    mechanistic model. *Proc. R. Soc. B.*, 287, 20201526.

608 Kirk, D., Luijckx, P., Stanic, A. & Krkošek, M. (2019). Predicting the Thermal and Allometric
609    Dependencies of Disease Transmission via the Metabolic Theory of Ecology. *The*
610    *American Naturalist*, 193, 661–676.

611 Koelle, K., Pascual, M. & Yunus, M. (2005). Pathogen adaptation to seasonal forcing and
612    climate change. *Proc. R. Soc. B*, 272, 971–977.

613 Kordas, R.L., Harley, C.D.G. & O'Connor, M.I. (2011). Community ecology in a warming world:
614    The influence of temperature on interspecific interactions in marine systems. *Journal of*
615    *Experimental Marine Biology and Ecology*, 400, 218–226.

616 Lafferty, K.D. (2009). The ecology of climate change and infectious diseases. *Ecology*, 90, 888–
617    900.

618 Lafferty, K.D. & Mordecai, E.A. (2016). The rise and fall of infectious disease in a warmer world.
619    *F1000Res*, 5, 2040.

620 Marcogliese, D.J. (2008). The impact of climate change on the parasites and infectious
621    diseases of aquatic animals. *Rev. scii. tech. Off. init. Epiz.*, 27, 467–484.

622 McCallum, H., Fenton, A., Hudson, P.J., Lee, B., Levick, B., Norman, R., *et al.* (2017). Breaking
623    beta: deconstructing the parasite transmission function. *Phil. Trans. R. Soc. B*, 372,
624    20160084.

625 Mideo, N., Nelson, W.A., Reece, S.E., Bell, A.S., Read, A.F. & Day, T. (2011). Bridging scales
626    in the evolution of infectious disease life histories: application. *Evolution*, 65, 3298–3310.

627 Mignatti, A., Boag, B. & Cattadori, I.M. (2016). Host immunity shapes the impact of climate
628    changes on the dynamics of parasite infections. *Proc Natl Acad Sci USA*, 113, 2970–
629    2975.

Molnár, P.K., Sckrabulis, J.P., Altman, K.A. & Raffel, T.R. (2017). Thermal Performance Curves and the Metabolic Theory of Ecology—A Practical Guide to Models and Experiments for Parasitologists. *Journal of Parasitology*, 103, 423.

Mordecai, E.A., Caldwell, J.M., Grossman, M.K., Lippi, C.A., Johnson, L.R., Neira, M., *et al.* (2019). Thermal biology of mosquito-borne disease. *Ecol Lett*, 22, 1690–1708.

Mordecai, E.A., Cohen, J.M., Evans, M.V., Gudapati, P., Johnson, L.R., Lippi, C.A., *et al.* (2017). Detecting the impact of temperature on transmission of Zika, dengue, and chikungunya using mechanistic models. *PLoS Negl Trop Dis*, 11, e0005568.

Mordecai, E.A., Paaijmans, K.P., Johnson, L.R., Balzer, C., Ben-Horin, T., de Moor, E., *et al.* (2013). Optimal temperature for malaria transmission is dramatically lower than previously predicted. *Ecol Lett*, 16, 22–30.

Pedersen, A.B. & Antonovics, J. (2013). Anthelmintic treatment alters the parasite community in a wild mouse host. *Biol. Lett.*, 9, 20130205.

R Core Team. (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Rohr, J.R., Civitello, D.J., Cohen, J.M., Roznik, E.A., Sinervo, B. & Dell, A.I. (2018). The complex drivers of thermal acclimation and breadth in ectotherms. *Ecol Lett*, 21, 1425–1439.

Rohr, J.R., Dobson, A.P., Johnson, P.T.J., Kilpatrick, A.M., Paull, S.H., Raffel, T.R., *et al.* (2011). Frontiers in climate change–disease research. *Trends in Ecology & Evolution*, 26, 270–277.

Shocket, M.S., Ryan, S.J. & Mordecai, E.A. (2018a). Temperature explains broad patterns of Ross River virus transmission. *eLife*, 7, e37762.

Shocket, M.S., Strauss, A.T., Hite, J.L., Šljivar, M., Civitello, D.J., Duffy, M.A., *et al.* (2018b). Temperature Drives Epidemics in a Zooplankton-Fungus Disease System: A Trait-Driven Approach Points to Transmission via Host Foraging. *The American Naturalist*, 191, 435–451.

Shocket, M.S., Verwillow, A.B., Numazu, M.G., Slamani, H., Cohen, J.M., El Moustaid, F., *et al.* (2020). Transmission of West Nile and five other temperate mosquito-borne viruses peaks at temperatures between 23°C and 26°C. *eLife*, 9, e58511.

Tesla, B., Demakovsky, L.R., Mordecai, E.A., Ryan, S.J., Bonds, M.H., Ngonghala, C.N., *et al.* (2018). Temperature drives Zika virus transmission: evidence from empirical and mathematical models. *Proc. R. Soc. B*, 285, 20180795.

Thomas, M.B. & Blanford, S. (2003). Thermal biology in insect-parasite interactions. *Trends in Ecology & Evolution*, 18, 344–350.

Villena, O.C., Ryan, S.J., Murdock, C.C. & Johnson, L.R. (2020). Temperature impacts the transmission of malaria parasites by Anopheles gambiae and Anopheles stephensi mosquitoes. *bioRxiv*, https://doi.org/10.1101/2020.07.08.194472.