Experimental evolution-driven identification of Arabidopsis rhizosphere competence genes in *Pseudomonas protegens*

Erqin Li[1,#], Henan Jiang[1], Corné M.J. Pieterse[1], Alexandre Jousset[2], Peter A.H.M. Bakker[1], Ronnie de Jonge[1]

[1]Plant-Microbe Interactions, Department of Biology, Science4Life, Utrecht University, Padualaan 8, 3584 CH, Utrecht, The Netherlands

[2]Ecology and Biodiversity, Department of Biology, Science4Life, Utrecht University, Padualaan 8, 3584 CH, Utrecht, The Netherlands.

[†]Corresponding author: Ronnie de Jonge; r.dejonge@uu.nl

#Current address: Freie Universität Berlin, Institut für Biologie, Dahlem Center of Plant Sciences, Plant Ecology, Altensteinstr. 6, D-14195, Berlin, Germany, erqinli22@gmail.com

Abstract

Beneficial plant root-associated microorganisms carry out a range of functions that are essential for plant performance. Establishment of a bacterium on plant roots, however, requires overcoming several challenges, including the ability to outcompete neighboring microorganisms and suppression of plant immunity. Forward and reverse genetics approaches have led to the identification of diverse mechanisms that are used by beneficial microorganisms to overcome these challenges such as the production of iron-chelating compounds, biofilm formation, or downregulation of plant immunity. However, how such mechanisms have developed from an evolutionary perspective is much less understood. In an attempt to study bacterial adaptation in the rhizosphere, we employed an experimental evolution approach to track the physiological and genetic dynamics of root-dwelling *Pseudomonas protegens* CHA0 in the *Arabidopsis thaliana* rhizosphere under axenic conditions. This simplified binary one plant, and one bacterium system allows for the amplification of key adaptive mechanisms for bacterial rhizosphere colonization. We found that mutations in global regulators, as well as in genes for siderophore production, cell surface decoration, attachment, and motility accumulated in parallel in our evolutionary experiment, underlining several different strategies of bacterial adaptation to the rhizosphere. In total we identified 35 mutations, including single-nucleotide polymorphisms, smaller indels and larger deletions, distributed over 28 genes in total. Altogether these results underscore the strength of experimental evolution to identify key genes and pathways for bacterial rhizosphere colonization, as well as highlighting a methodology for the development of elite beneficial microorganisms with enhanced root-colonizing capacities that can support sustainable agriculture in the future.

Introduction

Plants are associated with complex microbial communities assembled into a functional microbiome that safeguards optimal plant performance under harsh environmental conditions (1). The rhizosphere microbiome is a particularly interesting hotspot of plant-microbe interactions. Plants deposit up to 44% of their photosynthetically fixed carbon into the rhizosphere, fueling a specific microbial community (2). The microbial species pool in the bulk soil is the source from which the root microbiome is recruited, and plant genotype, immune responses, and environmental factors are postulated to affect this process (3–6). The establishment of beneficial microbial associations requires a high degree of coordination of both plant and microbial responses by means of a continuous molecular dialogue (7, 8). Plant-associated microorganisms can improve plant yield by protecting the plant from abiotic stresses (9), improving plant nutrition and growth (10–12), antagonizing soil-borne pathogens (13), or stimulating plant immunity (14). To exert their beneficial effects on plant performance, bacteria must colonize the roots efficiently and establish significant populations. For example, population densities above 10^5 cells per gram of root are required for efficient suppression of soil-borne plant pathogens by *Pseudomonas* spp. (15, 16). Therefore, bacterial adaptation to the plant root environment may be essential for the successful implementation of microbiome services in agriculture in order to support plant health.

Among all root-dwelling organisms, fluorescent *Pseudomonas* spp. are well characterized in terms of the traits required for their growth in the rhizosphere and for the establishment of beneficial plant-microbe interactions (17). To study bacterial traits involved in efficient root colonization, mutants defective in specific traits suspected to be involved in colonization, are compared to the ability of the parental strain to colonize roots. Such studies have highlighted a range of bacterial traits involved in efficient root colonization, including flagella (18), surface lipopolysaccharides (19), and amino acid synthesis (20). Using random mutagenesis and by determining the fitness of each mutant in competition with its parental strain in the rhizosphere, many other traits important for rhizosphere competence in *Pseudomonas* were discovered (17). Recently, random mutagenesis in *Pseudomonas capeferrum* WCS358 led to the identification of two genes important for gluconic acid biosynthesis which in turn is essential for the suppression of local, flagellin-induced root immune responses (21, 22). Such suppression was shown to be important for rhizosphere competence as gluconic acid-deficient mutants maintain reduced rhizospheric populations (22). In another recent study, genome-wide saturation mutagenesis of *Pseudomonas simiae* WCS417r revealed that 2% of the protein-coding genes are important for successful root colonization (23). Mutations that negatively affect rhizosphere competence, and mutations that

confer a competitive advantage were identified in this study. The observation that mutations can lead to

increased root colonization (23), suggests that there is room for improvement of bacterial fitness in the

rhizosphere.

In the present study we used an experimental evolution approach (24) to study how *Pseudomonas*

protegens CHA0 (CHA0) evolves during repeated colonization of the rhizosphere of the model plant

Arabidopsis thaliana (Arabidopsis). The model biological control agent CHA0 displays broad-spectrum

antagonistic activity against several plant pathogenic fungi and bacteria (11), and its complete genome

sequence is available (25). We performed highly controlled experimental evolution in a gnotobiotic and

carbon-free sand system in which bacteria depend solely on the plant for supply of carbon. Following

inoculation and establishment on the roots, bacterial populations were transferred to new plants, and

this cycle was repeated eight times. We hypothesized that the repeated colonization and establishment

of the bacterial population on the plant root would create an environment in which selection pressure

drives the accumulation of better colonizers. *In vitro* characterization of individual bacterial colonies from

these populations combined with sequencing analysis led to the identification of several evolutionary

trajectories involving 35 distinct mutations that impact social traits representing inter-population

communication and cooperation, carbon source utilization, motility, or biocontrol activity. By combining

experimental evolution with whole genome re-sequencing, we created a powerful screen for the

identification of adaptive mutations with positive effects on rhizosphere colonization.

Results

Mutational events in independent evolutionary lines

We previously studied five experimental evolutionary populations, referred to as lines, of CHA0 evolving

in the rhizosphere of Arabidopsis in a gnotobiotic system. Independent populations were introduced on

the roots and after four weeks of plant growth the populations were transferred to new plants. This cycle

of transferring was repeated eight times (26). To study adaptation at the individual level, here, we isolated

in total 90 bacterial colonies at the end of plant growth cycles 2, 4 and 6. These colonies, and six colonies

from the ancestral population that was initially introduced on the roots, were re-sequenced to an average

depth of 25-fold coverage (minimum 10, maximum 70) and used for the identification of single nucleotide

polymorphisms (SNPs), as well as small and large insertions or deletions (INDELs). The six ancestral

colonies were all identical, indicating that there was no genetic variation in the starting population and all

observed mutations are therefore *de novo* mutations. In total, one or more mutations were detected in

64 evolved colonies, which collectively represent 71% of the 90 characterized bacterial colonies (Table

S1). We identified 5 synonymous substitutions, 20 nonsynonymous substitutions, and 4 deletions ranging

in length from 1 base pair (bp) to about 400 bp, distributed over 22 genes, and 6 additional mutational

events located in intergenic regions (Table 1). Mutations located in intergenic regions possibly affect

transcription of nearby genes via affecting their regulatory proteins binding sites and subsequent changes

in their promoter activity (27, 28). Several mutations were found to be clustered in select genes and/or

regions on the CHA0 genome, e.g. those targeting the response regulator *gacA* (PFLCHA0_RS17965;

NC_021237:4,039,113-4,039,754) or the *OBC3* gene cluster involved in lipopolysaccharide (LPS)

biosynthesis (*NC_021237:2,173,707-2,196,028*)(29), but the majority of the mutations are spread across

the 6.1 Mbp CHA0 genome (Figure 1). Functional characterization of the mutated genes by analysis of

their Clusters of Orthologous Groups (COGs) annotation revealed that the majority of these genes are

involved in transcription (COG term 'K'), signal transduction mechanisms (COG term 'T'), amino acid

transport and metabolism (COG term 'E') and cell wall/membrane/envelope biogenesis (COG term 'M')

(Figure 1, Table 1).

Table 1. Mutations that occurred in 28 genes during Arabidopsis rhizosphere adaptation divided over five replicate CHA0 populations (lines)

Locus tag	Gene name	Description	COG[1]	Line	Total number of alleles	Number of mutations in protein-coding genes			Number of intergenic mutations
						Non-synonymous	Synonymous	Deletions	
PFLCHA0_RS02080	*hutI*	Imidazolonepropionase	F	4	1	0	1	0	0
PFLCHA0_RS03400	*accC*	Biotin carboxylase, acetyl-CoA carboxylase subunit	I	2	1	1	0	0	0
PFLCHA0_RS05510	*nudL*	Coenzyme A pyrophosphatase/nudix hydrolase NudL	L	4	1	0	1	0	0
PFLCHA0_RS06125	*rpoS*	RNA polymerase sigma factor RpoS	K	5	1	1	0	0	0
PFLCHA0_RS08340	*fleQ*	Sigma-54-dependent Fis family transcriptional regulator	T	5	1	1	0	0	0
PFLCHA0_RS08490	*flhA*	Flagellar biosynthesis protein FlhA	N	4	1	0	0	1	0
PFLCHA0_RS09880	*RS09880*	Glycosyltransferase (GT)	M	5	1	1	0	0	0
PFLCHA0_RS09890	*oafA*	O-antigen acetylase	I	1, 3	2	1	0	1	0
PFLCHA0_RS09920	*galE*	UDP-glucose 4-epimerase	M	2	1	1	0	0	0
PFLCHA0_RS11785	*RS11785*	LysR family transcriptional regulator	K	4	1	1	0	0	0
PFLCHA0_RS11820	*RS11820*	PaaI family thioesterase	Q	4	1	0	1	0	0
PFLCHA0_RS12070	*RS12070*	2OG-Fe(II) oxygenase superfamily	S	5	1	0	1	0	0
PFLCHA0_RS13000	*yvaQ2*	Methyl-accepting chemotaxis protein	NT	2	1	0	0	0	1
PFLCHA0_RS14960	*tetR*	TetR/AcrR family transcriptional regulator	K	5	1	1	0	0	0
PFLCHA0_RS17350	*RS17350*	Methyltransferase domain-containing protein	H	1	1	0	0	1	0
PFLCHA0_RS17965	*gacA*	UvrY/SirA/GacA family response regulator transcription factor	K	1, 2, 4	6	5	0	0	1
PFLCHA0_RS18525	*RS18525*	ABC transporter substrate-binding protein	E	5	1	1	0	0	0
PFLCHA0_RS21265	*pvdS*	RNA polymerase factor sigma-70	K	1	1	0	0	0	1
PFLCHA0_RS21275	*RS21275*	Transporter substrate-binding domain-containing protein	ET	1	1	1	0	0	0
PFLCHA0_RS21855	*wbpM*	Polysaccharide biosynthesis protein/NDP-sugar epimerase	GM	1	1	1	0	0	0
PFLCHA0_RS22600	*gacS*	Hybrid sensor histidine kinase/response regulator	T	3	1	1	0	0	0
PFLCHA0_RS22950	*argT5*	ABC transporter substrate-binding protein/Lysine-arginine-ornithine-binding periplasmic protein ArgT	ET	4	1	0	0	1	0
PFLCHA0_RS25175	*mraZ*	Division/cell wall cluster transcriptional repressor MraZ	K	2	1	0	0	0	1
PFLCHA0_RS26215	*osmY*	Osmotically-inducible protein OsmY/BON domain-containing protein	S	2	1	0	0	0	1
PFLCHA0_RS27515	*rpsH*	30S ribosomal protein S8	J	4	1	0	0	0	1
PFLCHA0_RS30075	*sadB*	Surface attachment defective (SadB) ortholog/HDOD domain-containing protein	T	1	2	2	0	0	0
PFLCHA0_RS30120	*tssM*	Type VI secretion system membrane subunit TssM	S	2	1	0	1	0	0
PFLCHA0_RS31060	*nlpD*	Lipoprotein nlpD/lppB/LysM domain-containing protein	D	3	1	1	0	0	0
Total	28				35	20	5	4	6

 [1]Clusters of Orthologous Groups (COGs); https://www.ncbi.nlm.nih.gov/COG/(30)

Identification of potential root colonization genes

Bacterial genes that are involved in colonization of plant roots can be revealed by identifying beneficial mutations that evolve during adaption of bacteria to the rhizosphere environment and have positive effects on root colonization. Over time, such mutants will outcompete the ancestral strain and become dominant in the bacterial population. The observation that only a limited number of mutations accumulated relative to the total number of genes within the genome across the entire experiment, makes it highly unlikely that the same gene would acquire several changes by chance in independent evolutionary lines. Nevertheless, we observe recurrent mutations in several of the same genes and/or pathways (Table 1), which is a strong indication for adaptive evolution. Genes or pathways that acquired mutations in multiple independent CHA0 populations included *gacA* and *gacS*, the *OBC3* gene cluster (*oafA*; *galE*; *PFLCHA0_RS09880* (29)), and a putative pyoverdine siderophore biosynthesis cluster (*pvdS*; *PFLCHA0_RS21275*). Other genes were targeted more than once in the same population, such as *sadB*. Because these genes were repeatedly identified in the CHA0 evolution experiment, they can be assumed to contribute significantly to bacterial fitness in the rhizosphere (Table 1, Figure 2). Moreover, it suggests that the independent evolutionary lines converge on similar evolutionary trajectories involving overlapping biological processes and molecular mechanisms.

To illustrate the diverse evolutionary trajectories, we constructed phylogenetic trees of each replicate population (line) using all mutations that were detected (Figure 3). In each line, both the number and depth of branches are very different (Figure 3, Table 2). In line 1 and 3, the populations were swept early by *oafA* mutants and later by specific *gac* mutants, while in line 2 and 4, co-existence of multiple genotypes characterized by *gac* mutations was observed. The evolved isolates contain one to four mutations in general, and in each line, 3 (line 3) to 9 (line 2 and 4) mutations were identified in total. All mutations were unique for each line, as is expected for independently evolving populations.

The frequency of each mutation (Table 2) was determined from the six bacterial colonies that were isolated and sequenced from each evolutionary line at cycles 2, 4 and 6. To further investigate and verify the changes in mutation frequency in the entire CHA0 population at the end of a given cycle, we set out to determine the frequency of three of the mutations, i.e. $gacA^{D49Y}$, $gacA^{D54Y}$, and $gacS^{G27D}$, with increased accuracy and resolution during long-term rhizosphere adaptation. To this end, we developed a PCR-based high-resolution melting (HRM) analysis using mutation-specific HRM probes (Table S2), that allow for the accurate quantification of mutant frequency across a wide range for all three mutations ($p<0.001$; Figure S1). Using this method, we found that the population-level frequency of these three mutations in the

three respective evolutionary lines at the end of each of the eight rhizosphere adaptation cycles
corresponded well with the frequency previously obtained from the six cultured and sequenced isolates
(Table 2, Figure S2). These findings indicate that the culture-based quantification of mutant frequency
provides a realistic measure for population-level frequency. Moreover, it can be seen that whereas the
$gacS^{G27D}$ mutation reaches fixation after eight cycles of rhizosphere adaptation, the $gacA^{D49Y}$ and $gacA^{D54Y}$
mutants stabilize at around 50% and 25%, respectively, indicative of frequency-dependent (FD) selection
(Figure S2). FD selection describes the phenomenon that the fitness of a particular genotype or phenotype
is dependent on its frequency. Such context-dependency has been frequently linked to cheating
behaviour in which microbial cells that lack or limit the production of certain costly compounds benefit
from other cells that do produce these compounds. When a minimal amount of such compound increases
genotype or phenotype fitness, FD selection can occur resulting at the stabilization of the mutation
frequency.

Table 2. Nomenclature for CHA0 mutations and variants and their frequency/cycle in each respective population.

Line	Locus tag	Gene name	DNA sequence change[1]	Amino acid change (three-letter code)[2]	Standard nomenclature (used in text)	Frequency in the population (x/6)			
						Cycle 2	Cycle 4	Cycle 6	RQ-HRM[3]
1	PFLCHA0_RS09890	*oafA*	c.1005C>A	p.Tyr335X	Y335X	4/6	6/6	6/6	
1	PFLCHA0_RS17350	*RS17350*	c.230delC	p.Ala77Ala.fsX14	A77A.fsX14	2/6	6/6	6/6	
1	PFLCHA0_RS17965	*gacA*	c.145G>T	p.Asp49Tyr	D49Y	-	2/6	4/6	x
1	PFLCHA0_RS21855	*wbpM*	c.235G>C	p.Gly79Arg	G79R	-	1/6	-	
1	PFLCHA0_RS21275	*RS21275*	c.233G>T	p.Arg78Leu	R78L	-	1/6	-	
1	PFLCHA0_RS21265	*pvdS*	c.-33T>C	NA	-33T>C	-	-	1/6	
1	PFLCHA0_RS30075	*sadB*	c.548G>C	p.Arg183Pro	R183P	-	-	1/6	
1			c.773T>A	p.Leu258Gln	L258Q	-	-	2/6	
2	PFLCHA0_RS17965	*gacA*	c.548A>C	p.Tyr183Ser	Y183S	2/6	1/6	-	
2	PFLCHA0_RS09920	*galE*	c.94G>A	p.Val32Met	V32M	2/6	5/6	6/6	
2	PFLCHA0_RS30120	*tssM*	c.1336C>T	p.Leu446Leu	1336C>T	1/6	-	-	
2	PFLCHA0_RS03400	*accC*	c.1237G>A	p.Glu413Lys	E413K	-	4/6	6/6	
2	PFLCHA0_RS13000	*yvaQ2*	c.-9G>T	NA	-9G>T	-	1/6	-	
2	PFLCHA0_RS17965	*gacA*	c.160G>T	p.As54Tyr	D54Y	-	-	2/6	x
2	PFLCHA0_RS17965	*gacA*	c.289G>A	p.Gly97Ser	G97S	-	-	3/6	
2	PFLCHA0_RS25175	*mraZ*	c.-211A>G	NA	-211A>G	-	-	2/6	
2	PFLCHA0_RS26215	*osmY*	c.-104G>A	NA	-104G>A	-	-	1/6	
3	PFLCHA0_RS09890	*oafA*	c.1013delA	p.Lys338Ser.fsX18	K338S.fsX18	1/6	5/6	6/6	
3	PFLCHA0_RS22600	*gacS*	c.80G>A	p.Gly27Asp	G27D	-	1/6	4/6	x
3	PFLCHA0_RS31060	*nlpD*	c.590A>C	p.Gln197Pro	Q197P	-	1/6	-	
4	PFLCHA0_RS22950	*argT5*	c.120-533del	p.Lys40-Leu185del.fsX6	Δ*argT5*	2/6	-	-	
4	PFLCHA0_RS02080	*hutI*	c.786C>T	p.Phe262Phe	786C>T	1/6	-	-	
4	PFLCHA0_RS17965	*gacA*	c.-40T>A	NA	-40T>A	-	2/6	1/4	
4	PFLCHA0_RS17965	*gacA*	c.112G>T	p.E38X	E38X	-	2/6	3/4	
4	PFLCHA0_RS05510	*nudL*	c.288C>T	p.Ser96Ser	288C>T	-	1/6	-	
4	PFLCHA0_RS11820	*RS11820*	c.33C>T	p.Ala11Ala	33C>T	-	1/6	-	
4	PFLCHA0_RS08490	*flhA*	c.1176delT	p.His393Gln.fsX15	H393Q.fsX15	-	-	3/4	
4	PFLCHA0_RS11785	*RS11785*	c.766A>T	p.Ser256Cys	S256C	-	-	3/4	
4	PFLCHA0_RS27515	*rpsH*	c.-90C>G	NA	-90C>G	-	-	1/4	
5	PFLCHA0_RS18525	*RS18525*	c.254delT	p.Val85Gly.fsX40	V85G.fsX40	1/6	-	-	
5	PFLCHA0_RS06125	*rpoS*	c.193C>T	p.Gln65X	Q65X	-	1/6	4/6	
5	PFLCHA0_RS08340	*fleQ*	c.959G>A	p.Arg320Gln	R320Q	-	-	1/6	
5	PFLCHA0_RS14960	*tetR*	c.381C>G	p.Tyr127X	Y127X	-	-	2/6	
5	PFLCHA0_RS09880	*RS09880*	c.246C>A	p.Tyr82X	Y82X	-	-	1/6	
5	PFLCHA0_RS12070	*RS12070*	c.1389C>G	p.Pro463Pro	1389C>G	-	-	1/6	

159 [1]DNA sequence change positions are relative to the cDNA (c); del represents a deletion

160 [2]X represents a stop codon (at its relative position in case of a shifted frame); fs represents a frame shift; del represents a deletion

161 [3]Mutations that are selected for the analysis of their frequency trajectories using RQ-HMR analysis

Early adaptations are driven by cell surface-related genes

In general, mutations that are fixed early on in the rhizosphere adaptation process tend to have a high selective advantage (31). Disruptive mutations in *oafA*, resulting in premature stops halfway the coding region, were detected as the first acquired mutations in two independent evolutionary lines (lines 1 and 3) and appear to have swept the population in later generations (Table 2). This gene, which is part of the previously described *OBC3* gene cluster (29), encodes an O-acetyltransferase, which is postulated to acetylate the O-antigen component of the outer membrane LPS (32). Another *OBC3* mutation that accumulated early on in the rhizosphere adaptation process was *galE*^c.94G>A leading to an amino acid substitution (V32M) in galE which swept through the population in evolutionary line 2, reaching fixation in cycle 6 (Table 2). *GalE* encodes an UDP-glucose 4-epimerase which is involved in O-antigen and LPS core biosynthesis (33–35). One colony with a mutation in a third *OBC3* cluster gene, *RS09880*, encoding a putative glycosyl transferase was found in evolutionary line 5. Thus, in four out of the five evolutionary lines mutations that likely affect bacterial LPS structure appeared during rhizosphere adaptation, and in three evolutionary lines such mutations even became dominant in the population. These results strongly suggest that modifying bacterial cell surface structure is an important bacterial strategy in early adaptation to the rhizosphere.

Adaption driven by global regulators

In the present study, six mutations were detected in *gacA* in three of the five evolutionary lines, representing approximately 20% of all missense mutations. Notably, in evolutionary lines 2 and 4, multiple *gacA* alleles accumulated, some of which were detected in early plant adaptation cycles (Table 2). Additionally, a *gacS* mutation accumulated in line 3. These two genes encode the main constituents of the conserved GacA/GacS two-component regulator system, i.e. the hybrid sensor histidine kinase GacS and the cognate response regulator GacA (Figure 2). In gram-negative bacteria, activation of GacS results in cross-phosphorylation of GacA via phosphotransfer which in turn leads to activation of the expression of the small RNA genes *rsmY* and *rsmZ* via its helix-turn-helix (HTH) domain-binding domain (36). In CHA0, this regulatory pathway is known to control quorum sensing as well as secondary metabolism and stress resistance (37–40). In the closely related strain *P. protegens* Pf-5 this pathway was shown to have a big impact on bacterial genome-wide gene expression, affecting the expression of more than 10% of the annotated genes (41). Similarly, *gacA* mutants that arose on the edge of swarming Pf-5 colonies showed dramatically altered genome-wide gene expression patterns (42).

Including *gacS* and *gacA*, about half of the mutated genes in this study are global regulators or sigma factors (Table 1, Figure 1). This high frequency suggests that global regulator-controlled networks are evolvable and play a major role in rapid bacterial adaptation. Pleiotropic adaptive mutations in global regulator genes have been shown to be important for bacterial adaption both in laboratory (43, 44), and in natural settings (45, 46). Remodeling and continuous optimization of existing regulatory networks by single mutations is an important strategy for bacterial adaptation to the host (47).

Bacterial motility

Bacterial motility is an important trait for rhizosphere competence, mediating colonization of distal parts of the root system (48), and both LPS and the GacS/GacA two-component regulator system affect this trait. The O-antigenic side chain of the LPS was reported to contribute to swimming and swarming motility in the plant pathogenic bacterium *Erwinia amylovora* (49, 50). The GacS/GacA two-component regulator system controls bacterial motility, for example by affecting transcription of genes related to flagella and biosurfactant biosynthesis (42, 48, 51). We also identified several other mutations across the various evolutionary lines that can be linked to bacterial motility. For instance, a disruptive mutation in the flagellar biosynthesis protein-coding gene *FlhA* (H393Q.fsX15) that is involved in the biogenesis of the bacterial flagellum (Table 1) appeared in evolutionary line 4 reaching up to a predicted frequency of 75% (Table 2). In *P. fluorescens* Pf-5, and in *P. aeruginosa* PA01 FlhA is reported to be essential for swimming motility (41).

Furthermore, we identified one amino-acid substitution, R320Q, in FleQ and two in SadB, R183P and L258Q respectively, that based on sequence similarity to well-studied, homologous proteins in other bacteria are likely to affect motility as well. FleQ is a σ^{54}-dependent Fis family transcriptional regulator which regulates flagellar motility, biofilm formation as well as Pel exopolysaccharide production in response to cellular c-di-GMP in *P. aeruginosa* (52). *P. protegens* FleQ shares 84% sequence identity and 98% sequence coverage with *P. aeruginosa* FleQ and similarly contains an N-terminal flagellar regulatory domain FleQ (PF06490) also known as the REC receiver domain, a central AAA+/ATPase σ^{54}-interaction domain (PF00158), and a C-terminal Fis-type HTH DNA-binding domain (PF02954). The R320Q substitution we identified here is found in the AAA+/ATPase σ^{54}-interacting domain in between the arginine (R) finger (300-303) and a c-di-GMP-binding motif, ExxxR (330-334) (52). The conserved arginine residues in FleQ, including the here mutated R320, are thought to be important for protein oligomerization, and substitution of any of them abolishes ATPase activity in *Vibrio cholerae* EpsE completely (53). SadB, finally,

encodes a HDOD domain-containing protein (Table 1) which shares 77% sequence identity and 99% sequence coverage with *P. aeruginosa* SadB (PA5346). In *P. aeruginosa*, SadB stimulates Pel exopolysaccharide production and the chemotaxis-like cluster CheIV, which in turn affect flagellar motility as well as biofilm formation (54). In *Pseudomonas fluorescens* F113, SadB, together with FleQ control flagellar motility, dependent and independent of the GacS/GacA two-component regulator system (55, 56).

Parallelism of targeted mutations on this functional motility pathway impelled us to assess bacterial motility and track its dynamics across multiple evolutionary lines. Two independent evolutionary trajectories to genotypes *galE*^{V32M} · *accC*^{E413K} · *gacA*^{D54Y} and *oafA*$^{K338S.fsX18}$ · *gacS*^{G27D} from evolutionary lines 2 and 3, respectively, were studied for changes in their swimming and swarming behavior. Both genotypes accumulated mutations in *OBC3* cluster genes (29) linked to LPS biogenesis (*galE* or *oafA*) and the GacS/GacA two-component regulator system (*gacA* or *gacS*). Interestingly, *galE*^{V32M} and *oafA*$^{K338S.fsX18}$ mutants displayed opposing phenotypes when their swimming and swarming behavior was examined. Whereas the *galE*^{V32M} mutant produces significantly larger swimming and swarming haloes as compared to the wild type, the *oafA*$^{K338S.fsX18}$ mutant produces significantly smaller swimming haloes and is even completely defective in swarming (Figure 4). The *gacA* and *gacS* mutations, however, consistently reduced swimming and swarming behavior.

Dynamics of global phenotypic change

Since natural selection eventually operates at the phenotypic level, revealing bacterial global phenotypic evolutionary dynamics can help us to identify traits that are under selection. Moreover, beneficial genetic mutations can be predicted if they are linked to well-known root colonization traits. A broad range of 30 bacterial traits including different aspects of bacterial life-history traits, were measured for the sequenced isolates, allowing genotype-phenotype association analysis.

As shown in Figure 5a, the 30 bacterial traits separated into four clusters that share a similar pattern across the different mutant genotypes and this clustering is supported by model-based clustering analysis (Figure S3). Growth of the bacteria in King's B (KB) medium was positively correlated with siderophore and exoprotease production, tryptophan side oxidase activity, and growth inhibition of the bacterial plant pathogen *Ralstonia solanacearum*. Thus, this cluster, designated cluster 1, contains traits associated with bacterial social behavior, related to within population communication and cooperation, such as the

production of public goods. In a principle component analysis (Figure S4a), the first principle component

(PC1) is strongly correlated with all five traits, with a total explanation of 64.5% for all variables. Cluster 2

contains traits linked to carbon source utilization. For this cluster PC1 is strongly correlated with all carbon

source usage-related traits, with a total explanation of 83.9% for all variables (Figure S4b). A third cluster

was observed for the bacterial ability to form a biofilm, to produce indole-3-acetic acid (IAA), and to inhibit

the growth of two fungal plant pathogens, with a total explanation of 82.4% by PC1 (Figure S4c). Finally,

cluster 4 contains all seven traits that are related to bacterial resistance to biotic and abiotic stresses. The

first principle component (PC1) is strongly correlated with all seven traits, with a total explanation of

53.9% for all variables (Figure S4d). For these four clusters, the PC1 (or -PC1) value is used as a proxy to

present the general performance of all the traits that clustered together.

As is shown in Figure 5b, all five evolutionary lines showed a parallel trend of accumulative decline of

social traits. Mutations in *gacA, gacS,* and *rpoS* resulted in significantly decreased bacterial social traits,

and this was also observed for the double mutant $RS11785^{S256C} \cdot flhA^{H393Q.fsX15}$ relative to the background

$gacA^{E38X}$ mutation. *RS11785*, like *gacA*, encodes a LysR-type transcriptional regulator. In addition to these

global regulators, earlier mutations in *oafA*, *galE* and $RS17350^{A77A.fsX14}$ encoding a putative

methyltransferase, also resulted in a significant but relatively small decrease in social traits. This parallel

decline of the bacterial social traits index suggests a negative selection of bacterial social behavior,

especially the production of costly public goods such as siderophores and exoproteases.

All traits related to the utilization of 14 different carbon sources, that were selected based on their

reported presence in root exudates of Arabidopsis (57) grouped in one cluster (cluster 2; Figure 5a),

suggesting that increase or decrease of the ability to utilize carbon sources is co-regulated. Different

mutations in *gacS* and *gacA* resulted in contrasting bacterial carbon source utilization. A mutation in the

first N-terminal transmembrane domain of GacS (G27D) (Figure 2) resulted in significant enhancement of

carbon source utilization (Figure 5a, c). A similar trend was observed for mutant $gacA^{G97S}$. In contrast, the

majority of evolved genotypes, including most mutations in *gacA* and *rpoS*, showed a reduced ability to

utilize carbon sources (Figure 5a, c).

In contrast to the previous two clusters, traits in clusters 3 (Figure 5d) and 4 (Figure 5e) were more stable,

in that most genotypes behave like the ancestor. Only two *gacA* mutants, $gacA^{E38X}$ and $gacA^{Y183S}$, were

associated with a significant decline of bacterial traits associated with biocontrol, such as antifungal

activity and biofilm formation, while the $rpoS^{Q65X}$ mutation dramatically increased these traits. The double

mutation $RS11785^{S256C} \cdot flhA^{H393Q.fsX15}$, in the background of the disruptive $gacA^{E38X}$ mutation (Figure 5e),

was the only mutational event that led to a significant increase of general resistance to various

environmental stresses. *FlhA*, as previously mentioned encodes the flagellar biosynthesis protein FlhA and

is linked to bacterial motility. The LysR-type transcriptional regulator RS11785 in addition to its N-terminal

HTH DNA-binding domain (PF00126) contains a predicted C-terminal substrate binding domain (PF03466),

which is linked to aromatic compound degradation and resembles that of the type 2 periplasmic binding

proteins or PBP2s. PBP2s are responsible for binding and uptake of several substrates including

polysaccharides and several amino acids. We used HHpred (58) for protein remote homology detection

and three-dimensional structure analysis to assess the possible consequence of the S256C amino acid

substitution. This analysis identified the remote homolog OxyR from *Escherichia coli* as the best template

for modeling (E-value: 2.9e-30, Score: 166.45). Intriguingly, it was previously shown that the activity of

OxyR, which represents the master peroxide sensor in gram-negative bacteria, is regulated by

intramolecular disulfide-bond formation in the regulatory PBP2-like domain. Notably, RS11785 is unlikely

to represent the master peroxide sensor in *P. protegens* CHA0, because another CHA0 protein encoded

by *PFLCHA0_RS030065* is much more similar (89% percent pairwise sequence identity versus 29%).

Nevertheless, it is tempting to speculate that the here observed S256C substitution might influence

regulatory activity by altered intramolecular disulfide-bond formation. Lastly, RS11785 bears resemblance

(37% sequence identity, 93% sequence coverage) to the LysR-type PBP2-domain containing regulator *alsR*

from *Bacillus amyloliquefaciens*. *AlsR* regulates the activity of the acetoin operon (*alsSD*) in response to

several signals such as glucose and acetate. Acetoin (3-hydroxy-2-butanone, a precursor for 2,3-

butanediol) in turn is known as a potent elicitor of induced systemic resistance (59), and it is linked to

general plant growth promotion in *Bacilli*. Given these observations from literature, but in the absence of

singular mutations in *flhA* and *RS11785*, the molecular mechanism underlying the observed enhanced

resistance in this double mutant remains to be clarified but we hypothesize that altered regulatory activity

by RS11785 is a likely driver of the increased stress resistance observed in this mutant.

Discussion

The rhizosphere is a nutrient-rich environment for root-associated bacteria. However, to access the available nutrients bacteria must overcome various challenges, including the plant immune system, presence of competing and/or predatory microorganisms, and abiotic stresses. In the gnotobiotic binary system used in this study we tracked changes in *Pseudomonas protegens* CHA0 in the rhizosphere of Arabidopsis under reproducible and controlled conditions without interference of complex interactions with other microbes. Mutations affecting global regulators, bacterial cell surface structure, and motility accumulated in parallel in our evolutionary experiment, revealing at least three important strategies of bacterial adaptation to the rhizosphere.

Global regulators and rhizosphere adaptation

The GacS/GacA two-component regulator system controls the production of antimicrobial secondary metabolites, exoenzymes, and siderophores, but also biofilm formation, stress responses, motility, and quorum sensing (13, 38, 41, 60, 61). In the present study, mutations in the GacS/GacA two-component regulator system caused dramatic changes in several bacterial phenotypic traits, including motility, carbon source utilization, and social traits (Figures 4 and 5). The observed loss of motility does not logically fit with adaptation to the rhizosphere as motility has been reported to be an important root colonization trait in Pseudomonas (17). In line with this, the *gac* mutations that accumulated in *Pseudomonas fluorescens* F113 colonizing the rhizosphere of alfalfa showed increased motility and resulted in an enhanced competitive colonization ability (48, 62). Our results fit observations by Song et al (2016)(42) in the closely related *P. protegens* strain Pf-5. During swarming motility, *gac* mutants emerged that lack production of the surfactant orfamide. These mutants cannot swarm, but they co-swarm with orfamide producing cells (42). Thus, the lack of motility in CHA0 may also be the result of loss of surfactant production and leads to a selective advantage in the presence of surfactant producers. Several other mutations that are identified in diverse regulatory, signaling and executionary genes, *flhA*, *fleQ* and *sadB* can be linked to bacterial motility. Although no data is currently available for the *flhA* and *sadB* mutants, preliminary analysis of swarming and swimming capacity *in vitro* shows that the *fleQ* mutant has also lost its ability to move. Possibly, in our experimental system motility is less important than the ability to adhere to the root surface and the formation of a biofilm that is frequently described alongside motility as two sides of the same coin. In many bacteria, these processes in fact appear inversely correlated and

controlled by the intracellular secondary messenger cyclic dimeric-GMP (c-di-GMP) (63–65). Future experiments should reveal whether indeed such trade-off between bacterial motility and root adherence underlies our observations in this evolutionary experiment.

The decreased production of public goods, such as siderophores and exoprotease, which were observed in all independent evolutionary lines (Figure 5), could be beneficial for bacterial fitness by saving energy and primary metabolites. For example, adaptation of *Pseudomonas aeruginosa* to the human host through mutations in regulators was accompanied by loss of siderophore production, secreted proteases, and biofilm formation (66, 67).

Cell surface structures as bacterial adaptation targets

Bacterial surface components are the first line of defense against environmental stress and interplay with hosts (68–70). Lipopolysaccharide (LPS) is a central outer membrane component for gram-negative bacteria, and exhibits structural adaptability that is contributed especially by its O-antigen part (68, 70). Bacterial O-antigen structure modification plays an important role in evasion of host immunity (71), and has the potential to change host-bacteria interactions (68, 70, 72). In plant pathogenic bacteria LPS components are important virulence determinants (73, 74), that activate a variety of defense-related responses (73, 75–77). In *P. fluorescens* the O-antigen component has been implicated to induce systemic resistance in radish (78) and Arabidopsis (79).

We observed parallel mutations in genes that are involved in LPS biosynthesis and structure modification. In three out of the five evolutionary lines, the first fixed mutations were identified in *oafA* and *galE* that are annotated as O-antigen biosynthesis and structure modification related (Table 1). *OafA* encodes an O-acetyltransferase, which is postulated to modify the O-antigen by acetylation (32). The enzyme GalE, UDP-galactose 4-epimerase, is involved in the interconversion of UDP-glucose to UDP-galactose, an essential intermediate for LPS core and O-antigen structures (33–35). Inactivation of *galE* in *Porphyromonas gingivalis* resulted in shortening of the O-antigenic side chain (80). And in *Bradyrhizobium japonicum* disruption of *galE* resulted in complete lack of the O-antigen (33). Thus, it is tempting to speculate that evasion of the plant's immune response plays a role in adaptation of CHA0 to the rhizosphere of Arabidopsis.

In conclusion, the observed bacterial genetic and phenotypic adaption dynamics emphasize an important role of global regulators, motility, and cell surface structures in bacterial adaptation to its host. The parallel

369 emergence of mutations in similar genes resulting in specific fitness advantages in the rhizosphere,
370 suggest that this evolutionary process is driven by the rhizosphere environment.

Materials and Methods

Experimental setup

We set up an experimental evolution experiment with *Arabidopsis thaliana* (Arabidopsis) as host plant
and *Pseudomonas protegens* CHA0 (CHA0) as the evolving bacterial strain. CHA0 (81) is a model strain
originally isolated from roots of tobacco plants grown in soil naturally suppressive to black root rot (82).
CHA0 was chromosomally tagged with GFP and a kanamycin resistance cassette (Jousset et al 2006) to
enable easy tracking of the strain and checking for contaminations. Briefly, the ancestral bacterial
population was inoculated on Arabidopsis roots in a gnotobiotic system. After a plant growth cycle of 4
weeks, evolved bacterial populations were transferred to new plants and this cycle was repeated 8 times.
Bacterial colonies were retrieved from the evolved populations at the end of each growth cycles by
plating.

We selected six bacterial isolates from each independent evolutionary line at the end of plant growth
cycles two, four and six. Together with six isolates from the ancestor population, for a total of 94 isolates
various life-history traits (as shown below) were measured and their genomes re-sequenced using the
NextSeq-500 Illumina platform (2 x 75 bp paired-end). We used the snippy pipeline
(https://github.com/tseemann/snippy), integrating reference genome-based mapping of the Illumina
reads by BWA-MEM, variant calling by SAMtools and FreeBayes, and variant impact analysis using SnpEff
(83), to identify single nucleotide polymorphisms (SNPs) and small indels and deletions (INDELs). Larger
INDELs were identified by calculating the breath of coverage of the mapped Illumina reads on the
reference genome in a sliding window using bedtools (84). Regions with reduced coverage (<99%) were
manually inspected in the Integrative Genome Viewer (IGV). For two isolates from experimental line four
re-sequencing failed.

Bacterial life-history traits

A variety of bacterial life-history traits, reflecting various aspects of bacterial physiological processes, were
measured. We measured bacterial growth yield and resistance to various stresses, including acidic (pH =

5) and alkaline (pH = 9) conditions, oxidative stress in 0.0025% H_2O_2, water potential stress (15%

polyethylene glycol (PEG)-6000)), and salt stress (2% NaCl), and resistance to the antibiotics streptomycin

(1 $\mu g \cdot ml^{-1}$), tetracycline (1 $\mu g \cdot ml^{-1}$), and penicillin (5 $\mu g \cdot ml^{-1}$). Bacterial growth yield was quantified in

modified Ornston and Stanier (OS) minimal medium (85) supplemented with single carbon sources that

have been reported to be abundant in Arabidopsis root exudates (Chaparro et al 2013). These included

the following carbon sources; alanine, arabinose, butyrolactam, fructose, galactose, glucose, glycerol,

glycine, lactic acid, putrescine, serine, succinic acid, threonine, and valine that were added to a final

concentration of 0.5 $g \cdot l^{-1}$. Growth yield was measured as OD_{600} (SPECTROstar Nano plate reader), and

this measure was used to determine carbon source utilization ability. In addition, we measured bacterial

auxin (indole-3-acetic acid or IAA) production with a colorimetric test (86), iron-chelating ability using a

modified chrome azurol S (CAS) assay (87), proteolytic activity by the adapted assay from Smeltzer et al.

(1993), tryptophan side chain oxidase activity using a colorimetric assay (88), and biofilm formation using

a modified Chrystal Violet staining assay (89). We further assessed bacterial antimicrobial activity by

quantifying their effect on growth of the fungi *Verticillium dahliae* and *Fusarium oxysporum,* and the

bacterium *Ralstonia solanacearum*. Bacterial motility assays were performed on Cook's Cytophaga (CC)

medium containing 0.2% tryptone (90). This medium was solidified with 0.3 or 0.5% agar to determine

swimming and swarming motility, respectively (42). Bacteria were first cultured in liquid CC medium

overnight at 28 °C, and subsequently inoculated in the middle of the motility plates using toothpicks.

Plates were incubated upside-down at 28 °C for 24 h, and the colony areas were calculated using Image J

as a measure of bacterial motility.

Hierarchical and Model-based clustering of bacterial traits

Hierarchical clustering

A heatmap to illustrate the association patterns of bacterial genotypes and their measured traits was

constructed in R using the *ggplot2* package. Hierarchical clustering was performed using the Ward.D2

method, that is based on the squared dissimilarities of the set of objects being clustered (91).

Model-based clustering

We applied a model-based clustering method to reveal the best fitting structures of trait covariance

patterns. For example, some traits might be either directly or indirectly co-regulated by the same gene,

which is expected for global regulators particularly which can co-regulate thousands of genes. We used

the *mclust* package in R to run the model simulation (92). This method assumes that the input data are

distributed as a mixture of two or more clusters. The advantage of the model-based clustering method is

that it avoids heuristic assumptions and uses a soft assignment that every data point has a possibility of

falling to each cluster, which facilitates the best clustering solution. The so-called Bayesian Information

Criterion (BIC) was used to select the best model. A large BIC score indicates a better fit of the model.

This result is in line with the outcome of hierarchical clustering with Adjusted Rand Index (ARI) set as 1,

and *k* set as 4 in "ward.D2" as indicated in Figure 5a. ARI is usually used to evaluate the match degree of

a given clustering solution comparing to the model-based clustering result, with 0 reflecting a random

partition and 1 the boundary of accuracy of a certain clustering solution (92).

Genotype-phenotype association analysis

Bacterial traits within each model-predicted cluster have similar data distribution patterns and covaried

together by the definition of the clustering method. Thus, we applied a linear regression-based method,

i.e. principle component analysis (PCA), to reduce the dimensionality of data and generate a proxy for

each model predicted cluster. These proxies were later used as the x-axis values in Figure 5b-e. We applied

the package *ggbiplot* in R to generate the PCA plots and PC1 index from the normalized datasets. The

proxies were normalized for further analysis.

To examine the effect of each mutation on bacterial phenotype, a Students' *t*-test was used to compare

cluster proxies of evolved genotypes with their direct ancestors. Only genotypes identified more than

once were included in this analysis.

Relative quantification of mutant frequency using HRM profile analysis

We used High-Resolution Melting (HRM) profile analysis with integrated LunaProbes to quantify the ratio

of mutant to wild type genotypes (93–95). The probes and primers used in this study are listed in Table

S2. Primers were designed using Primer3. Probes were designed with the single nucleotide polymorphism

(SNP) located in the middle of the sequence, and the 3' end was blocked by carbon spacer C3. The primer

asymmetry was set to 2:1 (excess primer: limiting primer) in all cases. Pre-PCR was performed in a 10-μl

reaction system, with 0.25 μM excess primer, 0.125 μM limiting primer, 0.25 μM probe, 0.5 μl bacterial

sample culture (100-fold diluted saved sample, OD_{600} is about 0.01), 1X LightScanner Master Mix (BioFire

Defense). DMSO with the final concentration 5% was supplemented in all reactions to ensure the targeted

melting domains are within the detection limit of the LightScanner (Idaho Technology Inc.). Finally, MQ

water was used to supplement up to 10 µl. A 96-well black microtiter plate with white wells was used to

minimize background fluorescence. Before amplification, 25 µl mineral oil was loaded in each well to

prevent evaporation, and the plate was covered with a foil seal to prevent the degradation of fluorescent

molecules. Amplification was initiated by a holding at 95 °C for 3 min, followed by 55 cycles of

denaturation at 95 °C for 30 s, annealing at 60 °C for 30 s and extension at 72 °C for 30 s and then kept at

72 °C for 10 min. After amplification, samples were heated in a ThermalCycler (Bio-Rad) shortly to 95 °C

for 30 s to denature all double-stranded structures followed by a rapid cooling to 25 °C for 30 s to facilitate

successful hybridization between probes and the target strands. The plate was then transferred to a

LightScanner (Idaho Technology Inc.). Melting profiles of each well were collected by monitoring the

continuous loss of fluorescence with a steady increase of the temperature from 35 °C to 97 °C with a ramp

rate of 0.1 °C /s. The relative quantification was based on the negative first derivative plots using software

MATLAB. The areas of probe-target duplexes melting peaks were auto-calculated by 'AutoFit Peaks I

Residuals' function in software PeakFit (SeaSolve Software Inc.). The mutant frequency X was calculated

using the formula shown below:

474
$$X = \frac{\text{Area}_{mutant}}{\text{Area}_{mutant} + \text{Area}_{WT}}$$

To validate the HRM method, standard curves were generated by measuring mixed samples with known

proportions of mutant templates: 0%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% and 100%.

Measurements for each sample were done in triplicate. Linear regression formula of each mutant

between actual frequencies and measured frequencies were shown in Figure S1. The high R^2 values, and

nearly equal to 1 slope values of these equations, confirmed that the HRM method can accurately detect

mutants' frequency in a mixed population.

Data availability

The *P. protegens* CHA0-GFP reference genome is deposited on GenBank: RCSR00000000.1. Raw

sequencing data used in this study are deposited at the NCBI database under BioProject PRJNA473919. A

conversion table for the CHA0-GFP to CHA0 gene annotations including recent NCBI accession codes is

available here: https://doi.org/10.6084/m9.figshare.13295828.v2.

Acknowledgments

This work was supported by a China Scholarship Council fellowship (to E.L.), a postdoctoral fellowship of the Research Foundation Flanders (FWO 12B8116RN) (to R.D.J.), and a European Research Council Advanced Grant 269072 (to C.M.J.P.).

Author contributions

E.L., A.J., C.M.J.P., P.A.H.M.B., and R.D.J. designed experiments; E.L. and H.J. performed experiments; E.L. and R.D.J. analyzed data. All authors discussed the results and contributed to the final manuscript.

Figures and Figure legends

a)

b)

Figure 1. Genomic distribution of *Pseudomonas protegens* CHA0 evolutionary adaptations. **a)** Rings from inside to outside: Ring 1, nucleotide position indicator; Ring 2, green/purple GC-skew (-/+); Ring 3, %GC; Ring 4, protein-coding genes; Ring 5, distribution of identified mutations arisen during the evolutionary experiment. Functional annotation of mutated genes are indicated by color representing Cluster of orthologous groups (COGs); the key of which can be seen in panel B. **b)** Frequency of mutations per COG class, highlighting enrichment in the classes T (signal transduction), K (transcription), M (cell wall) and E (amino acid transport and metabolism).

a)

b)

Figure 2. Localization of *gac* and *OBC3* mutations identified in the evolutionary experiment. **a)** Schematic view of the cell wall of the gram-negative bacterium *Pseudomonas protegens* highlighting the presence of the lipopolysaccharide (LPS) on the outside and the presence of the two-component regulator system GacS/GacA on the inner membrane and in the cytoplasm. The arrows indicate the location of the amino acid substitutions that were identified in this study in GacS (G27D) and GacA (G38X, D49Y, D54Y, G97S and Y183S). Note the travel of a phosphate group (P) from GacS to GacA upon signal perception which is accepted by either the Asp49 or Asp54 residue in GacA which are both found mutated in two different mutants. **b)** Genomic region harboring the *OBC3* gene cluster responsible for the synthesis of long O-polysaccharide (O-PS) on the LPS and indicated mutations that were identified in a glycosyl transferase (RS09880; Y82X), the O-antigen acetyltransferase oafA (K338S.fsX18 and Y335X) and the UDP-glucose 4-epimerase (galE; V32M).

Figure 3. Phylogenetic trees of five independently evolving *P. protegens* CHA0 populations. Phylogenies for 18 genomes from each population (16 for the population in experimental line 4), based on the sequential appearance of mutations. Synonymous mutations are underlined. Circles represent unique bacterial genotypes. Intensity of the circle fill represents the time point at which a genotype was firstly detected. *OBC3* and *gac* mutants specifically are highlighted in blue and purple respectively. The frequency of each mutation in the respective population can be found in Table 2.

Figure 4. Effects of *gacS/gacA* mutations on bacterial swimming and swarming behaviour. **a)** The top panels illustrate the experimental layout of motility assays, that are pictured in the middle and bottom panels for swarming and swimming respectively. **b)** The accumulative effects of mutations on swarming (left panel) and swimming (right panel) behaviour. Colony area as a measure for bacterial motility was determined using ImageJ. The number of acquired mutations compared to the ancestor is used to express the genetic distance.

a)

Figure 5. Genotype-phenotype association analysis of evolved bacterial populations. **a)** Heatmap representing the performance of 30 life-history traits of six ancestral and 63 evolved isolates (isolates without mutations or only with synonymous mutation were excluded for this association analysis). The "ward.D2" method was used for hierarchical clustering, based on the dissimilarities of the set of objects being clustered. The four clusters depicted on top are model-predicted. Each column illustrates one bacterial life-history trait, indicated on the bottom of the figure. Each row represents an isolate, indicated by its sample ID number and mutational events. **b-e)** Principle component analysis was applied to generate a general symbolic index for each model predicted cluster. The proxy values of each cluster were normalized, higher values represent better performance. To illustrate the effects of each mutational event, ANOVA was applied to reveal the phenotypic change between each genotype and its identified nearest ancestor in each line. Colour-filled circles represent bacterial genotypes, with different colours representing independent evolving populations and the size representing the count. The dashed line represents the average value of the ancestor.

Supplementary Figures:

Figure S1. Standard curves of measured mutant versus ancestor proportion as a function of the actual proportion,

using series of mixed samples with known proportions (0%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% and

100% of mutant frequency). Relative densities of mutants $gacA^{D49Y}$, $gacA^{D54Y}$, and $gacS^{G27D}$ were measured by PCR-

based high-resolution melting (HRM) analysis. Measurements for each sample were performed in triplicate. In each

plot, the black dots represent the measurements, the blue line the fit which was generated based on linear

regression modelling.

Figure S2. Frequency trajectories of $gacA^{D49Y}$, $gacA^{D54Y}$ and $gacS^{G27D}$ mutants during long-term rhizosphere adaption.

The X-axis represents the plant-to-plant transferring cycle of the bacterial population. Mutant frequency was

determined by quantifying the ratio of mutant allele relative to wild-type allele, using PCR-based high-resolution

melting (HRM) analysis. The data shown are the mean of two technical replicates, and error bars represent the

standard deviation of the mean.

Figure S3. Selection of the best clustering model using Bayesian Information Criterion (BIC) using the *mclust* R

package. A large BIC score indicates strong evidence for the corresponding model. The VEI model with 4 components

best fits our data. VEI indicates that the clusters have variable volume, the same shape and an orientation equal to

coordinate axes.

a

b

c

d

564 **Figure S4.** Coordinate axes transformation of each model predicted cluster using principal component analysis (PCA).

565 **a.** PCA plot of social trait cluster (cluster 1); **b.** PCA plot of carbon utilization trait cluster (cluster 2); **c.** PCA plot of

566   biocontrol trait cluster (cluster 3); **d.** PCA plot of stress resistance trait cluster (cluster 4). Each dot represents one

567   isolate.

Table S1 Genotypes of whole-genome sequenced CHA0 isolates.

Sample ID	Line	Cycle	Genotype
61	1	2	*WT*
169	1	2	*WT*
13	1	2	*oafA*^{Y335X}
121	1	2	*oafA*^{Y335X}
133	1	2	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$
157	1	2	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$
158	1	4	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$
182	1	4	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$
14	1	4	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *gacA*^{D49Y}
122	1	4	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *gacA*^{D49Y}
86	1	4	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *wbpM*^{G79R}
2	1	4	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *RS21275*^{R78L}
195	1	6	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *gacA*^{D49Y}
242	1	6	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *gacA*^{D49Y}
278	1	6	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *gacA*^{D49Y}
255	1	6	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *gacA*^{D49Y} · *sadB*^{R183P}
206	1	6	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *sadB*^{L258Q}
207	1	6	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *sadB*^{L258Q} · *pvdS*$^{-33T>C}$
87	2	2	*WT*
39	2	2	*gacA*^{Y183S}
183	2	2	*gacA*^{Y183S}
123	2	2	*galE*^{V32M}
135	2	2	*galE*^{V32M}
63	2	2	*tssM*$^{1336C>T}$
172	2	4	*gacA*^{Y183S}
52	2	4	*galE*^{V32M}
16	2	4	*galE*^{V32M} · *accC*^{E413K}
28	2	4	*galE*^{V32M} · *accC*^{E413K}
100	2	4	*galE*^{V32M} · *accC*^{E413K}
160	2	4	*galE*^{V32M} · *accC*^{E413K} · *yvaQ2*$^{-9G>T}$
245	2	6	*galE*^{V32M} · *accC*^{E413K}
220	2	6	*galE*^{V32M} · *accC*^{E413K} · *gacA*^{D54Y}

197	2	6	*galE*[V32M] · *accC*[E413K] · *gacA*[D54Y] · *osmY*[-104G>A]
221	2	6	*galE*[V32M] · *accC*[E413K] · *gacA*[G97S]
196	2	6	*galE*[V32M] · *accC*[E413K] · *gacA*[G97S] · *mraZ*[-211A>G]
268	2	6	*galE*[V32M] · *accC*[E413K] · *gacA*[G97S] · *mraZ*[-211A>G]
29	3	2	*WT*
65	3	2	*WT*
53	3	2	*WT*
77	3	2	*WT*
161	3	2	*WT*
137	3	2	*oafA*[K338S.fsX18]
90	3	4	*nlpD*[Q197P]
18	3	4	*oafA*[K338S.fsX18]
66	3	4	*oafA*[K338S.fsX18]
150	3	4	*oafA*[K338S.fsX18]
186	3	4	*oafA*[K338S.fsX18]
30	3	4	*oafA*[K338S.fsX18] · *gacS*[G27D]
235	3	6	*oafA*[K338S.fsX18]
270	3	6	*oafA*[K338S.fsX18]
199	3	6	*oafA*[K338S.fsX18] · *gacS*[G27D]
222	3	6	*oafA*[K338S.fsX18] · *gacS*[G27D]
234	3	6	*oafA*[K338S.fsX18] · *gacS*[G27D]
198	3	6	*oafA*[K338S.fsX18] · *gacS*[G27D]
57	4	2	*WT*
105	4	2	*WT*
81	4	2	*WT*
129	4	2	*ΔargT5*
141	4	2	*ΔargT5*
21	4	2	*hult*[786C>T]
80	4	4	*WT*
152	4	4	*WT*
188	4	4	*gacA*[-40T>A]
68	4	4	*gacA*[-40T>A] · *RS11820*[33C>T]
164	4	4	*gacA*[E38X]
20	4	4	*gacA*[E38X] · *nudL*[288C>T]
225	4	6	*gacA*[-40T>A] · *rpsH*[-90C>G]
212	4	6	*gacA*[E38X] · *RS11785*[S256C] · *flhA*[H393Q.fsX15]

237	4	6	*gacA*^{E38X} · *RS11785*^{S256C} · *flhA*$^{H393Q.fsX15}$
260	4	6	*gacA*^{E38X} · *RS11785*^{S256C} · *flhA*$^{H393Q.fsX15}$
11	6	2	*WT*
71	6	2	*WT*
191	6	2	*WT*
35	6	2	*WT*
83	6	2	*WT*
143	6	2	*RS18525*$^{V85G.fsX40}$
70	6	4	*WT*
106	6	4	*WT*
142	6	4	*WT*
58	6	4	*WT*
190	6	4	*WT*
34	6	4	*rpoS*^{Q65X}
251	6	6	*fleQ*^{R320Q}
262	6	6	*rpoS*^{Q65X}
263	6	6	*rpoS*^{Q65X}
203	6	6	*rpoS*^{Q65X} · *tetR*^{Y127X}
239	6	6	*rpoS*^{Q65X} · *tetR*^{Y127X}
226	6	6	*RS09880*^{Y82X} · *RS12070*$^{1389C>G}$
7	ancestor	0	*WT*
31	ancestor	0	*WT*
79	ancestor	0	*WT*
127	ancestor	0	*WT*
163	ancestor	0	*WT*
187	ancestor	0	*WT*

569 Note: X, represents a stop codon (at its relative position in case of a shifted frame); fs, frame shift; del, deletion; DNA sequence
570 change positions are relative to the cDNA.

Table S2 Primers and probes used for high-resolution melting (HRM) analysis. For the two *gacA* mutants the same set of primers was used. Underlined bases indicate the position of the single nucleotide point (SNP) mutations within the probe sequences. ΔTm (℃) indicates the melting temperature difference between WT-probe duplex and mutant-probe duplex.

Target gene	Strain ID	SNP locus	Forward primer (excess)	Reverse primer (limiting)	Amplicon size	Probe sequence	Probe length	Target strand	Perfect match/ mismatch	ₐTm (℃)
gacA[D49Y]	242	145G>T				5'-CATCAGGACCACATCGGGCTTCAGCTCCCG-/C3/3'	30nt	WT	G::C / T::C	5.31
gacA[D54Y]	220	160G>T	5'-ATCGATGGCCTGCAAGTAGT-3'	5'-CGGGTAGGAAAGGGATCTTC-3'	206 bp	5'-TGGCATCTTGACGTCCATCAGGACCACATC-/C3/3'	30nt	WT	G::C / T::C	4.92
gacS[G27D]	222	80G>A	5'-GCGTACTGTTGCTGACCTTG-3'	5'-AGCATCTGGGTGTTGTGGTT-3'	178bp	5'-AGGTGAAGTAGCCGCCCAGCACCAAAGCCA-/C3/3'	30nt	WT	G::C / A::C	4.83

References

1. Berendsen RL, Pieterse CMJ, Bakker PAHM. 2012. The rhizosphere microbiome and plant health. Trends Plant Sci 17:478–486.

2. Bais HP, Weir TL, Perry LG, Gilroy S, Vivanco JM. 2006. The role of root exudates in rhizosphere interations with plants and other organisms. Annu Rev Plant Biol 57:233–266.

3. Stringlis IA, Yu K, Feussner K, de Jonge R, Van Bentum S, Van Verk MC, Berendsen RL, Bakker PAHM, Feussner I, Pieterse CMJ. 2018. MYB72-dependent coumarin exudation shapes root microbiome assembly to promote plant health. Proc Natl Acad Sci U S A 115:E5213–E5222.

4. Bakker PAHM, Pieterse CMJ, de Jonge R, Berendsen RL. 2018. The soil-borne legacy. Cell 172:1178–1180.

5. Philippot L, Raaijmakers JM, Lemanceau P, van der Putten WH. 2013. Going back to the roots: the microbial ecology of the rhizosphere. Nat Rev Microbiol 11:789–799.

6. Sasse J, Martinoia E, Northen T. 2018. Feed your friends: do plant exudates shape the root microbiome? Trends Plant Sci 23:25–41.

7. Bais HP, Park SW, Weir TL, Callaway RM, Vivanco JM. 2004. How plants communicate using the underground information superhighway. Trends Plant Sci 9:26–32.

8. Zamioudis C, Pieterse CMJ. 2012. Modulation of host immunity by beneficial microbes. Mol Plant-Microbe Interact 25:139–150.

9. Yang J, Kloepper JW, Ryu C-M. 2009. Rhizosphere bacteria help plants tolerate abiotic stress. Trends Plant Sci 14:1–4.

10. Browne P, Rice O, Miller SH, Burke J, Dowling DN, Morrissey JP, O'Gara F. 2009. Superior inorganic phosphate solubilization is linked to phylogeny within the *Pseudomonas fluorescens* complex. Appl Soil Ecol 43:131–138.

11. Lugtenberg B, Kamilova F. 2009. Plant growth promoting rhizobacteria. Annu Rev Microbiol 63:541–556.

12. Ryu C-M, Farag MA, Hu C-H, Reddy MS, Wei H-X, Paré PW, Kloepper JW. 2003. Bacterial volatiles promote growth in *Arabidopsis*. Proc Natl Acad Sci U S A 100:4927–4932.

13. Haas D, Défago G. 2005. Biological control of soil-borne pathogens by fluorescent pseudomonads. Nat Rev Microbiol 3:307–319.

14. Pieterse CMJ, Zamioudis C, Berendsen RL, Weller DM, Van Wees SCM, Bakker PAHM. 2014. Induced systemic resistance by beneficial microbes. Annu Rev Phytopathol 52:347–375.

15. Raaijmakers JM, Bonsall RF, Weller DM. 1999. Effect of population density of *Pseudomonas fluorescens* on production of 2,4-Diacetylphloroglucinol in the rhizosphere of wheat. Phytopathology 89:470–475.

16. Raaijmakers JM, Leeman M, Oorschot van MM., et.al. 1995. Dose-response relationships in biological control of Fusarium wilt of radish by *Pseudomonas* spp.. Phytopathology 85:1075–1081.

17. Lugtenberg BJJ, Dekkers L, Bloemberg G V. 2001. Molecular determinants of rhizosphere colonization *Pseudomonas*. Annu Rev Phytopathol 39:461–490.

18. de Weger LA, van der Vlugt CI, Wijfjes AH, Bakker PA, Schippers B, Lugtenberg B. 1987. Flagella of a plant-growth-stimulating *Pseudomonas fluorescens* strain are required for colonization of potato roots. J Bacteriol 169:2769–2773.

19. de Weger LA, van Loosdrecht MC, Klaassen HE, Lugtenberg B. 1989. Mutational changes in physiochemical cell surface properties of plant-growth-stimulating *Pseudomonas* spp. do not influence the attachment properties of the cells. J Bacteriol 171:2756–2761.

20. Simons M, Permentier HP, de Weger LA, Wijffelman CA, Lugtenberg BJJ. 1997. Amino acid synthesis is necessary for tomato root colonization by *Pseudomonas fluorescens* strain WCS365. Mol Plant-Microbe Interact 10:102–106.

21. Berendsen RL, Vismans G, Yu K, Song Y, de Jonge R, Burgman WP, Burmølle M, Herschend J, Bakker PAHM, Pieterse CMJ. 2018. Disease-induced assemblage of a plant-beneficial bacterial consortium. ISME J 12:1496–1507.

22. Yu K, Liu Y, Tichelaar R, Savant N, Lagendijk E, van Kuijk SJL, Stringlis IA, van Dijken AJH, Pieterse CMJ, Bakker PAHM, Haney CH, Berendsen RL. 2019. Rhizosphere-associated *Pseudomonas* suppress local root immune responses by gluconic acid-mediated lowering of environmental pH. Curr Biol 29:3913-3920.e4.

23. Cole BJ, Feltcher ME, Waters RJ, Wetmore KM, Mucyn TS, Ryan EM, Wang G, Ul-Hasan S, McDonald M, Yoshikuni Y, Malmstrom RR, Deutschbauer AM, Dangl JL, Visel A. 2017. Genome-wide identification of

bacterial plant colonization genes. PLoS Biol 15:e2002860.

24. Kawecki TJ, Lenski RE, Ebert D, Hollis B, Olivieri I, Whitlock MC. 2012. Experimental evolution. Trends Ecol
Evol 27:547–560.

25. Jousset A, Schuldes J, Keel C, Maurhofer M, Daniel R, Scheu S, Thuermer A. 2014. Full-genome sequence of
the plant growth-promoting bacterium *Pseudomonas protegens* CHA0. Genome Announc 2:e00322-14.

26. Li E, de Jonge R, Chen L, Friman V, Bakker P, Pieterse C, Jousset A. 2020. Rapid evolution of plant-bacterium
mutualism in the rhizosphere. SSRN Electron J https://doi.org/10.2139/ssrn.3650572.

27. Oren Y, Smith MB, Johns NI, Kaplan Zeevi M, Biran D, Ron EZ, Corander J, Wang HH, Alm EJ, Pupko T. 2014.
Transfer of noncoding DNA drives regulatory rewiring in bacteria. Proc Natl Acad Sci U S A 111:16112–
16117.

28. Carroll SB. 2008. Evo-devo and an expanding evolutionary synthesis: a genetic theory of morphological
evolution. Cell 134:25–36.

29. Kupferschmied P, Chai T, Flury P, Blom J, Smits THM, Maurhofer M, Keel C. 2016. Specific surface glycan
decorations enable antimicrobial peptide resistance in plant-beneficial *pseudomonads* with insect-
pathogenic properties. Environ Microbiol 18:4265–4281.

30. Galperin MY, Makarova KS, Wolf YI, Koonin E V. 2015. Expanded Microbial genome coverage and improved
protein family annotation in the COG database. Nucleic Acids Res 43:D261–D269.

31. Barrett RDH, Maclean RC, Bell G. 2006. Mutations of intermediate effect are responsible for adaptation in
evolving *Pseudomonas fluorescens* populations. Biol Lett 2:236–238.

32. Allison GE, Verma NK. 2000. Serotype-converting bacteriophages and O-antigen modification in *Shigella
flexneri*. Trends Microbiol 8:17–23.

33. Chang W-S, Park K-M, Koh S-C, So J-S. 2008. Characterization of the *Bradyrhizobium japonicum galE* gene:
its impact on lipopolysaccharide profile and nodulation of soybean. FEMS Microbiol Lett 280:242–249.

34. Liu B, Knirel YA, Feng L, Perepelov A V., Senchenkova SN, Wang Q, Reeves PR, Wang L. 2008. Structure and
genetics of *Shigella* O antigens. FEMS Microbiol Rev 32:627–653.

35. Samuel G, Reeves P. 2003. Biosynthesis of O-antigens: genes and pathways involved in nucleotide sugar
precursor synthesis and O-antigen assembly. Carbohydr Res 338:2503–2519.

36. Brencic A, McFarland KA, McManus HR, Castang S, Mogno I, Dove SL, Lory S. 2009. The GacS/GacA signal
transduction system of *Pseudomonas aeruginosa* acts exclusively through its control over the transcription
of the RsmY and RsmZ regulatory small RNAs. Mol Microbiol 73:434–445.

37. Brown D. 2010. A mathematical model of the Gac/Rsm quorum sensing network in *Pseudomonas
fluorescens*. BioSystems 101:200–212.

38. Heeb S, Blumer C, Haas D. 2002. Regulatory RNA as mediator in GacA/RsmA-dependent global control of
exoproduct formation in *Pseudomonas fluorescens* CHA0. J Bacteriol 184:1046–1056.

39. Valverde C, Heeb S, Keel C, Haas D. 2003. RsmY, a small regulatory RNA , is required in concert with RsmZ
for GacA-dependent expression of biocontrol traits in *Pseudomonas fluorescens* CHA0. Mol Microbiol
50:1361–1379.

40. Heeb S, Valverde C, Haas D. 2005. Role of the stress sigma factor RpoS in GacA / RsmA-controlled
secondary metabolism and resistance to oxidative stress in *Pseudomonas fluorescens* CHA0. FEMS
Microbiol Lett 243:251–258.

41. Hassan KA, Johnson A, Shaffer BT, Ren Q, Kidarsa TA, Elbourne LDH, Hartney S, Duboy R, Goebel NC,
Zabriskie TM, Paulsen IT, Loper JE. 2010. Inactivation of the GacA response regulator in *Pseudomonas
fluorescens* Pf-5 has far-reaching transcriptomic consequences. Environ Microbiol 12:899–915.

42. Song C, Kidarsa TA, Mortel JE Van De, Loper JE, Jos M. 2016. Living on the edge: emergence of spontaneous
gac mutations in *Pseudomonas protegens* during swarming motility. Environ Microbiol 18:3453–3465.

43. Hunter P. 2008. The great leap forward. EMBO Rep 9:608–611.

44. Zambrano MM, Siegele DA, Almirón M, Tormo A, Kolter R. 1993. Microbial competition: *Escherichia coli
* mutants that take over stationary phase cultures. Science 259:1757–1760.

45. Sumby P, Whitney AR, Graviss EA, DeLeo FR, Musser JM. 2006. Genome-wide analysis of group A
streptococci reveals a mutation that modulates global phenotype and disease specificity. PLoS Pathog 2:e5.

46. Giraud A, Arous S, De Paepe M, Gaboriau-Routhiau V, Bambou JC, Rakotobe S, Lindner AB, Taddei F, Cerf-
Bensussan N. 2008. Dissecting the genetic components of adaptation of *Escherichia coli* to the mouse gut.
PLoS Genet 4:0052–0061.

47. Damkiaer S, Yang L, Molin S, Jelsbak L. 2013. Evolutionary remodeling of global regulatory networks during
long-term bacterial adaptation to human hosts. Proc Natl Acad Sci U S A 110:7766–7771.
48. Sánchez-Contreras M, Martín M, Villacieros M, O'Gara F, Bonilla I, Rivilla R. 2002. Phenotypic selection and
phase variation occur during alfalfa root colonization by *Pseudomonas fluorescens* F113. J Bacteriol
184:1587–1596.
49. Bowden SD, Hale N, Chung JCS, Hodgkinson JT, Spring DR, Welch M. 2013. Surface swarming motility by
Pectobacterium atrosepticum is a latent phenotype that requires O antigen and is regulated by quorum
sensing. Microbiology 159:2375–2385.
50. Berry MC, McGhee GC, Zhao Y, Sundin GW. 2009. Effect of a *waaL* mutation on lipopolysaccharide
composition, oxidative stress survival, and virulence in *Erwinia amylovora*. FEMS Microbiol Lett 291:80–87.
51. Lapouge K, Schubert M, Allain FH-T, Haas D. 2007. Gac/Rsm signal transduction pathway of γ-
proteobacteria: from RNA recognition to regulation of social behaviour. Mol Microbiol 67:241–253.
52. Matsuyama BY, Krasteva P V., Baraquet C, Harwood CS, Sondermann H, Navarro MVAS. 2016. Mechanistic
insights into c-di-GMP-dependent control of the biofilm regulator FleQ from *Pseudomonas aeruginosa*.
Proc Natl Acad Sci U S A 113:E209–E218.
53. Patrick M, Korotkov K V., Hol WGJ, Sandkvist M. 2011. Oligomerization of EpsE coordinates residues from
multiple subunits to facilitate ATPase activity. J Biol Chem 286:10378–10386.
54. Caiazza NC, O'Toole GA. 2004. SadB is required for the transition from reversible to irreversible attachment
during biofilm formation by *Pseudomonas aeruginosa* PA14. J Bacteriol 186:4476–4485.
55. Muriel C, Blanco-Romero E, Trampari E, Arrebola E, Durán D, Redondo-Nieto M, Malone JG, Martín M,
Rivilla R. 2019. The diguanylate cyclase AdrA regulates flagellar biosynthesis in *Pseudomonas fluorescens*
F113 through SadB. Sci Rep 9:1–9.
56. Martínez-Granero F, Navazo A, Barahona E, Redondo-Nieto M, Rivilla R, Martín M. 2012. The Gac-Rsm and
SadB signal transduction pathways converge on AlgU to downregulate motility in *Pseudomonas*
fluorescens. PLoS One 7:e31765.
57. Chaparro JM, Badri D V., Bakker MG, Sugiyama A, Manter DK, Vivanco JM. 2013. Root exudation of
phytochemicals in *Arabidopsis* follows specific patterns that are developmentally programmed and
correlate with soil microbial functions. PLoS One 8:e55731.
58. Zimmermann L, Stephens A, Nam SZ, Rau D, Kübler J, Lozajic M, Gabler F, Söding J, Lupas AN, Alva V. 2018.
A completely reimplemented MPI bioinformatics toolkit with a new HHpred server at its core. J Mol Biol
430:2237–2243.
59. Rudrappa T, Biedrzycki ML, Kunjeti SG, Donofrio NM, Czymmek KJ, Paré PW, Bais HP. 2010. The
rhizobacterial elicitor acetoin induces systemic resistance in *Arabidopsis thaliana*. Commun Integr Biol
3:130–138.
60. Haas D, Keel C. 2003. Regulation of antibiotic production in root-colonizing *Pseudomonas* spp. and
relavance for biological control of plant disease. Annu Rev Phytopathol 41:117–153.
61. Raaijmakers JM, De Bruijn I, Nybroe O, Ongena M. 2010. Natural functions of lipopeptides from *Bacillus*
and *Pseudomonas* : more than surfactants and antibiotics. FEMS Microbiol Rev 34:1037–1062.
62. Martínez-Granero F, Rivilla R, Martín M. 2006. Rhizosphere selection of highly motile phenotypic variants
of *Pseudomonas fluorescens* with enhanced competitive colonization ability. Appl Environ Microbiol
61:1946–1952.
63. Romling U, Galperin MY, Gomelsky M. 2013. Cyclic di-GMP: the first 25 years of a universal bacterial
second messenger. Microbiol Mol Biol Rev 77:1–52.
64. Caiazza NC, Merritt JH, Brothers KM, O'Toole GA. 2007. Inverse regulation of biofilm formation and
swarming motility by *Pseudomonas aeruginosa* PA14. J Bacteriol 189:3603–3612.
65. Simm R, Morr M, Kader A, Nimtz M, Römling U. 2004. GGDEF and EAL domains inversely regulate cyclic di-
GMP levels and transition from sessibility to motility. Mol Microbiol 53:1123–1134.
66. Smith EE, Buckley DG, Wu Z, Saenphimmachak C, Hoffman LR, D'Argenio DA, Miller SI, Ramsey BW, Speert
DP, Moskowitz SM, Burns JL, Kaul R, Olson M V. 2006. Genetic adaptation by *Pseudomonas aeruginosa* to
the airways of cystic fibrosis patients. Proc Natl Acad Sci U S A 103:8487–92.
67. Winstanley C, O'Brien S, Brockhurst MA. 2016. *Pseudomonas aeruginosa* evolutionary adaptation and
diversification in cystic fibrosis chronic lung infections. Trends Microbiol 24:327–337.
68. Lerouge I, Vanderleyden J. 2002. O-antigen structural variation: mechanisms and possible roles in

734   animal/plant–microbe interactions. FEMS Microbiol Rev 26:17–47.

735   69.   Whitfield C, Trent MS. 2014. Biosynthesis and export of bacterial lipopolysaccharides. Annu Rev Biochem
736   83:99–128.

737   70.   Maldonado RF, Sá-Correia I, Valvano MA. 2016. Lipopolysaccharide modification in Gram-negative bacteria
738   during chronic infection. FEMS Microbiol Rev 40:480–493.

739   71.   Murray GL, Attridge SR, Morona R. 2006. Altering the length of the lipopolysaccharide O antigen has an
740   impact on the interaction of *Salmonella enterica* serovar Typhimurium with macrophages and
741   complement. J Bacteriol 188:2735–2739.

742   72.   Ranf S. 2016. Immune sensing of lipopolysaccharide in plants and animals: same but different. PLoS Pathog
743   12:e1005596.

744   73.   Newman M-A, von Roepenack E, Daniels M, Dow M. 2000. Lipopolysaccharides and plant responses to
745   phytopathogenic bacteria. Mol Plant Pathol 1:25–31.

746   74.   Di Lorenzo F, Silipo A, Andersen Gersby LB, Palmigiano A, Lanzetta R, Garozzo D, Boyer C, Pruvost O,
747   Newman MA, Molinaro A. 2017. *Xanthomonas citri* pv. *citri* pathotypes: LPS structure and function as
748   microbe-associated molecular patterns. ChemBioChem 18:772–781.

749   75.   Newman MA, Daniels MJ, Dow JM. 1995. Lipopolysaccharide from *Xanthomonas campestris* induces
750   defense-related gene expression in *Brassica campestris*. Mol Plant-Microbe Interact 8:778–80.

751   76.   Barton-Willis PA, Wang MC, Holliday MJ, Long MR, Keen NT. 1984. Purification and composition of
752   lipopolysaccharides from *Pseudomonas syringae*. pv. *glycinea*. Physiol Plant Pathol 25:387–398.

753   77.   Erbs G, Newman M-A. 2003. The role of lipopolysaccharides in induction of plant defence responses. Mol
754   Plant Pathol 4:421–425.

755   78.   Leeman M, Pelt JA van, Ouden FM den, Heinsbroek M, Bakker PAHM, Schippers B. 1995. Induction of
756   systemic resistance against *Fusarium wilt* of radish by lipopolysaccharides of *Pseudomonas fluorescens*.
757   Phytopathol 85:1021–1027.

758   79.   Van Wees SCM, Pieterse CMJ, Trijssenaar A, Van 't Westende YAM, Hartog F, Van Loon LC. 1997.
759   Differential induction of systemic resistance in *Arabidopsis* by biocontrol bacteria. Mol Plant-Microbe
760   Interact 10:716–724.

761   80.   Nakao R, Senpuku H, Watanabe H. 2006. *Porphyromonas gingivalis galE* is involved in lipopolysaccharide
762   O-antigen synthesis and biofilm formation. Infect Immun 74:6145–53.

763   81.   Ramette A, Frapolli M, Saux MF-L, Gruffaz C, Meyer J-M, Défago G, Sutra L, Moënne-Loccoz Y. 2011.
764   *Pseudomonas protegens* sp. nov., widespread plant-protecting bacteria producing the biocontrol
765   compounds 2,4-diacetylphloroglucinol and pyoluteorin. Syst Appl Microbiol 34:180–188.

766   82.   Stutz EW, Défago G, Kern H. 1986. Naturally occurring fluorescent pseudomonads involved in suppression
767   of black root rot of tobacco. Phytopathology 76:181–185.

768   83.   Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012. A program for
769   annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of
770   *Drosophila melanogaster* strain *w1118; iso-2; iso-3*. Fly (Austin) 6:80–92.

771   84.   Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features.
772   Bioinformatics 26:841–842.

773   85.   Højberg O, Schnider U, Winteler H V, Sørensen J, Haas D. 1999. Oxygen-sensing reporter strain of
774   *Pseudomonas fluorescens* for monitoring the distribution of low-oxygen habitats in soil. Appl Environ
775   Microbiol 65:4085–4093.

776   86.   Glickmann E, Dessaux Y. 1995. A critical examination of the specificity of the salkowski reagent for indolic
777   compounds produced by phytopathogenic bacteria. Appl Environ Microbiol 61:793–796.

778   87.   Alexander DB, Zuberer DA. 1991. Use of chrome azurol S reagents to evaluate siderophore production by
779   rhizosphere bacteria. Biol Fertil Soils 12:39–45.

780   88.   Oberhänsli T, Défago G, Haas D. 1991. Indole-3-acetic acid (IAA) synthesis in the biocontrol strain CHA0 of
781   *Pseudomonas fluorescens*: role of tryptophan side chain oxidase. Microbiology 137:2273–2279.

782   89.   Moskowitz SM, Foster JM, Emerson J, Burns JL. 2004. Clinically feasible biofilm susceptibility assay for
783   isolates of *Pseudomonas aeruginosa* from patients with cystic fibrosis. J Clin Microbiol 42:1915–1922.

784   90.   Christensen PJ, Cook FD. 1972. The isolation and enumeration of cytophagas. Can J Microbiol 18:1933–
785   1940.

91.   Murtagh F, Legendre P. 2014. Ward's hierarchical agglomerative clustering method: which algorithms

implement ward's criterion? J Classif 31:274–295.
92.   Scrucca L, Fop M, Murphy TB, Raftery AE. 2016. Mclust 5: clustering, classification and density estimation
using gaussian finite mixture models. R J 8:289–317.
93.   Kai Lee H, Kiat Lee C, Ping Loh T, Wei-Tze Tang J, Anantharajah Tambyah P, Siew-Chuan Koay E. 2011. High-
resolution melting approach to efficient identification and quantification of H275Y mutant influenza
H1N1/2009 virus in mixed-virus-population samples. J Clin Microbiol 49:3555–3559.
94.   Zhong W-L, Wang L, Wu X, Zhang J, Chen X-F, Zhang W, Dou X, Yu B. 2016. Development of unlabeled
probe based high-resolution melting analysis for detection of filaggrin gene mutation c.3321delA. J Clin Lab
Anal 30:892–896.
95.   Capper RL, Jin YK, Lundgren PB, Peplow LM, Matz M V., van Oppen MJH. 2015. Quantitative high resolution
melting: two methods to determine SNP allele frequencies from pooled samples. BMC Genet 16:62.
798