Experimental evolution-driven identification of Arabidopsis rhizosphere competence genes in *Pseudomonas protegens*

competence genes in *Pseudomonas protegens*

Erqin Li[1,#], Henan Jiang[1], Corné M.J. Pieterse[1], Alexandre Jousset[2], Peter A.H.M. Bakker[1], Ronnie de

Jonge[1]

[1]Plant-Microbe Interactions, Department of Biology, Science4Life, Utrecht University, Padualaan 8, 3584
CH, Utrecht, The Netherlands

[2]Ecology and Biodiversity, Department of Biology, Science4Life, Utrecht University, Padualaan 8, 3584
CH, Utrecht, The Netherlands.

[†]Corresponding author: Ronnie de Jonge; r.dejonge@uu.nl

#Current address: Freie Universität Berlin, Institut für Biologie, Dahlem Center of Plant Sciences, Plant
Ecology, Altensteinstr. 6, D-14195, Berlin, Germany, erqinli22@gmail.com

Abstract

Beneficial plant root-associated microorganisms carry out a range of functions that are essential for plant performance. Establishment of a bacterium on plant roots, however, requires overcoming several challenges, including the ability to outcompete neighboring microorganisms and suppression of plant immunity. Forward and reverse genetics approaches have led to the identification of diverse mechanisms that are used by beneficial microorganisms to overcome these challenges such as the production of iron-chelating compounds, biofilm formation, or downregulation of plant immunity. However, how such mechanisms have developed from an evolutionary perspective is much less understood. In an attempt to study bacterial adaptation in the rhizosphere, we employed an experimental evolution approach to track the physiological and genetic dynamics of root-dwelling *Pseudomonas protegens* CHA0 in the *Arabidopsis thaliana* rhizosphere under axenic conditions. This simplified binary one plant, and one bacterium system allows for the amplification of key adaptive mechanisms for bacterial rhizosphere colonization. We found that mutations in global regulators, as well as in genes for siderophore production, cell surface decoration, attachment, and motility accumulated in parallel in our evolutionary experiment, underlining several different strategies of bacterial adaptation to the rhizosphere. In total we identified 35 mutations, including single-nucleotide polymorphisms, smaller indels and larger deletions, distributed over 28 genes in total. Altogether these results underscore the strength of experimental evolution to identify key genes and pathways for bacterial rhizosphere colonization, as well as highlighting a methodology for the development of elite beneficial microorganisms with enhanced root-colonizing capacities that can support sustainable agriculture in the future.

Introduction

Plants are associated with complex microbial communities assembled into a functional microbiome that safeguards optimal plant performance under harsh environmental conditions (1). The rhizosphere microbiome is a particularly interesting hotspot of plant-microbe interactions. Plants deposit up to 44% of their photosynthetically fixed carbon into the rhizosphere, fueling a specific microbial community (2). The microbial species pool in the bulk soil is the source from which the root microbiome is recruited, and plant genotype, immune responses, and environmental factors are postulated to affect this process (3–6). The establishment of beneficial microbial associations requires a high degree of coordination of both plant and microbial responses by means of a continuous molecular dialogue (7, 8). Plant-associated microorganisms can improve plant yield by protecting the plant from abiotic stresses (9), improving plant nutrition and growth (10–12), antagonizing soil-borne pathogens (13), or stimulating plant immunity (14). To exert their beneficial effects on plant performance, bacteria must colonize the roots efficiently and establish significant populations. For example, population densities above 10^5 cells per gram of root are required for efficient suppression of soil-borne plant pathogens by *Pseudomonas* spp. (15, 16). Therefore, bacterial adaptation to the plant root environment may be essential for the successful implementation of microbiome services in agriculture in order to support plant health.

Among all root-dwelling organisms, fluorescent *Pseudomonas* spp. are well characterized in terms of the traits required for their growth in the rhizosphere and for the establishment of beneficial plant-microbe interactions (17). To study bacterial traits involved in efficient root colonization, mutants defective in specific traits suspected to be involved in colonization, are compared to the ability of the parental strain to colonize roots. Such studies have highlighted a range of bacterial traits involved in efficient root colonization, including flagella (18), surface lipopolysaccharides (19), and amino acid synthesis (20). Using random mutagenesis and by determining the fitness of each mutant in competition with its parental strain in the rhizosphere, many other traits important for rhizosphere competence in *Pseudomonas* were discovered (17). Recently, random mutagenesis in *Pseudomonas capeferrum* WCS358 led to the identification of two genes important for gluconic acid biosynthesis which in turn is essential for the suppression of local, flagellin-induced root immune responses (21, 22). Such suppression was shown to be important for rhizosphere competence as gluconic acid-deficient mutants maintain reduced rhizospheric populations (22). In another recent study, genome-wide saturation mutagenesis of *Pseudomonas simiae* WCS417r revealed that 2% of the protein-coding genes are important for successful root colonization (23). Mutations that negatively affect rhizosphere

competence, and mutations that confer a competitive advantage were identified in this study. The
observation that mutations can lead to increased root colonization (23), suggests that there is room for
improvement of bacterial fitness in the rhizosphere.

In the present study we used an experimental evolution approach (24) to study how *Pseudomonas*
protegens CHA0 (CHA0) evolves during repeated colonization of the rhizosphere of the model plant
Arabidopsis thaliana (Arabidopsis). The model biological control agent CHA0 displays broad-spectrum
antagonistic activity against several plant pathogenic fungi and bacteria (11), and its complete genome
sequence is available (25). We performed highly controlled experimental evolution in a gnotobiotic and
carbon-free sand system in which bacteria depend solely on the plant for supply of carbon. Following
inoculation and establishment on the roots, bacterial populations were transferred to new plants, and
this cycle was repeated eight times. We hypothesized that the repeated colonization and establishment
of the bacterial population on the plant root would create an environment in which selection pressure
drives the accumulation of better colonizers. *In vitro* characterization of individual bacterial colonies
from these populations combined with sequencing analysis led to the identification of several
evolutionary trajectories involving 35 distinct mutations that impact social traits representing inter-
population communication and cooperation, carbon source utilization, motility, or biocontrol activity. By
combining experimental evolution with whole genome re-sequencing, we created a powerful screen for
the identification of adaptive mutations with positive effects on rhizosphere colonization.

Results

Mutational events in independent evolutionary lines

We previously studied five experimental evolutionary populations, referred to as lines, of CHA0 evolving
in the rhizosphere of Arabidopsis in a gnotobiotic system. Independent populations were introduced on
the roots and after four weeks of plant growth the populations were transferred to new plants. This
cycle of transferring was repeated eight times (26). To study adaptation at the individual level, here, we
isolated in total 90 bacterial colonies at the end of plant growth cycles 2, 4 and 6. These colonies, and six
colonies from the ancestral population that was initially introduced on the roots, were re-sequenced to
an average depth of 25-fold coverage (minimum 10, maximum 70) and used for the identification of
single nucleotide polymorphisms (SNPs), as well as small and large insertions or deletions (INDELs). The

six ancestral colonies were all identical, indicating that there was no genetic variation in the starting

population and all observed mutations are therefore *de novo* mutations. In total, one or more mutations

were detected in 64 evolved colonies, which collectively represent 71% of the 90 characterized bacterial

colonies (Table S1). We identified 5 synonymous substitutions, 20 nonsynonymous substitutions, and 4

deletions ranging in length from 1 base pair (bp) to about 400 bp, distributed over 22 genes, and 6

additional mutational events located in intergenic regions (Table 1). Mutations located in intergenic

regions possibly affect transcription of nearby genes via affecting their regulatory proteins binding sites

and subsequent changes in their promoter activity (27, 28). Several mutations were found to be

clustered in select genes and/or regions on the CHA0 genome, e.g. those targeting the response

regulator *gacA* (PFLCHA0_RS17965; *NC_021237:4,039,113-4,039,754*) or the *OBC3* gene cluster involved

in lipopolysaccharide (LPS) biosynthesis (*NC_021237:2,173,707-2,196,028*)(29), but the majority of the

mutations are spread across the 6.1 Mbp CHA0 genome (Figure 1). Functional characterization of the

mutated genes by analysis of their Clusters of Orthologous Groups (COGs) annotation revealed that the

majority of these genes are involved in transcription (COG term 'K'), signal transduction mechanisms

(COG term 'T'), amino acid transport and metabolism (COG term 'E') and cell wall/membrane/envelope

biogenesis (COG term 'M') (Figure 1, Table 1).

Table 1. Mutations that occurred in 28 genes during Arabidopsis rhizosphere adaptation divided over five replicate CHA0 populations (lines)

Locus tag	Gene name	Description	COG[1]	Line	Total number of alleles	Number of mutations in protein-coding genes			Number of intergenic mutations
						Non-synonymous	Synonymous	Deletions	
PFLCHA0_RS02080	hutI	Imidazolonepropionase	F	4	1	0	1	0	0
PFLCHA0_RS03400	accC	Biotin carboxylase, acetyl-CoA carboxylase subunit	I	2	1	1	0	0	0
PFLCHA0_RS05510	nudL	Coenzyme A pyrophosphatase/nudix hydrolase NudL	L	4	1	0	1	0	0
PFLCHA0_RS06125	rpoS	RNA polymerase sigma factor RpoS	K	5	1	1	0	0	0
PFLCHA0_RS08340	fleQ	Sigma-54-dependent Fis family transcriptional regulator	T	5	1	1	0	0	0
PFLCHA0_RS08490	flhA	Flagellar biosynthesis protein FlhA	N	4	1	0	0	1	0
PFLCHA0_RS09880	RS09880	Glycosyltransferase (GT)	M	5	1	1	0	0	0
PFLCHA0_RS09890	oafA	O-antigen acetylase	I	1, 3	2	1	0	1	0
PFLCHA0_RS09920	galE	UDP-glucose 4-epimerase	M	2	1	1	0	0	0
PFLCHA0_RS11785	RS11785	LysR family transcriptional regulator	K	4	1	1	0	0	0
PFLCHA0_RS11820	RS11820	PaaI family thioesterase	Q	4	1	0	1	0	0
PFLCHA0_RS12070	RS12070	2OG-Fe(II) oxygenase superfamily	S	5	1	0	1	0	0
PFLCHA0_RS13000	yvaQ2	Methyl-accepting chemotaxis protein	NT	2	1	0	0	0	1
PFLCHA0_RS14960	tetR	TetR/AcrR family transcriptional regulator	K	5	1	1	0	0	0
PFLCHA0_RS17350	RS17350	Methyltransferase domain-containing protein	H	1	1	0	0	1	0
PFLCHA0_RS17965	gacA	UvrY/SirA/GacA family response regulator transcription factor	K	1, 2, 4	6	5	0	0	1
PFLCHA0_RS18525	RS18525	ABC transporter substrate-binding protein	E	5	1	1	0	0	0
PFLCHA0_RS21265	pvdS	RNA polymerase factor sigma-70	K	1	1	0	0	0	1
PFLCHA0_RS21275	RS21275	Transporter substrate-binding domain-containing protein	ET	1	1	1	0	0	0
PFLCHA0_RS21855	wbpM	Polysaccharide biosynthesis protein/NDP-sugar epimerase	GM	1	1	1	0	0	0
PFLCHA0_RS22600	gacS	Hybrid sensor histidine kinase/response regulator	T	3	1	1	0	0	0
PFLCHA0_RS22950	argT5	ABC transporter substrate-binding protein/Lysine-arginine-ornithine-binding periplasmic protein ArgT	ET	4	1	0	0	1	0
PFLCHA0_RS25175	mraZ	Division/cell wall cluster transcriptional repressor MraZ	K	2	1	0	0	0	1
PFLCHA0_RS26215	osmY	Osmotically-inducible protein OsmY/BON domain-containing protein	S	2	1	0	0	0	1
PFLCHA0_RS27515	rpsH	30S ribosomal protein S8	J	4	1	0	0	0	1
PFLCHA0_RS30075	sadB	Surface attachment defective (SadB) ortholog/HDOD domain-containing protein	T	1	2	2	0	0	0
PFLCHA0_RS30120	tssM	Type VI secretion system membrane subunit TssM	S	2	1	0	1	0	0
PFLCHA0_RS31060	nlpD	Lipoprotein nlpD/lppB/LysM domain-containing protein	D	3	1	1	0	0	0
Total	28				35	20	5	4	6

[1]Clusters of Orthologous Groups (COGs); https://www.ncbi.nlm.nih.gov/COG/(30)

Identification of potential root colonization genes

Bacterial genes that are involved in colonization of plant roots can be revealed by identifying beneficial mutations that evolve during adaption of bacteria to the rhizosphere environment and have positive effects on root colonization. Over time, such mutants will outcompete the ancestral strain and become dominant in the bacterial population. The observation that only a limited number of mutations accumulated relative to the total number of genes within the genome across the entire experiment, makes it highly unlikely that the same gene would acquire several changes by chance in independent evolutionary lines. Nevertheless, we observe recurrent mutations in several of the same genes and/or pathways (Table 1), which is a strong indication for adaptive evolution. Genes or pathways that acquired mutations in multiple independent CHA0 populations included *gacA* and *gacS*, the *OBC3* gene cluster (*oafA*; *galE*; *PFLCHA0_RS09880* (29)), and a putative pyoverdine siderophore biosynthesis cluster (*pvdS*; *PFLCHA0_RS21275*). Other genes were targeted more than once in the same population, such as *sadB*. Because these genes were repeatedly identified in the CHA0 evolution experiment, they can be assumed to contribute significantly to bacterial fitness in the rhizosphere (Table 1, Figure 2). Moreover, it suggests that the independent evolutionary lines converge on similar evolutionary trajectories involving overlapping biological processes and molecular mechanisms.

To illustrate the diverse evolutionary trajectories, we constructed phylogenetic trees of each replicate population (line) using all mutations that were detected (Figure 3). In each line, both the number and depth of branches are very different (Figure 3, Table 2). In line 1 and 3, the populations were swept early by *oafA* mutants and later by specific *gac* mutants, while in line 2 and 4, co-existence of multiple genotypes characterized by *gac* mutations was observed. The evolved isolates contain one to four mutations in general, and in each line, 3 (line 3) to 9 (line 2 and 4) mutations were identified in total. All mutations were unique for each line, as is expected for independently evolving populations.

The frequency of each mutation (Table 2) was determined from the six bacterial colonies that were isolated and sequenced from each evolutionary line at cycles 2, 4 and 6. To further investigate and verify the changes in mutation frequency in the entire CHA0 population at the end of a given cycle, we set out to determine the frequency of three of the mutations, i.e. *gacA*^{D49Y}, *gacA*^{D54Y}, and *gacS*^{G27D}, with increased accuracy and resolution during long-term rhizosphere adaptation. To this end, we developed a PCR-based high-resolution melting (HRM) analysis using mutation-specific HRM probes (Table S2), that allow for the accurate quantification of mutant frequency across a wide range for all three mutations ($p < 0.001$; Figure S1). Using this method, we found that the population-level frequency of these three

mutations in the three respective evolutionary lines at the end of each of the eight rhizosphere

adaptation cycles corresponded well with the frequency previously obtained from the six cultured and

sequenced isolates (Table 2, Figure S2). These findings indicate that the culture-based quantification of

mutant frequency provides a realistic measure for population-level frequency. Moreover, it can be seen

that whereas the $gacS^{G27D}$ mutation reaches fixation after eight cycles of rhizosphere adaptation, the

$gacA^{D49Y}$ and $gacA^{D54Y}$ mutants stabilize at around 50% and 25%, respectively, indicative of frequency-

dependent (FD) selection (Figure S2). FD selection describes the phenomenon that the fitness of a

particular genotype or phenotype is dependent on its frequency. Such context-dependency has been

frequently linked to cheating behaviour in which microbial cells that lack or limit the production of

certain costly compounds benefit from other cells that do produce these compounds. When a minimal

amount of such compound increases genotype or phenotype fitness, FD selection can occur resulting at

the stabilization of the mutation frequency.

Table 2. Nomenclature for CHA0 mutations and variants and their frequency/cycle in each respective population.

| Line | Locus tag | Gene name | DNA sequence change[1] | Amino acid change (three-letter code)[2] | Standard nomenclature (used in text) | Frequency in the population (x/6) | | | |
						Cycle 2	Cycle 4	Cycle 6	RQ-HRM[3]
1	PFLCHA0_RS09890	*oafA*	c.1005C>A	p.Tyr335X	Y335X	4/6	6/6	6/6	
1	PFLCHA0_RS17350	*RS17350*	c.230delC	p.Ala77Ala.fsX14	A77A.fsX14	2/6	6/6	6/6	
1	PFLCHA0_RS17965	*gacA*	c.145G>T	p.Asp49Tyr	D49Y	-	2/6	4/6	x
1	PFLCHA0_RS21855	*wbpM*	c.235G>C	p.Gly79Arg	G79R	-	1/6	-	
1	PFLCHA0_RS21275	*RS21275*	c.233G>T	p.Arg78Leu	R78L	-	1/6	-	
1	PFLCHA0_RS21265	*pvdS*	c.-33T>C	NA	-33T>C	-	-	1/6	
1	PFLCHA0_RS30075	*sadB*	c.548G>C	p.Arg183Pro	R183P	-	-	1/6	
1			c.773T>A	p.Leu258Gln	L258Q	-	-	2/6	
2	PFLCHA0_RS17965	*gacA*	c.548A>C	p.Tyr183Ser	Y183S	2/6	1/6	-	
2	PFLCHA0_RS09920	*galE*	c.94G>A	p.Val32Met	V32M	2/6	5/6	6/6	
2	PFLCHA0_RS30120	*tssM*	c.1336C>T	p.Leu446Leu	1336C>T	1/6	-	-	
2	PFLCHA0_RS03400	*accC*	c.1237G>A	p.Glu413Lys	E413K	-	4/6	6/6	
2	PFLCHA0_RS13000	*yvaQ2*	c.-9G>T	NA	-9G>T	-	1/6	-	
2	PFLCHA0_RS17965	*gacA*	c.160G>T	p.As54Tyr	D54Y	-	-	2/6	x
2	PFLCHA0_RS17965	*gacA*	c.289G>A	p.Gly97Ser	G97S	-	-	3/6	
2	PFLCHA0_RS25175	*mraZ*	c.-211A>G	NA	-211A>G	-	-	2/6	
2	PFLCHA0_RS26215	*osmY*	c.-104G>A	NA	-104G>A	-	-	1/6	
3	PFLCHA0_RS09890	*oafA*	c.1013delA	p.Lys338Ser.fsX18	K338S.fsX18	1/6	5/6	6/6	
3	PFLCHA0_RS22600	*gacS*	c.80G>A	p.Gly27Asp	G27D	-	1/6	4/6	x
3	PFLCHA0_RS31060	*nlpD*	c.590A>C	p.Gln197Pro	Q197P	-	1/6	-	
4	PFLCHA0_RS22950	*argT5*	c.120-533del	p.Lys40-Leu185del.fsX6	ΔargT5	2/6	-	-	
4	PFLCHA0_RS02080	*hutI*	c.786C>T	p.Phe262Phe	786C>T	1/6	-	-	
4	PFLCHA0_RS17965	*gacA*	c.-40T>A	NA	-40T>A	-	2/6	1/4	
4	PFLCHA0_RS17965	*gacA*	c.112G>T	p.E38X	E38X	-	2/6	3/4	
4	PFLCHA0_RS05510	*nudL*	c.288C>T	p.Ser96Ser	288C>T	-	1/6	-	
4	PFLCHA0_RS11820	*RS11820*	c.33C>T	p.Ala11Ala	33C>T	-	1/6	-	
4	PFLCHA0_RS08490	*flhA*	c.1176delT	p.His393Gln.fsX15	H393Q.fsX15	-	-	3/4	
4	PFLCHA0_RS11785	*RS11785*	c.766A>T	p.Ser256Cys	S256C	-	-	3/4	
4	PFLCHA0_RS27515	*rpsH*	c.-90C>G	NA	-90C>G	-	-	1/4	
5	PFLCHA0_RS18525	*RS18525*	c.254delT	p.Val85Gly.fsX40	V85G.fsX40	1/6	-	-	
5	PFLCHA0_RS06125	*rpoS*	c.193C>T	p.Gln65X	Q65X	-	1/6	4/6	
5	PFLCHA0_RS08340	*fleQ*	c.959G>A	p.Arg320Gln	R320Q	-	-	1/6	
5	PFLCHA0_RS14960	*tetR*	c.381C>G	p.Tyr127X	Y127X	-	-	2/6	
5	PFLCHA0_RS09880	*RS09880*	c.246C>A	p.Tyr82X	Y82X	-	-	1/6	
5	PFLCHA0_RS12070	*RS12070*	c.1389C>G	p.Pro463Pro	1389C>G	-	-	1/6	

160 [1]DNA sequence change positions are relative to the cDNA (c); del represents a deletion

161 [2]X represents a stop codon (at its relative position in case of a shifted frame); fs represents a frame shift; del represents a

162 deletion

163 [3]Mutations that are selected for the analysis of their frequency trajectories using RQ-HMR analysis

Early adaptations are driven by cell surface-related genes

In general, mutations that are fixed early on in the rhizosphere adaptation process tend to have a high selective advantage (31). Disruptive mutations in *oafA*, resulting in premature stops halfway the coding region, were detected as the first acquired mutations in two independent evolutionary lines (lines 1 and 3) and appear to have swept the population in later generations (Table 2). This gene, which is part of the previously described *OBC3* gene cluster (29), encodes an O-acetyltransferase, which is postulated to acetylate the O-antigen component of the outer membrane LPS (32). Another *OBC3* mutation that accumulated early on in the rhizosphere adaptation process was *galE*$^{c.94G>A}$ leading to an amino acid substitution (V32M) in galE which swept through the population in evolutionary line 2, reaching fixation in cycle 6 (Table 2). *GalE* encodes an UDP-glucose 4-epimerase which is involved in O-antigen and LPS core biosynthesis (33–35). One colony with a mutation in a third *OBC3* cluster gene, *RS09880*, encoding a putative glycosyl transferase was found in evolutionary line 5. Thus, in four out of the five evolutionary lines mutations that likely affect bacterial LPS structure appeared during rhizosphere adaptation, and in three evolutionary lines such mutations even became dominant in the population. These results strongly suggest that modifying bacterial cell surface structure is an important bacterial strategy in early adaptation to the rhizosphere.

Adaption driven by global regulators

In the present study, six mutations were detected in *gacA* in three of the five evolutionary lines, representing approximately 20% of all missense mutations. Notably, in evolutionary lines 2 and 4, multiple *gacA* alleles accumulated, some of which were detected in early plant adaptation cycles (Table 2). Additionally, a *gacS* mutation accumulated in line 3. These two genes encode the main constituents of the conserved GacA/GacS two-component regulator system, i.e. the hybrid sensor histidine kinase GacS and the cognate response regulator GacA (Figure 2). In gram-negative bacteria, activation of GacS results in cross-phosphorylation of GacA via phosphotransfer which in turn leads to activation of the expression of the small RNA genes *rsmY* and *rsmZ* via its helix-turn-helix (HTH) domain-binding domain (36). In CHA0, this regulatory pathway is known to control quorum sensing as well as secondary metabolism and stress resistance (37–40). In the closely related strain *P. protegens* Pf-5 this pathway was shown to have a big impact on bacterial genome-wide gene expression, affecting the expression of more than 10% of the annotated genes (41). Similarly, *gacA* mutants that arose on the edge of swarming Pf-5 colonies showed dramatically altered genome-wide gene expression patterns (42).

Including *gacS* and *gacA*, about half of the mutated genes in this study are global regulators or sigma

factors (Table 1, Figure 1). This high frequency suggests that global regulator-controlled networks are

evolvable and play a major role in rapid bacterial adaptation. Pleiotropic adaptive mutations in global

regulator genes have been shown to be important for bacterial adaption both in laboratory (43, 44), and

in natural settings (45, 46). Remodeling and continuous optimization of existing regulatory networks by

single mutations is an important strategy for bacterial adaptation to the host (47).

Bacterial motility

Bacterial motility is an important trait for rhizosphere competence, mediating colonization of distal parts

of the root system (48), and both LPS and the GacS/GacA two-component regulator system affect this

trait. The O-antigenic side chain of the LPS was reported to contribute to swimming and swarming

motility in the plant pathogenic bacterium *Erwinia amylovora* (49, 50). The GacS/GacA two-component

regulator system controls bacterial motility, for example by affecting transcription of genes related to

flagella and biosurfactant biosynthesis (42, 48, 51). We also identified several other mutations across

the various evolutionary lines that can be linked to bacterial motility. For instance, a disruptive mutation

in the flagellar biosynthesis protein-coding gene *FlhA* (H393Q.fsX15) that is involved in the biogenesis of

the bacterial flagellum (Table 1) appeared in evolutionary line 4 reaching up to a predicted frequency of

75% (Table 2). In *P. fluorescens* Pf-5, and in *P. aeruginosa* PA01 FlhA is reported to be essential for

swimming motility (41).

Furthermore, we identified one amino-acid substitution, R320Q, in FleQ and two in SadB, R183P and

L258Q respectively, that based on sequence similarity to well-studied, homologous proteins in other

bacteria are likely to affect motility as well. FleQ is a σ^{54}-dependent Fis family transcriptional regulator

which regulates flagellar motility, biofilm formation as well as Pel exopolysaccharide production in

response to cellular c-di-GMP in *P. aeruginosa* (52). *P. protegens* FleQ shares 84% sequence identity and

98% sequence coverage with *P. aeruginosa* FleQ and similarly contains an N-terminal flagellar regulatory

domain FleQ (PF06490) also known as the REC receiver domain, a central AAA+/ATPase σ^{54}-interaction

domain (PF00158), and a C-terminal Fis-type HTH DNA-binding domain (PF02954). The R320Q

substitution we identified here is found in the AAA+/ATPase σ^{54}-interacting domain in between the

arginine (R) finger (300-303) and a c-di-GMP-binding motif, ExxxR (330-334) (52). The conserved arginine

residues in FleQ, including the here mutated R320, are thought to be important for protein

oligomerization, and substitution of any of them abolishes ATPase activity in *Vibrio cholerae* EpsE

completely (53). SadB, finally, encodes a HDOD domain-containing protein (Table 1) which shares 77%

sequence identity and 99% sequence coverage with *P. aeruginosa* SadB (PA5346). In *P. aeruginosa*, SadB

stimulates Pel exopolysaccharide production and the chemotaxis-like cluster CheIV, which in turn affect

flagellar motility as well as biofilm formation (54). In *Pseudomonas fluorescens* F113, SadB, together

with FleQ control flagellar motility, dependent and independent of the GacS/GacA two-component

regulator system (55, 56).

Parallelism of targeted mutations on this functional motility pathway impelled us to assess bacterial

motility and track its dynamics across multiple evolutionary lines. Two independent evolutionary

trajectories to genotypes *galE*^{V32M} · *accC*^{E413K} · *gacA*^{D54Y} and *oafA*$^{K338S.fsX18}$ · *gacS*^{G27D} from evolutionary

lines 2 and 3, respectively, were studied for changes in their swimming and swarming behavior. Both

genotypes accumulated mutations in *OBC3* cluster genes (29) linked to LPS biogenesis (*galE* or *oafA*) and

the GacS/GacA two-component regulator system (*gacA* or *gacS*). Interestingly, *galE*^{V32M} and *oafA*$^{K338S.fsX18}$

mutants displayed opposing phenotypes when their swimming and swarming behavior was examined.

Whereas the *galE*^{V32M} mutant produces significantly larger swimming and swarming haloes as compared

to the wild type, the *oafA*$^{K338S.fsX18}$ mutant produces significantly smaller swimming haloes and is even

completely defective in swarming (Figure 4). The *gacA* and *gacS* mutations, however, consistently

reduced swimming and swarming behavior.

Dynamics of global phenotypic change

Since natural selection eventually operates at the phenotypic level, revealing bacterial global phenotypic

evolutionary dynamics can help us to identify traits that are under selection. Moreover, beneficial

genetic mutations can be predicted if they are linked to well-known root colonization traits. A broad

range of 30 bacterial traits including different aspects of bacterial life-history traits, were measured for

the sequenced isolates, allowing genotype-phenotype association analysis.

As shown in Figure 5a, the 30 bacterial traits separated into four clusters that share a similar pattern

across the different mutant genotypes and this clustering is supported by model-based clustering

analysis (Figure S3). Growth of the bacteria in King's B (KB) medium was positively correlated with

siderophore and exoprotease production, tryptophan side oxidase activity, and growth inhibition of the

bacterial plant pathogen *Ralstonia solanacearum*. Thus, this cluster, designated cluster 1, contains traits

associated with bacterial social behavior, related to within population communication and cooperation,

such as the production of public goods. In a principle component analysis (Figure S4a), the first principle
component (PC1) is strongly correlated with all five traits, with a total explanation of 64.5% for all
variables. Cluster 2 contains traits linked to carbon source utilization. For this cluster PC1 is strongly
correlated with all carbon source usage-related traits, with a total explanation of 83.9% for all variables
(Figure S4b). A third cluster was observed for the bacterial ability to form a biofilm, to produce indole-3-
acetic acid (IAA), and to inhibit the growth of two fungal plant pathogens, with a total explanation of
82.4% by PC1 (Figure S4c). Finally, cluster 4 contains all seven traits that are related to bacterial
resistance to biotic and abiotic stresses. The first principle component (PC1) is strongly correlated with
all seven traits, with a total explanation of 53.9% for all variables (Figure S4d). For these four clusters,
the PC1 (or -PC1) value is used as a proxy to present the general performance of all the traits that
clustered together.

As is shown in Figure 5b, all five evolutionary lines showed a parallel trend of accumulative decline of
social traits. Mutations in *gacA, gacS,* and *rpoS* resulted in significantly decreased bacterial social traits,
and this was also observed for the double mutant $RS11785^{S256C} \cdot flhA^{H393Q.fsX15}$ relative to the background
$gacA^{E38X}$ mutation. *RS11785*, like *gacA*, encodes a LysR-type transcriptional regulator. In addition to
these global regulators, earlier mutations in *oafA*, *galE* and $RS17350^{A77A.fsX14}$ encoding a putative
methyltransferase, also resulted in a significant but relatively small decrease in social traits. This parallel
decline of the bacterial social traits index suggests a negative selection of bacterial social behavior,
especially the production of costly public goods such as siderophores and exoproteases.

All traits related to the utilization of 14 different carbon sources, that were selected based on their
reported presence in root exudates of Arabidopsis (57) grouped in one cluster (cluster 2; Figure 5a),
suggesting that increase or decrease of the ability to utilize carbon sources is co-regulated. Different
mutations in *gacS* and *gacA* resulted in contrasting bacterial carbon source utilization. A mutation in the
first N-terminal transmembrane domain of GacS (G27D) (Figure 2) resulted in significant enhancement
of carbon source utilization (Figure 5a, c). A similar trend was observed for mutant $gacA^{G97S}$. In contrast,
the majority of evolved genotypes, including most mutations in *gacA* and *rpoS*, showed a reduced ability
to utilize carbon sources (Figure 5a, c).

In contrast to the previous two clusters, traits in clusters 3 (Figure 5d) and 4 (Figure 5e) were more
stable, in that most genotypes behave like the ancestor. Only two *gacA* mutants, $gacA^{E38X}$ and $gacA^{Y183S}$,
were associated with a significant decline of bacterial traits associated with biocontrol, such as
antifungal activity and biofilm formation, while the $rpoS^{Q65X}$ mutation dramatically increased these traits.

The double mutation $RS11785^{S256C} \cdot flhA^{H393Q.fsX15}$, in the background of the disruptive $gacA^{E38X}$ mutation

(Figure 5e), was the only mutational event that led to a significant increase of general resistance to

various environmental stresses. *FlhA*, as previously mentioned encodes the flagellar biosynthesis protein

FlhA and is linked to bacterial motility. The LysR-type transcriptional regulator RS11785 in addition to its

N-terminal HTH DNA-binding domain (PF00126) contains a predicted C-terminal substrate binding

domain (PF03466), which is linked to aromatic compound degradation and resembles that of the type 2

periplasmic binding proteins or PBP2s. PBP2s are responsible for binding and uptake of several

substrates including polysaccharides and several amino acids. We used HHpred (58) for protein remote

homology detection and three-dimensional structure analysis to assess the possible consequence of the

S256C amino acid substitution. This analysis identified the remote homolog OxyR from *Escherichia coli*

as the best template for modeling (E-value: 2.9e-30, Score: 166.45). Intriguingly, it was previously shown

that the activity of OxyR, which represents the master peroxide sensor in gram-negative bacteria, is

regulated by intramolecular disulfide-bond formation in the regulatory PBP2-like domain. Notably,

RS11785 is unlikely to represent the master peroxide sensor in *P. protegens* CHA0, because another

CHA0 protein encoded by *PFLCHA0_RS030065* is much more similar (89% percent pairwise sequence

identity versus 29%). Nevertheless, it is tempting to speculate that the here observed S256C substitution

might influence regulatory activity by altered intramolecular disulfide-bond formation. Lastly, RS11785

bears resemblance (37% sequence identity, 93% sequence coverage) to the LysR-type PBP2-domain

containing regulator *alsR* from *Bacillus amyloliquefaciens*. *AlsR* regulates the activity of the acetoin

operon (*alsSD*) in response to several signals such as glucose and acetate. Acetoin (3-hydroxy-2-

butanone, a precursor for 2,3-butanediol) in turn is known as a potent elicitor of induced systemic

resistance (59), and it is linked to general plant growth promotion in *Bacilli*. Given these observations

from literature, but in the absence of singular mutations in *flhA* and *RS11785*, the molecular mechanism

underlying the observed enhanced resistance in this double mutant remains to be clarified but we

hypothesize that altered regulatory activity by RS11785 is a likely driver of the increased stress

resistance observed in this mutant.

Discussion

The rhizosphere is a nutrient-rich environment for root-associated bacteria. However, to access the available nutrients bacteria must overcome various challenges, including the plant immune system, presence of competing and/or predatory microorganisms, and abiotic stresses. In the gnotobiotic binary system used in this study we tracked changes in *Pseudomonas protegens* CHA0 in the rhizosphere of Arabidopsis under reproducible and controlled conditions without interference of complex interactions with other microbes. Mutations affecting global regulators, bacterial cell surface structure, and motility accumulated in parallel in our evolutionary experiment, revealing at least three important strategies of bacterial adaptation to the rhizosphere.

Global regulators and rhizosphere adaptation

The GacS/GacA two-component regulator system controls the production of antimicrobial secondary metabolites, exoenzymes, and siderophores, but also biofilm formation, stress responses, motility, and quorum sensing (13, 38, 41, 60, 61). In the present study, mutations in the GacS/GacA two-component regulator system caused dramatic changes in several bacterial phenotypic traits, including motility, carbon source utilization, and social traits (Figures 4 and 5). The observed loss of motility does not logically fit with adaptation to the rhizosphere as motility has been reported to be an important root colonization trait in Pseudomonas (17). In line with this, the *gac* mutations that accumulated in *Pseudomonas fluorescens* F113 colonizing the rhizosphere of alfalfa showed increased motility and resulted in an enhanced competitive colonization ability (48, 62). Our results fit observations by Song et al (2016)(42) in the closely related *P. protegens* strain Pf-5. During swarming motility, *gac* mutants emerged that lack production of the surfactant orfamide. These mutants cannot swarm, but they co-swarm with orfamide producing cells (42). Thus, the lack of motility in CHA0 may also be the result of loss of surfactant production and leads to a selective advantage in the presence of surfactant producers. Several other mutations that are identified in diverse regulatory, signaling and executionary genes, *flhA*, *fleQ* and *sadB* can be linked to bacterial motility. Although no data is currently available for the *flhA* and *sadB* mutants, preliminary analysis of swarming and swimming capacity *in vitro* shows that the *fleQ* mutant has also lost its ability to move. Possibly, in our experimental system motility is less important than the ability to adhere to the root surface and the formation of a biofilm that is frequently described alongside motility as two sides of the same coin. In many bacteria, these processes in fact appear

inversely correlated and controlled by the intracellular secondary messenger cyclic dimeric-GMP (c-di-
GMP) (63–65). Future experiments should reveal whether indeed such trade-off between bacterial
motility and root adherence underlies our observations in this evolutionary experiment.

The decreased production of public goods, such as siderophores and exoprotease, which were observed
in all independent evolutionary lines (Figure 5), could be beneficial for bacterial fitness by saving energy
and primary metabolites. For example, adaptation of *Pseudomonas aeruginosa* to the human host
through mutations in regulators was accompanied by loss of siderophore production, secreted
proteases, and biofilm formation (66, 67).

Cell surface structures as bacterial adaptation targets

Bacterial surface components are the first line of defense against environmental stress and interplay
with hosts (68–70). Lipopolysaccharide (LPS) is a central outer membrane component for gram-negative
bacteria, and exhibits structural adaptability that is contributed especially by its O-antigen part (68, 70).
Bacterial O-antigen structure modification plays an important role in evasion of host immunity (71), and
has the potential to change host-bacteria interactions (68, 70, 72). In plant pathogenic bacteria LPS
components are important virulence determinants (73, 74), that activate a variety of defense-related
responses (73, 75–77). In *P. fluorescens* the O-antigen component has been implicated to induce
systemic resistance in radish (78) and Arabidopsis (79).

We observed parallel mutations in genes that are involved in LPS biosynthesis and structure
modification. In three out of the five evolutionary lines, the first fixed mutations were identified in *oafA*
and *galE* that are annotated as O-antigen biosynthesis and structure modification related (Table 1). *OafA*
encodes an O-acetyltransferase, which is postulated to modify the O-antigen by acetylation (32). The
enzyme GalE, UDP-galactose 4-epimerase, is involved in the interconversion of UDP-glucose to UDP-
galactose, an essential intermediate for LPS core and O-antigen structures (33–35). Inactivation of *galE*
in *Porphyromonas gingivalis* resulted in shortening of the O-antigenic side chain (80). And in
Bradyrhizobium japonicum disruption of *galE* resulted in complete lack of the O-antigen (33). Thus, it is
tempting to speculate that evasion of the plant's immune response plays a role in adaptation of CHA0 to
the rhizosphere of Arabidopsis.

In conclusion, the observed bacterial genetic and phenotypic adaption dynamics emphasize an
important role of global regulators, motility, and cell surface structures in bacterial adaptation to its

host. The parallel emergence of mutations in similar genes resulting in specific fitness advantages in the

rhizosphere, suggest that this evolutionary process is driven by the rhizosphere environment.

Materials and Methods

Experimental setup

We set up an experimental evolution experiment with *Arabidopsis thaliana* (Arabidopsis) as host plant

and *Pseudomonas protegens* CHA0 (CHA0) as the evolving bacterial strain. CHA0 (81) is a model strain

originally isolated from roots of tobacco plants grown in soil naturally suppressive to black root rot (82).

CHA0 was chromosomally tagged with GFP and a kanamycin resistance cassette (Jousset et al 2006) to

enable easy tracking of the strain and checking for contaminations. Briefly, the ancestral bacterial

population was inoculated on Arabidopsis roots in a gnotobiotic system. After a plant growth cycle of 4

weeks, evolved bacterial populations were transferred to new plants and this cycle was repeated 8

times. Bacterial colonies were retrieved from the evolved populations at the end of each growth cycles

by plating.

We selected six bacterial isolates from each independent evolutionary line at the end of plant growth

cycles two, four and six. Together with six isolates from the ancestor population, for a total of 94 isolates

various life-history traits (as shown below) were measured and their genomes re-sequenced using the

NextSeq-500 Illumina platform (2 x 75 bp paired-end). We used the snippy pipeline

(https://github.com/tseemann/snippy), integrating reference genome-based mapping of the Illumina

reads by BWA-MEM, variant calling by SAMtools and FreeBayes, and variant impact analysis using SnpEff

(83), to identify single nucleotide polymorphisms (SNPs) and small indels and deletions (INDELs). Larger

INDELs were identified by calculating the breath of coverage of the mapped Illumina reads on the

reference genome in a sliding window using bedtools (84). Regions with reduced coverage (<99%) were

manually inspected in the Integrative Genome Viewer (IGV). For two isolates from experimental line

four re-sequencing failed.

Bacterial life-history traits

A variety of bacterial life-history traits, reflecting various aspects of bacterial physiological processes,

were measured. We measured bacterial growth yield and resistance to various stresses, including acidic

(pH = 5) and alkaline (pH = 9) conditions, oxidative stress in 0.0025% H_2O_2, water potential stress (15%

polyethylene glycol (PEG)-6000)), and salt stress (2% NaCl), and resistance to the antibiotics

streptomycin (1 µg · ml^{-1}), tetracycline (1 µg · ml^{-1}), and penicillin (5 µg · ml^{-1}). Bacterial growth yield was

quantified in modified Ornston and Stanier (OS) minimal medium (85) supplemented with single carbon

sources that have been reported to be abundant in Arabidopsis root exudates (Chaparro et al 2013).

These included the following carbon sources; alanine, arabinose, butyrolactam, fructose, galactose,

glucose, glycerol, glycine, lactic acid, putrescine, serine, succinic acid, threonine, and valine that were

added to a final concentration of 0.5 g · l^{-1}. Growth yield was measured as OD_{600} (SPECTROstar Nano

plate reader), and this measure was used to determine carbon source utilization ability. In addition, we

measured bacterial auxin (indole-3-acetic acid or IAA) production with a colorimetric test (86), iron-

chelating ability using a modified chrome azurol S (CAS) assay (87), proteolytic activity by the adapted

assay from Smeltzer et al. (1993), tryptophan side chain oxidase activity using a colorimetric assay (88),

and biofilm formation using a modified Chrystal Violet staining assay (89). We further assessed bacterial

antimicrobial activity by quantifying their effect on growth of the fungi *Verticillium dahliae* and *Fusarium*

oxysporum, and the bacterium *Ralstonia solanacearum*. Bacterial motility assays were performed on

Cook's Cytophaga (CC) medium containing 0.2% tryptone (90). This medium was solidified with 0.3 or

0.5% agar to determine swimming and swarming motility, respectively (42). Bacteria were first cultured

in liquid CC medium overnight at 28 °C, and subsequently inoculated in the middle of the motility plates

using toothpicks. Plates were incubated upside-down at 28 °C for 24 h, and the colony areas were

calculated using Image J as a measure of bacterial motility.

Hierarchical and Model-based clustering of bacterial traits

Hierarchical clustering

A heatmap to illustrate the association patterns of bacterial genotypes and their measured traits was

constructed in R using the *ggplot2* package. Hierarchical clustering was performed using the Ward.D2

method, that is based on the squared dissimilarities of the set of objects being clustered (91).

Model-based clustering

We applied a model-based clustering method to reveal the best fitting structures of trait covariance

patterns. For example, some traits might be either directly or indirectly co-regulated by the same gene,

which is expected for global regulators particularly which can co-regulate thousands of genes. We used

the *mclust* package in R to run the model simulation (92). This method assumes that the input data are

distributed as a mixture of two or more clusters. The advantage of the model-based clustering method is

that it avoids heuristic assumptions and uses a soft assignment that every data point has a possibility of

falling to each cluster, which facilitates the best clustering solution. The so-called Bayesian Information

Criterion (BIC) was used to select the best model. A large BIC score indicates a better fit of the model.

This result is in line with the outcome of hierarchical clustering with Adjusted Rand Index (ARI) set as 1,

and *k* set as 4 in "ward.D2" as indicated in Figure 5a. ARI is usually used to evaluate the match degree of

a given clustering solution comparing to the model-based clustering result, with 0 reflecting a random

partition and 1 the boundary of accuracy of a certain clustering solution (92).

Genotype-phenotype association analysis

Bacterial traits within each model-predicted cluster have similar data distribution patterns and covaried

together by the definition of the clustering method. Thus, we applied a linear regression-based method,

i.e. principle component analysis (PCA), to reduce the dimensionality of data and generate a proxy for

each model predicted cluster. These proxies were later used as the x-axis values in Figure 5b-e. We

applied the package *ggbiplot* in R to generate the PCA plots and PC1 index from the normalized

datasets. The proxies were normalized for further analysis.

To examine the effect of each mutation on bacterial phenotype, a Students' *t*-test was used to compare

cluster proxies of evolved genotypes with their direct ancestors. Only genotypes identified more than

once were included in this analysis.

Relative quantification of mutant frequency using HRM profile analysis

We used High-Resolution Melting (HRM) profile analysis with integrated LunaProbes to quantify the

ratio of mutant to wild type genotypes (93–95). The probes and primers used in this study are listed in

Table S2. Primers were designed using Primer3. Probes were designed with the single nucleotide

polymorphism (SNP) located in the middle of the sequence, and the 3' end was blocked by carbon

spacer C3. The primer asymmetry was set to 2:1 (excess primer: limiting primer) in all cases. Pre-PCR

was performed in a 10-μl reaction system, with 0.25 μM excess primer, 0.125 μM limiting primer, 0.25

μM probe, 0.5 μl bacterial sample culture (100-fold diluted saved sample, OD_{600} is about 0.01), 1X

LightScanner Master Mix (BioFire Defense). DMSO with the final concentration 5% was supplemented in

all reactions to ensure the targeted melting domains are within the detection limit of the LightScanner

(Idaho Technology Inc.). Finally, MQ water was used to supplement up to 10 μl. A 96-well black

microtiter plate with white wells was used to minimize background fluorescence. Before amplification,

25 μl mineral oil was loaded in each well to prevent evaporation, and the plate was covered with a foil

seal to prevent the degradation of fluorescent molecules. Amplification was initiated by a holding at 95

°C for 3 min, followed by 55 cycles of denaturation at 95 °C for 30 s, annealing at 60 °C for 30 s and

extension at 72 °C for 30 s and then kept at 72 °C for 10 min. After amplification, samples were heated in

a ThermalCycler (Bio-Rad) shortly to 95 °C for 30 s to denature all double-stranded structures followed

by a rapid cooling to 25 °C for 30 s to facilitate successful hybridization between probes and the target

strands. The plate was then transferred to a LightScanner (Idaho Technology Inc.). Melting profiles of

each well were collected by monitoring the continuous loss of fluorescence with a steady increase of the

temperature from 35 °C to 97 °C with a ramp rate of 0.1 °C /s. The relative quantification was based on

the negative first derivative plots using software MATLAB. The areas of probe-target duplexes melting

peaks were auto-calculated by 'AutoFit Peaks I Residuals' function in software PeakFit (SeaSolve

Software Inc.). The mutant frequency X was calculated using the formula shown below:

$$X = \frac{\text{Area}_{mutant}}{\text{Area}_{mutant} + \text{Area}_{WT}}$$

To validate the HRM method, standard curves were generated by measuring mixed samples with known

proportions of mutant templates: 0%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% and 100%.

Measurements for each sample were done in triplicate. Linear regression formula of each mutant

between actual frequencies and measured frequencies were shown in Figure S1. The high R^2 values, and

nearly equal to 1 slope values of these equations, confirmed that the HRM method can accurately

detect mutants' frequency in a mixed population.

Data availability

The *P. protegens* CHA0-GFP reference genome is deposited on GenBank: RCSR00000000.1. Raw

sequencing data used in this study are deposited at the NCBI database under BioProject PRJNA473919.

A conversion table for the CHA0-GFP to CHA0 gene annotations including recent NCBI accession codes is

available here: https://doi.org/10.6084/m9.figshare.13295828.v2.

Acknowledgments

This work was supported by a China Scholarship Council fellowship (to E.L.), a postdoctoral fellowship of the Research Foundation Flanders (FWO 12B8116RN) (to R.D.J.), and a European Research Council Advanced Grant 269072 (to C.M.J.P.).

Author contributions

E.L., A.J., C.M.J.P., P.A.H.M.B., and R.D.J. designed experiments; E.L. and H.J. performed experiments; E.L. and R.D.J. analyzed data. All authors discussed the results and contributed to the final manuscript.

Figures and Figure legends

a)

b)

Figure 1. Genomic distribution of *Pseudomonas protegens* CHA0 evolutionary adaptations. **a)** Rings from inside to outside: Ring 1, nucleotide position indicator; Ring 2, green/purple GC-skew (-/+); Ring 3, %GC; Ring 4, protein-coding genes; Ring 5, distribution of identified mutations arisen during the evolutionary experiment. Functional annotation of mutated genes are indicated by color representing Cluster of orthologous groups (COGs); the key of which can be seen in panel B. **b)** Frequency of mutations per COG class, highlighting enrichment in the classes T (signal transduction), K (transcription), M (cell wall) and E (amino acid transport and metabolism).

a)

b)

Figure 2. Localization of *gac* and *OBC3* mutations identified in the evolutionary experiment. **a)** Schematic view of the cell wall of the gram-negative bacterium *Pseudomonas protegens* highlighting the presence of the lipopolysaccharide (LPS) on the outside and the presence of the two-component regulator system GacS/GacA on the inner membrane and in the cytoplasm. The arrows indicate the location of the amino acid substitutions that were identified in this study in GacS (G27D) and GacA (G38X, D49Y, D54Y, G97S and Y183S). Note the travel of a phosphate group (P) from GacS to GacA upon signal perception which is accepted by either the Asp49 or Asp54 residue in GacA which are both found mutated in two different mutants. **b)** Genomic region harboring the *OBC3* gene cluster responsible for the synthesis of long O-polysaccharide (O-PS) on the LPS and indicated mutations that were identified in a glycosyl transferase (RS09880; Y82X), the O-antigen acetyltransferase oafA (K338S.fsX18 and Y335X) and the UDP-glucose 4-epimerase (galE; V32M).

Figure 3. Phylogenetic trees of five independently evolving *P. protegens* CHA0 populations. Phylogenies for 18 genomes from each population (16 for the population in experimental line 4), based on the sequential appearance of mutations. Synonymous mutations are underlined. Circles represent unique bacterial genotypes. Intensity of the circle fill represents the time point at which a genotype was firstly detected. *OBC3* and *gac* mutants specifically are highlighted in blue and purple respectively. The frequency of each mutation in the respective population can be found in Table 2.

Figure 4. Effects of *gacS/gacA* mutations on bacterial swimming and swarming behaviour. **a)** The top panels illustrate the experimental layout of motility assays, that are pictured in the middle and bottom panels for swarming and swimming respectively. **b)** The accumulative effects of mutations on swarming (left panel) and swimming (right panel) behaviour. Colony area as a measure for bacterial motility was determined using ImageJ. The number of acquired mutations compared to the ancestor is used to express the genetic distance.

a)

Figure 5. Genotype-phenotype association analysis of evolved bacterial populations. **a)** Heatmap representing the performance of 30 life-history traits of six ancestral and 63 evolved isolates (isolates without mutations or only with synonymous mutation were excluded for this association analysis). The "ward.D2" method was used for hierarchical clustering, based on the dissimilarities of the set of objects being clustered. The four clusters depicted on top are model-predicted. Each column illustrates one bacterial life-history trait, indicated on the bottom of the figure. Each row represents an isolate, indicated by its sample ID number and mutational events. **b-e)** Principle component analysis was applied to generate a general symbolic index for each model predicted cluster. The proxy values of each cluster were normalized, higher values represent better performance. To illustrate the effects of each mutational event, ANOVA was applied to reveal the phenotypic change between each genotype and its identified nearest ancestor in each line. Colour-filled circles represent bacterial genotypes, with different colours representing independent evolving populations and the size representing the count. The dashed line represents the average value of the ancestor.

Supplementary Figures:

Figure S1. Standard curves of measured mutant versus ancestor proportion as a function of the actual proportion, using series of mixed samples with known proportions (0%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% and 100% of mutant frequency). Relative densities of mutants $gacA^{D49Y}$, $gacA^{D54Y}$, and $gacS^{G27D}$ were measured by PCR-based high-resolution melting (HRM) analysis. Measurements for each sample were performed in triplicate. In each plot, the black dots represent the measurements, the blue line the fit which was generated based on linear regression modelling.

Figure S2. Frequency trajectories of $gacA^{D49Y}$, $gacA^{D54Y}$ and $gacS^{G27D}$ mutants during long-term rhizosphere adaption. The X-axis represents the plant-to-plant transferring cycle of the bacterial population. Mutant frequency was determined by quantifying the ratio of mutant allele relative to wild-type allele, using PCR-based high-resolution melting (HRM) analysis. The data shown are the mean of two technical replicates, and error bars represent the standard deviation of the mean.

559 **Figure S3.** Selection of the best clustering model using Bayesian Information Criterion (BIC) using the *mclust* R

560 package. A large BIC score indicates strong evidence for the corresponding model. The VEI model with 4

561 components best fits our data. VEI indicates that the clusters have variable volume, the same shape and an

562 orientation equal to coordinate axes.

a

Figure S4. Coordinate axes transformation of each model predicted cluster using principal component analysis
(PCA). **a.** PCA plot of social trait cluster (cluster 1); **b.** PCA plot of carbon utilization trait cluster (cluster 2); **c.** PCA

569   plot of biocontrol trait cluster (cluster 3); **d.** PCA plot of stress resistance trait cluster (cluster 4). Each dot

570   represents one isolate.

Table S1 Genotypes of whole-genome sequenced CHA0 isolates.

Sample ID	Line	Cycle	Genotype
61	1	2	*WT*
169	1	2	*WT*
13	1	2	*oafA*^{Y335X}
121	1	2	*oafA*^{Y335X}
133	1	2	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$
157	1	2	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$
158	1	4	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$
182	1	4	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$
14	1	4	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *gacA*^{D49Y}
122	1	4	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *gacA*^{D49Y}
86	1	4	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *wbpM*^{G79R}
2	1	4	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *RS21275*^{R78L}
195	1	6	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *gacA*^{D49Y}
242	1	6	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *gacA*^{D49Y}
278	1	6	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *gacA*^{D49Y}
255	1	6	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *gacA*^{D49Y} · *sadB*^{R183P}
206	1	6	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *sadB*^{L258Q}
207	1	6	*oafA*^{Y335X} · *RS17350*$^{A77A.fsX14}$ · *sadB*^{L258Q} · *pvdS*$^{-33T>C}$
87	2	2	*WT*
39	2	2	*gacA*^{Y183S}
183	2	2	*gacA*^{Y183S}
123	2	2	*galE*^{V32M}
135	2	2	*galE*^{V32M}
63	2	2	*tssM*$^{1336C>T}$
172	2	4	*gacA*^{Y183S}
52	2	4	*galE*^{V32M}
16	2	4	*galE*^{V32M} · *accC*^{E413K}
28	2	4	*galE*^{V32M} · *accC*^{E413K}
100	2	4	*galE*^{V32M} · *accC*^{E413K}
160	2	4	*galE*^{V32M} · *accC*^{E413K} · *yvaQ2*$^{-9G>T}$
245	2	6	*galE*^{V32M} · *accC*^{E413K}
220	2	6	*galE*^{V32M} · *accC*^{E413K} · *gacA*^{D54Y}

197	2	6	$galE^{V32M} \cdot accC^{E413K} \cdot gacA^{D54Y} \cdot osmY^{-104G>A}$
221	2	6	$galE^{V32M} \cdot accC^{E413K} \cdot gacA^{G97S}$
196	2	6	$galE^{V32M} \cdot accC^{E413K} \cdot gacA^{G97S} \cdot mraZ^{-211A>G}$
268	2	6	$galE^{V32M} \cdot accC^{E413K} \cdot gacA^{G97S} \cdot mraZ^{-211A>G}$
29	3	2	WT
65	3	2	WT
53	3	2	WT
77	3	2	WT
161	3	2	WT
137	3	2	$oafA^{K338S.fsX18}$
90	3	4	$nlpD^{Q197P}$
18	3	4	$oafA^{K338S.fsX18}$
66	3	4	$oafA^{K338S.fsX18}$
150	3	4	$oafA^{K338S.fsX18}$
186	3	4	$oafA^{K338S.fsX18}$
30	3	4	$oafA^{K338S.fsX18} \cdot gacS^{G27D}$
235	3	6	$oafA^{K338S.fsX18}$
270	3	6	$oafA^{K338S.fsX18}$
199	3	6	$oafA^{K338S.fsX18} \cdot gacS^{G27D}$
222	3	6	$oafA^{K338S.fsX18} \cdot gacS^{G27D}$
234	3	6	$oafA^{K338S.fsX18} \cdot gacS^{G27D}$
198	3	6	$oafA^{K338S.fsX18} \cdot gacS^{G27D}$
57	4	2	WT
105	4	2	WT
81	4	2	WT
129	4	2	$\Delta argT5$
141	4	2	$\Delta argT5$
21	4	2	$hult^{786C>T}$
80	4	4	WT
152	4	4	WT
188	4	4	$gacA^{-40T>A}$
68	4	4	$gacA^{-40T>A} \cdot RS11820^{33C>T}$
164	4	4	$gacA^{E38X}$
20	4	4	$gacA^{E38X} \cdot nudL^{288C>T}$
225	4	6	$gacA^{-40T>A} \cdot rpsH^{-90C>G}$
212	4	6	$gacA^{E38X} \cdot RS11785^{S256C} \cdot flhA^{H393Q.fsX15}$

237	4	6	_gacA_^{E38X} · _RS11785_^{S256C} · _flhA_$^{H393Q.fsX15}$
260	4	6	_gacA_^{E38X} · _RS11785_^{S256C} · _flhA_$^{H393Q.fsX15}$
11	6	2	_WT_
71	6	2	_WT_
191	6	2	_WT_
35	6	2	_WT_
83	6	2	_WT_
143	6	2	_RS18525_$^{V85G.fsX40}$
70	6	4	_WT_
106	6	4	_WT_
142	6	4	_WT_
58	6	4	_WT_
190	6	4	_WT_
34	6	4	_rpoS_^{Q65X}
251	6	6	_fleQ_^{R320Q}
262	6	6	_rpoS_^{Q65X}
263	6	6	_rpoS_^{Q65X}
203	6	6	_rpoS_^{Q65X} · _tetR_^{Y127X}
239	6	6	_rpoS_^{Q65X} · _tetR_^{Y127X}
226	6	6	_RS09880_^{Y82X} · _RS12070_$^{1389C>G}$
7	ancestor	0	_WT_
31	ancestor	0	_WT_
79	ancestor	0	_WT_
127	ancestor	0	_WT_
163	ancestor	0	_WT_
187	ancestor	0	_WT_

572 Note: X, represents a stop codon (at its relative position in case of a shifted frame); fs, frame shift; del, deletion; DNA sequence
573 change positions are relative to the cDNA.

575 **Table S2** Primers and probes used for high-resolution melting (HRM) analysis. For the two *gacA* mutants the same set of
576 primers was used. Underlined bases indicate the position of the single nucleotide point (SNP) mutations within the probe

Target gene	Strain ID	SNP locus	Forward primer (excess)	Reverse primer (limiting)	Amplicon size	Probe sequence	Probe length	Target strand	Perfect match/ mismatch	ΔTm (°C)
gacA[D49Y]	242	145G>T				5'-CATCAGGACCACAT<u>C</u>GGGCTTCAGCTCCCG-/C3/3'	30nt	WT	G::C / T::C	5.31
gacA[D54Y]	220	160G>T	5'-ATCGATGGCCTGCAAGTAGT-3'	5'-CGGGTAGGAAAGGGATCTTC-3'	206 bp	5'-TGGCATCTTGACGT<u>C</u>CATCAGGACCACATC-/C3/3'	30nt	WT	G::C / T::C	4.92
gacS[G27D]	222	80G>A	5'-GCGTACTGTTGCTGACCTTG-3'	5'-AGCATCTGGGTGTTGTGGTT-3'	178bp	5'-AGGTGAAGTAGCCG<u>C</u>CCAGCACCAAAGCCA-/C3/3'	30nt	WT	G::C / A::C	4.83

577 sequences. ΔTm (°C) indicates the melting temperature difference between WT-probe duplex and mutant-probe duplex.

References

1. Berendsen RL, Pieterse CMJ, Bakker PAHM. 2012. The rhizosphere microbiome and plant health. Trends Plant Sci 17:478–486.

2. Bais HP, Weir TL, Perry LG, Gilroy S, Vivanco JM. 2006. The role of root exudates in rhizosphere interations with plants and other organisms. Annu Rev Plant Biol 57:233–266.

3. Stringlis IA, Yu K, Feussner K, de Jonge R, Van Bentum S, Van Verk MC, Berendsen RL, Bakker PAHM, Feussner I, Pieterse CMJ. 2018. MYB72-dependent coumarin exudation shapes root microbiome assembly to promote plant health. Proc Natl Acad Sci U S A 115:E5213–E5222.

4. Bakker PAHM, Pieterse CMJ, de Jonge R, Berendsen RL. 2018. The soil-borne legacy. Cell 172:1178–1180.

5. Philippot L, Raaijmakers JM, Lemanceau P, van der Putten WH. 2013. Going back to the roots: the microbial ecology of the rhizosphere. Nat Rev Microbiol 11:789–799.

6. Sasse J, Martinoia E, Northen T. 2018. Feed your friends: do plant exudates shape the root microbiome? Trends Plant Sci 23:25–41.

7. Bais HP, Park SW, Weir TL, Callaway RM, Vivanco JM. 2004. How plants communicate using the underground information superhighway. Trends Plant Sci 9:26–32.

8. Zamioudis C, Pieterse CMJ. 2012. Modulation of host immunity by beneficial microbes. Mol Plant-Microbe Interact 25:139–150.

9. Yang J, Kloepper JW, Ryu C-M. 2009. Rhizosphere bacteria help plants tolerate abiotic stress. Trends Plant Sci 14:1–4.

10. Browne P, Rice O, Miller SH, Burke J, Dowling DN, Morrissey JP, O'Gara F. 2009. Superior inorganic phosphate solubilization is linked to phylogeny within the *Pseudomonas fluorescens* complex. Appl Soil Ecol 43:131–138.

11. Lugtenberg B, Kamilova F. 2009. Plant growth promoting rhizobacteria. Annu Rev Microbiol 63:541–556.

12. Ryu C-M, Farag MA, Hu C-H, Reddy MS, Wei H-X, Paré PW, Kloepper JW. 2003. Bacterial volatiles promote growth in *Arabidopsis*. Proc Natl Acad Sci U S A 100:4927–4932.

13. Haas D, Défago G. 2005. Biological control of soil-borne pathogens by fluorescent pseudomonads. Nat Rev Microbiol 3:307–319.

14. Pieterse CMJ, Zamioudis C, Berendsen RL, Weller DM, Van Wees SCM, Bakker PAHM. 2014. Induced systemic resistance by beneficial microbes. Annu Rev Phytopathol 52:347–375.

15. Raaijmakers JM, Bonsall RF, Weller DM. 1999. Effect of population density of *Pseudomonas fluorescens* on production of 2,4-Diacetylphloroglucinol in the rhizosphere of wheat. Phytopathology 89:470–475.

16. Raaijmakers JM, Leeman M, Oorschot van MM., et.al. 1995. Dose-response relationships in biological control of Fusarium wilt of radish by *Pseudomonas* spp.. Phytopathology 85:1075–1081.

17. Lugtenberg BJJ, Dekkers L, Bloemberg G V. 2001. Molecular determinants of rhizosphere colonization *Pseudomonas*. Annu Rev Phytopathol 39:461–490.

18. de Weger LA, van der Vlugt CI, Wijfjes AH, Bakker PA, Schippers B, Lugtenberg B. 1987. Flagella of a plant-growth-stimulating *Pseudomonas fluorescens* strain are required for colonization of potato roots. J Bacteriol 169:2769–2773.

19. de Weger LA, van Loosdrecht MC, Klaassen HE, Lugtenberg B. 1989. Mutational changes in physiochemical cell surface properties of plant-growth-stimulating *Pseudomonas* spp. do not influence the attachment properties of the cells. J Bacteriol 171:2756–2761.

20. Simons M, Permentier HP, de Weger LA, Wijffelman CA, Lugtenberg BJJ. 1997. Amino acid synthesis is necessary for tomato root colonization by *Pseudomonas fluorescens* strain WCS365. Mol Plant-Microbe Interact 10:102–106.

21. Berendsen RL, Vismans G, Yu K, Song Y, de Jonge R, Burgman WP, Burmølle M, Herschend J, Bakker PAHM, Pieterse CMJ. 2018. Disease-induced assemblage of a plant-beneficial bacterial consortium. ISME J 12:1496–1507.

22. Yu K, Liu Y, Tichelaar R, Savant N, Lagendijk E, van Kuijk SJL, Stringlis IA, van Dijken AJH, Pieterse CMJ, Bakker PAHM, Haney CH, Berendsen RL. 2019. Rhizosphere-associated *Pseudomonas* suppress local root immune responses by gluconic acid-mediated lowering of environmental pH. Curr Biol 29:3913-3920.e4.

23. Cole BJ, Feltcher ME, Waters RJ, Wetmore KM, Mucyn TS, Ryan EM, Wang G, Ul-Hasan S, McDonald M, Yoshikuni Y, Malmstrom RR, Deutschbauer AM, Dangl JL, Visel A. 2017. Genome-wide identification of

631   bacterial plant colonization genes. PLoS Biol 15:e2002860.

632   24.   Kawecki TJ, Lenski RE, Ebert D, Hollis B, Olivieri I, Whitlock MC. 2012. Experimental evolution. Trends Ecol
633   Evol 27:547–560.

634   25.   Jousset A, Schuldes J, Keel C, Maurhofer M, Daniel R, Scheu S, Thuermer A. 2014. Full-genome sequence of
635   the plant growth-promoting bacterium *Pseudomonas protegens* CHA0. Genome Announc 2:e00322-14.

636   26.   Li E, de Jonge R, Chen L, Friman V, Bakker P, Pieterse C, Jousset A. 2020. Rapid evolution of plant-bacterium
637   mutualism in the rhizosphere. SSRN Electron J https://doi.org/10.2139/ssrn.3650572.

638   27.   Oren Y, Smith MB, Johns NI, Kaplan Zeevi M, Biran D, Ron EZ, Corander J, Wang HH, Alm EJ, Pupko T. 2014.
639   Transfer of noncoding DNA drives regulatory rewiring in bacteria. Proc Natl Acad Sci U S A 111:16112–
640   16117.

641   28.   Carroll SB. 2008. Evo-devo and an expanding evolutionary synthesis: a genetic theory of morphological
642   evolution. Cell 134:25–36.

643   29.   Kupferschmied P, Chai T, Flury P, Blom J, Smits THM, Maurhofer M, Keel C. 2016. Specific surface glycan
644   decorations enable antimicrobial peptide resistance in plant-beneficial *pseudomonads* with insect-
645   pathogenic properties. Environ Microbiol 18:4265–4281.

646   30.   Galperin MY, Makarova KS, Wolf YI, Koonin E V. 2015. Expanded Microbial genome coverage and improved
647   protein family annotation in the COG database. Nucleic Acids Res 43:D261–D269.

648   31.   Barrett RDH, Maclean RC, Bell G. 2006. Mutations of intermediate effect are responsible for adaptation in
649   evolving *Pseudomonas fluorescens* populations. Biol Lett 2:236–238.

650   32.   Allison GE, Verma NK. 2000. Serotype-converting bacteriophages and O-antigen modification in *Shigella*
651   *flexneri*. Trends Microbiol 8:17–23.

652   33.   Chang W-S, Park K-M, Koh S-C, So J-S. 2008. Characterization of the *Bradyrhizobium japonicum galE* gene:
653   its impact on lipopolysaccharide profile and nodulation of soybean. FEMS Microbiol Lett 280:242–249.

654   34.   Liu B, Knirel YA, Feng L, Perepelov A V., Senchenkova SN, Wang Q, Reeves PR, Wang L. 2008. Structure and
655   genetics of *Shigella* O antigens. FEMS Microbiol Rev 32:627–653.

656   35.   Samuel G, Reeves P. 2003. Biosynthesis of O-antigens: genes and pathways involved in nucleotide sugar
657   precursor synthesis and O-antigen assembly. Carbohydr Res 338:2503–2519.

658   36.   Brencic A, McFarland KA, McManus HR, Castang S, Mogno I, Dove SL, Lory S. 2009. The GacS/GacA signal
659   transduction system of *Pseudomonas aeruginosa* acts exclusively through its control over the transcription
660   of the RsmY and RsmZ regulatory small RNAs. Mol Microbiol 73:434–445.

661   37.   Brown D. 2010. A mathematical model of the Gac/Rsm quorum sensing network in *Pseudomonas*
662   *fluorescens*. BioSystems 101:200–212.

663   38.   Heeb S, Blumer C, Haas D. 2002. Regulatory RNA as mediator in GacA/RsmA-dependent global control of
664   exoproduct formation in *Pseudomonas fluorescens* CHA0. J Bacteriol 184:1046–1056.

665   39.   Valverde C, Heeb S, Keel C, Haas D. 2003. RsmY, a small regulatory RNA , is required in concert with RsmZ
666   for GacA-dependent expression of biocontrol traits in *Pseudomonas fluorescens* CHA0. Mol Microbiol
667   50:1361–1379.

668   40.   Heeb S, Valverde C, Haas D. 2005. Role of the stress sigma factor RpoS in GacA / RsmA-controlled
669   secondary metabolism and resistance to oxidative stress in *Pseudomonas fluorescens* CHA0. FEMS
670   Microbiol Lett 243:251–258.

671   41.   Hassan KA, Johnson A, Shaffer BT, Ren Q, Kidarsa TA, Elbourne LDH, Hartney S, Duboy R, Goebel NC,
672   Zabriskie TM, Paulsen IT, Loper JE. 2010. Inactivation of the GacA response regulator in *Pseudomonas*
673   *fluorescens* Pf-5 has far-reaching transcriptomic consequences. Environ Microbiol 12:899–915.

674   42.   Song C, Kidarsa TA, Mortel JE Van De, Loper JE, Jos M. 2016. Living on the edge: emergence of spontaneous
675   gac mutations in *Pseudomonas protegens* during swarming motility. Environ Microbiol 18:3453–3465.

676   43.   Hunter P. 2008. The great leap forward. EMBO Rep 9:608–611.

677   44.   Zambrano MM, Siegele DA, Almirón M, Tormo A, Kolter R. 1993. Microbial competition: *Escherichia coli*
678   mutants that take over stationary phase cultures. Science 259:1757–1760.

679   45.   Sumby P, Whitney AR, Graviss EA, DeLeo FR, Musser JM. 2006. Genome-wide analysis of group A
680   streptococci reveals a mutation that modulates global phenotype and disease specificity. PLoS Pathog 2:e5.

681   46.   Giraud A, Arous S, De Paepe M, Gaboriau-Routhiau V, Bambou JC, Rakotobe S, Lindner AB, Taddei F, Cerf-
682   Bensussan N. 2008. Dissecting the genetic components of adaptation of *Escherichia coli* to the mouse gut.
683   PLoS Genet 4:0052–0061.

684 47. Damkiaer S, Yang L, Molin S, Jelsbak L. 2013. Evolutionary remodeling of global regulatory networks during
685   long-term bacterial adaptation to human hosts. Proc Natl Acad Sci U S A 110:7766–7771.

686 48. Sánchez-Contreras M, Martín M, Villacieros M, O'Gara F, Bonilla I, Rivilla R. 2002. Phenotypic selection and
687   phase variation occur during alfalfa root colonization by *Pseudomonas fluorescens* F113. J Bacteriol
688   184:1587–1596.

689 49. Bowden SD, Hale N, Chung JCS, Hodgkinson JT, Spring DR, Welch M. 2013. Surface swarming motility by
690   *Pectobacterium atrosepticum* is a latent phenotype that requires O antigen and is regulated by quorum
691   sensing. Microbiology 159:2375–2385.

692 50. Berry MC, McGhee GC, Zhao Y, Sundin GW. 2009. Effect of a *waaL* mutation on lipopolysaccharide
693   composition, oxidative stress survival, and virulence in *Erwinia amylovora*. FEMS Microbiol Lett 291:80–87.

694 51. Lapouge K, Schubert M, Allain FH-T, Haas D. 2007. Gac/Rsm signal transduction pathway of γ-
695   proteobacteria: from RNA recognition to regulation of social behaviour. Mol Microbiol 67:241–253.

696 52. Matsuyama BY, Krasteva P V., Baraquet C, Harwood CS, Sondermann H, Navarro MVAS. 2016. Mechanistic
697   insights into c-di-GMP-dependent control of the biofilm regulator FleQ from *Pseudomonas aeruginosa*.
698   Proc Natl Acad Sci U S A 113:E209–E218.

699 53. Patrick M, Korotkov K V., Hol WGJ, Sandkvist M. 2011. Oligomerization of EpsE coordinates residues from
700   multiple subunits to facilitate ATPase activity. J Biol Chem 286:10378–10386.

701 54. Caiazza NC, O'Toole GA. 2004. SadB is required for the transition from reversible to irreversible attachment
702   during biofilm formation by *Pseudomonas aeruginosa* PA14. J Bacteriol 186:4476–4485.

703 55. Muriel C, Blanco-Romero E, Trampari E, Arrebola E, Durán D, Redondo-Nieto M, Malone JG, Martín M,
704   Rivilla R. 2019. The diguanylate cyclase AdrA regulates flagellar biosynthesis in *Pseudomonas fluorescens*
705   F113 through SadB. Sci Rep 9:1–9.

706 56. Martínez-Granero F, Navazo A, Barahona E, Redondo-Nieto M, Rivilla R, Martín M. 2012. The Gac-Rsm and
707   SadB signal transduction pathways converge on AlgU to downregulate motility in *Pseudomonas*
708   *fluorescens*. PLoS One 7:e31765.

709 57. Chaparro JM, Badri D V., Bakker MG, Sugiyama A, Manter DK, Vivanco JM. 2013. Root exudation of
710   phytochemicals in *Arabidopsis* follows specific patterns that are developmentally programmed and
711   correlate with soil microbial functions. PLoS One 8:e55731.

712 58. Zimmermann L, Stephens A, Nam SZ, Rau D, Kübler J, Lozajic M, Gabler F, Söding J, Lupas AN, Alva V. 2018.
713   A completely reimplemented MPI bioinformatics toolkit with a new HHpred server at its core. J Mol Biol
714   430:2237–2243.

715 59. Rudrappa T, Biedrzycki ML, Kunjeti SG, Donofrio NM, Czymmek KJ, Paré PW, Bais HP. 2010. The
716   rhizobacterial elicitor acetoin induces systemic resistance in *Arabidopsis thaliana*. Commun Integr Biol
717   3:130–138.

718 60. Haas D, Keel C. 2003. Regulation of antibotic production in root-colonizing *Pseudomonas* spp. and
719   relavance for biological control of plant disease. Annu Rev Phytopathol 41:117–153.

720 61. Raaijmakers JM, De Bruijn I, Nybroe O, Ongena M. 2010. Natural functions of lipopeptides from *Bacillus*
721   and *Pseudomonas* : more than surfactants and antibiotics. FEMS Microbiol Rev 34:1037–1062.

722 62. Martínez-Granero F, Rivilla R, Martín M. 2006. Rhizosphere selection of highly motile phenotypic variants
723   of *Pseudomonas fluorescens* with enhanced competitive colonization ability. Appl Environ Microbiol
724   61:1946–1952.

725 63. Romling U, Galperin MY, Gomelsky M. 2013. Cyclic di-GMP: the first 25 years of a universal bacterial
726   second messenger. Microbiol Mol Biol Rev 77:1–52.

727 64. Caiazza NC, Merritt JH, Brothers KM, O'Toole GA. 2007. Inverse regulation of biofilm formation and
728   swarming motility by *Pseudomonas aeruginosa* PA14. J Bacteriol 189:3603–3612.

729 65. Simm R, Morr M, Kader A, Nimtz M, Römling U. 2004. GGDEF and EAL domains inversely regulate cyclic di-
730   GMP levels and transition from sessibility to motility. Mol Microbiol 53:1123–1134.

731 66. Smith EE, Buckley DG, Wu Z, Saenphimmachak C, Hoffman LR, D'Argenio DA, Miller SI, Ramsey BW, Speert
732   DP, Moskowitz SM, Burns JL, Kaul R, Olson M V. 2006. Genetic adaptation by *Pseudomonas aeruginosa* to
733   the airways of cystic fibrosis patients. Proc Natl Acad Sci U S A 103:8487–92.

734 67. Winstanley C, O'Brien S, Brockhurst MA. 2016. *Pseudomonas aeruginosa* evolutionary adaptation and
735   diversification in cystic fibrosis chronic lung infections. Trends Microbiol 24:327–337.

68. Lerouge I, Vanderleyden J. 2002. O-antigen structural variation: mechanisms and possible roles in

animal/plant–microbe interactions. FEMS Microbiol Rev 26:17–47.

69.   Whitfield C, Trent MS. 2014. Biosynthesis and export of bacterial lipopolysaccharides. Annu Rev Biochem
83:99–128.

70.   Maldonado RF, Sá-Correia I, Valvano MA. 2016. Lipopolysaccharide modification in Gram-negative bacteria
during chronic infection. FEMS Microbiol Rev 40:480–493.

71.   Murray GL, Attridge SR, Morona R. 2006. Altering the length of the lipopolysaccharide O antigen has an
impact on the interaction of *Salmonella enterica* serovar Typhimurium with macrophages and
complement. J Bacteriol 188:2735–2739.

72.   Ranf S. 2016. Immune sensing of lipopolysaccharide in plants and animals: same but different. PLoS Pathog
12:e1005596.

73.   Newman M-A, von Roepenack E, Daniels M, Dow M. 2000. Lipopolysaccharides and plant responses to
phytopathogenic bacteria. Mol Plant Pathol 1:25–31.

74.   Di Lorenzo F, Silipo A, Andersen Gersby LB, Palmigiano A, Lanzetta R, Garozzo D, Boyer C, Pruvost O,
Newman MA, Molinaro A. 2017. *Xanthomonas citri* pv. *citri* pathotypes: LPS structure and function as
microbe-associated molecular patterns. ChemBioChem 18:772–781.

75.   Newman MA, Daniels MJ, Dow JM. 1995. Lipopolysaccharide from *Xanthomonas campestris* induces
defense-related gene expression in *Brassica campestris*. Mol Plant-Microbe Interact 8:778–80.

76.   Barton-Willis PA, Wang MC, Holliday MJ, Long MR, Keen NT. 1984. Purification and composition of
lipopolysaccharides from *Pseudomonas syringae*. pv. *glycinea*. Physiol Plant Pathol 25:387–398.

77.   Erbs G, Newman M-A. 2003. The role of lipopolysaccharides in induction of plant defence responses. Mol
Plant Pathol 4:421–425.

78.   Leeman M, Pelt JA van, Ouden FM den, Heinsbroek M, Bakker PAHM, Schippers B. 1995. Induction of
systemic resistance against *Fusarium wilt* of radish by lipopolysaccharides of *Pseudomonas fluorescens*.
Phytopathol 85:1021–1027.

79.   Van Wees SCM, Pieterse CMJ, Trijssenaar A, Van 't Westende YAM, Hartog F, Van Loon LC. 1997.
Differential induction of systemic resistance in *Arabidopsis* by biocontrol bacteria. Mol Plant-Microbe
Interact 10:716–724.

80.   Nakao R, Senpuku H, Watanabe H. 2006. *Porphyromonas gingivalis galE* is involved in lipopolysaccharide
O-antigen synthesis and biofilm formation. Infect Immun 74:6145–53.

81.   Ramette A, Frapolli M, Saux MF-L, Gruffaz C, Meyer J-M, Défago G, Sutra L, Moënne-Loccoz Y. 2011.
Pseudomonas protegens sp. nov., widespread plant-protecting bacteria producing the biocontrol
compounds 2,4-diacetylphloroglucinol and pyoluteorin. Syst Appl Microbiol 34:180–188.

82.   Stutz EW, Défago G, Kern H. 1986. Naturally occurring fluorescent pseudomonads involved in suppression
of black root rot of tobacco. Phytopathology 76:181–185.

83.   Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012. A program for
annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of
Drosophila melanogaster strain *w1118; iso-2; iso-3*. Fly (Austin) 6:80–92.

84.   Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics 26:841–842.

85.   Højberg O, Schnider U, Winteler H V, Sørensen J, Haas D. 1999. Oxygen-sensing reporter strain of
Pseudomonas fluorescens for monitoring the distribution of low-oxygen habitats in soil. Appl Environ
Microbiol 65:4085–4093.

86.   Glickmann E, Dessaux Y. 1995. A critical examination of the specificity of the salkowski reagent for indolic
compounds produced by phytopathogenic bacteria. Appl Environ Microbiol 61:793–796.

87.   Alexander DB, Zuberer DA. 1991. Use of chrome azurol S reagents to evaluate siderophore production by
rhizosphere bacteria. Biol Fertil Soils 12:39–45.

88.   Oberhänsli T, Défago G, Haas D. 1991. Indole-3-acetic acid (IAA) synthesis in the biocontrol strain CHA0 of
Pseudomonas fluorescens: role of tryptophan side chain oxidase. Microbiology 137:2273–2279.

89.   Moskowitz SM, Foster JM, Emerson J, Burns JL. 2004. Clinically feasible biofilm susceptibility assay for
isolates of *Pseudomonas aeruginosa* from patients with cystic fibrosis. J Clin Microbiol 42:1915–1922.

90.   Christensen PJ, Cook FD. 1972. The isolation and enumeration of cytophagas. Can J Microbiol 18:1933–
1940.

91.   Murtagh F, Legendre P. 2014. Ward's hierarchical agglomerative clustering method: which algorithms

implement ward's criterion? J Classif 31:274–295.

92.   Scrucca L, Fop M, Murphy TB, Raftery AE. 2016. Mclust 5: clustering, classification and density estimation
using gaussian finite mixture models. R J 8:289–317.

93.   Kai Lee H, Kiat Lee C, Ping Loh T, Wei-Tze Tang J, Anantharajah Tambyah P, Siew-Chuan Koay E. 2011. High-
resolution melting approach to efficient identification and quantification of H275Y mutant influenza
H1N1/2009 virus in mixed-virus-population samples. J Clin Microbiol 49:3555–3559.

94.   Zhong W-L, Wang L, Wu X, Zhang J, Chen X-F, Zhang W, Dou X, Yu B. 2016. Development of unlabeled
probe based high-resolution melting analysis for detection of filaggrin gene mutation c.3321delA. J Clin Lab
Anal 30:892–896.

95.   Capper RL, Jin YK, Lundgren PB, Peplow LM, Matz M V., van Oppen MJH. 2015. Quantitative high resolution
melting: two methods to determine SNP allele frequencies from pooled samples. BMC Genet 16:62.