The indoor environment -

a potent source for intact human-associated anaerobes

Manuela-Raluca Pausan[1,#,±], Marcus Blohs[1,±], Alexander Mahnert[1], Christine

Moissl-Eichinger[1,*]

[1] Diagnostic and Research Institute of Hygiene, Microbiology and Environmental Medicine,

Medical University of Graz, Graz 8010, Austria

[±] Authors contributed equally

*Corresponding author

[#] Current affiliation: Steigerwald Arzneimittelwerk GmbH, Bayer Consumer Health,

Havelstrasse 5, D-64295 Darmstadt, Germany

Abstract

People in westernised countries spend most of their time indoors. A healthy human microbiome relies on the interaction with and exchange of microbes that takes place between the human body and its environment. For this reason, the built environment (BE) might represents a potent source of commensal microbes. Anaerobic microbes are of particular interest, as researchers have not yet sufficiently clarified how the human microbiome acquires oxygen-sensitive microbes, such as obligate or facultative anaerobes.

We sampled a total of ten bathroom floors and used propidium monoazide (PMA) to assess the viability of the collected prokaryotes. Quantitative and qualitative analysis revealed that most of the microbial taxa are of human origin. Less than 25% of the prokaryotic signatures originate from intact – and thus potentially living – cells, indicating that aerobic and stress resistant taxa display an apparent survival advantage. Although the dominant microbial fraction identified on the bathroom floors is composed of aerobes, we confirmed the presence of strictly anaerobic taxa, including methanogenic archaea, in PMA-treated samples. These results show that these taxa have a limited but substantial ability to survive in the BE. As methanogens are regarded as highly sensitive to aerobic conditions, oxygen-tolerance experiments were performed with human-associated isolates to validate their survival. We determined that human-associated methanogens can survive oxic conditions for at least 6 h.

This study enabled us to collect strong evidence that supports the hypothesis that obligate anaerobic taxa can survive in the BE for a limited amount of time. This suggests that the BE serves as a potential source of anaerobic human commensals.

Background

The development of a healthy human microbiome depends on extensive microbial transmission from other hosts, food, water, but also reservoirs in the built environment (BE; Browne et al., 2017). Especially for newborn infants, the acquisition of microbes is a crucial process, as they represent an empty niche for a variety of microorganisms. In this way, they differ from adults, who usually have well-established, complex and stable microbiomes. A fairly stable microbiome is normally achieved by the third year after birth (Stewart et al., 2018) and, up until this point, numerous microorganisms are acquired from other microbiomes. Various factors affect this maturation process, including its speed and quality. These factors include the birth mode (Caesarean section vs. vaginal birth), type of feeding (breast-fed vs. formula-fed), medication (e.g. antibiotics), time point of introduction of solid food, quality of social interactions and contact to animals and nature (Guaraldi & Salvatori, 2012; Moossavi & Azad, 2020; Pasolli et al., 2019; Rocchi et al., 2020). During the maturation of the intestinal microbiome, facultative anaerobic microorganisms, which initially thrive in the gastrointestinal tract (GIT) after birth, are subsequently replaced by obligate anaerobes. The latter support the optimal fermentation of solid food (Browne et al., 2017).

The abiotic BE together with its indoor microbiome (reviewed by Chase et al., 2016; Karen C. Dannemiller, 2019; Gilbert & Stephens, 2018; Horve et al., 2020; Kelley & Gilbert, 2013) is considered to have an important role in the development and maintenance of a healthy human microbiome (Browne et al., 2017). Populations in urban areas of high-income countries currently spend on average 90% of the day indoors; therefore, the abiotic BE and its indoor microbiome represent one of the major reservoirs of environmental microbes (Klepeis et al., 2001). This reservoir plays a vital role in feeding and shaping the human microbiome early in life (see A. W. Brooks et al., 2018; B. Brooks et al., 2014), but it also likely contributes to recolonisation after disease and infection or antibiotic treatment (Ng et al., 2019; Parajuli et al., 2020).

Cross-talk between the human being and the BE can have both positive and negative aspects.

On the one hand, indoor microbes could trigger immediate health issues under rare

circumstances. This aspect is of particular concern in the course of pathogen and resistance

transmission in hospital environments (Lax, Sangwan, et al., 2017; Mahnert et al., 2019; Mora

et al., 2016). On the other hand, several studies have pointed out the beneficial effects of a

healthy indoor microbiome and presented the observation that the presence of animal-

associated and farm-like indoor microorganisms in rural areas is linked to decreased asthma

and allergy risk (K. C. Dannemiller et al., 2014; Ege et al., 2012; Kirjavainen et al., 2019).

For microbes to be transmitted successful between hosts via the built environment, they have

to be able to survive outside of their natural habitat ("ex-host" survivability) (Browne et al.,

2017; Costello et al., 2012). The majority of human commensals are subject to a number of

different stressors, including UV radiation and desiccation (Browne et al., 2017). For gut

commensals, the exposure to atmospheric oxygen is a critical factor, as an estimated 99% of

all microorganisms that thrive in the distal colon are obligate anaerobes (Eckburg et al., 2005).

The extent of aero-tolerance among species varies, and strictly anaerobic taxa are usually

unable to actively replicate outside of the gastrointestinal tract (GIT; Ley et al., 2008; Meehan

& Beiko, 2014). To overcome this limitation, numerous (anaerobic) bacteria protect

themselves through sporulation, as this provides them with the opportunity to escape difficult

environmental situations for an extended period of time. Researchers believe that up to 30%

of all intestinal microorganisms can form spores, including *Lachnospiraceae*,

Ruminococcaceae, *Clostridiaceae* and *Peptostreptococcaceae* (Browne et al., 2017). After

being transferred to their new environment, spores react to trigger signals (such as bile acids)

by germinating, spreading and colonising in the new host thereafter (Browne et al., 2017).

Non-spore formers, however, must have other strategies that allow them to tolerate

environmental stress factors. Strictly anaerobic microorganisms such as *Roseburia* usually

die within few minutes of exposure to atmospheric oxygen, as they have not evolved

mechanisms that enable them to avoid and repair the damage caused by reactive oxygen

species (ROS); i.e. they lack catalases, peroxidases, or superoxide dismutases (Miller &

Britigan, 1997; Rolfe et al., 1978). Thus, the survival strategies and transmission routes of

most of the strictly anaerobic microbiome members, including methanogenic archaea, are

largely unknown.

Methanogenic archaea ("methanogens") represent important components of the human GIT,

along with bacteria, viruses and small eukaryotes (Maria Chibani et al., 2020). These archaea

are obligate anaerobic microorganisms and lack the ability to form spores. Despite their

oxygen sensitivity, the most common representative, *Methanobrevibacter smithii*, can be

detected in almost 96% of the adult population (Dridi et al., 2009). Interestingly, *M. smithii* is

rarely found in younger cohorts (Palmer et al., 2007), and especially uncommon in children

born via Caesarean section (Wampach et al., 2017).

In this communication, we address a major question: Can the BE be a potent reservoir for

anaerobic, commensal microorganisms? To answer this question, we analysed the indoor

microbiome of ten households and compared the microbiome data with the human microbiome

data. As the bathroom represents the richest source of GIT-derived microorganisms (Browne

et al., 2017; Dunn et al., 2013), we sampled surfaces near toilets in different households. One

major shortcoming of previous studies was that no distinction was made between viable

microorganisms (i.e. those with the potential to colonise a new host) and non-viable

microorganisms (Browne et al., 2017). We addressed this knowledge gap by combining a

propidium monoazide (PMA) treatment with high-throughput 16S rRNA gene sequencing. This

procedure enabled us to identify the signatures of intact, and thus probably living,

microorganisms. As archaeal cells were frequently detected amongst the surviving proportion

of the BE microbiome, we analysed the BE archaeome in more detail. The results of this

analysis show that human-associated methanogens can survive air exposure and, therefore,

could be transferred to another human microbiome through the BE.

Material and methods

Sample collection and DNA extraction

Samples were collected from ten different family houses in the vicinity of Graz, Austria (in March 2017). All houses were occupied by at least one adult and five of the houses by families with children. The sampling surface was selected by considering the richest source of GIT-derived microorganisms (Dunn et al., 2013), namely the bathroom. Per household, two areas of 30 cm^2 in the proximity of the toilet (example shown in Supplementary Fig. 1a) were cleaned with bleach and sterile water and left uncovered and untouched for 7 days. After 7 days, the areas were sampled using a sterile nylon swab (FLOQSwabsTM, Copan, Brescia, Italy) that has been dipped in 0.9% saline (NaCl) solution. Each area was sampled with the swab three times, by rotating the swab every time before sampling the area again (Supplementary Fig. 1b).

The samples were transported back to the laboratory on ice packs on the day of the sampling. One of the two samples from each house, which were collected in parallel, was treated with PMA (propidium monoazide) as previously described (Mora et al., 2016) to differentiate between cells with an intact membrane. PMA is a photo-reactive DNA binding dye which intercalates with free DNA inhibiting downstream DNA amplification thereof. As PMA is not permeable for intact cell walls, a molecular distinction between PMA-masked free DNA from disrupted, most probably dead cells, and DNA from intact, most probably living cells can be made (Nocker et al., 2007). In the following, we use the term "PMA" to indicate samples treated with PMA (= intact cells only), and the term "non-PMA" for samples not treated with PMA (= both, intact and dead cells/free DNA).

For genomic DNA extraction, the indoor samples were processed using the FastDNA Spin Kit (MP Biomedicals, Germany) according to manufacturer's instructions. The genomic DNA was

140 eluted in 100 µL elution buffer, and the concentration was measured using Qubit™ dsDNA HS

141 Assay Kit (ThermoFisher Scientific).

PCR and qPCR

The obtained genomic DNA was used to amplify the V4 region of the 16S rRNA gene using

Illumina-tagged primers 515F and 806R (Supplementary Tab. 1). In order to identify the

archaeal communities present in the samples, a nested PCR was performed using the primer

combination 344F-1041R/Illu519F-Illu806R as described previously (Pausan et al., 2019). The

PCR reaction was performed in a final volume of 25 µL containing: TAKARA Ex Taq® buffer

with $MgCl_2$ (10 X; Takara Bio Inc., Tokyo, Japan), primers 200 nM, dNTP mix 200 µM,

TAKARA Ex Taq® Polymerase 0.5 U, water (Lichrosolv®; Merck, Darmstadt, Germany), and

DNA template (1-2 µL of genomic DNA). The PCR amplification conditions are listed in

Supplementary Tab. 2.

The bacterial and archaeal 16S rRNA gene copies were determined using SYBR based

quantitative PCR (qPCR) with the primer pair Bac331F-Bac797R and A806f-A958r,

respectively (Supplementary Tab. 1). The reaction mix contained: 1x SsoAdvanced™

Universal SYBR® Green Supermix (Bio-Rad, Hercules, USA), 300 nM of forward and reverse

primer, 1 µL gDNA template, PCR grade water (Lichrosolv®; Merck, Darmstadt, Germany).

The qPCR was performed in triplicates using the CFX96 Touch™ Real-Time PCR Detection

System (Bio-Rad, Hercules, USA). The qPCR conditions are given in Supplementary Tab. 2.

Crossing point (Cp) values were determined using the regression method within the Bio-Rad

CFX Manager software version 3.1. Absolute copy numbers of bacterial 16S rRNA genes were

calculated using the Cp values and the reaction efficiencies based on standard curves

obtained from defined DNA samples from *Escherichia coli* and *Nitrososphaera viennensis*

(Probst et al., 2013). The qPCR efficiency was between 85-105%, and the R^2 values were

always above 0.9. Detection limits were defined based on the average Cp values of non-

template controls (triplicates) and the corresponding standard curves of the positive controls.

NGS-based 16S rRNA gene sequencing

Library preparation and sequencing of the amplicons were carried out at the Core Facility

Molecular Biology, Center for Medical Research at the Medical University Graz, Austria. In

brief, DNA concentrations were normalised using a SequalPrep™ normalisation plate

(Invitrogen), and each sample was indexed with a unique barcode sequence (8 cycles index

PCR). After pooling of the indexed samples, a gel cut was carried out to purify the products of

the index PCR. Sequencing was performed using the Illumina MiSeq device and MS-102-

3003 MiSeq® Reagent Kit v3-600cycles (2x150 cycles). The obtained fastq data is available

in the European Nucleotide Archive under the study accession number: PRJEB41618.

The fastq data analysis was performed using QIIME2 (Bolyen et al., 2019) as described

previously (Mahnert et al., 2020). After quality filtering, the DADA2 algorithm (Callahan et al.,

2016) was used to denoise truncated reads and generate amplicon sequence variants (ASVs).

Taxonomic classification (Bokulich et al., 2018) was based on the SILVA v132 database

(Quast et al., 2013) and the obtained feature table and taxonomy file were used for further

analysis (Supplementary Tab. 3). The overlapping features from negative controls (DNA

extraction and PCR negative controls) were manually subtracted or removed from both the

bacterial and archaeal dataset. The reads classified as chloroplast and mitochondria were

also removed.

Human microbiome data

In order to assess to which extent the microbial community indoors is affected and affects the

human microbiome, a representative dataset of the human microbiome from several body

sites was collected. We took advantage of several previous in-house projects, covering the

microbiome of skin, nasal cavity, saliva, urine, vagina, and stool (for more details, please see:

Supplementary Tab. 4). All samples were taken from healthy participants and processed in

our lab with methods similarly to the current study: DNA was extracted by a combination of

enzymatic and mechanical lysis; amplification of the 16S rRNA gene was done using the

"universal" primers based on Caporaso et al. (2011) or Walters et al. (2016) and library

preparation, as well as sequencing, were performed by the Core Facility Molecular Biology at

the Medical University Graz, Austria. Obtained raw reads were processed in parallel to the

indoor microbiome data of the current study (Supplementary Tab. 5 and 8).

Data analysis and statistics

In general, data analysis of the microbial data is based on the indoor data alone

(Supplementary Tab. 3 and 8). Comparative analysis between indoor and human microbiomes

are based on the mixed dataset (Supplementary Tab. 5 and 9). Bar charts, box plots, and

bubble plots are based on relative abundances and were constructed for both the bacterial

and archaeal communities at different taxonomic levels using the *phyloseq* (McMurdie &

Holmes, 2013) and *ggplot2* package in R (Wickham, 2016). Significant differences between

microbial taxa were calculated in R using Wilcoxon signed-rank tests for dependent samples

(e.g. PMA and non-PMA) or Mann–Whitney *U* test for independent samples (e.g. indoor and

human). P values were adjusted for multiple testing using the method after Benjamini &

Hochberg (1995). The online-tool Calypso (Zakrzewski et al., 2017) was used to calculate

alpha diversity metrics, Linear discriminant analysis Effect Size (LEfSe), the bar plots and

principal coordinates analysis (PCoA) plots based on Bray-Curtis. Before analysis in Calypso,

the data was normalised using total sum normalisation (TSS) combined with square root

transformation. For alpha diversity, reads were rarefied to 5,588 and analysed based on

Shannon and richness indices. To test for differences in the beta diversity between sample

categories, PERMANOVA analysis was performed in QIIME2 based on Bray-Curtis distances.

BugBase (Ward et al., 2017) was used to predict potential phenotypes such as aerobic,

anaerobic, facultative anaerobic and stress-tolerant communities which were present in the

PMA and non-PMA treated samples. SourceTracker2 analysis (Knights et al., 2011) was

performed with default settings. Sampling depth was set to 1,000 reads for source and sink,

respectively.

A phylogenetic tree was constructed to determine if the sequences belonging to methanogens

identified in the analysed samples are of human origin. All sequences classified within the

genera *Methanobacterium*, *Methanobrevibacter* and *Methanomassiliicoccus* from our

analysed samples were used for creating the phylogenetic tree. Additionally, 16S rRNA gene

sequences of species of *Methanobacterium*, *Methanobrevibacter* and *Methanomassiliicoccus*

previously identified in the human body or the environment were included from NCBI and two

other studies (Borrel et al., 2017; Pasolli et al., 2019). The alignment was performed using the

SILVA SINA alignment tool (Pruesse et al., 2012). The sequences were cropped to the same

length using BioEdit and used afterwards to construct a tree based on the maximum-likelihood

algorithm with a bootstrap value of 1,000 using MEGA7 (Kumar et al., 2016). The phylogenetic

tree was further processed using the online tool iTOL (Letunic & Bork, 2007).

Definition of obligate anaerobic taxa

Bacterial and archaeal taxa were annotated as obligate anaerobic on family and genus level

and marked with an asterix *. The categorisation is based on the "list of prokaryotes according

to their aerotolerant or obligate anaerobic metabolism" V1.3 (Million & Raoult, 2018) and

literature review.

Cultivation and oxygen tolerance test

An oxygen sensitivity test was implemented to determine if methanogenic archaeal strains can

survive under aerobic conditions for a certain period of time. Three human-associated

methanogenic strains (*Methanosphaera stadtmanae* DSM no. 3091, *Methanobrevibacter*

smithii DSM no. 2375, and *Methanomassiliicoccus luminyensis* DSM no. 25720) and a newly

isolated strain (*Methanobrevibacter* sp., unpublished) have been tested for their ability to

survive under aerobic conditions. The human-associated methanogen cultures were obtained

from DSMZ and both, *M. stadtmanae* and *M. luminyensis* were grown in MPT1 medium at

37 °C (Mauerhofer et al., unpublished). The new *Methanobrevibacter* strain was isolated in

our lab from a fresh stool sample using ATCC medium: 1340 MS medium for methanogens

(https://www.atcc.org/). Both *Methanobrevibacter* strains were grown in MS medium at 37°C.

Cultures were used after 3-7 days of growth as follows: 1 mL culture was transferred to sterile

1.5 mL Eppendorf tubes and centrifuged at 10,000 x g for 10 min, the cell pellet was washed

twice either anaerobically for the controls or aerobically for the oxygen exposed

microorganisms with 1 mL sterile 1x PBS and centrifuged again at 10,000 x g for 10 min. After

removing most of the PBS, the cell pellet was resuspended in the remaining liquid and

exposed to aerobic or anaerobic conditions for different time points: 0 h (30 minutes after

transfer), 6 h, 24 h, 48 h, and 168 h (7 days). After exposure, the liquid from the tubes was

transferred to a sterile growth medium in Hungate tubes and the cultures were left to grow for

2-3 weeks. Then, the first optical density (OD) measurements (600 nm) and methane

measurements were done using a spectrometer and methane sensor (BCP-CH4 sensor,

BlueSens). For the methane measurement, 10 mL of gas-phase was taken from the Hungate

tubes using an air-tight glass syringe. For each experiment, two sterile unopened Hungate

tubes with media served as control, and two inoculated Hungate tubes with 0.5 mL of the

culture served as a positive control. In addition, a tube of 1x PBS served as a control for 168h.

Each experiment was performed in duplicates. After OD and methane measurements,

microscopic observations were performed to determine the shape of the microorganisms

growing in the medium. Oxygen saturation of the PBS was monitored in a control experiment

using the FireStingO2 optical oxygen meter (PyroScience GmbH) and the oxygen probe

OXROB10. Using the identical methodology as described above we were able to confirm an

oxygen saturation in the medium of > 80% at any given time point (0 h, 10 min, 20 min, 30

min, 45 min, 1 h, 2 h, 4 h, 6 h and 24 h) for every strain.

Results

In this section, we present information about the anaerobic microbial communities detected in the bathrooms of the selected family houses. We wanted to determine whether the indoor environment serves as a source of anaerobic, commensal microorganisms, such as methanogens, which could potentially colonise the human body. To achieve this purpose, we collected samples from two surface areas (~30 cm^2) in the bathrooms of ten households.

One of the two samples collected from the BE was treated with PMA in order to mask the DNA of disrupted cells (Nocker et al., 2007). By comparing non-PMA (untreated) and PMA-treated samples, we could distinguish the overall (non-PMA) microbiome from the intact (i.e. probably alive) microbiome (PMA). This comparison enabled us to draw conclusions regarding the survival of strictly anaerobic microorganisms under exposed conditions. These findings were further supported by the results of subsequent experiments.

Although we present data on all detected microbial signatures, we focussed specifically on strictly anaerobic taxa. For convenience, we highlight taxa that mainly consist of strict anaerobes with an asterisk *.

The human microbiome is the predominant source of microbes in the BE

Using a universal 16S rRNA gene sequencing approach, it was possible to obtain overall more than 380,000 sequences (Bacteria: 99.86%, Archaea: 0.14%), corresponding to 3,684 ASVs and an average of 473 ± 110 ASVs per household (285 ± 50 and 245 ± 80 ASVs in non-PMA and PMA samples, respectively; Supplementary Tab. 3). We observed a high amount of heterogeneity amongst households, with only 26 shared ASVs (0.7%) amongst all households, and 351 ASVs (9.5%) that were shared between at least two households (Supplementary Tab. 6). The 26 ASVs present in all samples include typical skin-associated taxa, such as

Staphylococcus (11 ASVs) and *Corynebacterium* (8 ASVs), but also *Finegoldia** (6 ASVs) and

Peptoniphilus* (1 ASV).

Fig. 1: Distribution and relative abundance of bacterial taxa in samples from different households/bathrooms. Bar charts showing the microbial composition of non-PMA- and PMA-treated bathroom floor samples at the **(a)** phylum and **(b)** genus levels. Genera with < 2% rel. abundance are summarised in grey. Bubble plots display the 25 most abundant genera (bubble size reflects the relative abundance): **(c)** Household samples (H1–H10) are depicted individually for non-PMA and PMA treatment together with human samples (grey background) that represent reads from several body sites: nasal cavity, skin, vagina, urine, stool and oral samples. **(d)** Human samples compared to non-PMA (blue background) and PMA (red background) bathroom samples. Genera are coloured according to their taxonomic phylum, and taxa that predominantly contain strict anaerobes are marked by an asterisk.

A total of 32 phyla were observed, and most sequences were classified to four phyla: Firmicutes (52.2% non-PMA, 41.2% PMA), Actinobacteria (20.3% non-PMA, 26.0% PMA),

Proteobacteria (18.8% non-PMA, 23.3% PMA) and Bacteroidetes (6% non-PMA, 4.3% PMA) (Fig. 1a). The distribution of sequences classified to these phyla did not change significantly upon PMA treatment. The Linear discriminant analysis Effect Size (LEfSe) method and Wilcoxon signed-rank test were performed, revealing significant differences only in four low-abundance phyla, namely, an increase (Actinobacteria) or decrease (Tenericutes, Fusobacteria, Epsilonbacteraeota) in relative abundance upon PMA treatment ($P < 0.05$, Supplementary Fig. 2).

Genera associated with the human skin (e.g. *Staphylococcus, Corynebacterium, Streptococcus, Neisseria, Micrococcus, Cutibacterium*), gastrointestinal and genitourinary tract (e.g. *Lactobacillus, Finegoldia*, Bacteroides*, Anaerococcus*, Peptoniphilus*,* Lachnospiraceae*) were identified in both non-PMA and PMA samples (Fig. 1c). Notably, *Bifidobacterium*, Rothia, Bacteroides*, Blautia*, Dialister*, Faecalibacterium*, Gemella, Subdoligranulum** and *Haemophilus* were detected in every single household, but not in all PMA samples, indicating a potential impairment by environmental stressors.

Due to the high proportion of common human-associated commensals, we compared the BE microbiome data collected in this study with human microbiome data that were collected in other studies performed by our lab (see material and methods and Supplementary Tab. 4). The results of a bubble plot analysis of the most abundant genera (Fig. 1d) revealed a specific pattern with respect to the abundance of particular commensals in BE samples or human samples, retrieved from the nasal and oral cavity, skin, stool and vaginal samples. In particular, we observed that signatures of typical human commensals such as *Corynebacterium, Anaerococcus*, Dolosigranulum, Finegoldia*, Gardnerella* and *Staphylococcus* were highly enriched in non-PMA indoor samples as compared to in human samples ($P_{adj} < 0.001$; Mann–Whitney *U* test), indicating that an efficient transfer to indoor surfaces had occurred.

With respect to alpha diversity, the non-PMA indoor samples showed high Shannon indices, which were comparable to stool samples and even exceeded stool samples in richness by

more than two-fold (mean richness: 282 and 136 for non-PMA indoor and stool samples,

respectively; see Fig. 2). A beta diversity analysis confirmed that all analysed body sites

grouped separately ($P_{adj} < 0.01$, PERMANOVA), except for urine and vaginal samples ($P_{adj} =$

0.81; Fig. 2a). However, we did not observe a grouping of indoor samples with a specific

human body site, as the clusters differed significantly from one another ($P_{adj} < 0.01$).

Fig. 2: A comparison between the indoor and human microbiome. (a) Principal coordinates analysis plots

based on Bray-Curtis dissimilarity and **(b)** alpha diversity indices (Shannon, left; richness, right) are depicted for all

indoor samples (blue = untreated, red = PMA treated) together with representative human body site samples (green

= nasal cavity, light blue = oral, yellow = skin, brown = stool, pink = urine, purple = vaginal). Significant differences

To determine the level of impact of the human microbiome on our indoor samples, we performed a SourceTracking 2 analysis (Knights et al., 2011). Independent of the PMA treatment, about 75% of the indoor microbiome was associated with human sources. These findings indicate that the human body was the dominant source of the microbes detected in bathroom samples (Fig. 3a,b). Nevertheless, depending on the household, we observed large fluctuations in the proportion of human-associated taxa. Especially in PMA samples, this proportion ranges from 41–89%. This may be indicative of a large variation in the frequency and duration of exposure of the tested surfaces to a human being.

Upon PMA treatment, we observed a significant reduction in the vaginal, stool, and oral signatures in the surface samples, while the proportion of microbes from human skin significantly increased ($P < 0.05$, Wilcoxon signed-rank test; see Fig. 3d). These results indicate that skin-related microbes possess the greatest ability to survive in the bathroom environment.

Vice versa, microbial traces from the indoor environment could potentially serve as a source for the human microbiome as well (Fig. 3c). As has been identified in other studies, we identified that the highest level of exchange occurred between human skin and the indoor and human microbes, whereby the skin shared on average 29% and 32% of the microbial signatures with non-PMA and PMA samples from the bathroom. The overlap with other body sites was generally very low, and especially PMA-treated samples shared on average only about 6% of the taxa with urine, 5% with nasal cavity, 4% with vaginal and < 0.01% with oral

and stool samples. Therefore, the impact of the bathroom microbiome on human body sites,

with the exception of the skin, seems to be very small or even negligible.

Oxygen-tolerance primarily determines microbial survival indoors

Human commensals are exposed to several stress factors in the BE; therefore, many of them

are perceived as incapable of surviving in this environment over extended periods. We

performed PMA treatment to mask DNA from disrupted cells, and in fact, the absolute

abundance was significantly affected by PMA treatment (Fig. 3a). Results of the quantitative

PCR reveal a 4.2-fold decrease in microbial 16S rRNA gene copy number (Fig. 4c), indicating

that less than 25% of all microbial signatures originate from intact, probably living cells. This

is also reflected by the significant drop in both of the alpha diversity parameters in PMA

samples as compared to those in untreated indoor samples (Shannon: $P_{adj} = 0.007$, richness:

$P_{adj} = 0.0006$; Wilcoxon rank-sum tests, Fig. 2b). PCoA plots based on Bray-Curtis distances

also showed a significant separation based on the PMA treatment ($P_{adj} = 0.018$) even though

indoor samples were more similar to one another than to any human body site (Fig. 2a).

LEfSe was performed on PMA and non-PMA samples to determine which taxa are indicative

for PMA samples and, therefore, likely to survive in the indoor environment (Supplementary

Fig. 3). Representative taxa for PMA were found to be predominantly environmental and

aerobic (or aerotolerant) groups, such as *Acidobacteria*, *Micromonosporales*,

Solirubrobacterales, Rhizobiales, Nocardiaceae, Bacillus and *Sphingomonas,* but also

included human skin commensals, such as *Corynebacterium*_1, *Staphylococcus*, or

*Rombustia**. On the other hand, representative taxa for non-PMA samples were more diverse

in terms of their physiological traits and origins. They predominantly included (facultatively)

anaerobic, non-spore forming, human-associated groups, such as *Bacteroidales**,

Lactobacillaceae, multiple genera in the families *Lachnospiraceae** and *Ruminococcaceae**,

*Bifidobacteriaceae**, as well as in Christensenellaceae*, Veillonellaceae*, *Haemophilus*,

Neisseria and the *Eubacterium coprostanoligenes* group*. Interestingly, some opportunistic

pathogens (*Atopobium*, *Rothia* and *Campylobacter*) were also characteristic for and

significantly enriched in the non-PMA samples (Supplementary Fig. 4), which suggests that

these taxa might not survive on BE surfaces over extended periods of time.

Fig. 3 The effects of PMA treatment on prokaryotic abundance and composition. (a) Quantitative PCR
analysis of untreated and PMA-treated bathroom floor samples using universal and archaeal primer pairs. **(b)**
Wilcoxon signed-rank test analysis results, revealing the relative abundance of the 50 and 100 most abundant
bacterial **(b)** families and **(c)** genera, respectively, that were significantly affected by PMA treatment. Strict
anaerobic taxa are marked by an asterix. Sign.: FDR-adjusted Wilcoxon signed-rank test; *, $P_{adj} < 0.05$; ***, $P_{adj} <$
0.01. **(d)** The relative abundance of different phenotypes as determined through BugBase. Significances were
automatically defined by Mann-Whitney *U* test in BugBase.

After PMA treatment, we observed a significant change in the relative abundance of 53 taxa

among the most abundant bacterial genera and families ($P < 0.05$, $n_{genus} = 100$, $n_{family} = 50$,

Wilcoxon signed-rank test, Supplementary Tab. 7), although only 10 taxa remained significant

after performing an adjustment for multiple testing ($P_{adj} < 0.05$, Fig. 3b,c). In accordance with

our LEfSe analysis results, a strong reduction was seen in non-spore forming, (facultative)

anaerobic human commensals, and particularly for *Lactobacillus*, *Streptococcus*, *Gemella* and

the *Eubacterium coprostanoligenes* group*. *Ruminococcaceae* and several genera within this

family were also strongly affected by PMA-treatment and seemed to be susceptible to stress

factors in the BE, whereas taxa in the family *Acetobacteraceae* and other commensals such

as *Bacillus* and *Staphylococcus* appear to be extraordinarily well-suited for survival under BE

conditions.

BugBase (Ward et al., 2017) was used to validate the observed phenotypes for non-PMA and

PMA samples (Fig. 3d). As compared with non-PMA samples, the PMA samples showed an

increase in the relative abundance of aerobic ($P_{adj} = 0.089$) and facultative anaerobic

communities ($P_{adj} = 0.393$), and a slight decrease in the abundance of anaerobic communities

($P_{adj} = 0.217$) (Fig. 3d). Interestingly, the abundance of taxa with stress-tolerant phenotypes

increased in the PMA samples ($P_{adj} = 0.063$), indicating that microorganisms that could survive

in the indoor environment might be better equipped to deal with BE-associated stress factors.

Archaea are an integral component of the BE

An archaea-centric approach (Pausan et al., 2019) was taken to explore the archaeal

communities present on the sampled surfaces. Overall, we obtained about 376,000 archaeal

sequences in both PMA and non-PMA samples, which corresponded to 373 ASVs (on average

70.2 ± 22.8 ASVs/ house; 53.6 ± 16.8 ASVs/ non-PMA samples; 35.9 ± 22.1 ASVs/ PMA

samples; Supplementary Tab. 8). The archaeal ASVs were classified into four different phyla,

namely, Euryarchaeota (36.6% relative abundance in non-PMA samples, PMA: 35.0%),

Thaumarcheota (non-PMA: 62.2%, PMA: 64.1%), Crenarcheota (non-PMA: 0.8%, PMA:

0.5%) and Nanoarchaeota (non-PMA: 0.4%, PMA: 0.5%) (Fig. 4a).

On a family level, most reads were classified into the *Nitrososphaeraceae* (non-PMA: 62.1%,

PMA: 64.1%), which is also the only archaeal family that was found in every household

included in this study (Fig. 4c). In the *Nitrososphaeraceae*, the dominant genus identified was

Candidatus Nitrosocosmicus (non-PMA: 22.5%, PMA: 15.9%), but sequences classified within

Candidatus Nitrososphaera (non-PMA: 0.9%, PMA: 0.2%) were also identified (Fig. 4b). The

second-most abundant family was *Halococcaceae* (non-PMA: 19.7%, PMA: 28.7%), with

Halococcus (non-PMA: 13.7%, PMA: 28.6%) identified as the predominant genus. One

particularly fascinating finding was that oxygen-sensitive methanogenic taxa were detected in

every household, but at a low relative abundance (*Methanobacteria** non-PMA: 3.7%, PMA:

3.5%; *Methanomicrobia** non-PMA: 4.0%, PMA: 2.6%), with *Methanosarcina** (non-PMA:

3.2%, PMA: 1.8%), *Methanobrevibacter** (non-PMA: 2.4%, PMA: 0.2%), *Methanobacterium**

(non-PMA: 1.3%, PMA: 3.3%) and *Methanomassiliicoccus** (non-PMA: 0.4%, PMA: 0.2%)

identified as the most abundant genera.

Humans are potentially the source of most archaeal taxa in the BE

The archaeal community composition suggests that some of the archaeal signatures identified

on the bathroom surface could be of human origin, as *Thaumarchaeota*, *Haloarchaea* and

methanogens have been previously associated with the human body (Borrel et al., 2020).

Thus, we analysed the data obtained from these surfaces together with data obtained by

applying the same archaea-centric approach to samples collected from the human body,

namely, to stool ($n = 38$), urine ($n = 43$), nasal ($n = 30$), oral ($n = 26$), vaginal ($n = 16$) and skin

($n = 7$) samples (see Supplementary Tab. 4 and 8).

We recognised that indoor and human samples can be distinguished by their abundance and

prevalence of *Thaumarchaeota* and *Euryarchaeota*, as *Thaumarchaeota* are highly enriched

Fig. 4. Archaeal community composition in indoor samples. The relative abundance of archaeal taxa among 16S rRNA gene reads from bathroom floor surfaces at the **(a)** phylum and **(b)** genus levels. Bubble plot showing relative abundances of the 25 most-abundant archaeal genera found in bathroom and human samples: **(c)** Household samples (H1-H10) are depicted individually for non-PMA and PMA treatment of human samples (grey background) that represent reads from several body sites including nasal cavity, skin, vagina, urine, stool and oral samples. **(d)** Bubble plots of the 25 most-abundant archaeal genera, displayed according to their original samples. Relative abundance is reflected by the size of the bubbles. Genera are coloured according to their taxonomic phylum, and taxa that predominantly contain strict anaerobes are marked by an asterisk. Venn diagram of shared ASVs in **(e)** non-PMA indoor and **(f)** PMA indoor samples compared to nasal, oral, skin and stool samples (body sites that show the highest numbers of shared ASVs).

in indoor samples (P_{adj} = 3.5*10^{-11}; see Supplementary Fig. 5; Mann–Whitney U test), and *Euryarchaeota* are highly enriched in human samples (P_{adj} = 1.3*10^{-5}). This especially applies to signatures from *Candidatus Nitrososphaera* (P_{adj} = 2.9*10^{-26}) and *Cand. Nitrosocosmicus* (P_{adj} = 1.1*10^{-24}) that were rarely abundant in samples from the human body, but were often found in the analysed indoor samples. In addition, the relative abundance of unassigned *Nitrososphaeraceae* that were found in skin, urine and nasal samples was significantly higher (P_{adj} = 1.0*10^{-20}) in indoor samples.

Nevertheless, several archaeal genera were found in both indoor samples and human-associated samples. For example, reads classified as *Methanobacterium**, *Methanobrevibacter**, *Methanosarcina**, *Halococcus*, unassigned *Nitrososphaeraceae* and *Woesearchaeia* were present in both kinds of samples, indicating an overlap and that exchange is occurring between both microbiomes.

Venn diagrams were created to illustrate the number of shared archaeal ASVs between the indoor environments and all analysed body sites. Each sample type contained a certain number of specific ASVs (e.g. nasal samples: 489 ASVs, oral: 41 ASVs, skin: 5 ASVs, stool: 24 ASVs, indoor samples: 267 ASVs). The vaginal samples revealed only a small overlap with other samples with respect to the archaeal ASVs (15 unique ASVs, two shared with urine samples). Notably, nine ASVs (*Methanobrevibacter**) were shared across all sites, and the

skin samples shared the most (20) ASVs with the indoor environment (four *Candidatus Nitrosocosmicus* ASVs and 16 ASVs uncultured *Nitrososphaeraceae*) (Fig. 4e,f).

In the next step, we examined the signatures of methanogenic archaea in more detail, as we had used them as specific model microorganisms in our study. We were interested in determining whether the methanogens identified in the indoor samples were of human origin or were associated with archaeal signatures from the natural environment. Therefore, we first extracted methanoarchaeal sequences from our indoor datasets (PMA and non-PMA) and used these together with reference sequences from different environments that were available from official databases to generate a phylogenetic tree (Fig. 5).

Most of the *Methanobrevibacter* sequences identified in the indoor samples clustered together with host-associated (human (*M. smithii**, *M. oralis**), bovine/ovine (*M. millerae**), or biogas-reactor/anaerobic-digestor-derived sequences, which are believed to be of holobiont origin. Due to the fact that *Methanobrevibacter** is considered to be a strongly host-associated taxon (Maria Chibani et al., 2020), we hypothesise that our detected *Methanobrevibacter** signatures are mostly derived from human body-sites and only rarely from other, non-human environments.

Borrel et al. (2017) showed that nearly all human-associated *Methanomassiliicoccales** DNA sequences clustered in a host-associated clade together with other *Methanomassiliicoccales** sequences identified in animals, with the exception of two taxa, namely *Candidatus Methanomassiliicoccus intestinalis** and *Methanomassiliicoccus luminyensis**. These two taxa they have been identified and isolated from human samples, but they cluster together with sequences of microorganisms that are mainly associated with the soil and sediment environment ("environmental clade"). This clearly shows that some environmental *Methanomassiliicoccales** taxa can be transferred from the environment to the human body and can colonise the human gut. The sequences assigned to the order of *Methanomassiliicoccales** in our samples clustered only in the environment-associated clade and were closer to the *Methanomassiliicoccales** sequences that have been identified in soils,

sediments and anaerobic digesters. Therefore, the external environment appears to be the

most probable source of the *Methanomassiliicoccales** sequences identified in our bathroom

floor samples.

Signatures of *Methanobacterium** are being more closely associated with human origin (Borrel

et al., 2020; Maria Chibani et al., 2020). Regarding *Methanobacterium**, we included

sequences from environment- and human-associated taxa in order to determine whether the

Methanobacterium* sequences identified on the bathroom floor were of human origin or not.

Most *Methanobacterium** sequences identified in our human dataset clustered apart from all

other *Methanobacterium** sequences, with the exception of one ASV. This ASV clustered

together with *Methanobacterium ferruginis** and ASV49, which was identified in the non-PMA

samples. Seven ASVs from the PMA samples clustered with *Methanobacterium oryzae** and

shared a node with the phylogenetic group formed by the human-associated ASVs.

The results of phylogenetic and abundance-based analyses indicate that most of the identified

methanogenic sequences (and particularly *Methanobrevibacter** and *Methanobacterium**) in

the bathroom floor samples are probably indeed of human origin. This especially applies to

the identified sequences in indoor samples.

Human-associated methanogens can survive oxygen exposure for up to 48 hours

Methanogenic archaea are strict anaerobes and are regarded as highly oxygen-sensitive. Nevertheless, in households with methanogenic taxa, we have frequently observed them in both non-PMA and PMA samples, indicating that these taxa display a certain level of tolerance towards stress factors in the BE. Therefore, we performed an experimental analysis to determine whether methanogenic archaea, serving as models for strict anaerobic gastrointestinal microorganisms, were able to survive under aerial oxygen conditions and, therefore, potentially colonise the human body.

Three methanogen strains previously isolated from the human body, namely,

Methanobrevibacter smithii* (DSM no. 2375), *Methanosphaera stadtmanae** (DSM no. 3091),

Methanomassiliicoccus luminyensis* (DSM no. 25720) and one recently obtained isolate

(*Methanobrevibacter sp.**; unpublished) from human faeces were exposed to aerobic and

anoxic conditions over the period of up to 7 days (0 h, 6 h, 24 h, 48 h and 168 h). After

exposure, the cells were transferred to fresh anoxic media and were allowed to grow for 2–3

weeks. Growth was then tested by measuring OD_{600} and methane production. All four strains

could survive in the aerobic environment for at least 6 h (Fig. 6). *Methanomassiliicoccus*

luminyensis* was even able to survive for more than 48 h of exposure to aerobic conditions.

No growth of *Methanomassiliicoccus luminyensis** was detected after 7 d of cultivation, under

either anoxic or oxic conditions, indicating that other negative influences (e.g. starvation)

potentially hindered the normal outgrowth.

Fig. 6. Observed growth of selected human-associated methanogens after exposure to 21% oxygen (air).
Plots indicate the growth as determined by **(a)** OD_{600} measurements and **(b)** methane production of the tested
strains after exposure to an aerobic (red) or anaerobic (blue) environment for different periods of time ($n = 2$).

To measure the genomic capacity of methanogens to resist oxygenic stress, we analysed the available representative genomes of *M. stadtmanae**, *M. smithii**, and *Methanomassiliicoccus** (Tab. 1). We specifically searched for keystone genes, as identified in Lyu and Lu (2018), and used the Microbial Genome Annotation and Analysis Platform (MaGe) for detailed comparative genomics (Vallenet et al., 2006).

Tab. 1. Genes for O$_2$/reactive oxygen species elimination enzymes and their presence in genomes of human-associated methanogens (Lyu & Lu, 2018). **Explanation of letters in brackets: (a)** protein inferred from homology, **(b)** predicted protein, **(c)** putative, **(d)** ferritin-like, **(e)** function from experimental evidence in other organisms, **(f)** ferritin, dps family protein. Specific gene identifiers derived from MaGe microscope database (Vallenet et al., 2006) and UniProtKB (Bateman, 2019) are displayed in each column when present. n.d.: not detected.

Enzyme	*Methanosphaera stadtmanae* DSM 3091	*Methanosphaera stadtmanae* DEW79	*Methanobrevibacter smithii* DSM2374	*Methanobrevibacter smithii* DSM861	*Methanobrevibacter smithii* DSM 2375 (ALI)	*Methanomassiliicoccus* M. luminyensis B10	*Methanomassiliicoccus* M. intestinalis strain Issoire-Mx1
Catalase (EC 1.11.1.6)	n.d.	n.d.	n.d.	n.d.	n.d.	CAJEv1_180010 (e)	MMINT_12710 (a) MMINT_12690 (a)
Peroxiredoxin (EC 1.11.1.24)	n.d.	n.d.	n.d.	n.d.	n.d.	CAJEv1_50026	MMINT_19160 (a) MMINT_05530 (b) MMINT_15900 (b)
Rubrerythrin	Msp_0789 (b)	n.d.	METSMIF1_02444 (b)	Msm_1733 (b) Msm_1348 (b)	METSMIALI_01290 (b)	CAJEv1_240034 (e)	MMINT_05220 (b) MMINT_00570 (b)
Superoxide dismutase (EC 1.15.1.1)	n.d.	n.d.	n.d.	n.d.	n.d.	CAJEv1_120763 (e)	n.d.
Superoxide reductase	n.d.	n.d.	METSMIF1_03626 (a,c)	n.d.	METSMIALI_00110 (a,c)	n.d.	n.d.
F$_{420}$H$_2$ oxidase	n.d.	n.d.	n.d.	Msm_1349 (b)	n.d.	n.d.	n.d.
NO/O$_2$ reductase	n.d.	n.d.	n.d.	n.d.	n.d.	CAJEv1_120660 (e)	n.d.
Ferritin	Msp_0852 (a, b)	n.d.	METSMIF1_03296 (a,d)	Msm_1712 (a, b)	METSMIALI_00411 (a,d)	CAJEv1_120620 (e, f)	MMINT_08340 (a, f)
Thioredoxin	Msp_0103 (a,c) Msp_0820 (a,c)	n.d.	METSMIF1_02203 (a)	Msm_0838 (a, c)	METSMIALI_01538 (a)	CAJEv1_200051 CAJEv1_230019 (c,) CAJEv1_240002	MMINT_17530 (b) MMINT_19530 (b)
Rubredoxin (EC 1.4.3)	Msp_0445 (a) Msp_0444 (a)	n.d.	METSMIF1_03551 (a) METSMIF1_03550 (b) METSMIF1_03395 (a) METSMIF1_03318 (b)	Msm_0187 (b) Msm_0188 (a)	METSMIALI_00388 (b) METSMIALI_00192 (b) METSMIALI_00191 (a) METSMIALI_00344 (a)	CAJEv1_120430 (e)	MMINT_08590 (a)
UniProtKB	UniProtKB	UniProtKB	UniProtKB	UniProtKB	UniProtKB	MAGE	UniProtKB

All investigated methanogens (except DEW79) revealed the presence of rubrerythrin, ferritin, thioredoxin and rubredoxin in their genomic inventory. The extended survival under oxygenated conditions for *Methanomassiliicoccus** was supported by the detection of catalase and peroxiredoxin, as well as superoxide dismutase in the case of *M. luminyensis**.

576 Overall, the experimental and database-derived results show that human-associated archaea

577 have profound capacities to survive under oxygenated conditions for a minimum of 6 hours.

Discussion

The viability of microorganisms that exist in indoor environments is of great interest (Gilbert &

Stephens, 2018), as a substantial number of microorganisms are in constant exchange with

the human body (B. Brooks et al., 2014; Lax et al., 2014; Lax, Smith, et al., 2017; Qian et al.,

2012). These interactions between humans and (viable) microbes are arguably of crucial

importance for training the immune system and sensitisation in infants (Shan et al., 2019), as

well as the recolonisation of the human microbiome after antibiotic intervention or disease (Ng

et al., 2019; Parajuli et al., 2020).

We are aware, that our study results are limited by the fact that a small sample size was used;

the high level of heterogeneity among the different households serves as a source of potential

bias. In this study, we did not analyse other surfaces/house areas (such as showers or sinks,

door handles, dust etc.), which would be of interest in future studies to obtain a better overall

picture (Adams et al., 2015; Dunn et al., 2013). Due to the limitations of amplicon sequencing,

it was not possible to retrieve functional data for the analyses, i.e. to speculate on the abilities

of the microbiome to deal with environmental stress factors. Other techniques, such as

shotgun metagenomics, that allow for taxonomic and functional analyses could be employed

in future, PMA-based BE surveys to obtain such functional data. In addition, environmental,

physical parameters such as humidity, temperature, or the frequency of ventilation in the BE

should be tracked, as these could help explain microbial survival (Adams et al., 2017; Xu et

al., 2020).

Our study findings support the hypothesis that the indoor surfaces of normal houses harbour

unique microbial communities (Lax et al., 2014). The single household profiles are largely

dominated by human microbial signatures and environmental taxa play more minor roles (for

comparison, please see Adams et al., 2015; Chase et al., 2016; Hewitt et al., 2012; Jeon et

al., 2013; Rintala et al., 2008), (Fig. 3). This finding is in agreement with the concept of a

personalised microbial cloud that mediates the transfer to surfaces (Meadow et al., 2015). In

particular, we identified the skin microbiome as a major contributor to this cloud (~40%), even

in areas close to the toilet, as has been shown in previous studies (Adams et al., 2015; Flores

et al., 2011; Gibbons et al., 2015). Although we observed substantial variation among the

microbial communities in the ten households, approximately 30% of the overall microbial

community detected on bathroom floors consisted of microbiomes from faecal, vaginal and

urinary origins. The presence of those microbial taxa can be explained by the production of

aerosols by toilet flushing (Prussin & Marr, 2015). However, closing the toilet lid prior to

flushing, as is practised in households H4 and H10, did not substantially alter the proportions

of contributing body sites. This result raises the question of whether other possible sources of

microbes exist, such as the shower (Feazel et al., 2009) or the sink (Dunn et al., 2013).

Molecular surveys of the BE microbiome usually do not discriminate between viable and dead

microbes. As DNA is highly stable in the environment, and microbial signatures remain

detectable for a long time, the presence of DNA signatures alone does not serve as a good

indicator for viability and metabolic activity per se. Once dispersed, DNA traces of human gut-

and skin-associated taxa can still be tracked on public restroom floors and walls after a period

of several weeks (Gibbons et al., 2015). Thus, the proportion of relevant microorganisms in

the BE might be overestimated when conventional microbiome approaches are taken.

Cultivation-based assays, however, are hindered by their technological limitations and might

only reflect a non-representative proportion of living indoor microbiota (Gilbert & Stephens,

2018).

In this study, we analysed specifically the intact floor microbial community and determined the

quality and quantity of the microbial ex-host survivors by using PMA treatment, which has

rarely been applied to BE environment samples so far (Emerson et al., 2017; Mahnert et al.,

2015; Moissl-Eichinger et al., 2015). One past specific application was the assessment of living microorganisms in the vicinity of spacecraft, a study that was carried out to understand the risk of contamination for extra-terrestrial research targets (planetary protection; Mahnert et al., 2015; Moissl-Eichinger et al., 2015). Notably, a proportion of 1–45% of the detected microbial community was deemed to be alive under (stringent) cleanroom conditions (Moissl-Eichinger et al., 2015; Vaishampayan et al., 2012).

Our experiment revealed that the majority of detected microbial signatures originated from disintegrated cells or free DNA. More specifically, we observed a more than 3- and 4-fold decrease in the absolute number of 16S rRNA genes analysed via qPCR for archaea and bacteria, respectively, upon the PMA-treatment of bathroom floor samples (Fig. 3). However, these results also imply that less than 25% of the microbial signatures originate from intact, and likely living, cells that are potentially able to colonise the human body after seven days of regular contact to a human host. The potentially living cells detected in PMA samples could be associated with a predominantly aerobic lifestyle and increased stress tolerance, and these cells could include aerobic, human- and environment-associated, but also spore-forming taxa. The predominance of aerotolerant and spore-forming taxa in PMA samples supports the hypothesis that microbes do not proliferate, but instead persist in the BE as a result of accumulation and dormancy (Gibbons, 2016).

Whether obligately anaerobic human commensals can be transmitted between individuals via the BE is still an open question. Notably, microbial taxa such as clostridia might not be directly transmitted from mother to child, even though strains of clostridia are the most abundant bacterial group in the maternal gut (Korpela et al., 2018). Korpela and de Vos (2018) argued that clostridia may rely on transmission via relatives or non-direct transmission between hosts (i.e. transmission via the environment). This argument is supported by our observation that spore formers, including clostridia, have an increased survival rate, and seem to be able to survive oxygen exposure and potentially the acidic environment of the stomach.

This kind of horizontal, indirect transfer may be a very rare event in healthy adults, as the

transmission success depends on several factors, such as transmission routes, dispersal

efficiency, the ability to survive "ex-host", human colonisation resistance, and the ability of gut

commensals to survive gastric and bile acids (Browne et al., 2017). These factors are largely

unexplored for commensals, but it can be assumed that pathogens and commensals use

similar mechanisms. For example, the transmission routes of many pathogens are well-known

and primarily include the direct inhalation of aerosols and dust (e.g. *Bacillus anthracis*,

Mycobacterium tuberculosis) and surface contact (e.g. *Clostridium difficile**, *Staphylococcus*

aureus, *Enterococcus faecalis*) (Gilbert & Stephens, 2018). It is likely that human commensals

also use such mechanisms for transmission. However, this has to be evaluated in further

studies and could be validated by performing a microbial genomic sequence variation analysis

to gather evidence for the transfer of commensals between individuals and the BE.

One intriguing results of this study was that we also detected archaea as integral components

of the BE microbiome. Even though archaea were outnumbered by bacteria nearly 100-fold,

they were found in every PMA-treated household sample. Especially taxa of the

Nitrososphaeraceae were highly predominant in PMA and non-PMA samples and seem to be

relatively stable components of the BE. In contrast, *Euryarchaeota* appeared to be only

transient occupants of the indoor microbiome, as many genera were limited to a few

households, and their signals frequently disappeared in PMA samples.

An exception to this pattern was observed for the taxon *Methanobrevibacter**, which is the

most abundant human archaeal genus in the GIT (Borrel et al., 2020). *Methanobrevibacter**

signatures were found on all analysed human body sites and in seven out of ten household

samples in both PMA and non-PMA samples. These results appear to contradict the general

assumption that methanogens are highly oxygen-sensitive (Fetzer & Conrad, 1993; Jarrell,

1985; Thauer et al., 2010; Yuan et al., 2011). However, although *Methanobacteriales** seem

to be susceptible to ROS (Lyu & Lu, 2018), analyses of experimental data reveal that

methanogens can survive oxygen exposure for periods of several hours to days (Ma & Lu,

2011; Ueki et al., 1997; Yuan et al., 2009); they also have even been found in typical

oxygenated environments (Angel et al., 2011, 2012).

By performing experiments with representative human GIT methanogens, we could confirm

that *M. smithii** *species* and *M. stadtmanae** survived aeration conditions for more than 6

hours, while *M. luminyensis** even endured oxygen exposure for more than 48 hours under

our experimental conditions (see Fig. 6). The ability of methanogens to endure aerobic

conditions was supported by a genomic analysis of common pathways to detoxify ROS. All

analysed human-associated methanogens, except *Methanosphaera stadtmanae* strain

DEW79, possessed key genes for enzymes such as rubrerythrin, ferritin, thioredoxin and

rubredoxin, all of which enable organisms to survive in aerobic environments (see Tab. 1).

Methanomassiliicoccus sp. further exhibits genes that encode for putative peroxidases and

catalases, indicating that this taxon has the potential to deal with ROS, as indicated by our

experimental approach.

By taking both genomic and experimental approaches, we gathered evidence that supports

the hypothesis that methanogens can endure aerobic conditions for a limited amounts of time.

This introduces a new potential transmission pathway, which is especially interesting with

respect to infants, as it presents another way that they can acquire microbes aside from direct

seeding during vaginal birth. It is still not know whether methanogens such as

Methanobrevibacter* are acquired perinatal and escape detection early in life or colonise the

GIT via another source, such as the BE or dietary products like yoghurt, organic milk and

vegetables (Brusa et al., 1998; Taffner, Bergna, et al., 2020; Taffner, Laggner, et al., 2020;

van de Pol et al., 2017). The latter scenario is plausible, as methanogens co-evolved with

animals (Borrel et al., 2017; Youngblut et al., 2020) and, thus, have adapted to the human

GIT.

In conclusion, this study enabled us to use PMA treatment together with molecular quantitative

and qualitative methods to successfully assess the survival of Bacteria and Archaea on indoor

surfaces. The results clearly indicate that one-fifth of the bathroom surface microbiome is

intact or even alive. As even strict anaerobes, such as oxygen-sensitive methanogenic

archaea, were found to be intact and alive despite their exposure to the fully aerated

environment, we conclude that these microorganisms potentially represent a valid source for

human microbiome constituents.

Availability of data and materials

713 The datasets supporting the conclusions of this article are available in the European

714 Nucleotide Archive (ENA) repository, Primary Accession: PRJEB41618 in

715 https://www.ebi.ac.uk/. Further details can be found in Supplementary Table 4. The R script

716 for the generation of bubble plots was used and adapted from https://github.com/alex-

717 bagnoud/OTU-table-to-bubble-plot.

Abbreviations

ASV: amplicon sequence variant

BE: built environment

Cp: crossing point

GIT: gastrointestinal tract

LEfSe: Linear discriminant analysis Effect Size

MaGe: Microbial Genome Annotation and Analysis Platform

OD: optical density

PCoA: principal coordinates analysis

PMA: propidium monoazide

qPCR: quantitative polymerase chain reaction

ROS: reactive oxygen species

TSS: total sum normalisation

Acknowledgements

We thank the participants of our study by providing samples from their bathroom floors. We

thank Charlotte Neumann and Christina Kumpitsch for critical proof-reading. Manuela Pausan

and Marcus Blohs are students of the local PhD doctoral program "MolMed". The authors

acknowledge the support of the ZMF Galaxy Team: Core Facility Computational Bioanalytics,

Medical University of Graz, funded by the Austrian Federal Ministry of Education, Science and

Research, Hochschulraum-Strukturmittel 2016 grant as part of BioTechMed Graz.

Funding

741   The project was supported financially by the local PhD doctoral program "MolMed".

Author information

Affiliations

Diagnostic and Research Institute of Hygiene, Microbiology and Environmental

Medicine, Medical University of Graz, Graz 8010, Austria

M. Blohs, A. Mahnert, C. Moissl-Eichinger

Steigerwald Arzneimittelwerk GmbH, Bayer Consumer Health, Havelstrasse 5, D-64295

Darmstadt, Germany

M.-R. Pausan

Authors´ contributions

MRP and MB contributed equally to this work. MRP, CME: study design and phylogenetic

analysis. MRP: sampling, biological samples processing, collection of metadata, microbiome

data analysis, contributed to manuscript writing. MB: analysis of microbiome data and

statistical analysis, wrote manuscript. AM: support of microbiome data and statistical analysis.

CME: genome mining, wrote manuscript.

Corresponding author

Correspondence to C. Moissl-Eichinger.

760 Ethics declarations

Ethics approval and consent to participate

Ethical approval for the study was obtained from the ethics committee of the Medical University

Graz. Written informed consent was obtained from all participants. Ethic approvals for the

respective body sites are listed in the Supplementary Table 4.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

References

Adams, R. I., Bateman, A. C., Bik, H. M., & Meadow, J. F. (2015). Microbiota of the indoor

environment: a meta-analysis. *Microbiome*, *3*, 49. https://doi.org/10.1186/s40168-015-

0108-3

Adams, R. I., Lymperopoulou, D. S., Misztal, P. K., De Cassia Pessotti, R., Behie, S. W., Tian,

Y., Goldstein, A. H., Lindow, S. E., Nazaroff, W. W., Taylor, J. W., Traxler, M. F., & Bruns,

T. D. (2017). Microbes and associated soluble and volatile chemicals on periodically wet

household surfaces. *Microbiome*, *5*(1), 128. https://doi.org/10.1186/s40168-017-0347-6

Amann, R. I., Ludwig, W., & Schleifer, K. H. (1995). Phylogenetic identification and in situ

detection of individual microbial cells without cultivation. In *Microbiological Reviews* (Vol.

59, Issue 1, pp. 143–169). American Society for Microbiology.

https://doi.org/10.1128/mmbr.59.1.143-169.1995

Angel, R., Claus, P., & Conrad, R. (2012). Methanogenic archaea are globally ubiquitous in

aerated soils and become active under wet anoxic conditions. *ISME Journal*, *6*(4), 847–

862. https://doi.org/10.1038/ismej.2011.141

Angel, R., Matthies, D., & Conrad, R. (2011). Activation of Methanogenesis in Arid Biological

Soil Crusts Despite the Presence of Oxygen. *PLoS ONE*, *6*(5), e20453.

https://doi.org/10.1371/journal.pone.0020453

Bateman, A. (2019). UniProt: A worldwide hub of protein knowledge. *Nucleic Acids Research*,

47(D1), D506–D515. https://doi.org/10.1093/nar/gky1049

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. In *Source: Journal of the Royal Statistical Society.*

Series B (Methodological) (Vol. 57, Issue 1).

Bokulich, N. A., Kaehler, B. D., Rideout, J. R., Dillon, M., Bolyen, E., Knight, R., Huttley, G. A.,

& Gregory Caporaso, J. (2018). Optimizing taxonomic classification of marker-gene

amplicon sequences with QIIME 2's q2-feature-classifier plugin. *Microbiome*, *6*(1), 90.

https://doi.org/10.1186/s40168-018-0470-z

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A.,

Alexander, H., Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger, K.,

Brejnrod, A., Brislawn, C. J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A. M.,

Chase, J., … Caporaso, J. G. (2019). Reproducible, interactive, scalable and extensible

microbiome data science using QIIME 2. In *Nature Biotechnology* (Vol. 37, Issue 8, pp.

852–857). Nature Publishing Group. https://doi.org/10.1038/s41587-019-0209-9

Borrel, G., Brugère, J. F., Gribaldo, S., Schmitz, R. A., & Moissl-Eichinger, C. (2020). The

host-associated archaeome. In *Nature Reviews Microbiology* (Vol. 18, Issue 11, pp. 622–

636). Nature Research. https://doi.org/10.1038/s41579-020-0407-y

Borrel, G., McCann, A., Deane, J., Neto, M. C., Lynch, D. B., Brugère, J. F., & O'Toole, P. W.

(2017). Genomics and metagenomics of trimethylamine-utilizing Archaea in the human

gut microbiome. *ISME Journal*, *11*(9), 2059–2074. https://doi.org/10.1038/ismej.2017.72

Brooks, A. W., Priya, S., Blekhman, R., & Bordenstein, S. R. (2018). Gut microbiota diversity

across ethnicities in the United States. *PLOS Biology*, *16*(12), e2006842.

https://doi.org/10.1371/journal.pbio.2006842

Brooks, B., Firek, B. A., Miller, C. S., Sharon, I., Thomas, B. C., Baker, R., Morowitz, M. J., &

Banfield, J. F. (2014). Microbes in the neonatal intensive care unit resemble those found

in the gut of premature infants. *Microbiome*, *2*(1), 1. https://doi.org/10.1186/2049-2618-

2-1

Browne, H. P., Neville, B. A., Forster, S. C., & Lawley, T. D. (2017). Transmission of the gut

microbiota: Spreading of health. *Nature Reviews Microbiology*, *15*(9), 531–543.

https://doi.org/10.1038/nrmicro.2017.50

Brusa, T., Ferrari, F., & Canzi, E. (1998). Methanogenic bacteria: Presence in foodstuffs.

Journal of Basic Microbiology, *38*(2), 79–84. https://doi.org/10.1002/(SICI)1521-

4028(199805)38:2<79::AID-JOBM79>3.0.CO;2-J

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P.

(2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature*

Methods, *13*(7), 581–583. https://doi.org/10.1038/nmeth.3869

Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens,

S. M., Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J. A., Smith, G., & Knight, R.

(2012). Ultra-high-throughput microbial community analysis on the Illumina HiSeq and

MiSeq platforms. *ISME Journal*, *6*(8), 1621–1624. https://doi.org/10.1038/ismej.2012.8

Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C. A., Turnbaugh,

P. J., Fierer, N., & Knight, R. (2011). Global patterns of 16S rRNA diversity at a depth of

millions of sequences per sample. *Proceedings of the National Academy of Sciences of*

the United States of America, *108*(SUPPL. 1), 4516–4522.

https://doi.org/10.1073/pnas.1000080107

Chase, J., Fouquier, J., Zare, M., Sonderegger, D. L., Knight, R., Kelley, S. T., Siegel, J., &

Caporaso, J. G. (2016). Geography and Location Are the Primary Drivers of Office

Microbiome Composition. *MSystems*, *1*(2), 1–18.

https://doi.org/10.1128/msystems.00022-16

Costello, E. K., Stagaman, K., Dethlefsen, L., Bohannan, B. J. M., & Relman, D. A. (2012).

The application of ecological theory toward an understanding of the human microbiome.

Science, *336*(6086), 1255–1262. https://doi.org/10.1126/science.1224203

Dannemiller, K. C., Mendell, M. J., Macher, J. M., Kumagai, K., Bradman, A., Holland, N.,

Harley, K., Eskenazi, B., & Peccia, J. (2014). Next-generation DNA sequencing reveals

that low fungal diversity in house dust is associated with childhood asthma development.

Indoor Air, *24*(3), 236–247. https://doi.org/10.1111/ina.12072

Dannemiller, Karen C. (2019). Moving towards a Robust Definition for a "Healthy" Indoor

Microbiome. *MSystems*, *4*(3), 1–5. https://doi.org/10.1128/msystems.00074-19

DeLong, E. F. (1992). Archaea in coastal marine environments. *Proceedings of the National*

Academy of Sciences of the United States of America, *89*(12), 5685–5689.

https://doi.org/10.1073/pnas.89.12.5685

Dridi, B., Henry, M., El Khéchine, A., Raoult, D., & Drancourt, M. (2009). High prevalence of

Methanobrevibacter smithii and Methanosphaera stadtmanae detected in the human gut

using an improved DNA detection protocol. *PLoS ONE*, *4*(9).

https://doi.org/10.1371/journal.pone.0007063

Dunn, R. R., Fierer, N., Henley, J. B., Leff, J. W., & Menninger, H. L. (2013). Home Life:

Factors Structuring the Bacterial Diversity Found within and between Homes. *PLoS ONE*,

8(5). https://doi.org/10.1371/journal.pone.0064133

Eckburg, P. B., Bik, E. M., Bernstein, C. N., Purdom, E., Dethlefsen, L., Sargent, M., Gill, S.

R., Nelson, K. E., & Relman, D. A. (2005). Microbiology: Diversity of the human intestinal

microbial flora. *Science*, *308*(5728), 1635–1638.

https://doi.org/10.1126/science.1110591

Ege, M. J., Mayer, M., Schwaiger, K., Mattes, J., Pershagen, G., Van Hage, M., Scheynius,

A., Bauer, J., & Von Mutius, E. (2012). Environmental bacteria and childhood asthma.

Allergy: European Journal of Allergy and Clinical Immunology, *67*(12), 1565–1571.

https://doi.org/10.1111/all.12028

El Fantroussi, S., Verschuere, L., Verstraete, W., & Top, E. M. (1999). Effect of phenylurea herbicides on soil microbial communities estimated by analysis of 16S rRNA gene fingerprints and community-level physiological profiles. *Applied and Environmental Microbiology*, *65*(3), 982–988. https://doi.org/10.1128/aem.65.3.982-988.1999

Emerson, J. B., Adams, R. I., Román, C. M. B., Brooks, B., Coil, D. A., Dahlhausen, K., Ganz, H. H., Hartmann, E. M., Hsu, T., Justice, N. B., Paulino-Lima, I. G., Luongo, J. C., Lymperopoulou, D. S., Gomez-Silvan, C., Rothschild-Mancinelli, B., Balk, M., Huttenhower, C., Nocker, A., Vaishampayan, P., & Rothschild, L. J. (2017). Schrödinger's microbes: Tools for distinguishing the living from the dead in microbial ecosystems. In *Microbiome* (Vol. 5, Issue 1, p. 86). BioMed Central. https://doi.org/10.1186/s40168-017-0285-3

Feazel, L. M., Baumgartner, L. K., Peterson, K. L., Frank, D. N., Harris, J. K., & Pace, N. R. (2009). Opportunistic pathogens enriched in showerhead biofilms. *Proceedings of the National Academy of Sciences of the United States of America*, *106*(38), 16393–16398. https://doi.org/10.1073/pnas.0908446106

Fetzer, S., & Conrad, R. (1993). Effect of redox potential on methanogenesis by Methanosarcina barkeri. *Archives of Microbiology*, *160*(2), 108–113. https://doi.org/10.1007/BF00288711

Flores, G. E., Bates, S. T., Knights, D., Lauber, C. L., Stombaugh, J., Knight, R., & Fierer, N. (2011). Microbial biogeography of public restroom surfaces. *PLoS ONE*, *6*(11). https://doi.org/10.1371/journal.pone.0028132

Gibbons, S. M. (2016). The Built Environment Is a Microbial Wasteland. *MSystems*, *1*(2), 1–4. https://doi.org/10.1128/msystems.00033-16

Gibbons, S. M., Schwartz, T., Fouquier, J., Mitchell, M., Sangwan, N., Gilbert, J. A., & Kelley, S. T. (2015). Ecological succession and viability of human-associated microbiota on

restroom surfaces. *Applied and Environmental Microbiology*, *81*(2), 765–773.

https://doi.org/10.1128/AEM.03117-14

Gilbert, J. A., & Stephens, B. (2018). Microbiology of the built environment. *Nature Reviews*

Microbiology, *16*(11), 661–670. https://doi.org/10.1038/s41579-018-0065-5

Guaraldi, F., & Salvatori, G. (2012). Effect of Breast and Formula Feeding on Gut Microbiota

Shaping in Newborns. *Frontiers in Cellular and Infection Microbiology*, *2*, 94.

https://doi.org/10.3389/fcimb.2012.00094

Hewitt, K. M., Gerba, C. P., Maxwell, S. L., & Kelley, S. T. (2012). Office space bacterial

abundance and diversity in three metropolitan areas. *PLoS ONE*, *7*(5).

https://doi.org/10.1371/journal.pone.0037849

Horve, P. F., Lloyd, S., Mhuireach, G. A., Dietz, L., Fretz, M., MacCrone, G., Van Den

Wymelenberg, K., & Ishaq, S. L. (2020). Building upon current knowledge and techniques

of indoor microbiology to construct the next era of theory into microorganisms, health,

and the built environment. *Journal of Exposure Science and Environmental*

Epidemiology, *30*(2), 219–235. https://doi.org/10.1038/s41370-019-0157-y

Jarrell, K. F. (1985). Extreme Oxygen Sensitivity in Methanogenic Archaebacteria.

BioScience, *35*(5), 298–302. https://doi.org/10.2307/1309929

Jeon, Y. S., Chun, J., & Kim, B. S. (2013). Identification of household bacterial community and

analysis of species shared with human microbiome. *Current Microbiology*, *67*(5), 557–

563. https://doi.org/10.1007/s00284-013-0401-y

Kelley, S. T., & Gilbert, J. A. (2013). Studying the microbiology of the indoor environment. In

Genome Biology (Vol. 14, Issue 2). BioMed Central Ltd. https://doi.org/10.1186/gb-2013-

14-2-202

Kirjavainen, P. V., Karvonen, A. M., Adams, R. I., Täubel, M., Roponen, M., Tuoresmäki, P.,

Loss, G., Jayaprakash, B., Depner, M., Ege, M. J., Renz, H., Pfefferle, P. I., Schaub, B.,

Lauener, R., Hyvärinen, A., Knight, R., Heederik, D. J. J., von Mutius, E., & Pekkanen, J.

(2019). Farm-like indoor microbiota in non-farm homes protects children from asthma

development. *Nature Medicine*, *25*(7), 1089–1095. https://doi.org/10.1038/s41591-019-

0469-4

Klepeis, N. E., Nelson, W. C., Ott, W. R., Robinson, J. P., Tsang, A. M., Switzer, P., Behar, J.

V., Hern, S. C., & Engelmann, W. H. (2001). The National Human Activity Pattern Survey

(NHAPS): A resource for assessing exposure to environmental pollutants. *Journal of*

Exposure Analysis and Environmental Epidemiology, *11*(3), 231–252.

https://doi.org/10.1038/sj.jea.7500165

Klindworth, A., Pruesse, E., Schweer, T., Peplies, J., Quast, C., Horn, M., & Glöckner, F. O.

(2013). Evaluation of general 16S ribosomal RNA gene PCR primers for classical and

next-generation sequencing-based diversity studies. *Nucleic Acids Research*, *41*(1), e1–

e1. https://doi.org/10.1093/nar/gks808

Knights, D., Kuczynski, J., Charlson, E. S., Zaneveld, J., Mozer, M. C., Collman, R. G.,

Bushman, F. D., Knight, R., & Kelley, S. T. (2011). Bayesian community-wide culture-

independent microbial source tracking. *Nature Methods*, *8*(9), 761–765.

https://doi.org/10.1038/nmeth.1650

Korpela, K., Costea, P., Coelho, L. P., Kandels-Lewis, S., Willemsen, G., Boomsma, D. I.,

Segata, N., & Bork, P. (2018). Selective maternal seeding and environment shape the

human gut microbiome. *Genome Research*, *28*(4), 561–568.

https://doi.org/10.1101/gr.233940.117

Koskinen, K., Reichert, J. L., Hoier, S., Schachenreiter, J., Duller, S., Moissl-Eichinger, C., &

Schöpf, V. (2018). The nasal microbiome mirrors and potentially shapes olfactory

function. *Scientific Reports*, *8*(1), 1–11. https://doi.org/10.1038/s41598-018-19438-3

Kumar, S., Stecher, G., & Tamura, K. (2016). MEGA7: Molecular Evolutionary Genetics

Analysis Version 7.0 for Bigger Datasets. *Molecular Biology and Evolution*, *33*(7), 1870–

1874. https://doi.org/10.1093/molbev/msw054

Lax, S., Sangwan, N., Smith, D., Larsen, P., Handley, K. M., Richardson, M., Guyton, K.,

Krezalek, M., Shogan, B. D., Defazio, J., Flemming, I., Shakhsheer, B., Weber, S.,

Landon, E., Garcia-Houchins, S., Siegel, J., Alverdy, J., Knight, R., Stephens, B., &

Gilbert, J. A. (2017). Bacterial colonization and succession in a newly opened hospital.

Science Translational Medicine, *9*(391), 1–12.

https://doi.org/10.1126/scitranslmed.aah6500

Lax, S., Smith, D. P., Hampton-Marcell, J., Owens, S. M., Handley, K. M., Scott, N. M.,

Gibbons, S. M., Larsen, P., Shogan, B. D., Weiss, S., Metcalf, J. L., Ursell, L. K.,

Vázquez-Baeza, Y., Van Treuren, W., Hasan, N. A., Gibson, M. K., Colwell, R., Dantas,

G., Knight, R., & Gilbert, J. A. (2014). Longitudinal analysis of microbial interaction

between humans and the indoor environment. *Science*, *345*(6200), 1048–1052.

https://doi.org/10.1126/science.1254529

Lax, S., Smith, D., Sangwan, N., Handley, K., Larsen, P., Richardson, M., Taylor, S., Landon,

E., Alverdy, J., Siegel, J., Stephens, B., Knight, R., & Gilbert, J. A. (2017). Colonization

and succession of hospital-associated microbiota. *Science Translational Medicine*,

9(391), eaah6500. https://doi.org/10.1126/scitranslmed.aah6500.Colonization

Letunic, I., & Bork, P. (2007). Interactive Tree Of Life (iTOL): An online tool for phylogenetic

tree display and annotation. *Bioinformatics*, *23*(1), 127–128.

https://doi.org/10.1093/bioinformatics/btl529

Ley, R. E., Lozupone, C. A., Hamady, M., Knight, R., & Gordon, J. I. (2008). Worlds within

worlds: Evolution of the vertebrate gut microbiota. *Nature Reviews Microbiology*, *6*(10),

776–788. https://doi.org/10.1038/nrmicro1978

Lyu, Z., & Lu, Y. (2018). Metabolic shift at the class level sheds light on adaptation of methanogens to oxidative environments. *ISME Journal*, *12*(2), 411–423. https://doi.org/10.1038/ismej.2017.173

Ma, K., & Lu, Y. (2011). Regulation of microbial methane production and oxidation by intermittent drainage in rice field soil. *FEMS Microbiology Ecology*, *75*(3), 446–456. https://doi.org/10.1111/j.1574-6941.2010.01018.x

Mahnert, A., Moissl-Eichinger, C., Zojer, M., Bogumil, D., Mizrahi, I., Rattei, T., Martinez, J. L., & Berg, G. (2019). Man-made microbial resistances in built environments. *Nature Communications*, *10*(1), 1–12. https://doi.org/10.1038/s41467-019-08864-0

Mahnert, A., Vaishampayan, P., Probst, A. J., Auerbach, A., Moissl-Eichinger, C., Venkateswaran, K., & Berg, G. (2015). Cleanroom Maintenance Significantly Reduces Abundance but Not Diversity of Indoor Microbiomes. *PLOS ONE*, *10*(8), e0134848. https://doi.org/10.1371/journal.pone.0134848

Mahnert, A., Verseux, C., Schwendner, P., Koskinen, K., Kumpitsch, C., Blohs, M., Wink, L., Brunner, D., Goessler, T., Billi, D., & Moissl-Eichinger, C. (2020). *Microbiome dynamics during the HI-SEAS IV mission, and implications for future crewed missions beyond Earth*. https://doi.org/10.21203/rs.3.rs-44929/v1

Maria Chibani, C., Mahnert, A., Borrel, G., Almeida, A., Werner, A., Brugère, J.-F., Gribaldo, S., Finn, R. D., Schmitz, R. A., & Moissl-Eichinger, C. (2020). A comprehensive analysis of the global human gut archaeome from a thousand genome catalogue. *BioRxiv*, 2020.11.21.392621. https://doi.org/10.1101/2020.11.21.392621

Mauerhofer, L.-M., Zwirtmayr, S., Pappenreiter, P., Bernacchi, S., Seifert, A. H., Reischl, B., Schmider, T., Taubner, R.-S., Paulik, C., & Rittmann, S. .-M. R. (n.d.). Hyperthermophilic methanogenic archaea are high-pressure cell factories for H2/CO2 conversion. *Communications Biology*.

McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. *PLoS ONE*, *8*(4), e61217.
https://doi.org/10.1371/journal.pone.0061217

Meadow, J. F., Altrichter, A. E., Bateman, A. C., Stenson, J., Brown, G. Z., Green, J. L., &
Bohannan, B. J. M. (2015). Humans differ in their personal microbial cloud. *PeerJ*,
2015(9). https://doi.org/10.7717/peerj.1258

Meehan, C. J., & Beiko, R. G. (2014). A phylogenomic view of ecological specialization in the
lachnospiraceae, a family of digestive tract-associated bacteria. *Genome Biology and*
Evolution, *6*(3), 703–713. https://doi.org/10.1093/gbe/evu050

Miller, R. A., & Britigan, B. E. (1997). Role of oxidants in microbial pathophysiology. In *Clinical*
Microbiology Reviews (Vol. 10, Issue 1, pp. 1–18). American Society for Microbiology.
https://doi.org/10.1128/cmr.10.1.1

Million, M., & Raoult, D. (2018). Linking gut redox to human microbiome. *Human Microbiome*
Journal, *10*(July), 27–32. https://doi.org/10.1016/j.humic.2018.07.002

Moissl-Eichinger, C., Auerbach, A. K., Probst, A. J., Mahnert, A., Tom, L., Piceno, Y.,
Andersen, G. L., Venkateswaran, K., Rettberg, P., Barczyk, S., Pukall, R., & Berg, G.
(2015). Quo vadis? Microbial profiling revealed strong effects of cleanroom maintenance
and routes of contamination in indoor environments. *Scientific Reports*, *5*(1), 1–13.
https://doi.org/10.1038/srep09156

Moossavi, S., & Azad, M. B. (2020). Origins of human milk microbiota: new evidence and
arising   questions.   *Gut   Microbes*,   *12*(1),   1667722.
https://doi.org/10.1080/19490976.2019.1667722

Mora, M., Mahnert, A., Koskinen, K., Pausan, M. R., Oberauner-Wappis, L., Krause, R.,
Perras, A. K., Gorkiewicz, G., Berg, G., & Moissl-Eichinger, C. (2016). Microorganisms

in confined habitats: Microbial monitoring and control of intensive care units, operating

rooms, cleanrooms and the international space station. *Frontiers in Microbiology*,

7(October), 1–20. https://doi.org/10.3389/fmicb.2016.01573

Ng, K. M., Aranda-Díaz, A., Tropini, C., Frankel, M. R., Van Treuren, W., O'Laughlin, C. T.,

Merrill, B. D., Yu, F. B., Pruss, K. M., Oliveira, R. A., Higginbottom, S. K., Neff, N. F.,

Fischbach, M. A., Xavier, K. B., Sonnenburg, J. L., & Huang, K. C. (2019). Recovery of

the Gut Microbiota after Antibiotics Depends on Host Diet, Community Context, and

Environmental Reservoirs. *Cell Host and Microbe*, *26*(5), 650-665.e4.

https://doi.org/10.1016/j.chom.2019.10.011

Nocker, A., Sossa-Fernandez, P., Burr, M. D., & Camper, A. K. (2007). Use of propidium

monoazide for live/dead distinction in microbial ecology. *Applied and Environmental*

Microbiology, *73*(16), 5111–5117. https://doi.org/10.1128/AEM.02987-06

Palmer, C., Bik, E. M., DiGiulio, D. B., Relman, D. A., & Brown, P. O. (2007). Development of

the human infant intestinal microbiota. *PLoS Biology*, *5*(7), 1556–1573.

https://doi.org/10.1371/journal.pbio.0050177

Parajuli, A., Hui, N., Puhakka, R., Oikarinen, S., Grönroos, M., Selonen, V. A. O., Siter, N.,

Kramna, L., Roslund, M. I., Vari, H. K., Nurminen, N., Honkanen, H., Hintikka, J.,

Sarkkinen, H., Romantschuk, M., Kauppi, M., Valve, R., Cinek, O., Laitinen, O. H., …

Sinkkonen, A. (2020). Yard vegetation is associated with gut microbiota composition.

Science of the Total Environment, *713*, 136707.

https://doi.org/10.1016/j.scitotenv.2020.136707

Pasolli, E., Asnicar, F., Manara, S., Zolfo, M., Karcher, N., Armanini, F., Beghini, F., Manghi,

P., Tett, A., Ghensi, P., Collado, M. C., Rice, B. L., DuLong, C., Morgan, X. C., Golden,

C. D., Quince, C., Huttenhower, C., & Segata, N. (2019). Extensive Unexplored Human

Microbiome Diversity Revealed by Over 150,000 Genomes from Metagenomes Spanning

Age, Geography, and Lifestyle. *Cell*, *176*(3), 649-662.e20.

https://doi.org/10.1016/j.cell.2019.01.001

Pausan, M. R., Csorba, C., Singer, G., Till, H., Schöpf, V., Santigli, E., Klug, B., Högenauer,

C., Blohs, M., & Moissl-Eichinger, C. (2019). Exploring the Archaeome: Detection of

Archaeal Signatures in the Human Body. *Frontiers in Microbiology*, *10*, 2796.

https://doi.org/10.3389/fmicb.2019.02796

Probst, A. J., Auerbach, A. K., & Moissl-Eichinger, C. (2013). Archaea on Human Skin. *PLoS*

ONE, *8*(6). https://doi.org/10.1371/journal.pone.0065388

Pruesse, E., Peplies, J., & Glöckner, F. O. (2012). SINA: Accurate high-throughput multiple

sequence alignment of ribosomal RNA genes. *Bioinformatics*, *28*(14), 1823–1829.

https://doi.org/10.1093/bioinformatics/bts252

Prussin, A. J., & Marr, L. C. (2015). Sources of airborne microorganisms in the built

environment. *Microbiome*, *3*, 78. https://doi.org/10.1186/s40168-015-0144-z

Qian, J., Hospodsky, D., Yamamoto, N., Nazaroff, W. W., & Peccia, J. (2012). Size-resolved

emission rates of airborne bacteria and fungi in an occupied classroom. *Indoor Air*, *22*(4),

339–351. https://doi.org/10.1111/j.1600-0668.2012.00769.x

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., & Glöckner,

F. O. (2013). The SILVA ribosomal RNA gene database project: Improved data

processing and web-based tools. *Nucleic Acids Research*, *41*(D1), D590.

https://doi.org/10.1093/nar/gks1219

Raskin, L., Stromley, J. M., Rittmann, B. E., & Stahl, D. A. (1994). Group-specific 16S rRNA

hybridization probes to describe natural communities of methanogens. *Applied and*

Environmental Microbiology, *60*(4), 1232–1240. https://doi.org/10.1128/aem.60.4.1232-

1240.1994

Rintala, H., Pitkäranta, M., Toivola, M., Paulin, L., & Nevalainen, A. (2008). Diversity and
seasonal dynamics of bacterial community in indoor environment. *BMC Microbiology*, *8*.
https://doi.org/10.1186/1471-2180-8-56

Rocchi, S., Reboux, G., Scherer, E., Laboissière, A., Zaros, C., Rouzet, A., Valot, B., Khan,
S., Dufourg, M. N., Leynaert, B., Raherison, C., & Millon, L. (2020). Indoor microbiome:
Quantification of exposure and association with geographical location, meteorological
factors, and land use in France. *Microorganisms*, *8*(3).
https://doi.org/10.3390/microorganisms8030341

Rolfe, R. D., Hentges, D. J., Campbell, B. J., & Barrett, J. T. (1978). Factors related to the
oxygen tolerance of anaerobic bacteria. *Applied and Environmental Microbiology*, *36*(2),
306–313. https://doi.org/10.1128/aem.36.2.306-313.1978

Shan, Y., Wu, W., Fan, W., Haahtela, T., & Zhang, G. (2019). House dust microbiome and
human health risks. *International Microbiology*, *22*(3), 297–304.
https://doi.org/10.1007/s10123-019-00057-5

Stewart, C. J., Ajami, N. J., O'Brien, J. L., Hutchinson, D. S., Smith, D. P., Wong, M. C., Ross,
M. C., Lloyd, R. E., Doddapaneni, H. V., Metcalf, G. A., Muzny, D., Gibbs, R. A., Vatanen,
T., Huttenhower, C., Xavier, R. J., Rewers, M., Hagopian, W., Toppari, J., Ziegler, A. G.,
… Petrosino, J. F. (2018). Temporal development of the gut microbiome in early
childhood from the TEDDY study. *Nature*, *562*(7728), 583–588.
https://doi.org/10.1038/s41586-018-0617-x

Taffner, J., Bergna, A., Cernava, T., & Berg, G. (2020). Tomato-Associated Archaea Show a
Cultivar-Specific Rhizosphere Effect but an Unspecific Transmission by Seeds.
Phytobiomes Journal, *4*(2), 133–141. https://doi.org/10.1094/PBIOMES-01-20-0017-R

Taffner, J., Laggner, O., Wolfgang, A., Coyne, D., & Berg, G. (2020). Exploring the Microbiota
of East African Indigenous Leafy Greens for Plant Growth, Health, and Resilience.

Frontiers in Microbiology, *11*. https://doi.org/10.3389/fmicb.2020.585690

Thauer, R. K., Kaster, A.-K., Goenrich, M., Schick, M., Hiromoto, T., & Shima, S. (2010).

Hydrogenases from Methanogenic Archaea, Nickel, a Novel Cofactor, and H 2 Storage .

Annual Review of Biochemistry, *79*(1), 507–536.

https://doi.org/10.1146/annurev.biochem.030508.152103

Ueki, A., Ono, K., Tsuchiya, A., & Ueki, K. (1997). Survival of methanogens in air-dried paddy

field soil and their heat tolerance. *Water Science and Technology*, *36*(6–7), 517–522.

https://doi.org/10.1016/S0273-1223(97)00563-5

Vaishampayan, P. A., Rabbow, E., Horneck, G., & Venkateswaran, K. J. (2012). Survival of

bacillus pumilus spores for a prolonged period of time in real space conditions.

Astrobiology, *12*(5), 487–497. https://doi.org/10.1089/ast.2011.0738

Vallenet, D., Labarre, L., Rouy, Z., Barbe, V., Bocs, S., Cruveiller, S., Lajus, A., Pascal, G.,

Scarpelli, C., & Médigue, C. (2006). MaGe: A microbial genome annotation system

supported by synteny results. *Nucleic Acids Research*, *34*(1), 53–65.

https://doi.org/10.1093/nar/gkj406

van de Pol, J. A. A., van Best, N., Mbakwa, C. A., Thijs, C., Savelkoul, P. H., Arts, I. C. W.,

Hornef, M. W., Mommers, M., & Penders, J. (2017). Gut Colonization by Methanogenic

Archaea Is Associated with Organic Dairy Consumption in Children. *Frontiers in*

Microbiology, *8*(MAR), 355. https://doi.org/10.3389/fmicb.2017.00355

Walters, W., Hyde, E. R., Berg-Lyons, D., Ackermann, G., Humphrey, G., Parada, A., Gilbert,

J. A., Jansson, J. K., Caporaso, J. G., Fuhrman, J. A., Apprill, A., & Knight, R. (2016).

Improved Bacterial 16S rRNA Gene (V4 and V4-5) and Fungal Internal Transcribed

Spacer Marker Gene Primers for Microbial Community Surveys. *MSystems*, *1*(1).

https://doi.org/10.1128/msystems.00009-15

Wampach, L., Heintz-Buschart, A., Hogan, A., Muller, E. E. L., Narayanasamy, S., Laczny, C.

C., Hugerth, L. W., Bindl, L., Bottu, J., Andersson, A. F., de Beaufort, C., & Wilmes, P.

(2017). Colonization and succession within the human gut microbiome by archaea,

bacteria, and microeukaryotes during the first year of life. *Frontiers in Microbiology*,

8(MAY), 1–21. https://doi.org/10.3389/fmicb.2017.00738

Ward, T., Larson, J., Meulemans, J., Hillmann, B., Lynch, J., Sidiropoulos, D., Spear, J. R.,

Caporaso, G., Blekhman, R., Knight, R., Fink, R., & Knights, D. (2017). BugBase predicts

organism-level microbiome phenotypes. In *bioRxiv* (p. 133462). bioRxiv.

https://doi.org/10.1101/133462

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

https://ggplot2.tidyverse.org

Wolf, A., Moissl-Eichinger, C., Perras, A., Koskinen, K., Tomazic, P. V., & Thurnher, D. (2017).

The salivary microbiome as an indicator of carcinogenesis in patients with oropharyngeal

squamous cell carcinoma: A pilot study. *Scientific Reports*, *7*(1).

https://doi.org/10.1038/s41598-017-06361-2

Xu, Y., Tandon, R., Ancheta, C., Arroyo, P., Gilbert, J. A., Stephens, B., & Kelley, S. T. (2020).

Quantitative profiling of built environment bacterial and fungal communities reveals

dynamic material dependent growth patterns and microbial interactions. *Indoor Air*.

https://doi.org/10.1111/ina.12727

Youngblut, N. D., Reischer, G. H., Dauser, S., Walzer, C., Farnleitner, A. H., & Ley, R. E.

(2020). Environmental Microbiology and Molecular Diagnostics 166/5/3, Gumpendorfer

Straße 1a. *BioRxiv*, 2020.11.10.376293. https://doi.org/10.1101/2020.11.10.376293

Yuan, Y., Conrad, R., & Lu, Y. (2009). Responses of methanogenic archaeal community to

oxygen exposure in rice field soil. *Environmental Microbiology Reports*, *1*(5), 347–354.

https://doi.org/10.1111/j.1758-2229.2009.00036.x

Yuan, Y., Conrad, R., & Lu, Y. (2011). Transcriptional response of methanogen mcrA genes

to oxygen exposure of rice field soil. *Environmental Microbiology Reports*, *3*(3), 320–328.

https://doi.org/10.1111/j.1758-2229.2010.00228.x

Zakrzewski, M., Proietti, C., Ellis, J. J., Hasan, S., Brion, M. J., Berger, B., & Krause, L. (2017).

Calypso: A user-friendly web-server for mining and visualizing microbiome-environment

interactions. *Bioinformatics*, *33*(5), 782–783.

https://doi.org/10.1093/bioinformatics/btw725

1143

1144 Supplementary files

Supplementary Figure 1. Sampling area of a representative bathroom floor.

Supplementary Figure 2. Analysis of indoor microbiome samples, displaying differences

upon PMA treatment on bacterial phyla level. Depicted are taxa that were characteristic to

nonPMA and PMA indoor samples. (A) LEfSe analysis, and (B) boxplots of significant different

taxa defined by Wilcoxon signed-rank test: *, $P < 0.05$; **, $P < 0.01$.

Supplementary Figure 3. LEfSe analysis of bacterial taxa (phylum, class, order, family,

genus-level) depicting differences of PMA and non-PMA indoor samples.

Supplementary Figure 4. PMA treated indoor samples show a significantly decreased

relative abundance for some opportunistic pathogens.

Supplementary Figure 5. Comparison of archaeal signatures found on different human body

sites (blue) and in the 10 analysed bathrooms (red) on phylum (top) level and for the 25 most

abundant genera (bottom). Only samples without PMA treatment were included. Sign.: *, $P <$

0.05; ****, $P < 0.0001$; Wilcoxon signed-rank test.

Supplementary Table 1. Primer pairs used for archaea and bacteria PCR and qPCR.

Supplementary Table 2. The PCR and qPCR conditions for the primer pairs used.

Supplementary Table 3. ASV table and data overview for indoor microbiomes.

Supplementary Table 4. Summary of additional datasets (human microbiome) used in this

study. The study including the saliva samples (bacterial dataset), as well as the study on the

nasal cavity (archaeal dataset) have already been published (Koskinen et al., 2018; Wolf et

al., 2017).

Supplementary Table 5. ASV table of all indoor and human samples, universal dataset.

Supplementary Table 6. List of shared taxa among different households.

Supplementary Table 7. List of most abundant bacterial genera, families and orders and the

respective impact of PMA treatment on their relative abundance, analysed by Wilcoxon

signed-rank test.

Supplementary Table 8. ASV table of all indoor samples, archaeal dataset.

Supplementary Table 9. ASV table of all indoor and human samples, archaeal dataset.

1172