Quantifying bacterial evolution in the wild: a birthday problem for *Campylobacter*

lineages

Jessica K. Calland[1], Ben Pascoe[1], Sion C. Bayliss[1], Evangelos Mourkas[1], Elvire Berthenet[2],

Harry A. Thorpe[1,3], Matthew D. Hitchings[4], Edward J. Feil[1], Jukka Corander[3,5,6], Martin J.

Blaser[7], Daniel Falush[8]# & Samuel K. Sheppard[1,9]#

[1]The Milner Centre for Evolution, University of Bath, Claverton down, Bath, UK; [2]French
National Reference Center for Campylobacters and Helicobacters, University of Bordeaux,
33076, Bordeaux, France; [3]Department of Biostatistics, University of Oslo, Oslo, Norway;
[4]Institute of Life Sciences, Swansea University Medical School, Swansea University,
Singleton Park, Swansea, UK; [5]Department of Mathematics and Statistics, Helsinki Institute
for Information Technology, University 18 of Helsinki, Helsinki, Finland; [6]Parasites and
Microbes, Wellcome Sanger Institute, Cambridge, UK; [7]Center for Advanced Biotechnology
and Medicine, Rutgers University, New Brunswick, New Jersey, USA; [8]Centre for Microbes,
Development and Health, Institute Pasteur Shanghai, China; [9]Department of Zoology,
University of Oxford, South Parks Road, Oxford OX1 3PS, UK.
18

19 #Authors to which correspondence should be addressed: s.k.sheppard@bath.ac.uk &

20 daniel.falush@ips.ac.cn

22 **Short title**: *Campylobacter* molecular clock

24 **Keywords:** Evolution / Pathogenesis / Molecular clock / Population structure / Population

25 genomics.

Abstract

Measuring molecular evolution in bacteria typically requires estimation of the rate at which nucleotide changes accumulate in strains sampled at different times that share a common ancestor. This approach has been useful for dating ecological and evolutionary events that coincide with the emergence of important lineages, such as outbreak strains and obligate human pathogens. However, in multi-host (niche) transmission scenarios, where the pathogen is essentially an opportunistic environmental organism, sampling is often sporadic and rarely reflects the overall population, particularly when concentrated on clinical isolates. This means that approaches that assume recent common ancestry are not applicable. Here we present a new approach to estimate the molecular clock rate in *Campylobacter* that draws on the popular probability conundrum known as the 'birthday problem'. Using large genomic datasets and comparative genomic approaches, we use isolate pairs that share recent common ancestry to estimate the rate of nucleotide change for the population. Identifying synonymous and non-synonymous nucleotide changes, both within and outside of recombined regions of the genome, we quantify clock-like diversification to estimate synonymous rates of nucleotide change for the common pathogenic bacteria *Campylobacter coli* (2.4×10^{-6} s/s/y) and *Campylobacter jejuni* (3.4×10^{-6} s/s/y). Finally, using estimated total rates of nucleotide change, we infer the number of effective lineages within the sample time-frame – analogous to a shared birthdays – and assess the rate of turnover of lineages in our sample set over short evolutionary timescales. This provides a generalizable approach to calibrating rates in populations of environmental bacteria and shows that multiple lineages are maintained, implying that large-scale clonal sweeps may take hundreds of years or more in these species.

Author Summary

Growth and reproduction in living organisms require DNA replication but this process is error prone. Along with variation introduced by horizontal gene transfer, it can lead to alterations in the nucleotide sequence. These nucleotide changes accumulate over time in successive generations at an approximately constant rate termed the molecular clock. Therefore, if this rate is known, one can estimate the date when two or more lineages diverged. In bacteria, this can be informative for understanding the time-scale of emergence and spread of pathogenic strains. Such analyses are robust when the ancestral population is known, such as for obligate pathogens that only infect humans. However, when the bacterium inhabits multiple hosts or niches it is difficult to infer direct ancestry from one strain to another, reducing the accuracy of molecular clock estimates. Here we focus on one such multi-host organism, *Campylobacter,* a leading cause of food-borne gastroenteritis. Reconstructing the population history by estimating empirical nucleotide change rates from carefully selected isolate pairs, and evaluating the maintenance of multiple lineages over time, we provide information about strain diversification. Our method is a new addition to the bacterial genomics toolkit that will help in understanding the spread of opportunistic pathogens.

Introduction

Theoretical models of a relatively constant rate of molecular change over time [1], the molecular clock, have become fundamental to explaining the evolution in bacteria [2,3]. Spurred by the increasing availability of population-scale genome datasets, it is now common for comparative genomic studies to describe not only the relatedness of isolates but also how long ago they diverged [4-8]. This can provide valuable information when combined with host, habitat or ecosystem data. For example, it is possible to investigate how events such as host transitions or global dissemination have influenced the emergence and spread of lineages that may display important phenotypes, including pathogenicity.

There are significant challenges when applying molecular clocks to date lineage diversification in natural bacterial populations. In particular, it is necessary to determine the rate at which the clock 'ticks' and the uniform accumulation of nucleotide change (NC) over time. However, this is not simply a reflection of the background point mutation rate (associated with replication error) and the generation time of the bacterium [9,10], but is also influenced by horizontal gene transfer (HGT) that can introduce several NCs in a single event [11]. Furthermore, the rate at which NCs accumulate in the population is influenced by the population size [12] and selection (positive and stabilizing) on different fitness effects [13].

While debate continues about NCs that are effectively neutral, and hence provide accurate clock estimates [14], there is clear utility for even approximations of the rate of genome change over time [15,16]. This has allowed the development of time-calibrated phylogenies explaining molecular evolution in numerous well-known pathogen species [4-7]. However, even with large genome datasets and increasingly sophisticated models [17,18], the accuracy

of molecular evolution estimates is dependent upon the data from which they are derived, and

two important considerations remain. First, the data should represent a longitudinal sample set

[15,19]. Second, the data should be representative of the population as a whole.

It is conceptually simple to understand how a long time-frame between collection of the

earliest and latest sample would increase the number of NCs recorded, and how sampling at

consistent intervals could help to determine if accumulation was linear over time.

Comparisons between modern samples and DNA from the stomach of a 5,300 year old frozen

iceman 'Otzi' have been used to investigate the emergence of modern *Helicobacter pylori*

lineages [20]. However, ancient pathogen samples are rarely available. More frequently,

molecular clock rates are estimated using collections of contemporary isolates that often share

a common ancestor older than the sample frame. Convincing estimations have been possible

for medically important bacteria, through comparison of large numbers of closely related

isolates [21-24] but for many pathogens sampling of outbreaks may not provide an adequate

representation of the bacterial population.

Most disease-causing bacteria are not obligate human pathogens. In this case, large reservoirs

of isolates from which infection can arise may be infrequently sampled, despite their potential

importance as emergent pathogenic strains. For example, *Campylobacter jejuni* and *C. coli*

are among the most common causes of bacterial gastroenteritis worldwide but exist

principally as commensal organisms in the gut of mammals and birds [25-29]. Human

infection results primarily via food contaminated with strains from wild and agricultural

animals, especially chickens [30-35]. In multi-host (niche) transmission scenarios such as this,

where the pathogen is essentially an environmental organism, sampling is often sporadic and

rarely reflects the overall population, particularly when concentrated on clinical isolates [36].

Overcoming the problem of sporadic or unrepresentative sampling for molecular clock

estimation requires that sufficient numbers of isolates are collected to ensure that there are

pairs that share a recent common ancestor (within the sampling period). However, with the

enormous effective population size of environmental bacteria populations, questions remain

about how many isolates need to be sampled to achieve this. This is analogous to the well-

known probability theory conundrum known as the birthday problem [37]. This puzzle asks

how many randomly chosen people need to be sampled so that a pair of them will share the

same birthday. To be sure, requires a sample size of 366 (the number of possible birthdays),

assuming that all birthdays are equally common, but a 99.9% probability is achieved with just

70 people and 50% with 23 people. This may seem counter intuitive but can be explained by

considering that rather than comparing the birthday of a single individual to everyone else's,

in fact comparisons are made between every pair of individuals, 23 x 22/2 = 253. The result is

greater than half the number days in the year, hence the 50% probability. Clearly, there are

challenges in relating this conceptual model to bacteria. First, it is not known how many

possible lineages (here equivalent to birthdays) there are in natural bacterial populations.

Second, how to define lineages or isolate pairs with recent common ancestry. Third, just as

with birthdays, some lineages are far more common than others. For example, of >72,000 *C.*

jejuni and *C. coli* isolates archived in the pubMLST database [38], >50% belong to just 5

clonal complexes (out of 45).

Together, factors relating to isolate sampling and genome analysis conspire such that it may

be difficult to distinguish NCs that reflect the passage of time [16,39]. Here, we take a multi-

layered approach to estimate the rate of molecular evolution of *C. jejuni* and *C. coli* using a

large genome collection (2,425 genomes) representing isolates sampled over a 46-year period.

We begin by identifying closely related isolate pairs in which the most recently sampled

isolate has accumulated NCs over time. We then quantify synonymous and non-synonymous

polymorphisms to take (some) account of selection, both within and outside of recombinant

regions of the genome, and use synonymous polymorphisms to quantify clock-like

diversification in *Campylobacter* [40,41]. Finally, using estimated rates of nucleotide change

we assess the rate of turnover of lineages in our sample sets over short evolutionary

timescales. This provides a generalizable approach to calibrating rates in populations of

environmental bacteria and clues about lineage diversification in two important enteric

pathogens.

Results

There is a weak temporal signal in *C. coli* and *C. jejuni* phylogenies

Core genome phylogenies revealed little evidence of clustering by collection date (**Fig 1**). Isolates belonging to common sequence types (STs) and clonal complexes were sampled over the 46-year period. For *C. coli* and *C. jejuni* respectively, 1, 16, 3, 211, 370 and 41, 3, 34, 469, 1277 isolates were sampled over 50, 40, 30, 20, and 10 years ago. These included poultry associated ST-353, ST-354 and ST-257 complexes, cattle associated ST-61 and ST-42 complexes, and host generalist ST-21, ST-45, ST-828 (*C. coli*) complexes [42] (**Fig 1 and S1 Table**). Linear regression of root-to-tip distances and sampling dates of *C. coli* and *C. jejuni* phylogenies **(S1 and S2 Fig)**, using TempEst software, provided very weak evidence of a temporal signal when the best-fitting root was estimated. The R^2 values were low for both *C. coli* ($R^2 = 0.176$, slope $= 3 \times 10^{-5}$) and *C. jejuni* ($R^2 = 9.5 \times 10^{-2}$, slope $= 6.4 \times 10^{-5}$) phylogenies (**S3 and S4 Tables**). Root-to-tip regression analysis was also run for *C. coli* and *C. jejuni* on three separate phylogenetic trees which were built from core gene alignments with the top 1, 5 and 10% most and least variable core genes (alleles / locus) filtered and removed. However, temporal signal remained poor for both *C. coli* (Avg. $R^2 = 2.4 \times 10^{-2}$) and *C. jejuni* (Avg. $R^2 = 1.3 \times 10^{-2}$) (**S3 Fig**). Consistent with some other studies [43], this poor branch-length to isolation date correlation suggests that estimation of the molecular clock rate from the entire dataset may be difficult. However, the accumulation of polymorphisms exhibited a positive correlation with sampling date in all datasets (**S1 and S2 Figs**) implying the maintenance of multiple STs and clonal complexes through time.

Analyses of temporal signal can be improved in bacterial genomes by the removal of recombined regions where multiple genetic variations may be introduced in a single

evolutionary event [44]. Masking recombination in this way is challenging for large genome

datasets such as those used in this study. Therefore, we conducted root-to-tip regression

analysis on sub-lineages (tree clusters) within *C. coli* and *C. jejuni* where recombination

events could be efficiently excluded (**S3 and S4 Tables**). Consistent with previous studies

[44], any lineage with an R^2 value >0.5 was described as having a strong temporal signal and

was used in subsequent Bayesian evolutionary analysis. There were large differences in the

strength of the temporal signal across all sub-lineages with the lowest R^2 value found in the

host generalist *C. jejuni* ST-45 clonal complex ($R^2 = 0.0229$, slope = 1.89 x 10^{-5}) and highest

in *C. coli* ST-1090 ($R^2 = 0.8603$, slope = 7.07 x 10^{-5}) (**S3 and S4 Tables**). However, only 3

out of 8 *C. coli* and 5 out of 18 *C. jejuni* sub-lineages exhibited strong temporal signal with R^2

>0.5.

Bayesian evolutionary analyses were performed on sub-lineages where temporal signal was

strong ($R^2 > 0.5$) using BEAST2 [45]. This excluded species-wide datasets but included 3 *C.*

coli and 5 *C. jejuni* sub-lineages for which rate estimates were obtained (**S3 and S4 Tables**).

Mean rate estimates were similar for all 3 *C. coli* lineages averaging at 7.82 x 10^{-4} s/s/y but

varied from 8.20 x 10^{-5} (ST-661 clonal complex) to 1.00 x 10^{-3} s/s/y (ST-22 clonal complex)

for *C. jejuni* (**S3 and S4 Tables**). In part because of the poor temporal signal in the species-

wide analyses and most sub-lineages, we developed an alternative method using paired

isolates.

Sampling matched isolate pairs allows estimation of the rate of nucleotide change

Nucleotide change (NC) is introduced into the bacterial genome by recombination resulting

from HGT, and point mutation. For clarification, we use the empirical term 'nucleotide

change' to describe any nucleotide variation resulting from these two processes, consistent

with previous studies [46]. Estimation of molecular clock rates requires comparison of

isolates from related, or preferably the same, lineages that have accumulated NCs over time.

To achieve this there is a necessary balance between maximizing the time between sampling

and accumulated NCs whilst ensuring comparisons are made between related strains.

Therefore, we plotted NC difference against time difference to determine criteria for choosing

comparable isolate pairs (**Fig 2**). The sample time difference was chosen to maximize the time

between sampling and the number of comparable pairs belonging to the same lineage. Pair

selection criteria were standardised for both species so that isolate pairs were included where

the sampling time difference was >8 years and there were <5000 SNPs between them (**Fig 2,**

S2 and S5 Tables). Based upon these criteria, there were 18 *C. coli* and 74 *C. jejuni* isolate

pairs (**S5 Table**). However, for consistency between the species we chose the 20 *C. jejuni*

pairs with the highest nucleotide identity, hence those with the strongest evidence of recent

common ancestry. Therefore 18 *C. coli* and 20 *C. jejuni* pairs comprised the dataset for NC

rate calibration. These belonged to the ST-21, ST-22, ST-45, ST-1332, ST-828 clonal

complexes and isolate pairs had a difference in sampling date of 8 to 11 years (*C. coli*) and 8

to 36 years (*C. jejuni*) (**Fig 3, S2 and S5 Tables**).

Estimation of a molecular clock rate requires that NCs accumulate over time, defined here as

NCs per site per year (s/s/y). It is also possible that branch shortening can occur where there

are fewer NCs in the more recent isolate of a pair. While not specifically describing branch

shortening, negative rates of NC have previously been observed [44]. In this study, 13 out 18

C. coli and 11 out of 20 *C. jejuni* isolate pairs exhibited branch lengthening, that is to say

more total NCs (within and outside recombined regions) were found in the more recent isolate

(**S2 and S6 Tables**). Only pairs having undergone measurable evolution (branch lengthening)

were included in further analysis of the accumulation of NCs over time. While this may

inflate our estimate of the NC rate, it was necessary to ensure a positive rate for calculating

the number of effective lineages. For measurably evolving isolate pairs, the total NC rate was

calculated as well as the rates within and outside of recombined regions (**Table 1, S7 Table**).

The mean NC rate for non-recombined regions was 6.36×10^{-6} and 8.45×10^{-6} s/s/y but

ranged from 1.60×10^{-6} – 1.50×10^{-6} and 1.00×10^{-7} – 3.60×10^{-5} s/s/y for *C. coli* and *C.*

jejuni respectively, or 11.46 and 13.53 average NCs per genome per year (s/g/y) (**Table 1**).

Rates of nucleotide change	*C. coli*			*C. jejuni*			Units**
	Min	Mean	Max	Min	Mean	Max	
Total*	1.7×10^{-6}	6.3×10^{-5}	3.0×10^{-4}	4.8×10^{-8}	8.8×10^{-6}	2.3×10^{-5}	s/s/y
S and N (exc. rec)	1.6×10^{-6}	6.4×10^{-6}	1.5×10^{-6}	1.0×10^{-7}	8.5×10^{-6}	3.6×10^{-5}	s/s/y
S (inc. rec)	4.9×10^{-6}	3.1×10^{-4}	1.8×10^{-3}	2.1×10^{-7}	1.9×10^{-6}	7.6×10^{-6}	s/s/y
S (exc. rec) (molecular clock)	2.1×10^{-7}	2.4×10^{-6}	7.7×10^{-6}	3.8×10^{-8}	3.4×10^{-6}	1.7×10^{-5}	s/s/y
N (inc. rec)	4.4×10^{-8}	2.4×10^{-5}	1.1×10^{-4}	5.0×10^{-8}	1.4×10^{-6}	4.8×10^{-6}	s/s/y
N (exc. rec)	4.8×10^{-7}	3.2×10^{-6}	7.8×10^{-6}	3.1×10^{-7}	4.8×10^{-6}	1.6×10^{-5}	s/s/y

Table 1. Average rate calibrations for *C. coli* and *C. jejuni*

*all NCs from including (inc) and excluding (exc) recombination (rec); **NCs per site per

year (*C. jejuni* = 1.6 Mbp, *C. coli* = 1.8 Mbp); S = synonymous NCs, N = nonsynonymous

NCs

Recombination drives molecular evolution in *Campylobacter*

NCs in coding sequence based on gene definitions in the reference *C. coli* (YH501) and *C.*

jejuni (NCTC 11168) isolate genomes introduced an average of 1569 and 242 NCs in *C. coli*

and *C. jejuni* paired genome datasets respectively. Of these, an average of only 222 (*C. coli*)

and 106 (*C. jejuni*) were inferred to be the result of point mutation, with the remainder

resulting from recombination (**S2 Table**). Recombination is therefore the major source of

sequence variation in both species (**Fig 4, S2 Table**), introducing nearly six times as many

NCs in *C. coli* than in *C. jejuni* – consistent with previous estimates based upon MLST [47].

The effects of recombination on effective genotypes over successive generations were

simulated for *C. coli* and *C. jejuni.* For both species, simulations provided results consistent

with observations using our method on real data (Fig S5) in several ways. First, simulation of

Campylobacter evolution under different recombination rates demonstrates how elevated

nucleotide change (due to recombination) effects the number genotypes in successive

generations. Second, the increase in the number of genotypes (and isolate pairs) is observed in

both real and simulated data with evidence that rate slows over time. Third, the number of

genotypes carried over to the next generation was higher for *C. coli* than *C. jejuni* in

simulations at different recombination rates.

To assess the effect of NCs on amino acid sequences we quantified non-synonymous (N) and

synonymous (S) NCs and determined the ratio per site (*dN/dS*) for all isolate pairs in

recombined and non-recombined sequence (**S2 Table**). Point mutation on average accounted

for an unequal amount of N and S polymorphism both within and between species (*C. coli,* N

= 99, S = 123; *C. jejuni,* N = 63, S = 43) (**S2 Table**). While recombination introduced many

more NCs than point mutation, in both species these were biased towards synonymous

changes. Specifically, around six times as many S than N NCs were introduced by

recombination in *C. coli* and approximately twice more in *C. jejuni* (*C. coli,* N = 546, S =

801; *C. jejuni,* N = 59, S = 77) (**S2 Table**). Overall, average *dN/dS* ratios were consistent

between species within recombined (*C. coli* 0.492, *C. jejuni* 0.490) and non-recombined (*C.*

coli 0.594, *C. jejuni* 0.509) portions of the genome. However, because of the relative

importance of recombination (*r/m* = 37.240 (*C. coli*), *r/m* = 5.098 (*C. jejuni*)), on average N

NCs were similar for *C. jejuni* from recombination and point mutation (59 and 63

respectively). However, recombination introduced 5.5 times more N NCs than point mutation

in *C. coli* (**S2 Table**). Variation in *dN/dS* was observed between isolate pairs but was mostly

indicative of purifying selection (*dN/dS*<1). Evidence of positive selection (*dN/dS*>1) was

only observed within recombined sequence in 6 isolate pairs (**S2 Table**). It is important to

note that, while *dN/dS* comparisons have been made between closely related bacteria within

the same species [48,49], this method was originally intended for between species

comparisons [50].

Additional analysis of the distribution of recombination events revealed that an average of

13% (*C. coli*) and 2% (*C. jejuni*) of the genome has undergone recombination in at least one

isolate pair since divergence from the common ancestor of each sub-tree (**S2 Table**).

Recombination was distributed across the genome in both species but was elevated in certain

regions of *C. coli* introducing more NCs at potential recombination hotspots [51]. However,

recombination remained the main source of variation in both species (**Fig 4**).

Molecular clock estimates for *C. coli* and *C. jejuni*

Molecular clock estimates require that NCs accumulate at a consistent rate over time. We

maximized the chance of identifying this signal in several ways. First, genomic variation

within recombined regions was discounted as multiple NCs can be introduced in a single

evolutionary event – distorting clock estimates [39,47,52]. Second, non-synonymous NCs

were discounted as selection may be more likely to influence the frequency of variation at

these sites. Third, only pairs in which the most recently sampled isolate contained more NCs

(branch lengthening) were used as they displayed measurable evolution over time. Based on

these criteria, a similar average molecular clock rate was obtained for *C. coli,* 2.4 x 10^{-6} s/s/y

(4.27 s/g/y), and *C. jejuni,* 3.4 x 10^{-6} s/s/y (5.42 s/g/y) (**Table 1**) but ranged from 2.1 x 10^{-7} –

7.7 x 10^{-6} and 3.8 x 10^{-8} – 1.7 x 10^{-5} s/s/y.

Coalescence and maintenance of lineages over time

Molecular clock estimates can vary within a population. Therefore the applicability of

generalized clocks depend upon how much of the population has been sampled. To quantify

this we estimated the average total NC rate (**μ**) (*C. coli* = 77.292 s/g/y, *C. jejuni* = 14.101

s/g/y), including all NCs within and outside recombined sequence. These rates were used to

determine the number of coalescences in the population at a given time point (here referred to

as '*effective lineages*') within the dataset. The maximum timeframe for comparison was 37

years for *C. coli* and 46 years for *C. jejuni* (short in evolutionary terms). This provided

information about the number of ancestral strains and the rate of turnover of lineages within

the dataset. The total number of potential pairs without accounting for genetic similarity (**Y**),

was equal to the square of the total number of isolates (n^2) divided by two (to avoid double

counting of isolate pairs), 180,600 and 1,663,488 for *C. coli* and *C. jejuni* respectively.

Having determined the total NC rate, we were able to predict the expected number of NCs

over a given period of time. For example, 14 in 1 year for *C. jejuni.* We then subsampled all

isolate pairs (**Y**) to determine how many isolate pairs had $\leq$14 NCs between them – 76 isolate

pairs. This is the possible number of isolate pairs that have arisen in 1 year. This process was

repeated for each time cut-off, up to a maximum of 37 and 46 years for *C. coli* and *C. jejuni*

respectively (**S8 Table**), to give the number of possible pairs for every time cut-off (*X*) (**Figs**

2B and 2D*)*. Dividing *Y/X* resulted in the number of coalescences (*effective lineages*) at a

given time interval in the past (*Z*) (**S8 Table**). For example, if the total NC rate was 14 s/g/y

and we were interested in the number of birthdays within 5 years of our dataset, we would

multiply the NC rate by 5 to result in 70 NCs of evolution over 5 years. The number of

potential pairs (*Y* = 1,663,488) / *possible pairs* (*X* = 174) = ~9,560 coalescences (ancestors)

within this time period (**S4 Fig**, **S8 Table**).

The number of effective lineages at a given time-point can also be interpreted as the number

of lineages that gave rise to those that are seen today. This provides valuable information

about how the population is maintained over time and the extent to which it has diversified.

For example, 1,263 *C. coli* lineages 37 years ago gave rise to an estimated 22,575 one year

ago and 4,726 *C. jejuni* lineages 46 years ago gave rise to 21,888 lineages one year ago. This

equates to an average increase in the number of effective lineages of 576 and 373 per year for

C. coli and *C. jejuni* respectively. For *C. jejuni* it is clear that a considerable proportion (22%)

of all lineages have been maintained throughout the 46 year sampling period and probably

much longer (**Fig 5**). In contrast, only 6% of all effective lineages were present in the *C. coli*

population 37 years ago. Perhaps the most striking finding is that the *C. coli* population has

rapidly diversified in recent years. For example, there has been an 800% increase in the

number of effective lineages in the last 10 years, over 3 times the rate of increase observed in

C. jejuni

(**Fig 5, S4 Fig**). This provides a basis for considering evolution in the wild but a longer

sample time frame and more varied collection of isolates will improve representation of the

natural *Campylobacter* population.

Discussion

The increasing availability of large genome datasets has great potential for improving molecular clock estimates in bacteria. However, significant challenges remain. While it is clear that the frequency of NCs can vary between different species and strains [6,7,10,24,44,53,54], the extent to which nucleotide variation represents an intrinsic molecular clock is often less apparent. Biological factors such as generation time, population size and recombination rate, and ecological factors including cellular responses to habitat variation or stress and the strength of natural selection, influence the rate at which NCs accumulate in populations [55]. Therefore, obtaining a robust molecular clock estimate from natural bacterial populations requires an appropriate sample frame and careful consideration of the nature of observed sequence variation.

In cases where there is a clear temporal signal among isolates, it may be possible to obtain a robust molecular clock estimate by applying models to large genome datasets [24]. However, analysing all *C. coli* and *C. jejuni* genomes gave a weak temporal signal. This is likely related to the population structure and biology of these organisms that is in stark contrast to many obligate human pathogens [24]. Consistent with many other zoonotic or environmental bacteria, *Campylobacter* is a diverse genus with multiple lineages (STs and clonal complexes) inhabiting multiple hosts/niches. This required a more targeted approach to microevolutionary analysis consistent with that used to investigate transmission in similarly variable organisms [21].

Sub-sampling within the isolate collection, sampled over 46 years, identified closely related pairs of isolates with divergent sampling dates. Clearly, calibration of the molecular clock

requires that NCs accumulate over time. This was not the case in all isolate pairs. In some

cases, the most recently sampled isolate had accumulated fewer NCs than the comparator

strain leading to a negative NC rate. A negative slope of the root-to-tip regression line has

been interpreted as evidence for a lack of temporal signal or for a large dispersion of the rate

of change [24,44]. Furthermore other studies have described the time-dependency of

molecular evolution [56,57] and it can be the case that deleterious NCs in the older isolate

have been purged leading to differences in long and short term molecular clock estimates

[48,58]. To simplify interpretation in our study, isolate pairs where the most recent isolate had

not accumulated NCs were excluded from the analysis. While this might inflate the NC rate

estimate, in organisms with complex ecology such as *Campylobacter,* it is also possible that

closely related isolates occupy different sub-niches and experience different selection

pressures even when sampled from the same host.

Returning to the birthday problem analogy, considering the number of isolate pairs

(equivalent to people with the same birthday) obtained from the original genome dataset can

provide clues about the extent of lineage diversity in the natural population. Using total NC

rates, we were able to assess the nature of coalescence across the sample time frame for each

species. The coalescence we refer to here is equivalent to the number of ancestral strains at a

particular time point (effective lineages) in the natural environment from which contemporary

strains emerged. Effective population size (Ne) is commonly used to reflect the number of

individuals in a population that contribute to subsequent generations [59]. This has been used

to investigate bacteria but contrasting approaches can provide different estimates depending

on the method used [53,60]. The idea of effective lineages, described in this study, is related

to Ne but is more specific for organisms that reproduce clonally. Rather than typical Ne

estimates for sexual populations, where the mating of two individuals is largely independent

of what happened in previous generations, the number of effective lineages in a bacterial

population reflects the number of distinct lineages that will survive and therefore contribute to

future generations. This provides information on the genetic inertia of the population, i.e. the

limitations for future evolutionary pathways based on the number of successful ancestors at a

particular point in time.

These analyses highlighted the importance of appropriate sampling when calibrating NC rates

and can help in determining the extent to which samples represent the population as a whole.

Specifically, by considering the number of coalescences in a random population, we can look

back through the sample time frame to estimate the number of effective lineages across a

randomly sampled dataset. For example, suppose we would like to know if our contemporary

isolates have a common ancestor in 1980. We know that a proportion of these ancestors gave

rise to the diversity we see today but many lineages would go extinct and therefore not

contribute [61]. Based on an average NC rate of 14 s/g/y for *C. jejuni*, there would be 560

NCs over 40 years total evolution between a strain pair. So, one can then ask how many pairs

are close enough genetically for that to be the case. This gives an estimate of the effective

number of ancestors in 1980 that gave rise to the contemporary dataset - equivalent to the

number of birthdays.

Our estimate of effective lineages aims to better account for both unobserved lineages and

lineages that co-exist over a long period of time. Both of these estimates are strongly affected

by the sample frame and, while we have relatively large isolate collections from livestock and

humans, the same cannot be said for the vast number of other potential host and

environmental sources. Therefore, our estimate of effective lineages is more useful as an

estimate of the number of ancestors that have given rise to the diversity of strains that we see

today. For *Campylobacter,* it is clear that multiple lineages have persisted over a long period

of time. This indicates that although the population is large, the strains are not turning over at

a particularly fast rate. The absence of lineage replacement is inconsistent with some models

of bacterial evolution that predict periodic population bottlenecks [62] but this can be

explained in several ways. First, it is possible that the 37/46 year sampling period in this study

is not sufficient time to out-compete a rival strain. Second, bacteria occupy different niches

that are sustained so strains are not in direct competition. Third, the fitness differences among

strains are not great enough for one lineage to out-compete another.

As well as the maintenance of multiple lineages, there is also evidence for variation in the

number of effective lineages that contributed to successive generations between the two major

pathogenic *Campylobacter* species. While, demographic inference and estimates of the

number of generations using BEAST were similar for *C. coli* and *C. jejuni* (**S3 and S4**

Tables), the number of effective lineages was shown to be consistently higher for *C. jejuni*

throughout much of the sample frame. Furthermore, there was evidence for a rapid increase in

the number of *C. coli* lineages that began around 8 years ago (**Fig 5**). The reason for this is

unclear. The average synonymous NC rate estimates were similar for *C. jejuni* and *C. coli,* 3.4

x 10^{-6} and 2.4 x 10^{-6} s/s/y respectively, equating to approximately 5.4 (*C. jejuni*) and 4.3 (*C.*

coli) NCs per genome per year. This is somewhat lower than previous estimates for *C. jejuni*

calculated from 7-locus MLST (2.79 x 10^{-5} s/s/y) [47] but is within the range of molecular

clock estimates calculated from genomic variation for *Enterococcus faecium* (9.35 x 10^{-6}

s/s/y) and *Y. pestis* (1.57 x 10^{-8} s/s/y) [44]. Although average synonymous NC rates were

consistent with other estimates, rates ranged from 2.10 x 10^{-7} to 7.70 x 10^{-6} s/s/y in *C. coli* and

3.80 x 10^{-8} to 1.70 x 10^{-5} s/s/y for *C. jejuni.* This implies uncertainty around the estimate.

However, rate heterogeneity among lineages is not uncommon in bacterial species [63]

potentially reflecting differences in the evolution and ecology of different species and strains.

While the average NC rate was consistent for *C. coli* and *C. jejuni,* the relative number of

NCs introduced by homologous recombination and mutation (*r/m*) differed markedly, with on

average 37-fold (*C. coli*), compared to 5-fold (*C. jejuni*), greater impact on sequence

variation. HGT is known to be an important driver of genome evolution in *Campylobacter*

[47,64] but these estimates are considerably higher than previous ones using 7-locus MLST

[11]. Recombination introduced nearly twice as many synonymous than non-synonymous

NCs, but even taking this into account, recombined sequence accounted for around 79% of all

non-synonymous variation. This highlights the importance of HGT in rapidly evolving

Campylobacter genomes and provides evidence that recombination may have been an

important factor in the recent diversification of *C. coli* [26,65,66], potentially associated with

an adaptive radiation [67,68] linked to the colonization of agricultural niches [69]. However,

this should be balanced against the evidence of purifying selection within recombined

sequence (*dN/dS* = 0.492 for *C. coli* and 0.49 for *C. jejuni)* and the removal of non-

synonymous NCs through negative selection [48].

Finally, throughout this study we have emphasized the importance of sampling so that

measures of molecular evolution are obtained by comparing recent samples with a true

ancestor. The uneven distribution of lineages within the population and the possibility that

they differ in key evolutionary measures (*r/m* and *dN/dS*), means that our molecular clock

estimate may not be applicable to all *Campylobacter* lineages [21,71-72]. Perhaps this is best

illustrated by considering two host-specialist *C. jejuni* lineages, one associated with chickens

and the other with cattle [8,26]. There are 19 billion chickens on earth compared to 1.3 billion

cattle [73] and *C. jejuni* colonizes up to 80% of chickens [74] with much lower rates in cattle.

As the efficiency by which natural selection acts on sequence variation is related to effective

population size [41], the rate of fixation and removal of NCs will be much faster in *C. jejuni*

in chickens. Furthermore, chickens have a higher body temperature than cattle therefore the

C. jejuni will grow faster, have a shorter generation time, and accumulate NCs at a higher rate

[9]. From this simple example, which ignores many important factors (eg. subniche structure,

host transition bottle necking, resident microbiome) it is clear molecular evolution can be

influenced by population-scale forces down to the physiology of the individual cell. The

approach employed in this study goes some way towards mitigating effects that confound

generalized molecular clock estimates. Focussing on well-defined closely related isolate pairs

inevitably reduces the number of comparisons from which the mean molecular clock rate is

estimated. However, consideration of the distribution of effective lineages within the

population is essential for identifying robust molecular clock estimates in environmental

bacteria with complex multi-host ecology and massive effective population sizes.

Materials and Methods

Isolate sampling, genome sequencing and assembly

The accuracy of molecular clock estimates are improved by analysing large numbers of isolates sampled over long time periods. While very large isolate genome collections exist for *Campylobacter* (see below)*,* many of these were sampled within the last 20 years. Therefore, to extend the sample time frame we assembled an isolate collection comprising 53 isolates sampled between 1978 and 1985 (12 *C. coli*, 41 *C. jejuni*) derived from multiple sources (human, duck, cattle, dog, turkey, wild bird and pig (**S1 Table**)). These samples were streaked onto mCCDA (PO0119A Oxoid Ltd, Basingstoke, UK) with CCDA Selective Supplement (SR0155E Oxoid Ltd, Basingstoke, UK) and incubated at 37˚C for 48h in a microaerobic atmosphere (85% N_2, 10% CO_2, and 5% O_2) using CampyGen Compact sachets (Thermo Fisher Scientific Oxoid Ltd, Basingstoke UK). Single colonies from each plate was then sub-cultured onto Mueller Hinton (MH) (CM0337 Oxoid Ltd, Basingstoke, UK) agar and grown for an additional 48h at 37˚C and stored in 20% glycerol stocks at -80˚C.

DNA was extracted using the QIAamp DNA Mini Kit (QIAGEN, Crawley, UK), according to manufacturer's instructions. DNA was quantified using a Nanodrop spectrophotometer before sequencing on an Illumina MiSeq sequencer using the Nextera XT library preparation kits with standard protocols. Paired end libraries were sequenced using 2×300 bp 3rd generation reagent kits (Illumina). Short read data was assembled using the *de novo* assembly algorithm, SPAdes (version 3.10.0 35) [75] generating an average of 49 contigs (range: 2 -115) for a total average assembled genome size of 1.69 Mbp (range: 1.62-1.80). The average N50 was 189,430 bp (range: 81,283-974,529). These isolate genomes were augmented with 1,783 *C. jejuni* and 589 *C. coli* genomes archived in BIGSdb [76] representing isolates sampled from

multiple sources (human, cattle, chicken, cat, dog, duck, environmental waters, farm

environments, geese, lamb, rabbit, sand, seal, wild birds, turkey, pig) between 1970 and 2016

(**S1 Table).** The total isolate collection comprised 2,425 *Campylobacter* genomes, including

C. jejuni belonging to 286 STs and 36 clonal complexes, and *C. coli* to 125 STs and 1 clonal

complex. For *C. coli*, much of the genetic variation is within three ancestral clades, thought to

have diversified before major introgressions with *C. jejuni* [66, 77-78]. However, isolates

from these clades are not a major cause of human infection. In fact, ~96% of all disease-

causing *C. coli* strains (PubMLST, 16/08/2021), belong to the (introgressed) ST-828 clonal

complex analysed in our study. For this reason we focused on the ST-828 complex [38]). All

assembled genomes and raw reads have been deposited in the NCBI repository associated

with BioProject: PRJNA524315. Individual accession numbers can be found in **S1 Table**.

Assembled genomes of all isolates used in the study are available in FigShare DOI:

10.6084/m9.figshare.7886810.

Simulating evolution in *Campylobacter* genomes

We performed forward-time simulations of neutral evolution on populations of *C. coli* and *C.*

jejuni using a Wright-Fisher model in order to test the impact of recombination and point

mutation on the effective genotypes over successive generations (time) using Bacmeta [79].

Starting population sizes of 1000 bacteria represented by 10 loci of length 1 kb were

simulated for 60,000 generations to represent ten years of *Campylobacter* doubling time [80].

Mutation rates were set at 1 x 10^{-5} and 1 x 10^{-6} base^{-1} generation^{-1} for *C. coli* and for *C. jejuni*

respectively and simulations were run three times with high (35.641), medium (13.058) and

low (0.191) *r/m* values. Recombination events were exchanged as complete loci (**S5 Fig**).

C. coli **and** *C. jejuni* **phylogenies and assessing temporal signal and 'clock-likeness'**

Phylogenies were constructed for 601 *C. coli* and 1,824 *C. jejuni* isolates (**S1 Table, Fig 1**).

Gene-by-gene alignments were produced with comparison to reference *C. coli* (YH501,

accession number NZ_CP015528.1) and *C. jejuni* (NCTC11168, accession number

NC_002163.1) isolate genomes, sampled in 2016 and 1977 respectively. These reference

genomes, belonging to the ST-828 and ST-21 clonal complexes, were chosen as those most

commonly used in *Campylobacter* comparative population genomics studies [81,82]. For

some bacteria the use of a lineage specific reference genome may increase the pool of core

genes identified. However, we do not to take this approach for two reasons. First, all *C. coli* in

this study belong to a single lineage, the ST-828 clonal complex, so there would be little

benefit to creating separate reference pangenomes for already closely related isolates. Second,

much of the advantage of creating lineage-specific pangenomes in *Campylobacter* is lost due

to extensive recombination between divergent lineages [47]. For example, after comparison of

all isolate genomes to the ST-21 complex NTCT11168 *C. jejuni* reference, the average

number of core genome SNPs identified for non-ST-21 complex strains (255) was not

significantly different compared to ST-21 complex strains (217) (T-test, Welch's correction,

p=0.683).

Homology to the reference genome (*C. coli* YH501; *C. jejuni* NCTC11168) was determined

using MAFFT, with default parameters of minimum nucleotide identity of 70% over >50% of

the gene and a BLAST-n word size of 20. Core genes (2,014 for *C. coli* and 1,668 for *C.*

jejuni), shared by all isolates within a species were concatenated and used to construct

Maximum likelihood (ML) trees using FastTree version 2.1.8 and the Generalised time-

reversible (*GTR*) model of nucleotide evolution [83]. Isolates were analysed to test for a

temporal signal of the accumulation of genetic variation over time (**S1 and S2 Figs**). This was

carried out prior to NC rate analysis using a phylogeny of genetic distances and sampling

dates, and root-to-tip regression implemented in the software TempEst v1.5.1 [84] with the

best fitting root selected to maximise the coefficient, R^2. Core genome phylogenies contained

dated-tip isolates sampled between 1970 and 2016 for *C. coli* and *C. jejuni*.

Root-to-tip regression analyses to identify a temporal signal in bacteria rely on the removal of

recombination as this is often the main source of genetic variation. Such analyses have been

performed on small genome collections (15-189 isolates) of multiple bacterial species [44].

However, for *Campylobacter*, this would not be sufficient to represent diversity within the

species as there are >40 known clonal complexes within *C. jejuni* alone. Using existing

methodology, it was not possible to exclude recombining regions and perform root-to-tip

regression on the entire *C. jejuni* (1824 isolates) and *C. coli* (601 isolates) genome collections.

To account for this, we also assessed temporal signal in the whole datasets after removal of

outlier genes that likely reflect recent recombination. Specifically, using a comparative gene-

by-gene approach [85], we determined the number of alleles per locus in all genomes through

comparison with the reference genomes (*C. coli* YH501*; C. jejuni* NCTC 11168). Three new

concatenated genome alignments were constructed for each species after removal of 1%, 5%

and 10% of the most variable loci (most/least alleles per locus) of respective core genome

length of 1522 (1.11 Mbp), 1398 (1.02 Mbp), 1238 (0.89 Mbp) (*C. coli*) and 1268 (1.2 Mbp),

1164 (1.04 Mbp), 1034 (0.91 Mbp) (*C. jejuni*) core genes. Consistent with analyses of

complete masked core genome alignments, TempEst analysis of reduced alignments (1%, 5%,

10%) also revealed poor temporal signal. Therefore, we conducted analyses on sub-lineages

within *C. coli* and *C. jejuni*.

In order to identify temporal signal using TempEst, individual subsets of *Campylobacter* lineages were chosen from our datasets for *C. coli* (ST-825, ST-827, ST-828, ST-830, ST-872, ST-1090 and ST-1541 of the clonal complex, ST-828) and *C. jejuni* (ST-1325, ST-1034, ST-206, ST-21, ST-22, ST-257, ST-353, ST-354, ST-42, ST-433, ST-45, ST-464, ST-48, ST-52, ST-574, ST-61, ST-658 and ST-661 clonal complexes). Core genes were identified for each sub-lineage by comparison to *C. coli* (YH501) and *C. jejuni (*NCTC 11168) reference genomes using MAFFT (as described above). Individual ML phylogenies were constructed from core gene alignments using FastTree. Recombined regions were inferred using ClonalFrameML with basic model parameters [86] and removed using a cfml-mask script and replaced with gaps (https://github.com/kwongj/cfml-maskrc). Recombination-masked trees of individual lineages were subsequently tested for temporal signal using TempEst software v1.5.1.

Bayesian evolutionary analysis

Analysis with BEAST2 v2.5.0 [45] was performed using a method previously described for *Campylobacter* [8]. SNP alignments were constructed from variable sites of the whole datasets of 601 (*C. coli*) and 1824 (*C. jejuni*) isolates and for the sub-lineages which exhibited the strongest temporal signal using snp-sites v2.5.1 [87]. A time-scale phylogeny was constructed using BEAST2 v2.5.0 [45] based on variable sites, using the GTR +G4 model of DNA substitution. The relaxed log-normal clock with Bayesian skyline model was used as previously described [8]. Input xml files were prepared using BEAUti2 v2.5.0 [45]. A prior on the clock rate was set as a log-normal distribution with a mean value of 1×10^{-6} mutations per site per year with a lower value of 1×10^{-8} and an upper value of 1×10^{-3}. Markov chains

were run for 50 million generations, sampled every 10,000 generations with the first

5,000,000 generations (10%) discarded as burn-in.

Selection of closely related isolate pairs

An ideal dataset for rate analysis would include isolate pairs with divergent sampling dates,

sufficient to measure NC rates over time, while remaining close enough (clustering on the

tree) to share reliable recent common ancestry. Furthermore we required as many pairs as

possible for confidence in average rates. In order to achieve this, pairwise nucleotide identity

and year between isolation date matrices were constructed separately for 601 (*C. coli*) and

1,824 (*C. jejuni*) isolates. Using a bespoke R script

(https://github.com/SionBayliss/CallandMolClock), the distribution of nucleotide identity was

determined for isolate pairs within sequential isolation date categories of 1 year or more (1-37

for *C. coli*, 1-46 for *C. jejuni*) by comparing every isolate to all other isolates (**Fig 2**). In each

analysis, isolates were used only once as the ancestral or derived strain. Large numbers of

divergent isolate pairs can be identified from distant time points reflecting genome evolution

over time (**Fig 2**). However, our analysis required inference of recent common ancestry

(<5,000 SNPs) to define isolate pairs. Specifically, we needed closely related isolates with a

large difference between sample dates. To provide a quantitative basis for selecting pairs we

compared the nucleotide identity of each isolate pair at given sampling time difference

thresholds. Isolates with sample dates >8 years apart (**Fig 2**) were chosen for calibrating a rate

of nucleotide change because there was an enrichment of closely related pairs at this

threshold. An arbitrary threshold of <5000 SNPs was selected for candidate isolate pairs to be

included in the NC rate calibration analysis based on the pair selection curves. The final

number of pairs used in the analysis included the 20 most closely match pairs for *C. coli* and

C. jejuni (**Fig 2A**).

Recombination and nucleotide change inference

The raw reads of genomes (**S1 Table**) of isolate pairs (**S2 Table**) were mapped to the

complete reference genomes: *C. coli* YH501 (accession: NZ_CP015528.1) and *C. jejuni*

NCTC 11168 (accession: NC_002163.1) using the BWA-MEM algorithm [88]. Variants were

called using Freebayes v1.1.0-dirty [89] and NC effects predicted and annotated using SnpEff

version 4.3 [90] (**S6 Table**). These tools were included in the haploid variant calling pipeline,

'snippy' v3.0 (https://github.com/tseemann/snippy). Core genome sub-tree alignments were

constructed using snippy-core. NCs introduced by point mutation and recombination were

inferred on the alignments using Gubbins v2.4.1 (default settings) [91] for each isolate pair

(**S6 Table**). The snippy pipeline was used to identify synonymous and non-synonymous NCs

within and outside of inferred recombinant regions [90]. *dN/dS* ratios were calculated for sites

across the core genome using the synonymous/non-synonymous analysis program (SNAP)

v2.1.1 based on the Nei and Gojobori 1986 method [92,93] (www.hiv.lanl.gov). Rates were

calculated by reconstructing the NCs on internal branches (inferred using Gubbins) leading to

the MRCA shared by a pair of isolates. By quantifying point mutation and recombination and

synonymous and nonsynonymous NCs per branch, we were able to infer different molecular

evolution rate estimates based on the difference in isolation time between isolates pairs (**S4**

Fig C)**.** These included (i) the total NC rate, used to calculate the number of effective lineages

and (ii) the rate of accumulation of synonymous NCs occurring outside of recombinant

regions, used to estimate the molecular clock. Hotspots of recombination occurring across

multiple isolate pairs were observed.

Estimating the number of coalescences at yearly intervals (Birthday problem)

To consider the extent to which a given sample set represented genetic diversity within the population we developed a pipeline that calculated the number of coalescences (*effective lineages,* $\mathbf{Z}$) at yearly time intervals ($Z1$, $Z2$, $Z3$….Zn) within the datasets. This is described by the equation $\mathbf{Z} = \mathbf{Y/X,}$ Where: $\mathbf{Y}$ = all potential isolate pairs ($n^2/2$); $\mathbf{X}$ = the number of possible pairs for each time interval (*t1, t2,t3….tn*) that is less than the predicted number of NCs that have occurred over a given time interval (*$\mu(t(1-n))$*); $\boldsymbol{\mu}$ = rate of nucleotide change; *t* = time interval between sampling dates, 1-46 and 1-37 years for *C. jejuni* and *C. coli* respectively. The resultant $\mathbf{Z}$ value for each time period is the estimated number of effective lineages (Birthdays) at each time cut-off, equivalent to the number of lineages sharing a common ancestor at a particular time interval (**S4 Fig**).

Acknowledgements

All high-performance computing was performed on MRC CLIMB, funded by the Medical Research Council (MR/L015080/1). This publication made use of the PubMLST website (http:// pubmlst. org/) developed by Keith Jolley and Martin Maiden [38] and sited at the University of Oxford. The development of that website was funded by the Wellcome Trust.

Author contributions

JKC, SKS and DF designed the study and wrote the paper with BP. JKC, BP performed genomic analysis with input from HT, EM, SCB, EJF and JC. MB provided historical collection of isolates that EB cultured isolates for sequencing. EM, MDH and BP sequenced and assembled genomes. All authors contributed and approved the final manuscript.

References

1. Kimura M. Evolutionary rate at the molecular level. Nature 1968;217(5129):624-626.

2. Kuo CH, Ochman H. Inferring clocks when lacking rocks: the variable rates of molecular evolution in bacteria. Biol Direct. 2009;4:35.

3. Didelot X, Walker AS, Peto TE, Crook DW, Wilson DJ. Within-host evolution of bacterial pathogens. Nat Rev Microbiol. 2016;14:150-162.

4. Kidgell C, Reichard U, Wain J, Linz B, Torpdahl M, Dougan G, Achtman M. Salmonella typhi, the causative agent of typhoid fever, is approximately 50,000 years old. Infect Genet Evol. 2002;2(1):39–45.

5. Mutreja A, Kim DW, Thomson NR, Connor TR, Lee JH, Kariuki S, Croucher NJ,

Choi SY, Harris SR, Lebens M, et al. Evidence for several waves of global

transmission in the seventh cholera pandemic. Nature 2011;477(7365):462–465.

6. Mcadam PR, Templeton KE, Edwards GF, Holden MTG, Feil EJ, Aanensen DM,

Bargawi HJA, Spratt BG, Bentley SD, Parkhill J, et al. Molecular tracing of the

emergence, adaptation, and transmission of hospital-associated methicillin-resistant

Staphylococcus aureus. Proc Natl Acad Sci U S A. 2012;109(23):9107–9112.

7. Cui Y, Yu C, Yan Y, Li D, Li Y, Jombart T, Weinert LA, Wang Z, Guo Z, Xu L, et al.

Historical variations in mutation rate in an epidemic pathogen, Yersinia pestis. Proc

Natl Acad Sci U S A. 2013;110(2):577–582.

8. Mourkas E, Taylor AJ, Meric G, Bayliss SC, Pascoe B, Mageiros L, Calland JK,

Hitchings MD, Ridley A, Vidal A, et al. Agricultural intensification and the evolution

of host specialism in the enteric pathogen Campylobacter jejuni. PNAS

2020;117(20):11018-11028.

9. Weller C, Wu M. A generation-time effect on the rate of molecular evolution in

bacteria. Evolution (N Y). 2015;69(3):643–652.

10. Gibson B, Wilson D, Feil E, Eyre-Walker A. The Distribution of Bacterial Doubling

Times in the Wild. Proc Biol Sci. 2018;285(1880).

11. Vos M, Didelot X. A comparison of homologous recombination rates in bacteria and

archaea. ISME J. 2009;3(2):199–208.

12. Bromham L. Why do species vary in their rate of molecular evolution? Biol Lett.

2009;5:401-404.

13. Eyre-Walker A, Keightley PD. The distribution of fitness effects of new mutations.

Nat Rev Gen. 2007;8:610-618.

14. Gibson B, Eyre-Walker A. Investigating evolutionary rate variation in bacteria. J Mol
Evol. 2019;87:317-326.

15. Drummond AJ, Pybus OG, Rambaut A, Forsberg R, Rodrigo AG. Measurably
evolving populations. Trends Ecol Evol. 2003;18:481–488.

16. Biek R, Pybus OG, Lloyd-Smith JO, Didelot X. Measurably evolving pathogens in the
genomic era. Trends Ecol Evol. 2015;30(6):306–313.

17. Drummond AJ, Rambaut A. BEAST: Bayesian evolutionary analysis by sampling
trees. BioMed Cent. 2007;7(214):1–8.

18. Suchard MA, Lemey P, Baele G, Ayres DL, Drummond AJ, Rambaut A. Bayesian
phylogenetic and phylodynamic data integration using BEAST 1.10. Virus Evol
2018;8:4(1):vey016.

19. Arnold BJ, Hanage WP. Longitudinal samples of bacterial genomes potentially bias
evolutionary   analyses.   Unpublished   data   bioRxiv   2017
https://www.biorxiv.org/content/10.1101/103465v1, last accessed October 18, 2020

20. Maixner F, Krause-Kyora B, Turaev D, Herbig A, Hoopmann MR, Hallows JL,
Kusebauch U, Egarter Vigl E, Malfertheiner P, Megraud F, et al. The 5,300-year-old
Helicobacter pylori genome of the Iceman HHS Public Access. Science
2016;351(6269):162–165.

21. Didelot X, Eyre DW, Cule M, Ip CLC, Ansari MA, Griffiths D, Vaughan A,
O'Connor L, Golubchik T, Batty EM, et al. Microevolutionary analysis of Clostridium
difficile genomes to investigate transmission. Genome Biol. 2012;13(12):R118.

22. Walker TM, C Ip CL, Harrell RH, Evans JT, Kapatai G, Dedicoat MJ, Eyre DW,
Wilson DJ, Hawkey PM, Crook DW, et al. Whole-genome sequencing to delineate

726 Mycobacterium tuberculosis outbreaks: a retrospective observational study. Lancet

727 Infect Dis. 2013;13:137–146.

728 23. Mathers AJ, Stoesser N, Sheppard AE, Pankhurst L, Giess A, Yeh AJ, Didelot X,

729 Turner SD, Sebra R, Kasarskis A, et al. Klebsiella pneumoniae Carbapenemase

730 (KPC)-Producing K. pneumoniae at a Single Institution: Insights into Endemicity

731 from Whole-Genome Sequencing. Antimicrob Agents Chemother. 2015;59:1656–

732 1663.

733 24. Menardo F, Duchêne S, Brites D, Gagneux S. The molecular clock of Mycobacterium

734 tuberculosis. PLoS Pathog. 2019;15(9):e1008067.

735 25. Waldenström J, Broman T, Carlsson I, Hasselquist D, Achterberg RP, Wagenaar JA,

736 Olsen B. Prevalence of Campylobacter jejuni, Campylobacter lari, and Campylobacter

737 coli in different ecological guilds and taxa of migrating birds. Appl Environ

738 Microbiol. 2002;68(12):5911–5917.

739 26. Sheppard SK, Colles FM, Mccarthy ND, Strachan NJC, Ogden ID, Forbes KJ, Dallas

740 JF, Maiden MCJ. Niche segregation and genetic structure of Campylobacter jejuni

741 populations from wild and agricultural host species. Mol Ecol. 2011;20(16):3484–

742 3490.

743 27. Bronowski C, James CE, Winstanley C. Role of environmental survival in

744 transmission of *Campylobacter jejuni*. FEMS Microbiol Lett. 2014;356(1):8–19.

745 28. Cody AJ, McCarthy ND, Bray JE, Wimalarathna HML, Colles FM, Jansen van

746 Rensburg MJ, Dingle KE, Waldenstrom J, Maiden MCJ. Wild bird-associated

747 Campylobacter jejuni isolates are a consistent source of human disease, in

748 Oxfordshire, United Kingdom. Env Micro Reports. 2015;7(5)782-788.

29. Sheppard SK, Maiden MCJ. The evolution of Campylobacter jejuni and

Campylobacter coli. Cold Spring Harb Perspec Biol. 2015;7(8):a018119.

30. Wilson DJ, Gabriel E, Leatherbarrow AJH, Cheesbrough J, Gee S, Bolton E, Fox A,

Fearnhead P, Hart CA, Diggle PJ. Tracing the Source of Campylobacteriosis. PLoS

Genet. 2008;4(9):e1000203.

31. Sheppard SK, Dallas JF, Strachan NJC, MacRae M, McCarthy ND, Wilson DJ,

Gormley FJ, Falush D, Ogden ID, Maiden MCJ, et al. Campylobacter genotyping to

determine the source of human infection. Clin Infect Dis. 2009;48(8):1072–1078.

32. Strachan NJC, Gormley FJ, Rotariu O, Ogden ID, Miller G, Dunn GM, Sheppard SK,

Dallas JF, Reid TMS, Howie H, et al. Attribution of *Campylobacter* Infections in

Northeast Scotland to Specific Sources by Use of Multilocus Sequence Typing. J

Infect Dis. 2009;199(8):1205–1208.

33. Dearlove BL, Cody AJ, Pascoe B, Méric G, Wilson DJ, Sheppard SK. Rapid host

switching in generalist Campylobacter strains erodes the signal for tracing human

infections. 2016;ISME J. 10(3):721–729.

34. Rosner BM, Schielke A, Didelot X, Kops F, Breidenbach J, Willrich N, Gölz G, Alter

T, Stingl K, Josenhans C, et al. A combined case-control and molecular source

attribution study of human Campylobacter infections in Germany, 2011–2014. Sci

Rep. 2017;7(1):5139.

35. Thépault A, Méric G, Rivoal K, Pascoe B, Mageiros L, Touzain F, Rose V, Béven V,

Chemaly M, Sheppard SK. Genome-Wide Identification of Host- Segregating

Epidemiological Markers for Source Attribution in Campylobacter jejuni. Appl

Environ Microbiol. 2017;83(7).

36. Marin J, Hedges SB. Undersampling genomes has biased time and rate estimates

throughout the tree of life. Mol Biol Evol. 2018;35(10):2595.

37. Mathis, FH. A generalized birthday problem. SIAM review. 1991;33(2):265-270.

38. Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics:

BIGSdb software, the PubMLST.org website and their applications. Wellcome Open

Res. 2018;24(3):124.

39. Didelot X, Falush D. Inference of Bacterial Microevolution Using Multilocus

Sequence Data. Genetics. 2007;175(3):1251–1266.

40. Kimura M. Molecular evolutionary clock and the neutral theory. J Mol Evol.

1987;26(1–2):24–33.

41. Gojobori T, Moriyama EN, Kimura M. Molecular clock of viral evolution, and the

neutral theory. Proc Natl Acad Sci U S A. 1990;87(24):10015-10018.

42. Rieux A, Balloux F. Inferences from tip-calibrated phylogenies: a review and a

practical guide. Mol Ecol. 2016;25(9):1911–1924.

43. Sheppard SK, Cheng L, Méric G, De Haan CPA, Llarena AK, Marttinen P, Vidal A,

Ridley A, Clifton-Hadley F, Connor TR, et al. Cryptic ecology among host generalist

Campylobacter jejuni in domestic animals. Mol Ecol. 2014;23(10):2442–2451.

44. Duchêne S, Holt KE, Weill F-X, Le Hello S, Hawkey J, Edwards DJ, Fourment M,

Holmes EC. Genome-scale rates of evolutionary change in bacteria. Microb Genomic.

2016;2(11).

45. Bouckaert R, Vaughan TG, Barido-Sottani J, Duchene S, Fourment M, Gavryushkina

A, Heled J, Jones G, Kuhnert D, De Maio N, et al. BEAST 2.5: An advanced software

platform   for   Bayesian   evolutionary   analysis.   PLoS   Comput   Biol.

2019;15(4):e1006650.

46. Ho SYW, Phillips MJ, Cooper A, Drummond AJ. Time dependency of molecular rate

estimates and systematic overestimation of recent divergence times. Mol Biol Evol.

2005;22(7):1561-1568.

47. Wilson DJ, Gabriel E, Leatherbarrow AJH, Cheesbrough J, Gee S, Bolton E, Fox A,

Hart CA, Diggle PJ, Fearnhead P. Rapid evolution and the importance of

recombination to the gastroenteric pathogen Campylobacter jejuni. Mol Biol Evol.

2009;26(2):385–397.

48. Rocha EPC, Smith JM, Hurst LD, Holden MTG, Cooper JE, Smith NH, Feil EJ.

Comparisons of dN/dS are time dependent for closely related bacterial genomes. J

Theor Biol. 2006;239:226–235.

49. Castillo-Ramirez S, Harris SR, Holden MTG, He M, Parkhill J, Bentley SD, Feil EJ.

The impact of recombination on dN/dS within recently emerged bacterial clones.

PLoS Pathog. 2011;(7):e1002129.

50. Kryazhimskiy S, Plotkin JB. The population genetics of dN/dS. PLoS genet.

2008;4(12), e1000304.

51. Yahara K, Didelot X, Ansari MA, Sheppard SK, Falush D. Efficient inference of

recombination hot regions in bacterial genomes. Molecular Biology and Evolution

2014;31(6): 1593-1605.

52. Croucher NJ, Harris SR, Fraser C, Quail MA, Burton J, Van Der Linden M, Mcgee L,

Von Gottberg A, Song JH, Ko KS, et al. Rapid Pneumococcal Evolution in Response

to Clinical Interventions. Science 2011;331(6016):430–434.

53. von Mering C, Hugenholtz P, Raes J, Tringe SG, Doerks T, Jensen LJ, Ward N, Bork

P. Quantitative Phylogenetic Assessment of Microbial Communities in Diverse

Environments. Science 2007;315(5815):1126–1130.

54. Li S-J, Hua Z-S, Huang L-N, Li J, Shi S-H, Chen L-X, Kuang J-L, Liu J, Hu M, Shu

W-S. Microbial communities evolve faster in extreme environments. Sci Rep.

2015;4(1):6205.

55. Denamur E, Matic I. Evolution of mutation rates in bacteria. Mol Microbiol.

2006;60(4).

56. Ho SYW, Shapiro B, Phillips MJ, Cooper A, Drummond AJ, Sullivan J. Evidence for

Time Dependency of Molecular Rate Estimates. Sullivan J, editor. Syst Biol.

2007;56(3):515–522.

57. Ho SYW, Lanfear R, Bromham L, Phillips MJ, Soubrier J, Rodrigo AG, Cooper A.

Time-dependent rates of molecular evolution. Mol Ecol. 2011;20(15):3087–3101.

58. Duchêne S, Holmes EC, Ho SYW. Analyses of evolutionary dynamics in viruses are

hindered by a time-dependent bias in rate estimates. Proceedings Biol Sci.

2014;281(1786).

59. Kirchberger PC, Schmidt ML, Ochman H. The ingenuity of bacterial genomes. Annu

Rev Microbiol. 2020;8;74:815-834.

60. Cui Y, Yang X, Didelot X, Guo C, Li D, Yan Y, Zhang Y, Yuan Y, Yang H, Wang J

et al. Epidemic clones, oceanic gene pools, and eco-LD in the free-living marine

pathogen Vibrio parahaemolyticus. Mol Biol Evol. 2015;32(6):1396-410.

61. Louca S, Shih PM, Pennell MW, Fischer WW, Parfrey LW, Doebeli M. Bacterial

diversification through geological time. Nat Ecol Evol. 2018;2, 1458-1467.

62. Koeppel A, Perry EB, Sikorski J, Krizanc D, Warner A, Ward DM, Rooney AP,

Brambilla E, Connor N, Ratcliff RM, et al. Identifying the fundamental units of

bacterial diversity: A paradigm shift to incorporate ecology into bacterial systematics.

PNAS 2008;105(7):2504-2509.

63. den Bakker HC, Didelot X, Fortes ED, Nightingale KK, Wiedman M. Lineage

specific recombination rates and microevolution in Listeria monocytogenes. BMC

Evol Biol. 2008;8;277. doi: 10.1186/1471-2148-8-277.

64. Sheppard SK, Dallas JF, Wilson DJ, Strachan NJC, McCarthy ND, Jolley KA, Colles

FM, Rotariu O, Ogden ID, Forbes KJ, et al. Evolution of an Agriculture-Associated

Disease Causing Campylobacter coli Clade: Evidence from National Surveillance

Data in Scotland. Hartskeerl RA, editor. PLoS One. 2010;5(12):e15708.

65. Sheppard SK, McCarthy ND, Falush D, Maiden MCJ. Convergence of Campylobacter

species: implications for bacterial evolution. Science 2008;320(5873):237-239.

66. Sheppard SK, Didelot X, Jolley KA, Darling AE, Pascoe B, Meric G, Kelly DJ, Cody

A, Colles FM, Strachan NJC, et al. Progressive genome-wide introgression in

agricultural Campylobacter coli. Mol Ecol. 2012;22(4)1051-1064.

67. Rainey PB, Travisano M. Adaptive radiation in a heterogeneous environment. Nature.

1998;394(6688):69-72.

68. Flohr RCE, Blom CJ, Rainey PB, Beaumont HJE. Founder niche constrains

evolutionary adaptive radiation. PNAS. 2013;110(51):20663-20668.

69. Thakur S, Morrow WEM, Funk JA, Bahnson PB, Gebreyes WA. Molecular

epidemiologic investigation of Campylobacter coli in swine production systems, using

multilocus sequence typing. Appl Environ Microbiol. 2006;72(8):5666-5669.

70. Croucher NJ, Finkelstein JA, Pelton SI, Mitchell PK, Lee GM, Parkhill J, Bentley SD,

Hanage WP, Lipsitch M. Population genomics of post-vaccine changes in

pneumococcal epidemiology. Nat Genet. 2013;45:656–663.

71. Didelot X, Nell S, Yang I, Woltemate S, Van Der Merwe S, Suerbaum S. Genomic

evolution and transmission of Helicobacter pylori in two South African families. Proc

Natl Acad Sci U S A. 2013;110(34):13880–13885.

72. Everitt RG, Didelot X, Batty EM, Miller RR, Knox K, Young BC, Bowden R, Auton

A, Votintseva A, Larner-Svensson H, et al. Mobile elements drive recombination

hotspots in the core genome of Staphylococcus aureus. Nat Commun. 2014;5(3956).

73. Bar-On YM, Phillips R, Milo R. The biomass distribution on Earth. Proc Natl Acad

Sci. 2018;115(25):6506–6511.

74. Dhillon AS, Shivaprasad HL, Schaberg D, Wier F. Campylobacter jejuni in broiler

chickens. Avian Diseases. 2006;50(1):55-58.

75. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,

Nikolenko SI, Pham S, Prjibelski AD, et al. SPAdes: A new genome assembly

algorithm   and   its   applications   to   single-cell   sequencing.   J   Comput   Biol.

2012;19(5):455-477.

76. Jolley KA, Maiden MC. BIGSdb: Scalable analysis of bacterial genome variation at

the population level. BMC Bioinformatics. 2010;11:595.

77. Sheppard SK, McCarthy ND, Falush D, Maiden MCJ. Convergence of Campylobacter

species: implications for bacterial evolution. Science 2008. 320(5873):237-239.

78. Skarp-de Haan CPA, Culebro A, Schott T, Revez J, Schweda EKH, Hanninen ML, et

al. Comparative genomics of unintrogressed Campylobacter coli clades 2 and 3. BMC

Genomics 2014. 15(129)

79. Sipola A, Marttinen P, Corander J. Bacmeta: a simulator for genomic evolution in

bacterial metapopulations. Bioinformatics 2018;34(13):2308-2310.

80. Konkel ME, Christensen JE, Singh Dillon A, Lane AB, Hare-Sanford R, Schaberg

DM, Larson CL. Campylobacter jejuni strains compete for colonisation in broiler

chicks. Appl Environ Microbiol 2001. 73(7):2297-2305.

81. Ghatak S, He Y, Reed S, Strobaugh Jr T, Irwin P. Whole genome sequencing and

analysis of Campylobacter coli YH502 from retail chicken reveals a plasmid-borne

type VI secretion system. Genom Data. 2017;11:128-131.

82. Pascoe B, Williams LK, Calland JK, Meric G, Hitchings MD, Dyer M, Ryder J, Shaw

S, Lopes BS, Chintoan-Uta C, et al. Domestication of Campylobacter jejuni NCTC

11168. Microb Genom. 2019;5(7):e000279.

83. Price MN, Dehal PS, Arkin AP. FastTree 2 – Approximately Maximum-Likelihood

Trees for Large Alignments. Poon AFY, editor. PLoS One. 2010;5(3):e9490.

84. Rambaut A, Lam TT, Carvalho LM, Pybus OG. Exploring the temporal structure of

heterochronous sequences using TempEst (formerly Path-O-Gen). Virus Evol.

2016;2(1).

85. Sheppard SK, Jolley KA, Maiden MCJ. A gene-by-gene approach to bacterial

population genomics: whole genome MLST of Campylobacter. 2012. Genes 3(2):261-

277.

86. Didelot X, Wilson DJ. ClonalFrameML: Efficient inference of recombination in whole

bacterial genomes. PLoS Comput Biol. 2015;11(2): e1004041.

87. Page AJ, Taylor B, Delaney AJ, Soares J, Seeman T, Keane JA, Harris SR. SNP-sites:

rapid efficient extraction of SNPs from multi-FASTA alignments. Microb Genom.

2016;2(4):e000056.

88. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. 2013;Unpublished data, arXiv:1303.3997v2 [q-bio.GN]. Last accessed on

October 18 2020

89. Garrison E, Marth G. Haplotype-based variant detection from short-read sequencing.

2012;Unpublished data, arXiv preprint arXiv:1207.3907 [q-bio.GN]. Last accessed on

November 24 2020.

90. Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Luan Wang SJL, Ruden DM. A

program for annotating and predicting the effects of single nucleotide polymorphisms,

SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3.

Fly. 2012;6(2):80–92.

91. Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, Parkhill J,

Harris SR. Rapid phylogenetic analysis of large samples of recombinant bacterial

whole genome sequences using Gubbins. Nucleic Acids Res. 2015;43(3):e15.

92. Nei M, Gojobori T. Simple methods for estimating the numbers of synonymous and

nonsynonymous nucleotide substitutions. Mol Biol Evol. 1986;3(5):418-426.

93. Korber B. HIV Signature and Sequence Variation Anaylsis. Computational Analysis

of HIV Molecular Sequences, Allen G. Rodrigo and Gerald H. Learn, eds. Dordrecht,

Netherlands: Kluwer Academic Publishers, 2000. Chapter 4, pages 55-72.

Supporting Information

S1 Table. Isolate list information

S2 Table. Effects of NC from point mutation and recombination on isolate pairs

S3 Table. TempEst root-tip regression and BEAST analysis estimates (*C. coli*)

S4 Table. TempEst root-to-tip regression and BEAST analysis estimates (*C. jejuni*)

S5 Table. List of all possible pairs >8 years apart and <5000 NCs in difference for both *C. jejuni* and *C. coli*

S6 Table. All annotated NC effects

S7 Table. Individual rate of nucleotide change estimates per pair for *C. jejuni* and *C. coli* (SNPs/year)

S8 Table. *C. coli* and *C. jejuni* "birthday problem" data and estimates of coalescence across sample time frame

Fig 1. Little evidence of clustering of isolate sampling dates in *Campylobacter* phylogenies. Maximum likelihood (ML) core genome phylogenetic trees of *C. coli* (A) (n = 601) and *C. jejuni* (B) (n = 1824) constructed using FastTree version 2.1.8 [79] and the *GTR* model of nucleotide evolution. Both phylogenies show the distribution of the sample time frame used in this study with major *Campylobacter* clonal complexes (CCs) identified and terminal nodes coloured according to isolation decade (orange = 1970s, yellow = 1980s, white = 1990s, green = 2000s, blue = 2010s). Scale bars represent the estimated number of NCs per site. Terminal nodes sampled from different decades can be seen scattered throughout both trees with little evidence of clustering by decade. Isolates sampled from the 2000s and 2010s are most abundant within each dataset. The position of the *C. jejuni* (NCTC11168) and *C. coli* (YH501) reference genomes are indicated on the phylogeny. These were sampled in 1977 and 2016 respectively.

Fig 2. Pair selection criteria curves for inclusion in rate estimates. Visual representation of possible pairs of isolates at all time cut-offs across the sample time frame for *C. jejuni* (B)

and *C. coli* (D). As time difference between pairs increases, distinguishing between individual

curves becomes distorted. Therefore, a selection of years were plotted (A and C) (black = all

pairs >1 year difference, blue = >2 years, pink = >4 years, purple = >6 years, orange = >8

years, red = >10 years, green = >15 years). All isolates were paired with the nearest isolate

(genetic distance), matched according to difference in year of isolation (coloured lines) for

both *C. jejuni* (A) and *C. coli* (C) (orange line). Dashed boxes (A and C) show magnified

images of the closest pairs from all curves. The 20 and 18 pairs used in rate calibration are

highlighted by a black oval (A) and every orange pair in dashed box (C). Grey scale bars (B

and D) indicate the time difference cut-off of each curve for every time point in the sample

date frame.

Fig 3. Scatter plots of individual pair NC rates for *C. coli* and *C. jejuni*. Three different

NC rates were determined per pair for *C. coli* (grey circles) (18 pairs) and *C. jejuni* (black

circles) (20 pairs): (A) all S (synonymous) and N (nonsynonymous) NCs, excluding those in

recombined regions; (B) S NCs only, excluding those in recombined regions (molecular

clock); (C) all NCs including those resulting from recombination (total NC rate). Variation

was gratest among total NC rates. This demonstrates the impact of recombination, introducing

the majority of NC's into *Campylobacter* genomes. All remaining information on isolate pairs

can be found in S2 Table.

Fig 4. Mutation and recombination in *C. coli* and *C. jejuni*. Average genome-wide NC

positions (red dots = synonymous NCs, blue dots = nonsynonymous NCs) per isolate pair in

relation to inferred recombined regions (grey blocks). Each plot represents one pair of isolates

considered in rate calibration for *C. coli* (A) and *C. jejuni* (B) and are ordered according to S2

Table. *y axis* = number of NCs in relation to particular bp position of the reference genome

(*C. coli* = YH501, *C. jejuni* = NCTC11168) and varies between pairs. *x axis* = position of

reference genome in bins of 10,000 bp. The cladogram shows the relatedness of isolate pairs

based on nucleotide identity, scale bar indicates NCs per site. It is evident from both A and B

that recombination is the main source of variation in *C. coli* and *C. jejuni*.

Fig 5. Lineage expansion in *C. jejuni* and *C. coli*. (A) Number of effective lineages (y axis)

at each time point within the sample time frame (x axis) for *C. coli* (grey) and *C. jejuni*

(black). (B) Diagrammatic representation of lineage expansion in *C. coli* and *C. jejuni*

showing contrasting lineage diversification scenarios.

S1 Fig. Root-to-tip linear regression of *C. coli* implemented in the software, TempEst.

Root-to-tip genetic distance (y axis) is correlated with sampling time (x axis) for phylogenies

of (A) 601 *C. coli* and (B) 8 sub-lineages of the ST-828 clonal complex. Only 3 out of 8 sub-

lineages had strong temporal signal ($R^2 > 0.5$).

S2 Fig. Root-to-tip linear regression of *C. jejuni* implemented in the software, TempEst.

Root-to-tip genetic distance (y axis) is correlated with sampling time (x axis) for phylogenies

of (A) 1824 *C. jejuni* and (B) 18 sub-lineages representing *C. jejuni* clonal complexes. Only 5

out of 18 sub-lineages had strong temporal signal ($R^2 > 0.5$).

1003

S3 Fig. Root-to-tip linear regression of *C. coli* and *C. jejuni* using TempEst software,

after removal of the most variable core genes. Core gene alignments were constructed from

601 (*C. coli*) and 1824 (*C. jejuni*) isolates. The most and least 1, 5 and 10% of variable loci

(alleles/loci) were removed and phylogenies were analysed with Tempest. Root-to-tip genetic

distance (y axis) is correlated with sampling time (x axis) to reveal poor temporal signal ($R^2 <$

0.5) in all six scenarios.

1010

S4 Fig. Methods for calculating the number of effective lineages within the population.

(A) The total number of *C. jejuni* and *C. coli* isolates in the population and all potential

pairwise comparisons between putative ancestral (black) and contemporary (white) strains to

give the total number of potential isolate pairs, Y. (B) Isolate pair selection based on

divergent sampling date (>8 years) and a nucleotide identity threshold <5000 SNPs. (C) Total

rate of nucleotide change (μ) calculated for all chosen pairs. The rate of accumulation of all

synonymous (S), nonsynonymous (N) NCs, within (rec) and outside (mut) of recombined

regions, was estimated since the most recent common ancestor (MRCA, red circle). The

difference in NCs between each pair was divided by the difference in isolation years to give

μ. (D) The total NC rate was used to estimate the number of NCs that were to accumulate

over a time period and the number of possible isolate pairs at given time intervals (t*1,*

t2,t3....tn) for each species.

1023

S5 Fig. Simulated migration of *Campylobacter* genotypes over successive generations

using Bacmeta software. Plots show the increase in the number of genotypes (y axis) over

60,000 generations (x axis) representing ten years of *Campylobacter* doubling time for *C. coli*

(**A**) and *C. jejuni* (**B**). Forward simulations were run with mutation rates reflecting estimates

from this study and three different *r/m* values (high (35.641) (dotted line), medium (13.058)

(dashed line), low (0.191) (block line)) to monitor the effects of recombination on genotype

frequency from one generation to the next. Simulated data was compared to the number of

possible isolate pairs with fewer SNPs than predicted based on the total NC rate (X) at a given

time for *C. coli* and *C. jejuni* (red dashed line).

1033

1034

Fig 1.

A

C. coli CC828

ref: YH501

B

CC257

CC443

CC353

CC45

CC61

CC22

ref: NCTC 11168

CC21

0.0005

0.001

	1970s
	1980s
	1990s
	2000s
	2010s

Fig 2.

Fig 3.

Fig 4.

A *C. coli*

No. of NCs

1.5

B *C. jejuni*

No. of NCs

3.0

Genome position (bp)

Fig 5.