**Leukodystrophy resembling Vanishing White Matter Disease is**

**recapitulated by brain-specific depletion of apoptosis regulator MCL-1**

**Running title**: MCL-1 is required for white matter integrity

Abigail H. Cleveland[1,2], Alejandra Romero[3], Laurent Alfonso Azcona[2], Melisa Herrero[4],

Sheryl Moy[5,6], Orna Elroy-Stein[4,7], Vivian Gama[3,8,9] and Timothy R. Gershon[2,10,11]

[1]UNC Cancer Cell Biology Training Program, University of North Carolina, Chapel Hill,

North Carolina 27599, USA

[2]Department of Neurology, UNC School of Medicine, University of North Carolina,

Chapel Hill, North Carolina 27599, USA

[3]Vanderbilt University, Cell and Developmental Biology, Nashville, Tennessee 37232,

USA .

[4]Shmunis School of Biomedicine and Cancer Research, Faculty of Life Sciences, Tel

Aviv University, Tel Aviv, Israel

[5]UNC Carolina Institute for Developmental Disabilities

[6]Department of Psychiatry, UNC School of Medicine, University of North Carolina,

Chapel Hill, North Carolina 27599, USA

[7]Sagol School of Neuroscience, Tel Aviv University, Tel Aviv, Israel

[8]Vanderbilt University, Vanderbilt Center for Stem Cell Biology, Nashville, Tennessee

37232, USA .

[9]Vanderbilt University, Vanderbilt Brain Institute, Nashville, Tennessee 37232, USA .

[10]UNC Neuroscience Center, University of North Carolina, Chapel Hill, North Carolina

27599, USA

[11]Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill,

North Carolina 27599, USA

**Co-corresponding authors:** Vivian Gama (vivian.gama@vanderbilt.edu) and

Timothy R. Gershon (gershont@neurology.med.unc.edu)

33
34

**SUMMARY**

Diverse mutations cause leukodystrophies through unresolved processes. Developing leukodystrophy therapies requires identifying mechanisms that operate downstream of causative mutations to produce white matter degeneration. We have identified the apoptosis regulator MCL-1 as required for white matter stability and depleted by leukodystrophy-causing mutations. Brain-specific deletion of *Mcl-1* in mice recapitulates progressive leukodystrophy, with MRI changes, decreased oligodendrocytes, and astrocytic and microglial changes typical of Vanishing White Matter Disease (VWMD). Disabling apoptosis through co-deletion of *Bax* or *Bak* rescued white matter *Mcl-1*-dependence, broadly implicating the intrinsic apoptotic pathway in leukodystrophy pathogenesis. *Bax* and *Bak* co-deletions rescued different aspects of the *Mcl-1*-deleted phenotype, demonstrating their discrete roles in white matter stability. MCL-1 protein abundance was reduced in *eif2b5*-mutant mice that model VWMD, suggesting a mechanistic relationship between MCL-1 and VWMD. Our data show that oligodendrocytes require MCL-1 to suppress apoptosis, implicate MCL-1 loss in VWMD pathogenesis, and suggest inhibiting apoptosis for leukodystrophy therapy.

**Introduction**

Leukodystrophies are a heterogeneous group of severe neurodegenerative disorders marked by progressive white matter disruption (Raymond, 2017, van der Knaap and Bugiani, 2017). The age of onset varies in each leukodystrophy, but patients typically become symptomatic during childhood, after a latent, asymptomatic period (Kohler et al., 2018, Kohlschutter and Eichler, 2011). Patients with leukodystrophies may experience progressive symptoms including ataxia, dementia,

hallucinations, seizures, coma, and death (Vanderver et al., 1993). There is currently

no specific targeted therapy that has been clinically implemented to arrest the

progression of any leukodystrophy.

Diverse pathogenic mutations have been linked to different leukodystrophies

(van der Knaap and Bugiani, 2018), but the mechanistic processes through which

specific mutations cause white matter degeneration are incompletely understood.

Mutations in the apoptotic regulator *APOPT1* (aka *COA8*) cause Progressive

Cavitating Leukoencephalopathy, suggesting a role for inappropriate apoptosis in this

disease (Melchionda et al., 2014, Yasuda et al., 2006). However, it is not clear why

this mutation specifically affects cells of the white matter. Mutations in *eIF2B* subunits

result in Vanishing White Matter Disease (VWMD), another leukodystrophy similarly

marked by white matter cavitation (Bugiani et al., 2018, van der Knaap et al., 1997,

van der Knaap et al., 1993). The causative role of *eIF2B* mutations implicates

impaired mRNA translational regulation in VWMD pathogenesis, associated with

activation of the Integrated Stress Response (ISR) (van der Knaap et al., 2006, Fogli

et al., 2002, Leegwater et al., 2001, Bugiani et al., 2010, Abbink et al., 2019, Kantor

et al., 2005). Prior studies have not identified a role for apoptosis in VWMD causation,

as seen in Progressive Cavitating Leukoencephalopathy, and whether a common

pathogenic process unites these similar phenotypes is unclear.

Studies in *eIf2b5* $^{R132H/R132H}$ mice provide insight into the disproportionate effect

of *eIF2B* mutations on white matter. These mice develop leukodystrophy resembling

VWMD and show impaired mitochondrial function in fibroblasts, astrocytes, and

oligodendrocytes (Herrero et al., 2019, Raini et al., 2017, Geva et al., 2010). These

studies suggest that *eIF2B* mutations disproportionately affect white matter because

glia are more dependent than other types cells on accurate translational regulation

and mitochondrial function (Elroy-Stein, 2017). However, several unresolved steps

are required to connect the biological processes that have been found to be disrupted

in Progressive Cavitating Leukoencephalopathy and VWMD to the observed disease

process of progressive leukodystrophy.

Identifying proteins that are both required for white matter integrity and

affected by *eIF2B* mutations may show specific mechanisms that link VWMD

genotype and phenotype. Defining these mechanistic links may show why diverse

leukodystrophy mutations produce the final common pathology of white matter

degeneration. Identifying proximal mediators of white matter degeneration may also

provide important insight into leukodystrophy pathogenesis and treatment.

Here we report that mice with conditional deletion of the anti-apoptotic

regulator *Mcl-1* targeted to brain stem cells develop progressive postnatal white

matter degeneration that recapitulates the phenotype of human leukodystrophies,

particularly VWMD. MCL-1 promotes the survival of different cell lineages through its

ability to inhibit the pro-apoptotic proteins BAK and BAX, the main effectors of cell

death in mammalian cells. Importantly, MCL-1 is a mitochondrial protein that is

translationally suppressed by ISR activity (Fritsch et al., 2007), while VWMD disrupts

mitochondria and activates the ISR, raising the possibility the MCL-1 deficiency could

play a role in the VWMD phenotype.

*Mcl-1*-deleted mice were born with normal brain anatomy and show normal

motor control and behavior at postnatal day 7 (P7), but then develop increasingly

severe ataxia, abnormal motor responses, and extensive white matter atrophy. This

white matter-predominant neurodegenerative process closely matches the course of

human leukodystrophies with similar postnatal onset of symptoms, MRI changes,

progressive white matter degeneration, reduced oligodendrocyte populations,

astrocytic abnormalities, and microglial accumulation. Mutations that block the

intrinsic apoptotic pathway rescued the phenotype of *Mcl-1* deletion, implicating

apoptosis in leukodystrophy pathogenesis, and suggesting a potential approach to

therapy.

**Results**

**Conditional *Mcl-1* deletion in the CNS causes progressive white matter**
**degeneration.**

We generated mice with conditional, brain-specific *Mcl-1* deletion by breeding

*Mcl-1$^{loxP/loxP}$* mice that harbor loxP sites around exon 1 of the *Mcl1* locus (Dzhagalov

et al., 2007), with *GFAP-Cre* mice that express Cre recombinase in stem cells that

give rise to the neurons and glia of the cerebrum and cerebellum, excluding the

Purkinje cells (Supp fig. 1A-B) (Zhuo et al., 2001, Kuang et al., 2012). The resulting

*hGFAP-Cre/Mcl-1$^{loxP/loxP}$* (*Mcl-1$^{cKO}$*) mice were born at the expected Mendelian ratio.

We compared *Mcl-1$^{cKO}$* mice to littermate controls, including *Mcl-1$^{loxP/loxP}$* mice

without Cre and heterozygously deleted *hGFAP-Cre/Mcl-1$^{loxP/+}$*. We found no

differences between control genotypes and therefore pooled them for analysis.

During the first week of life, *Mcl-1$^{cKO}$* mice showed no deficits and gained weight

similarly to controls (Fig. 1A). However, by P14, *Mcl-1$^{cKO}$* mice showed reduced

weight gain (Fig. 1A) and progressive neurologic impairment. While control mice

showed normal ability to be weaned at P21, *Mcl-1$^{cKO}$* mice were unable to be weaned

and all died by P23.

*Mcl-1^{cKO}* mice demonstrated progressive abnormalities of motor function. We

used developmentally adjusted tests to compare behavior of *Mcl-1^{cKO}* mice and

controls at P7, P14, and P18. To examine behavior at P7, when mice do not yet have

open eyes and do not venture from the nest, we used a test of geotaxis by placing

mice on a tilted screen and measuring latency to turn. We also measured latency to

turn over in a righting test, and we counted pivots during a 3-min open field test. We

found no genotype-specific differences at P7 (Table 1).

| Test | WT | *Mcl-1^{cKO}* |
|---|---|---|
| Negative geotaxis (latency, s) | | |
| Trial 1 | 15.4 ± 2 | 14.3 ± 3 |
| Trial 2 | 15.1 ± 2 | 16.7 ± 3 |
| Trial 3 | 15.3 ± 2 | 20.9 ± 3 |
| Righting (latency, s) | 8.2 ± 1 | 10.0 ± 3 |
| Number of pivots | 1.7 ± 0.3 | 1.3 ± 0.5 |

**Table 1. Neonatal behavior testing showed no effects of *Mcl-1^{cKO}* genotype**

**at P7.** Each test was performed on 48 replicate WT pups and 13 replicate *Mcl-*

*1^{cKO}* pups at P7. Measures included latency to turn on a tilted screen in a test for

geotaxis, and latency to turn over in a righting test. Pivots were counted by an

experimenter during a 3-min open field test. One-way ANOVA found no

statistically significant differences.

In contrast to the similar behaviors at P7, by P14 *Mcl-1^{cKO}* mice showed

decreased rearing when placed in an open field and spent more time in the center of

the open field, while traveling similar distances compared to controls (Fig. 1B; Supp

fig. 1C). P18 *Mcl-1$^{cKO}$* mice also showed a marked increase in limb clasping when

vertically suspended (Fig. 1C), a commonly used marker of disease progression in a

number of mouse models of neurodegeneration and ataxia (Guyenet et al., 2010,

Nakamura et al., 2016).

Responses to sensory input were abnormal in *Mcl-1$^{cKO}$* mice and changed over

time. At P14, mice showed increased amplitude of acoustic startle response (Fig. 1D).

By P21, when motor function was significantly impaired, the amplitude of acoustic

startle response was markedly diminished compared to controls. These data show

reduced inhibition in response to sensory stimuli at P14, and reduced motor

responses at later ages.

Neuropathology studies demonstrated progressive white matter degeneration,

that began after the first week of life. At P7, *Mcl-1$^{cKO}$* brains showed no abnormalities

and appeared similar to controls. At P14, *Mcl-1$^{cKO}$* mice showed white matter

rarefaction and ventricular dilation and these changes were more pronounced at P21

(Fig. 1E). In addition to white matter loss, we also noted migration abnormalities in

the cerebellum, with ectopic cerebellar granule cells in the molecular layer of the

cerebellum (Fig. 1E). *Mcl-1$^{cKO}$* thus showed progressive neuropathology that

corresponded temporally with the onset of neurologic symptoms, beginning after P7

and increasing over time.

**Figure 1. *Mcl-1* deletion causes progressive motor impairment and white matter degeneration.** (**A**) Reduced weight gain in *Mcl-1^cKO* mice compared to controls. (**B**) *Mcl-1^cKO* mice show normal motor function at P7, with altered motor function by P14 and (**C**) increased hind-limb clasping at P18. (**D**) *Mcl-1^cKO* mice show increased startle in response to auditory stimuli at P14, and reduced elicited movement at P21. (**E**) Brains of representative *Mcl-1^cKO* mice shown in H&E-stained sagittal sections appear similar to controls at P7. By P14, *Mcl-1^cKO* mice develop white matter degeneration with ventricular enalargment (*) and abnormal migration of cerebellar granule neurons (arrows). White matter loss increases by P21. (**F**) Deleting *Mcl-1* in the neuron-

182    specific *Emx* lineage in *EMX1-Cre/Mcl-1$^{loxP/loxP}$* mice does not recapitulate the *Mcl-*

183    *1$^{cKO}$* phenotype. *, ** and *** denote p<0.05, p<0.01 and p<0.001 respectively, relative

184    to controls.

**185    *Mcl-1* deletion must occur in glia to cause white matter degeneration.**

The *GFAP-Cre* transgenic line induces recombination in a broad lineage that

includes both glia and neurons (Zhuo et al., 2001). In order to determine whether *Mcl-*

*1* loss in neurons drives the *Mcl-1$^{cKO}$* phenotype, we deleted *Mcl-1* with the neuron-

specific *EMX1-Cre* by crossing *Mcl-1$^{loxP/loxP}$* and *EMX1-Cre* mouse lines. The

resulting *EMX1-Cre/Mcl-1$^{loxP/loxP}$* mice did not develop white matter degeneration or

ventriculomegaly (Fig. 1F). In contrast, deletion of *Mcl-1* in *Nestin-Cre/Mcl-1$^{loxP/loxP}$*

mice produced white matter degeneration as seen in *Mcl-1$^{cKO}$* mice (Supp Fig. 2).

*Nestin-Cre*, like *GFAP-Cre*, drives recombination in stem cells with neural and glial

potential. The phenotype of white matter degeneration thus required *Mcl-1* deletion

in the glial lineage.

**196    Blocking apoptosis prevents white matter degeneration in *Mcl-1*-deleted mice**

To determine if white matter degeneration in *Mcl-1$^{cKO}$* mice resulted from

increased apoptosis, we examined cell death through cleaved caspase-3

immunohistochemistry and TUNEL assays. We did not detect an increase in cleaved

caspase-3-stained cells at P7, prior to the onset of white matter loss, or at P14, after

white matter loss was evident (data not shown). However, TUNEL staining, performed

at P11 when white matter loss was developing, demonstrated increased cell death in

*Mcl-1$^{cKO}$* brains, consistent with a pro-apoptotic effect of *Mcl-1* deletion (Fig. 2A).

To determine whether the increased cell death in *Mcl-1*-deleted mice was

causally related to the white matter degeneration, we tested whether blocking

apoptosis through co-deletion of either *Bax* or *Bak* rescued the *Mcl-1$^{cKO}$* phenotype.

We repeatedly intercrossed *Mcl-1$^{cKO}$* mice with *Bax$^{loxP/LoxP}$/Bak$^{-/-}$* mice to generate

animals with conditional deletion of *Mcl-1* and heterozygous or homozygous deletion

of *Bax* or *Bak*. We then raised pups of each genotype and analyzed survival,

neurologic function, and neuropathology.

We found that co-deletion of *Bax* or *Bak* significantly suppressed the *Mcl-1$^{cKO}$*

phenotype, but resulted in different, mitigated phenotypes. *Mcl-1*-deleted mice with

heterozygous co-deletion of *Bak*, with the genotype *hGFAP-Cre/Mcl1$^{loxP/loxP}$/Bak$^{-/+}$*

(*Mcl-1$^{cKO}$/Bak$^{-/+}$*), showed normal survival without overt neurologic deficits and normal

white matter and cerebellar migration (Fig. 2B). In contrast, *Mcl-1*-deleted mice with

heterozygous co-deletion of *Bax*, with the genotype *hGFAP-Cre/Mcl1$^{loxP/loxP}$/Bax$^{loxP/+}$*

(*Mcl-1$^{cKO}$/Bax$^{fl/+}$*), showed a progressive leukoencephalopathy phenotype similar *Mcl-*

*1$^{cKO}$* mice (Fig. 2B). Immunohistochemistry demonstrated markedly lower BAK

protein in *Bak$^{+/-}$* animals (Fig. 2C), indicating that that heterozygous *Bak* deletion was

sufficient to alter BAK expression. The rescued phenotype of *Mcl-1$^{cKO}$/Bak$^{-/+}$* mice

therefore demonstrated reducing BAK protein expression was sufficient to prevent

white matter degeneration.

**Figure 2. *Mcl-1* deletion causes white matter degeneration by increasing apoptosis.** (**A**) TUNEL demonstrates increased cell death in *Mcl-1^cKO* brains. For cell counts, one sagittal section from each replicate was scanned at 20x and TUNEL+ cells were counted across the entire portion of each designated region that was contained within the section. (**B**) Representative H&E sections show that heterozygous *Bak* co-deletion and homozygous *Bax* co-deletion both rescue the white matter loss with ventricular enlargement (*) and cerebellar abnormalities caused by *Mcl-1* deletion. (**C**) Quantification of BAK immunohistochemistry in corpus callosum shows that heterozygous *Bak* deletion significantly reduces the number of BAK+ cells. Quantification of western blot of whole cerebella shows that homozygous *Bax* deletion reduces BAX protein abundance. (**D**) Representative images of the corpus callosum in P14 mice of the indicated genotypes shows mild white matter rarefaction in *Mcl-1/Bax^dKO* mice, not seen in *Mcl-1^cKO/Bak^+/-* mice. *, ** and *** denote p<0.05, p<0.01 and p<0.001 respectively, relative to controls.

*Mcl-1*-deleted mice with homozygous *Bax* co-deletion, with the genotype *hGFAP-Cre/Mcl1^loxP/loxP/Bax^loxP/loxP* (*Mcl-1/Bax^dKO*) showed normal survival, similar to *Mcl-1^cKO/Bak^-/+* mice. Compared to *Mcl-1^cKO* mice, *Mcl-1/Bax^dKO* mice showed greatly reduced neurologic deficits; however, we noted that by 200 days, *Mcl-1/Bax^dKO* showed a slight delay in movement on opening the cage that was not seen in *Mcl-*

*1$^{cKO}$/Bak$^{+/-}$* mice. The extensive white matter loss in *Mcl-1$^{cKO}$* mice was rescued in

*Mcl-1/Bax$^{dKO}$* mice (Fig. 2B). Close examination of brain sections from *Mcl-1/Bax$^{dKO}$*

mice, however, showed mild white matter loss (Fig 2D). The rescued phenotypes of

*Mcl-1$^{cKO}$/Bak$^{-/+}$* and *Mcl-1/Bax$^{dKO}$* mice show that disrupting either *Bax* or *Bak* is

sufficient to prevent the severe white matter degeneration that otherwise results from

*Mcl-1* deletion, implicating apoptosis in the pathogenesis of the *Mcl-1$^{cKO}$* phenotype.

The mild neurologic deficits and white matter loss in *Mcl-1/Bax$^{dKO}$* mice showed that

co-deletion of *Bax* and *Bak* do not have identical effects in *Mcl-1*-deleted animals.

**Figure 3. Abnormal neuroimaging and myelination in *Mcl-1*-deleted mice, with different rescue effects caused by co-deletion of *Bak* or *Bax*.** (**A**) Representative MRIs show white matter hyperintensity in *Mcl-1cKO* mice. (**B**) Total ventricular volume is abnormal in *Mcl-1cKO* mice and increases from P7-P21, while the volume of the 4th ventricle decreases significantly by P21. (**C,D**) Representative images and quantitative analysis of MBP IHC show that myelination is markedly decreased in *Mcl-1cKO* mice, rescued in *Mcl-1cKO/Bak+/-* mice, and incompletely rescued in *Mcl-1/BaxdKO*

mice. **(D)** Graphs show MBP+ area/total area within individual sections, normalized to the mean value in the control mice. *, ** and *** denote p<0.05, p<0.01 and p<0.001 respectively, relative to controls.

**Mcl-1 deletion phenocopies human leukodystrophies**

Leukoencephalopathies in patients produce characteristic MRI findings, including high water signal in white matter tracts (Vanderver et al., 1993), and the replacement of white matter by cerebrospinal fluid (van der Knaap et al., 1993). We analyzed whether *Mcl-1^cKO* mice showed similar white matter changes on MRI. Similar to VWMD patients, *Mcl-1^cKO* mice showed progressive ventriculomegaly (Fig. 3A). The increase in ventricle size was not evenly distributed, however, as the volume of the 4^th ventricle was significantly smaller in *Mcl-1^cKO* mice at P21 (Fig. 3B). Increased lateral ventricles and reduced 4^th ventricle was consistent with loss of periventricular cerebral white matter and an overall reduction in intracranial pressure, indicating hydrocephalus *ex vacuo*.

To determine whether myelination was disrupted in *Mcl-1^cKO* mice, we analyzed the distribution of Myelin Basic Protein (MBP). In control brains, the MBP+ area increased markedly from P7 to P14 (Fig 3C). In contrast, myelination was delayed in *Mcl-1^cKO* brains, which showed significantly less MBP+ area at P14 in all brain regions analyzed (Fig. 3D).

Analysis of MBP+ area in *Mcl-1*-deleted mice with *Bax* or *Bak* co-deletions showed that myelination required MCL-1 specifically to regulate BAK. We compared brains in P21 animals, when normal brains show widespread myelination. Heterozygous *Bak* co-deletion largely rescued myelination in *Mcl-1^cKO/Bak^+/-* mice. In these animals, MBP+ area was over 2-fold greater than in *Mcl-1^cKO* mice, although

we noted a small, statistically significant decrease in myelination compared to *Mcl-1*

intact controls. Homozygous *Bax* deletion, however, did not rescue myelination in

*Mcl-1/Bax^dKO* mice (Fig. 3D). The differential effects of *Bak* or *Bax* deletion on

myelination support the interpretation that BAX and BAK act in cells with different

functions, with both BAX and BAK affecting white matter maintenance, and only BAK

affecting myelination.

**_Mcl-1_ deletion reduces oligodendrocyte and Bergmann glia populations**

To determine if impaired myelination in *Mcl-1^cKO* mice resulted from

oligodendrocyte loss, we compared the numbers of cells expressing the

oligodendrocyte marker SOX10 in *Mcl-1^cKO* and control mice (Fig. 4A). SOX10 marks

the oligodendrocyte lineage from the initial fate commitment in oligodendrocyte

precursors to the maturation of myelinating oligodendrocytes (Glasgow et al., 2014,

Nishiyama et al., 2016); SOX10 cell counts thus reflect the total oligodendrocytic

population across the differentiation spectrum. At P7, oligodendrocyte populations in

*Mcl-1^cKO* mice were not significantly different from controls. By P14, *Mcl-1^cKO* mice

showed significantly fewer SOX10+ cells in diverse brain regions (Fig. 4B), indicating

a global decrease in oligodendrocytic cells. At P21, oligodendrocytic populations

remained reduced in the cerebellum, corpus callosum, and superior colliculus, but

were not statistically significantly reduced in the hippocampus and cerebral cortex,

raising the possibility of a dynamic, homeostatic response in these gray matter

regions.

**Figure 4. *Mcl-1* deletion causes loss of oligodendrocytes and Bergmann glia that is rescued with regional variation by co-deletion of *Bak* or *Bax*.** (**A,B**) Representative images and quantification of SOX10 IHC in the indicated genotypes. (**C**) Representative immunofluorescence for GFAP and SOX2 IHC identify Bergmann glia in P14 mice of indicated genotypes. (**D**) SOX2+ cells are significantly reduced in *Mcl-1cKO* mice. * and ** denote p<0.05 and p<0.01 respectively, relative to controls.

To determine if apoptosis mediates the decline in the oligodendrocyte populations in *Mcl-1*-deleted mice, we examined the effect of *Bax* or *Bak* co-deletion in *Mcl-1cKO/Bak+/-*and *Mcl-1/BaxdKO* mice. Heterozygous *Bak* co-deletion in *Mcl-*

*1<sup>cKO</sup>/Bak<sup>+/-</sup>* mice normalized oligodendrocyte population size, but homozygous *Bax*

deletion in *Mcl-1/Bax<sup>cKO</sup>* mice did not (Fig. 4B). *Bak* co-deletion thus rescued the

effects of *Mcl-1* deletion on white matter volume, myelination, and total

oligodendrocyte populations. In contrast, *Bax* co-deletion rescued only white matter

volume, but myelination and oligodendrocyte numbers remained affected in *Mcl-*

*1/Bax<sup>dKO</sup>* mice. The discordant effects of *Bak* and *Bax* co-deletions suggest that MCL-

1 discretely controls BAK-dependent apoptosis or BAX-dependent apoptosis in

different types of white matter cells.

In contrast to the discordant effects of *Bax* and *Bak* co-deletions in

oligodendrocytes, Bergmann glia were rescued by either co-deletion. We examined

Bergmann glia because of their known involvement in VWMD (Dooves et al., 2016),

and because of the migration abnormalities in *Mcl-1<sup>cKO</sup>* mice, as Bergmann glia guide

cerebellar granule neuron progenitors as they migrate to the internal granule cell layer

to become cerebellar granule neurons (Hatten, 1990, Rakic, 1971). We identified

Bergmann glia as cells with SOX2+ nuclei and GFAP+ processes (Fig. 4C). At P7,

the numbers of Bergmann glia in *Mcl-1<sup>cKO</sup>* were similar to control mice (data not

shown), but by P14, the Bergmann glia populations in *Mcl-1<sup>cKO</sup>* mice were significantly

reduced (Fig. 4D). The loss of Bergmann glia during the period of CGN migration

account for the observed cerebellar migration defects. Co-deletion of either one copy

of *Bak* in *Mcl-1<sup>cKO</sup>/Bak<sup>+/-</sup>* mice or both copies of *Bax* in *Mcl-1/Bax<sup>dKO</sup>* mice was

sufficient to rescue cerebellar migration, indicating that MCL-1 functions in Bergmann

glia to restrict an apoptotic program that requires both BAX AND BAK in order to drive

cell death.

### *Mcl-1* deletion alters the astrocytic populations

While *Mcl-1$^{cKO}$* mice contained fewer Bergmann glia*,* we noted increased numbers of GFAP+ astrocytic processes throughout the brain, that was most conspicuous in the relatively hypocellular molecular layer of the cerebellum (Fig. 5A). To determine if this increase in GFAP+ cells indicated an increase in the total number of astrocytes or an increase in the fraction of astrocytes expressing GFAP, we compared the number of cells expressing the astrocytic nuclear marker SOX9 in *Mcl-1$^{cKO}$* mice and controls (Fig. 5B). SOX9+ cells were not significantly increased in *Mcl-1$^{cKO}$* mice in any brain region at either P7 or P14, and rather were decreased in the corpus callosum at P14 (Fig. 5B,C). The increase in GFAP+ processes in *Mcl-1$^{cKO}$* mice without the increase in SOX9+ cells shows the *Mcl-1* deletion did not increase the total astrocyte population, but rather increased the fraction of astrocytes expressing GFAP, indicating a reactive response to injury (Eddleston and Mucke, 1993, Dahl and Bignami, 1974).

The intermediate filament protein NESTIN is known to be up-regulated in reactive astrocytes (Tamagno and Schiffer, 2006, Schmidt-Kastner et al., 2005, Sahin Kaya et al., 1999, Clarke et al., 1994), and prior studies report increased Nestin-expressing astrocytes in white matter from humans with VWMD (Bugiani et al., 2011). Similarly, *eIF2B5 $^{R191H/R191H}$* mice show increased astrocytes expressing both Nestin and GFAP in the corpus callosum, compared to wildtype controls (Dooves et al., 2016, Geva et al., 2010). To determine if *Mcl1*-deleted mice also showed similar astrocytic changes, we compared Nestin+/GFAP+ populations in P14 *Mcl-1$^{cKO}$* mice and controls (Fig. 5D). *Mcl-1$^{cKO}$* mice demonstrated increased fractions of

Nestin+/GFAP+ cells in the corpus callosum (Fig 5E), with a trend toward increased

fractions in other white matter regions, consistent with astrocytic changes in VWMD

patients and *eIF2B5* $^{R191H/R191H}$ mice. *Mcl-1$^{cKO}$* mice thus recapitulated both the

astrocytic and oligodendrocytic abnormalities seen in VWMD in patients and the

VWMD mouse models.

**Figure 5. Mcl-1 deletion causes reactive astrocytosis.** (**A**) Representative GFAP

IHC in sections of P14 cerebella in the indicated genotypes shows Bergmann glia

processes in the control mice that are absent in the *Mcl-1$^{cKO}$*, and numerous astrocyte

processes in the *Mcl-1$^{cKO}$* mice. (**B**) Representative SOX9 IHC in sections of P14

corpus callosum in the indicated genotypes show a significant decrease of SOX9+

cells in the *Mcl-1$^{cKO}$* mice. (**C**) Quantification of SOX9 IHC at the indicated ages and

genotypes, showing that astrocyte numbers are not increased in *Mcl-1$^{cKO}$* mice. (**D,E**)

Nestin/GFAP IHC, with representative images and quantification in the indicated

genotypes. * and ** denote $p<0.05$ and $p<0.01$ respectively, relative to controls.

**Mcl-1 deletion causes neuroinflammation**

Brains of *Mcl-1$^{cKO}$* mice contained increased cells expressing the myeloid

marker IBA1 (Fig. 6A), indicating microglial activation (Ito et al., 1998, Imai et al.,

1996). The microglial changes in *Mcl-1^cKO^* mice began later than the astrocytic and

oligodendrocytic changes, beginning at the corpus callosum at P14; by P21, *Mcl-1^cKO^*

mice showed increased IBA+ cells in diverse brain regions (Fig. 6B). The increased

IBA1+ microglia in *Mcl-1^cKO^* mice demonstrates a non-cell autonomous effect in which

*Mcl-1* deletion in the glio-neuronally restricted GFAP lineage induced inflammatory

changes in myeloid cells.

Co-deletion of either *Bak* or *Bax* prevented microgliosis in different brain

regions. Heterozygous co-deletion of *Bak* in *Mcl-1^cKO^; Bak^+/-^* mice significantly

reduced IBA1+ cells in the cerebellum, cerebral cortex, colliculus, and corpus

callosum, relative to *Mcl-1^cKO^* mice. In contrast, homozygous co-deletion of *Bax* in

*Mcl-1^/^Bax^dKO^* mice showed a significant decrease in IBA1+ cells in the hippocampus,

relative to both *Mcl-1^cKO^* and *Mcl-1^cKO^/Bak^+/-^* mice (Fig. 6B). The rescue of neuro-

inflammation by co-deletion of *Bax* or *Bak* implicates increased regional apoptosis as

the cause of inflammation, while the region-specific pattern of rescue further supports

the conclusion that white matter cells in different areas of the brain alternatively

regulate apoptosis through interactions of MCL-1 with either BAX or BAK.

**Decreased brain MCL-1 protein in a VWMD mouse model**

Based on the similarity of the *Mcl-1^cKO^* phenotype to VWMD, the role of the ISR in

VWMD, and the regulation of MCL-1 abundance by the ISR, we investigated whether

changes in MCL-1 protein might mediate the effect of mutant eIF2B on glial cells. To

begin to test this possibility, we compared MCL-1 abundance in brain lysates from

wildtype and *eIF2B5 ^R132H/R132H^* mice at 7 and 10 months old (Fig. 6C). We found that

by 10 months, *eIF2B5 ^R132H/R132H^* mice expressed reduced MCL-1 protein in the brain,

indicating that the *eIF2B5 ^R132H/R132H^* mutation results in reduced MCL-1 expression

over time. In *eIF2B5 $^{R132H/R132H}$* mice, the temporal course of reduced brain MCL-1

protein correlated with the development of VWMD symptoms. Thus, along with our

finding that MCL-1 deficiency in *Mcl-1$^{cKO}$* mice was sufficient to cause a VWMD

phenotype, we found declining MCL-1 expression in *eIF2B5 $^{R132H/R132H}$* mice that

model VWMD.

**Figure 6. Progressive neuroinflammation in *Mcl-1$^{cKO}$* mice and reduced MCL-1**

**protein in mice with VWMD.** (**A,B**) IBA1 IHC, with representative images and

quantification in the indicated genotypes, showing a progressive increase in microglia

in *Mcl-1$^{cKO}$* mice, beginning in the corpus callosum at P14 and expanding to include

diverse regions by P21. Co-deletion of *Bak* or *Bax* variably rescue microgliosis with

different effects in different regions. *, ** and *** denote p<0.05, p<0.01 and p<0.001

respectively, relative to controls. (**C**) Western blot comparing MCL-1 protein

abundance in brain lysates of 7- and 10-months old WT and *eIF2B5$^{R132H/R132H}$* mice,

demonstrating reduced MCL-1 in 10-month old mutant mice. * denotes p<0.03,

relative to controls.

**Discussion**

Our data show that conditional deletion of anti-apoptotic *Mcl-1* in neuro-glial

stem cells results in a progressive encephalopathy with post-natal white matter

degeneration. While prior studies have shown increased apoptosis in embryonic and

adult neural progenitors resulting from *Mcl-1* deletion (Fogarty et al., 2019, Hasan et

al., 2013, Malone et al., 2012), our data show that the predominant effect of *Mcl-1*

deletion in the postnatal brain is white matter degeneration. Although MCL-1

mutations have not been described in any human leukodystrophy, the mechanisms

of white matter injury in human leukodystrophies are unresolved. Our studies show

that altered MCL-1-dependent regulation of apoptosis can drive the pathogenesis of

leukodystrophy.

The *Mcl-1^cKO* phenotype presented striking similarities to human

leukodystrophies, and most specifically to VWMD. Brain function and white matter

appeared normal in *Mcl-1^cKO* mice until after P7, when mice showed motor and

sensory impairment, delayed myelination, and rarefaction of the periventricular white

matter and corpus callosum, eventually leading to white matter loss and

hydrocephalus *ex vacuo*. Similar to VWMD, mature myelinating oligodendrocytes

were markedly reduced, Bergmann glia of the cerebellum were also affected, and

astrocytes showed reactive changes that are also seen in astrocytes in VWMD,

including up-regulation of GFAP and NESTIN. The similarity of the *Mcl-1^cKO* and

VWMD phenotypes, including the overall histopathology of vanishing white matter

and the specific loss of terminally differentiated oligodendrocytes and Bergmann glia,

show that *eIF2B* mutations and *Mcl-1* deletion affect specific cells of the brain in a

similar manner.

Taken together, reduced MBP and SOX10 demonstrate that within global

oligodendrocyte depletion of *Mcl-1^cKO* mice, mature, myelinating oligodendrocytes are

particularly affected. Studies in VWMD patients and mouse models show that while

mature  oligodendrocyte  populations  are  reduced,  immature  oligodendrocyte

precursors may actually be increased (Herrero et al., 2019, Bugiani et al., 2013). A

similar increase in immature oligodendrocytes may occur in the *Mcl-1^cKO* mice and

would not be inconsistent with the observed overall reductions in SOX10+ brain cells

and brain myelination.

The similarity of the *Mcl-1^cKO* phenotype to VWMD raises the possibility that

reduced MCL-1 protein abundance specifically contributes to VWMD pathology. The

known activation of the ISR in VWMD (Abbink et al., 2019, Kantor et al., 2005) and

the suppression of MCL-1 translation by the ISR together provide a potential

molecular mechanism to link the *Mcl-1^cKO* phenotype to VWMD. Consistent with this

proposed mechanism, we found reduced MCL-1 protein in *eIF2B5* $^{R132H/R132H}$ mice,

which correlated with the development of white matter pathology. While studies in

patient-derived samples could potentially test whether MCL-1 abundance is altered

in VWMD patients, these studies would require samples that preceded fulminant

white matter loss, in order to include oligodendrocyte populations that may be lost

through apoptosis. Our data show that MCL-1 deficiency is sufficient to reproduce the

VWMD phenotype in mice. Studies of *Bax* and *Bak* deletion in *eIF2B5* $^{R132H/R132H}$ mice

would allow testing of whether apoptosis is required for VWMD.

The rescue of the *Mcl-1^cKO* phenotype by co-deletion of *Bak* or *Bax* implicate

apoptosis in the causation of white matter loss in *Mcl-1^cKO* mice. We previously found

that conditional *Mcl-1* deletion in cerebellar granule neuron progenitors does not

produce a phenotype of gray matter loss as these cells regulate BAX via the

homologous anti-apoptotic protein BCL-xL *(Crowther et al., 2013)*. The specific

dependencies of different cell types on different anti-apoptotic factors is also seen

outside the nervous system, as in the myeloid lineage where neutrophils undergo

spontaneous apoptosis when *Mcl-1* is deleted, but macrophages survive (Dzhagalov

et al., 2007). Different cell types within broad lineages may thus require different anti-

apoptotic family proteins to regulate apoptosis, and the survival of neurons and

astrocytes in *Mcl-1$^{cKO}$* mice shows that within the broad lineage derived from glio-

neuronal stem cells, MCL-1 protein is specifically required by a subset of cells at some

point along the more narrow lineage of the oligodendrocytes.

In addition to demonstrating the requirement for apoptosis in white matter

degeneration in *Mcl-1$^{cKO}$* mice, the *Bax* and *Bak* rescue studies demonstrate that

myelination defects and white matter loss can be dissociated. Co-deletion of *Bak*

rescued both myelination, oligodendrocyte populations, and white matter loss. In

contrast co-deletion of *Bax* reduced white matter loss without rescuing myelination

and oligodendrocyte populations. These differences show that different cell types that

can be differentiated by their apoptosis regulation contribute to different aspects of

the *Mcl-1$^{cKO}$* phenotype.

The possibility that BAX and BAK act in different cells with different sensitivities

does not require differential expression patterns. For example, prior studies show that

both BAX and BAK can drive apoptosis in cerebellar granule neuron progenitors, but

under different conditions. BAX-dependent apoptosis in these cells can be engaged

either by radiation-induced DNA damage (Chong et al., 2000), or by chromosomal

fragmentation in the setting of *Atr* deletion (Lang et al., 2016). In contrast, BAK-

dependent apoptosis is not engaged by radiation, as radiation induced apoptosis is

blocked by *Bax* deletion(Crowther et al., 2013, Chong et al., 2000); BAK-dependent

apoptosis requires the more intense trigger of *Atr* deletion and resulting chromosomal

fragmentation (Lang et al., 2016). These prior studies show that cells in the CNS that

express both BAX and BAK can show different mechanisms and sensitivities for

triggering BAX-dependent or BAK-dependent apoptosis. Similarly, specific subsets of

glia may regulate BAX-dependent and BAK-dependent apoptosis through different

mechanisms, with different, cell-type specific sensitivities and dependencies on MCL-

1.

The rescue myelination by co-deletion of *Bak* but not *Bax* shows that

myelinating oligodendrocytes or their precursors regulate apoptosis specifically

through MCL-1:BAK interactions. The different rescue pattern with *Bax* co-deletion,

which prevented white matter degeneration without normalizing myelination, shows

that a different type of cell regulates white matter loss and requires both BAK and

BAX to be present for *Mcl-1* deletion to produce phenotype changes. Clearly, diverse

cell types are affected in both *Mcl-1$^{cKO}$* mice and VWMD patients (Bugiani et al., 2018,

Zhou et al., 2019). Our genetic rescue studies show that different types of glial cells

depend on MCL-1 to regulate different pro-apoptotic proteins, and that loss of this

regulation in different types of cells reproduces different aspects of the overall

pathology. The loss of multiple subsets of glial cells may similarly contribute to

different aspects of the overall VWMD phenotype.

While multiple mutations can cause leukodystrophies, for many of these

genetic diseases, the mechanistic steps from mutation to white matter loss remain to

be specified in detail. Our findings that MCL-1 deletion in neuro-glial stem cells

produces white matter degeneration, and that this degeneration is rescued by co-

deletion of *Bax* or *Bak* implicates the process of apoptosis in the causation of

leukodystrophy. Moreover, our rescue experiments show that apoptosis in multiple

cell types derived from neuro-glial stem cells contribute to different aspects of white

matter pathology. Treatments that inhibit the intrinsic apoptotic pathway in the

postnatal brain may provide an avenue for reducing the impact of diverse

leukodystrophies.

**Acknowledgments:** supported by R01NS088219 (TRG), R01NS102627 (TRG), and
R01NS1062 (TRG), T32CA071341 (AHG), 4R00CA178190 (VG), R35GM128915
(VG),  and R21CA227483 (VG).
529

**Author Contributions**: Conceptualization, T.R.G. and V.G.; Methodology, A.H.C.,
T.R.G., V.G. and O.E-S.; Investigation, A.H.C, A.R., L.A.A., M.H.,; Writing – Original
Draft, A.H.C. and T.R.G.; Writing – Review & Editing, O.E-S., V.G. and T.R.G;
Funding Acquisition, V.G., and T.R.G.;; Resources, O.E-S., V.G., and T.R.G.;
Supervision, O.E-S., V.G., and T.R.G.

**<u>Methods</u>**

**Mice**

We generated *Mcl-1cKO* mice, by crossing *GFAP-Cre* mice (Jackson Labs, stock #

004600) with *Mcl-1LoxP/LoxP* mice (Dzhagalov et al., 2007), generously shared by Dr.

You-Wen He. We interbred GFAP-Cre, *Mcl-1loxP/loxP*, and *BaxLoxP/LoxP/Bak-/-* (Jackson

Labs, stock # 006329) to generate *Mcl-1;BaxdKO*, *Mcl-1;BakdKO*, and various

heterozygous controls. We generated *EMX1-Cre/Mcl-1loxP/loxP* mice by breeding

*EMX1-Cre* mice, generously provided by Dr. William Snide, into the *Mcl-1 f/f* line. All

mice were of species *Mus musculus* and crossed into the C57BL/6 background

through at least five generations. All animal studies were carried out with the approval

of the University of North Carolina Institutional Animal Care and Use Committee

under protocols (19-099).

Wild type (C57BL/6 strain) and *eIF2B5$^{R132H/R132H}$* mice (Geva et al., 2010) were

generated, bred, and housed in Tel Aviv University animal facility (Lab Products Inc.,

Seaford, DE, USA). All experimental procedures were approved by the Tel Aviv

University Animal Care Committee according to national guidelines (permit #04-17-

022). Breeding and genotyping were performed as previously described. (Geva et al.,

2010) Each generation was established by back cross of homozygous

*eIF2B5$^{R132H/R132H}$* mice with WT C57BL/6J (Harlan Labs, Jerusalem, Israel) to prevent

genetic drift.

**Histology and immunohistochemistry (IHC)**

Mouse brains were processed and immunostained as previously described (Ocasio

et al., 2019, Tech et al., 2017, Lang et al., 2016, Crowther et al., 2016). Primary

antibodies used were: BAK diluted 1:200 (Cell Signaling, #12105), MBP diluted

1:1000 (Abcam, #ab7349), SOX10 diluted 1:100 (Santa Cruz, #sc-17342), cleaved-

Caspase 3 (cC3) diluted 1:400 (Biocare Medical, #CP229C), glial fibrillary acidic

protein (GFAP) diluted 1:2000 (Dako, Z0334), SOX2 1:200 (Cell Signaling, #4900S),

SOX9 1:200 (R&D Systems, #AF3075), NESTIN 1:500 (Cell Signaling, #4760), IBA1

1:2000 (Wako Chemicals, #019-19741). Stained images were counterstained with

DAPI, digitally imaged using an Aperio Scan Scope XT (Aperio) and subjected to

automated cell counting using Tissue Studio (Definiens).

**Magnetic resonance imaging**

For MRI studies, brains from P7, P14, and P21 mice were harvested and fixed in 4%

formaldehyde in PBS for 48hrs, then transferred to PBS. Brains were then embedded

in 2% agarose in 15mL plastic vials and imaged in pairs as previously described

(Williams et al., 2015).

**Western blot**

Whole brains or cerebella were harvested and homogenized in lysis buffer (Cell

Signaling, #9803). Lysate protein concentrations were quantified using a

bicinchoninic acid (BCA) protein assay kit (Thermo scientific, #23225). Equal total

protein concentrations were resolved on SDS-polyacrylamide gels (BioRad, #

4561105, # 4568094) and transferred onto polyvinylidene difluoride membranes.

Membranes were blotted using a SNAP i.d. 2.0 Protein Detection System (Millipore).

Membranes were imaged using a chemiluminescent SuperSignal West Femto

Maximum Sensitivity Substrate (34095, Thermo Fisher Scientific) and the C-DiGit blot

scanner (LI-COR Biosciences). Signal was quantified using the Image Studio Lite

software (LI-COR). The following antibodies were used for western blot: MCL-1 (Cell

Signaling, #94296; 1:500 dilution), BAX (Cell Signaling, #14796; 1:500 dilution), β-

actin (Cell Signaling, #3700; 1:5000 dilution), Actin (Sigma #A3853; 1:240000

dilution), Anti-rabbit IgG, HRP-linked antibody (Cell Signaling, #7074; 1.5:1000

dilution), Anti-mouse IgG, HRP-linked antibody (Cell Signaling, #7076; 1.5:1000

dilution). For the eIF2B5$^{R132H/R132H}$ studies, cerebrums were harvested from 7 and 10

month old mice and protein was extracted using a lysis buffer containing 1% triton,

0.5% NaDOC, 0.1% SDS, 50 mM Tris pH 8, 100 mM NaCl, 10 mM β-

glycerophosphate, 5 mM NaF, 1 mM DTT, 1 mM vanadate, and EDTA-free complete

TM protease inhibitor cocktail (#11-836-170-001; ROCHE). Protein was quantified

using a BCA protein assay kit (#23227 Pierce), diluted in 1X sample buffer (0.05 M

HEPES pH 7.4, 2% (w/v) SDS, 10% (v/v) Glycerol, traces of Bromophenol Blue, 100

mM Dithiothreitol (DTT)), and separated on a 10% SDS-PAGE gel. Blots were imaged

using an AI600 Imager (Amersham) and quantified by ImageQuant TL (GE

Healthcare, Pittsburgh, PA, United States).

**Behavioral studies**

**Animals included.** Subjects were *Mcl-1$^{cKO}$* mice or littermate controls with intact Mcl-

1, including genotypes Mcl1$^{f/+}$ without Cre, Mcl1$^{f/f}$ without Cre, and Gfap-Cre/Mcl1$^{f/+}$.

A total of 8 litters were needed to generate all of the replicates. Day of birth was

considered postnatal day 0 (P0). Subject numbers at the beginning of the study (P7)

were 13 (21%) *Mcl-1$^{cKO}$* and 48 (79%) control. All litters completed the first 2 weeks

of testing. Three litters (n=23) were removed from the study for histology assays

during the third week of testing: one on P15, after the second open field test; one on

PD16, after the first acoustic startle test; and one on P21, after the tail suspension

test.

**Behavioral testing regimen**

**Developmentally-calibrated behavioral tests:** For motor tests at P7, mice were

removed from the home cage and maintained in a beaker warmed to 35°C. Mice were

then subjected to the neurobehavioral negative geotaxis, righting, and open field

tests, adapted from published studies (Cheh et al., 2006).

**Negative geotaxis**. A 20 cm × 20 cm screen with 0.5 cm wire mesh squares was set

at a 25° angle. Each pup was placed downward on the screen with its head facing

toward the bottom, allowed to grip the mesh, and then released. The time it took for

each pup to turn 180° (head and upper torso vertical), was recorded, with a 30 s

maximum and each pup was tested 3 times.

**Righting.** Each pup was placed onto its back on a flat Plexiglas surface. The

experimenter gently held the pup in place for 2 s and then released it. The latency to

righting (all 4 feet on the ground) was timed, with a maximum of 30 s.

**Open field activity and pivoting.** Each pup was placed into the center of a Plexiglas

chamber (PhenoTyper; 30.5 cm × 30.5 cm × 43.5 cm; Noldus Information

Technology, Wageningen, The Netherlands). A dim yellow light (30 lx) at the top of

the PhenoTyper box was illuminated throughout the 3 min trial. Measures were taken

of distance traveled and time in the center region (4 cm × 4 cm square in the center

of the arena) by an image-tracking system (EthoVision, Noldus Information

Technology). In addition, the experimenter recorded the number of vertical rearing

movements and number of pivots (circling or lateral movement, as opposed to

forward locomotion).

Each mouse was screened through the 3 assays in the order presented here. After

finishing the neonatal motor screen, each pup was weighed, marked with a non-toxic

Sharpie, and placed back into the warm beaker. The litter was returned to the home

cage after all pups had been tested.

**Open field activity at P14 and P19.** Additional open field activity tests were

performed at P14 and P19. The distance traveled, the number of vertical rearing

movements, and time spent in the center region was recorded in each 10 min test.

**Acoustic startle responses.** This test was used to assess auditory function,

reactivity to environmental stimuli, and sensorimotor gating. Mice were evaluated at

2 time points, P14 and P21. The procedure was based on the reflexive whole-body

flinch, or startle response, that follows exposure to the acoustic stimuli. Measures

were taken of startle magnitude and prepulse inhibition, which occurs when a weak

stimulus leads to a reduced startle in response to a subsequent louder noise.

Mice were placed into individual Plexiglas cylinders within larger, sound-

attenuating chambers. Each cylinder was seated upon a piezoelectric transducer,

which allowed vibrations to be quantified and displayed on a computer (San Diego

Instruments SR-Lab system). The chambers included a ceiling light, fan, and a

loudspeaker for the acoustic stimuli. Background sound levels (70 dB) and calibration

of the acoustic stimuli were confirmed with a digital sound level meter (San Diego

Instruments). Each session began with a 5-min habituation period, followed by 42

trials of 7 different types: no-stimulus (NoS) trials, trials with the acoustic startle

stimulus (AS; 40 msec, 120 dB) alone, and trials in which a prepulse stimulus (20

msec; either 74, 78, 82, 86, or 90 dB) occurred 100 ms before the onset of the startle

stimulus. Measures were taken of the startle amplitude for each trial across a 65-

msec sampling window, and an overall analysis was performed for each subject's

data for levels of prepulse inhibition at each prepulse sound level (calculated as 100

- [(response amplitude for prepulse stimulus and startle stimulus together / response

amplitude for startle stimulus alone) x 100].

**Limb clasping.** At PD18-21, a subset of mice (n=39) was assessed for limb clasping

during a tail-suspension test. Each mouse was lifted by the base of the tail, and

suspended above cage bedding for 15 sec, with ventral (stomach) side facing the

experimenter. Response to tail suspension was coded using the following scoring

system:

Forepaw movement

-2 = paws are clasped together, little movement, may be drawn up against

chest

0 = paws move quickly in periodic flailing or kicking motions

2 = vigorous movement, may resemble rapid digging

Hindlimb placement and movement

-4 = both hindlimbs are retracted inwards toward abdomen, with little

movement

-2 = hindlimbs are partially or intermittently retracted inwards toward abdomen

0 = hindlimbs are splayed outward from the abdomen.  Moderate kicking behavior

2 = vigorous rapid kicking

**Statistical analysis of behavioral tests**

All testing was conducted by experimenters blinded to mouse genotype. Statview

(SAS, Cary, NC) was used for data analyses. To reduce the number of $Mcl\text{-}1^{cKO}$

subjects required, behavioral data from males and females were combined.  One-

way or repeated measures analysis of variance (ANOVA) were used to determine

effects of genotype. Post-hoc analyses were conducted using Fisher's Protected

Least Significant Difference (PLSD) tests following a significant ANOVA F value. For

all comparisons, significance was set at $p < 0.05$.

**Supplementary Figures**

**Supplementary Figure 1. *hGFAP-Cre* conditionally deletes *Mcl-1* in regions**
**throughout the brain. A)** *hGFAP-Cre* induces expression of the fluorescent reporter
DENDRA2 in *hGFAP-cre/Pham* mice, indicating regions of GFAP expression. *hGFAP-*
*Cre* targets cells in diverse regions throughout the brain, including the cerebellum,
hippocampus, and cortex. **B)** MCL-1 protein levels are significantly reduced in
cerebella of *Mcl-1$^{cKO}$* mice. **C)** *Mcl-1$^{cKO}$* mice displayed decreased rearing at both P14
and P18.

**Supplementary Figure 2. *Nestin-Cre/Mcl-1$^{f/f}$* mice develop white matter degeneration similar to that of *Mcl-1$^{cKO}$* mice.** Similar to GFAP-Cre, Nestin-Cre targets cells of both the neural and glial lineage. Brains harvested from *Nestin-Cre/Mcl-1$^{f/f}$* mice showed ventricular dilation and white matter rarefaction by P30. Deletion of one copy of *Bak* rescued this phenotype in *Nestin-Cre/Mcl-1$^{f/f}$/Bak$^{-/+}$* mice.

711     **<u>References</u>**

712     ABBINK, T. E. M., WISSE, L. E., JAKU, E., THIECKE, M. J., VOLTOLINI-GONZALEZ, D., FRITSEN,
713             H., BOBELDIJK, S., TER BRAAK, T. J., POLDER, E., POSTMA, N. L., BUGIANI, M.,
714             STRUIJS, E. A., VERHEIJEN, M., STRAAT, N., VAN DER SLUIS, S., THOMAS, A. A. M.,
715             MOLENAAR, D. & VAN DER KNAAP, M. S. 2019. Vanishing white matter: deregulated
716             integrated stress response as therapy target. *Ann Clin Transl Neurol,* 6**,** 1407-1422.
717     BUGIANI, M., BOOR, I., POWERS, J. M., SCHEPER, G. C. & VAN DER KNAAP, M. S. 2010.
718             Leukoencephalopathy with vanishing white matter: a review. *J Neuropathol Exp*
719             *Neurol,* 69**,** 987-96.
720     BUGIANI, M., BOOR, I., VAN KOLLENBURG, B., POSTMA, N., POLDER, E., VAN BERKEL, C.,
721             VAN KESTEREN, R. E., WINDREM, M. S., HOL, E. M., SCHEPER, G. C., GOLDMAN, S. A.
722             & VAN DER KNAAP, M. S. 2011. Defective glial maturation in vanishing white matter
723             disease. *J Neuropathol Exp Neurol,* 70**,** 69-82.
724     BUGIANI, M., POSTMA, N., POLDER, E., DIELEMAN, N., SCHEFFER, P. G., SIM, F. J., VAN DER
725             KNAAP, M. S. & BOOR, I. 2013. Hyaluronan accumulation and arrested
726             oligodendrocyte progenitor maturation in vanishing white matter disease. *Brain,*
727             136**,** 209-22.
728     BUGIANI, M., VUONG, C., BREUR, M. & VAN DER KNAAP, M. S. 2018. Vanishing white
729             matter: a leukodystrophy due to astrocytic dysfunction. *Brain Pathol,* 28**,** 408-421.
730     CHEH, M. A., MILLONIG, J. H., ROSELLI, L. M., MING, X., JACOBSEN, E., KAMDAR, S. &
731             WAGNER, G. C. 2006. En2 knockout mice display neurobehavioral and neurochemical
732             alterations relevant to autism spectrum disorder. *Brain Res,* 1116**,** 166-76.
733     CHONG, M. J., MURRAY, M. R., GOSINK, E. C., RUSSELL, H. R. C., SRINIVASAN, A., KAPSETAKI,
734             M., KORSMEYER, S. J. & MCKINNON, P. J. 2000. Atm and Bax cooperate in ionizing
735             radiation-induced apoptosis in the central nervous system. *Proceedings of the*
736             *National Academy of Sciences,* 97**,** 889-894.
737     CLARKE, S. R., SHETTY, A. K., BRADLEY, J. L. & TURNER, D. A. 1994. Reactive astrocytes
738             express the embryonic intermediate neurofilament nestin. *Neuroreport,* 5**,** 1885-8.
739     CROWTHER, A. J., GAMA, V., BEVILACQUA, A., CHANG, S. X., YUAN, H., DESHMUKH, M. &
740             GERSHON, T. R. 2013. Tonic activation of Bax primes neural progenitors for rapid
741             apoptosis through a mechanism preserved in medulloblastoma. *The Journal of*
742             *neuroscience : the official journal of the Society for Neuroscience,* 33**,** 18098-108.
743     CROWTHER, A. J., OCASIO, J. K., FANG, F., MEIDINGER, J., WU, J., DEAL, A. M., CHANG, S. X.,
744             YUAN, H., SCHMID, R., DAVIS, I. & GERSHON, T. R. 2016. Radiation Sensitivity in a
745             Preclinical Mouse Model of Medulloblastoma Relies on the Function of the Intrinsic
746             Apoptotic Pathway. *Cancer Res,* 76**,** 3211-23.
747     DAHL, D. & BIGNAMI, A. 1974. Heterogeneity of the glial fibrillary acidic protein in gliosed
748             human brains. *J Neurol Sci,* 23**,** 551-63.
749     DOOVES, S., BUGIANI, M., POSTMA, N. L., POLDER, E., LAND, N., HORAN, S. T., VAN DEIJK, A.
750             L., VAN DE KREEKE, A., JACOBS, G., VUONG, C., KLOOSTER, J., KAMERMANS, M.,
751             WORTEL, J., LOOS, M., WISSE, L. E., SCHEPER, G. C., ABBINK, T. E., HEINE, V. M. &
752             VAN DER KNAAP, M. S. 2016. Astrocytes are central in the pathomechanisms of
753             vanishing white matter. *J Clin Invest,* 126**,** 1512-24.
754     DZHAGALOV, I., ST JOHN, A. & HE, Y. W. 2007. The antiapoptotic protein Mcl-1 is essential
755             for the survival of neutrophils but not macrophages. *Blood,* 109**,** 1620-6.

EDDLESTON, M. & MUCKE, L. 1993. Molecular profile of reactive astrocytes--implications for
their role in neurologic disease. *Neuroscience,* 54**,** 15-36.
ELROY-STEIN, O. 2017. Mitochondrial malfunction in vanishing white matter disease: a
disease of the cytosolic translation machinery. *Neural Regen Res,* 12**,** 1610-1612.
FOGARTY, L. C., FLEMMER, R. T., GEIZER, B. A., LICURSI, M., KARUNANITHY, A., OPFERMAN,
J. T., HIRASAWA, K. & VANDERLUIT, J. L. 2019. Mcl-1 and Bcl-xL are essential for
survival of the developing nervous system. *Cell Death Differ,* 26**,** 1501-1515.
FOGLI, A., WONG, K., EYMARD-PIERRE, E., WENGER, J., BOUFFARD, J. P., GOLDIN, E., BLACK,
D. N., BOESPFLUG-TANGUY, O. & SCHIFFMANN, R. 2002. Cree leukoencephalopathy
and CACH/VWM disease are allelic at the EIF2B5 locus. *Ann Neurol,* 52**,** 506-10.
FRITSCH, R. M., SCHNEIDER, G., SAUR, D., SCHEIBEL, M. & SCHMID, R. M. 2007. Translational
repression of MCL-1 couples stress-induced eIF2 alpha phosphorylation to
mitochondrial apoptosis initiation. *J Biol Chem,* 282**,** 22551-62.
GEVA, M., CABILLY, Y., ASSAF, Y., MINDROUL, N., MAROM, L., RAINI, G., PINCHASI, D. &
ELROY-STEIN, O. 2010. A mouse model for eukaryotic translation initiation factor 2B-
leucodystrophy reveals abnormal development of brain white matter. *Brain,* 133**,**
2448-61.
GLASGOW, S. M., ZHU, W., STOLT, C. C., HUANG, T. W., CHEN, F., LOTURCO, J. J., NEUL, J. L.,
WEGNER, M., MOHILA, C. & DENEEN, B. 2014. Mutual antagonism between Sox10
and NFIA regulates diversification of glial lineages and glioma subtypes. *Nat
Neurosci,* 17**,** 1322-9.
GUYENET, S. J., FURRER, S. A., DAMIAN, V. M., BAUGHAN, T. D., LA SPADA, A. R. & GARDEN,
G. A. 2010. A simple composite phenotype scoring system for evaluating mouse
models of cerebellar ataxia. *J Vis Exp*.
HASAN, S. M., SHEEN, A. D., POWER, A. M., LANGEVIN, L. M., XIONG, J., FURLONG, M., DAY,
K., SCHUURMANS, C., OPFERMAN, J. T. & VANDERLUIT, J. L. 2013. Mcl1 regulates the
terminal mitosis of neural precursor cells in the mammalian brain through p27Kip1.
*Development,* 140**,** 3118-27.
HATTEN, M. E. 1990. Riding the glial monorail: a common mechanism for glial-guided
neuronal migration in different regions of the developing mammalian brain. *Trends
Neurosci,* 13**,** 179-84.
HERRERO, M., MANDELBOUM, S. & ELROY-STEIN, O. 2019. eIF2B Mutations Cause
Mitochondrial Malfunction in Oligodendrocytes. *Neuromolecular Med,* 21**,** 303-313.
IMAI, Y., IBATA, I., ITO, D., OHSAWA, K. & KOHSAKA, S. 1996. A novel gene iba1 in the major
histocompatibility complex class III region encoding an EF hand protein expressed in
a monocytic lineage. *Biochem Biophys Res Commun,* 224**,** 855-62.
ITO, D., IMAI, Y., OHSAWA, K., NAKAJIMA, K., FUKUUCHI, Y. & KOHSAKA, S. 1998. Microglia-
specific localisation of a novel calcium binding protein, Iba1. *Brain Res Mol Brain Res,*
57**,** 1-9.
KANTOR, L., HARDING, H. P., RON, D., SCHIFFMANN, R., KANESKI, C. R., KIMBALL, S. R. &
ELROY-STEIN, O. 2005. Heightened stress response in primary fibroblasts expressing
mutant eIF2B genes from CACH/VWM leukodystrophy patients. *Hum Genet,* 118**,** 99-
106.
KOHLER, W., CURIEL, J. & VANDERVER, A. 2018. Adulthood leukodystrophies. *Nat Rev
Neurol,* 14**,** 94-105.
KOHLSCHUTTER, A. & EICHLER, F. 2011. Childhood leukodystrophies: a clinical perspective.
*Expert Rev Neurother,* 11**,** 1485-96.

KUANG, Y., LIU, Q., SHU, X., ZHANG, C., HUANG, N., LI, J., JIANG, M. & LI, H. 2012. Dicer1 and
MiR-9 are required for proper Notch1 signaling and the Bergmann glial phenotype in
the developing mouse cerebellum. *Glia,* 60**,** 1734-46.
LANG, P. Y., NANJANGUD, G. J., SOKOLSKY-PAPKOV, M., SHAW, C., HWANG, D., PARKER, J.
S., KABANOV, A. V. & GERSHON, T. R. 2016. ATR maintains chromosomal integrity
during postnatal cerebellar neurogenesis and is required for medulloblastoma
formation. *Development,* 143**,** 4038-4052.
LEEGWATER, P. A., VERMEULEN, G., KONST, A. A., NAIDU, S., MULDERS, J., VISSER, A.,
KERSBERGEN, P., MOBACH, D., FONDS, D., VAN BERKEL, C. G., LEMMERS, R. J.,
FRANTS, R. R., OUDEJANS, C. B., SCHUTGENS, R. B., PRONK, J. C. & VAN DER KNAAP,
M. S. 2001. Subunits of the translation initiation factor eIF2B are mutant in
leukoencephalopathy with vanishing white matter. *Nat Genet,* 29**,** 383-8.
MALONE, C. D., HASAN, S. M., ROOME, R. B., XIONG, J., FURLONG, M., OPFERMAN, J. T. &
VANDERLUIT, J. L. 2012. Mcl-1 regulates the survival of adult neural precursor cells.
*Mol Cell Neurosci,* 49**,** 439-47.
MELCHIONDA, L., HAACK, T. B., HARDY, S., ABBINK, T. E., FERNANDEZ-VIZARRA, E.,
LAMANTEA, E., MARCHET, S., MORANDI, L., MOGGIO, M., CARROZZO, R., TORRACO,
A., DIODATO, D., STROM, T. M., MEITINGER, T., TEKTURK, P., YAPICI, Z., AL-
MURSHEDI, F., STEVENS, R., RODENBURG, R. J., LAMPERTI, C., ARDISSONE, A.,
MORONI, I., UZIEL, G., PROKISCH, H., TAYLOR, R. W., BERTINI, E., VAN DER KNAAP,
M. S., GHEZZI, D. & ZEVIANI, M. 2014. Mutations in APOPT1, encoding a
mitochondrial protein, cause cavitating leukoencephalopathy with cytochrome c
oxidase deficiency. *Am J Hum Genet,* 95**,** 315-25.
NAKAMURA, A., SWAHARI, V., PLESTANT, C., SMITH, I., MCCOY, E., SMITH, S., MOY, S. S.,
ANTON, E. S. & DESHMUKH, M. 2016. Bcl-xL Is Essential for the Survival and Function
of Differentiated Neurons in the Cortex That Control Complex Behaviors. *J Neurosci,*
36**,** 5448-61.
NISHIYAMA, A., BOSHANS, L., GONCALVES, C. M., WEGRZYN, J. & PATEL, K. D. 2016. Lineage,
fate, and fate potential of NG2-glia. *Brain Res,* 1638**,** 116-128.
OCASIO, J. K., BATES, R. D. P., RAPP, C. D. & GERSHON, T. R. 2019. GSK-3 modulates SHH-
driven proliferation in postnatal cerebellar neurogenesis and medulloblastoma.
*Development,* 146.
RAINI, G., SHARET, R., HERRERO, M., ATZMON, A., SHENOY, A., GEIGER, T. & ELROY-STEIN,
O. 2017. Mutant eIF2B leads to impaired mitochondrial oxidative phosphorylation in
vanishing white matter disease. *J Neurochem,* 141**,** 694-707.
RAKIC, P. 1971. Neuron-glia relationship during granule cell migration in developing
cerebellar cortex. A Golgi and electronmicroscopic study in Macacus Rhesus. *J Comp*
*Neurol,* 141**,** 283-312.
RAYMOND, G. V. 2017. Leukodystrophy: Basic and Clinical. *Adv Neurobiol,* 15**,** 365-382.
SAHIN KAYA, S., MAHMOOD, A., LI, Y., YAVUZ, E. & CHOPP, M. 1999. Expression of nestin
after traumatic brain injury in rat brain. *Brain Res,* 840**,** 153-7.
SCHMIDT-KASTNER, R., AGUIRRE-CHEN, C., SAUL, I., YICK, L., HAMASAKI, D., BUSTO, R. &
GINSBERG, M. D. 2005. Astrocytes react to oligemia in the forebrain induced by
chronic bilateral common carotid artery occlusion in rats. *Brain Res,* 1052**,** 28-39.
TAMAGNO, I. & SCHIFFER, D. 2006. Nestin expression in reactive astrocytes of human
pathology. *J Neurooncol,* 80**,** 227-33.

TECH, K., TIKUNOV, A. P., FAROOQ, H., MORRISSY, A. S., MEIDINGER, J., FISH, T., GREEN, S.
C., LIU, H., LI, Y., MUNGALL, A. J., MOORE, R. A., MA, Y., JONES, S. J. M., MARRA, M.
A., VANDER HEIDEN, M. G., TAYLOR, M. D., MACDONALD, J. M. & GERSHON, T. R.
2017. Pyruvate Kinase Inhibits Proliferation during Postnatal Cerebellar
Neurogenesis and Suppresses Medulloblastoma Formation. *Cancer Res,* 77**,** 3217-
3230.
VAN DER KNAAP, M. S., BARTH, P. G., GABREELS, F. J., FRANZONI, E., BEGEER, J. H., STROINK,
H., ROTTEVEEL, J. J. & VALK, J. 1997. A new leukoencephalopathy with vanishing
white matter. *Neurology,* 48**,** 845-55.
VAN DER KNAAP, M. S. & BUGIANI, M. 2017. Leukodystrophies: a proposed classification
system based on pathological changes and pathogenetic mechanisms. *Acta*
*Neuropathol,* 134**,** 351-382.
VAN DER KNAAP, M. S. & BUGIANI, M. 2018. Leukodystrophies - much more than just
diseases of myelin. *Nat Rev Neurol,* 14**,** 747-748.
VAN DER KNAAP, M. S., FOGLI, A., BOESPFLUG-TANGUY, O., ABBINK, T. E. M. &
SCHIFFMANN, R. 1993. Childhood Ataxia with Central Nervous System
Hypomyelination / Vanishing White Matter. *In:* ADAM, M. P., ARDINGER, H. H.,
PAGON, R. A., WALLACE, S. E., BEAN, L. J. H., STEPHENS, K. & AMEMIYA, A. (eds.)
*GeneReviews((R)).* Seattle (WA).
VAN DER KNAAP, M. S., PRONK, J. C. & SCHEPER, G. C. 2006. Vanishing white matter disease.
*Lancet Neurol,* 5**,** 413-23.
VANDERVER, A., TONDUTI, D., SCHIFFMANN, R., SCHMIDT, J. & VAN DER KNAAP, M. S. 1993.
Leukodystrophy Overview - ARCHIVED CHAPTER, FOR HISTORICAL REFERENCE ONLY.
*In:* ADAM, M. P., ARDINGER, H. H., PAGON, R. A., WALLACE, S. E., BEAN, L. J. H.,
STEPHENS, K. & AMEMIYA, A. (eds.) *GeneReviews((R)).* Seattle (WA).
YASUDA, O., FUKUO, K., SUN, X., NISHITANI, M., YOTSUI, T., HIGUCHI, M., SUZUKI, T.,
RAKUGI, H., SMITHIES, O., MAEDA, N. & OGIHARA, T. 2006. Apop-1, a novel protein
inducing cyclophilin D-dependent but Bax/Bak-related channel-independent
apoptosis. *J Biol Chem,* 281**,** 23899-907.
ZHOU, L., LI, P., CHEN, N., DAI, L. F., GAO, K., LIU, Y. N., SHEN, L., WANG, J. M., JIANG, Y. W.
& WU, Y. 2019. Modeling vanishing white matter disease with patient-derived
induced pluripotent stem cells reveals astrocytic dysfunction. *CNS Neurosci Ther,* 25**,**
759-771.
ZHUO, L., THEIS, M., ALVAREZ-MAYA, I., BRENNER, M., WILLECKE, K. & MESSING, A. 2001.
hGFAP-cre transgenic mice for manipulation of glial and neuronal function in vivo.
*genesis,* 31**,** 85-94.
885

891