Leukodystrophy resembling Vanishing White Matter Disease is

recapitulated by brain-specific depletion of apoptosis regulator MCL-1

Running title: MCL-1 is required for white matter integrity

Abigail H. Cleveland[1,2], Alejandra Romero[3], Laurent Alfonso Azcona[2], Melisa Herrero[4],

Orna Elroy-Stein[4,5], Vivian Gama[3,6,7] and Timothy R. Gershon[2,8,9]

[1]UNC Cancer Cell Biology Training Program, University of North Carolina, Chapel Hill,

North Carolina 27599, USA

[2]Department of Neurology, UNC School of Medicine, University of North Carolina,

Chapel Hill, North Carolina 27599, USA

[3]Vanderbilt University, Cell and Developmental Biology, Nashville, Tennessee 37232,

USA .

[4]Shmunis School of Biomedicine and Cancer Research, Faculty of Life Sciences, Tel

Aviv University, Tel Aviv, Israel

[5]Sagol School of Neuroscience, Tel Aviv University, Tel Aviv, Israel

[6]Vanderbilt University, Vanderbilt Center for Stem Cell Biology, Nashville, Tennessee

37232, USA .

[7]Vanderbilt University, Vanderbilt Brain Institute, Nashville, Tennessee 37232, USA .

[8]UNC Neuroscience Center, University of North Carolina, Chapel Hill, North Carolina

27599, USA

[9]Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill,

North Carolina 27599, USA

Co-corresponding authors: Vivian Gama (vivian.gama@vanderbilt.edu) and

Timothy R. Gershon (gershont@neurology.med.unc.edu)

31 **Acknowledgments:** supported by R01NS088219 (TRG), R01NS102627 (TRG), and
32 R01NS1062 (TRG), T32CA071341 (AHG), 4R00CA178190 (VG), R35GM128915
33 (VG), and R21CA227483 (VG).

SUMMARY

Diverse mutations cause leukodystrophies through unresolved processes. Developing leukodystrophy therapies requires identifying mechanisms that operate downstream of causative mutations to produce white matter degeneration. We have identified the apoptosis regulator MCL-1 as required for white matter stability and depleted by leukodystrophy-causing mutations. Brain-specific deletion of *Mcl-1* in mice recapitulates progressive leukodystrophy, with MRI changes, decreased oligodendrocytes, and astrocytic and microglial changes typical of Vanishing White Matter Disease (VWMD). Disabling apoptosis through co-deletion of *Bax* or *Bak* rescued white matter *Mcl-1*-dependence, broadly implicating the intrinsic apoptotic pathway in leukodystrophy pathogenesis. *Bax* and *Bak* co-deletions rescued different aspects of the *Mcl-1*-deleted phenotype, demonstrating their discrete roles in white matter stability. MCL-1 protein abundance was reduced in *eif2b5*-mutant mice that model VWMD, suggesting a mechanistic relationship between MCL-1 and VWMD. Our data show that oligodendrocytes require MCL-1 to suppress apoptosis, implicate MCL-1 loss in VWMD pathogenesis, and suggest inhibiting apoptosis for leukodystrophy therapy.

Introduction

Leukodystrophies are a heterogeneous group of severe neurodegenerative disorders marked by progressive white matter disruption [1,2]. The age of onset varies in each leukodystrophy, but patients typically become symptomatic during childhood, after a latent, asymptomatic period [3,4]. Patients with leukodystrophies may experience progressive symptoms including ataxia, dementia, hallucinations, seizures, coma,

and death [5]. There is currently no specific targeted therapy that has been clinically

implemented to arrest the progression of any leukodystrophy.

Diverse pathogenic mutations have been linked to different leukodystrophies

[6], but the mechanistic processes through which specific mutations cause white matter

degeneration are incompletely understood. Mutations in the apoptotic regulator

APOPT1 (aka *COA8*) cause Progressive Cavitating Leukoencephalopathy,

suggesting a role for inappropriate apoptosis in this disease [7,8]. However, it is not

clear why this mutation specifically affects cells of the white matter. Mutations in

eIF2B subunits result in Vanishing White Matter Disease (VWMD), another

leukodystrophy similarly marked by white matter cavitation [9-11]. The causative role of

eIF2B mutations implicates impaired mRNA translational regulation in VWMD

pathogenesis, associated with activation of the Integrated Stress Response (ISR) [12-

17]. Prior studies have not identified a role for apoptosis in VWMD causation, as seen

in Progressive Cavitating Leukoencephalopathy, and whether a common pathogenic

process unites these similar phenotypes is unclear.

Studies in *eIf2b5* $^{R132H/R132H}$ mice provide insight into the disproportionate effect

of *eIF2B* mutations on white matter. These mice develop leukodystrophy resembling

VWMD and show impaired mitochondrial function in fibroblasts, astrocytes, and

oligodendrocytes [18-20]. These studies suggest that *eIF2B* mutations disproportionately

affect white matter because glia are more dependent than other types cells on

accurate translational regulation and mitochondrial function [21]. However, several

unresolved steps are required to connect the biological processes that have been

found to be disrupted in Progressive Cavitating Leukoencephalopathy and VWMD to

the observed disease process of progressive leukodystrophy.

Identifying proteins that are both required for white matter integrity and

affected by *eIF2B* mutations may show specific mechanisms that link VWMD

genotype and phenotype. Defining these mechanistic links may show why diverse

leukodystrophy mutations produce the final common pathology of white matter

degeneration. Identifying proximal mediators of white matter degeneration may also

provide important insight into leukodystrophy pathogenesis and treatment.

Here we report that mice with conditional deletion of the anti-apoptotic

regulator *Mcl-1* targeted to brain stem cells develop progressive postnatal white

matter degeneration that recapitulates the phenotype of human leukodystrophies,

particularly VWMD. MCL-1 promotes the survival of different cell lineages through its

ability to inhibit the pro-apoptotic proteins BAK and BAX, the main effectors of cell

death in mammalian cells. Importantly, MCL-1 is a mitochondrial protein that is

translationally suppressed by ISR activity [22], while VWMD disrupts mitochondria and

activates the ISR, raising the possibility the MCL-1 deficiency could play a role in the

VWMD phenotype.

Mcl-1-deleted mice were born with normal brain anatomy and show normal

motor control and behavior at postnatal day 7 (P7), but then develop increasingly

severe ataxia, abnormal motor responses, and extensive white matter atrophy. This

white matter-predominant neurodegenerative process closely matches the course of

human leukodystrophies with similar postnatal onset of symptoms, MRI changes,

progressive white matter degeneration, reduced oligodendrocyte populations,

astrocytic abnormalities, and microglial accumulation. Mutations that block the

intrinsic apoptotic pathway rescued the phenotype of *Mcl-1* deletion, implicating

apoptosis in leukodystrophy pathogenesis, and suggesting a potential approach to

therapy.

Results

Conditional *Mcl-1* deletion in the CNS causes progressive white matter degeneration.

We generated mice with conditional, brain-specific *Mcl-1* deletion by breeding *Mcl-1^loxP/loxP^* mice that harbor loxP sites around exon 1 of the *Mcl1* locus [23], with *GFAP-Cre* mice that express Cre recombinase in stem cells that give rise to the neurons and glia of the cerebrum and cerebellum, excluding the Purkinje cells (Supp fig. 1A-B) [24,25]. The resulting *hGFAP-Cre/Mcl-1^loxP/loxP^* (*Mcl-1^cKO^*) mice were born at the expected Mendelian ratio.

We compared *Mcl-1^cKO^* mice to littermate controls, including *Mcl-1^loxP/loxP^* mice without Cre and heterozygously deleted *hGFAP-Cre/Mcl-1^loxP/+^*. We found no differences between control genotypes and therefore pooled them for analysis. During the first week of life, *Mcl-1^cKO^* mice showed no deficits and gained weight similarly to controls (Fig. 1A). However, by P14, *Mcl-1^cKO^* mice showed reduced weight gain (Fig. 1A) and progressive neurologic impairment. While control mice showed normal ability to be weaned at P21, *Mcl-1^cKO^* mice were unable to be weaned and all died by P23.

*Mcl-1^cKO^* mice demonstrated progressive abnormalities of motor function. We used developmentally adjusted tests to compare behavior of *Mcl-1^cKO^* mice and controls at P7, P14, and P18. To examine behavior at P7, when mice do not yet have open eyes and do not venture from the nest, we used a test of geotaxis by placing

132   mice on a tilted screen and measuring latency to turn. We also measured latency to

133   turn over in a righting test, and we counted pivots during a 3-min open field test. We

134   found no genotype-specific differences at P7 (Table 1).

Test	WT	Mcl-1cKO
Negative geotaxis (latency, s)		
Trial 1	15.4 ± 2	14.3 ± 3
Trial 2	15.1 ± 2	16.7 ± 3
Trial 3	15.3 ± 2	20.9 ± 3
Righting (latency, s)	8.2 ± 1	10.0 ± 3
Number of pivots	1.7 ± 0.3	1.3 ± 0.5

Table 1. Neonatal behavior testing showed no effects of *Mcl-1cKO* genotype

at P7. Each test was performed on 48 replicate WT pups and 13 replicate *Mcl-*

1cKO pups at P7. Measures included latency to turn on a tilted screen in a test for

geotaxis, and latency to turn over in a righting test. Pivots were counted by an

experimenter during a 3-min open field test. One-way ANOVA found no

statistically significant differences.

In contrast to the similar behaviors at P7, by P14 *Mcl-1cKO* mice showed

decreased rearing when placed in an open field and spent more time in the center of

the open field, while traveling similar distances compared to controls (Fig. 1B; Supp

fig. 1C). P18 *Mcl-1cKO* mice also showed a marked increase in limb clasping when

vertically suspended (Fig. 1C), a commonly used marker of disease progression in a

number of mouse models of neurodegeneration and ataxia [26,27].

Responses to sensory input were abnormal in *Mcl-1^(cKO)* mice and changed over

time. At P14, mice showed increased amplitude of acoustic startle response (Fig. 1D).

By P21, when motor function was significantly impaired, the amplitude of acoustic

startle response was markedly diminished compared to controls. These data show

reduced inhibition in response to sensory stimuli at P14, and reduced motor

responses at later ages.

Neuropathology studies demonstrated progressive white matter degeneration,

that began after the first week of life. At P7, *Mcl-1^(cKO)* brains showed no abnormalities

and appeared similar to controls. At P14, *Mcl-1^(cKO)* mice showed white matter

rarefaction and ventricular dilation and these changes were more pronounced at P21

(Fig. 1E). In addition to white matter loss, we also noted migration abnormalities in

the cerebellum, with ectopic cerebellar granule cells in the molecular layer of the

cerebellum (Fig. 1E). *Mcl-1^(cKO)* thus showed progressive neuropathology that

corresponded temporally with the onset of neurologic symptoms, beginning after P7

and increasing over time.

Figure 1. *Mcl-1* **deletion causes progressive motor impairment and white matter degeneration.** (**A**) Reduced weight gain in *Mcl-1^{cKO}* mice compared to controls. (**B**) *Mcl-1^{cKO}* mice show normal motor function at P7, with altered motor function by P14 and (**C**) increased hind-limb clasping at P18. (**D**) *Mcl-1^{cKO}* mice show increased startle in response to auditory stimuli at P14, and reduced elicited movement at P21. (**E**) Brains of representative *Mcl-1^{cKO}* mice shown in H&E-stained sagittal sections appear similar to controls at P7. By P14, *Mcl-1^{cKO}* mice develop white matter degeneration with ventricular enalargment (*) and abnormal migration of cerebellar granule neurons (arrows). White matter loss increases by P21. (**F**) Deleting *Mcl-1* in the neuron-

176   specific *Emx* lineage in *EMX1-Cre/Mcl-1^loxP/loxP* mice does not recapitulate the *Mcl-*

177   *1^cKO* phenotype. *, ** and *** denote $p<0.05$, $p<0.01$ and $p<0.001$ respectively, relative

178   to controls.

Mcl-1 deletion must occur in glia to cause white matter degeneration.

The *GFAP-Cre* transgenic line induces recombination in a broad lineage that

includes both glia and neurons [24]. In order to determine whether *Mcl-1* loss in neurons

drives the *Mcl-1^cKO* phenotype, we deleted *Mcl-1* with the neuron-specific *EMX1-Cre*

by crossing *Mcl-1^loxP/loxP* and *EMX1-Cre* mouse lines. The resulting *EMX1-Cre/Mcl-*

1^loxP/loxP mice did not develop white matter degeneration or ventriculomegaly (Fig. 1F).

In contrast, deletion of *Mcl-1* in *Nestin-Cre/Mcl-1^loxP/loxP* mice produced white matter

degeneration as seen in *Mcl-1^cKO* mice (Supp Fig. 2). *Nestin-Cre*, like *GFAP-Cre*,

drives recombination in stem cells with neural and glial potential. The phenotype of

white matter degeneration thus required *Mcl-1* deletion in the glial lineage.

Blocking apoptosis prevents white matter degeneration in *Mcl-1*-deleted mice

To determine if white matter degeneration in *Mcl-1^cKO* mice resulted from

increased apoptosis, we examined cell death through cleaved caspase-3

immunohistochemistry and TUNEL assays. We did not detect an increase in cleaved

caspase-3-stained cells at P7, prior to the onset of white matter loss, or at P14, after

white matter loss was evident (data not shown). However, TUNEL staining, performed

at P11 when white matter loss was developing, demonstrated increased cell death in

Mcl-1^cKO brains, consistent with a pro-apoptotic effect of *Mcl-1* deletion (Fig. 2A).

To determine whether the increased cell death in *Mcl-1*-deleted mice was

causally related to the white matter degeneration, we tested whether blocking

apoptosis through co-deletion of either *Bax* or *Bak* rescued the *Mcl-1^cKO* phenotype.

We repeatedly intercrossed *Mcl-1^cKO* mice with *Bax^loxP/LoxP/Bak^-/-* mice to generate

animals with conditional deletion of *Mcl-1* and heterozygous or homozygous deletion

of *Bax* or *Bak*. We then raised pups of each genotype and analyzed survival,

neurologic function, and neuropathology.

We found that co-deletion of *Bax* or *Bak* significantly suppressed the *Mcl-1^cKO*

phenotype, but resulted in different, mitigated phenotypes. *Mcl-1*-deleted mice with

heterozygous co-deletion of *Bak*, with the genotype *hGFAP-Cre/Mcl1^loxP/loxP/Bak^-/+*

(*Mcl-1^cKO/Bak^-/+*), showed normal survival without overt neurologic deficits and normal

white matter and cerebellar migration (Fig. 2B). In contrast, *Mcl-1*-deleted mice with

heterozygous co-deletion of *Bax*, with the genotype *hGFAP-Cre/Mcl1^loxP/loxP/Bax^loxP/+*

(*Mcl-1^cKO/Bax^fl/+*), showed a progressive leukoencephalopathy phenotype similar *Mcl-*

1^cKO mice (Fig. 2B). Immunohistochemistry demonstrated markedly lower BAK

protein in *Bak^+/-* animals (Fig. 2C), indicating that that heterozygous *Bak* deletion was

sufficient to alter BAK expression. The rescued phenotype of *Mcl-1^cKO/Bak^-/+* mice

therefore demonstrated reducing BAK protein expression was sufficient to prevent

white matter degeneration.

Figure 2. *Mcl-1* deletion causes white matter degeneration by increasing apoptosis. (**A**) TUNEL demonstrates increased cell death in *Mcl-1cKO* brains. For cell counts, one sagittal section from each replicate was scanned at 20x and TUNEL+ cells were counted across the entire portion of each designated region that was contained within the section. (**B**) Representative H&E sections show that heterozygous *Bak* co-deletion and homozygous *Bax* co-deletion both rescue the white matter loss with ventricular enlargement (*) and cerebellar abnormalities caused by *Mcl-1* deletion. (**C**) Quantification of BAK immunohistochemistry in corpus callosum shows that heterozygous *Bak* deletion significantly reduces the number of BAK+ cells. Quantification of western blot of whole cerebella shows that homozygous *Bax* deletion reduces BAX protein abundance. (**D**) Representative images of the corpus callosum in P14 mice of the indicated genotypes shows mild white matter rarefaction in *Mcl-1/BaxdKO* mice, not seen in *Mcl-1cKO/Bak+/-* mice. *, ** and *** denote p<0.05, p<0.01 and p<0.001 respectively, relative to controls.

Mcl-1-deleted mice with homozygous *Bax* co-deletion, with the genotype *hGFAP-Cre/Mcl1loxP/loxP/BaxloxP/loxP* (*Mcl-1/BaxdKO*) showed normal survival, similar to *Mcl-1cKO/Bak-/+* mice. Compared to *Mcl-1cKO* mice, *Mcl-1/BaxdKO* mice showed greatly reduced neurologic deficits; however, we noted that by 200 days, *Mcl-1/BaxdKO* showed a slight delay in movement on opening the cage that was not seen in *Mcl-*

1^{cKO}/Bak$^{+/-}$ mice. The extensive white matter loss in *Mcl-1^{cKO}* mice was rescued in

Mcl-1/Bax^{dKO} mice (Fig. 2B). Close examination of brain sections from *Mcl-1/Bax^{dKO}*

mice, however, showed mild white matter loss (Fig 2D). The rescued phenotypes of

Mcl-1^{cKO}/Bak$^{-/+}$ and *Mcl-1/Bax^{dKO}* mice show that disrupting either *Bax* or *Bak* is

sufficient to prevent the severe white matter degeneration that otherwise results from

Mcl-1 deletion, implicating apoptosis in the pathogenesis of the *Mcl-1^{cKO}* phenotype.

The mild neurologic deficits and white matter loss in *Mcl-1/Bax^{dKO}* mice showed that

co-deletion of *Bax* and *Bak* do not have identical effects in *Mcl-1*-deleted animals.

Figure 3. Abnormal neuroimaging and myelination in *Mcl-1*-deleted mice, with different rescue effects caused by co-deletion of *Bak* or *Bax*. (**A**) Representative MRIs show white matter hyperintensity in *Mcl-1cKO* mice. (**B**) Total ventricular volume is abnormal in *Mcl-1cKO* mice and increases from P7-P21, while the volume of the 4th ventricle decreases significantly by P21. (**C,D**) Representative images and quantitative analysis of MBP IHC show that myelination is markedly decreased in *Mcl-1cKO* mice, rescued in *Mcl-1cKO/Bak+/-* mice, and incompletely rescued in *Mcl-1/BaxdKO*

253   mice. **(D)** Graphs show MBP+ area/total area within individual sections, normalized

254   to the mean value in the control mice. *, ** and *** denote p<0.05, p<0.01 and

255   p<0.001 respectively, relative to controls.

Mcl-1 deletion phenocopies human leukodystrophies

Leukoencephalopathies in patients produce characteristic MRI findings,

including high water signal in white matter tracts [5], and the replacement of white

matter by cerebrospinal fluid [11]. We analyzed whether *Mcl-1^{cKO}* mice showed similar

white matter changes on MRI. Similar to VWMD patients, *Mcl-1^{cKO}* mice showed

progressive ventriculomegaly (Fig. 3A). The increase in ventricle size was not evenly

distributed, however, as the volume of the 4th ventricle was significantly smaller in

Mcl-1^{cKO} mice at P21 (Fig. 3B). Increased lateral ventricles and reduced 4th ventricle

was consistent with loss of periventricular cerebral white matter and an overall

reduction in intracranial pressure, indicating hydrocephalus *ex vacuo*.

To determine whether myelination was disrupted in *Mcl-1^{cKO}* mice, we

analyzed the distribution of Myelin Basic Protein (MBP). In control brains, the MBP+

area increased markedly from P7 to P14 (Fig 3C). In contrast, myelination was

delayed in *Mcl-1^{cKO}* brains, which showed significantly less MBP+ area at P14 in all

brain regions analyzed (Fig. 3D).

Analysis of MBP+ area in *Mcl-1*-deleted mice with *Bax* or *Bak* co-deletions

showed that myelination required MCL-1 specifically to regulate BAK. We compared

brains in P21 animals, when normal brains show widespread myelination.

Heterozygous *Bak* co-deletion largely rescued myelination in *Mcl-1^{cKO}/Bak$^{+/-}$* mice. In

these animals, MBP+ area was over 2-fold greater than in *Mcl-1^{cKO}* mice, although

we noted a small, statistically significant decrease in myelination compared to *Mcl-1*

intact controls. Homozygous *Bax* deletion, however, did not rescue myelination in

Mcl-1/Bax^dKO mice (Fig. 3D). The differential effects of *Bak* or *Bax* deletion on

myelination support the interpretation that BAX and BAK act in cells with different

functions, with both BAX and BAK affecting white matter maintenance, and only BAK

affecting myelination.

Mcl-1 deletion reduces oligodendrocyte and Bergmann glia populations

To determine if impaired myelination in *Mcl-1^cKO* mice resulted from

oligodendrocyte loss, we compared the numbers of cells expressing the

oligodendrocyte marker SOX10 in *Mcl-1^cKO* and control mice (Fig. 4A). SOX10 marks

the oligodendrocyte lineage from the initial fate commitment in oligodendrocyte

precursors to the maturation of myelinating oligodendrocytes [31,32]; SOX10 cell counts

thus reflect the total oligodendrocytic population across the differentiation spectrum.

At P7, oligodendrocyte populations in *Mcl-1^cKO* mice were not significantly different

from controls. By P14, *Mcl-1^cKO* mice showed significantly fewer SOX10+ cells in

diverse brain regions (Fig. 4B), indicating a global decrease in oligodendrocytic cells.

At P21, oligodendrocytic populations remained reduced in the cerebellum, corpus

callosum, and superior colliculus, but were not statistically significantly reduced in the

hippocampus and cerebral cortex, raising the possibility of a dynamic, homeostatic

response in these gray matter regions.

Figure 4. *Mcl-1* deletion causes loss of oligodendrocytes and Bergmann glia that is rescued with regional variation by co-deletion of *Bak* or *Bax*. (**A,B**) Representative images and quantification of SOX10 IHC in the indicated genotypes. (**C**) Representative immunofluorescence for GFAP and SOX2 IHC identify Bergmann glia in P14 mice of indicated genotypes. (**D**) SOX2+ cells are significantly reduced in *Mcl-1cKO* mice. * and ** denote p<0.05 and p<0.01 respectively, relative to controls.

To determine if apoptosis mediates the decline in the oligodendrocyte populations in *Mcl-1*-deleted mice, we examined the effect of *Bax* or *Bak* co-deletion in *Mcl-1cKO/Bak+/-*and *Mcl-1/BaxdKO* mice. Heterozygous *Bak* co-deletion in *Mcl-*

1cKO/Bak+/- mice normalized oligodendrocyte population size, but homozygous *Bax*

deletion in *Mcl-1/BaxcKO* mice did not (Fig. 4B). *Bak* co-deletion thus rescued the

effects of *Mcl-1* deletion on white matter volume, myelination, and total

oligodendrocyte populations. In contrast, *Bax* co-deletion rescued only white matter

volume, but myelination and oligodendrocyte numbers remained affected in *Mcl-*

1/BaxdKO mice. The discordant effects of *Bak* and *Bax* co-deletions suggest that MCL-

1 discretely controls BAK-dependent apoptosis or BAX-dependent apoptosis in

different types of white matter cells.

In contrast to the discordant effects of *Bax* and *Bak* co-deletions in

oligodendrocytes, Bergmann glia were rescued by either co-deletion. We examined

Bergmann glia because of their known involvement in VWMD [33], and because of the

migration abnormalities in *Mcl-1cKO* mice, as Bergmann glia guide cerebellar granule

neuron progenitors as they migrate to the internal granule cell layer to become

cerebellar granule neurons [34,35]. We identified Bergmann glia as cells with SOX2+

nuclei and GFAP+ processes (Fig. 4C). At P7, the numbers of Bergmann glia in *Mcl-*

1cKO were similar to control mice (data not shown), but by P14, the Bergmann glia

populations in *Mcl-1cKO* mice were significantly reduced (Fig. 4D). The loss of

Bergmann glia during the period of CGN migration account for the observed

cerebellar migration defects. Co-deletion of either one copy of *Bak* in *Mcl-1cKO/Bak+/-*

mice or both copies of *Bax* in *Mcl-1/BaxdKO* mice was sufficient to rescue cerebellar

migration, indicating that MCL-1 functions in Bergmann glia to restrict an apoptotic

program that requires both BAX AND BAK in order to drive cell death.

Mcl-1 deletion alters the astrocytic populations

While _Mcl-1cKO_ mice contained fewer Bergmann glia, we noted increased

numbers of GFAP+ astrocytic processes throughout the brain, that was most

conspicuous in the relatively hypocellular molecular layer of the cerebellum (Fig. 5A).

To determine if this increase in GFAP+ cells indicated an increase in the total number

of astrocytes or an increase in the fraction of astrocytes expressing GFAP, we

compared the number of cells expressing the astrocytic nuclear marker SOX9 in _Mcl-_

1cKO mice and controls (Fig. 5B). SOX9+ cells were not significantly increased in _Mcl-_

1cKO mice in any brain region at either P7 or P14, and rather were decreased in the

corpus callosum at P14 (Fig. 5B,C). The increase in GFAP+ processes in _Mcl-1cKO_

mice without the increase in SOX9+ cells shows the _Mcl-1_ deletion did not increase

the total astrocyte population, but rather increased the fraction of astrocytes

expressing GFAP, indicating a reactive response to injury [36,37].

The intermediate filament protein NESTIN is known to be up-regulated in

reactive astrocytes [38-41], and prior studies report increased Nestin-expressing

astrocytes in white matter from humans with VWMD [42]. Similarly, _eIF2B5_ [R191H/R191H]

mice show increased astrocytes expressing both Nestin and GFAP in the corpus

callosum, compared to wildtype controls [20,33]. To determine if _Mcl1_-deleted mice also

showed similar astrocytic changes, we compared Nestin+/GFAP+ populations in P14

Mcl-1cKO mice and controls (Fig. 5D). _Mcl-1cKO_ mice demonstrated increased fractions

of Nestin+/GFAP+ cells in the corpus callosum (Fig 5E), with a trend toward increased

fractions in other white matter regions, consistent with astrocytic changes in VWMD

patients and _eIF2B5_ [R191H/R191H] mice. _Mcl-1cKO_ mice thus recapitulated both the

astrocytic and oligodendrocytic abnormalities seen in VWMD in patients and the

VWMD mouse models.

Figure 5. Mcl-1 deletion causes reactive astrocytosis. (**A**) Representative GFAP
IHC in sections of P14 cerebella in the indicated genotypes shows Bergmann glia
processes in the control mice that are absent in the *Mcl-1^cKO^*, and numerous astrocyte
processes in the *Mcl-1^cKO^* mice. (**B**) Representative SOX9 IHC in sections of P14
corpus callosum in the indicated genotypes show a significant decrease of SOX9+
cells in the *Mcl-1^cKO^* mice. (**C**) Quantification of SOX9 IHC at the indicated ages and
genotypes, showing that astrocyte numbers are not increased in *Mcl-1^cKO^* mice. (**D,E**)
Nestin/GFAP IHC, with representative images and quantification in the indicated
genotypes. * and ** denote $p<0.05$ and $p<0.01$ respectively, relative to controls.

Mcl-1 deletion causes neuroinflammation

367   Brains of *Mcl-1^cKO^* mice contained increased cells expressing the myeloid
marker IBA1 (Fig. 6A), indicating microglial activation [43,44]. The microglial changes in
*Mcl-1^cKO^* mice began later than the astrocytic and oligodendrocytic changes,
beginning at the corpus callosum at P14; by P21, *Mcl-1^cKO^* mice showed increased
IBA+ cells in diverse brain regions (Fig. 6B). The increased IBA1+ microglia in *Mcl-*

1^{cKO} mice demonstrates a non-cell autonomous effect in which *Mcl-1* deletion in the glio-neuronally restricted GFAP lineage induced inflammatory changes in myeloid cells.

Co-deletion of either *Bak* or *Bax* prevented microgliosis in different brain regions. Heterozygous co-deletion of *Bak* in *Mcl-1^{cKO}; Bak$^{+/-}$* mice significantly reduced IBA1+ cells in the cerebellum, cerebral cortex, colliculus, and corpus callosum, relative to *Mcl-1^{cKO}* mice. In contrast, homozygous co-deletion of *Bax* in *Mcl-1$^{/}$Bax^{dKO}* mice showed a significant decrease in IBA1+ cells in the hippocampus, relative to both *Mcl-1^{cKO}* and *Mcl-1^{cKO}/Bak$^{+/-}$* mice (Fig. 6B). The rescue of neuro-inflammation by co-deletion of *Bax* or *Bak* implicates increased regional apoptosis as the cause of inflammation, while the region-specific pattern of rescue further supports the conclusion that white matter cells in different areas of the brain alternatively regulate apoptosis through interactions of MCL-1 with either BAX or BAK.

Decreased brain MCL-1 protein in a VWMD mouse model

Based on the similarity of the *Mcl-1^{cKO}* phenotype to VWMD, the role of the ISR in VWMD, and the regulation of MCL-1 abundance by the ISR, we investigated whether changes in MCL-1 protein might mediate the effect of mutant eIF2B on glial cells. To begin to test this possibility, we compared MCL-1 abundance in brain lysates from wildtype and *eIF2B5 $^{R132H/R132H}$* mice at 7 and 10 months old (Fig. 6C). We found that by 10 months, *eIF2B5 $^{R132H/R132H}$* mice expressed reduced MCL-1 protein in the brain, indicating that the *eIF2B5 $^{R132H/R132H}$* mutation results in reduced MCL-1 expression over time. In *eIF2B5 $^{R132H/R132H}$* mice, the temporal course of reduced brain MCL-1 protein correlated with the development of VWMD symptoms. Thus, along with our finding that MCL-1 deficiency in *Mcl-1^{cKO}* mice was sufficient to cause a VWMD

phenotype, we found declining MCL-1 expression in *eIF2B5* $^{R132H/R132H}$ mice that

model VWMD.

Figure 6. Progressive neuroinflammation in *Mcl-1^cKO* mice and reduced MCL-1

protein in mice with VWMD. (**A,B**) IBA1 IHC, with representative images and

quantification in the indicated genotypes, showing a progressive increase in microglia

in *Mcl-1^cKO* mice, beginning in the corpus callosum at P14 and expanding to include

diverse regions by P21. Co-deletion of *Bak* or *Bax* variably rescue microgliosis with

different effects in different regions. *, ** and *** denote $p<0.05$, $p<0.01$ and $p<0.001$

respectively, relative to controls. (**C**) Western blot comparing MCL-1 protein

abundance in brain lysates of 7- and 10-months old WT and *eIF2B5*$^{R132H/R132H}$ mice,

demonstrating reduced MCL-1 in 10-month old mutant mice. * denotes $p<0.03$,

relative to controls.

Discussion

Our data show that conditional deletion of anti-apoptotic *Mcl-1* in neuro-glial stem cells results in a progressive encephalopathy with post-natal white matter degeneration. While MCL-1 mutations have not been described in any human leukodystrophy, our data show that *Mcl-1* loss is sufficient to cause white matter degeneration. The mechanisms of white matter injury in human leukodystrophies are unresolved; our studies suggest that altered regulation of apoptosis, potentially through disruption of MCL-1 expression, may play a central role in the pathogenesis of leukodystrophies.

The *Mcl-1^{cKO}* phenotype presented striking similarities to human leukodystrophies, and most specifically to VWMD. Brain function and white matter appeared normal in *Mcl-1^{cKO}* mice until after P7, when mice showed motor and sensory impairment, delayed myelination, and rarefaction of the periventricular white matter and corpus callosum, eventually leading to white matter loss and hydrocephalus *ex vacuo*. Similar to VWMD, mature myelinating oligodendrocytes were markedly reduced, Bergmann glia of the cerebellum were also affected, and astrocytes showed reactive changes that are also seen in astrocytes in VWMD, including up-regulation of GFAP and NESTIN. The similarity of the *Mcl-1^{cKO}* and VWMD phenotypes, including the overall histopathology of vanishing white matter and the specific loss of terminally differentiated oligodendrocytes and Bergmann glia, show that *eIF2B* mutations and *Mcl-1* deletion affect specific cells of the brain in a similar manner.

Taken together, reduced MBP and SOX10 demonstrate that within global oligodendrocyte depletion of *Mcl-1^{cKO}* mice, mature, myelinating oligodendrocytes are

435   particularly affected. Studies in VWMD patients and mouse models show that while

436   mature oligodendrocyte populations are reduced, immature oligodendrocyte

437   precursors may actually be increased [18,45]. A similar increase in immature

438   oligodendrocytes may occur in the *Mcl-1^cKO* mice and would not be inconsistent with

439   the observed overall reductions in SOX10+ brain cells and brain myelination.

440   The similarity of the *Mcl-1^cKO* phenotype to VWMD raises the possibility that

reduced MCL-1 protein abundance specifically contributes to VWMD pathology. The

known activation of the ISR in VWMD [16,17] and the suppression of MCL-1 translation

by the ISR together provide a potential molecular mechanism to link the *Mcl-1^cKO*

phenotype to VWMD. Consistent with this proposed mechanism, we found reduced

MCL-1 protein in *eIF2B5 ^R132H/R132H* mice, which correlated with the development of

white matter pathology. While studies in patient-derived samples could potentially test

whether MCL-1 abundance is altered in VWMD patients, these studies would require

samples that preceded fulminant white matter loss, in order to include

oligodendrocyte populations that may be lost through apoptosis. Our data show that

MCL-1 deficiency is sufficient to reproduce the VWMD phenotype in mice. Studies of

Bax and *Bak* deletion in *eIF2B5 ^R132H/R132H* mice would allow testing of whether

apoptosis is required for VWMD.

453   The rescue of the *Mcl-1^cKO* phenotype by co-deletion of *Bak* or *Bax* implicate

apoptosis in the causation of white matter loss in *Mcl-1^cKO* mice. We previously found

that conditional *Mcl-1* deletion in cerebellar granule neuron progenitors does not

produce a phenotype of gray matter loss as these cells regulate BAX via the

homologous anti-apoptotic protein BCL-xL [30]. The specific dependencies of different

cell types on different anti-apoptotic factors is also seen outside the nervous system,

as in the myeloid lineage where neutrophils undergo spontaneous apoptosis when

Mcl-1 is deleted, but macrophages survive [23]. Different cell types within broad

lineages may thus require different anti-apoptotic family proteins to regulate

apoptosis, and the survival of neurons and astrocytes in *Mcl-1^cKO* mice shows that

within the broad lineage derived from glio-neuronal stem cells, MCL-1 protein is

specifically required by a subset of cells at some point along the more narrow lineage

of the oligodendrocytes.

In addition to demonstrating the requirement for apoptosis in white matter

degeneration in *Mcl-1^cKO* mice, the *Bax* and *Bak* rescue studies demonstrate that

myelination defects and white matter loss can be dissociated. Co-deletion of *Bak*

rescued both myelination, oligodendrocyte populations, and white matter loss. In

contrast co-deletion of *Bax* reduced white matter loss without rescuing myelination

and oligodendrocyte populations. These differences show that different cell types that

can be differentiated by their apoptosis regulation contribute to different aspects of

the *Mcl-1^cKO* phenotype.

The possibility that BAX and BAK act in different cells with different sensitivities

does not require differential expression patterns. For example, prior studies show that

both BAX and BAK can drive apoptosis in cerebellar granule neuron progenitors, but

under different conditions. BAX-dependent apoptosis in these cells can be engaged

either by radiation-induced DNA damage [28], or by chromosomal fragmentation in the

setting of *Atr* deletion [29]. In contrast, BAK-dependent apoptosis is not engaged by

radiation, as radiation induced apoptosis is blocked by *Bax* deletion[28,30]; BAK-

dependent apoptosis requires the more intense trigger of *Atr* deletion and resulting

chromosomal fragmentation [29]. These prior studies show that cells in the CNS that

express both BAX and BAK can show different mechanisms and sensitivities for

triggering BAX-dependent or BAK-dependent apoptosis. Similarly, specific subsets of

glia may regulate BAX-dependent and BAK-dependent apoptosis through different

mechanisms, with different, cell-type specific sensitivities and dependencies on MCL-

1.

The rescue myelination by co-deletion of *Bak* but not *Bax* shows that

myelinating oligodendrocytes or their precursors regulate apoptosis specifically

through MCL-1:BAK interactions. The different rescue pattern with *Bax* co-deletion,

which prevented white matter degeneration without normalizing myelination, shows

that a different type of cell regulates white matter loss and requires both BAK and

BAX to be present for *Mcl-1* deletion to produce phenotype changes. Clearly, diverse

cell types are affected in both *Mcl-1^cKO* mice and VWMD patients [9,46]. Our genetic

rescue studies show that different types of glial cells depend on MCL-1 to regulate

different pro-apoptotic proteins, and that loss of this regulation in different types of

cells reproduces different aspects of the overall pathology. The loss of multiple

subsets of glial cells may similarly contribute to different aspects of the overall VWMD

phenotype.

While multiple mutations can cause leukodystrophies, for many of these

genetic diseases, the mechanistic steps from mutation to white matter loss remain to

be specified in detail. Our findings that MCL-1 deletion in neuro-glial stem cells

produces white matter degeneration, and that this degeneration is rescued by co-

deletion of *Bax* or *Bak* implicates the process of apoptosis in the causation of

leukodystrophy. Moreover, our rescue experiments show that apoptosis in multiple

cell types derived from neuro-glial stem cells contribute to different aspects of white

matter pathology. Treatments that inhibit the intrinsic apoptotic pathway in the
postnatal brain may provide an avenue for reducing the impact of diverse
leukodystrophies.

Methods

Mice

We generated *Mcl-1^cKO* mice, by crossing *GFAP-Cre* mice (Jackson Labs, stock #
004600) with *Mcl-1^LoxP/LoxP* mice [23], generously shared by Dr. You-Wen He. We
interbred GFAP-Cre, *Mcl-1^loxP/loxP*, and *Bax^LoxP/LoxP/Bak^-/-* (Jackson Labs, stock #
006329) to generate *Mcl-1;Bax^dKO*, *Mcl-1;Bak^dKO*, and various heterozygous controls.
We generated *EMX1-Cre/Mcl-1^loxP/loxP* mice by breeding *EMX1-Cre* mice, generously
provided by Dr. William Snide, into the *Mcl-1 f/f* line. All mice were of species *Mus*
musculus and crossed into the C57BL/6 background through at least five generations.
All animal studies were carried out with the approval of the University of North
Carolina Institutional Animal Care and Use Committee under protocols (19-099).

Wild type (C57BL/6 strain) and *eIF2B5^R132H/R132H* mice [20] were generated, bred,
and housed in Tel Aviv University animal facility (Lab Products Inc., Seaford, DE,
USA). All experimental procedures were approved by the Tel Aviv University Animal
Care Committee according to national guidelines (permit #04-17-022). Breeding and
genotyping were performed as previously described. [20] Each generation was
established by back cross of homozygous *eIF2B5^R132H/R132H* mice with WT C57BL/6J
(Harlan Labs, Jerusalem, Israel) to prevent genetic drift.

Histology and immunohistochemistry (IHC)

Mouse brains were processed and immunostained as previously described [29,47-49].

Primary antibodies used were: BAK diluted 1:200 (Cell Signaling, #12105), MBP

diluted 1:1000 (Abcam, #ab7349), SOX10 diluted 1:100 (Santa Cruz, #sc-17342),

cleaved-Caspase 3 (cC3) diluted 1:400 (Biocare Medical, #CP229C), glial fibrillary

acidic protein (GFAP) diluted 1:2000 (Dako, Z0334), SOX2 1:200 (Cell Signaling,

#4900S), SOX9 1:200 (R&D Systems, #AF3075), NESTIN 1:500 (Cell Signaling,

#4760), IBA1 1:2000 (Wako Chemicals, #019-19741). Stained images were

counterstained with DAPI, digitally imaged using an Aperio Scan Scope XT (Aperio)

and subjected to automated cell counting using Tissue Studio (Definiens).

Magnetic resonance imaging

For MRI studies, brains from P7, P14, and P21 mice were harvested and fixed in 4%

formaldehyde in PBS for 48hrs, then transferred to PBS. Brains were then embedded

in 2% agarose in 15mL plastic vials and imaged in pairs as previously described

(Williams et al., 2015).

Western blot

Whole brains or cerebella were harvested and homogenized in lysis buffer (Cell

Signaling, #9803). Lysate protein concentrations were quantified using a

bicinchoninic acid (BCA) protein assay kit (Thermo scientific, #23225). Equal total

protein concentrations were resolved on SDS-polyacrylamide gels (BioRad, #

4561105, # 4568094) and transferred onto polyvinylidene difluoride membranes.

Membranes were blotted using a SNAP i.d. 2.0 Protein Detection System (Millipore).

Membranes were imaged using a chemiluminescent SuperSignal West Femto

Maximum Sensitivity Substrate (34095, Thermo Fisher Scientific) and the C-DiGit blot

scanner (LI-COR Biosciences). Signal was quantified using the Image Studio Lite

software (LI-COR). The following antibodies were used for western blot: MCL-1 (Cell

Signaling, #94296; 1:500 dilution), BAX (Cell Signaling, #14796; 1:500 dilution), β-

actin (Cell Signaling, #3700; 1:5000 dilution), Actin (Sigma #A3853; 1:240000

dilution), Anti-rabbit IgG, HRP-linked antibody (Cell Signaling, #7074; 1.5:1000

dilution), Anti-mouse IgG, HRP-linked antibody (Cell Signaling, #7076; 1.5:1000

dilution). For the eIF2B5$^{R132H/R132H}$ studies, cerebrums were harvested from 7 and 10

month old mice and protein was extracted using a lysis buffer containing 1% triton,

0.5% NaDOC, 0.1% SDS, 50 mM Tris pH 8, 100 mM NaCl, 10 mM β-

glycerophosphate, 5 mM NaF, 1 mM DTT, 1 mM vanadate, and EDTA-free complete

TM protease inhibitor cocktail (#11-836-170-001; ROCHE). Protein was quantified

using a BCA protein assay kit (#23227 Pierce), diluted in 1X sample buffer (0.05 M

HEPES pH 7.4, 2% (w/v) SDS, 10% (v/v) Glycerol, traces of Bromophenol Blue, 100

mM Dithiothreitol (DTT)), and separated on a 10% SDS-PAGE gel. Blots were imaged

using an AI600 Imager (Amersham) and quantified by ImageQuant TL (GE

Healthcare, Pittsburgh, PA, United States).

Behavioral studies

Animals included. Subjects were *Mcl-1^{cKO}* mice or littermate controls with intact Mcl-

1, including genotypes Mcl1$^{f/+}$ without Cre, Mcl1$^{f/f}$ without Cre, and Gfap-Cre/Mcl1$^{f/+}$.

A total of 8 litters were needed to generate all of the replicates. Day of birth was

considered postnatal day 0 (P0). Subject numbers at the beginning of the study (P7)

were 13 (21%) *Mcl-1^cKO* and 48 (79%) control. All litters completed the first 2 weeks

of testing. Three litters (n=23) were removed from the study for histology assays

during the third week of testing: one on P15, after the second open field test; one on

PD16, after the first acoustic startle test; and one on P21, after the tail suspension

test.

Behavioral testing regimen

Developmentally-calibrated behavioral tests: For motor tests at P7, mice were

removed from the home cage and maintained in a beaker warmed to 35°C. Mice were

then subjected to the neurobehavioral negative geotaxis, righting, and open field

tests, adapted from published studies [50].

Negative geotaxis. A 20 cm × 20 cm screen with 0.5 cm wire mesh squares was set

at a 25° angle. Each pup was placed downward on the screen with its head facing

toward the bottom, allowed to grip the mesh, and then released. The time it took for

each pup to turn 180° (head and upper torso vertical), was recorded, with a 30 s

maximum and each pup was tested 3 times.

Righting. Each pup was placed onto its back on a flat Plexiglas surface. The

experimenter gently held the pup in place for 2 s and then released it. The latency to

righting (all 4 feet on the ground) was timed, with a maximum of 30 s.

Open field activity and pivoting. Each pup was placed into the center of a Plexiglas

chamber (PhenoTyper; 30.5 cm × 30.5 cm × 43.5 cm; Noldus Information

Technology, Wageningen, The Netherlands). A dim yellow light (30 lx) at the top of

the PhenoTyper box was illuminated throughout the 3 min trial. Measures were taken

of distance traveled and time in the center region (4 cm × 4 cm square in the center of the arena) by an image-tracking system (EthoVision, Noldus Information Technology). In addition, the experimenter recorded the number of vertical rearing movements and number of pivots (circling or lateral movement, as opposed to forward locomotion).

Each mouse was screened through the 3 assays in the order presented here. After finishing the neonatal motor screen, each pup was weighed, marked with a non-toxic Sharpie, and placed back into the warm beaker. The litter was returned to the home cage after all pups had been tested.

Open field activity at P14 and P19. Additional open field activity tests were performed at P14 and P19. The distance traveled, the number of vertical rearing movements, and time spent in the center region was recorded in each 10 min test.

Acoustic startle responses. This test was used to assess auditory function, reactivity to environmental stimuli, and sensorimotor gating. Mice were evaluated at 2 time points, P14 and P21. The procedure was based on the reflexive whole-body flinch, or startle response, that follows exposure to the acoustic stimuli. Measures were taken of startle magnitude and prepulse inhibition, which occurs when a weak stimulus leads to a reduced startle in response to a subsequent louder noise.

Mice were placed into individual Plexiglas cylinders within larger, sound-attenuating chambers. Each cylinder was seated upon a piezoelectric transducer, which allowed vibrations to be quantified and displayed on a computer (San Diego Instruments SR-Lab system). The chambers included a ceiling light, fan, and a loudspeaker for the acoustic stimuli. Background sound levels (70 dB) and calibration

of the acoustic stimuli were confirmed with a digital sound level meter (San Diego

Instruments). Each session began with a 5-min habituation period, followed by 42

trials of 7 different types: no-stimulus (NoS) trials, trials with the acoustic startle

stimulus (AS; 40 msec, 120 dB) alone, and trials in which a prepulse stimulus (20

msec; either 74, 78, 82, 86, or 90 dB) occurred 100 ms before the onset of the startle

stimulus. Measures were taken of the startle amplitude for each trial across a 65-

msec sampling window, and an overall analysis was performed for each subject's

data for levels of prepulse inhibition at each prepulse sound level (calculated as 100

- [(response amplitude for prepulse stimulus and startle stimulus together / response

amplitude for startle stimulus alone) x 100].

Limb clasping. At PD18-21, a subset of mice (n=39) was assessed for limb clasping

during a tail-suspension test. Each mouse was lifted by the base of the tail, and

suspended above cage bedding for 15 sec, with ventral (stomach) side facing the

experimenter. Response to tail suspension was coded using the following scoring

system:

Forepaw movement

-2 = paws are clasped together, little movement, may be drawn up against

chest

0 = paws move quickly in periodic flailing or kicking motions

2 = vigorous movement, may resemble rapid digging

Hindlimb placement and movement

-4 = both hindlimbs are retracted inwards toward abdomen, with little

movement

657   -2 = hindlimbs are partially or intermittently retracted inwards toward abdomen

658   0 = hindlimbs are splayed outward from the abdomen. Moderate kicking behavior

659   2 = vigorous rapid kicking

Statistical analysis of behavioral tests

All testing was conducted by experimenters blinded to mouse genotype. Statview

(SAS, Cary, NC) was used for data analyses. To reduce the number of $Mcl\text{-}1^{cKO}$

subjects required, behavioral data from males and females were combined. One-

way or repeated measures analysis of variance (ANOVA) were used to determine

effects of genotype. Post-hoc analyses were conducted using Fisher's Protected

Least Significant Difference (PLSD) tests following a significant ANOVA F value. For

all comparisons, significance was set at $p < 0.05$.

Supplementary Figures

Supplementary Figure 1. ***hGFAP-Cre*** **conditionally deletes** ***Mcl-1*** **in regions**
throughout the brain. A) *hGFAP-Cre* induces expression of the fluorescent reporter
DENDRA2 in *hGFAP-cre/Pham* mice, indicating regions of GFAP expression. *hGFAP-*
Cre targets cells in diverse regions throughout the brain, including the cerebellum,
hippocampus, and cortex. **B)** MCL-1 protein levels are significantly reduced in
cerebella of *Mcl-1^{cKO}* mice. **C)** *Mcl-1^{cKO}* mice displayed decreased rearing at both P14
and P18.

**Supplementary Figure 2. *Nestin-Cre/Mcl-1^f/f* mice develop white matter
degeneration similar to that of *Mcl-1^cKO* mice.** Similar to GFAP-Cre, Nestin-Cre
targets cells of both the neural and glial lineage. Brains harvested from *Nestin-
Cre/Mcl-1^f/f* mice showed ventricular dilation and white matter rarefaction by P30.
Deletion of one copy of *Bak* rescued this phenotype in *Nestin-Cre/Mcl-1^f/f/Bak^-/+* mice.

References

1   Raymond, G. V. Leukodystrophy: Basic and Clinical. *Adv Neurobiol* **15**, 365-382,
doi:10.1007/978-3-319-57193-5_14 (2017).
2   van der Knaap, M. S. & Bugiani, M. Leukodystrophies: a proposed classification
system based on pathological changes and pathogenetic mechanisms. *Acta*
Neuropathol **134**, 351-382, doi:10.1007/s00401-017-1739-1 (2017).
3   Kohler, W., Curiel, J. & Vanderver, A. Adulthood leukodystrophies. *Nat Rev Neurol*
14, 94-105, doi:10.1038/nrneurol.2017.175 (2018).
4   Kohlschutter, A. & Eichler, F. Childhood leukodystrophies: a clinical perspective.
Expert Rev Neurother **11**, 1485-1496, doi:10.1586/ern.11.135 (2011).
5   Vanderver, A., Tonduti, D., Schiffmann, R., Schmidt, J. & van der Knaap, M. S. in
GeneReviews((R)) (eds M. P. Adam *et al.*) (1993).
6   van der Knaap, M. S. & Bugiani, M. Leukodystrophies - much more than just diseases
of myelin. *Nat Rev Neurol* **14**, 747-748, doi:10.1038/s41582-018-0093-9 (2018).
7   Melchionda, L. *et al.* Mutations in APOPT1, encoding a mitochondrial protein, cause
cavitating leukoencephalopathy with cytochrome c oxidase deficiency. *Am J Hum*
Genet **95**, 315-325, doi:10.1016/j.ajhg.2014.08.003 (2014).
8   Yasuda, O. *et al.* Apop-1, a novel protein inducing cyclophilin D-dependent but
Bax/Bak-related channel-independent apoptosis. *J Biol Chem* **281**, 23899-23907,
doi:10.1074/jbc.M512610200 (2006).
9   Bugiani, M., Vuong, C., Breur, M. & van der Knaap, M. S. Vanishing white matter: a
leukodystrophy due to astrocytic dysfunction. *Brain Pathol* **28**, 408-421,
doi:10.1111/bpa.12606 (2018).
10   van der Knaap, M. S. *et al.* A new leukoencephalopathy with vanishing white matter.
Neurology **48**, 845-855, doi:10.1212/wnl.48.4.845 (1997).
11   van der Knaap, M. S., Fogli, A., Boespflug-Tanguy, O., Abbink, T. E. M. & Schiffmann,
R. in *GeneReviews((R))* (eds M. P. Adam *et al.*) (1993).
12   van der Knaap, M. S., Pronk, J. C. & Scheper, G. C. Vanishing white matter disease.
Lancet Neurol **5**, 413-423, doi:10.1016/S1474-4422(06)70440-9 (2006).
13   Fogli, A. *et al.* Cree leukoencephalopathy and CACH/VWM disease are allelic at the
EIF2B5 locus. *Ann Neurol* **52**, 506-510, doi:10.1002/ana.10339 (2002).
14   Leegwater, P. A. *et al.* Subunits of the translation initiation factor eIF2B are mutant in
leukoencephalopathy with vanishing white matter. *Nat Genet* **29**, 383-388,
doi:10.1038/ng764 (2001).
15   Bugiani, M., Boor, I., Powers, J. M., Scheper, G. C. & van der Knaap, M. S.
Leukoencephalopathy with vanishing white matter: a review. *J Neuropathol Exp*
Neurol **69**, 987-996, doi:10.1097/NEN.0b013e3181f2eafa (2010).
16   Abbink, T. E. M. *et al.* Vanishing white matter: deregulated integrated stress
response as therapy target. *Ann Clin Transl Neurol* **6**, 1407-1422,
doi:10.1002/acn3.50826 (2019).
17   Kantor, L. *et al.* Heightened stress response in primary fibroblasts expressing mutant
eIF2B genes from CACH/VWM leukodystrophy patients. *Hum Genet* **118**, 99-106,
doi:10.1007/s00439-005-0024-x (2005).

18 Herrero, M., Mandelboum, S. & Elroy-Stein, O. eIF2B Mutations Cause Mitochondrial
Malfunction in Oligodendrocytes. *Neuromolecular Med* **21**, 303-313,
doi:10.1007/s12017-019-08551-9 (2019).
19 Raini, G. *et al.* Mutant eIF2B leads to impaired mitochondrial oxidative
phosphorylation in vanishing white matter disease. *J Neurochem* **141**, 694-707,
doi:10.1111/jnc.14024 (2017).
20 Geva, M. *et al.* A mouse model for eukaryotic translation initiation factor 2B-
leucodystrophy reveals abnormal development of brain white matter. *Brain* **133**,
2448-2461, doi:10.1093/brain/awq180 (2010).
21 Elroy-Stein, O. Mitochondrial malfunction in vanishing white matter disease: a
disease of the cytosolic translation machinery. *Neural Regen Res* **12**, 1610-1612,
doi:10.4103/1673-5374.217329 (2017).
22 Fritsch, R. M., Schneider, G., Saur, D., Scheibel, M. & Schmid, R. M. Translational
repression of MCL-1 couples stress-induced eIF2 alpha phosphorylation to
mitochondrial apoptosis initiation. *J Biol Chem* **282**, 22551-22562,
doi:10.1074/jbc.M702673200 (2007).
23 Dzhagalov, I., St John, A. & He, Y. W. The antiapoptotic protein Mcl-1 is essential for
the survival of neutrophils but not macrophages. *Blood* **109**, 1620-1626,
doi:10.1182/blood-2006-03-013771 (2007).
24 Zhuo, L. *et al.* hGFAP-cre transgenic mice for manipulation of glial and neuronal
function in vivo. *Genesis* **31**, 85-94 (2001).
25 Kuang, Y. *et al.* Dicer1 and MiR-9 are required for proper Notch1 signaling and the
Bergmann glial phenotype in the developing mouse cerebellum. *Glia* **60**, 1734-1746,
doi:10.1002/glia.22392 (2012).
26 Guyenet, S. J. *et al.* A simple composite phenotype scoring system for evaluating
mouse models of cerebellar ataxia. *J Vis Exp*, doi:10.3791/1787 (2010).
27 Nakamura, A. *et al.* Bcl-xL Is Essential for the Survival and Function of Differentiated
Neurons in the Cortex That Control Complex Behaviors. *J Neurosci* **36**, 5448-5461,
doi:10.1523/JNEUROSCI.4247-15.2016 (2016).
28 Chong, M. J. *et al.* Atm and Bax cooperate in ionizing radiation-induced apoptosis in
the central nervous system. *Proceedings of the National Academy of Sciences* **97**,
889-894, doi:10.1073/pnas.97.2.889 (2000).
29 Lang, P. Y. *et al.* ATR maintains chromosomal integrity during postnatal cerebellar
neurogenesis and is required for medulloblastoma formation. *Development* **143**,
4038-4052, doi:10.1242/dev.139022 (2016).
30 Crowther, A. J. *et al.* Tonic activation of Bax primes neural progenitors for rapid
apoptosis through a mechanism preserved in medulloblastoma. *The Journal of
neuroscience : the official journal of the Society for Neuroscience* **33**, 18098-18108,
doi:10.1523/JNEUROSCI.2602-13.2013 (2013).
31 Glasgow, S. M. *et al.* Mutual antagonism between Sox10 and NFIA regulates
diversification of glial lineages and glioma subtypes. *Nat Neurosci* **17**, 1322-1329,
doi:10.1038/nn.3790 (2014).
32 Nishiyama, A., Boshans, L., Goncalves, C. M., Wegrzyn, J. & Patel, K. D. Lineage, fate,
and fate potential of NG2-glia. *Brain Res* **1638**, 116-128,
doi:10.1016/j.brainres.2015.08.013 (2016).
33 Dooves, S. *et al.* Astrocytes are central in the pathomechanisms of vanishing white
matter. *J Clin Invest* **126**, 1512-1524, doi:10.1172/JCI83908 (2016).

34   Hatten, M. E. Riding the glial monorail: a common mechanism for glial-guided
782   neuronal migration in different regions of the developing mammalian brain. *Trends
783   Neurosci* **13**, 179-184, doi:10.1016/0166-2236(90)90044-b (1990).
784   35   Rakic, P. Neuron-glia relationship during granule cell migration in developing
785   cerebellar cortex. A Golgi and electronmicroscopic study in Macacus Rhesus. *J Comp
786   Neurol* **141**, 283-312, doi:10.1002/cne.901410303 (1971).
787   36   Eddleston, M. & Mucke, L. Molecular profile of reactive astrocytes--implications for
788   their role in neurologic disease. *Neuroscience* **54**, 15-36, doi:10.1016/0306-
789   4522(93)90380-x (1993).
790   37   Dahl, D. & Bignami, A. Heterogeneity of the glial fibrillary acidic protein in gliosed
791   human brains. *J Neurol Sci* **23**, 551-563, doi:10.1016/0022-510x(74)90027-6 (1974).
792   38   Tamagno, I. & Schiffer, D. Nestin expression in reactive astrocytes of human
793   pathology. *J Neurooncol* **80**, 227-233, doi:10.1007/s11060-006-9181-6 (2006).
794   39   Schmidt-Kastner, R. *et al.* Astrocytes react to oligemia in the forebrain induced by
795   chronic bilateral common carotid artery occlusion in rats. *Brain Res* **1052**, 28-39,
796   doi:10.1016/j.brainres.2005.06.018 (2005).
797   40   Sahin Kaya, S., Mahmood, A., Li, Y., Yavuz, E. & Chopp, M. Expression of nestin after
798   traumatic brain injury in rat brain. *Brain Res* **840**, 153-157, doi:10.1016/s0006-
799   8993(99)01757-6 (1999).
800   41   Clarke, S. R., Shetty, A. K., Bradley, J. L. & Turner, D. A. Reactive astrocytes express
801   the embryonic intermediate neurofilament nestin. *Neuroreport* **5**, 1885-1888,
802   doi:10.1097/00001756-199410000-00011 (1994).
803   42   Bugiani, M. *et al.* Defective glial maturation in vanishing white matter disease. *J
804   Neuropathol Exp Neurol* **70**, 69-82, doi:10.1097/NEN.0b013e318203ae74 (2011).
805   43   Ito, D. *et al.* Microglia-specific localisation of a novel calcium binding protein, Iba1.
806   *Brain Res Mol Brain Res* **57**, 1-9, doi:10.1016/s0169-328x(98)00040-0 (1998).
807   44   Imai, Y., Ibata, I., Ito, D., Ohsawa, K. & Kohsaka, S. A novel gene iba1 in the major
808   histocompatibility complex class III region encoding an EF hand protein expressed in
809   a monocytic lineage. *Biochem Biophys Res Commun* **224**, 855-862,
810   doi:10.1006/bbrc.1996.1112 (1996).
811   45   Bugiani, M. *et al.* Hyaluronan accumulation and arrested oligodendrocyte progenitor
812   maturation in vanishing white matter disease. *Brain* **136**, 209-222,
813   doi:10.1093/brain/aws320 (2013).
814   46   Zhou, L. *et al.* Modeling vanishing white matter disease with patient-derived induced
815   pluripotent stem cells reveals astrocytic dysfunction. *CNS Neurosci Ther* **25**, 759-771,
816   doi:10.1111/cns.13107 (2019).
817   47   Ocasio, J. K., Bates, R. D. P., Rapp, C. D. & Gershon, T. R. GSK-3 modulates SHH-driven
818   proliferation in postnatal cerebellar neurogenesis and medulloblastoma.
819   *Development* **146**, doi:10.1242/dev.177550 (2019).
820   48   Tech, K. *et al.* Pyruvate Kinase Inhibits Proliferation during Postnatal Cerebellar
821   Neurogenesis and Suppresses Medulloblastoma Formation. *Cancer Res* **77**, 3217-
822   3230, doi:10.1158/0008-5472.CAN-16-3304 (2017).
823   49   Crowther, A. J. *et al.* Radiation Sensitivity in a Preclinical Mouse Model of
824   Medulloblastoma Relies on the Function of the Intrinsic Apoptotic Pathway. *Cancer
825   Res* **76**, 3211-3223, doi:10.1158/0008-5472.CAN-15-0025 (2016).

826   50   Cheh, M. A. *et al.* En2 knockout mice display neurobehavioral and neurochemical
827   alterations relevant to autism spectrum disorder. *Brain Res* **1116**, 166-176,
828   doi:10.1016/j.brainres.2006.07.086 (2006).

835