Implementation of a Bayesian Secondary Structure Estimation Method for the SESCA Circular Dichroism Analysis Package

Gabor Nagy[1] & Helmut Grubmüller[1]

October 2020

[1]: Department of Theoretical and Computational Biophysics, Max Planck Institute for Biophysical Chemistry, Am Fassberg 11, 37077 Göttingen, Germany

Keywords: CD spectroscopy, SS estimation, Bayesian statistics, model validation, accuracy improvement, SESCA

1 Abstract

Circular dichroism spectroscopy is a structural biology technique frequently
applied to determine the secondary structure composition of soluble proteins.
Our recently introduced computational analysis package SESCA aids the in-
terpretation of protein circular dichroism spectra and enables the validation of
proposed corresponding structural models. To further these aims, we present
the implementation and characterization of a new Bayesian secondary structure
estimation method in SESCA, termed SESCA_bayes. SESCA_bayes samples
possible secondary structures using a Monte Carlo scheme, driven by the like-
lihood of estimated scaling errors and non-secondary-structure contributions of
the measured spectrum. SESCA_bayes provides an estimated secondary struc-
ture composition and separate uncertainties on the fraction of residues in each
secondary structure class. It also assists efficient model validation by providing
a posterior secondary structure probability distribution based on the measured
spectrum. Our presented study indicates that SESCA_bayes estimates the sec-
ondary structure composition with a significantly smaller uncertainty than its
predecessor, SESCA_deconv, which is based on spectrum deconvolution. Fur-
ther, the mean accuracy of the two methods in our analysis is comparable, but
SESCA_bayes provides more accurate estimates for circular dichroism spectra
that contain considerable non-SS contributions.

21 1 Introduction

22 Circular dichroism (CD) spectroscopy in the far ultraviolet (UV) range (175-
23 260 nm) is an established method to study the structure of proteins in solution
24 [1, 2], because of the conformation-dependent characteristic CD signal of pep-
25 tide bonds that comprise the backbone of all proteins and oligo-peptides. In
26 particular, the CD spectrum is known to change with the secondary structure

(SS) of proteins, and markedly different spectra are observed for proteins rich in α-helices, β-sheets, and disordered regions [3, 4]. Because of these characteristic signals, it is common to interpret CD spectra by decomposing them into a set of basis spectra that each represent the average CD signal of pure (secondary) structure elements.

The CD analysis package SESCA (Structure-based Empirical Spectrum Calculation Approach) [5] allows for using several empirical basis spectrum sets in two methods. The first method predicts a theoretical CD spectrum from a proposed SS composition, which is typically obtained from a model structure or structural ensemble. The second method fits a measured CD spectrum to estimate the protein SS composition. Both methods can be used to validate protein structural models. The accuracy and precision of validation methods is mainly limited by scaling errors due to the uncertainty of the measured protein concentration and non-SS contributions that are not represented in the basis spectra. We have quantified the uncertainty caused by these deviations between measured CD spectra and their predicted SS signals previously [6].

The same study also revealed a potential caveat in the current SS estimation method used in SESCA. In this deconvolution method, a linear combination of selected basis spectra is used to approximate a measured CD spectrum of the protein of interest. The coefficients of the approximation with the smallest deviation are used to estimate the fraction of SS elements in the protein under the measurement conditions. Unfortunately, the interference caused by non-SS contributions may increase the deviation from the measured spectrum for some SS compositions and decrease it for others, which may lead to significant errors in deconvolution-based SS estimates.

To alleviate this problem, we developed and implemented a new SS estimation method for SESCA. The Python module, SESCA_bayes determines the

likelihood of putative SS compositions using a Bayesian inference framework for
a given measured CD spectrum and a basis spectrum set. This method uses
the expected joint probability distribution of deviations caused by scaling errors
and non-SS contributions, and thus fully accounts for the uncertainty caused by
these two experimental factors. Here, we describe the theoretical background,
general workflow, as well as input and output parameters of this implementa-
tion. Further, we will assess the accuracy and precision of this method through
a series of sample applications.

62 2   Theory: Bayesian SS probabilities

63 Our goal using this method is to determine the conditional probability $P(SS|CD)$
64 of SS compositions given a previously measured CD spectrum. According to
65 Bayes' rule [7], this probability can be inferred according to

$$P(SS_j|CD) \propto P(CD|SS_j) \cdot P(SS_j), \tag{1}$$

where $P(CD|SS_j)$ is the probability of observing the measured spectrum for a
protein with a given SS composition j (*i.e*, the likelihood function) and $P(SS_j)$
is the prior probability of the given SS composition of the protein. As shown
in Fig. 1 (top), the likelihood $P(CD|SS_j)$ is determined in five steps. First,
the SS signal is predicted from the SS composition of interest (C_{ji}) using an
appropriate basis spectrum set (B_{il}), as discussed in our previous study [5].
Second, if the basis set provides side chain corrections based on the protein
sequence, they are added to the predicted spectrum. Third, the measured CD
spectrum is rescaled to minimize the root-mean-square deviation (RMSD) from
the predicted spectrum. The obtained scaling factor α_j quantifies and eliminates
deviations from scaling errors of the measured spectrum, whereas the RMSD

from the rescaled spectrum ($RMSD_j$) quantifies the average deviation due to

unaccounted non-SS contributions. Once $RMSD_j$ and α_j (collectively CD de-

viations) are computed, the likelihood of such deviations is determined from

the joint probability distribution ($P_{RMSD,\alpha}$, see below), which also estimates

the likelihood of observing the measured CD spectrum for the given SS compo-

sition $P(CD|SS_j)$. Finally, to compute the posterior probability $P(SS_j|CD)$

of SS composition j, the CD spectrum likelihood is multiplied by the prior SS

probability.

3   Methods

3.1   Joint probability distributions

We computed discrete joint-probability distribution functions for SESCA_bayes

that can be used to determine CD spectrum likelihoods. These probability

distributions were computed from CD deviations extracted from SS estimations

of previously measured CD spectra. Reference CD spectra were taken from the

SP175 reference set [8], which contains 71 synchrotron radiation CD (SR-CD)

spectra of globular proteins with varying SS compositions. The CD spectrum

of Jacalin (SP175/41) was discarded from the data set due to issues reported

during the measurement and its unusually large estimated CD deviations.

The joint probability distribution functions of CD deviations were con-

structed as the sum of 70 two-dimensional Gaussian functions, each representing

the estimated scaling factors and non-SS contributions of a reference spectrum

from the SP175 set. The mean and the variance of these Gaussian functions

was determined by averaging over multiple $RMSD_j$ and α_j values obtained for

each CD spectrum from SS estimations using four different basis spectrum sets.

This approach yielded likelihood functions that were defined for a wide range

of possible CD deviations, and took the uncertainty due to discretization errors

of the basis spectrum determination into account.

In SESCA there are two types of basis sets, those that are solely based on on

SS compositions, and those that also include side chain corrections. Because the

average size of CD deviations differ for these two basis set types, we determined

two probability distributions shown in Fig. 2. The joint probability distribution

function for basis set without side-chain corrections (left) was calculated from

CD deviations estimated using the basis sets DS-dT, DSSP-1, HBSS-3, and DS5-

4. For basis sets including side-chain corrections, the joint probability of CD

deviations (right) were computed using the basis sets DS-dTSC3, DSSP-1SC3,

HBSS-3SC1, and DS5-4SC1. For clarity, the Figure shows both a linear (top

row) as well as logarithmic (bottom row) representation of the CD deviation

likelihood. For both likelihoods, the one-dimensional probability distribution

of $RMSD_j$ was also calculated, which can be used to estimate the secondary

structure from CD spectra without regards to the applied scaling factors, albeit

these estimates naturally have a lower precision.

3.2   Synthetic spectra

To test the accuracy of the Bayesian SS estimation method, six synthetic CD

spectra were created using a linear combination of the three basis spectra from

the DS-dT basis set (as discussed in our previous study [5]). To this aim,

the coefficients shown in Table 1 for the basis spectra α-helix, β-strand, and

Other for each spectrum were used. For five of six synthetic spectra (k= 1

to 5), random coefficients were generated from uniformly distributed random

numbers between zero and one, subsequently normalized to sum up to one. For

the sixth synthetic spectrum (k= 6), the coefficients 0.3, 0.4, and 0.3 as well as

the non-SS contributions (see below) were adopted from our previous study [6]

for comparison.

To model the effects of experimental deviations from the ideal SS signal, the

CD spectra were modified by adding non-SS signals and scaling errors. The size

of these CD deviations for each synthetic spectrum was quantified by the scaling

factors α_k and the root-mean-squared intensities of non-SS signals $RMSI_k$ listed

in Table 1. Synthetic spectrum 1 (k= 1) was a positive control without any

CD deviations (α_k= 1.0, $RMSDI_k$= 0.0 kMRE), spectra 2 and 6 included

small (0.4 kMRE) and large (3.5 kMRE) non-SS deviations, respectively, but

no scaling errors. CD deviations for spectra 3, 4, and 5 were drawn from the

marginal distributions of experimentally observed scaling factors and non-SS

contributions using rejection sampling.

The shapes of the non-SS signals were chosen as sums of Gaussian functions

$$S_{jl}^{\text{nonSS}} = \sum_{g=1}^{G} \frac{I_g}{\sqrt{2\pi\sigma_g^2}} \times e^{-\frac{(\lambda_l - \mu_g)^2}{2\sigma_g^2}}, \tag{2}$$

where the non-SS signal S_{jl} of protein j at wavelengths λ_l from 178 to 269

nm was computed from the following randomly chosen parameters. The number

of Gaussians G was chosen from the range 1 to 5, the relative peak intensity for

Gaussian g I_g was chosen between -20.0 and 20.0, with a peak position μ_g chosen

from 178 to 241 nm, and peak half-widths σ_g chosen between 2 and 37 nm. Once

the parameters were determined, the non-SS signal at every wavelength (using

1 nm spacing) was calculated, and the non-SS signal intensity was rescaled to

match the previously defined RMSI values in Table 1.

The final synthetic spectra were computed by determining the SS signals

first, by adding the appropriately scaled non-SS signal contributions in a second

set, and finally by rescaling the resulting CD spectrum according to the indicated

scaling factor.

152   4   Algorithm overview

153   Our newly implemented Python module SESCA_bayes.py performs a Monte-
154   Carlo (MC) sampling in SS space to determine the most probable SS compo-
155   sition of a protein based on its measured CD spectrum. Figure 3 shows the
156   flowchart of the algorithm that is divided into three phases: preparation, sam-
157   pling, and evaluation.

158   4.1   Preparation and input parameters

In the preparation phase, input, output, and run parameters are read based
on the user-provided command line arguments. If SESCA_bayes.py is used as
a Python module, an array of arguments can be processed by the function
Read_Args and passed to the *Main* function to run the algorithm. Arguments in
SESCA are identified by preceding command flags (marked by the "@" character
in the first position. There are four input files – shown as blue parallelograms
in Fig. 2 – that SESCA_bayes accepts, each read in white-space separated data
blocks stored as simple *ascii* text files.

The CD spectrum file (specified using the @spect flag) should contain two
columns, wavelength in nanometers (nm) and CD signal intensity in 1000 mean
residue ellipticity (kMRE) units. This file must be specified for SESCA_bayes,
and if no command flags are provided, the first argument is automatically rec-
ognized as a CD spectrum file.

The side-chain correction file (specified by @corr) is an optional file to add
baseline or sequence-dependent side-chain correction to the predicted CD spec-
trum, which are independent of the SS composition. If the basis spectrum
set has basis spectra to calculate side-chain contributions, these signals can be
computed before running SESCA_bayes, and added as a correction.

The Bayesian parameter file (@par) contains several data blocks, most im-

portantly, the binned probability distribution function of CD deviations $P_{RMSD,\alpha}$

(likelihood function), prior SS probability distributions for the SS composition

$P(SS_j)$ and scaling factors $P(\alpha_j)$, as well as the MC step parameters. If no

parameter file is provided by the user, SESCA_bayes.py uses one of two default

parameter files (Bayes_2D_SC.dat and Bayes_2D_noSC.dat) found in the "libs"

sub-directory of SESCA, depending on whether a side chain correction file was

provided or not. These files contain one of the two likelihood functions shown

in Fig. 2, and uniform prior SS probability distributions. A more detailed

description of the parameter blocks is provided in the examples sub-directory

(examples 5).

The basis set file (@lib) contains several data blocks for CD spectrum cal-

culations, including a block where the CD intensity of 3-6 basis spectra at each

wavelength (175-269 nm) is provided. Several derived basis sets are available

in libs sub-directory, and a detailed description of the data blocks is given in

example 1.

In addition to the input files, SESCA_bayes recognizes several additional

command flags to modify program behavior. The number of initial SS composi-

tions for MC sampling phase is specified by @size. The number of MC steps per

initial SS composition is set by @iter. The @scale flag allows the user to control

whether the measured CD spectrum is rescaled before determining the deviation

from the predicted CD spectra or not. In the absence of these command flags,

the values 100, 500, and 1 (yes) are used for the SS estimation.

Finally, three command flags control the output of SESCA_bayes.py; provid-

ing a "0" argument to any of these flags disables writing the associated output.

The command flag @write specifies the file name for the primary output, and

if no command flags are given, SESCA_bayes automatically recognizes the sec-

ond argument as primary output file. This file contains a summary of the

input parameters, binned posterior probability distributions for the SS compo-
sitions and scaling factors, as well as the most probable SS fractions and their
uncertainties. The command flags @proj and @data allows the user to print
secondary output files. The @proj flag specifies a file name for heatmap-style
two-dimensional projection of the posterior SS distribution. The projection is
made along two SS fractions selected using the @pdim flag Finally, the flag
@data specifies a file name for printing all the sampled SS compositions the
primary output is computed from, along with their estimated CD deviations,
prior and posterior probabilities. By default, only the primary output file is
printed into 'SS_est.out', and no secondary output is written.

215 4.2 Monte Carlo sampling

To determine the most probable SS composition of the protein based on its CD
spectrum, sampling of the SS space is required. To this aim, SESCA_bayes uses
a MC sampling scheme starting from N (set by @size) initial SS compositions,
drawn from the prior SS distribution using rejection sampling. As the center
part of Fig. 3 shows, at every step t of the MC sampling phase, a change
on each of the SS compositions ($C_{ji,t}$) is attempted. The change is realized
by transferring a given SS fraction between two randomly chosen SS classes,
yielding a new SS composition ($C'_{ji,t}$). The amount of the transferred SS fraction
from the donor class to the acceptor class is determined based on the distribution
specified in the Bayesian parameter file. If no distribution is provided, the
fraction is drawn from a Gaussian distribution with a mean of 0.05 and variance
of 0.1. To remain in the space of possible SS compositions, the transferred SS
fraction cannot exceed the current fraction assigned to the donor class, and
classes that currently have a fraction of zero assigned to them cannot be selected
as donors.

After the changes are attempted, the posterior probabilities P'_{jt} of the new
SS compositions are calculated (see Section 2) and compared to the posterior
probabilities (P_{jt}) of the SS compositions before the change. The attempted
change is accepted or rejected by applying the Metropolis criterion to the ratio
of posterior probabilities, *i.e.* the change is accepted if the ratio P'_{jt}/P_{jt} is
larger than a randomly generated number between zero and one. If the change
is accepted, $C'_{ji,t}$ is added to the sampled SS distribution and used as the initial
SS composition $C_{ji,t+1}$ in the next MC step, otherwise $C_{ji,t}$ is added to the
sampled SS distribution (again) and is used in the next MC step. This procedure
is repeated until the specified number of MC attempts is reached, and yields
$N \times t_{max}$ sampled SS compositions. The sampled SS compositions resemble the
prior SS distribution during the initial MC steps but converge towards an SS
distribution weighted by the posterior SS probabilities.

244 4.3   Sample evaluation

The sampled SS distribution is analysed in the evaluation phase, as shown in
the bottom part of Fig 3. To avoid the over-representation of very low posterior
probability SS compositions, a fraction of the initially sampled SS compositions
may be discarded from final SS distribution. This fraction can be set by the
user through the @discard flag, otherwise, the initial 5% of SS compositions is
discarded. The remaining probability-weighted ensemble of possible SS compo-
sitions is used to compute the estimated SS composition C_{ji}^{est} for the protein, the
estimated scaling factor α_j^{est}, as well as to approximate the discrete posterior
probability distribution for both quantities.
The estimated SS composition is determined by computing the mean and
standard deviation (SD) of each SS fraction over the sampled SS compositions.
Similarly, the most probable scaling factor is computed as the mean and SD of

scaling factors estimated for the sampled SS compositions. The discrete prob-

ability distribution for both scaling factors and SS compositions are computed

by binning all sampled SS compositions and scaling factors using the parame-

ters extracted from the prior distributions provided in the Bayesian parameter

file. The number of sampled SS compositions and scaling factors in each bin is

normalized by the final sample size to obtain the discrete probability distribu-

tions. The computed estimates, their uncertainties and the discrete probability

distributions are all written in the primary output file (defined by the @write

flag) and returned as output by the SESCA_bayes module. If requested (@proj

flag), the sampled SS compositions can be printed in a separate file. Finally,

the two-dimensional projection of posterior SS distribution along two chosen SS

fractions can also written into a separate output file (@proj flag), formatted as

a human readable heat map, that can be easily processed into images using *e.g.*

Python's Matplotlib module [9] or external visualization programs.

271   5   Testing the Algorithm

272   5.1   Accuracy and precision

The accuracy and precision of the Bayesian SS estimation was tested using the

10 CD spectra listed in Table 1. Six of these spectra (k= 1-6) are synthetic spec-

tra that were generated from a given SS composition, but modified by adding

artificial non-SS signals and scaling errors (see Section 3.2) to emulate CD devi-

ations in real measured spectra. The remaining four CD spectra (k= 7-10) are

measured spectra from the SP175 set [8], for which the estimated SS composi-

tions are compared to those extracted from the (protein data bank) structure

of the reference protein. Table 1 also lists the (estimated) CD deviations of all

ten CD spectra, quantified by the scaling factors α_k and the root-mean-square

intensity ($RMSI_k$) of non-SS signals in each spectrum.

To test the accuracy of SESCA_bayes, we estimated the SS composition

of the above ten CD spectra using the same DS-dT basis set with three SS

classes (α-helix,β-strand, and Other) that was used to generate the synthetic

spectra. The obtained Bayesian estimates for the test set are summarized in

Table 2. This table includes the mean and SD (in parentheses) of SS fractions

of the sampled posterior distributions, as well as the total SS deviation from

the reference SS compositions, computed according to

$$\Delta SS_k = \frac{1}{2} \sum_{i=1}^{F} |C_{ki}^{est} - C_{ki}^{ref}|, \tag{3}$$

where C_{ki}^{est} are the estimated SS fractions and C_{ki}^{ref} are the reference SS fractions

listed in Table 1.

The obtained SS fractions show a fairly consistent 0.03 to 0.06 uncertainty.

As expected, 27 of 30 SS fractions are within two SD of their reference value,

with no significant difference in accuracy between synthetic and measured CD

spectra. In addition, the calculated total SS deviations (ΔSS) from the reference

structures range between 0.03 and 0.12, and eight of ten values are also smaller

than the estimated uncertainty of the estimation (two SD) that was calculated

from the individual SD of SS fractions (σ_{ki}) according to

$$\sigma_k = \frac{1}{2} \sqrt{\sum_{i=1}^{F} \sigma_{ki}^2}. \tag{4}$$

5.2 Comparison to deconvolution

Next, we compare the accuracy and precision of the Bayesian estimates to that

of SS estimates obtained through spectrum deconvolution. To this aim, we esti-

mated SS compositions with the deconvolution module of SESCA (SESCA_deconv)

for the same ten CD spectra (Table 1), using the same DS-dT basis spectrum set.

The deconvolution was carried out using the most accurate protocol (method

D2) tested previously [6]. This method constrains the basis spectrum coefficients

to positive values, but normalizes them to unity only after the search for the best

approximation. The SS compositions obtained using SESCA_deconv are listed

in Table 3, along with the total SS deviations from reference SS compositions

(found in Table 1). The total SS deviation of deconvolution estimates (ΔSS_k)

ranges from 0.0 to 0.29. The mean SS deviation for the whole set (0.08) is

similar to that of the Bayesian estimates (0.07), but shows a significantly larger

scatter (0.9 vs. 0.03). All three CD spectra with larger than average SS devi-

ations (k= 3,4,8) have large non-SS contributions (2.0-2.9 kMRE), which is in

line with our previous findings that non-SS contributions may be detrimental

to the accuracy of deconvolution methods.

Although the SESCA_deconv module does not provide information on the

uncertainty of individual SS fractions, many SESCA basis sets (including DS-

dT) include a calibration curve to estimate the expected total SS deviation if

the true SS composition is unknown. This curve was computed from $4.9 \times$

10^5 synthetic spectrum-structure combinations, which were binned according to

their estimated non-SS contributions ($RMSD_j$), to provide an expected mean

and SD of SS deviations for a given (rescaled) RMSD. Comparing the true SS

deviations of the deconvolution results with their estimated values shows that

these estimates correctly describe the precision of the deconvolution method:

six of ten ΔSS_k values are within 1 SD of the estimated total deviation, and

all ten fall within 2 SD. However, the average uncertainty of the deconvolution

(0.09) is again considerably larger than that of the Bayesian SS estimates (0.04),

and it increases with increasing non-SS contributions.

In summary, Bayesian SS estimation and spectrum deconvolution provides

SS estimates that – in most cases – have a similar accuracy. However, Bayesian

SS estimates are considerably more precise when significant non-SS contribu-

tions are present in the measured spectrum. Further, the Bayesian approach

provides uncertainties for each individual SS fraction as well as the optimal scal-

ing factor for the measured CD spectrum, which is an additional advantage of

the new method.

336   5.3   Example spectrum analysis

To further investigate the differences between the two methods, we analysed the

SS estimates for the CD spectrum with the largest difference between the de-

convolution and Bayesian SS estimates. Figure 4A shows the obtained posterior

SS probability distribution for synthetic spectrum 3, which contains larger than

average non-SS contributions (2.02 kMRE). The heatmap shown in Fig. 4A

illustrates that the most likely SS compositions are indeed clustered around the

SS composition the synthetic spectrum was created from (shown as a red cross),

with the highest posterior probability regions (shown in dark green) located in

the immediate ($\Delta SS_k < 0.05$) vicinity of correct SS composition. However, the

SS composition determined by deconvolution (purple cross) has a much higher

α-helix content and it is not in a high-probability region in the Bayesian SS

estimation.

To examine why the two algorithms evaluate the proposed SS compositions

differently, in Fig. 4B we computed the predicted CD signals of the two es-

timated SS compositions, rescaled them, and compared them to the synthetic

spectrum, as is done during the deconvolution process. The figure shows that

with the proper scaling factor both SS compositions approximate the synthetic

spectrum well, but the deconvolution estimate (purple dashed line, $RMSD_j$:

1.31 kMRE) fits slightly better than the Bayesian estimate (blue dashed lines,

$RMSD_j$: 1.71 kMRE).

In contrast, the Bayesian SS estimation rescales the synthetic CD spectrum
to match the predicted spectra, and evaluates the likelihood of the SS com-
positions based on the joint probability of their non-SS contributions $RMSD_j$
and scaling factor α_j, as shown in Fig 4C. Although the two estimates have a
comparable RMSD in this method as well, the deconvolution estimate requires
a scaling factor (α_j: 1.99) to achieve a good agreement that is shown to be very
unlikely according to the joint-probability map in Fig. 3. Comparing the two
estimated SS signals (dashed lines) to the SS signal of the true SS composition
(in red) illustrates how considering scaling factors improves the precision of the
SESCA_bayes. In this case, eliminating SS compositions with unlikely scaling
factors from the sampled distribution allowed a fairly accurate (RMSD: 0.99
kMRE) approximation of the true SS signal.

Declarations

Funding: This research project was funded and supported by the Alexander
von Humboldt Foundation and the Max Planck Society.

Conflicts of interest: The authors declare no conflict of interest.

Code availability: the new SESCA implementation based on this study is
available at: https://www.mpibpc.mpg.de/sesca

License: SESCA is free available under GNU general public license 3 (GPLv3)

Authors' contributions: G.N. designed and performed the computational
analysis, and implemented code improvements. H.G. is the corresponding au-
thor, and assisted with the conceptualisation. Both authors contributed to
writing the manuscript.

Acknowledgements: The authors would like to thank K. Blom and P. Kellers
for the aid in editing the manuscript.

References

[1] Gerald D. Fasman, editor. *Circular Dichroism and the Conformational Analysis of Biomolecules.* Springer US, Boston, MA, 1996.

[2] Norma J. Greenfield. Methods to estimate the conformation of proteins and polypeptides from circular dichroism data. *Analytical biochemistry*, 235(1):1–10, 1996.

[3] Sharon M. Kelly, Thomas J. Jess, and Nicholas C. Price. How to study proteins by circular dichroism. *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics*, 1751(2):119–139, August 2005.

[4] S. Brahms and J. Brahms. Determination of Protein Secondary Structure in Solution by Vacuum Ultraviolet Circular Dichroism. *Journal of Molecular Biology*, 138:147–178, 1980.

[5] Gabor Nagy, Maxim Igaev, Nykola C. Jones, Søren V. Hoffmann, and Helmut Grubmüller. SESCA : Predicting Circular Dichroism Spectra from Protein Molecular Structures. *Journal of Chemical Theory and Computation*, August 2019.

[6] Gabor Nagy and Helmut Grubmüller. How accurate is circular dichroism-based model validation? *European Biophysics Journal*, 49(6):497–510, September 2020.

[7] Andrew Gelman and John B. Carlin. *Bayesian Data Analysis.* Texts in Statistical Science. Chapman and Hall/CRC, 3rd edition, 2014.

[8] J. G. Lees, A. J. Miles, F. Wien, and B. A. Wallace. A reference database for circular dichroism spectroscopy covering fold and secondary structure space. *Bioinformatics*, 22(16):1955–1962, August 2006.

[9] J. D. Hunter. Matplotlib: A 2d graphics environment. *Computing in Science & Engineering*, 9(3):90–95, 2007.

Tables

Table 1: SS compositions and CD deviations of model proteins. Columns show the index and name of the respective model protein, the fraction of its amino acids assigned to the SS classes α-helix, β-strand, and Other, as well as scaling factors α_k and root-mean squared intensities $RMSI_k$ of non-SS signals to quantify scaling errors and non-SS contributions in the protein CD spectrum, respectively. Synth denotes synthetic spectrum in the proteins name, whereas Lysm, Dqd-1, and Sub-C abbreviate Lysozyme, Dehydroquinate dehydratase I, and Subtilisin Carlsberg, respectively. Note that SS fractions, scaling factors, and non-SS contributions for all synthetic proteins (k= 1-6) were parameters used to generate their CD spectrum, whereas for real reference proteins (k= 7-10), all values were computed based on their measured spectra and protein data bank structures (193L, 2DHQ, 1KU8, and 1SCD, respectively).

k	protein	α-helix	β-strand	Other	α_k	$RMSI_k$
1	Synth-1	0.11	0.40	0.49	1.0	0.0
2	Synth-2	0.41	0.20	0.39	1.0	0.4
3	Synth-3	0.43	0.10	0.47	1.2	2.0
4	Synth-4	0.27	0.26	0.47	1.6	2.7
5	Synth-5	0.00	0.33	0.67	1.4	0.7
6	Synth-6	0.30	0.40	0.30	1.0	3.6
7	Lysm	0.35	0.03	0.62	1.1	1.0
8	Dqd-1	0.43	0.18	0.39	1.1	2.9
9	Jacalin	0.01	0.28	0.71	0.3	3.2
10	Sub-C	0.30	0.12	0.58	0.4	1.2

Table 2: Bayesian secondary structure estimates. The table lists the index and name of the model protein, the estimated fraction of its amino acids assigned to SS classes α-helix, β-strand, and Other, as well as the total SS deviation ΔSS_k from the reference SS compositions shown in Table 1. The uncertainty (standard deviation) of each SS fraction and deviation is given in parentheses. Estimates that are more than 2 SD away from their reference value are highlighted in red.

k	protein	α-helix	β-strand	Other	ΔSS_k
1	Synth-1	0.14 (0.05)	0.44 (0.06)	0.43 (0.04)	0.06 (0.04)
2	Synth-2	0.49 (0.06)	0.19 (0.06)	0.32 (0.06)	0.08 (0.05)
3	Synth-3	0.45 (0.06)	0.12 (0.05)	0.43 (0.05)	0.03 (0.04)
4	Synth-4	0.22 (0.04)	0.21 (0.05)	0.57 (0.04)	0.09 (0.04)
5	Synth-5	0.03 (0.04)	0.26 (0.03)	0.71 (0.05)	0.07 (0.04)
6	Synth-6	0.36 (0.05)	0.32 (0.04)	0.31 (0.06)	0.08 (0.04)
7	Lysm	0.38 (0.05)	0.04 (0.05)	0.57 (0.05)	0.05 (0.04)
8	Dqd-2	0.48 (0.06)	0.06 (0.05)	0.47 (0.05)	0.12 (0.05)
9	Jacalin	0.01 (0.04)	0.31 (0.06)	0.68 (0.06)	0.03 (0.05)
10	Sub-C	0.26 (0.05)	0.13 (0.04)	0.61 (0.04)	0.04 (0.04)

Table 3: Secondary structure estimates based on spectrum deconvolution. The table lists the index and name of the model protein, the estimated fraction of its amino acids assigned to SS classes α-helix, β-strand, and Other, as well as the total SS deviation ΔSS_k from the reference SS compositions shown in Table 1. The values in parentheses after ΔSS_k show the mean and SD of the estimated total SS deviation computed from the rescaled RMSD between the measured (generated) CD spectrum and predicted spectrum of the SS estimate.

k	protein	α-helix	β-strand	Other	ΔSS_k
1	Synth-1	0.11	0.40	0.49	0.00 (0.00 ± 0.02)
2	Synth-2	0.41	0.20	0.39	0.00 (0.05 ± 0.03)
3	Synth-3	0.72	0.03	0.24	0.29 (0.16 ± 0.09)
4	Synth-4	0.19	0.22	0.59	0.12 (0.08 ± 0.05)
5	Synth-5	0.01	0.33	0.66	0.01 (0.06 ± 0.04)
6	Synth-6	0.31	0.31	0.37	0.08 (0.14 ± 0.09)
7	Lysm	0.34	0.06	0.60	0.03 (0.07 ± 0.05)
8	Dqd-2	0.51	0.04	0.45	0.14 (0.09 ± 0.06)
9	Jacalin	0.00	0.35	0.65	0.07 (0.20 ± 0.09)
10	Sub-C	0.25	0.13	0.62	0.05 (0.07 ± 0.05)

Figures

Figure 1: Secondary structure probability calculation scheme. The figure depicts the algorithm to compute the posterior probability of a given secondary structure j, based on its prior probability, and the deviations between its predicted CD signal and a given measured CD spectrum. Input data are depcited as blue parallelograms, operations as white rectangles, and decisions as white diamonds.

Figure 2: The panels depict the heat map representation of two likelihood functions provided for Bayesian SS estimation with SESCA. The estimated joint-probability distributions are shown for basis spectra that A) predict CD signals solely from SS information (left) and B) also include CD corrections from sequence-based side-chain information (right). Panels on the top and bottom show the same probability distributions using a linear and logarithmic color scale, respectively.

Figure 3: Schematic workflow of the Bayesian secondary structure estimation module in SESCA. The scheme depicts input data files as blue parallelograms, data on the sampled SS compositions are shown as a red parallelogram. Operations are depicted as white rectangles, and decisions are shown as white diamonds. Posterior probability calculation operations (see Fig. 1) are highlighted as yellow rectangles on the scheme.

Figure 4: SS estimation for a synthetic CD spectrum. The figure compares the true SS composition (shown in red) with SS compositions obtained from Bayesian SS estimation (in blue) and spectrum deconvolution (in purple). A) shows the posterior probability distribution of sampled SS compositions in a heat map representation and indicates the true SS composition and the deconvolution SS estimate as crosses. The SS compositions, estimated scaling factors, and SS deviations are also listed in a tabulated format on the top. The difference on how the two estimates are evaluated by B) the deconvolution and C) Bayesian SS estimation are also shown. During deconvolution, the predicted CD signal of SS estimates is rescaled to match the measured CD spectrum, and the measure of quality is solely the RMSD. In the Bayesian approach, the measured spectrum is rescaled to match the predicted SS signals, and both the RMSD-s and the scaling factors are used to determine the most likely SS composition.