Normalizing the brain connectome for communication through synchronization

S. Petkoski[1], V.K. Jirsa[1]*†

[1]Aix-Marseille Univ, Inserm, INS, Institut de Neurosciences des Systèmes, 13005 Marseille, France.

*Correspondence to: spase.petkoski@univ-amu.fr & viktor.jirsa@univ-amu.fr

†Both authors equally contributed to the work.

Keywords: Synchronization, Time-delays, Normalized connectome, Spectral activation patterns, Oscillations, Resting state networks.

ABSTRACT:

Networks in neuroscience determine how brain function unfolds, and their perturbations lead to psychiatric disorders and brain disease. Brain networks are characterized by their connectomes, which comprise the totality of all connections, and are commonly described by graph theory. This approach is deeply rooted in a particle view of information processing, based on the quantification of informational bits such as firing rates. Oscillations and brain rhythms demand, however, a wave perspective of information processing based on synchronization. We extend traditional graph theory to a dual particle-wave-perspective, integrate time delays due to finite transmission speeds and derive a normalization of the connectome. When applied to the data base of the Human Connectome project, we explain the emergence of frequency-specific network cores including the visual and default mode networks. These findings are robust across human subjects (N=100) and are a fundamental network property within the wave picture. The

normalized connectome comprises the particle view in the limit of infinite transmission speeds

and opens the applicability of graph theory to a wide range of novel network phenomena,

including physiological and pathological brain rhythms. These two perspectives are orthogonal,

but not incommensurable, when understood within the novel here proposed generalized

framework of structural connectivity.

AUTHOR SUMMARY:

All networks are composed of nodes and links, forming the structural frame, in which

communication occurs. We demonstrate that graph theoretical tools make the implicit

assumption of information transmission via exchange of bits, suggesting that the stronger

connected nodes are more impactful upon the remainder of the network. This corollary does not

extend to communication through oscillations, which is the prominent information carrier in

brain networks. We extend traditional network analysis to the oscillatory domain and derive a

novel network normalization including descriptive metrics. Along the prototypical example of

the brain as a network, we illustrate the consequences of this novel approach and demonstrate

that the normalization robustly explains the emergence of the prominent frequency-specific

network cores, which cannot be understood within the traditional framework.

INTRODUCTION

Network theory significantly advanced our understanding of complex processes in nature,

ranging from gene regulation of protein interactions (*1*), coordination of brain activity (*2*) to

social networks (*3*). Connectivity is the dominant concept that shapes the capacity of a network

to transmit information (*4*) and is described by its topological and statistical properties (*5*). In

neuroscience, network theory has been applied on several scales, including microscopic

neocortical microcircuitry (*6*), macroscopic structural connectivity – connectome (*7*), and

networks of coordinated brain activity, also called functional connectivity (*8*). Models of brain

networks based on connectomes (*9*) have demonstrated individual predictive power in resting

state paradigms (*10*), cognitive tasks (*11*) and brain disease such as epilepsy (*12*). In all these

applications, there remains a deeply rooted understanding of signal transmission in which bits of

information are transmitted between network nodes, independently of the underlying oscillations

and brain rhythms. But, as the particle picture is complemented by a wave picture in physics,

similarly in neuroscience there are oscillatory processes deeply implicated in healthy and

pathological brain activation (*13*), such as cognitive functions (*14*) or aberrant discharges in

epilepsy (*12*). How such rhythms support the communication through the phenomena such as

phase and frequency dependance does not have a fundamental description within the particle

view of information processing, which describes network processes by activations, but does so in

the wave picture that imposes a language in terms of synchronization.

The importance of the particle-wave duality in network neuroscience becomes evident when

considering signal transmission delays, present in large scale brain networks and ranging on the

order of 10-200 ms (*15*). As physiological rhythms in the brain lie in the same range (*16*), shifts

in arrival times due to delays have minimal to no effects in the particle view, but result in

frequency dependent effects in the wave view, where a synchronized pair of oscillators may

switch from full synchrony to anti-synchrony. As such, frequency and time delays become

inseparable properties of the network and together with the connectome determine the

synchronization and nodes' activity (*17*, *18*). This also confirms previous computational studies

about the effects of the space-time structure of the brain on its emergent dynamics (*19*), and

could also determine the spectrally dependent information processing capacities based on

synchrony (*20*).

Current state of the art network neuroscience is mainly focused on the topological aspects in

describing the connectivity of the brain (*2*, *21*), thus limits itself to the particle view only and

omits space and physical distances between the interacting units (*22*). Possible exceptions from

this are the cases when structural network motifs are shown to be related with the functional hubs

due to the time-delays (*23*), but without spectral specificity. In the rare cases when lengths of the

links are studied (*24*), it is still done in static manner, without consideration for the impact of the

delays on the emerging dynamics, thus critically demanding the extension of network

neuroscience to the wave picture. The only attempts to study negative links between brain

regions were limited to the clustering of functional data with negative correlations (*22*), without

any application to the structural links, for which there was never a mechanistic explanation that

would render them effectively inhibitory for the network synchronization.

RESULTS

Particle versus wave representation for network dynamics. The particle-wave dichotomy for

network communication appears in networks dynamics due to the finite transmission velocities

for the signals between spatially distributed nodes. The governing equation reads

85
$$\dot{x}_i = F(x_i) + \sum_j w_{ij} h(x_j(t - \tau_{ij})),$$
(1)

where x_i is a vector of states, F is a non-linear function, and h is the coupling function. Within

the particle view, x is a one-dimensional scalar variable such as firing rate, rendering the

interactions to be frequency-independent, Fig. 1 (A). Consequently, the strongest flow of

information and the activity is generally along the links with the strongest coupling weight (*25*),

although there are cases when hubs are significantly more influential (*26, 27*). In addition, time-

delays are less important during single events that occur independently of the possible

underlying oscillatory activity (middle plot of Fig. 1(A)), such as the action potentials that make

up subsequent metrics such as firing rate. Similar examples of particle-like interactions happen

during strong perturbations and stimulations (*28*), or when the coupling is no longer weak and

bifurcations occur, such as during seizure propagation in epilepsy (*29*).

Networks of oscillators, on the other hand, are often used to study dynamical systems for which

the local activity is multidimensional and nonlinear (*30*). They have been conceptualized to be

responsible for the communication in the brain through coherence (*20*) and synchronization (*31*),

Fig. 1 (A), but this aspect is still unlinked on the network level to the features of the structural

links, i.e. weights and time delays, that shape the synchronization of the brain network (*18*). For

weak couplings, dynamics of the nonlinear oscillatory system in Eq. (1) are captured by phase

models, of which the simplest and the most elaborate due to the analytical tractability is the

Kuramoto model (*32*). For synchronization at frequency Ω, the model reads (*33*)

104
$$\dot{\theta}_i = \omega_i - \sum_j w_{ij} \sin\left(\theta_i - \theta_j + \Omega\tau_{ij}\right), \tag{2}$$

where θ_i denotes the phase of the *i*-th node oscillating with a natural frequency ω_i. Note that if

time-delays were short so that they would not appear explicitly in the phase model (*33*), then Eq.

(2) would have still represented the phase model for the case of synchronization.

From Eq. (2) follows that when delays are negligible compared to the time-scale of the system,

the interactions are governed only by the weights, as in the particle case. Due to better

tractability and the readily applicable tools from the graph theory, this has been the default

network representation in neuroscience. However, when the time-delays are comparable to the

time-scale of the intrinsic oscillations, they need to be included in the analysis of network

dynamics. This system can only be solved numerically, in order to unveil the activation patterns

at different frequencies as there is no alternative graph theoretical approach that integrates the

topology and the impact of the time-delays on the oscillations. Hence the systematic effect of the

time-delays on the emergent dynamics in Eq. (1) is concealed, beyond the synchronizabillity and

the order parameter which can be calculated using the Ott and Antonsen ansatz (*34*), but only for

all to all couplings and spatially decomposed delays (*35*).

Fig. 1. **Spatio-temporal network organization and wave coupling weights for synchronized networks**. (A) Particle (static or frequency-independent) and wave (synchronization-dependent) interactions with delay τ, due to distance l, and propagation velocity v. (top) Transmission of packets over non-oscillatory local dynamics; (middle) a particle type of transmission that does not depend on the coherence (e.g. during strong perturbation) and is thus independent of the underlying oscillatory dynamics; (bottom) the communication is dependent on the synchronization. (B-D) In-and anti-phase synchronization for fixed frequency and different time-delays for 2 oscillators (B), and for networks with a resonant (C) and non-resonant (D) spatio-temporal alignment of the oscillators. Nodes with same color are in-phase with each other, and anti-phase with others, as also illustrated with the waveform. Time-delays are shown relatively to the period of oscillations T, and the normalization factor is shown for some of the links. (D) Midpoints of links with in-phase oscillations are highlighted yellow, and green if anti-phase, and the sum of all positively and negatively contributing links (yellow minus green links) to each node is also shown. (E-G) Wave coupling weights for a particle (or static) case (E) and for an optimal, (F) and suboptimal (G) phase arrangements of nodes. Links contributing positively to the synchronization are yellow, and green are negative contributions. Lines have width proportional to $|w_{ij}^{(f)}|$ (shown next to the links) and the size of the circles corresponds to the nodes spectral capacity (dark) and strength (colored), both shown for each node.

To go beyond the static representation of networks, we use the insight that the impact of the direct link in the phase difference between oscillators can be separated from the rest of the network (see the Model section), leading to solutions given as

$$\Delta\theta_{ij} = \sin^{-1}\frac{\Delta\omega_{ij} - I_{ij}}{w_{ij}\cos\Omega\tau_{ij}}. \tag{3}$$

Here $\Delta\omega_{ij} = (\omega_i - \omega_j)/2$, and normalized wave coupling $w_{ij}^{(f)} = w_{ij}\cos\Omega\tau_{ij}$ describes the

impact of the direct link between the oscillators. The influence of all the other links of these two

nodes is contained in I_{ij}. The cosine term vanishes for particle-like communication $\Omega\tau_{ij} \to 0$,

simplifying Eq. (3) to the case of no delays, when weights are the only factor shaping the

network dynamics. For interactions of synchronized oscillations, wave coupling $w_{ij}^{(f)}$ unifies

spatio-temporal aspects of each link by modulating weights with the frequency-specific impact

of the time-delays. This reflects duality in the network interactions, which in the wave case is

reliant on the timings of the waveforms arriving from the distant nodes, Fig. 1 (B-D).

As illustrated in Fig. 1 (B), two delay-coupled oscillators can synchronize either in-phase or anti-

phase (*17*, *30*), depending on the sign of $\cos\Omega\tau$. The former is stable for positive wave

couplings, the latter for negative. The same is true for networks (*17*) and not limited to phase

oscillators (*18*). Time-delays can be perfectly spatially distributed to cause maximum

synchronization at a given frequency Fig. 1 (C), but a more realistic scenario is when their

distribution causes some links to decrease the network synchronization, i.e. if $w_{ij}^{(f)} < 0$ for in-

phase nodes, or if $w_{ij}^{(f)} > 0$ for anti-phase. One possibility here is the oscillators rearranging

their phases to minimize this disturbance Fig. 1 (D). From here, to identify the skeleton of the

wave synergy we define a spectral strength and capacity for each network node

159
$$S_i(f, Q_i) = \sum_{j \in Q_i} w_{ij}^{(f)} - \sum_{j \notin Q_i} w_{ij}^{(f)}, \tag{4}$$

160
$$C_i^{(f)} = \sum_j |w_{ij}^{(f)}|. \tag{5}$$

In the static case, both metrics yield the topological node strength, Fig. 1 (E). $S_i(f, Q_i)$ is adapted

from the particle case node strength (*22*) by accounting for possible inhibitory contribution of

some links to the overall synchronization, i.e. negative wave coupling from an in-phase node, or

a positive one from an anti-phase node, Fig. 1 (D, F, G). The cluster Q_i thus contains all the

nodes that are in phase with the node i. Related to this, $C_i^{(f)}$ gives the upper bound of node's

synchronizabillity and hence of its $S_i(f, Q_i)$, Fig. 1 (F, G). Therefore, it shows the strength of

the node when all its links contribute positively to its dynamics. As shown in the same plots,

different phase arrangements of the nodes change their spectral strength. A hub node can become

more peripheral, or vice-versa, while still being constrained by its spectral capacity, which does

not change as different arrangements are realized for a fixed frequency.

For simplicity, in the rest of the manuscript we will omit the explicit dependence of the spectral

strength on the node-specific cluster Q_i, but instead we will refer to a specific in/anti phase

arrangements on the network level that imply specific clustering for the nodes. For example, in

most of the results we will show results for $S_i^{(f)}$ from a clustering that maximizes its sum over

the whole network.

<u>Normalized Connectome and Spectral Dependent Activity of Brain Regions.</u> The wave

coupling weights are frequency dependent, and in the case of the connectome this causes

activations of different network cores. We assume homogeneous propagation velocity, which is

often used as a first approximation, resulting in time delays being defined by the lengths of the

links (*9, 17–19*), i.e. $\tau_{ij} = l_{ij}/v$. The cosine term modulates the strength of the links, which can

even change the sign, reverting once positive interactions depending on the frequency. This is

demonstrated for connectomes of a human (*36*) and mice (*37*), Fig. 2 (A).

The spectral strength and capacity are illustrated in more details in Fig. 2 (B, C) for three

different frequencies, and they are shown for all the brain regions ordered by lobes in Fig. 3. The

phase arrangement of the nodes is obtained by maximizing the overall spectral strength of the

network (see Methods for more details). For example, the two spectral graph theoretical metrics

in Fig. 2 (B, C) illustrate that the highest spectral capacity for regions of cingulate are at 20 Hz

and their strength is lowest at 10 Hz, while the occipital lobe has highest spectral strength and

capacity at 10 Hz. Large spectral capacity implies possibility for a given node to get stronger

strength during different phase arrangements, since as illustrated in Fig. 1 (B), even for a

network of several nodes it can already become impossible not to have negatively contributing

links that inhibit the synchronization. The ratio $S_i^{(f)}/C_i^{(f)}$ shows what proportion of the weighted

links are contributing positively to the spectral strength of the node for the optimal phase

arrangement. Hence, the superior parietal (SupPa) region is by far the strongest at 40 Hz due to

the more positively contributing links, as compared to the precuneus, which is much weaker at

this arrangement, besides having a similar capacity.

Fig. 2. **Normalized connectomes at different frequencies**. (A) Wave couplings for a human (top) and averaged mice (bottom) connectomes. (B) Spectral strength and capacity, and their ratio for regions of the left hemisphere ordered by lobes for 3 different frequencies. Full names of the regions are shown in Tab. S14. (C) Spectral strength (blue or red, indicating anti-phase arrangements) and capacity (grey) for cortical regions of a human brain. The links with positive contribution to the optimal phase arrangement are yellow, and others are green (c.f. Fig. 1 (B)).

At each frequency only the links with spectral strengths above 0.33 of the mean strength are

shown for simplicity.

In Fig. 3 relative activation patterns of all the brain regions are shown for the frequencies up to

80 Hz. They are more heterogeneous for the frequencies between alpha and lower gamma, and

homotopic brain regions generally show similar activation patterns, due to the high symmetry of

the weights and the delays of the connectome. Frequency-normalized values of the spectral

strength and capacity unveil the regions with relatively strongest activity at each frequency,

whereas the relative change of the activation compared with the particle case is also shown (right

plots in Fig. 3 (C, D)). The strength of each region's activity relative to the particle case, as well

as the mean relative activity for the whole brain are also shown in left plots in Fig. 3 (C, D). For

every region the strength generally decreases with frequencies, as it is clearly seen in their mean,

while the capacity has much steeper decline up to alpha frequencies, when it slightly recovers

and stays largely constant.

Fig. 3. **Spectral strength and capacity for brain regions.** Mean values are shown over 100

subjects and for frequencies in a step of 1Hz. Brain regions are ordered by lobe with exact

ordering within lobes given in Tab. S.14, each right hemisphere region is placed right after its

left-hemisphere counterpart. (A) $S^{(f)}$ and $C^{(f)}$ normalized across each frequency, (B, C) their

ratio and difference compared to the case with no delays, where the former is for the non-

normalized metrics, to reveal the frequency dependent mean activity.

<u>Spectral Dependent activity of RSNs and brain lobes.</u> Studies of functional connectivity have

demonstrated that signals in brain regions that are related in different tasks, can also correlate

even in the absence of an apparent task, hence the notion of resting state networks (RSNs). The

most consistent RSNs are default mode network (DMN), visual, sensory/motor (SensMot),

auditory, executive control (ExecCont) and frontal-parietal (FronPar) (*38*). In Fig. 4 we show the

average spectral strength and capacity of brain regions of 100 subjects, further projected on

RSNs and anatomical lobes. Regions with the strongest activity averaged over the subjects are

shown in Fig. 4 (A), through the mean $S_i^{(f)}$, which is normalized at different frequency bands.

The same panel contains the boxplots of the strengths of the RSNs of different subjects. The

panel (B) of Fig. 4 contains the same metric, but the in-/anti-phase arrangements of the nodes are

taken into account through the sign of $S_i^{(f)}$, which is now plotted in red-white-blue color code

with red regions being anti-phase with the blue. In Fig. 4 (C) the strength at each RSN is

projected across the bands, complementing the information about the spectral affinity of RSN.

For example, Visual network is by far the most active during alpha frequencies, Fig. 4 (C), even

though the alpha frequency also contains a strong activity of the Frontal Parietal network, Fig. 4

(A). Fig. 4 (D) shows the mean and the standard deviation over the 100 subjects of the spectral

strength and capacity of each lobe normalized at each frequency. It shows that the robustness

over the subjects is much stronger for lower frequency bands (up to around 30 Hz), similarly

with the most robust empirically observed spectral activation patterns (*39–42*).

244
Fig. 4. **Cortical activation patterns of 100 subjects for different frequencies**. (A) Mean

spectral strength is shown for different cortical regions (higher for warmer colors) and frequency

bands (Tab. S13), and activation of RSNs is projected in the frequencies. (B) In-/anti-phase

arrangements of mean $S_i^{(f)}$ are indicated with red and blue. (C) Relative contribution of the

frequency bands per RSN. Boxplots contain the scatter plots of all 100 subjects. (D) Mean

spectral strength and capacity (lines) and regions of two standard deviations (shaded) calculated

in step of 1Hz for each lobe. The values per given RSN and lobe are normalized by the number

of regions in that subnetwork. Brain regions of different RSNs are given in Tab. S.15, and EEG

frequency bands are given in Tab. S.13.

Some of the most remarkable and statistically significant (see Tab. S1-S12) activation patterns

predicted by the spectral graph theoretical metrics in Fig. 4 are in agreement with the strongest

empirically observed phenomena. It is the frequency dependent normalized links that support the

visual network areas, which are mainly located in the occipital lobe, to be the most active at

alpha frequencies, as it is observed during rest with closed eyes and absence of visual inputs (*43*,

44). Similarly, while activity is dominant in the delta/theta frequencies for the anterior-cognitive

state, the posterior-cognitive state is dominated by alpha (*39*). In contrary, the sensory motor and

auditory networks are more active at gamma (*45*), whilst DMN, which is mainly in the frontal

lobe, is more prominent at beta band (*46*), associated with the idleness of the cognitive and motor

setup.

Numerical confirmation for forward modeling and Linear Stability Analysis (LSA) with

Stuart Landau oscillators. The activation patterns predicted by the spectral graph theoretical

metrics of the normalized connectome are numerically validated with a linear stability analysis

and a forward simulation of brain network model of Stuart-Landau oscillators (*18*, *30*) over the

same 100 human connectomes, Fig. 5. The model is explored at critical, sub-critical and super-

critical regimes, along with noise (see Methods). For consistency, in the simulated results,

powers get signs that are based on the in-/anti- phase arrangements of the simulated data, which

is obtained from calculating the average phase difference between the regions and checking

whether it is larger than π. Similarly, for consistent comparison between the metrics, the signs

from the phase arrangements either from the data, from the connectome, or from the eigenvalues,

are explicitly assigned to the spectral strengths and capacities. Thus, for instance, the spectral

strengths respectively become $S_{\text{data}}^{(f)}$, $S_{\text{conn}}^{(f)}$ or $S_{eig}^{(f)}$.

With the generative model we confirm that these spectral patterns of activity are predicted by the

two graph theoretical metrics on the normalized connectome, i.e. the spectral strength and

capacity. The former depends on the actual phase arrangement of the oscillators. For this besides

the arrangement that maximizes the overall $S^{(f)}$ (that is based only on the normalized

connectome) we also use the signs of the eigenvectors to inform the spectral strength metric

about the in/anti phase organization for the specific model. In this dynamical setting the

amplitude and the sign of the observed activation pattern for the network coupled through the

normalized connectome is guaranteed to be fully captured by the values of the first eigenvector

of the LSA (*19*, *29*). The signs of the different directions of the eigenvector capture the in-/anti-

phase arrangements of the nodes, with the opposite sign regions corresponding to anti-phase

synchronized regions.

In Fig. 5 (A), we show the correlation of the leading eigenvectors with spectral strength,

calculated using both arrangements, for all 100 subjects. The arrangement implied by the signs of

the eigenvectors takes into account the specificity of the model, and hence this case better

predicts the activation of a given node. This means that the network is not always arranged in

such a manner to yield maximal possible synchronization. The patterns are similar across

subjects, and for frequencies larger than 40Hz, there seem to be specific frequencies for different

subjects, for which the arrangement of the signs of the eigenvectors is quite different from those

maximizing $S^{(f)}$. Spectral capacity on the other hand, has generally lower predictability than

$S_{conn}^{(f)}$, but it is never insignificant or lower than the case without normalization. As shown in the

third column of the panel (A), all three spectral metrics exhibit much better correlation than the

null metric, which is constructed assuming particle-like interactions at 0Hz. Thus, only the

weights of the links are considered, giving the in-strength of the nodes $C_i^{(0)} = \sum_j w_{ij}$. Since the

null model cannot yield anti-phase nodes, for more conservative comparison we assign it signs

predicted by the arrangements of the nodes with which it is being compared.

Fig. 5. **Comparison with forward models based on the normalized and the full connectome.**

The power across regions from the forward model is compared with the spectral capacity and

strength, where the latter is calculated either using the signs of the eigenvectors or using

arrangements that maximize its value, and with the null metrics assuming particle-like

interactions. The power is calculated (A) using LSA of the brain network model at criticality,

constructed over the normalized connectome or (B) with numerical simulations performed for

the full connectome, with the brain network model being, sub- (G=17, D=0) and super-critical

(G=9, D=1e-5), and at the criticality with noise (G=9, D=1e-5) and without (G=9, D=0). (B) For

the full simulations, the most prominent phase arrangement of the simulated data is also used to

set the in-/anti-phase arrangements. Correlations of the powers across frequencies (A) for all 100

subjects, as well as the p values, and the difference with the correlations assuming particle-like

interactions, and (B) for one subject. Correlations with p<0.01 are transparent in the first 2

columns of (A), and thinner lines for (B).

In Fig. 5 (B) the predictivity of the spectral metrics is tested by comparing them with the power

obtained by simulating the model built with the same oscillators, Eq. (17), but over the full

connectome containing both weights and delays, Eq. (18). Moreover, this is extended for the sub-

critical and super-critical regimes of the system, along with noise. The comparison shows that

each of the wave interactions metrics has consistently higher predictability than its particle-like

counterpart across all the frequencies. This holds even for noisy systems far from the criticality,

where the correlation is generally larger than 0.5 if the phase arrangements are taken from the

data. It is worth noting that for the full system the spectral capacity seems to be more informative

than the spectral strength, because the dynamics often becomes non-stationary and multi-stable

due to the explicit time-delays (35). Nodes might switch from in- to anti-phase and back, and

different phase arrangements lead to different spectral strengths over time. As a result, each

node's average relative strength within the network might be better captured by its capacity, than

by any given realization of the phase arrangements that has occurred at some point, even if that is

the most dominant one. This is especially pronounced when noise is added to the system, making

the spectral capacity more informative for the relative activity of the nodes almost for all the

frequencies.

__Robustness of the spectral strength.__ The consistency of the spectral strength of the normalized

connectome is tested against different types of uncertainties, which are modeled as rewiring and

noise in the connectome, all scaled to have a comparable impact. Boxplots that show the

statistics of the leading eigenvector v and its absolute value, are shown in Fig. 6 (A) for one

frequency and for two different levels of noise. At lower noise, the means of the regional activity

shown by the eigenvectors of the null model are still following the original values. The variance

of the null models' activity in this case would sufficiently describe the impact of the noise, and

the error that it causes. As the noise increases, the variance of the null model also increases, but

in addition for some regions the mean starts significantly differing from the original values.

These suggests that to better account the impact of the noise, and the network's resilience, we

need to take into account both the standard deviation σ_i of all realizations of v_i^{null}, and the error

of their mean $\bar{v}_i^{null}$ defined as

$$\widetilde{err}_i = \left| v_i - \bar{v}_i^{null} \right|. \tag{6}$$

344   Moreover, the impact of the above error seems to be inversely proportional with the spread of the

345   results for the null model, i.e. smaller the spread, larger the chance for the null model to give an

346   error around its mean. Hence, we arrive to the following definition of the error for each node

$$err_i = \frac{\widetilde{err}_i}{\sigma_i} + \frac{\sigma_i}{2}. \tag{7}$$

The mean of those errors over the whole network is shown in the first column of Fig. 6 (B).

Finally, not all the nodes are equally affected by the noise, and to describe this in the right

column plots we depict the proportion of the nodes that have error larger than the threshold set at

$(v_i + \bar{v})/2$. In this way the deviation at each node is evaluated against both, the local, i.e. its

own, and the global activity.

Results in Fig. 6 (B) show that the robustness against the noise and rewiring decreases with the

frequency, and this is especially evident for the uncertainties of the delays. For frequencies

below ~35 Hz less than half of the nodes have deviations that are considered large for all three

types of uncertainties, with rewiring being more sensitive than the noise of the same impact.

Fig. 6. **Robustness of the spectral strength.** (A) Boxplots showing the statistics of the leading

eigenvector and its absolute value for 1000 realizations of 2 different levels of noise added to the

lengths at $f = 20Hz$ on the connectome of the first subject. Black circles are the values for the

original connectome, which are filled red for the cases when the error terms Eq. (7), exceed the

error threshold. (B) Mean of the errors Eq. (7), over all the regions shown for different noise and

frequencies (left column), as well as the ratio of the nodes whose error exceeds the threshold

(right column).

DISCUSSION

Wave metrics offer a new perspective for characterizing dynamic networks. They integrate the

wave component of information processing, which was missing in network metrics that were

based solely on the connection topology (*2*). The incorporation of signal transmission delays in

the connectome's metrics completes the characterization of the spatiotemporal skeleton, within

which oscillatory brain activity can be amplified by the properties of the medium supporting it,

i.e. it provides a *corpus resonantiae*. We propose that the activation of certain parts of the brain

that are related to different tasks can be explained as being prewired in the anatomy, and we have

demonstrated here that the brain connectome has such properties and allows for selective and

frequency dependent information processing that could support the differentiation of the brain

activity for various processes and frequency bands (*47*, *48*). Notably, the wave perspective

naturally leads to the formation of overlapping, albeit functionally independent, subnetworks,

which cannot be derived within the particle picture. Besides revealing the frequency specific

hubness, as another consequence the wave couplings also support the general reduction of

activity with increasing the frequencies, which is among the most consistent and well-established

features of the brain dynamics (*16*).

The normalization of the connectome that we propose is derived from the expressions for the

phase lags between synchronized phase oscillators. These are the only dynamical feature by

which functional heterogeneity can arise in this set-up (*17*, *49*), due to the structural symmetry

breaking of the network interactions (weights and time-delays), or due to the different natural

frequencies. Both these aspects are present in Eq. 3, but in this study, we focus on the former. In

that respect we show that the in presence of time-delays the weights are effectively scaled (or

normalized) by $\cos \Omega \tau_{ij}$, even though the rest of the network is also needed to fully describe the

phase lags and hence the full dynamics. For the case of 2 oscillators, however, the normalized

weights are the sole determinant for the synchronized interactions.

Importantly, however, the network models that we introduced are not intended to serve as toy

brain models, but to illustrate the meaning of the here derived metrics. We use established

models to expose and justify open and hidden assumptions underlying the traditional and

extended metrics (the latter being normalized for time delays in the wave perspective). Every

metrics relies on an underlying theory/model, even if it is not made explicit. For instance,

functional connectivity relies via covariance on second order metrics (*8*) adapted to linear

models (fitting an ellipsoid to the corresponding data cloud), but this does not preclude its

application to nonlinear network models, in fact in practice this is often the rule. In the same

spirit, we motivate the particle and wave metrics by using rate and Kuramoto oscillator

equations, without the claim that these actually represent brain network models.

It is worth noting that studying the synchronizabillity of the brain networks with time-delays

could complement our framework for the study of spectrally dependent emergent functional

heterogeneity, beyond the forward modelling. A possible approach here would be to use recent

extensions of the Ott and Antonsen ansatz (*34*) that allow studying noisy networks, either

through circular cumulants (*50*), or by using Daido order parameters (*51*). These could be

applied on the spatially decomposed time-delays as a first approximation of the spatio-temporal

structure of the connectome (*35*). Another direction would be to study the stability of the

synchronized states over the network of wave couplings which can be positive and negative (*52*).

During tasks the spectrally dependent activation patterns are also constrained by stimuli that in

case of working memory visual tasks for example, cause gamma synchronization in frontal

regions (*53*), which is missing from our predictions at rest and in absence of stimuli. The offered

framework also do not account for communication across different frequencies, even though it

could be further generalized to other types of *n:m* synchronization (*30*). Having data informed

different intrinsic dynamics is also a feature that would increase the predictive value of any

generative model for the brain dynamics, but this would need to be biased by the data, which is

specific of the brain activity (*54–56*). This is however still to be generalized in a graph

theoretical framework. Similarly, our framework considers the activity at each frequency as fixed

(even though they are often time-dependent (*57–60*) and intermittent (*61*)) and as a separate

process. The latter might lead to omitting certain emergent properties in the cases when the same

process has clear multimodal frequency content as often seen in frequency and scale dependent

metrics from information theory (*62–64*) and time-frequency representation (*65*).

Beyond synchronization, frequency-dependent approaches to information processing offer data

driven insight into causality of network interactions (*66*). Applying transfer entropy (*67*) reveals

that high degree neurons feed cortical computations by receiving more information from high

out-degree neurons (*26*), indicating an asymmetry of information transfer between hubs and low-

degree nodes (*27*). Spectral and multiscale Granger causality (*64*, *68*) are similarly used to

describe the information flow in networks, and they demonstrate the nonlinear effects of the

structural weights, which can be seen through signatures of the law of diminishing marginal

returns (*69*, *70*). It would be interesting to compare the network flow that these metrics reveal for

the case of synchronization, with the spectrally dependent network core as defined by the

normalized connectome.

The predictive value of the normalized connectome for the functional in-strength is further

validated through comparisons with numerical simulations of Stuart Landau model. This model

is often used to describe mesoscopic neuronal activity (*18*, *55*, *71*, *72*) due to the fact that it

encompasses the working points of population rate models (*73*). Correlating the spatial activation

patterns with the predictions from the spectral in-strength and capacity shows that the normalized

metrics captures the dynamics of the model better than using only the weights, besides the

derivations being made on the Kuramoto model that does not contain amplitude dynamics, and

that assumes difference instead of additive coupling. Better performance of the wave coupling is

consistent across the frequencies, and holds for different working points, such as sub and

supercritical, with or without added noise.

The normalization of the connectome unifies its structural weights and time-delays, and

consequently it still inherits the known problems of these two modalities. What the number of

streamlines mean for the actual effective coupling between brain regions regardless of the task is

still lacking a clear response (*8*, *74*). In addition, diffusion tractography is known to under

sample the interhemispheric links (*75*), and the topology of the derived connectomes is still

dependent on the parcellation (*76*). The lack of personalized estimations for the time-delays of

the connectome is still unavoidable problem. A possible way forward here could be using

averaged delays derived from stimulation (*77*), combined with the personalized lengths of the

tracts, even if these datasets are still incomplete. The lack of directionality in the human

connectomes that rely on diffusion tractography is also a confounding factor of those

connectomes (*10*), which has led to introduction of concepts such as effective connectivity (*8*,

74), even though for our framework the derivations are simplified by the symmetry in the

coupling strengths.

Finally, we have only introduced two metrics for the node activity based on the normalized

weights, whereas other graph theoretical metrics for the wave interactions could be more

informative. However, even for the spectral capacity and strength, it is still not clear which one

would have higher predictive value in describing the activation patterns of the brain at rest.

Moreover, their systematic and more thorough comparison against empirical data could provide

with the answers about which brain regions operate closer to the limits of their capacity.

Similarly, the phase arrangements of the brain do not necessarily need to be such that maximizes

the overall information transfer across the whole brain, but maybe instead it gives priority at

certain areas, at expense of the others.

Transient synchronization between brain areas coupled through delayed interactions has been

proposed as mechanism that could support flexible information routing (*78*), which plays a

fundamental role for the cognition. Similarly, the implementation of selective communication

dependent on coherence has been generalized to Communication Through Coherence (*20*), as

one of the possible mechanisms of communication between neuronal networks (*79*), but with no

explanations of the origin of the selection of frequency bands and the spatial organization of the

involved neuronal populations. The normalization of the connectome completes this view

mechanistically, hypothesizing that the spatio-temporal brain structure itself could be at least

partly responsible for the observed patterns. Through the notion of wave-type communication,

the offered normalization lends legitimacy to concepts of synchronization and coherence in

network science and opens up a many new applications ways beyond brain sciences.

MODEL & METHODS

Particle versus wave representation for network dynamics

477   The particle-wave dichotomy appears in networks dynamics due to the finite transmission

478   velocities for the signals between spatially distributed network nodes. Within the particle view,

479   the general network dynamics is governed by

$$\dot{r}_i = g_1(r_i) + \sum_j w_{ij} g_2(r_j(t - \tau_{ij})). \tag{8}$$

Here r_i is any scalar variable such as the firing rate, w_{ij} are the weights and τ_{ij} are the time-

delays of the links, and g_1 and g_2 are linear functions for the intrinsic dynamics and coupling

respectively. This type of particle-like interactions is frequency-independent and here the

information and the activity always flow along the links with the strongest coupling weight.

On the other hand, networks of oscillators are often used to conceptualize and to study dynamical

systems for which the local activity is multidimensional and nonlinear (*30*). They have been

conceptualized to be responsible for the communication in the brain through coherence (*20*) and

synchronization (*31, 78, 80*). In this case the dynamics of the oscillatory network nodes are

governed by Eq. (1). When delays are negligible compared to the time-scale of the system, the

interactions are still governed only by the weights, as in the particle case. However, when the

time-delays are comparable to the time-scale of the intrinsic oscillations, they need to be

included in the analysis of network dynamics. As a consequence, this system can only be solved

numerically, as there is no alternative graph theoretical approach that integrates the topology and

the impact of the time-delays on the oscillations at a given frequency. Hence the systematic

effect of the time-delays on the emergent dynamics is concealed and even the full representation

of network interactions in Eq. (1), with separated w_{ij} and τ_{ij} cannot adequately identify the

skeleton of the wave synergy.

For weak couplings, dynamics of the system in Eq. (1) is captured by phase models, which are

often used to study the interactions in the brain at different levels (*58, 61*). The simplest and

hence most elaborate is the Kuramoto model (*32*). It is widely utilized for describing emergent

phenomena in complex systems (*81*), such as the brain (*82–84*), with a structure often

represented via complex networks (*85*). For synchronization at frequency Ω, the model reads

(*33*)

504
$$\dot{\theta}_i = \omega_i - \sum_j w_{ij} \sin\left(\theta_i - \theta_j\left(t - \tau_{ij}\right)\right) = \omega_i - \sum_j w_{ij} \sin\left(\theta_i - \theta_j + \Omega\tau_{ij}\right), \tag{9}$$

505   where θ_i denotes the phase of the *i*-th node oscillating with a natural frequency ω_i, the

506   dependence on time is implicit, and due to the synchronization $\theta_j\left(t - \tau_{ij}\right) = \theta_j(t) + \Omega\tau_{ij}$.

507   From here, the phase difference between each pair of Kuramoto oscillators is given as

508   $\dot{\theta}_{ij=}\dot{\theta}_i - \dot{\theta}_j = \Omega - \Omega = \omega_i - \omega_j - \sum_k w_{ik}\sin(\theta_i - \theta_k + \Omega\tau_{ik}) + \sum_k w_{jk}\sin\left(\theta_j - \theta_k + \Omega\tau_{jk}\right) =$

509
$$= 2\Delta\omega_{ij} - \sum_k(w_{ik}\sin(\theta_i - \theta_k + \Omega\tau_{ik}) + w_{jk}\sin(\theta_j - \theta_k + \Omega\tau_{jk})), \tag{10}$$

510   where $\Delta\omega_{ij} = (\omega_i - \omega_j)/2$. Next we introduce $\Delta\theta_{ij} = \theta_i - \theta_j$ and we separate the sum, such

511   that

512
$$0 = 2\Delta\omega_{ij} - w_{ij}\sin\left(\Delta\theta_{ij} + \Omega\tau_{ij}\right) + w_{ji}\sin\left(\Delta\theta_{ji} + \Omega\tau_{ji}\right) -$$

513
$$- \sum_{k \neq i,j}\left(w_{ik}\sin(\Delta\theta_{ik} + \Omega\tau_{ik}) - w_{jk}\sin\left(\Delta\theta_{jk} + \Omega\tau_{jk}\right)\right) =$$

514
$$= 2\Delta\omega_{ij} - w_{ij}\sin\left(\Delta\theta_{ij} + \Omega\tau_{ij}\right) + w_{ji}\sin\left(\Delta\theta_{ji} + \Omega\tau_{ji}\right) - 2I_{ij}, \tag{11}$$

515   where

516
$$I_{ij} = \frac{1}{2}\sum_{k \neq i,j}\left(w_{ik}\sin(\Delta\theta_{ik} + \Omega\tau_{ik}) - w_{jk}\sin\left(\Delta\theta_{jk} + \Omega\tau_{jk}\right)\right). \tag{12}$$

517   Note that I_{ij} contains all the other links towards the nodes *i* and *j* apart their direct link.

518   For symmetric coupling, $w_{ij} = w_{ji}$ and $\tau_{ij} = \tau_{ji}$, while $\Delta\theta_{ij} = \Delta\theta_{ij}$ by definition, hence Eq. (11)

519   becomes

$$0 = 2\Delta\omega_{ij} - w_{ij}\left[\sin\left(\Delta\theta_{ij} + \Omega\tau_{ij}\right) - \sin\left(-\Delta\theta_{ij} + \Omega\tau_{ij}\right)\right] - 2I_{ij} =$$

$$= 2\Delta\omega_{ij} - 2\,w_{ij}\sin\Delta\theta_{ij}\cos\Omega\tau_{ij} - 2I_{ij} \tag{13}$$

522   Finally, for the phase difference between the oscillators we get

$$\Delta\theta_{ij} = \sin^{-1}\frac{\Delta\omega_{ij} - I_{ij}}{w_{ij}\cos\Omega\tau_{ij}} = \sin^{-1}\frac{\Delta\omega_{ij} - I_{ij}}{w_{ij}^{(f)}}.$$

where $w_{ij}^{(f)} = w_{ij}\cos\Omega\,\tau_{ij}$ is the normalized weight containing the impact of the direct link

between the pair of oscillators. For particle-like communication in the limit $\Omega\tau_{ij} \to 0$ the cosine

term and I_{ij} vanish making the coupling weights the only factor shaping the network dynamics,

consistent with the case of no delays. Numerical simulations validate the analytical results for

$\Delta\theta_{ij}$ by showing that the difference between the analytical and numerical values is small and

decreases for smaller integration step, and increase with added noise (Fig. S1).

In the case of asymmetric weights, Eq. (13) would also include a term that accounts for the

asymmetry in the interactions $w_{ij} - w_{ji}$, and the exact derivation of the expression in Eq. (3) will

not be possible anymore. However, if the asymmetry is relatively small, as it is the case for the

mice tracer connectome (37) for example, then the same expression would approximately hold

up to a leading term. As for the human connectomes, they almost exclusively rely on diffusion

tractography imaging, which produces links with symmetric weights.

Limits of the approximation

If we assume normalized connectome, the governing equation of the phase dynamics becomes

$$537 \qquad \dot{\theta}_i = \omega_i - \sum_j w_{ij} \cos \Omega \, \tau_{ij} \sin(\theta_i - \theta_j), \qquad (14)$$

538 Following the same steps as above, the phase difference between a pair of oscillators reads

$$539 \qquad \dot{\theta}_i - \dot{\theta}_j = \Omega - \Omega = 2\Delta\omega_{ij} - \sum_k \left(w_{ik} \cos \Omega \, \tau_{ik} \sin \Delta\theta_{\text{ik}} - w_{jk} \cos \Omega \, \tau_{jk} \sin \Delta\theta_{\text{jk}} \right) =$$

$$540 \qquad 2\Delta\omega_{ij} - 2w_{ij} \cos \Omega \, \tau_{ij} \sin \Delta\theta_{\text{ij}} - \sum_{k \neq i,j} \left(w_{ik} \cos \Omega \, \tau_{ik} \sin \Delta\theta_{\text{ik}} - w_{jk} \cos \Omega \, \tau_{jk} \sin \Delta\theta_{\text{jk}} \right) =$$

$$541 \qquad 2\Delta\omega_{ij} - 2w_{ij} \cos \Omega \, \tau_{ij} \sin \Delta\theta_{\text{ij}} - 2J_{ij}. \qquad (15)$$

542 As for the difference between the full model and the approximation using the normalized

543 weights, we get

$$544 \qquad I_{ij} - J_{ij} = \frac{1}{2}\sum_{k \neq i,j} \left(w_{ik}\sin(\Delta\theta_{\text{ik}} + \Omega\tau_{\text{ik}}) - w_{jk}\sin(\Delta\theta_{\text{jk}} + \Omega\tau_{\text{jk}}) - w_{ik} \cos \Omega \, \tau_{ik} \sin \Delta\theta_{\text{ik}} + $$

$$545 \qquad + w_{jk} \cos \Omega \, \tau_{jk} \, \sin \Delta\theta_{\text{jk}} \right) =$$

$$546 \qquad = \frac{1}{2}\sum_{k \neq i,j} \left(w_{ik} \sin \Omega \, \tau_{ik} \cos \Delta\theta_{\text{ik}} - w_{jk} \sin \Omega \, \tau_{jk} \cos \Delta\theta_{\text{jk}} \right) \qquad (16)$$

547 For many symmetric networks, such as the delay-imposed clusters with identical weights (*35*)

548 this difference vanishes.

549 Forward modeling and Linear Stability Analysis (LSA)

The normalization of the connectome also allows straightforward application of LSA for the

dynamics close to a critical state for a given forward model of the oscillatory activity. To build

our brain network model (*17, 18, 29, 55*) over the same 100 human connectomes, we choose

Landau Stuart oscillators, as a canonical model for oscillations near Andronov - Hopf bifurcation

(*32*), which is also the working point of many neuronal mass models (*86, 87*). The local

dynamics from Eq. (1) is then given as

$$556 \qquad F(X_i) = X_i(r + j\omega - |X_i|^2), \qquad (17)$$

where X_i is a complex number, and the coupling is linear additive with

$$w_{ij} h\left(X_j\left(t - \tau_{ij}\right)\right) = w_{ij}^{(f)} X_j. \tag{18}$$

All the parameters, including the frequency ω and the distance from the local bifurcation r are

identical for every oscillator. The stability of this system can be analyzed by following the

evolution of infinitesimal perturbations in the tangent space, whose dynamics is ruled by the

linearization of Eq. 1 with Eqs. (17 − 18). If the eigenvalues of the Jacobian matrix all have real

parts less than zero, then the steady state is stable. For each frequency for which the system is

analyzed, we set the parameter r to keep the system at the edge of criticality. This is done in an

iterative procedure, where r is decreased until the system is just above the criticality. In this case

there are two identical positive eigenvalues, which contain the same eigenvectors of opposite

signs, due to the way how the two dimensions of the local model are coupled.

In addition to the LSA, we carried a numerical simulation over the full connectome of the same

model, with the coupling term reading

$$w_{ij} h\left(X_j\left(t - \tau_{ij}\right)\right) = G w_{ij} X_j\left(t - \tau_{ij}\right), \tag{19}$$

where G is a global coupling parameter. Numerical integration is performed using stochastic

Heun integration scheme implemented in Matlab, with an integration time-step that is inversely

proportional of the natural frequencies and the global coupling, but it is never larger than 0.005s.

In some of the analyzed scenarios the dynamics also contains white additive Gaussian, with

noise intensity D. The critical value of r is different for each connectome and each frequency,

and sub- and super- criticality are set such that r is at 0.2 below or above criticality, respectively.

Setting the weights for the coupling

Human connectomes were derived from the first release of Diffusion Tractography Imaging of

100 healthy subjects, which are publicly available as part of the Human Connectome project

(*36*). The subjects were scanned on a customized 3T scanner at Washington University and the

structural connectivity was constructed using a publicly available pipeline (*88*) that applies

spherical deconvolution method to a probabilistic streamlines tracking algorithm. The obtained

connectome consists of few million tracts spatially averaged to connect 68 cortical regions

defined according to Desikan-Kiliany atlas (*89*). For each link, weights are numbers of

individual tracts between the regions of that link, and lengths are their averages.

The mouse connectome in Fig. 2 (A) was obtained from the Allen Institute Mouse Brain

Connectivity data (*37*), which is integrated in The Virtual Mouse Brain (*90*), and which was

already used to model mice brain dynamics (*10*, *84*).

The spectral graph theoretical results (Fig. 2-4) and the LSA analysis (Fig. 5 (A) and 6) use the

natural logarithms of number of streamlines as weights of the couplings. This is done for better

illustration of the findings in order to decrease the heterogeneity between the different nodes. For

the full forward model (Fig. 5 (B)), we use the number of streamlines, as it has been more

common in the numerical simulations (*17*, *18*, *29*, *55*).

Phase arrangement of the nodes that maximizes the spectral strength

When the in-/anti-phase arrangement of the nodes is such that $S^{(f)}$ is maximized, this is referred

to as $S_{conn}^{(f)}$, since the arrangement takes into account only the connectome, but not the model. If

not specified otherwise in the text, then by the spectral strength $S^{(f)}$ we refer to $S_{conn}^{(f)}$. The

clusters of in-/anti-phase nodes can be of any size, but the assigning of the clusters is not

important (e.g. $i \in Q_i \equiv \{1,2,\dots N\} \backslash i \in Q_i$, and similarly swapping all the red and blue colors in

Fig. 1 does not change the results). As a consequence the total number of combinations is

$\sum_{k=1}^{N/2} \binom{N}{k} \approx \frac{N!}{2}$ and a brute force procedure cannot be used to identify the clustering that yields the

maximum $S_{conn}^{(f)}$. Hence, we developed a heuristic procedure that seems to be able to identify the

clusters of maximum spectral strength. In short, it starts with random initial arrangements and

then uses three types of cost functions to identify the node that needs to change the cluster. These

are:

-   The node with smallest $S^{(f)}$ changes the cluster.

-   The node with a smaller $S^{(f)}$ from the 2 nodes of the link that has the smallest (most

608   negative) internal (from the links within each cluster) $w_{ij}^{(f)}$.

609   -   The node with a smaller $S^{(f)}$ from the 2 nodes of the link that has the smallest (most

610   negative) external (from the links between the clusters) $w_{ij}^{(f)}$.

For better sampling of the different arrangements, a randomness is also systematically added in

some of the loops, so that the cost function chooses a suboptimal node or link among some

proportion of them that were identified to have the largest cost.

The procedure is run over multiple initializations, until it recovers a cluster that has been already

identified. At the end of the runs, the largest spectral capacity that was obtained is set to be

$S_{conn}^{(f)}$. In particular, we start from 10 random initialization for the phases and for each of them

the three procedures are repeated one after another until an already visited configuration is

achieved, after which the loop is repeated 10 more times with added randomness. Every loop is

stopped after arriving to an already visited configuration, and all the visited configurations are

then classified according to the highest spectral strength. We have verified that the convergence

to the selected configuration occurs even after 10-100 times less samples than we generate with

our procedure.

Projecting the spectral strengths and capacities into RSNs and brain lobes

To better illustrate the patterns of spectral activity we project the spectral strength and capacity

in each brain lobe and RSN, Figs. 2-4. For each subject, after calculating them for a given

frequency (from 0 to 80 Hz with a step of 1Hz), we calculate the average (per number of regions

in the given subnetwork) activity of a given subnetwork (RSN or lobe). In addition, for the

results in Fig. 3 and 4 the values of each subnetwork are normalized at each frequency, to focus

on the relative activation patterns. The activity of different frequency bands contains the

averaged activity of the frequencies within that band (see Tab. S.13). Mean and the standard

deviation over the subjects are calculated for each frequency for the activity of the brain lobes.

Note that the averaging over subjects in Fig. 4 (B) is done over the spectral strengths, which

explicitly take into account the in-/anti-phase arrangements by assigning them opposite signs. In

particular, the regions which are in-phase with regions that have strongest spectral strength over

the subjects are set to be always of a positive sign (blue color), as compared with the others

which are negative (red). Since during the averaging some nodes can be of opposite phase

arrangements over the subjects, the absolute value of the observed activity here, still can be still

different from the one shown in Fig. 4 (A).

Robustness of the spectral strength

To test the robustness of the results, we modified connectomes by either introducing noise or

rewiring. We use 3 types of modified connectomes:

-    Noise is added to the lengths of the links, with the strength proportional to the mean

length of over the links.

-    Noise is added to the number of streamlines that are used for calculating the weights. The

level of the noise is proportional to the mean number of streamlines of over the links.

-    A proportion of the links equivalent to the level of noise is rewired while preserving the

distribution of in-strengths (*91*).

For each of this and for each frequency we perform 1000 realizations of the null networks, and

then we compute the LSA for each frequency, to obtain the patterns of activity.

<u>Acknowledgment</u>

This research was supported by the European Union's Horizon 2020 research and innovation

program under grant agreement No. 785907 (SGA2) and 945539 (SGA3) Human Brain Project,

and by grant agreement No. 826421 Virtual Brain Cloud.

<u>Data availability statement</u>

All the data, the codes, and all the relevant information to reproduce the results in the paper will

be made available in a public repository.

References:

1.   S. Maslov, K. Sneppen, Specificity and stability in topology of protein networks. *Science*

(80-.). **296**, 910–913 (2002).

2.   D. S. Bassett, O. Sporns, Network neuroscience. *Nat. Neurosci.* **20**, 353–364 (2017).

3.   A. Mislove, M. Marcon, K. P. Gummadi, P. Druschel, B. Bhattacharjee, in *Proceedings of*

the ACM SIGCOMM Internet Measurement Conference, IMC (2007).

4.   E. Bullmore, O. Sporns, The economy of brain network organization. *Nat. Rev. Neurosci.*

13, 336–349 (2012).

5.   R. Albert, A. L. Barabasi, Statistical mechanics of complex networks. *Rev. Mod. Phys.* **74**,

47–97 (2002).

6.   H. Markram, E. Muller, S. Ramaswamy, M. W. Reimann, M. Abdellah, C. A. Sanchez, A.

Ailamaki, L. Alonso-Nanclares, N. Antille, S. Arsever, G. A. A. Kahou, T. K. Berger, A.

Bilgili, N. Buncic, A. Chalimourda, G. Chindemi, J. D. Courcol, F. Delalondre, V.

Delattre, S. Druckmann, R. Dumusc, J. Dynes, S. Eilemann, E. Gal, M. E. Gevaert, J. P.

Ghobril, A. Gidon, J. W. Graham, A. Gupta, V. Haenel, E. Hay, T. Heinis, J. B.

Hernando, M. Hines, L. Kanari, D. Keller, J. Kenyon, G. Khazen, Y. Kim, J. G. King, Z.

Kisvarday, P. Kumbhar, S. Lasserre, J. V. Le Bé, B. R. C. Magalhães, A. Merchán-Pérez,

J. Meystre, B. R. Morrice, J. Muller, A. Muñoz-Céspedes, S. Muralidhar, K. Muthurasa,

D. Nachbaur, T. H. Newton, M. Nolte, A. Ovcharenko, J. Palacios, L. Pastor, R. Perin, R.

Ranjan, I. Riachi, J. R. Rodríguez, J. L. Riquelme, C. Rössert, K. Sfyrakis, Y. Shi, J. C.

Shillcock, G. Silberberg, R. Silva, F. Tauheed, M. Telefont, M. Toledo-Rodriguez, T.

Tränkler, W. Van Geit, J. V. Díaz, R. Walker, Y. Wang, S. M. Zaninetta, J. Defelipe, S. L.

Hill, I. Segev, F. Schürmann, Reconstruction and Simulation of Neocortical

Microcircuitry. *Cell*. **163**, 456–492 (2015).

7.   O. Sporns, G. Tononi, R. Kötter, The human connectome: A structural description of the

human brain. *PLoS Comput. Biol.* **1**, 0245–0251 (2005).

8.   K. J. Friston, Functional and effective connectivity: a review. *Brain Connect.* **1**, 13–36
(2011).

9.   P. Sanz-Leon, S. A. Knock, A. Spiegler, V. K. Jirsa, Mathematical framework for large-
scale brain network modeling in The Virtual Brain. *Neuroimage*. **111**, 385–430 (2015).

10.   F. Melozzi, E. Bergmann, J. A. Harris, I. Kahn, V. Jirsa, C. Bernard, Individual structural
features constrain the mouse functional connectome. *PNAS*. **116**, 26961–26969 (2019).

11.   A. Brovelli, M. Ding, A. Ledberg, Y. Chen, R. Nakamura, S. L. Bressler, Beta oscillations
in a large-scale sensorimotor cortical network: Directional influences revealed by Granger
causality. *PNAS*. **101**, 9849–9854 (2004).

12.   T. Proix, F. Bartolomei, M. Guye, V. K. Jirsa, Individual brain structure and modelling
predict seizure propagation. *Brain*. **140**, 641–654 (2017).

13.   G. Buzsáki, *Rhythms of the Brain* (2009).

14.   S. Haegens, V. Nácher, R. Luna, R. Romo, O. Jensen, α-Oscillations in the monkey
sensorimotor network influence discrimination performance by rhythmical inhibition of
neuronal spiking. *PNAS*. **108**, 19377–82 (2011).

15.   P. L. Nunez, R. Srinivasan, *Electric fields of the brain : the neurophysics of EEG*
(Oxford University Press, 2006).

16.   G. Buzsáki, D. Andreas, A. Draguhn, Neuronal Oscillations in Cortical Networks. *Science*
(80-.). **304**, 1926 (2004).

17.   S. Petkoski, J. M. Palva, V. K. Jirsa, Phase-lags in large scale brain synchronization:
Methodological considerations and in-silico analysis. *PLoS Comput. Biol.* **14**, 1–30
(2018).

18.   S. Petkoski, V. K. Jirsa, Transmission time delays organize the brain network

709   synchronization. *Philos. Trans. R. Soc. A*. **377**, 20180132 (2019).

710   19.   A. Ghosh, Y. Rho, A. R. McIntosh, R. Kötter, V. K. Jirsa, Noise during rest enables the

711   exploration of the brain's dynamic repertoire. *PLoS Comput. Biol.* **4**, e1000196 (2008).

712   20.   P. Fries, Rhythms for Cognition: Communication through Coherence. *Neuron*. **88**, 220–

713   235 (2015).

714   21.   D. S. Bassett, P. Zurn, J. I. Gold, On the nature and use of models in network

715   neuroscience. *Nat. Rev. Neurosci.* (2018), pp. 1–13.

716   22.   M. Rubinov, O. Sporns, Complex network measures of brain connectivity: Uses and

717   interpretations. *Neuroimage*. **52**, 1059–1069 (2010).

718   23.   L. L. Gollo, A. Zalesky, R. Matthew Hutchison, M. Van Den Heuvel, M. Breakspear,

719   Dwelling quietly in the rich club: Brain network determinants of slow cortical

720   fluctuations. *Philos. Trans. R. Soc. B*. **370**, 20140165–20140165 (2015).

721   24.   J. A. Roberts, A. Perry, A. R. Lord, G. Roberts, P. B. Mitchell, R. E. Smith, F. Calamante,

722   M. Breakspear, The contribution of geometry to the human connectome. *Neuroimage*.

723   **124**, 379–393 (2016).

724   25.   E. T. Bullmore, O. Sporns, Complex brain networks: graph theoretical analysis of

725   structural and functional systems. *Nat. Rev. Neurosci.* **10**, 186–98 (2009).

726   26.   N. M. Timme, S. Ito, M. Myroshnychenko, S. Nigam, M. Shimono, F. C. Yeh, P.

727   Hottowy, A. M. Litke, J. M. Beggs, High-Degree Neurons Feed Cortical Computations.

728   *PLoS Comput. Biol.* **12**, 1–31 (2016).

27.   L. Novelli, F. M. Atay, J. Jost, J. T. Lizier, Deriving pairwise transfer entropy

fromnetwork structure and motifs. *Proc. R. Soc. A Math. Phys. Eng. Sci.* **476** (2020),

doi:10.1098/rspa.2019.0779.

28.   A. Spiegler, E. C. A. Hansen, C. Bernard, A. R. McIntosh, V. K. Jirsa, *eNeuro*, in press,

doi:10.1523/ENEURO.0068-16.2016.

29.   S. Olmi, S. Petkoski, M. Guye, F. Bartolomei, V. Jirsa, Controlling seizure propagation in

large-scale brain networks. *PLoS Comput. Biol.* **15**, e1006805 (2019).

30.   A. Pikovsky, M. Rosenblum, J. Kurths, *Synchronization : a universal concept in*

nonlinear sciences (Cambridge University Press, 2001).

31.   S. Palva, J. M. Palva, Discovering oscillatory interaction networks with M/EEG:

Challenges and breakthroughs. *Trends Cogn. Sci.* **16**, 219–229 (2012).

32.   Y. Kuramoto, *Chemical Oscillations, Waves, and Turbulence* (Springer Berlin Heidelberg,

Heidelberg, 1984; http://link.springer.com/10.1007/978-3-642-69689-3), vol. 19 of

Springer Series in Synergetics.

33.   E. Izhikevich, Phase models with explicit time delays. *Phys. Rev. E.* **58**, 905–908 (1998).

34.   E. Ott, T. M. Antonsen, Low dimensional behavior of large systems of globally coupled

oscillators. *Chaos.* **18**, 037113 (2008).

35.   S. Petkoski, A. Spiegler, T. Proix, P. Aram, J.-J. J. Temprado, V. K. Jirsa, Heterogeneity

of time delays determines synchronization of coupled oscillators. *Phys. Rev. E.* **94**, 1–7

(2016).

36.   D. C. Van Essen, S. M. Smith, D. M. Barch, T. E. J. Behrens, E. Yacoub, K. Ugurbil, The

WU-Minn Human Connectome Project: An overview. *Neuroimage.* **80**, 62–79 (2013).

37.   S. W. Oh, J. A. Harris, L. Ng, B. Winslow, N. Cain, S. Mihalas, Q. Wang, C. Lau, L.

Kuan, A. M. Henry, M. T. Mortrud, B. Ouellette, T. N. Nguyen, S. A. Sorensen, C. R.

Slaughterbeck, W. Wakeman, Y. Li, D. Feng, A. Ho, E. Nicholas, K. E. Hirokawa, P.

Bohn, K. M. Joines, H. Peng, M. J. Hawrylycz, J. W. Phillips, J. G. Hohmann, P.

Wohnoutka, C. R. Gerfen, C. Koch, A. Bernard, C. Dang, A. R. Jones, H. Zeng, A

mesoscale connectome of the mouse brain. *Nature*. **508**, 207–214 (2014).

38.   J. S. Damoiseaux, S. A. R. B. Rombouts, F. Barkhof, P. Scheltens, C. J. Stam, S. M.

Smith, C. F. Beckmann, Consistent resting-state networks across healthy subjects. *PNAS*.

103, 13848–13853 (2006).

39.   D. Vidaurre, L. T. Hunt, A. J. Quinn, B. A. E. Hunt, M. J. Brookes, A. C. Nobre, M. W.

Woolrich, Spontaneous cortical activity transiently organises into frequency specific

phase-coupling networks. *Nat. Commun.* **9** (2018), doi:10.1038/s41467-018-05316-z.

40.   F. De Pasquale, S. Della Penna, A. Z. Snyder, C. Lewis, D. Mantini, L. Marzetti, P.

Belardinelli, L. Ciancetta, V. Pizzella, G. L. Romani, M. Corbetta, Temporal dynamics of

spontaneous MEG activity in brain networks. *PNAS*. **107**, 6040–6045 (2010).

41.   F. de Pasquale, M. Corbetta, V. Betti, S. Della Penna, Cortical cores in network dynamics.

Neuroimage, 1–13 (2017).

42.   A. Hillebrand, P. Tewarie, E. van Dellen, M. Yu, E. W. S. Carbo, L. Douw, A. A. Gouw,

E. C. W. van Straaten, C. J. Stam, Direction of information flow in large-scale resting-

state networks is frequency-dependent. *PNAS*. **113**, 3867–3872 (2016).

43.   D. Mantini, M. G. Perrucci, D. Gratta, R. C., G.l, M. Corbetta, C. Del Gratta, G. L.

Romani, C. Del Gratta, G. L. Romani, M. Corbetta, Electrophysiological signatures of

resting state networks in the human brain. *PNAS*. **104**, 13170–13175 (2007).

44.   J. Samogin, M. Marino, C. Porcaro, N. Wenderoth, P. Dupont, S. P. Swinnen, D. Mantini,

Frequency-dependent functional connectivity in resting state networks. *Hum. Brain Mapp.*

41, 5187–5198 (2020).

45.   D. Cheyne, P. Ferrari, MEG studies of motor cortex gamma oscillations: Evidence for a

gamma "fingerprint" in the brain? *Front. Hum. Neurosci.* **7**, 1–7 (2013).

46.   H. Laufs, K. Krakow, P. Sterzer, E. Eger, A. Beyerle, A. Salek-Haddadi, A. Kleinschmidt,

Electroencephalographic signatures of attentional and cognitive default modes in

spontaneous brain activity fluctuations at rest. *PNAS*. **100**, 11053–11058 (2003).

47.   J. F. Mejias, J. D. Murray, H. Kennedy, X. J. Wang, Feedforward and feedback

frequency-dependent interactions in a large-scale laminar network of the primate cortex.

Sci. Adv. **2**, e1601335 (2016).

48.   J. Liu, H. J. Lee, A. J. Weitz, Z. Fang, P. Lin, M. K. Choy, R. Fisher, V. Pinskiy, A.

Tolpygo, P. Mitra, N. Schiff, J. H. Lee, Frequency-selective control of cortical and

subcortical networks by central thalamus. *Elife*. **4**, e09215 (2015).

49.   J. Y. Moon, U. C. Lee, S. Blain-Moraes, G. A. Mashour, General relationship of global

topology, local dynamics, and directionality in large-scale brain networks. *PLoS Comput.*

Biol. **11**, 1–30 (2015).

50.   I. V. Tyulkina, D. S. Goldobin, L. S. Klimenko, A. Pikovsky, Dynamics of Noisy

Oscillator Populations beyond the Ott-Antonsen Ansatz. *Phys. Rev. Lett.* **120**, 264101

(2018).

51.   K. M. Hannay, D. B. Forger, V. Booth, Macroscopic models for networks of coupled

biological oscillators. *Sci. Adv.* **4**, 1–10 (2018).

52.   D. Iatsenko, S. Petkoski, P. V. E. E. McClintock, A. Stefanovska, Stationary and traveling

wave states of the Kuramoto model with an arbitrary distribution of frequencies and

coupling strengths. *Phys. Rev. Lett.* **110**, 1–5 (2013).

53.   F. Roux, M. Wibral, H. M. Mohr, W. Singer, P. J. Uhlhaas, Gamma-band activity in

human prefrontal cortex codes for the number of relevant items maintained in working

memory. *J. Neurosci.* **32**, 12411–12420 (2012).

54.   P. Wang, R. Kong, X. Kong, R. Liégeois, C. Orban, G. Deco, M. P. van den Heuvel, B. T.

Thomas Yeo, Inversion of a large-scale circuit model reveals a cortical hierarchy in the

dynamic resting human brain. *Sci. Adv.* **5**, eaat7854 (2019).

55.   J. Courtiol, M. Guye, F. Bartolomei, S. Petkoski, V. K. Jirsa, Dynamical mechanisms of

interictal resting-state functional connectivity in epilepsy. *J. Neurosci.* **40**, 5572–5588

(2020).

56.   M. Schirner, A. R. McIntosh, V. Jirsa, G. Deco, P. Ritter, Inferring multi-scale neural

mechanisms with brain network modelling. *Elife*. **7**, 1–30 (2018).

57.   T. Stankovski, A. Duggento, P. V. E. McClintock, A. Stefanovska, Inference of time-

evolving coupled dynamical systems in the presence of noise. *Phys. Rev. Lett.* **109**, 1–5

(2012).

58.   T. Stankovski, S. Petkoski, A. F. Smith, P. V. E. E. McClintock, A. Stefanovska, J.

Raeder, A. F. Smith, P. V. E. E. McClintock, A. Stefanovska, Alterations in the coupling

functions between cortical and cardio-respiratory oscillations due to anaesthesia with

propofol and sevoflurane. *Philos. Trans. R. Soc. A*. **374**, 20150186 (2016).

59.   P. Clemson, G. Lancaster, A. Stefanovska, Reconstructing Time-Dependent Dynamics.

818   *Proc. IEEE*. **104**, 223–241 (2016).

819   60.   S. Petkoski, A. Stefanovska, Kuramoto model with time-varying parameters. *Phys. Rev. E*.

820   **86**, 1–7 (2012).

821   61.   L. W. Sheppard, A. C. Hale, S. Petkoski, P. V. E. McClintock, A. Stefanovska,

822   Characterizing an ensemble of interacting oscillators: The mean-field variability index.

823   *Phys. Rev. E*. **87**, 1–11 (2013).

824   62.   J. Courtiol, D. Perdikis, S. Petkoski, V. Müller, R. Huys, R. Sleimen-Malkoun, V. K.

825   Jirsa, The multiscale entropy: Guidelines for use and interpretation in brain signal

826   analysis. *J. Neurosci. Methods*. **273**, 175–190 (2016).

827   63.   D. Marinazzo, M. Pellicoro, S. Stramaglia, Kernel method for nonlinear Granger

828   causality. *Phys. Rev. Lett.* **100**, 1–4 (2008).

829   64.   M. J. Kaminski, K. J. Blinowska, A new method of the description of the information flow

830   in the brain structures. *Biol. Cybern.* **65**, 203–210 (1991).

831   65.   D. Iatsenko, P. V. E. McClintock, A. Stefanovska, Linear and synchrosqueezed time-

832   frequency representations revisited: Overview, standards of use, resolution,

833   reconstruction, concentration, and algorithms. *Digit. Signal Process. A Rev. J.* **42**, 1–26

834   (2015).

835   66.   A. T. Reid, D. B. Headley, R. D. Mill, R. Sanchez-Romero, L. Q. Uddin, D. Marinazzo,

836   D. J. Lurie, P. A. Valdés-Sosa, S. J. Hanson, B. B. Biswal, V. Calhoun, R. A. Poldrack,

837   M. W. Cole, Advancing functional connectivity research from association to causation.

838   *Nat. Neurosci.* **22**, 1751–1760 (2019).

67. T. Schreiber, Measuring information transfer. *Phys. Rev. Lett.* **85**, 461–464 (2000).

68. L. Faes, G. Nollo, S. Stramaglia, D. Marinazzo, Multiscale Granger causality. *Phys. Rev.*
E. **96**, 1–7 (2017).

69. D. Marinazzo, G. Wu, M. Pellicoro, L. Angelini, S. Stramaglia, Information Flow in
Networks and the Law of Diminishing Marginal Returns: Evidence from Modeling and
Human Electroencephalographic Recordings. *PLoS One*. **7**, 1–9 (2012).

70. D. Marinazzo, M. Pellicoro, G. Wu, L. Angelini, J. M. Cortés, S. Stramaglia, Information
transfer and criticality in the ising model on the human connectome. *PLoS One*. **9**, 1–7
(2014).

71. G. Deco, J. Cabral, M. W. Woolrich, A. B. A. Stevner, T. J. van Hartevelt, M. L.
Kringelbach, Single or multiple frequency generators in on-going brain activity: A
mechanistic whole-brain model of empirical MEG data. *Neuroimage*. **152**, 538–550
(2017).

72. G. Deco, M. L. Kringelbach, Turbulent-like Dynamics in the Human Brain. *Cell Rep.* **33**,
108471 (2020).

73. G. Deco, V. Jirs, A. R. McIntosh, O. Sporns, R. Kötter, Key role of coupling, delay, and
noise in resting brain fluctuations. *PNAS*. **106**, 10302–10307 (2009).

74. M. Gilson, N. E. Kouvaris, G. Deco, J. F. Mangin, C. Poupon, S. Lefranc, D. Rivière, G.
Zamora-López, Network analysis of whole-brain fMRI dynamics: A new framework
based on dynamic communicability. *Neuroimage*. **201**, 25–27 (2019).

75. M. Schirner, S. Rothmeier, V. K. Jirsa, A. R. McIntosh, P. Ritter, An automated pipeline
for constructing personalized virtual brains from multimodal neuroimaging data.

Neuroimage. **117**, 343–357 (2015).

76. J. W. M. Domhof, K. Jung, S. B. Eickhoff, O. V. Popovych, Parcellation-induced

variation of empirical and simulated brain connectomes at group and subject levels. *Netw.*

Neurosci. **5**, 798–830 (2021).

77. L. Trebaul, P. Deman, V. Tuyisenge, M. Jedynak, E. Hugues, D. Rudrauf, M.

Bhattacharjee, F. Tadel, B. Chanteloup-Foret, C. Saubat, G. C. Reyes Mejia, C. Adam, A.

Nica, M. Pail, F. Dubeau, S. Rheims, A. Trébuchon, H. Wang, S. Liu, T. Blauwblomme,

M. Garcés, L. De Palma, A. Valentin, E.-L. Metsähonkala, V. Bouilleret, E. Landré, W.

Szurhaj, E. Hirsch, L. Valton, R. Rocamora, A. Schulze-Bonhage, I. Mindruta, S.

Francione, L. Maillard, D. Taussig, P. Kahane, O. David, Probabilistic functional

tractography of the human cortex revisited. *Neuroimage*. **181**, 414–429 (2018).

78. A. Palmigiano, T. Geisel, F. Wolf, D. Battaglia, Flexible information routing by transient

synchrony. *Nat. Neurosci.* **20**, 1014–1022 (2017).

79. G. Hahn, A. Ponce-Alvarez, G. Deco, A. Aertsen, A. Kumar, Portraits of communication

in neuronal networks. *Nat. Rev. Neurosci.* **20**, 117–127 (2019).

80. C. Kirst, M. Timme, D. Battaglia, Dynamic information routing in complex networks.

Nat. Commun. **7** (2016), doi:10.1038/ncomms11061.

81. S. H. Strogatz, *Sync: How Order Emerges from Chaos in the Universe* (Hyperion, New

York, 2003).

82. M. Breakspear, S. Heitmann, A. Daffertshofer, Generative models of cortical oscillations:

neurobiological implications of the kuramoto model. *Front. Hum. Neurosci.* **4**, 190

(2010).

83. J. Cabral, E. Hugues, O. Sporns, G. Deco, Role of local network oscillations in resting-
state functional connectivity. *Neuroimage*. **57**, 130–139 (2011).

84. A. L. Allegra Mascaro, E. Falotico, S. Petkoski, M. Pasquini, L. Vannucci, N. Tort-Colet,
E. Conti, F. Resta, C. Spalletti, S. T. Ramalingasetty, A. von Arnim, E. Formento, E.
Angelidis, C. H. Blixhavn, T. B. Leergaard, M. Caleo, A. Destexhe, A. Ijspeert, S. Micera,
C. Laschi, V. Jirsa, M. O. Gewaltig, F. S. Pavone, Experimental and Computational Study
on Motor Control and Recovery After Stroke: Toward a Constructive Loop Between
Experimental and Virtual Embodied Neuroscience. *Front. Syst. Neurosci.* **14**, 1–34
(2020).

85. F. A. Rodrigues, T. K. D. M. Peron, P. Ji, J. Kurths, The Kuramoto model in complex
networks. *Phys. Rep.* **610**, 1–98 (2016).

86. G. Deco, V. K. Jirsa, Ongoing cortical activity at rest: Criticality, multistability, and ghost
attractors. *J Neurosci*. **32**, 3366–3375 (2012).

87. M. L. Kringelbach, G. Deco, Brain States and Transitions: Insights from Computational
Neuroscience. *Cell Rep.* **32**, 108128 (2020).

88. J. D. Tournier, R. Smith, D. Raffelt, R. Tabbara, T. Dhollander, M. Pietsch, D.
Christiaens, B. Jeurissen, C. H. Yeh, A. Connelly, MRtrix3: A fast, flexible and open
software framework for medical image processing and visualisation. *Neuroimage*. **202**
(2019), , doi:10.1016/j.neuroimage.2019.116137.

89. R. S. Desikan, F. Ségonne, B. Fischl, B. T. Quinn, B. C. Dickerson, D. Blacker, R. L.
Buckner, A. M. Dale, R. P. Maguire, B. T. Hyman, M. S. Albert, R. J. Killiany, An
automated labeling system for subdividing the human cerebral cortex on MRI scans into

gyral based regions of interest. *Neuroimage*. **31**, 968–980 (2006).

90.   F. Melozzi, M. M. Woodman, V. K. Jirsa, C. Bernard, *eNeuro*, in press,

doi:10.1523/ENEURO.0111-17.2017.

91.   M. Rubinov, O. Sporns, Weight-conserving characterization of complex functional brain

networks. *Neuroimage*. **56**, 2068–2079 (2011).

910