Evaluating coarse-grained MARTINI force-fields for capturing the ripple phase of lipid membranes

Pradyumn Sharma[1,‡], Rajat Desikan[1,‡,†], K. Ganapathy Ayappa[1,2,]*

[1]Department of Chemical Engineering, Indian Institute of Science, Bengaluru, India

[2]Centre for BioSystems Science and Engineering, Indian Institute of Science, Bengaluru, India

[‡]Co-first authors

[†]Present address: Certara QSP, Certara UK Limited, Sheffield, UK

[*]Corresponding author: K. Ganapathy Ayappa; E-mail: ayappa@iisc.ac.in

Manuscript details:

Title: 96 characters with spaces; Abstract: 272 words; Figures: 6; Tables: 2; References: 76; Supplementary figures: 11

Keywords:

Molecular dynamics; multiscale simulations; CHARMM36; MARTINI; lipid membrane; DPPC; tilted ripple phase; gel phase; phase transition

Abstract

Phospholipids, which are an integral component of cell membranes, exhibit a rich variety of lamellar phases modulated by temperature and composition. Molecular dynamics (MD) simulations have greatly enhanced our understanding of phospholipid membranes by capturing experimentally observed phases and phase transitions at molecular resolution. However, the ripple ($P_{\beta'}$) membrane phase, observed as an intermediate phase below the main gel-to-liquid crystalline transition with some lipids, has been challenging to capture with MD simulations, both at all-atom and coarse-grained resolution. Here, we systematically assess the ability of five coarse-grained MARTINI 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC) lipid force-field (FF) variants, parametrized to reproduce the DPPC gel and fluid phases, for their ability to capture the $P_{\beta'}$ phase. Upon cooling from the fluid phase to below the phase transition temperature with smaller (380-lipid) and larger (> 2200-lipid) MARTINI and all-atom (CHARMM36 FF) DPPC lipid bilayers, we observed that smaller bilayers with both all-atom and MARTINI FFs sampled interdigitated $P_{\beta'}$ and ripple-like states, respectively. However, while all-atom simulations of the larger DPPC membranes exhibited the formation of the $P_{\beta'}$ phase, similar to previous studies, MARTINI membranes did not sample interdigitated ripple-like states at larger system sizes. We then demonstrated that the ripple-like states in smaller MARTINI membranes were kinetically-trapped structures caused by finite size effects rather than being representative of true $P_{\beta'}$ phases. We showed that even a MARTINI FF variant that could capture the tilted $L_{\beta'}$ gel phase, a prerequisite for stabilizing the $P_{\beta'}$ phase, could not capture the rippled phase upon cooling. Our study reveals that the current MARTINI FFs may require specific re-parametrization of the interaction potentials to stabilize lipid interdigitation, a characteristic of the ripple phase.

1 Introduction

Lamellar lipid membranes, composed of two layers of amphiphilic lipids with solvated hydrophilic 'heads' and buried hydrophobic 'tails', exhibit a wide variety of thermotropic phases such as crystalline (L_c), tilted gel ($L_{\beta'}$) or straight gel (L_β), tilted ripple ($P_{\beta'}$), and liquid-crystalline (L_α).[1,2] Lipid bilayers are important constituents of biomembranes which are ubiquitous in living organisms[3], and nanoparticle pharmaceuticals such liposomal drugs[4–6]. Specifically, lipid membrane phases influence: (i) biological processes at the cellular level such as membrane fusion and budding[7], permeation of drugs[8,9], effect of anaesthetics[10], vulnerability to pathogenic toxins[11,12], lipid flip-flop[13,14], neural signalling[15], and the function of membrane proteins[3,7], and (ii) industrial applications such as the design, storage and usage of liposomal drugs[4–6,16], cosmetic formulations[1], and lipid-rich foods[1]. Molecular simulations of membranes at various levels of detail have yielded atomistic insights into the rich phase behaviour of phospholipid membranes while complementing experimental observations[1,17]. All-atom molecular dynamics (MD) simulations have successfully captured the low temperature $L_{\beta'}$ or L_β phases, the liquid ordered and liquid disordered phases, and the main gel-to-fluid ($L_{\beta'}$ to L_α) phase transition in simple lipid bilayers while providing detailed insights into the rate-limiting formation of the melting seed that nucleates the transition[18]. Similarly, the molecular structure of lipid domains in the elusive $P_{\beta'}$ phase, which commonly occurs upon cooling membranes from the L_α phase to below the gel-to-fluid transition temperature (pre-transition region), and challenging to decipher at atomistic resolution through experiments, was unravelled via detailed MD simulations[19–21], consistent with observations.

While these *in silico* studies illustrate the wide applicability of all-atom lipid force-fields (FFs), all-atom MD simulations are typically limited by computational bottlenecks to nanoscales, and complex membrane processes typically require microscale MD simulations[17,22]. Therefore, coarse-grained (CG) lipid models such as those used in the dissipative particle dynamics (DPD), MARTINI, and other variants have been developed[22–26]. Among the CG models, the MARTINI

framework has been successfully used to study a variety of complex biological systems outside

the purview of all-atom MD simulations[22]. For example, MARTINI simulations have been used to

carry out microsecond simulations of an idealized 20000-lipid plasma membrane comprising 63

different species of lipids[27], a difficult feat with typical all-atom simulations. The versatility and

computational tractability of CG models such as MARTINI render them as attractive options for

capturing the microscopic details of complex membrane phenomena[22,24]. However, since CG

models by definition possess reduced molecular detail[28], their interaction potentials must be

continuously assessed and refined, often by systematic comparison with corresponding detailed

all-atom models[25,26,28–31], to enable increasingly accurate CGMD simulations.

DPPC possesses a large headgroup and favourable inter-head interactions that stabilizes

a tilted gel ($L_{\beta'}$) phase at lower temperatures, unlike lipids with smaller headgroups such as

phosphatidylethanolamine that favour the straight gel (L_{β}) phase[20,21,32]. At higher temperatures,

DPPC exhibits a main $L_{\beta'}$ to L_{α} phase transition, with the $P_{\beta'}$ phase occurring at pre-transition

temperatures[32–38]. The $P_{\beta'}$ phase has interesting properties such as the influence of thermal

history on the symmetry of the ripple domains[20], and therefore has been the subject of extensive

experimental[33,37–40] and theoretical[18–21,25,30,35,36,41–45] studies. In Table 1, we summarize some of

the previous studies where the rippled phase has been observed, including details of lipid type,

resolution (all-atom, united atom or coarse-grained) and FF. Despite extensive *in silico*

investigations, MD simulations that capture the $P_{\beta'}$ phase have proven to be challenging since

the ripple domain structure and formation is not only a function of the specific lipid FF (Table 1),

but also depends on the cooling/heating protocols[18,34] and system size effects[20,34,43]. In spite of

these challenges, MD simulations of lipid bilayers in the ripple phase have revealed elusive details

of molecular organization that were challenging to probe via experiments, such as the asymmetric

sawtooth-like molecular organization of lipids in the DPPC ripple domains[19–21].

Table 1. List of previous studies which have explored the $P_{\beta'}$ phase using MD simulations.

Lipid	Year	Resolution	Force-field	Comparison with experiments	Citation
DPPC	2005	United atom	Modified Berger lipid	Qualitative comparison with DMPC electron density profiles	Ref.[19]
DMPC	2005	Coarse-grained	DPD	–	Ref.[21]
DPPC	2007	Coarse-grained	Custom (single tailed)	Qualitative comparison with DPPC electron density maps	Ref.[20]
DMPC, DPPC	2012	Coarse-grained	DPD, MARTINI	Enthalpic signatures of phase transitions from MD compared with differential scanning calorimetry	Ref.[34]
BTMAC+SA	2014	United atom	GROMOS 87, OPLS	–	Ref.[41]
DPPC	2014	Coarse-grained	MARTINI	–	Ref.[46]
DMPC	2016	All-atom	CHARMM36, Lipid14	–	Ref.[42]
DPPC	2017	Coarse-grained	Dry MARTINI	–	Ref.[47]
DMPC, DPPC	2018	All-atom	CHARMM36	Comparison of bilayer properties with X-ray diffraction experiments	Ref.[44]
DPPC	2018	All-atom	CHARMM36	–	Ref.[18]
DPPC	2018	All-atom, coarse-grained	CHARMM36, MARTINI	–	Ref.[30]
DPPC	2019	All-atom	CHARMM36	–	Ref.[43]
BTMAC+SA	2020	All-atom, coarse-grained	CHARMM36, MARTINI	–	Ref.[25]

Abbreviations: DPD – dissipative particle dynamics; DPPC – 1,2-dipalmitoyl-sn-glycero-3-phosphocholine; DMPC – 1,2-dimyristoyl-sn-glycero-3-phosphocholine; BTMAC – benzyltrimethylammonium chloride; SA – stearyl alcohol.

While DPPC membranes described by the standard MARTINI FF and its variants have been shown to capture the gel and fluid states[22,30], there are conflicting reports of their ability to capture the ripple phase. Some MARTINI studies have reported ripple formation both with and without enhanced sampling[30,46,47], while others have reported that MARTINI membranes cannot capture the ripple phase[34], with the interdigitated ripple-like domains observed upon quasi-static cooling from the L_α phase reported to be artefacts caused by finite size effects[34]. Therefore, the ability of MARTINI membranes to capture the $P_{\beta'}$ phase still remains an unresolved problem. To systematically address this issue, we assess the ability of commonly employed[18,19,23,29,30,34,43,44,46]

DPPC bilayers, described by five MARTINI lipid FF variants (described subsequently), to capture the interdigitated tilted ripple membrane phase ($P_{\beta'}$).

We first establish a standard simulated-annealing-based cooling MD protocol, built on previous studies[18–20,30,43,44], that consistently yielded a ripple phase upon cooling from the fluid phase to below the gel-to-fluid phase transition temperature with all-atom DPPC membranes across multiple simulation replicates. By employing the same protocol with MARTINI DPPC membranes, and by fixing other important variables such as initial configurations (L_α phase), system sizes and cooling rates, we checked whether the MARTINI membranes yielded stable ripple phase structures, similar to those observed in all-atom simulations. Five popular MARTINI lipid and water FF combinations were tested: (1) standard MARTINI version 2.2 lipid FF with normal ('MARTINI2n') or polarizable[48] ('MARTINI2p') water, (2) optimized MARTINI FF for DPPC – has refined bond parameters for better reproduction of DPPC membrane phase behaviour and can capture the tilted $L_{\beta'}$ gel phase – with either normal ('MARTINI2On') or polarizable ('MARTINI2Op') water, and, (3) MARTINI 3 beta (version 3.0.b.3.2) with its in-built water model ('MARTINI3b'). For each FF variant, 10 independent replicates of simulated cooling were performed, ensuring sufficient sampling for robust statistical estimates[18,49]. We observed that smaller (380-lipid) MARTINI DPPC bilayers sampled interdigitated ripple-like states that were not observed with larger DPPC membranes (> 2200-lipid). The ripple-like states in these smaller MARTINI DPPC membranes were demonstrated to be kinetically-trapped structures caused due to finite size effects, rather than being representative of a true ripple phase. However, all-atom simulations of both the smaller and larger DPPC membranes exhibited the formation of the $P_{\beta'}$ phase, similar to previous studies. Our observations thus assess the ability of MARTINI lipid FFs to capture the ripple phase and serve as a guide for future CG lipid FF development.

2 Methods

2.1 All-atom MD simulations of DPPC membranes

The MD protocols employed in this study for simulating DPPC membranes in the gel, ripple and fluid phases are briefly described below. All the simulations performed in this study are summarized in Table 2.

For the small membrane atomistic simulations, two charge-neutral, solvated, fluid phase (L_α), 368-lipid membrane configurations at 350 K, with similar spatial coordinates but different starting velocities, were set up using CHARMM-GUI[50–52] as independent replicates (Figs. 1 and S1). The size of the lipid patches was approximately 15.2 nm (x-axis) x 7.6 nm (y-axis) x 9.4 nm (z-axis), with the characteristic dimension of the simulation box in the membrane plane in the range of the periodicity of 12-16 nm corresponding to previously observed ripple profiles in DPPC membranes[19,53]. This was important because the quality of the observed ripples was found to depend on the characteristic box dimension along the ripple formation[20]. Systems whose characteristic box length in the membrane plane were non-integer multiples of ripple periodicity resulted in frustrated and interconnected structures, unlike the clean ripples observed in systems with integer multiple box lengths[20]. Each system contained $>10^5$ atoms. The water:lipid ratio was set at 50:1, ensuring sufficient hydration normal to the membrane plane (z), so that the systems could be approximated as isolated lipid bilayers along the z-axis.

Atomistic simulations were performed using the CHARMM36 FF[42,54,55] implementation (March 2019 version from http://mackerell.umaryland.edu/charmm_ff.shtml) in Gromacs (version 2018.6)[56], according to the currently advocated best practices for lipid bilayer simulations[17,57–59]. (See Refs.[42,60] for a brief discussion of all-atom and united atom lipid FFs.) A leapfrog integrator (2 fs integration time step) was used along with Verlet buffered lists and three-dimensional periodic boundary conditions. Electrostatic interactions were computed with the particle mesh Ewald algorithm[61], and van der Waal interactions were smoothly switched to zero between 1.0 to

1.2 nm. Covalent bonds involving hydrogen atoms were constrained using the LINCS algorithm[62].

Simulations were performed in the NPT ensemble with semi-isotropic pressure coupling along

the lateral bilayer plane, by employing the stochastic velocity-rescaling thermostat[63] for

temperature control, and the Parrinello-Rahman barostat[64] for pressure control, with coupling

constants of 0.1 and 2.0 ps, respectively. After energy minimization and equilibration (100 ns) of

the initial configurations in the L_α phase (350 K) according to standard CHARMM-GUI protocols[51],

the systemic temperatures of the replicates were continuously reduced from 350 K to 290 K at

the rate of 1 K per 6 ns, or -0.167 K/ns, by employing the stochastic simulated annealing MD

(SAMD) algorithm implemented in Gromacs[56] (360 ns per SAMD replicate). We chose to cool

slightly beyond the pre-transition temperature of 300-310 K at which the $P_{\beta'}$ phase is typically

observed[18,32,34–36], because super-cooling is often required in kinetic MD simulations for eliciting

a phase transition[18,65,66]. The resulting cooled bilayers at 290 K were further equilibrated at a

constant temperature of 290 K for 200 ns for thermal and structural stability.

To test for finite size effects and to compare with MARTINI simulations (described next),

two large 2208-lipid membranes measuring 47.0 nm x 15.7 nm x 15.0 nm and possessing a high

water:lipid ratio of 128:1 ($>10^6$ atoms total; Table 2) were equilibrated in the L_α phase at 350 K.

Subsequently, both replicates were cooled to 290 K at the rate of -0.167 K/ns (chosen based on

Ref.[18]) and then equilibrated at 290 K for an additional 200 ns.

2.2 MARTINI simulations of DPPC membranes

380-lipid MARTINI DPPC membranes of size 15 nm x 7.5 nm x 10 nm were prepared using

the 'INSANE' python module[67]. The water-bead:lipid ratio was ~13.3:1. Since each water bead is

equivalent to 4 water molecules, this corresponds to a water:lipid ratio of ~53:1, similar to the

small all-atom membranes. As described in the Introduction, five MARTINI FF variants were used:

MARTINI2n, MARTINI2p, MARTINI2On, MARTINI2Op and MARTINI3b. MARTINI2n and

MARTINI2On systems also contained anti-freeze solvent beads to prevent artefactual freezing of

MARTINI water beads.

Input CGMD parameters were fixed based on previously reported parameter settings for

improved performance[68]. All CGMD simulations were performed with a 20 fs timestep in the NPT

ensemble by employing the stochastic velocity-rescaling thermostat[63] (coupling constant of 1.0

ps) and the Parrinello-Rahman semi-isotropic barostat[64] (coupling constant of 12.0 ps).

Electrostatics were treated with the reaction-field algorithm[69] with a cut-off of 1.1 nm, relative

dielectric constant of 2.5 and 15 for systems with and without polarizable water, respectively, and

the relative dielectric constant of the reaction field set to infinity. The van der Waal forces were

computed with a cut-off of 1.1 nm. Post energy minimization, five CGMD membranes, each

described by a MARTINI FF, were equilibrated in the L_α phase at 350 K for 100 ns, followed by

simulated cooling from 350 K to 280 K at -0.167 K/ns (420 ns per SAMD replicate). For each FF,

10 independent SAMD replicates were simulated for ensuring robustness of observations[49],

resulting in a total of 50 CG-SAMD simulations (Figs. 2, S2-S6). The replicates which yielded

ripple-like phases were further equilibrated for 1-5 µs at 280 K to ensure that the ripples had

attained thermal and structural equilibrium (Fig. 2).

To test for finite-size effects, many large MARTINI DPPC bilayer configurations, with both

rectangular and square geometries, were constructed and simulated (Table 2).

Table 2. Summary of all-atom and MARTINI simulations performed in this study.

Forcefield	Time (µs)[#]	Number of replicas	System size (L_x x L_y x L_z nm³)	Number of lipids	Number of particles	Type of MD simulation[a]	Temp. (K)	Figures
			Small membrane simulations					
CHARMM36	0.10	1	15.7 x 7.8 x 8.4	368	103040	EQMD[&]	350	
CHARMM36	0.36	2	13.4 x 6.7 x 10.6	368	103040	SAMD	350-290	1, S1
CHARMM36	0.20	2	13.4 x 6.7 x 10.6	368	103040	EQMD	290	
MARTINI2n	0.10	1	15.8 x 7.9 x 9.6	380	9626	EQMD[&]	350	2
MARTINI2On	0.10	1	15.8 x 7.9 x 9.6	380	9620	EQMD[&]	350	-
MARTINI3b	0.10	1	15.9 x 7.9 x 9.1	380	9626	EQMD[&]	350	-

MARTINI2p	0.10	1	16.0 x 8.0 x 9.0	380	19761	EQMD[&]	350	-
MARTINI2Op	0.10	1	15.5 x 7.8 x 9.5	380	19764	EQMD[&]	350	-
MARTINI2n	0.42	10	14.2 x 7.1 x 11.1	380	9626	SAMD	350-280	S2
MARTINI2On	0.42	10	13.9 x 7.0 x 11.5	380	9620	SAMD	350-280	S3
MARTINI3b	0.42	10	14.2 x 7.1 x 10.5	380	9626	SAMD	350-280	S4
MARTINI2p	0.42	10	14.4 x 7.2 x 10.2	380	19761	SAMD	350-280	S5
MARTINI2Op	0.42	10	13.8 x 6.9 x 10.9	380	19764	SAMD	350-280	S6
MARTINI2n	5	2[b]	14.3 x 7.1 x 11.0	380	9626	EQMD	280	2, 6
MARTINI2On	5	3[c]	13.9 x 7.0 x 11.4	380	9620	EQMD	280	2, 6, S11
MARTINI3b	5	1[d]	14.0 x 7.0 x 10.7	380	9626	EQMD	280	2
MARTINI2p	5	1[e]	14.3 x 7.2 x 10.2	380	19761	EQMD	280	2
MARTINI2Op	5	1[f]	13.8 x 6.9 x 10.9	380	19764	EQMD	280	2
Large membrane simulations								
CHARMM36	0.02	1	46.4 x 15.4 x 16.1	2208	1135872	EQMD[&]	350	
CHARMM36	0.36	2	40.3 x 13.4 x 19.9	2208	1135872	SAMD	350-290	3, S7
CHARMM36	0.20	2	40.4 x 13.5 x 19.8	2208	1135872	EQMD	290	
MARTINI2n	0.10	2	46.5 x 15.5 x 20.6	2204	112719	EQMD[&]	350	4
MARTINI2On	0.10	2	46.3 x 15.4 x 20.8	2204	112719	EQMD[&]	350	-
MARTINI3b	0.10	2	47.1 x 15.7 x 18.8	2204	112717	EQMD[&]	350	-
MARTINI2n	0.42	2	39.9 x 13.3 x 26.0	2204	112719	SAMD	350-280	4, S9a
MARTINI2On	0.42	2	39.9 x 13.1 x 26.6	2204	112719	SAMD	350-280	
MARTINI3b	0.42	2	41.3 x 13.8 x 22.7	2204	112717	SAMD	350-280	
MARTINI2n	3	1	39.6 x 13.2 x 12.7	2280	57756	EQMD	280	5, S10
MARTINI2On	5, 8, 10	3	39.7 x 13.2 x 12.6	2280	57720	EQMD	280	5, 6, S10, S11
MARTINI3b	10	1	40.0 x 13.3 x 11.8	2280	57756	EQMD	280	5, S10
MARTINI2On	0.10	1[g]	46.5 x 46.5 x 20.2	6728	333604	EQMD[&]	350	-
MARTINI2On	0.42	2[g]	39.5 x 39.5 x 26.0	6728	333604	SAMD	350-280	S9b

[#]Simulation times from MARTINI CGMD were <u>not</u> scaled. Total simulation time in this study: ~2.5 μs (all-atom MD)

and ~87 μs (MARTINI CGMD). [&]Initial simulations to equilibrate starting configurations. [a]EQMD – equilibrium MD

simulations in the NPT ensemble; SAMD – simulated annealing MD simulations at constant pressure. [b]Replicates

with the 'best' ripple-like (Repl. 1 in Fig. S2) and straight gel (Repl. 7 in Fig. S2) phases were further equilibrated for

5000 ns. [c]Replicates with the 'best' ripple-like (Repl. 5 in Fig. S3), tilted ripple-like (Repl. 7 in Fig. S3) and tilted gel

(Repl. 3 in Fig. S3) phases were further equilibrated for 5000 ns. [d]Replicate with the 'best' ripple-like phase (Repl. 9

in Fig. S4) was further equilibrated for 5000 ns. [e]Replicate with the 'best' ripple-like phase (Repl. 5 in Fig. S5) was

further equilibrated for 5000 ns. [f]Replicate with the 'best' ripple-like phase (Repl. 10 in Fig. S6) was further equilibrated

for 5000 ns. [g]Square membrane patch; all others are rectangular membrane patches

2.3 Analysis and visualization

MD snapshots are illustrated using the visual molecular dynamics 1.9.3 (VMD)[70] software.

The DPPC tails in the upper and lower leaflets of both all-atom and MARTINI membranes are

consistently colored cyan and magenta, respectively. In all-atom membranes, phosphorus atoms

(P) are colored orange, and nitrogen atoms in the choline group are colored blue. Similarly, in

MARTINI membranes, the phosphate bead is colored orange and the choline bead is colored

blue. The minimum inter-leaflet P-P distance (ξ), which is an order parameter to track leaflet

interdigitation (Fig. 1b), was computed using the inbuilt 'gmx mindist' tool in Gromacs. The ξ from

the fluid and interdigitated regimes are fit to a four-parameter piecewise-linear function (Eq. 1) to

accurately estimate the time of the onset of membrane interdigitation, τ.

$$\xi = (m_1 t + c)\big(1 - H(t - \tau)\big) + (m_2 t + (m_1 - m_2)\tau + c)H(t - \tau) \qquad\qquad -(1)$$

Here, m_1 and m_2 are the slopes of the two lines corresponding to the fluid and interdigitated

regimes (when it occurs), respectively, c is the intercept of the fluid-phase line (the intercept of

the other line is $(m_1 - m_2)\tau + c$ from the point of intersection), and $H(t - \tau)$ is the Heaviside

function which is zero for $t < \tau$, and one for $t \geq \tau$. If interdigitation occurs, $m_2 < 0$, else $m_2 > 0$

(illustrated in Fig. S2). The fitting was performed using the 'lsqcurvefit' function (optimization

toolbox) in conjunction with 'multistart' (global optimization toolbox) implemented in MATLAB

2018a (https://www.mathworks.com). All other plots are also created with MATLAB. The

combined Voronoi-thickness modulation heatmaps are plotted using the 'voronoi' MATLAB

function on the lipid phosphorus atom x, y coordinates (upper leaflet) from the corresponding MD

snapshot. The membrane thickness[41] is computed using the 'Membplugin' 1.1 tool[71] in VMD and

visualized using MATLAB. Similar to a previous study[72], lipid tilt angles from MARTINI

membranes (L_β vs. $L_{\beta'}$) were computed from the last 10 ns of the respective MD trajectories,

using the inbuilt 'gmx gangle' tool in Gromacs. The lipid tilt was defined as the angle between the

lipid vector and the leaflet normal, with the lipid vector defined as the vector joining the center of

mass of the terminal tail beads with the head phosphate bead.

3 Results and Discussion

3.1 Ripple formation in atomistic simulations of DPPC membranes

To robustly capture the ripple phase in all-atom DPPC membranes[26], well-equilibrated, fluid, 368-lipid DPPC membranes at 350 K (time point I in Fig. 1*a, c*) were cooled to 290 K, lower than the experimentally observed pre-transition temperature of 300-310 K[32,34–36] and sufficient to ensure leaflet interdigitation[18,44], using the stochastic simulated annealing MD algorithm, followed by constant temperature equilibration MD (EQMD) in the NPT ensemble at 290 K to ensure thermal equilibration and structural convergence (details in Methods; 2 independent MD replicates). Along with the initial state (time point I), three other time points are important in the combined SAMD+EQMD trajectory (Fig. 1*a, c*) – the fluid membrane configuration where interdigitation starts (time point II), the fully interdigitated ripple phase membranes at the end of simulated cooling (time point III), and the converged systems at the end of constant temperature equilibration (time point IV). The extent of interdigitation was tracked by the minimum inter-leaflet P-P distance with time (ξ; Methods and Eq. 1; Fig. 1*b, c*). A higher initial value marked the baseline corresponding to the fluid phase (350 K; time point I), and an abruptly decreasing trend indicated significant interdigitation of lipids from opposite leaflets, with the onset of interdigitation marked by the inflection point (time point II – τ; see Methods, Fig. S2, and Eq. 1). By the end of simulated cooling (290 K; time point III), interdigitation was complete and a membrane with an 'asymmetric'[20] 1D ripple was formed, similar in structure to previous studies[19,20,43,44], structurally stable upon further 200 ns equilibration at 290 K (time point IV), and with clearly distinct major and minor arms (illustrated in Fig. S1*a*).

Fig. 1. All-atom Lα DPPC membranes consistently form ripples upon cooling. (a) MD snapshots (colouring described in Methods) during the cooling of a 368-lipid, all-atom, DPPC membrane from 350 K to 290 K. Time points: I – initial equilibrated L_α phase membrane at 350 K before the start of cooling, II – start of interdigitation, II' –time at which the average inter-leaflet distance in the time series is at a minimum (300 K), III – end of simulated cooling (290 K), and IV – end of equilibration MD at constant temperature (290 K). Interdigitated domains are highlighted by yellow dashed circles. **(b)** Schematic illustrating the minimum inter-leaflet P-P distance, ξ, which is used as an order parameter to track leaflet interdigitation (Eq. 1). **(c)** Time trace (red) and biphasic linear fits (black) of ξ in cooling simulations of atomistic DPPC membranes. **(d)** Voronoi diagram from the x-y coordinates of the upper leaflet P atoms depicting the area per head group of individual lipids is superimposed on the thickness modulations of membranes at time points I-IV, corresponding to the MD snapshots in (a) (viewing angle pictorially depicted).

Voronoi-thickness membrane maps, which are Voronoi diagrams of the upper membrane leaflet superimposed onto P-P thickness heatmaps[25,41], are shown in Fig. 1*d*, corresponding to the top views of MD snapshots in Fig. 1*a*. The trends were similar[41] when the Voronoi diagrams were plotted using the lower leaflets (not shown). Compared to the uniform topography in the fluid phase, the formation of interdigitated (blue) and gel (yellow) domains in the ripple phase can be clearly observed. There is negligible difference in membrane topographies between time points III and IV except for the lateral movement of membrane across period boundaries, indicating that

the ripple phase membrane configuration at the end of simulated cooling is at equilibrium. An

interesting observation was that the interdigitated ripple domain (Fig. 1d) appeared to be tilted in

the lateral plane, whereas ripple domains in previous studies[19,20,41] were typically parallel to one

of the lateral box edges. Previous mesoscopic lipid bilayer simulations[21] have revealed that at the

highest value of the lipid head-head repulsion parameter explored, lower and higher values of

which either abrogated or stabilized the ripple phase, respectively, the bilayer formed a laterally

tilted ripple domain and a stable $P_{\beta'}$ membrane phase similar to our observations.

Another independent all-atom MD replicate also yielded a similar interdigitated DPPC

ripple phase membrane upon cooling (Fig. S1b), thus validating the robustness of the above MD

protocol to produce the $P_{\beta'}$ phase from fluid membrane configurations. We next employed this

MD cooling protocol to fluid-phase MARTINI membranes (all five FF variants) and evaluated their

ability to capture the ripple phase.

3.2 Ripple formation in smaller MARTINI membranes

Fluid phase, 380-lipid, DPPC MARTINI membranes belonging to five MARTINI FF variants

were cooled from 350 K to 280 K, and evaluated for their ability to form the interdigitated $P_{\beta'}$

membrane phase (see Introduction and Methods). The lower bound for temperature was set to

280 K, compared to 290 K for all-atom, because a previous CGMD study showed that ripple-like

interdigitated phases in MARTINI DPPC membranes were formed between 280-290 K[34]. The

trajectory of an SAMD+EQMD replicate with the MARTINI2n FF is shown with time points I-IV

highlighted (Fig. 2a-c). The constant temperature EQMD phase post cooling was chosen to be 5

µs for MARTINI (Fig. 2b), compared to 200 ns for the all-atom simulations (Fig. 1c). Similar to the

time course seen in atomistic DPPC membranes, a majority (70%), but notably not all,

MARTINI2n replicates showed interdigitation of membrane leaflets (time points I-IV on the $\xi(t)$

plot, Voronoi-thickness maps pertaining to MD snapshots at time points I and IV), thus apparently

capturing ripple-like membrane states. This is in accord with previous single replicate studies,

where MARTINI membranes of similar sizes captured the ripple phase[30,34,46,47]. In the rest of the

MARTINI2n replicates (30%), the membranes bypassed the interdigitated ripple-like state and

directly froze into L_β configurations (Fig. S2). This emphasizes that with simulated cooling MD,

rippling in MARTINI membranes depended on the initial configuration and starting velocities, and

therefore necessitated multiple MD replicates for obtaining robust trends.[49]

The orientation of the interdigitated domains in the MARTINI2n membranes were

perpendicular to the x-axis (Fig. 2*c*) in contrast with the laterally tilted interdigitated domains seen

in all-atom simulations (Figs. 1*d* and S1*b*). Similar domains were observed in a previous MARTINI

study.[34]

Formation of ripple-like phases in DPPC membranes described by the other FF variants –

MARTINI2On, MARTINI3b, MARTINI2p, and MARTINI2Op – are summarized in Fig. 2*d-g* (all MD

replicates for each FF variant displayed in Figs. S3-S6, respectively). MARTINI2On is similar to

MARTINI2n, with 70% of the replicates forming ripple-like (Figs. 2*d* and S3). MARTINI3b, the beta

version of the improved[73] next generation MARTINI3 FF (unpublished and currently under

development), showed ripple-like formation in a minority (30%) of the replicates (Figs. 2*e* and S4).

The MARTINI FF combinations with polarizable water, MARTINI2p (Figs. 2*f* and S5) and

MARTINI2Op (Figs. 2*g* and S6), showed ripple-like states in only 10% and 30% of replicates,

respectively, despite the reparameterization of polarizable water to improve the accuracy of

solvent electrostatic effects[48], important for capturing macroscopic membrane properties[54,55]. Our

observations with MARTINI2p are to be contrasted with previous single-replicate MARTINI[34]

simulations, where quasi-static cooling of fluid-phase membranes resulted in the formation of the

ripple-like domains only with polarizable water. While differences in MD protocols and differences

in the initial configurations may explain the disparities, careful future investigations, with systems

of varying size, may be necessary to determine the predictive accuracy of polarizable water-lipid

interactions.

Fig. 2. Smaller MARTINI membranes form a ripple-like interdigitated phase upon cooling.
(a) Similar to all-atom MD (Fig. 1), snapshots from time points I-IV from the simulated cooling of
a MARTINI2n membrane replicate from 350 K – 280 K are shown (Replicate 1 from Fig. S2).
Interdigitated regions are highlighted by a yellow dashed circle. **(b)** Minimum inter-leaflet P-P
distance with time in the 420 ns cooling and 5000 ns constant temperature equilibration phases.
(c) Voronoi-thickness maps contrasting the initial fluid membrane and the final interdigitated
ripple-like membrane (time points I and IV, respectively). Cooling of 7/10 independent MARTINI2n
MD replicates resulted in the formation of a ripple-like phase (Fig. S2). Other MARTINI FF
variants: **(d)** MARTINI2On (Replicate 5 from Fig. S3), **(e)** MARTINI3b (Replicate 9 from Fig. S4),
(f) MARTINI2p (Replicate 5 from Fig. S5) and **(g)** MARTINI2Op (Replicate 10 from Fig. S6), show
the formation of ripple-like phases in 7, 2, 1 and 3 out of 10 replicates each, respectively (Figs.
S3 – S6). These snapshots correspond to time point IV.

3.3 Testing for ripple formation in large atomistic and MARTINI DPPC membranes upon simulated cooling

In a previous study, Rodgers et al.[34] observed the interdigitation of lipids in relatively small

512-lipid MARTINI DPPC membranes upon quasi-equilibrium, stepwise, simulated cooling of fluid

membranes from 325 K to less than 280 K. These small membranes apparently sampled an

interdigitated ripple-like phase, similar to our observations (Fig. 2). However, when the exercise

was repeated with 4-times-larger (2048-lipid) DPPC membranes, the ripple-like phase could not

be reproduced. Thus, Rodgers et al.[34] concluded that the interdigitated ripple-like states in smaller

membranes were actually 'intermediate' kinetically trapped membrane states rather than a true

ripple phase. In contrast to these MARTINI simulations, a large all-atom 3200-lipid DPPC

membrane has been shown to form an interdigitated ripple phase[30], albeit with a different MD

protocol. Therefore, we next tested ripple formation in large all-atom DPPC membranes with our

simulated cooling MD protocol (similar to Fig. 1), that would re-establish a benchmark to test for

ripple formation and system size effects in large MARTINI membranes.

We set up two 2208-lipid DPPC bilayer configurations, which were 6-times-larger than the smaller

atomistic DPPC membranes shown in Fig. 1, and larger than the membranes employed by

Rodgers et al.[34], and pre-equilibrated them in the L_α phase at 350 K (Fig. 3a). Subsequently,

these membranes were cooled to 290 K with the same SAMD protocol. One of the all-atom MD

replicates in the fluid phase formed an interdigitated ripple phase membrane by 290 K (Fig. 3b).

The other replicate (Fig. S7) formed a non-interdigitated 'symmetric' ripple[20] in the larger

dimension, which is typically observed upon slow cooling from the fluid phase to approximately

the gel-to-fluid transition temperature[20], and an asymmetric interdigitated ripple (similar to Figs.

1, 3) in the smaller dimension. Therefore, both smaller and larger all-atom DPPC membranes,

described by the CHARMM36 FF, sample the thermodynamic ripple phase upon simulated

cooling MD without any evidence of finite size artefacts. In contrast to smaller membranes (Figs.

1d and S8), the Voronoi-thickness map of the first replicate (Figs. 3b and S8) shows reticulated

ripple domains. Our observations are in accord with previous studies[30,43], where simulated

quenching of 1280-lipid and 3200-lipid all-atom DPPC membranes to below the main transition

 temperature also resulted in reticulated ripple domains. It is conceivable that significantly longer

 runs would be required to anneal and reorganize the ripples along a given direction.

Fig. 3. Ripple formation from all-atom MD of a large DPPC membrane. (a) Front and top views of a large 2208-lipid DPPC membrane which was pre-equilibrated in the fluid phase at 350 K. For comparison, a representative simulation box of a smaller all-atom membrane (Fig. 1) is shown in red. The large membrane is 6-times (3X times 2Y) larger than the smaller membrane. **(b)** Front view of the large membrane after cooling to 290 K and EQMD (corresponds to time point IV), and its Voronoi-thickness modulation map (blue represents interdigitated regions) shows the formation of an aperiodic and reticulated interdigitated ripple domain (also see Figs. S7-S8).

We next tested whether large, fluid, rectangular, 2204-lipid MARTINI DPPC membranes, with normal (not polarizable) MARTINI water, also sampled the interdigitated ripple-like phase upon cooling from 350 K to 280 K. Two initial configurations each for MARTINI2n, MARTINI2On and MARTINI3b FFs were set up analogous to the large all-atom membranes in Fig. 3, and pre-equilibrated at 350 K, followed by cooling to 280 K (Figs. 4 and S9). However, none of the large MARTINI membranes captured a ripple-like phase upon cooling to 280 K. MARTINI2n and MARTINI3b membranes directly froze into the L_β phase, while the MARTINI2On membranes transitioned to the $L_{\beta'}$ phase (Fig. 4b); confirmed by independent MD replicates (Fig. S9*a*). To ensure that membrane geometry did not influence these results, we repeated these simulations with large 6728-lipid square MARTINI2On membranes, and these also did not sample ripple-like states (Fig. S9*b*). Since only smaller (380-lipid) but not larger (>2204-lipid) DPPC MARTINI membranes captured the interdigitated ripple-like states, in contrast to all-atom membranes, we inferred that the smaller MARTINI ripple-like membranes (Figs. 2, S2-S6) represent kinetically-trapped states caused by finite box-size effects rather than true ripple states.

3.4 Equilibrium MD simulations of large MARTINI membranes initialized in the ripple-like phase show loss of interdigitation

To confirm the aforementioned inference, we employed an orthogonal approach where multi-μs equilibrium simulations of large MARTINI membranes, initialized in periodic interdigitated ripple-like configurations, were performed at 280 K. These large systems were created from the

final, interdigitated, ripple-like, smaller (380-lipid) MARTINI membrane configurations after 1 μs of constant temperature EQMD at 280 K. The smaller membranes were repeated 6-times along the lateral bilayer plane across periodic boundaries (3x times 2y), and the initial large 2204-lipid ripple-like membrane configurations thus created are displayed in Fig. 5*a, c, d* and correspond to MARTINI2On, MARTINI2n and MARTINI3b, respectively.

The rationale behind performing equilibrium MD of these large systems was simply to test if the ripple-like states were stabilized due to finite size effects. Upon 3-10 μs of constant temperature NPT-MD at 280 K, we observed that the large interdigitated MARTINI membranes spontaneously froze into the gel phase, as shown by both MD snapshots (Fig. 5) and Voronoi-thickness maps (Fig. S10), indicating that the initial rippled state configuration was unfavourable at larger system sizes. This phenomenon was observed for all the three MARTINI FF variants, although the time scale over which interdigitation disappeared leading to formation of the final gel phase configuration varied amongst the 3 FFs. MARTINI2On showed progressive loss of interdigitation which was nearly complete by 8 μs (Figs. 5*b* and S10*a*). Longer simulations may have shown a complete loss of interdigitation, but we did not further pursue these computationally intensive MD simulations since the progressive loss of interdigitation with time was apparent (Fig. S10*a*). MARTINI2n showed a rapid and complete loss of interdigitation by 3 μs (Figs. 5*c* and S10*b*), and MARTINI3b showed near-complete loss of interdigitation by 10 μs (Figs. 5*d* and S10*c*), similar to MARTINI2On. In contrast, none of the smaller membranes showed any loss of interdigitation upon 5 μs of constant temperature EQMD at 280 K (Fig. 2).

Together with observations from the simulated cooling of large fluid membranes, these two orthogonal MD approaches (Figs. 4 and 5) strongly suggest that MARTINI membranes (all FF variants) do not capture the thermodynamic ripple phase unlike all-atom membranes. Our systematic study implies that ripple-like phase formation observed in smaller MARTINI membranes was due to kinetic trapping of interdigitated leaflets at small system sizes. The few studies which indeed report MARTINI ripple phase membranes, both with[47] and without[34,46]

enhanced sampling, are with smaller membranes composed of 390[47] and 512[34,46] lipids,

respectively. Therefore, based on our finding, the formation of rippled bilayers in these studies

could be attributed to finite size artefacts[34]. Thus, MARTINI FF parameters may have to be tuned

to reproduce the lipid membrane phase diagram, in particular the $P_{\beta'}$ phase, more accurately. FF

approaches that tune the bonded parameters of MARTINI lipids based on comparison with all-

atom simulations[25,30,34] may result in models that better capture membrane phase behaviour.

a   MARTINI2n, pre-equilibrated 2204-lipid L_α membrane at 350 K (initial)

Simulated cooling
350 K − 280 K, 0.167 K/ns

No ripple, straight gel phase (L_β) membrane at 280 K (final)

b   MARTINI2On, tilted gel phase ($L_{\beta'}$) membrane at 280 K (after cooling)

c   MARTINI3b, straight gel phase (L_β) membrane at 280 K (after cooling)

Fig. 4. Lack of ripple formation in large MARTINI DPPC membranes. (a) A representative
2204-lipid, rectangular, MARTINI2n DPPC membrane, pre-equilibrated at 350 K in the L_α phase,
was cooled directly into the gel phase at 280 K without the membrane sampling a ripple-like
phase, unlike all-atom systems (see Fig. 3). This behaviour was also observed with **(b)**
MARTINI2On and **(c)** MARTINI3b FFs. Independent MD replicates are shown in Fig. S9*a*. Two
independent simulation replicates, where large 6728-lipid, square, MARTINI2On membranes
were cooled to 290 K, also showed a similar lack of sampling of a ripple-like phase (Fig. S9*b*),
suggesting that system geometry did not influence our observations.

a MARTINI2On, 2280-lipid rippled membrane at 280 K (initial, top view)

Small rippled membrane patch, repeated 3x2 across PBC to create the larger membrane

Front view

Interdigitation

b Final snapshot after 8 µs MD at 280 K

c MARTINI2n (initial, front view)   3 µs MD at 280 K

d MARTINI3b (initial, front view)   10 µs MD at 280 K

Fig. 5. MD simulations of large MARTINI DPPC membranes starting with interdigitated ripple-like states. (a) To test for spontaneous loss of kinetically trapped domains (yellow dashed rectangles), a larger 2204-lipid MARTINI2On DPPC membrane configuration with periodic ripples (top and front views) was created by repeating smaller rippled membranes (from time point IV, Fig. 2d) six-times across periodic boundary conditions (thrice in X, twice in Y). **(b)** 8 µs MD at 280 K almost completely abolished the interdigitated ripple-like configuration and led to the establishment of the L_β phase (Voronoi-thickness maps in Fig. S10a). Similar loss of the ripple-like domains was seen for large MARTINI2n **(c),** and MARTINI3b **(d),** membranes (Voronoi-thickness maps in Fig. S10b, c).

3.5 The optimized MARTINI DPPC FF (MARTINI2On) forms the $L_{\beta'}$, but not $P_{\beta'}$, phase

DPPC, with its large headgroup and strong head-head interactions, favours a tilted gel ($L_{\beta'}$) phase at lower temperatures. Therefore, during the $L_\alpha \rightarrow L_{\beta'}$ transition upon simulated cooling, the tilted ripple ($P_{\beta'}$) state occurs. Lipids exhibiting an L_β state at lower temperatures, such as phosphatidylethanolamines[74], show a single phase transition and do not form the ripple phase[20]. In our small membrane simulations, MARTINI2n (Fig. S2) and MARTINI3b (Fig. S4) froze into L_β states below transition temperatures, in accord with previous studies[34,47]. (MARTINI2n was shown to sample the $L_{\beta'}$ phase only upon enhanced sampling[75].) This may explain to some extent their inability to capture the $P_{\beta'}$ phase. However, MARTINI2On, possessing optimized DPPC bond parameters that closely match macroscopic membrane properties from all-atom simulations, spontaneously captured the $L_{\beta'}$ phase in many of the small membrane replicates upon cooling with classical MD simulations (Fig. S3). This is in agreement with the previous study[30] that developed the MARTINI2On DPPC parameters, where the authors also observed the formation of the $L_{\beta'}$ phase without enhanced sampling. Even in replicates that formed interdigitated ripple-like configurations (Fig. S3), the non-interdigitated gel-like domains often resembled the $L_{\beta'}$ phase.

To further contrast the lipid tilt in the gel phase configurations of DPPC MARTINI2n and MARTINI2On membranes, we chose representative replicates (#7 in Fig. S2, #3 and #7 in Fig. S3) that had frozen into gel states after SAMD without sampling interdigitated ripple-like states,

and further equilibrated them at 280 K for 5 µs in the NPT ensemble. The final MD snapshots for

MARTINI2n (#7 from Fig. S2) and MARTINI2On (#7 from Fig. S3) are illustrated in Fig. 6*a, b*.

Clearly, MARTINI2n and MARTINI2On form stable L_β and $L_{\beta'}$ configurations, respectively, at 280

K. For both systems, we quantified the extent of lipid tilt from the tilt angle (θ°) distributions of

lipids with their respective leaflet normal (Methods; Fig. 6*c*). The MARTINI2n and MARTINI2On

gel phase tilt distributions were observed to have modes at 7.3° and 15.2°, respectively.

To ensure that the MARTINI2On $L_{\beta'}$ configuration was not a consequence of finite size

artefacts, we created a 6-times-larger membrane by repeating the membrane from Fig. 6*b* across

periodic boundaries (3x times 2y; similar to Fig. 5). This initial configuration is shown in Fig. 6*d*,

and was simulated at 280 K for 5 µs. From the final snapshot (Fig. 6*d*), the $L_{\beta'}$ configuration is

completely preserved. Additionally, simulated cooling of a large L_α MARTINI2On membrane from

350 K – 280 K also resulted in a $L_{\beta'}$ configuration (Fig. 4*b*). These observations strongly suggest

that MARTINI2On membranes can sample the true $L_{\beta'}$ phase at lower temperatures, without any

finite size artefacts, and in agreement with previous studies[30]. However, our multi-replicate

simulations show that MARTINI2On membranes also stochastically sample the L_β phase, as

shown by replicates #2 and #8 in smaller membranes (Fig. S3), and from the simulations of large

membranes (Figs. 5*b* and S9*a, b*). Therefore, there may be room for further optimization of the

MARTINI2On DPPC FF for accurate reproduction of low temperature phase behaviour.

Interestingly, when replicate #3 from Fig. S3 was equilibrated for 5 µs at 280 K, it showed

the formation of a weakly interdigitated titled domain (Fig. S11*a*), a potential precursor for the

formation of a true ripple phase. To assess this further, we set up a 6-times larger membrane

(Fig. S11*b*) by repeating the smaller system across periodic boundaries along the membrane

plane (similar to Fig. 5*a*). After 10 µs of constant temperature EQMD at 280 K, the interdigitation

almost completely disappeared as shown by the final MD snapshot (Fig. S11*b*) and corresponding

Voronoi-thickness maps (Fig. S11*c*).

Fig. 6. Comparison of gel-phase structures of MARTINI membranes. Gel phase DPPC bilayer structures with **(a)** the standard MARTINI2n FF, and **(b)** the optimized MARTINI2On FF are shown. MARTINI2n membranes freeze into the straight gel phase (L_β) whereas MARTINI2On membranes often sample the tilted gel phase ($L_{\beta'}$), quantified by the lipid tilt-angle distributions **(c)**. **(d)** To confirm that this was a true thermodynamic phase and not a finite size effect, the smaller $L_{\beta'}$ membrane in (a) was replicated 6x (initial snapshot) and then equilibrated for 5 µs at 280 K. The membrane remained in the $L_{\beta'}$ phase thus ruling out finite size artefacts.

4 Summary and Conclusion

Molecular simulations at various levels of FF descriptions, ranging from all-atom and united atom models at the finer scales to DPD and MARTINI models at higher levels of coarse graining, have been extensively used to study the rich variety of equilibrium phases observed in model phospholipid bilayer membranes[1,17,76]. In particular, the rippled $P_{\beta'}$ phase, sensitive to initial configurations, annealing protocols, and finite-size effects, has been challenging to capture using MD simulations[20,25,34,41]. Although all-atom FFs have successfully captured the formation of the ripple phase (Table 1), the ability of the widely used MARTINI force-fields to stabilize this intermediate phase, which occurs below the main gel-to-liquid crystalline transition, has been a matter of some debate[34,47].

In this manuscript, we assessed the ability of MARTINI DPPC lipid membranes described by five MARTINI FF variants (see Introduction and Methods) towards capturing the ripple phase. Previous experiments[33,37–40] and all-atom simulations[19,30,43,44] have shown that DPPC membranes sample the tilted ripple phase at pre-transition temperatures. Previous MARTINI studies yielded conflicting observations, with some inferring that MARTINI DPPC lipid membranes could not capture the ripple state[34], while other studies observed otherwise[30,46,47]. Therefore, we established a standard simulated annealing-based MD protocol, built on insights from previous studies[18–20,30,43,44], for reliably producing the tilted ripple phase in multiple all-atom DPPC membrane MD replicates. By cooling from the fluid phase to below the phase transition followed by constant temperature equilibration. The structure of these rippled membranes resembled those from previous studies[19,20,43,44]. We then applied the same MD protocol to fluid phase MARTINI DPPC membranes with five MARTINI FF variants and ten independent MD replicates per variant. Simulations with all the five MARTINI FFs showed that a fraction of replicates formed interdigitated ripple-like states. These simulations also reinforced the importance of using multiple independent MD replicates for robustly observing phase behaviour[18,49]. The lateral orientation of

interdigitated domains in these ripple-like membranes were different compared to those from all-atom simulations and resembled a membrane with co-existing interdigitated L_α and gel phases.

Next, we explored whether the MARTINI ripple-like membranes were kinetically trapped states enforced by finite system size effects. To this end, we first re-established the all-atom protocols by employing the aforementioned MD cooling protocol on large all-atom DPPC membranes. We observed that these larger membranes also formed the tilted ripple phase with reticulated interdigitated domains, in agreement with previously reported large-membrane all-atom DPPC simulations[30,43]. However, when similarly large MARTINI DPPC membranes (3 FF variants, 2 replicates each) were cooled from the fluid phase, they failed to capture ripple-like states and froze directly into the gel phase . To confirm this behaviour via an orthogonal approach, we created larger membranes from previously formed smaller ripple-like membranes, by repeating them across periodic boundaries. These larger MARTINI periodic ripple-like membranes (3 FF variants), with an initial configuration bias towards the ripple phase, were then simulated at a constant temperature of 280 K, below the transition temperature. Strikingly, all the membranes showed a progressive loss of the interdigitated ripple-like domains, and either transitioned, or were in the process of transitioning, into the gel phase. Based on these findings we attribute the ripple-like states observed in the smaller MARTINI membranes to finite size artefacts. In the case of the MARTINI2On DPPC FF, the inability to form the tilted ripple was despite its ability to capture the tilted gel phase at lower temperatures without the need for enhanced sampling.

In summary, our observations systematically display that the current versions of MARTINI DPPC FFs may not be able to capture the $P_{\beta'}$ phase, which involves an intricate interplay of lipid tail interdigitation, tilt and headgroup hydration in the membrane. Specifically reparametrizing the MARTINI FFs to better reproduce membrane thermodynamics and phase behaviour is expected to yield CG models that may bridge the qualitative and quantitative gap towards all-atom MD

558 simulations, while unlocking the speed and scaling advantages associated with the versatile and

559 powerful MARTINI framework.

Acknowledgements

562 We thank the Supercomputer Education and Research Center (SERC) computational

563 facility at the Indian Institute of Science, Bengaluru, for access to supercomputing resources.

564 K.G.A. acknowledges funding by a grant from the Department of Science and Technology,

565 Government of India.

Author contributions

Conceptualization: R.D.; Data curation: P.S.; Formal analysis: R.D. and P.S.; Funding acquisition:

K.G.A.; Investigation: P.S. and R.D.; Methodology: R.D. and P.S.; Project administration: P.S.

and R.D.; Resources: P.S., R.D. and K.G.A.; Software (simulations, analysis code): P.S. and

R.D.; Supervision: K.G.A.; Validation: P.S., R.D. and K.G.A.; Visualization: P.S. and R.D.; Writing

– original draft: R.D. and P.S.; Writing – review & editing: R.D., P.S. and K.G.A.

All authors have given approval to the final version of the manuscript.

[‡]These authors (P.S. and R.D.) contributed equally.

Conflict of interest

The authors declare no conflicts of interest.

TOC/Abstract Graphic

References

(1)　Kulkarni, C. V. Lipid Crystallization: From Self-Assembly to Hierarchical and Biological
Ordering. *Nanoscale* **2012**, *4* (19), 5779–5791.

(2)　Nagle, J. F. Theory of the Main Lipid Bilayer Phase Transition. *Ann. Rev. Phys. Chem.*
1980, *31* (1), 157–196.

(3)　Harayama, T.; Riezman, H. Understanding the Diversity of Membrane Lipid Composition.
Nat. Rev. Mol. Cell Biol. **2018**, *19* (5), 281–296.

(4)　Kohli, A. G.; Kierstead, P. H.; Venditto, V. J.; Walsh, C. L.; Szoka, F. C. Designer Lipids
for Drug Delivery: From Heads to Tails. *J. Control. Release* **2014**, *190*, 274–287.

(5)　Anderson, M.; Omri, A. The Effect of Different Lipid Components on the in Vitro Stability
and Release Kinetics of Liposome Formulations. *Drug. Deliv.* **2004**, *11* (1), 33–39.

(6)　Kulkarni, A.; Chandrasekar, V.; Natarajan, S. K.; Ramesh, A.; Pandey, P.; Nirgud, J.;
Bhatnagar, H.; Ashok, D.; Ajay, A. K.; Sengupta, S. A Designer Self-Assembled
Supramolecule Amplifies Macrophage Immune Responses against Aggressive Cancer.
Nat. Biomed. Eng. **2018**, *2* (8), 589–599.

(7)　Phillips, R.; Ursell, T.; Wiggins, P.; Sens, P. Emerging Roles for Lipids in Shaping
Membrane-Protein Function. *Nature* **2009**, *459* (7245), 379–385.

(8)　Blicher, A.; Wodzinska, K.; Fidorra, M.; Winterhalter, M.; Heimburg, T. The Temperature
Dependence of Lipid Membrane Permeability, Its Quantized Nature, and the Influence of
Anesthetics. *Biophys. J.* **2009**, *96* (11), 4581–4591.

(9)　Sharma, P.; Parthasarathi, S.; Patil, N.; Waskar, M.; Raut, J. S.; Puranik, M.; Ayappa, K.
G.; Basu, J. K. Assessing Barriers for Antimicrobial Penetration in Complex Asymmetric
Bacterial Membranes: A Case Study with Thymol. *Langmuir* **2020**, *36* (30), 8800–8814.
https://doi.org/10.1021/acs.langmuir.0c01124.

(10) Perez-Isidoro, R.; Sierra-Valdez, F. J.; Ruiz-Suarez, J. C. Anesthetic Diffusion through
Lipid Membranes Depends on the Protonation Rate. *Sci. Rep.* **2014**, *4*, 7534.

(11) Yilmaz, N.; Kobayashi, T. Assemblies of Pore-Forming Toxins Visualized by Atomic Force
Microscopy. *Biochim. Biophys. Acta* **2016**, *1858* (3), 500–511.

(12) Rojko, N.; Anderluh, G. How Lipid Membranes Affect Pore Forming Toxin Activity. *Acc.*
Chem. Res. **2015**, *48* (12), 3073–3079.

(13) Gerelli, Y.; Porcar, L.; Lombardi, L.; Fragneto, G. Lipid Exchange and Flip-Flop in Solid
Supported Bilayers. *Langmuir* **2013**, *29* (41), 12762–12769.

(14) Pomorski, T. G.; Menon, A. K. Lipid Somersaults: Uncovering the Mechanisms of Protein-
Mediated Lipid Flipping. *Prog. Lipid Res.* **2016**, *64*, 69–84.

(15) Andersen, S. S.; Jackson, A. D.; Heimburg, T. Towards a Thermodynamic Theory of
Nerve Pulse Propagation. *Prog. Neurobiol.* **2009**, *88* (2), 104–113.

(16) de Smet, M.; Langereis, S.; van den Bosch, S.; Grull, H. Temperature-Sensitive
Liposomes for Doxorubicin Delivery under MRI Guidance. *J. Control. Release* **2010**, *143*
(1), 120–127.

(17) Enkavi, G.; Javanainen, M.; Kulig, W.; Róg, T.; Vattulainen, I. Multiscale Simulations of
Biological Membranes: The Challenge To Understand Biological Phenomena in a Living
Substance. *Chem. Rev.* **2019**, *119* (9), 5607–5774.
https://doi.org/10.1021/acs.chemrev.8b00538.

(18) Sun, L.; Böckmann, R. A. Membrane Phase Transition during Heating and Cooling:
Molecular Insight into Reversible Melting. *European Biophysics Journal* **2018**, *47* (2), 151–
164.

(19) Vries, A. H. de; Yefimov, S.; Mark, A. E.; Marrink, S. J. Molecular Structure of the Lecithin
Ripple Phase. *PNAS* **2005**, *102* (15), 5392–5396.
https://doi.org/10.1073/pnas.0408249102.

(20) Lenz, O.; Schmid, F. Structure of Symmetric and Asymmetric ``Ripple'' Phases in Lipid
Bilayers. *Phys. Rev. Lett.* **2007**, *98* (5), 058104.
https://doi.org/10.1103/PhysRevLett.98.058104.

(21) Kranenburg, M.; Smit, B. Phase Behavior of Model Lipid Bilayers. *J. Phys. Chem. B* **2005**,
109 (14), 6553–6563. https://doi.org/10.1021/jp0457646.

(22) Marrink, S. J.; Tieleman, D. P. Perspective on the Martini Model. *Chem. Soc. Rev.* **2013**,
42 (16), 6801–6822.

(23) Marrink, S. J.; Risselada, H. J.; Yefimov, S.; Tieleman, D. P.; de Vries, A. H. The
MARTINI Force Field: Coarse Grained Model for Biomolecular Simulations. *The Journal of*
Physical Chemistry B **2007**, *111* (27), 7812–7824.

(24) Bruininks, B. M. H.; Souza, P. C. T.; Marrink, S. J. A Practical View of the Martini Force
Field. In *Biomolecular Simulations: Methods and Protocols*; Bonomi, M., Camilloni, C.,
Eds.; Methods in Molecular Biology; Springer: New York, NY, 2019; pp 105–127.
https://doi.org/10.1007/978-1-4939-9608-7_5.

(25) Srivastava, A.; Debnath, A. Asymmetry and Rippling in Mixed Surfactant Bilayers from All-
Atom and Coarse-Grained Simulations: Interdigitation and Per Chain Entropy†. *The*
Journal of Physical Chemistry B **2020**. https://doi.org/10.1021/acs.jpcb.0c03761.

(26) Vaiwala, R.; Sharma, P.; Puranik, M.; Ayappa, K. G. Developing a Coarse-Grained Model
for Bacterial Cell Walls: Evaluating Mechanical Properties and Free Energy Barriers. *J.*
Chem. Theory Comput. **2020**, *16* (8), 5369–5384.
https://doi.org/10.1021/acs.jctc.0c00539.

(27) Ingólfsson, H. I.; Melo, M. N.; van Eerden, F. J.; Arnarez, C.; Lopez, C. A.; Wassenaar, T.
A.; Periole, X.; de Vries, A. H.; Tieleman, D. P.; Marrink, S. J. Lipid Organization of the
Plasma Membrane. *J. Am. Chem. Soc.* **2014**, *136* (41), 14554–14559.
https://doi.org/10.1021/ja507832e.

(28) Alessandri, R.; Souza, P. C. T.; Thallmair, S.; Melo, M. N.; de Vries, A. H.; Marrink, S. J.
Pitfalls of the Martini Model. *J. Chem. Theory Comput.* **2019**, *15* (10), 5448–5460.
https://doi.org/10.1021/acs.jctc.9b00473.

(29) Desikan, R.; Patra, S. M.; Sarthak, K.; Maiti, P. K.; Ayappa, K. G. Comparison of Coarse-
Grained (MARTINI) and Atomistic Molecular Dynamics Simulations of α and β Toxin
Nanopores in Lipid Membranes. *J. Chem. Sci.* **2017**, *129* (7), 1017–1030.
https://doi.org/10.1007/s12039-017-1316-0.

(30) Carpenter, T. S.; López, C. A.; Neale, C.; Montour, C.; Ingólfsson, H. I.; Di Natale, F.;
Lightstone, F. C.; Gnanakaran, S. Capturing Phase Behavior of Ternary Lipid Mixtures
with a Refined Martini Coarse-Grained Force Field. *J. Chem. Theory Comput.* **2018**, *14*
(11), 6050–6062. https://doi.org/10.1021/acs.jctc.8b00496.

(31) Globisch, C.; Krishnamani, V.; Deserno, M.; Peter, C. Optimization of an Elastic Network
Augmented Coarse Grained Model to Study CCMV Capsid Deformation. *PLoS ONE*
2013, *8* (4), e60582.

(32) Biltonen, R. L.; Lichtenberg, D. The Use of Differential Scanning Calorimetry as a Tool to
Characterize Liposome Preparations. *Chem. Phys. Lipids* **1993**, *64* (1), 129–142.

(33) Tardieu, A.; Luzzati, V.; Reman, F. C. Structure and Polymorphism of the Hydrocarbon
Chains of Lipids: A Study of Lecithin-Water Phases. *Journal of Molecular Biology* **1973**, *75*
(4), 711–733. https://doi.org/10.1016/0022-2836(73)90303-3.

(34) Rodgers, J. M.; Sorensen, J.; de Meyer, F. J.; Schiott, B.; Smit, B. Understanding the
Phase Behavior of Coarse-Grained Model Lipid Bilayers through Computational
Calorimetry. *J. Phys. Chem. B* **2012**, *116* (5), 1551–1569.

(35) Riske, K. A.; Barroso, R. P.; Vequi-Suplicy, C. C.; Germano, R.; Henriques, V. B.; Lamy,
M. T. Lipid Bilayer Pre-Transition as the Beginning of the Melting Process. *Biochimica et*

Biophysica Acta (BBA) - Biomembranes **2009**, *1788* (5), 954–963.
https://doi.org/10.1016/j.bbamem.2009.01.007.

(36) Heimburg, T. A Model for the Lipid Pretransition: Coupling of Ripple Formation with the
Chain-Melting Transition. *Biophys. J.* **2000**, *78* (3), 1154–1165.
https://doi.org/10.1016/S0006-3495(00)76673-2.

(37) Sun, W. J.; Tristram-Nagle, S.; Suter, R. M.; Nagle, J. F. Structure of the Ripple Phase in
Lecithin Bilayers. *PNAS* **1996**, *93* (14), 7008–7012.
https://doi.org/10.1073/pnas.93.14.7008.

(38) Katsaras, J.; Tristram-Nagle, S.; Liu, Y.; Headrick, R. L.; Fontes, E.; Mason, P. C.; Nagle,
J. F. Clarification of the Ripple Phase of Lecithin Bilayers Using Fully Hydrated, Aligned
Samples. *Phys. Rev. E* **2000**, *61* (5), 5668–5677.
https://doi.org/10.1103/PhysRevE.61.5668.

(39) Sengupta, K.; Raghunathan, V. A.; Katsaras, J. Structure of the Ripple Phase of
Phospholipid Multibilayers. *Phys. Rev. E* **2003**, *68* (3), 031710.
https://doi.org/10.1103/PhysRevE.68.031710.

(40) Kamal, M. A.; Raghunathan, V. Naphthalene Derivatives Induce Acyl Chain Interdigitation
in Dipalmitoylphosphatidylcholine Bilayers. *The Journal of Physical Chemistry B* **2016**,
120 (1), 164–172.

(41) Debnath, A.; Thakkar, F. M.; Maiti, P. K.; Kumaran, V.; Ayappa, K. G. Laterally Structured
Ripple and Square Phases with One and Two Dimensional Thickness Modulations in a
Model Bilayer System. *Soft Matter* **2014**, *10* (38), 7630–7637.
https://doi.org/10.1039/C4SM01031K.

(42) Pluhackova, K.; Kirsch, S. A.; Han, J.; Sun, L.; Jiang, Z.; Unruh, T.; Böckmann, R. A. A
Critical Comparison of Biomembrane Force Fields: Structure and Dynamics of Model
DMPC, POPC, and POPE Bilayers. *J. Phys. Chem. B* **2016**, *120* (16), 3888–3903.
https://doi.org/10.1021/acs.jpcb.6b01870.

(43) Kirsch, S. A.; Böckmann, R. A. Coupling of Membrane Nanodomain Formation and
Enhanced Electroporation near Phase Transition. *Biophysical Journal* **2019**, *116* (11),
2131–2148. https://doi.org/10.1016/j.bpj.2019.04.024.

(44) Khakbaz, P.; Klauda, J. B. Investigation of Phase Transitions of Saturated
Phosphocholine Lipid Bilayers via Molecular Dynamics Simulations. *Biochimica et
Biophysica Acta (BBA) - Biomembranes* **2018**, *1860* (8), 1489–1501.
https://doi.org/10.1016/j.bbamem.2018.04.014.

(45) Debnath, A.; Ayappa, K. G.; Kumaran, V.; Maiti, P. K. The Influence of Bilayer
Composition on the Gel to Liquid Crystalline Transition. *J. Phys. Chem. B* **2009**, *113* (31),
10660–10668. https://doi.org/10.1021/jp901551d.

(46) Lin, X.; Gu, N. Surface Properties of Encapsulating Hydrophobic Nanoparticles Regulate
the Main Phase Transition Temperature of Lipid Bilayers: A Simulation Study. *Nano Res.*
2014, *7* (8), 1195–1204. https://doi.org/10.1007/s12274-014-0482-3.

(47) Stelter, D.; Keyes, T. Enhanced Sampling of Phase Transitions in Coarse-Grained Lipid Bilayers. *J. Phys. Chem. B* **2017**, *121* (23), 5770–5780. https://doi.org/10.1021/acs.jpcb.6b11711.

(48) Yesylevskyy, S. O.; Schäfer, L. V.; Sengupta, D.; Marrink, S. J. Polarizable Water Model for the Coarse-Grained MARTINI Force Field. *PLoS Comput Biol* **2010**, *6* (6), 1–17.

(49) Grossfield, A.; Patrone, P. N.; Roe, D. R.; Schultz, A. J.; Siderius, D.; Zuckerman, D. M. Best Practices for Quantification of Uncertainty and Sampling Quality in Molecular Simulations [Article v1.0]. *Living Journal of Computational Molecular Science* **2018**, *1* (1), 5067. https://doi.org/10.33011/livecoms.1.1.5067.

(50) Jo, S.; Kim, T.; Iyer, V. G.; Im, W. CHARMM-GUI: A Web-Based Graphical User Interface for CHARMM. *J Comput Chem* **2008**, *29* (11), 1859–1865. https://doi.org/10.1002/jcc.20945.

(51) Lee, J.; Cheng, X.; Swails, J. M.; Yeom, M. S.; Eastman, P. K.; Lemkul, J. A.; Wei, S.; Buckner, J.; Jeong, J. C.; Qi, Y.; Jo, S.; Pande, V. S.; Case, D. A.; Brooks, C. L.; MacKerell, A. D.; Klauda, J. B.; Im, W. CHARMM-GUI Input Generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM Simulations Using the CHARMM36 Additive Force Field. *J. Chem. Theory Comput.* **2016**, *12* (1), 405–413. https://doi.org/10.1021/acs.jctc.5b00935.

(52) Wu, E. L.; Cheng, X.; Jo, S.; Rui, H.; Song, K. C.; Dávila-Contreras, E. M.; Qi, Y.; Lee, J.; Monje-Galvan, V.; Venable, R. M.; Klauda, J. B.; Im, W. CHARMM-GUI Membrane Builder toward Realistic Biological Membrane Simulations. *J Comput Chem* **2014**, *35* (27), 1997–2004. https://doi.org/10.1002/jcc.23702.

(53) Meyer, H. W. Pretransition-Ripples in Bilayers of Dipalmitoylphosphatidylcholine: Undulation or Periodic Segments? A Freeze-Fracture Study. *Biochimica et Biophysica Acta (BBA) - Lipids and Lipid Metabolism* **1996**, *1302* (2), 138–144. https://doi.org/10.1016/0005-2760(96)00054-9.

(54) Pastor, R. W.; MacKerell, A. D. Development of the CHARMM Force Field for Lipids. *J. Phys. Chem. Lett.* **2011**, *2* (13), 1526–1532. https://doi.org/10.1021/jz200167q.

(55) Klauda, J. B.; Venable, R. M.; Freites, J. A.; O'Connor, J. W.; Tobias, D. J.; Mondragon-Ramirez, C.; Vorobyov, I.; MacKerell, A. D.; Pastor, R. W. Update of the CHARMM All-Atom Additive Force Field for Lipids: Validation on Six Lipid Types. *J. Phys. Chem. B* **2010**, *114* (23), 7830–7843.

(56) Abraham, M. J.; Murtola, T.; Schulz, R.; Páll, S.; Smith, J. C.; Hess, B.; Lindahl, E. GROMACS: High Performance Molecular Simulations through Multi-Level Parallelism from Laptops to Supercomputers. *SoftwareX* **2015**, *1*, 19–25.

(57) Braun, E.; Gilmer, J.; Mayes, H. B.; Mobley, D. L.; Monroe, J. I.; Prasad, S.; Zuckerman, D. M. Best Practices for Foundations in Molecular Simulations [Article v1.0]. *Living Journal of Computational Molecular Science* **2018**, *1* (1), 5957. https://doi.org/10.33011/livecoms.1.1.5957.

758 (58) Smith, D. J.; Klauda, J. B.; Sodt, A. J. Simulation Best Practices for Lipid Membranes
759    [Article v1.0]. *Living Journal of Computational Molecular Science* **2018**, *1* (1), 5966.
760    https://doi.org/10.33011/livecoms.1.1.5966.

761 (59) Lemkul, J. From Proteins to Perturbed Hamiltonians: A Suite of Tutorials for the
762    GROMACS-2018 Molecular Simulation Package [Article v1.0]. *Living Journal of*
763    *Computational Molecular Science* **2018**, *1* (1), 5068.
764    https://doi.org/10.33011/livecoms.1.1.5068.

765 (60) Desikan, R.; Behera, A.; Maiti, P. K.; Ayappa, K. G. Using Multiscale Molecular Dynamics
766    Simulations to Obtain Insights into Pore Forming Toxin Mechanisms. *arXiv:2010.13854*
767    *[cond-mat]* **2020**.

768 (61) Darden, T.; York, D.; Pedersen, L. Particle Mesh Ewald: An N.Log(N) Method for Ewald
769    Sums in Large Systems. *J. Chem. Phys.* **1993**, *98* (12), 10089–10092.

770 (62) Hess, B. P-LINCS: A Parallel Linear Constraint Solver for Molecular Simulation. *J. Chem.*
771    *Theory. Comput.* **2008**, *4* (1), 116–122.

772 (63) Bussi, G.; Donadio, D.; Parrinello, M. Canonical Sampling through Velocity Rescaling. *The*
773    *Journal of chemical physics* **2007**, *126* (1), 014101.

774 (64) Parrinello, M.; Rahman, A. Polymorphic Transitions in Single Crystals: A New Molecular
775    Dynamics Method. *Journal of Applied Physics* **1981**, *52* (12), 7182–7190.

776 (65) Alavi, S.; Thompson, D. L. Molecular Dynamics Studies of Melting and Some Liquid-State
777    Properties of 1-Ethyl-3-Methylimidazolium Hexafluorophosphate [Emim][PF6]. *J. Chem.*
778    *Phys.* **2005**, *122* (15), 154704.

779 (66) Zhang, Y.; Maginn, E. J. A Comparison of Methods for Melting Point Calculation Using
780    Molecular Dynamics Simulations. *J. Chem. Phys.* **2012**, *136* (14), 144116.
781    https://doi.org/10.1063/1.3702587.

782 (67) Wassenaar, T. A.; Ingólfsson, H. I.; Böckmann, R. A.; Tieleman, D. P.; Marrink, S. J.
783    Computational Lipidomics with Insane: A Versatile Tool for Generating Custom
784    Membranes for Molecular Simulations. *Journal of chemical theory and computation* **2015**,
785    *11* (5), 2144–2155.

786 (68) de Jong, D. H.; Baoukina, S.; Ingólfsson, H. I.; Marrink, S. J. Martini Straight: Boosting
787    Performance Using a Shorter Cutoff and GPUs. *Comput. Phys. Commun.* **2016**, *199*, 1–7.
788    https://doi.org/10.1016/j.cpc.2015.09.014.

789 (69) Tironi, I. G.; Sperb, R.; Smith, P. E.; van Gunsteren, W. F. A Generalized Reaction Field
790    Method for Molecular Dynamics Simulations. *The Journal of chemical physics* **1995**, *102*
791    (13), 5451–5459.

792 (70) Humphrey, W.; Dalke, A.; Schulten, K. VMD: Visual Molecular Dynamics. *Journal of*
793    *Molecular Graphics* **1996**, *14* (1), 33–38.

794   (71)   Guixà-González, R.; Rodriguez-Espigares, I.; Ramírez-Anguita, J. M.; Carrió-Gaspar, P.;
795   Martinez-Seara, H.; Giorgino, T.; Selent, J. MEMBPLUGIN: Studying Membrane
796   Complexity in VMD. *Bioinformatics* **2014**, *30* (10), 1478–1480.
797   https://doi.org/10.1093/bioinformatics/btu037.

798   (72)   Varadarajan, V.; Desikan, R.; Ayappa, K. G. Assessing the Extent of the Structural and
799   Dynamic Modulation of Membrane Lipids Due to Pore Forming Toxins: Insights from
800   Molecular Dynamics Simulations. *Soft Matter* **2020**, *16* (20), 4840–4857.
801   https://doi.org/10.1039/D0SM00086H.

802   (73)   Souza, P. C. T.; Thallmair, S.; Conflitti, P.; Ramírez-Palacios, C.; Alessandri, R.; Raniolo,
803   S.; Limongelli, V.; Marrink, S. J. Protein–Ligand Binding with the Coarse-Grained Martini
804   Model. *Nature Communications* **2020**, *11* (1), 3714. https://doi.org/10.1038/s41467-020-
805   17437-5.

806   (74)   Koynova, R.; Caffrey, M. Phases and Phase Transitions of the Hydrated
807   Phosphatidylethanolamines. *Chemistry and Physics of Lipids* **1994**, *69* (1), 1–34.
808   https://doi.org/10.1016/0009-3084(94)90024-8.

809   (75)   Nagai, T.; Okamoto, Y. Replica-Exchange Molecular Dynamics Simulation of a Lipid
810   Bilayer System with a Coarse-Grained Model. *Molecular Simulation* **2012**, *38* (5), 437–
811   441. https://doi.org/10.1080/08927022.2011.564172.

812   (76)   Manna, M.; Nieminen, T.; Vattulainen, I. Understanding the Role of Lipids in Signaling
813   Through Atomistic and Multiscale Simulations of Cell Membranes. *Annu. Rev. Biophys.*
814   **2019**, *48* (1), 421–439. https://doi.org/10.1146/annurev-biophys-052118-115553.

Supplementary information

Evaluating coarse-grained MARTINI force-fields for capturing the ripple

phase of lipid membranes

Pradyumn Sharma[1,‡], Rajat Desikan[1,‡,†], K. Ganapathy Ayappa[1,2,]*

[1]Department of Chemical Engineering, Indian Institute of Science, Bengaluru, India

[2]Centre for BioSystems Science and Engineering, Indian Institute of Science, Bengaluru, India

[‡]Co-first authors

[†]Present address: Certara QSP, Certara UK Limited, Sheffield, UK

[*]Corresponding author: K. Ganapathy Ayappa; E-mail: ayappa@iisc.ac.in

Fig. S1. An independent all-atom MD replicate reproduces the ripple phase. (a) MD snapshot at time point IV from Fig. 1*a* is periodically repeated along the x-axis, and the major and minor arms of the ripple in the upper leaflet are visually annotated for clarity. **(b)** Final snapshot (time point IV in Fig. 1) and corresponding Voronoi-thickness diagram from the independent simulated cooling replicate, using the same MD protocol as for the replicate in Fig. 1, shows robust reproduction of the tilted ripple phase from all-atom simulations.

Fig. S2. Summary of 10 independent cooling MD replicates with the MARTINI2n FF. Temporal evolution of the minimum inter-leaflet P-P distance (ξ), a measure of the extent of interdigitation (Fig. 1b; $m_2 < 0$ – interdigitation, and $m_2 > 0$ – no interdigitation (Methods, Eq. 1), are highlighted in replicates 1 and 2, respectively). Final snapshots corresponding to time point III at the end of cooling simulations with MARTINI2n DPPC membranes are shown for all the 10 independent MD replicates. The interdigitated domains in replicates which exhibit ripple-like structures are marked with yellow dashed circles.

Fig. S3. Summary of 10 independent cooling MD replicates with the MARTINI2On FF. Same as Fig. S2, but with the MARTINI2On FF variant.

Fig. S4. Summary of 10 independent cooling MD replicates with the MARTINI3b FF. Same as Fig. S2, but with the MARTINI3b FF variant.

Fig. S5. Summary of 10 independent cooling MD replicates with the MARTINI2p FF. Same as Fig. S2, but with the MARTINI2p FF variant. Note that while none of the ξ plots show ripple-like domain formation during the cooling phase, replicate 5 formed shallow interdigitated ripple-like domains during the 5 µs EQMD (Fig. 2*f*).

Fig. S6. Summary of 10 independent cooling MD replicates with the MARTINI2Op FF. Same as Fig. S2, but with the MARTINI2Op FF variant.

Fig. S7. Independent all-atom simulation replicate also shows ripple formation in a large DPPC membrane. Similar to Fig. 3, the final snapshot (front view) of a large DPPC membrane after being cooled to 290 K, and its corresponding Voronoi-thickness map, show the formation of symmetric, and asymmetric-interdigitated, ripple structures along the X-axis (shorter box dimension) and Y-axis (shorter box dimension), respectively. Note that the non-interdigitated symmetric ripple along the X-axis is indiscernible from the Voronoi-thickness map.

Fig. S8. Periodically repeated Voronoi-thickness modulation heatmaps of atomistic DPPC membranes. Voronoi-thickness modulation heatmaps after simulated cooling of atomistic **(a)** 368-lipid (Fig. 1), and **(b)** 2208-lipid (Fig. 3) DPPC membranes are shown with periodic images (original simulation box in the middle shown with a red border; same thickness colorbar applies for both systems). The reticulated interdigitated domains in the larger membrane contrast with the island-like isolated domains in the smaller membrane.

a MARTINI2n, after cooling a 2204-lipid rectangular membrane from 350 K to 280 K

MARTINI2On

MARTINI3b

b MARTINI2On, a 6728-lipid square L_α membrane, equilibrated at 350 K for 100 ns

Replicate 1, after cooling to 280 K

Replicate 2, after cooling to 280 K

Fig. S9 Independent MD replicates also exhibit lack of ripple formation in large MARTINI DPPC membranes. (a) Independent MD replicates similar to Fig. 4, where large, rectangular, MARTINI2n, MARTINI2On and MARTINI3b DPPC membranes, initially pre-equilibrated in the fluid phase at 350 K, exhibited direct transformation into the straight gel phase upon cooling to 280 K without forming either interdigitated or symmetric ripple-like structures (unlike the behaviour for smaller MARTINI membranes in Fig. 2). **(b)** The influence of membrane geometry was also tested by cooling two large, square, MARTINI2On membrane replicates, from 350 K to 280 K. Similar to the rectangular membrane patches, both square membrane replicates directly froze into the gel phase without forming ripple-like structures.

Fig. S10. Voronoi-thickness maps from MD of large MARTINI membranes. Voronoi-thickness map from constant temperature MD of large **(a)** MARTINI2On, **(b)** MARTINI2n, and **(c)** MARTINI3b membranes at 280 K, corresponding to MD snapshots in Fig. 5. Simulations starting with a periodic ripple-like configuration (0 µs), shows progressive loss of the interdigitated ripple-like domains and formation of the gel phase by 8 µs and 10 8 µs for the MARTINI2On and MARTINI3b membranes, respectively. Notably, the MARTINI2n membrane shows complete loss of interdigitation by 3 µs. Note that while the blue regions at the initial time points (0 µs) correspond to interdigitated domains, they could correspond to either interdigitated, or non-interdigitated but partially melted domains, at the final time points. The distinction can be made by simultaneously comparing the MD snapshots (Fig. 5) with the Voronoi-thickness maps above.

Fig. S11. Tilted ripple-like state of MARTINI2On membranes is also a finite size artefact.
Simulated cooling of replicate 3 with the MARTINI2On FF (from Fig. S3) showed the formation of an interdigitated domain in an otherwise $L_{\beta'}$ membrane at 280 K. This structure was enhanced upon performing constant temperature MD for 5000 ns: final snapshot shown in **(a)**. **(b)** To explore whether this was a finite size effect or a true ripple phase, the smaller membrane was replicated 6x across periodic boundaries to create a larger membrane with repeated ripple-like domains (snapshot at 0 µs), similar to Fig. 5*a*. After 10 µs constant temperature MD, the interdigitated ripple-like regions were greatly diminished, thus confirming that the ripple-like state in (a) was indeed a finite size artefact. **(c)** Voronoi-thickness maps pertaining to MD snapshots in (b), where regions of lower thickness in blue indicate the ripple-like domains.