DNA transduction in *Sodalis* species: implications for the genetic

modification of uncultured endosymbionts of insects

Chelsea M. Keller, Christopher G. Kendra, Roberto E. Bruna, David Craft, Mauricio H.

Pontes*

Department of Pathology and Laboratory Medicine, Department of Microbiology and

Immunology, Pennsylvania State University College of Medicine, Hershey, PA 17033,

USA.

*Corresponding author:

Mauricio H. Pontes

Penn State College of Medicine

Departments of Pathology, and

Microbiology and Immunology

500 University Drive, C6818A

Hershey, PA 17033

mpontes@pennstatehealth.psu.edu

717-531-0003 ext. 320524

Running title: Bacteriophage P1-mediated transduction in *Sodalis*

Abstract

Bacteriophages (phages) are ubiquitous in nature. These viruses play a number of

central roles in microbial ecology and evolution by, for instance, promoting horizontal

gene transfer (HGT) among bacterial species. The ability of phages to mediate HGT

through transduction has been widely exploited as an experimental tool for the genetic

study of bacteria. As such, bacteriophage P1 represents a prototypical generalized

transducing phage with a broad host range that has been extensively employed in the

genetic manipulation of *Escherichia coli* and a number of other model bacterial species.

Here we demonstrate that P1 is capable of infecting, lysogenizing and promoting

transduction in members of the bacterial genus *Sodalis*, including the maternally

inherited insect endosymbiont *Sodalis glossinidius*. While establishing new tools for the

genetic study of these bacterial species, our results suggest that P1 may be used to

deliver DNA to many Gram negative endosymbionts in their insect host, thereby

circumventing a culturing requirement to genetically manipulate these organisms.

Summary

A large number of economically important insects maintain intimate associations with

maternally inherited endosymbiotic bacteria. Due to the inherit nature of these

associations, insect endosymbionts cannot be usually isolated in pure culture nor

genetically manipulated. Here we use a broad-host range bacteriophage to deliver

exogenous DNA to an insect endosymbiont and a closely related free-living species.

Our results suggest that broad host range bacteriophages can be used to genetically

alter insect endosymbionts in their insect host and, as a result, bypass a culturing

requirement to genetically alter these bacteria.

Introduction

Bacteriophages (phages) are the most abundant and diverse biological entities on the planet. With an estimated population size greater than 1×10^{31} (Hendrix *et al*. 1999), these bacterial viruses play essential ecological and evolutionary functions. Phages control the size of bacterial populations and shape the diversity of microbial communities by modulating the abundance of bacterial lineages and promoting, directly and indirectly, the exchange of genetic information among species (Touchon *et al*., 2017; Soucy *et al*. 2015). Historically, phages have played a central role in the development of molecular biology, enabling, for instance, the establishment of DNA as the genetic material of living cells (Hershey and Chase, 1952). Today, phages are widely used as tools in the study of bacteria. For instance, generalized transducing phages such as P1 allow the rapid transfer of DNA among bacterial strains, greatly facilitating genetic dissection of biological processes (Thomason *et al*. 2007).

P1 is a temperate bacteriophage capable of alternating between lytic and lysogenic infection. P1 was initially described in studies involving lysogenic strains of *Escherichia coli* (Bertani, 1951). This phage is capable of mediating generalized transduction (Lennox, 1955), a property that has fostered its adoption as an important experimental tool for the genetic analysis and manipulation of *E. coli* (Thomason *et al*. 2007, Yarmolinsky and Sternberg, 1988). Notably, in addition to its habitual *E. coli* host, P1 can also infect a large number of Gram negative bacterial species (Yarmolinsky and Sternberg, 1988; Goldberg *et al*., 1974; Ornellas and Stocker, 1974; Kaiser and Dworkin, 1975; Murooka and Harada, 1979). This broad host range, along with its well-characterized molecular biology and established experimental procedures, has prompted the use of this phage as an experimental tool for the delivery of DNA to a large number of bacterial species (Streicher *et al*., 1971; O'Connor *et al*., 1983; Downard,

1988; Wolf-Watz *et al.*, 1985; Westwater *et al.*, 2002; Butela and Lawrence, 2012). Here

we establish that P1 is capable of infecting two members of the bacterial genus *Sodalis*,

including *Sodalis glossinidius* (Dale and Maudlin, 1999; Chari *et al.*, 2015).

Sodalis glossinidius is a maternally inherited, Gram negative bacterial

endosymbiont of tsetse flies (*Glossina* spp.; Diptera: *Glossinidae*). Similar to other insect

endosymbionts, *S. glossinidius* exists in a stable, chronic association with its insect host,

and undergoes a predominantly maternal mode of transmission (Aksoy *et al.*, 1997;

Cheng and Aksoy, 1997; De Vooght *et al.*, 2015). Notably, like other insect

endosymbionts, this bacterium has undergone an extensive process of genome

degeneration as a result of a recent evolutionary transition from free-living existence to

permanent host association (McCutcheon *et al.*, 2019; Moran *et al.*, 2008). Because this

process is accompanied by the loss of metabolic capability and stress response

pathways (McCutcheon *et al.*, 2019; Moran *et al.*, 2008; Toh *et al.*, 2006; Pontes *et al.* 2011;

Clayton *et al.*, 2012; Pontes and Dale, 2006), *S. glossinidius* has proven refractory to harsh

artificial DNA transformation procedures that are commonly employed in model

organisms such as *Escherichia coli* (Pontes and Dale, 2006). Consequently, this bacterium

has remained genetically intractable (Kendra *et al.* 2020).

In this study, we demonstrate that the bacteriophage P1 is capable of infecting,

lysogenizing and promoting transduction in *Sodalis glossinidius*, and its free-living close

relative, the plant-associated and opportunistic pathogen *Sodalis praecaptivus* (Dale and

Maudlin, 1999; Chari *et al.*, 2015). We demonstrate that P1 can be used to mediate

generalized transduction of chromosomal and extrachromosomal DNA in *S.*

praecaptivus. We use P1 to transduce autonomous replicating phagemids containing an

array of reporter genes and Tn*7* transposition systems harboring fluorescent proteins

for chromosomal tagging. Finally, we developed a suicide phagemid containing a

mariner transposase for random mutagenesis of bacterial strains susceptible to P1

infection. This study establishes a new efficient method for genetic manipulation of

Sodalis species (Fig. 1) that can be readily adapted to other Gram negative bacteria.

Furthermore, these results provide a potential means for the genetic modification of

bacterial endosymbionts, in their insect host, through the use of P1 as a DNA-delivery

system.

Results

Bacteriophage P1 infects, lysogenizes and forms phage particles in *Sodalis*

glossinidius* and *Sodalis praecaptivus

P1CM*clr*-100(ts) is a thermo-inducible P1 variant harboring a chloramphenicol resistant

marker. P1CM*clr*-100(ts) forms chloramphenicol resistant lysogens at low temperatures

(≤30°C) but produces phage particles at higher temperatures (≥37°C) (Rosner, 1972).

Consequently, infection of *E. coli* by P1CM*clr*-100(ts) yields chloramphenicol resistant

lysogens at 30°C. We took advantage of these P1CM*clr*-100(ts) properties to test if *S.*

glossinidius and *S. praecaptivus* were susceptible to P1 infection. We exposed cultures of

these bacteria to increasing concentrations of P1CM*clr*-100(ts) phage particles and

subsequently plated dilutions on solid medium containing chloramphenicol. We

established that, similar to the *E. coli* control (Fig. 2A), exposure to increasing

concentrations of P1CM*clr*-100(ts) particles yielded increasing numbers of

chloramphenicol resistant colonies in both *S. glossinidius* (Fig. 2B) and *S. praecaptivus*

(Fig. 2C). Importantly, no chloramphenicol resistant colonies were observed in cultures

that were not exposed to P1CM*clr*-100(ts) particles (Fig. 2B and C).

That these colonies were P1 lysogens, as opposed to recombinants harboring

only the P1-derived chloramphenicol resistant marker, was supported by several lines

of evidence. First, the presence of a P1 DNA fragment was detected by polymerase

chain reaction (PCR) in both chloramphenicol resistant *S. glossinidius* (Fig. 3A and B)

and *S. praecaptivus* clones (Fig. 3C and D), but not in the wild-type strains (Fig. 3A-D,

left side). This indicated that chloramphenicol resistant cells harbor at least part of the

P1CM*clr*-100(ts) genome. Second, lysates prepared from *S. glossinidius* and *S.*

praecaptivus chloramphenicol resistant clones, but not their wild-type counterparts,

formed plaques in soft agar cultures of *E. coli* grown at 37°C, a temperature that induces

P1CM*clr*-100(ts) lytic replication (Rosner, 1972) (Fig. 3E and F). This indicated that

chloramphenicol resistant *S. glossinidius* and *S. praecaptivus* clones can produce phage

particles that are lytic to *E. coli* grown at 37°C. Third, the lytic activity of lysates derived

from chloramphenicol resistant *S. praecaptivus* cultures propagated at 37°C was 10,000

times higher than those maintained at 30°C (Fig. 3G). This established that higher titers

of phage particles were being produced in *S. praecaptivus* chloramphenicol resistant

clones at a temperature where P1CM*clr*-100(ts) becomes lytic. Finally, lysates derived

from *S. glossinidius* and *S. praecaptivus* chloramphenicol resistant clones, but not their

wild-type isogenic counterparts, promoted the formation of chloramphenicol resistant

E. coli cells at 30°C (Fig. 3H). This indicated that the chloramphenicol resistant marker

can be transduced from *S. glossinidius* and *S. praecaptivus* back to *E. coli*.

In *S. glossinidius*, the frequency of chloramphenicol resistant colonies arising

following P1CM*clr*-100(ts) exposure was similar to those observed for the *E. coli* control

cells, indicating that P1 infection occurs efficiently in this bacterium. By contrast,

chloramphenicol resistant *S. praecaptivus* colonies emerged at a lower frequency, and

higher concentrations of bacterial cells were typically used in P1 infection experiments

(see Materials and Methods). Notably, in P1-resistant *Salmonella enterica*, the efficiency

of P1 infection can be drastically increased by mutations in either *galU* or *galE* (Ornellas

and Stocker, 1974). Because these mutations remove the O-antigen by truncating the

core region of the lipopolysaccharide (LPS) (Wilkinson and Stocker, 1968; Osborn,

1968), they presumably facilitate access of P1 to its host receptor—conserved structural

motifs within the LPS core (Yarmolinsky and Sternberg, 1988; Ornellas and Stocker,

1974). In particular, while the LPS of *S. praecaptivus* contains structural components

attached to its core region, *S. glossinidius* is devoid of such structures (Fig. S1A).

Nonetheless, the lower infectivity of P1 does not appear to be related to the physical

occlusion of the P1 receptor by components present in the outer portion of the *S.*

praecaptivus LPS. This is because a mutation in *galU* results in a truncated LPS in *S.*

praecaptivus, but does not affect P1 infectivity (Fig. S1). Hence, unlike *S. enterica*, this

phenotype is not due to the presence of P1-antagonizing structure(s) in the outer

portion of *S. praecaptivus* LPS. Taken together, these results indicate that phage P1 is

capable of infecting and lysogenize in *S. glossinidius* and *S. praecaptivus*.

P1 generalized transduction in *Sodalis praecaptivus*

During the formation of P1 virions, approximately 0.05-0.5% of infective phage particles

package random DNA fragments derived from the bacterial host (Ikeda and Tomizawa,

1965). These particles can mediate the transfer of bacterial DNA across P1 susceptible

strains through generalized transduction. In the laboratory, generalized transduction of

DNA can be identified by virtue of genetic markers that are packaged in these phage

particles and transferred between bacterial strains. Accordingly, we sought to

determine if P1 could mediate generalized transduction in *S. praecaptivus*. First, we

exposed wild-type *S. praecaptivus* to phage lysates derived from a *S. praecaptivus*

P1CM*clr*-100(ts) lysogen harboring the ampicillin resistant (Amp^R) plasmid pSIM6

(Datta *et al.*, 2006). Following lysate exposure, we were able to retrieve Amp^R *S.*

praecaptivus transductants. Importantly, Amp^R cells were absent from both phage

lysates alone and cultures of wild-type *S. praecaptivus* that were not exposed to phage

(data not shown). In agreement with the notion that these Amp^R clones were P1

transductants, diagnostic PCR revealed the presence of a pSIM6 fragment in these cells

(Fig. 4A).

Next, we attempted to transduce a chromosomal chloramphenicol resistant

marker (*rpoS-HA::Cm*) using the P1 lytic strain P1*vir*. [This P1 strain is widely used as a

transducing agent in *E. coli* due to its inability to lysogenize cells upon infection and the

ease with which transducing lysates can be generated (Thomason *et al*. 2007; Ikeda and

Tomizawa, 1965)]. We infected wild-type *S. praecaptivus* cells with P1*vir* lysates grown

in a *S. praecaptivus rpoS-HA::Cm* pSIM6 strain. Whereas chloramphenicol resistant

(Cm^R) cells emerged from wild-type *S. praecaptivus* exposed to phage, no Cm^R cells were

obtained from phage lysates or cultures of naïve wild-type *S. praecaptivus* alone (data

not shown). Notably, diagnostic PCR indicated that chloramphenicol resistant clones

were transductants harboring the *rpoS-HA::Cm* genetic modification (Fig. 4B).

Importantly, all of these *rpoS-HA::Cm* transductants were sensitive to ampicillin,

indicating that P1*vir* mediated the transduction of a discrete portion of the *S.*

praecaptivus genome. Taken together, these results indicate that bacteriophage P1 can be

used to mediate generalized transduction in *S. praecaptivus*.

Introduction of exogenous DNA in *Sodalis glossinidius* and *Sodalis praecaptivus* by

P1-mediated guided transduction

Whereas up to 0.5% of P1 particles can contain random fragments of bacterial host DNA

(Ikeda and Tomizawa, 1965), the vast majority of virions harbor P1 DNA. This is

because the packaging of P1 genome into phage particles is guided by elements

encoded within its DNA sequence (Sternberg and Coulby, 1987a and 1987b).

Particularly, this packaging element can be cloned into plasmids (to produce

phagemids) or incorporated into the bacterial chromosome to increase the frequency of

P1-mediated transduction of adjacent DNA (Westwater *et al.*, 2002; Kittleson *et al.*, 2012;

Huang and Masters, 2014). Indeed, the P1 packaging element can increase the

transduction of linked DNA by 1,600-fold above the levels obtained in generalized

transduction (Kittleson *et al.*, 2012). Hence, this DNA element can be used to increase

the number of transducing particles and, consequently, the efficiency of DNA transfer

among bacterial strains that are susceptible to P1 infection.

The lack of genetic tools available for the manipulation of *Sodalis* species,

specifically *S. glossinidius*, prompted us to explore P1 as a plasmid DNA-delivery tool

for these bacteria. As a proof of principle, we used the aforementioned general

technique to transfer a number of P1 phagemids (Kittleson *et al.*, 2012) (Fig. 5A) into *S.*

glossinidius. We were able to recover transductants expressing an array of phenotypic

traits encoded in the phagemids. These included light production (*luxCDABE* genes),

violacein pigment synthesis (*vioABCE*), β-galactosidase activity (*lacZ*) or green

fluorescence (*gfp*) (Fig. 5B and 5C). To expand the tool set available for the modification

of *Sodalis* species, we constructed (1) two phagemids for tagging bacterial chromosomes

with fluorescent genes at the Tn7 attachment site (Peters and Craig, 2011), and (2) a

suicide phagemid encoding a Himar1 transposition system for random mutagenesis

(Rubin *et al.*, 1999) (Fig. S2). Following packaging into P1 virions in an *E. coli* P1CM*clr*-

100(ts) lysogen, these phagemids were efficiently delivered to *S. glossinidius* and *S.*

praecaptivus (Fig. 5D and 5E). Together, these results establish that bacteriophage P1 can

be used to efficiently deliver replication-competent and suicide vectors into *S.*

glossinidius and *S. praecaptivus* through a "guided transduction" strategy.

Discussion

In the current study, we demonstrate that bacteriophage P1 can infect, lysogenize and

promote transduction in two species of the genus *Sodalis*. We show that P1 can mediate

generalized transduction in *S. praecaptivus* (Fig. 4), and we establish that this

bacteriophage can be used for the delivery of plasmids and suicide vectors for the

genetic manipulation of *S. glossinidius* and *S. praecaptivus* (Fig. 5). While these results

constitute a significant advance in the development of genetic modification tools to

study these bacterial species, they also clear the way for the implementation of P1-based

DNA-delivery systems to uncultured *Sodalis* species (Heddi *et al.*, 1998; Fukatsu *et al.*,

2007; Nováková and Hypsa, 2007; Chrudimský *et al.*, 2012; Smith *et al.*, 2013; Santos-

Garcia *et a*l., 2017; Šochová *et al.*, 2017) and Gram negative insect endosymbionts

belonging to other genera.

Whereas *S. praecaptivus* can be genetically engineered with relative ease, the

ability of P1 to mediate generalized transduction provides a number of applications for

the manipulation of this bacterium. For instance, although *S. praecaptivus* can be readily

modified by recombineering functions of phage λ (λ-Red; Clayton *et al.* 2017; Enomoto

et al., 2017; Thomason *et al.*, 2014), the use of this technique has two major drawbacks.

First, the expression of recombineering functions can be mutagenic (Murphy and

Campellone, 2003). This can potentially produce confounding results in subsequent

experiments as phenotypes associated with a particular engineered modification may

actually results from secondary mutation(s). Second, typical temperature-sensitive

plasmids (rep_{pSC101}^{ts} *ori*) harboring recombineering functions cannot be cured from *S.*

praecaptivus by propagating cells at non-permissive temperatures ($\geq 37^\circ$C) in the absence

of plasmid selection (our unpublished results). The inability to cure these plasmids can

increase the chances of secondary mutations through leaky expression of

recombineering functions, and hinder the use of plasmids from the same

incompatibility group in downstream genetic analyses. Importantly, both of these

issues can be overcome by P1-mediated generalized transduction. That is, genomic

DNA fragments engineered using λ-Red can be transferred to naïve *S. praecaptivus* cells

that lack recombineering plasmids and, therefore, have not been exposed to potential

mutagenic events (Fig. 4B).

By contrast, the establishment of P1 mediated transduction provides a

considerable advancement in our ability to genetically manipulate *S. glossinidius*. This

is because *S. glossinidius* is recalcitrant to DNA transformation by standard techniques

such as heat-shock and electroporation (Pontes and Dale, 2006 and 2011; Kendra *et al*.,

2020). Whereas we have recently developed a method for DNA transfer to *S.*

glossinidius via conjugation (Kendra *et al*., 2020), P1-mediated transduction provides an

alternative, simpler method for the introduction of exogenous DNA into this bacterium.

Altered chromosomal fragments, replication competent plasmids carrying an array of

functions, and suicide vectors engineered for allelic replacement or containing

transposition systems can be quickly transduced into *S. glossinidius* in a simple protocol.

Given the large DNA packaging capability of P1 (up to 100 Kbp) (Thomason *et al*. 2007),

this bacteriophage can be efficiently used for a variety of applications, including the

delivery of bacterial artificial chromosomes (BAC vectors) or large plasmids encoding

multiple genome editing CRISPR systems (Adiego-Pérez *et al*., 2019) that are not easily

transferred by conjugation (Kendra *et al*., 2020). Additionally, the P1 packing sequence

can be incorporated into DNA fragments used in insertional mutagenesis (Huang and

Masters, 2014), enabling rapid and efficient combination of mutations via P1 "guided

transduction." This approach can greatly facilitate the implementation of several

analyses (e.g. complementation and epistasis) to identify and dissect genetic

components and pathways governing bacterial behaviors and interactions with

eukaryotic hosts.

Beyond the genus *Sodalis*, the results highlighted in this study have potential

broad implications for the genetic modification of uncultured Gram negative insect

endosymbionts. In Gram negative bacteria, the LPS is the major structural constituent

of the outer leaflet of the outer membrane. The LPS is composed of a highly conserved

lipid A "anchor," a conserved core polysaccharide and, sometimes, a hypervariable

outer component designated O-antigen (Fig. S1A; Silipo and Molinaro, 2010).

Bacteriophage P1 has a broad host range, in part, because it recognizes, as its host

receptor, structural features of the conserved LPS core (Yarmolinsky and Sternberg,

1988). In addition to *E. coli* and several species of the Gammaproteobacteria, P1 has

been shown to be capable of infecting various members within the Alpha, Beta- and

Deltaproteobacteria, and even bacterial species residing outside the proteobacteria

Phylum such as *Flavobacterium* sp. M64 (Yarmolinsky and Sternberg, 1988; Goldberg *et*

al., 1974; Ornellas and Stocker, 1974; Kaiser and Dworkin, 1975; Murooka and Harada,

1979).

Notably, a large number of economically important insect species—including

several disease vectors of animals and plants—harbor maternally inherited, Gram

negative bacterial endosymbionts (McCutcheon *et al.*, 2019; Moran *et al.*, 2008).

However, unlike *S. glossinidius* and a handful of other species, the vast majority of these

endosymbionts have not been isolated in pure culture (Dale and Maudlin, 1999;

Chrudimský *et al.*, 2012; Hypsa and Dale, 1997; Sabri *et al.*, 2011; Dale *et al.*, 2006; Brandt

et al., 2017; Masson *et al.*, 2018). This is because these bacteria undergo a process of

genome degeneration and size reduction during the course of long-term evolution and

specialization within their eukaryotic insect hosts. This process leads to the loss of

many physiological functions that are required for replication outside the host

(McCutcheon *et al.*, 2019; Moran *et al.*, 2008; Pontes *et al.*, 2011; Pontes and Dale, 2006).

Notably, classical protocols of bacterial genetics require the manipulation of large

numbers of cells and the subsequent isolation of rare genetic events as bacterial colonies

on selective agar plates. Consequently, the implementation of genetics for the study of

insect endosymbionts has remained scarce and limited to species that can grow in

axenic culture, form colonies on agar plates and are receptive to exogenous DNA

(Pontes and Dale, 2006; Kendra *et al.*, 2020; Masson *et al.*, 2018).

The ability of bacteriophage P1 to deliver DNA to a large number of bacterial

species, suggests a clear method in which the requirement for culturing may be

bypassed. That is, similar to viral vectors that are commonly utilized in gene therapy in

mammalians (Kay *et al.*, 2011), P1 could be used to deliver DNA to bacterial

endosymbionts inside their insect hosts. Specifically, insects could be microinjected (De

Vooght *et al.*, 2015; Pontes *et al.*, 2011; Boyle *et al.*, 1993; Oliver *et al.*, 2003; Doremus *et al.*,

2018) with P1 virions packaged with recombinant DNA. The establishment of

successful P1 infections and subsequent enrichment of transductant endsymbionts to

near-homogeneous or clonal populations could be attained by making use of

phenotypic markers (Fig. 5) and implementing antibiotic selection regiments in insects

(Wilkinson, 1998; Dale and Welburn, 2001; Dobson and Rattanadechakul, 2001; Koga *et*

al., 2007; Pais *et al.*, 2008; McLean *et al.*, 2011). Whereas this approach would

preferentially target recently acquired endosymbionts by virtue of their ability to exist

intra and extracellularly in various insect tissues (Aksoy *et al.*, 1997 ; Cheng and Aksoy,

1999; De Vooght *et al.*, 2015; Moran *et al.*, 2008), ancient obligate intracellular

endosymbionts could also be subjected to infection and genetic modification via P1, if

they transiently exit host cells (Zaidman-Rémy *et al.*, 2018).

The results presented in this study pave the way for the development of tractable

genetic systems for *S. glossinidius* and, potentially, a myriad of Gram negative bacterial

endosymbionts of insects. While this may empower the use of genetics to study these

obscure bacteria, it has also clear translational applications. P1-mediated DNA delivery

into insect endosymbionts may allow the engineering of bacterial traits aimed at

modifying aspects of insect ecology (Pontes and Dale, 2006; Kendra *et al.*, 2020),

mitigating their burden on economic activities and human health.

Material and Methods

Microbial strains, phages, plasmids and growth conditions

Microbial strains, phages and plasmids used in this study are presented in Table S1.

Unless indicated, all *E. coli* strains were propagated at 30, 37 or 42°C in Luria Bertani

(LB) broth or agar (1.5% w/v). *Sodalis glossinidius* was grown at 27°C in brain-heart

infusion broth supplemented with 10 mM $MgCl_2$ (BHI) or on brain-heart infusion agar

(1.2% w/v) supplemented with 10 mM $MgCl_2$ (BHI agar). *Sodalis glossinidius* was also

propagated on BHI agar plates supplemented with 10% defibrinated horse blood

(BHIB). *Sodalis praecaptivus* was grown at 30, 39 or 42°C in Luria Bertani (LB) broth or

agar (1.5% w/v) lacking sodium chloride. For experiments involving P1 infection or

generation of lysates, the growth media was supplemented with $CaCl_2$ and $MgCl_2$ to a

final concentration of 10 mM respectively. Growth of *Sodalis glossinidius* on BHIB agar

plates was carried out under microaerophilic conditions, which was achieved either

using BD GasPak EZ Campy Gas Generating sachets or a gas mixture (5% oxygen 95%

CO_2). For all strains, growth in liquid medium was carried out in shaking water bath

incubators with aeration (250 rpm). When required, medium was supplemented with

ampicillin (100 μg/mL), chloramphenicol (20 μg/mL for *E. coli* or *S. praecaptivus*, and 10

μg/mL for *S. glossinidius*), kanamycin (50 μg/mL for *E. coli* and 25 μg/mL for *S.*

glossinidius or *S. praecaptivus*). Arabinose was used at a concentration of 0.5 or 1%

(w/v); 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-gal) was used at a

concentration of 100 μg/mL.

Lipopolysaccharide extraction and detection

Extraction of lipopolysaccharide (LPS) from *S. glossinidius* and *S. praecaptivus* cultures

were carried out as described (Davis and Goldberg, 2012). Extracted samples were

separated in a NuPAGE 10% Bis-Tris gel in NuPAGE MES SDS Running (ThermoFisher

Scientific). LPS in gels were stained with ProteoSilver Silver Stain Kit (Sigma Aldrich).

Construction of phagemid pP1-Tn7-mCardinal

Oligonucleotide sequences used in this study are presented in Table S2. Phusion®

High-Fidelity DNA Polymerase (New England BioLabs) was used in PCR reactions

with primers 469 and 470 and plasmid BBa_J72113-BBa_J72152 (Kittleson *et al.*, 2012) as

template. The PCR product was ligated into pMRE-Tn7-163 (Schlechter *et al.*, 2018),

previously digested with SbfI, using NEBuilder® HiFi DNA Assembly (New England

BioLabs). The integrity of the construct was verified by DNA sequencing and the

ability to be efficiently transduced by P1 particles.

Construction of phagemid pP1-Tn7-mVenus

Oligonucleotide sequences used in this study are presented in Table S2. Phusion®

High-Fidelity DNA Polymerase (New England BioLabs) was used in PCR reactions

with primers 469 and 470 and plasmid BBa_J72113-BBa_J72152 (Kittleson *et al.*, 2012) as

template. The PCR product was ligated into pMRE-Tn7-166 (Schlechter *et al.*, 2018),

previously digested with SbfI, using NEBuilder® HiFi DNA Assembly (New England

BioLabs). The integrity of the construct was verified by DNA sequencing and the

ability to be efficiently transduced by P1 particles.

Construction of phagemid pP1-Himar

Oligonucleotide sequences used in this study are presented in Table S2. Phusion®

High-Fidelity DNA Polymerase (New England BioLabs) was used in PCR reactions

with primers 475 and 476 and plasmid BBa_J72113-BBa_J72152 (Kittleson *et al.*, 2012) as

template. The PCR product was ligated into pMarC9-R6k (Lee *et al.*, 2019), previously

digested with EcoRI and HinDIII, using NEBuilder® HiFi DNA Assembly (New

England BioLabs). The integrity of the construct was verified by DNA sequencing and

the ability to be efficiently transduced by P1 particles.

Recombineering procedure for *Sodalis praecaptivus*

Oligonucleotide sequences used in this study are presented in Table S2. A *S.*

praecaptivus strain harboring plasmid pSIM6 (Datta *et al.*, 2006) was grown overnight in

LB broth supplemented with 100 μg/ml of ampicillin at 30°C and 250 rpm.   Cells were

diluted (1:100) in 30 ml of the same medium and grown to an OD_{600} between 0.45-0.5.

The culture flask was then grown in a water bath at 42°C and 250 rpm for 25 min. Cells

were immediately transferred to a 50 ml conical tube, collected by centrifugation (7,000

rpm for 2.5 min at 4°C), and resuspended in 40 ml of ice-cold dH_2O. Cells were

collected again by centrifugation, and this washing procedure was repeated a second

time. Finally, cells were resuspended in 150 μl of ice-cold dH_2O. Homologous

recombination was obtained by electroporating 70 µl of cell suspension with 10 µl of

purified PCR products generated with primers 251 and 252 (*galU::Kn*) or 84 and 85

(*rpoS-HA::Cm*) and plasmids pKD4 and pKD3 (Datsenko and Wanner, 2000) as

templates, respectively.

Preparation of phage lysates derived from EMG16 P1 lysogens

Lysates from *E. coli* EMG16 harboring selected phagemids were prepared following

arabinose induction as described (Kittleson *et al.*, 2012).

Preparation of stocks P1*vir* phage lysates

Stocks of P1*vir* phage lysates were prepared by infecting *E. coli* MG1655 (Blattner *et al.*,

1997), as described (Thomason *et al.* 2007).

Preparation of phage lysates derived from P1CM*clr*-100(ts) lysogens

Escherichia coli P1CM*clr*-100(ts) lysogens were grown overnight at 30°C. Cultures were

diluted (1:100, v/v) into fresh medium and grown to and OD_{600} value of 0.3-0.4.

Subsequently, cultures were shifted to 42°C and propagated until extensive cell lysis (3-

4 h). At times, cultures were allowed to grow at 42°C for 16 h prior to the preparation of

lysates. Partially lysed cells were disrupted by vortexing the cultures following the

addition of chloroform (1 volume of chloroform per 100 volumes of culture). Cell

debris was removed by centrifugation (12 min, 4,000 x rpms, room temperature), and

the supernatant was passed through a 0.22 µm polyethersulfone membrane filter.

Sodalis praecaptivus P1CM*clr*-100(ts) lysogens were prepared as described above, except

that cells were grown for 2 h at 42°C and 16 h at 37°C prior to lysate preparation. *Sodalis*

glossinidius P1CM*clr*-100(ts) lysogens were grown in BHI to an OD600 of 0.4. Cultures

were heat shocked at 37 °C for 2 h. The cultures were treated with chloroform (1

volume of chloroform per 100 volumes of culture) and processed as described for *E. coli*

and *S. praecaptivus*.

Infection by P1*vir*, P1CM*clr*-100(ts) and P1 transducing particles

Following overnight grow in LB, *E. coli* cultures were diluted in fresh LB supplemented

with 10 mM $CaCl_2$ and $MgCl_2$ to an OD_{600} of 1. 1 mL aliquots of these cell solutions

were incubated for 30 minutes at 30°C in the absence or presence of various

concentrations of P1 lysate. Cells were subsequently collected by centrifugation (1 min,

13,000 x rpms, room temperature), and the supernatants were replaced by 1 ml of LB

containing 5 mM sodium citrate. Cells were grown for 1 h at 30°C and 250 rpm prior to

plating. *Sodalis praecaptivus* cells grown overnight in LB were collected by

centrifugation (1 min, 13,000 x rpms, room temperature) and resuspended in fresh LB

supplemented with 10 mM $CaCl_2$ and $MgCl_2$. One ml of these resuspended solutions

was incubated for 30 minutes at 30°C in the absence or presence of various

concentrations of P1 lysate. Cells were collected by centrifugation (1 min, 13,000 x

rpms, room temperature), and the supernatants were replaced by 1 ml of LB containing

5 mM sodium citrate. Cells were plated following 1 h of growth at 30°C and 250 rpm.

Sodalis glossinidius cells were grown in BHI for 3-5 days, to an $OD_{600} \approx 0.5$. Cells were

collected by centrifugation and concentrated to an OD_{600} of 1. One ml of concentrated

cultures was incubated for 60 minutes at 30°C in the absence or presence of various

concentrations of P1 lysate. Cells were collected by centrifugation (1 min, 13,000 x

rpms, room temperature), and the supernatants were replaced by 10 ml of BHI.

Cultures were incubated overnight at 27°C overnight with shaking prior to plating.

Curing of pP1-Tn7 phagemids

Transduction of pP1-Tn7 phagemids into *S. glossinidius* and *S. praecaptivus* were initially

selected on plates containing ampicillin (Fig. 5D, Fig. S2). Because episomes harboring

rep$_{pSC101}$^{ts} origins of replication are not easily cured from *S. praecaptivus* (see discussion),

the curing of phagemids was only performed in *S. glossinidius*. The strategy used to

identify *S. glossinidius* clones lacking phagemids was similar to the one adopted

elsewhere (Pontes and Dale, 2001). Briefly, to identify *S. glossinidius* clones that

contained the chloramphenicol resistant marker at the Tn7 attachment site and had loss

the ampicillin resistant plasmid, cultures were propagated in BHI containing

chloramphenicol and 1% arabinose. After 4 passages, cells were diluted and plated.

Single colonies were screened for sensitivity to ampicillin. Transposon insertion at the

Tn7 attachment site was verified by PCR with primers 1018 and 1019.

Image Acquisition, Analysis and Manipulation

DNA agarose gel electrophoresis and bacterial colonies, with the exception of *S.*

glossinidsius macrocolonies (were detected using an Amersham Imager 680 (GE

Healthcare). *Sodalis glossinidsius* macrocolonies expressing GFP were detected using a

dark reader (Clare Chemical Research) and documented with an iPhone. When

oversaturated, the intensity of signals in images were adjusted across the entire images

using Preview (Apple).

Acknowledgement

We would like to thank Serap Aksoy (Yale University) for kindly providing us with a

culture of *Sodalis glossinidius*, and Hubert Salvail (Yale University) for assistance

obtaining an *E. coli* P1CM*clr*-100(ts) lysogen. MHP is supported by grant AI148774

from the National Institutes of Health and startup funds from The Pennsylvania State

University College of Medicine.

Conflict of Interest

The authors declare no conflict of interest.

Supplemental Material

Figure S1.

Figure S2.

Table S1.

Table S2.

References

Adiego-Pérez, B., Randazzo, P., Daran, J.M., Verwaal, R., Roubos, J.A., Daran-Lapujade,

P., and van der Oost, J. (2019) Multiplex genome editing of microorganisms using

CRISPR-Cas. *FEMS Microbiol Lett* **366**:fnz086.

Aksoy, S., Chen, X., and Hypsa, V. (1997) Phylogeny and potential transmission routes

of midgut-associated endosymbionts of tsetse (Diptera:Glossinidae). *Insect Mol Biol*

6:183-190.

Bertani, G. (1951) Studies on lysogenesis. I. The mode of phage liberation by lysogenic

Escherichia coli. *J Bacteriol* **62**:293-300.

Blattner, F.R., Plunkett, G. 3rd, Bloch, C.A., Perna, N.T., Burland, V., Riley, M., *et al.*

(1997) The complete genome sequence of *Escherichia coli* K-12. *Science* **277**:1453-1462.

Boyle, L., O'Neill, S.L., Robertson, H.M., and Karr, T.L. (1993) Interspecific and

intraspecific horizontal transfer of *Wolbachia* in *Drosophila*. *Science* **260**:1796-1799.

Brandt, J.W., Chevignon, G., Oliver, K.M., and Strand, M.R. (2017) Culture of an aphid

heritable symbiont demonstrates its direct role in defence against parasitoids. *Proc Biol*

Sci **284**:20171925.

Butela, K., and Lawrence, J.G. (2012) Genetic manipulation of pathogenicity loci in non-

Typhimurium *Salmonella*. *J Microbiol Methods* **91**:477-482.

Chari, A., Oakeson, K.F., Enomoto, S., Jackson, D.G., Fisher, M.A., and Dale, C. (2015)

Phenotypic characterization of *Sodalis praecaptivus* sp. nov., a close non-insect-associated

member of the *Sodalis*-allied lineage of insect endosymbionts. *Int J Syst Evol Microbiol*

65:1400-1405.

Cheng, Q., and Aksoy, S. (1999) Tissue tropism, transmission and expression of foreign

genes *in vivo* in midgut symbionts of tsetse flies. *Insect Mol Biol* **8**:125-132.

Chrudimský, T., Husník, F., Nováková, E., and Hypša, V. (2012) *Candidatus* Sodalis

melophagi sp. nov.: phylogenetically independent comparative model to the tsetse fly

symbiont *Sodalis glossinidius*. *PLoS One* **7**:e40354.

Clayton, A.L., Oakeson, K.F., Gutin, M., Pontes, A., Dunn, D.M., von Niederhausern,

A.C., *et al*. (2012) A novel human-infection-derived bacterium provides insights into the

evolutionary origins of mutualistic insect-bacterial symbioses. *PLoS Genet* **8**:e1002990.

Clayton, A.L., Enomoto, S., Su, Y., and Dale, C. (2017) The regulation of antimicrobial

peptide resistance in the transition to insect symbiosis. *Mol Microbiol* **103**:958-972.

Dale, C., and Maudlin, I. (1999) *Sodalis* gen. nov. and *Sodalis glossinidius* sp. nov., a

microaerophilic secondary endosymbiont of the tsetse fly *Glossina morsitans morsitans*.

Int J Syst Bacteriol **49**:267-275.

Dale, C., and Welburn, S.C. (2001) The endosymbionts of tsetse flies: manipulating host-

parasite interactions. *Int J Parasitol* **31**:628-631.

Dale, C., Beeton, M., Harbison, C., Jones, T., and Pontes M. (2006) Isolation, pure

culture, and characterization of "*Candidatus* Arsenophonus arthropodicus," an

intracellular secondary endosymbiont from the hippoboscid louse fly *Pseudolynchia*

canariensis. *Appl Environ Microbiol* **72**:2997-3004.

Datsenko, K.A., Wanner, B.L. (2000) One-step inactivation of chromosomal genes in

Escherichia coli K-12 using PCR products. *Proc Natl Acad Sci USA* **97**:6640-6645.

Datta, S., Costantino, N., and Court, D.L. (2006) A set of recombineering plasmids for

gram-negative bacteria. *Gene* **379**:109-115.

Davis, M.R. Jr., and Goldberg, J.B. (2012) Purification and visualization of

lipopolysaccharide from Gram-negative bacteria by hot aqueous-phenol extraction. *J*

Vis Exp **63**:3916.

De Vooght, L., Caljon, G., Van Hees, J., Van Den Abbeele, J. (2015) Paternal transmission

of a secondary symbiont during mating in the viviparous tsetse fly. *Mol Biol Evol*

32:1977-1980.

Dobson, S.L., and Rattanadechakul, W. (2001) A novel technique for removing *Wolbachia*

infections from *Aedes albopictus* (Diptera: Culicidae). *J Med Entomol* **38**:844-849.

Doremus, M.R., Smith, A.H., Kim, K.L., Holder, A.J., Russell, J.A., and Oliver, K.M.

(2018) Breakdown of a defensive symbiosis, but not endogenous defences, at elevated

temperatures. *Mol Ecol* **27**:2138-2151.

Downard, J.S. (1988) Tn*5*-mediated transposition of plasmid DNA after transduction to

Myxococcus xanthus. *J Bacteriol* **170**:4939-4941.

Enomoto, S., Chari, A., Clayton, A.L., and Dale, C. (2017) Quorum sensing attenuates

virulence in *Sodalis praecaptivus*. *Cell Host Microbe* **21**:629-636.e5.

Fukatsu, T., Koga, R., Smith, W.A., Tanaka, K., Nikoh, N., Sasaki-Fukatsu, K.,

Yoshizawa, K., *et al.* (2007) Bacterial endosymbiont of the slender pigeon louse,

Columbicola columbae, allied to endosymbionts of grain weevils and tsetse flies. *Appl*

Environ Microbiol **73**:6660-6668.

Goldberg, R.B., Bender, R.A., Streicher, S.L. (1974) Direct selection for P1-sensitive

mutants of enteric bacteria. *J. Bacteriol.* **118**:810-814.

Heddi, A., Charles, H., Khatchadourian, C., Bonnot, G., and Nardon, P. (1998)

Molecular characterization of the principal symbiotic bacteria of the weevil *Sitophilus*

oryzae: a peculiar G+C content of an endocytobiotic DNA. *J Mol Evol* **47**:52-61.

Hendrix, R.W., Smith, M.C., Burns, R.N., Ford, M.E., Hatfull, G.F. (1999) Evolutionary

relationships among diverse bacteriophages and prophages: all the world's a phage.

Proc Natl Acad Sci USA **96**:2192-2197.

Hershey, A.D., and Chase, M. (1952) Independent functions of viral protein and nucleic

acid in growth of bacteriophage. *J Gen Physiol* **36**:39-56.

Huang, H., and Masters, M. (2014) Bacteriophage P1 *pac* sites inserted into the

chromosome greatly increase packaging and transduction of *Escherichia coli* genomic

DNA. *Virology* **468-470**:274-282.

Hypsa, V., and Dale, C. (1997) *In vitro* culture and phylogenetic analysis of "*Candidatus*

Arsenophonus triatominarum," an intracellular bacterium from the triatomine bug,

Triatoma infestans. *Int J Syst Bacteriol* **47**:1140-1144.

Ikeda, H., and Tomizawa, J.I. (1965) Transducing fragments in generalized transduction

by phage P1. I. Molecular origin of the fragments. *J Mol Biol* **14**:85-109.

Kaiser, D., and Dworkin, M. (1975) Gene transfer to myxobacterium by *Escherichia coli*

phage P1. *Science* **187**:653-653.

Kay, M.A., Glorioso, J.C., and Naldini, L. (2011) Viral vectors for gene therapy: the art of

turning infectious agents into vehicles of therapeutics. *Nat Med* **7**:33-40.

Kendra, C.G., Keller, C.M., Bruna, R.E., and Pontes, M.H. (2020) Conjugal DNA transfer

in the maternally inherited symbiont of tsetse flies *Sodalis glossinidius*. *mSphere* **5**:e00864-

20.

Kittleson, J.T., DeLoache, W., Cheng, H.Y., and Anderson, J.C. (2012) Scalable plasmid

transfer using engineered P1-based phagemids. *ACS Synth Biol* **1**:583-589.

Koga, R., Tsuchida, T., Sakurai, M., and Fukatsu, T. (2007) Selective elimination of aphid

endosymbionts: effects of antibiotic dose and host genotype, and fitness consequences.

FEMS Microbiol Ecol **60**:229-239.

Lee, H.H., Ostrov, N., Wong, B.G., Gold, M.A., Khalil, A.S., Church, G.M. (2019)

Functional genomics of the rapidly replicating bacterium *Vibrio natriegens* by CRISPRi.

Nat Microbiol **4**:1105-1113.

Lennox, E.S. (1955) Transduction of linked genetic characters of the host by

bacteriophage P1. *Virology* **1**:190-206.

Masson, F., Calderon Copete, S., Schüpfer, F., Garcia-Arraez, G., and Lemaitre, B. (2018)

In vitro culture of the insect endosymbiont *Spiroplasma poulsonii* highlights bacterial

genes Iinvolved in host-symbiont interaction. *mBio* **9**:e00024-18.

McCutcheon, J.P., Boyd, B.M., and Dale, C. (2019) The life of an insect endosymbiont

from the cradle to the grave. *Curr Biol* **29**:R485-R495.

McLean, A.H., van Asch, M., Ferrari, J., and Godfray, H.C. (2011) Effects of bacterial

secondary symbionts on host plant use in pea aphids. *Proc Biol Sci* **278**:760-766.

Moran, N.A., McCutcheon, J.P., and Nakabachi, A. (2008) Genomics and evolution of

heritable bacterial symbionts. *Annu Rev Genet* **42**:165-190.

Murooka, Y., and Harada, T. (1979) Expansion of the host range of coliphage P1 and

gene transfer from enteric bacteria to other gram-negative bacteria. *Appl Environ*

Microbiol **38**:754-757.

Murphy, K.C., and Campellone K.G. (2003) Lambda Red-mediated recombinogenic

engineering of enterohemorrhagic and enteropathogenic *E. coli*. *BMC Mol Biol* **4**:11.

Nováková, E., and Hypsa, V. (2007) A new *Sodalis* lineage from bloodsucking fly

Craterina melbae (Diptera, Hippoboscoidea) originated independently of the tsetse flies

symbiont *Sodalis glossinidius*. *FEMS Microbiol Lett* **269**:131-135.

O'Connor, K.A., and Zusman, D.R. (1983) Coliphage P1-mediated transduction of

cloned DNA from *Escherichia coli* to *Myxococcus xanthus*: use for complementation and

recombinational analyses. *J Bacteriol* **155**:317-329.

Oliver, K.M., Russell, J.A., Moran, N.A., and Hunter, M.S. (2003) Facultative bacterial

symbionts in aphids confer resistance to parasitic wasps. *Proc Natl Acad Sci USA*

100:1803-1807.

Ornellas, E.P., and Stocker, B.A. (1974) Relation of lipopolysaccharide character to P1

sensitivity in *Salmonella typhimurium*. *Virology* **60**:491-502.

Osborn, M.J. (1968) Biochemical characterization of mutants of *Salmonella typhimurium*

lacking glucosyl or galactosyl lipopolysaccharide transferases. *Nature* **217**:957-960.

Pais, R., Lohs, C., Wu, Y., Wang, J., and Aksoy, S. (2008) The obligate mutualist

Wigglesworthia glossinidia influences reproduction, digestion, and immunity processes of

its host, the tsetse fly. *Appl Environ Microbiol* **74**:5965-5974.

Peters, J.E., and Craig, N.L. (2011) Tn7: smarter than we thought. *Nat Rev Mol Cell Biol*

2:806-814.

Pontes, M.H., Dale, C. (2006) Culture and manipulation of insect facultative symbionts.

Trends Microbiol **14**:406-412.

Pontes, M.H., and Dale, C. (2011) Lambda red-mediated genetic modification of the

insect endosymbiont *Sodalis glossinidius*. *Appl Environ Microbiol* **77**:1918-1920.

Pontes, M.H., Smith, K.L., De Vooght, L., Van Den Abbeele, J., and Dale, C. (2011)

Attenuation of the sensing capabilities of PhoQ in transition to obligate insect-bacterial

association. *PLoS Genet* **7**:e1002349.

Rosner, J.L. (1972) Formation, induction, and curing of bacteriophage P1 lysogens.

Virology **48**:679-689.

Rubin, E.J., Akerley, B.J., Novik, V.N., Lampe, D.J., Husson, R.N., and Mekalanos, J.J.

(1999) *In vivo* transposition of mariner-based elements in enteric bacteria and

mycobacteria. *Proc Natl Acad Sci USA*. **96**:1645-1650.

Sabri, A., Leroy, P., Haubruge, E., Hance, T., Frère, I., Destain, J., and Thonart P. (2011)

Isolation, pure culture and characterization of *Serratia symbiotica* sp. nov., the R-type of

secondary endosymbiont of the black bean aphid *Aphis fabae. Int J Syst Evol Microbiol*

61:2081-2088.

Santos-Garcia, D., Silva, F.J., Morin, S., Dettner, K., and Kuechler, S.M. (2017) The all-

rounder *Sodalis*: a new bacteriome-associated endosymbiont of the lygaeoid bug

Henestaris halophilus (Heteroptera: Henestarinae) and a critical examination of its

evolution. *Genome Biol Evol* **9**:2893-2910.

Schlechter, R.O., Jun, H., Bernach, M., Oso, S., Boyd, E., Muñoz-Lintz, D.A., *et al*. (2018)

Chromatic bacteria - A broad host-range plasmid and chromosomal insertion toolbox

for fluorescent protein expression in bacteria. *Front Microbiol* **9**:3052.

Silipo, A., and Molinaro, A. (2010) The diversity of the core oligosaccharide in

lipopolysaccharides. *Subcell Biochem* **53**:69-99.

Smith, W.A., Oakeson, K.F., Johnson, K.P., Reed, D.L., Carter, T., Smith, K.L., *et al*.

(2013). Phylogenetic analysis of symbionts in feather-feeding lice of the genus

Columbicola: evidence for repeated symbiont replacements. *BMC Evol Biol* **13**:109.

Šochová, E., Husník, F., Nováková, E., Halajian, A., and Hypša, V. (2017) *Arsenophonus*

and *Sodalis* replacements shape evolution of symbiosis in louse flies. *PeerJ* **5**:e4099.

Soucy, S.M., Huang, J., Gogarten, J.P. (2015) Horizontal gene transfer: building the web

of life. *Nat Rev Genet* **16**:472-482.

Sternberg, N., and Coulby, J. (1987a) Recognition and cleavage of the bacteriophage P1

packaging site (*pac*). I. Differential processing of the cleaved ends *in vivo*. *J Mol Biol*

194:453-468.

Sternberg, N., and Coulby J. (1987b) Recognition and cleavage of the bacteriophage P1

packaging site (*pac*). II. Functional limits of *pac* and location of *pac* cleavage termini. *J*

Mol Biol **194**:469-479.

Streicher, S., Gurney, E., and Valentine, R.C. (1971) Transduction of the nitrogen-

fixation genes in *Klebsiella pneumoniae*. *Proc Natl Acad Sci USA*. **68**:1174-1177.

Thomason, L.C., Costantino, N., and Court, D.L. (2007) *E. coli* genome manipulation by

P1 transduction. *Curr Protoc Mol Biol* Chapter 1:Unit 1.17.

Thomason, L.C., Sawitzke, J.A., Li, X., Costantino, N., and Court, D.L. (2014)

Recombineering: genetic engineering in bacteria using homologous recombination. *Curr*

Protoc Mol Biol **106**:1.16.1-39.

Toh, H., Weiss, B.L., Perkin, S.A., Yamashita, A., Oshima, K., Hattori, M., and Aksoy, S.

(2006) Massive genome erosion and functional adaptations provide insights into the

symbiotic lifestyle of *Sodalis glossinidius* in the tsetse host. *Genome Res* **16**:149-156.

Touchon, M., Moura de Sousa, J.A., and Rocha, E.P. (2017) Embracing the enemy: the

diversification of microbial gene repertoires by phage-mediated horizontal gene

transfer. *Curr Opin Microbiol* **38**:66-73.

Westwater, C., Schofield, D.A., Schmidt, M.G., Norris, J.S., and Dolan, J.W. (2002)

Development of a P1 phagemid system for the delivery of DNA into Gram-negative

bacteria. *Microbiology* **148:**943-950.

Wilkinson, R.G., and Stocker, B.A. (1968) Genetics and cultural properties of mutants of

Salmonella typhimurium lacking glucosyl or galactosyl lipopolysaccharide transferases.

Nature **217:**955-957.

Wilkinson, T.L. (1998) The elimination of intracellular microorganisms from insects: an

analysis of antibiotic-treatment in the pea aphid (*Acyrthosiphon pisum*). *Comp Biochem*

*Physiol Ser A***119:**871-881.

Wolf-Watz, H., Portnoy, D.A., Bölin, I., and Falkow, S. (1985) Transfer of the virulence

plasmid of *Yersinia pestis* to *Yersinia pseudotuberculosis*. *Infect Immun* **48:**241-253.

Yarmolinsky, M.B., Sternberg, N. (1988) Bacteriophage P1. In The Bacteriophages.

Calendar, R. (ed). New York: Plenum Press, vol **1**. pp. 291-438.

Zaidman-Rémy, A., Vigneron, A., Weiss, B.L., and Heddi, A. (2018) What can a weevil

teach a fly, and reciprocally? Interaction of host immune systems with endosymbionts

in *Glossina* and *Sitophilus*. *BMC Microbiol* **18**(Suppl 1):150.

 Figure Legends

Fig. 1. Cartoon representation depicting a workflow of the transduction procedure

developed for introduction of phagemids in *Sodalis* species. Following the direction of

the arrows: An *E. coli* P1 lysogen host is transformed with a P1 phagemid. The

phagemid is packaged following induction of the P1 prophage, and lysates derived

from culture supernatant are used to infect a *Sodalis* recipient strain. Cells receiving the

phagemid are subsequently isolated on plates containing a selective agent.

Fig. 2. Infection of bacterial strains by phage P1. Lysates derived from an *E. coli*

P1CM*clr*-100(ts) lysogen (KL463) were used to infect *E. coli* MG1655 (A), *Sodalis*

glossinidius (B) and *Sodalis praecaptivus* (C). Plates depict the formation of

chloramphenicol resistant colonies as functions of the concentration of bacteria (vertical

axis) and the concentration of P1CM*clr*-100(ts) lysates (horizontal axis). Note that P1

infection conditions for the strains are different (see Materials and Methods), and

images do not reflect efficiency of P1 infection. Images show representative plates of at

least 3 routine experiments.

Fig. 3. Lysogenization and production of infective phage particles by *Sodalis glossinidius*

and *Sodalis praecaptivus* P1 lysogens. (A) Detection of P1 *pacB* gene by PCR and agarose

gel electrophoresis in *S. glossinidius* chloramphenicol resistant clones that emerged

following exposure to an *E. coli* P1CM*clr*-100(ts) lysogen (KL463). (B) Detection of a *S.*

glossinidius specific DNA fragment in clones depicted in (A) by PCR and agarose gel

electrophoresis. (C) Detection of P1 *pacB* gene by PCR and agarose gel electrophoresis

in *S. praecaptivus* chloramphenicol resistant clones that emerged following exposure to

an *E. coli* P1CM*clr*-100(ts) lysogen (KL463). (D) Detection of a *S. praecaptivus* specific

DNA fragment in clones depicted in (C) by PCR and agarose gel electrophoresis. (E-G)

Formation of phage plaques on soft agar embedded with *E. coli* MG1655. Soft agar

plates were spotted with dilutions of lysates derived from wild-type and *S. glossinidius*

chloramphenicol resistant (Cm^R) P1CM*clr*-100(ts) lysogen (MP1705) (E), wild-type and

S. praecaptivus Cm^R P1CM*clr*-100(ts) lysogen (MP1703) (F), and *S. praecaptivus* Cm^R

P1CM*clr*-100(ts) lysogen (MP1703) grown either at 37°C or 30°C. Plates are

representative of routine experiments. (H) Emergence of Cm^R *E. coli* MG1655 following

exposure to lysates derived from wild-type *S. glossinidius*, *S. glossinidius* Cm^R P1CM*clr*-

100(ts) lysogen (MP1705), wild-type *S. praecaptivus*, and *S. praecaptivus* Cm^R P1CM*clr*-

100(ts) lysogen (MP1703).   Plate is representative of at least 3 experiments.

Fig. 4. Bacteriophage P1-mediated generalized transduction in *S. praecaptivus*. (A)

Detection of a pSIM6 DNA region by PCR and agarose gel electrophoresis in ampicillin

resistant (Amp^R) *S. praecaptivus* transductants following exposure to lysates derived

from *S. praecaptivus* Cm^R P1CM*clr*-100(ts) (MP1703) harboring Amp^R plasmid pSIM6.

(B) Detection of the *rpoS*-HA::Cm chromosomal insertion by PCR and agarose gel

electrophoresis in *S. praecaptivus* transductants following exposure to lysates of P1*vir*

grown in *S. praecaptivus rpoS*-HA::Cm pSIM6 (MP1522) strain.

Fig. 5. Transduction of P1 phagemids into *Sodalis* species. (A) Schematic representation

of replication competent P1 phagemids encoding a number of phenotypic markers

(Kittleson *et al.*, 2012). (B) Comparison of wild-type (top quadrants) and *S. glossinidius*

transductants (lower quadrants) carrying P1 phagemid BBa_J72114-BBa_J72104

(Kittleson *et al.*, 2012). Transductant colonies are purple due to the expression of

violacein biosynthetic genes (lower, left quadrant), and produce light due to the

expression of bioluminescence genes (lower, right quadrant). (C) Macrocolonies

derived from wild-type *S. glossinidius* (bottom row) and transductants carrying P1

phagemids BBa_J72114-BBa_J72104 (*luxCDABE*+ and *vioABCE*+), BBa_J72114-BBa_J72100

(*lacZ*+) or BBa_J72113-BBa_J72152 (*gfp*+) (upper row). (D) Transduction of pP1-Tn7-

mVenus into *S. praecaptivus* (left-hand side plate) and pP1-Tn7-mCardinal into *S.*

glossinidius (right-hand side plate). (E). Transduction of phagemid encoding a Himar1

transposition system (pP1-Himar) into *S. praecaptivus* (left-hand side plate) and *S.*

glossinidius (right-hand side plate).

Fig. S1. Effect of a *galU* mutation on LPS composition and P1 infectivity in *S.*

praecaptivus. (A) Schematic representation depicting the inner membrane, the

peptidoglycan and the outer membrane of a prototypical Gram negative bacterium.

Inlet shows a more detailed schematic of structural components of the LPS (left-hand

side). Silver-stained SDS-PAGE of LPS purified from *S. praecaptivus galU* (CMK36),

wild-type *S. praecaptivus* and wild-type *S. glossinidius* (right-hand side). (B) Plates

depicting the appearance of chloramphenicol resistant colonies as functions of the

concentration of bacteria (vertical axis) and the concentration of P1CM*clr*-100(ts) lysates

(horizontal axis). Cm^R colonies emerge at similar frequencies in wild-type *S.*

praecaptivus (left-hand side plate) and *S. praecaptivus galU* (CMK36) (right-hand side

plate) when cells are exposed to equal amounts of P1CM*clr*-100(ts) particles.

Fig. S2. Schematic representation of P1 phagemids used for tagging the bacterial

chromosome at the Tn*7* attachment site(s) with genes encoding the fluorescent proteins

mCardinal (pP1-Tn*7*-mCardinal) or mVenus (pP1-Tn*7*-mVenus); and the suicide P1

phagemid for random mutagenesis with a Himar1 transposition system (pP1-Himar).

Figure 1

Figure 2

Figure 3

A P1 specific primers

(-) (+) WT

Kbp
0.6
0.4
0.2

S. glossinidius Cm^R clones

(+) (-) 1 2 3 4 5 6

Kbp
0.6
0.4
0.2

B *S. glossinidius* specific primers
Cm^R clones

(+) (-) 1 2 3 4 5 6

Kbp
0.5
0.3
0.1

C P1 specific primers

(-) (+) WT

Kbp
0.6
0.4
0.2

S. praecaptivus Cm^R clones

(+) (-) 1 2 3 4 5 6

Kbp
0.6
0.4
0.2

D *S. praecaptivus* specific primers
Cm^R clones

(+) (-) 1 2 3 4 5 6

Kbp
0.2
0.1

E WT Cm^R
[*S. glossinidius* lysate]
10^0
10^{-1}
10^{-2}
10^{-3}
10^{-4}

F WT Cm^R
[*S. praecaptivus* lysate]
10^0
10^{-1}
10^{-2}
10^{-3}
10^{-4}

G 37°C 30°C
[*S. praecaptivus* Cm^R lysate]
10^0
10^{-1}
10^{-2}
10^{-3}
10^{-4}

H *Phage lysates*

S. glossinidius   *S. praecaptivus*

E. coli cells | Cm^R lysate | Cm^R lysate + E.coli cells | wt lysate + E.coli cells | Cm^R lysate | Cm^R lysate + E.coli cells | wt lysate + E.coli cells

[*E. coli*]
10^0
10^{-1}
10^{-2}
10^{-3}

Figure 4

A

P1CM*clr*100(ts) grown in
S. praecaptivus pSIM6

B

P1*vir* grown in
S. praecaptivus rpoS-HA::Cm pSIM6

Figure 5

Figure S1

A

B

Figure S2

pP1-Tn7-mCardinal

pP1-Tn7-mVenus

pP1-Himar